Note on the evaluation of one type scalar one loop three-point amplitude inspired by $H \rightarrow gg$ decay in the standard model

Jin Zhang*

School of Physics and Engineering, Yuxi Normal University, Yuxi, Yunnan, 653100, P. R. China

Abstract

Motivated by the Higgs boson decaying to gg at one loop approximation, the amplitude of scalar one loop three-point diagram with two different internal masses are evaluated and fully analytic results are obtained. The main ingredient of the evaluation is a integral in which the integrand is product of the reciprocal of the integral variable and a logarithm whose argument is a quadratic function of the general form. The results depend on the choice of the masses of the propagators and the massive external line. In the first case, in addition to the elementary functions, the amplitude contains a term which is a summation over an infinite series where each term is a hypergeometric function. In the second case the result is expressed through dilogarithms. In particular, if the three internal lines are taking the same mass, the results will reduce to the known functions in one loop evaluation of Higgs decaying to gg or $\gamma\gamma$.

PACS numbers:

^{*} jinzhang@yxnu.edu.cn

I. INTRODUCTION

In the framework of the Standard Model(SM), evaluation of the scalar one loop three-point amplitudes[1, 2] play a fundamental role in deciphering the property of the Higgs boson[3, 4] though its decaying to $gg(\text{or }\gamma\gamma)$ and the inverse process, i.e., production of the Higgs boson by gluon fusion[5–9]. Owing to the coupling of the Higgs boson to the fermions $g_{Hf\bar{f}}$ and the coupling of fermions to gluon, the three propagators take the same mass at the leading order of perturbative theory. In addition to the top quark, there is considerable mass difference between the Higgs boson and other quarks, an economic way to compute the amplitude is taking the limit that $m_H \to \infty$ such that the amplitude tends to a constant[11, 12]. The advantage of this manner is convenient for evaluation, but we lose the chance to compare the different contributions from various competing processes to the amplitude. Fully analytic results are essential to analyze $H \to gg$. The evaluated amplitude, if the high energy approximation is not exploited, will be expressed as function of m_f/m_H , where m_f and m_H are the masses of the internal fermion and the Higgs boson, respectively. At the final stage of the evaluation, a integral of the following type necessarily arises and must be handled carefully

$$I_1 = \int_0^1 dx \, \frac{\ln(ax^2 - ax + 1 - i\varepsilon)}{x}, \quad a > 0$$
 (1)

where ε is a positive infinitesimal. Then the results will be distinguished the cases 0 < a < 4 from the case a > 4 as detailed in the later works[7, 13, 15–19].

As a natural generalization of Eq.(1), let us consider the following integral

$$I_2 = \int_0^1 dx \, \frac{\ln(ax^2 - bx + 1 - i\varepsilon)}{x},$$
(2)

the parameters a and b satisfy

$$a > 0, \quad b > 0, \quad a \neq b \tag{3}$$

It is obvious that if a = b, Eq.(2) reduce to Eq.(1). In fact, this integral can be derived from the evaluation of the scalar one loop three-point diagram depicted in fig.1, in this case a and bwill be functions of the masses of the propagators ω_1 and ω_1 as well as the mass of external line m_1 . However, a close inspection to fig.1 indicates that it does not connect with real decaying processes of the Higgs boson even though the contribution from Higgs-Kibble ghosts associated with the W^{\pm} and Z bosons are taken into account. Maybe it is the reason that rarely can we find



FIG. 1: Massive triangle with two massless external lines. The solid lines and dashed lines denote massive and massless particles, respectively.

a specialized analysis on the integral displayed in Eq.(2) the Standard Model¹. In the minimal supersymmetric extension of the Standard Model, the integral in Eq.(2) can be extracted from the evaluation of radiative neutralino decay at one loop[20]². We notice that an obvious and systematic analysis on the integral in Eq.(3) is still lacking, hence in this paper we will present a thorough and specialized study on Eq.(2), fully analytic results are derived. Although the integral in Eq.(2) may be evaluated by computer algebra[?], a complete analytic evaluation is helpful, especially for the beginner of the Higgs physics from the pedagogical perspective. We hope that the results can be employed to some decaying process under reasonable approximation addition to $H \rightarrow gg$, but also enrich the results of scalar one loop three-point diagrams from the viewpoint of analytic evaluation.

The paper is organized as follows. In section II we introduce the integral in Eq.(2) by evaluation the amplitude depicted in fig.1 in a scalar field theory, some general results are derived. In section III the analytic results of fig.1 are obtained and the special case a = b are discussed in details. A short summary are presented in IV. Some useful formulas are listed in the appendix.

¹ Integral of this type has been computed in Eq.(23) of Ref.[10], but only the case $b^2 - 4a > 0$ is considered.

² There were typographical errors in Eq.(D.6) and Eq.(D.9) of the paper, which are corrected in Appendix D of a note which can be found in the webpage: http://scipp.ucsc.edu/~haber/webpage/oneloopfun4.pdf.

II. THE FORMULAS

A. The massive triangle with two massless external lines

To start with, we write down the amplitude corresponding to fig.1

$$I = \int \frac{\mathrm{d}^4 k}{(2\pi)^4} \frac{1}{A_1 A_2 A_3},\tag{4}$$

where the three denominators are defined by

$$A_{1} = k^{2} - \omega_{1}^{2} + i\varepsilon$$

$$A_{2} = (p_{1} - k)^{2} - \omega_{2}^{2} + i\varepsilon$$

$$A_{3} = (p_{1} - p_{2} - k)^{2} - \omega_{2}^{2} + i\varepsilon,$$
(5)

and ε is real positive infinitesimal, the three external momentum satisfy

$$p_1^2 = m_1^2, \quad p_2^2 = p_3^2 = 0.$$
 (6)

Using the Feynman's trick, Eq.(4) can be written as

$$I = \int \frac{\mathrm{d}^4 k}{(2\pi)^4} \int \mathrm{d}x \mathrm{d}y \mathrm{d}z \frac{2! \,\delta(1 - x - y - z)}{[D(x, y, z)]^3},\tag{7}$$

where

$$D(x, y, z) = x(k^2 - \omega_1^2 + i\varepsilon) + y[(p_1 - k)^2 - \omega_2^2 + i\varepsilon] + z[(p_1 - p_2 - k)^2 - \omega_2^2 + i\varepsilon], \quad (8)$$

Since the amplitude given by Eq.(4) is both ultraviolet and infrared finite, thus regularization is unnecessary, the evaluation can be carried out in the four-dimensional space-time. We first perform the integral over k and z, obtaining

$$I = -\frac{i}{16\pi^2} \int_0^1 \mathrm{d}x \int_0^{1-x} \mathrm{d}y \frac{1}{-yxm_1^2 + x(\omega_1^2 - \omega_2^2) + \omega_2^2 - i\varepsilon}$$
(9)

The integral over y in Eq.(9) is trivial, combining with Eq.(A1), we arrive at the following intermediate result

$$I = \frac{i}{16\pi^2 m_1^2} \int_0^1 \mathrm{d}x \frac{1}{x} \Big[\ln\left(\frac{m_1^2}{\omega_2^2} x^2 - \frac{m_1^2 - \omega_1^2 + \omega_2^2}{\omega_2^2} x + 1 - i\varepsilon\right) - \ln\left(\frac{\omega_1^2 - \omega_2^2}{\omega_2^2} x + 1 - i\varepsilon\right) \Big]$$

$$= \frac{i}{16\pi^2 m_1^2} \Big[\mathrm{Li}_2 \Big(1 - \frac{\omega_1^2}{\omega_2^2} \Big) + \int_0^1 \mathrm{d}x \frac{1}{x} \ln\left(\frac{m_1^2}{\omega_2^2} x^2 - \frac{m_1^2 - \omega_1^2 + \omega_2^2}{\omega_2^2} x + 1 - i\varepsilon\right) \Big]. \tag{10}$$

The remaining work is the evaluation of the last integral in Eq.(10). For brevity, in the forthcoming sections the pre-factor $i/(16\pi^2)$ will be suppressed while $1/m_1^2$ will be preserved so as to maintain the correct dimension of the primitive amplitude displayed in Eq.(4).

B. evaluation of integral with logarithms

The evaluation of the last term in Eq.(10) motivates a general investigation on the integral of the following type

$$F = \int_0^1 \mathrm{d}x \, \frac{\ln(ax^2 - bx + 1 - i\varepsilon)}{x}, \quad a > 0, \quad b > 0 \tag{11}$$

Since the argument of the logarithm is quadratic in x, we first consider the case $b^2 - 4a < 0$, in this case the argument of the logarithm is positive definite thus the $i\varepsilon$ term can be safely dropped. A feasible way to calculate the integral of Eq.(11) turns out to be expressing it as

$$F = \int_{0}^{1} dx \int_{0}^{1} dz \frac{ax - b}{1 + zx(ax - b)}$$

= $a \int_{0}^{1} dz \int_{0}^{1} dx \frac{x}{1 + zx(ax - b)} - b \int_{0}^{1} dz \int_{0}^{1} dx \frac{1}{1 + zx(ax - b)}$
= $\frac{1}{2} \int_{0}^{1} dz \frac{\ln[1 + z(a - b)]}{z} - \frac{b}{2} \int_{0}^{1} dz \int_{0}^{1} dx \frac{1}{1 + zx(ax - b)}$
= $-\frac{1}{2} \text{Li}_{2}(b - a) - \frac{b}{2} \int_{0}^{1} dz \int_{0}^{1} dx \frac{1}{1 + zx(ax - b)}.$ (12)

Now we concentrate on the last integral in Eq.(12), for later convenience we label it as A, the integral over x can be calculated[21]

$$A = \int_{0}^{1} dz \int_{0}^{1} dx \frac{1}{1 + zx(ax - b)}$$

=
$$\int_{0}^{1} dz \frac{2}{\sqrt{4az - b^{2}z^{2}}} \left(\arctan \frac{bz}{\sqrt{4az - b^{2}z^{2}}} + \arctan \frac{2az - bz}{\sqrt{4az - b^{2}z^{2}}} \right),$$
(13)

In deriving Eq.(13) we employ the property that $\arctan(x)$ is odd

$$\arctan(-x) = -\arctan(x),\tag{14}$$

To proceed we separate the integral in Eq.(13) into two parts

$$A = 2(A_1 + A_2), (15)$$

where

$$A_1 = \int_0^1 dz \frac{1}{\sqrt{4az - b^2 z^2}} \arctan \frac{bz}{\sqrt{4az - b^2 z^2}},$$

$$A_2 = \int_0^1 dz \frac{1}{\sqrt{4az - b^2 z^2}} \arctan \frac{2az - bz}{\sqrt{4az - b^2 z^2}}.$$
 (16)

It is not difficult to demonstrate that

$$\left[\arcsin\left(b\sqrt{\frac{z}{4a}}\right)\right]' = \left(\arctan\frac{bz}{\sqrt{4az - b^2 z^2}}\right)' = \frac{b}{2}\frac{1}{\sqrt{4az - b^2 z^2}},\tag{17}$$

Hence, the integral in A_1 is easy to calculate

$$A_1 = \frac{1}{b} \left(\arctan\frac{b}{\sqrt{4a - b^2}}\right)^2,\tag{18}$$

Using the identity [23]

$$\arctan x = \arcsin \frac{x}{\sqrt{1+x^2}},\tag{19}$$

leads to the final result for A_1

$$A_1 = \frac{1}{b} \left(\arcsin \frac{b}{2\sqrt{a}} \right)^2. \tag{20}$$

Next, we evaluate A_2 , by making use integration by parts, getting

$$A_{2} = \frac{2}{b} \arcsin(\frac{b}{2\sqrt{a}}) \arctan\frac{2a-b}{\sqrt{4a-b^{2}}} - \frac{2a-b}{b} \int_{0}^{1} \frac{\arcsin\left(b\sqrt{\frac{z}{4a}}\right)}{[1+(a-b)z]\sqrt{4az-b^{2}z^{2}}} \,\mathrm{d}z, \quad (21)$$

We label the second term in Eq.(21) as B and define

$$u = b\sqrt{\frac{z}{4a}},\tag{22}$$

such that this term casts into the following form

$$B = \frac{2a-b}{b^2} \int_0^{b/(2\sqrt{a})} \frac{\arcsin u}{(1+\alpha u^2)\sqrt{1-u^2}} \,\mathrm{d}u, \qquad \alpha = \frac{4a(a-b)}{b^2}$$
(23)

Since 0 < u < 1, we employ the following expansion[24]

$$\frac{\arcsin u}{\sqrt{1-u^2}} = \sum_{n=0}^{+\infty} \frac{2^n n!}{(2n+1)!!} u^{2n+1},\tag{24}$$

Plugging Eq.(24) into Eq.(23), interchanging the order of integration and summation, yields

$$B = \frac{2a-b}{b^2} \sum_{n=0}^{+\infty} \frac{2^n n!}{(2n+1)!!} \int_0^{b/(2\sqrt{a})} \frac{u^{2n+1}}{1+\alpha u^2} \,\mathrm{d}u,$$
(25)

It is convenient to make variable transformation for the second time

$$\xi = u^2, \tag{26}$$

Then we obtain

$$B = \frac{2a-b}{b^2} \sum_{n=0}^{+\infty} \frac{2^n n!}{(2n+1)!!} \int_0^{b^2/(4a)} \frac{\xi^n}{1+\alpha\xi} \mathrm{d}\xi,$$
(27)

Changing the variable further

$$\xi = \frac{b^2}{4a}\rho,\tag{28}$$

Eq.(27) can be written as

$$B = \frac{2a-b}{b^2} \sum_{n=0}^{+\infty} \frac{2^n n!}{(2n+1)!!} \left(\frac{b^2}{4a}\right)^{n+1} \int_0^1 \rho^n \left(1 + \frac{\alpha b^2}{4a}\rho\right)^{-1} \mathrm{d}\rho, \tag{29}$$

Comparing with Eq.(A3), we identify

$$\alpha = 1, \quad \beta = n+1, \quad \gamma = n+2, \quad z = b-a,$$
(30)

which generates the following result for B

$$B = \frac{2a-b}{2a} \sum_{n=0}^{+\infty} \frac{2^n n!}{(n+1)(2n+1)!!} \left(\frac{b^2}{4a}\right)^n {}_2F_1(1,n+1;n+2;b-a),$$
(31)

Combining Eq.(20), Eq.(21) and Eq.(31), we arrive at the final result of F in the case $b^2 - 4a < 0$

$$F = -\frac{1}{2} \text{Li}_2(b-a) - \left(\arcsin\frac{b}{2\sqrt{a}}\right)^2 - 2\arcsin\left(\frac{b}{2\sqrt{a}}\right) \arctan\frac{2a-b}{\sqrt{4a-b^2}} + \frac{(2a-b)b}{2a} \sum_{n=0}^{+\infty} \frac{2^n n!}{(n+1)(2n+1)!!} \left(\frac{b^2}{4a}\right)^n {}_2F_1(1,n+1;n+2;b-a).$$
(32)

Next, we consider the case $b^2 - 4a > 0$. In this case there are two zeros of the logarithm in the range [0, 1], the $i\varepsilon$ prescription must be retained appropriately. By exploiting integration by parts, it is easy to get

$$F = -\int_0^1 \frac{(2ax - b)\ln x}{a(x - x_1)(x - x_2)} \,\mathrm{d}x,$$
(33)

where x_1 and x_2 are the two roots of the argument of the logarithm

$$x_1 = x_+ + i\varepsilon, \qquad x_+ = \frac{b + \sqrt{b^2 - 4a}}{2a},$$

 $x_2 = x_- - i\varepsilon, \qquad x_- = \frac{b - \sqrt{b^2 - 4a}}{2a},$
(34)

Making use partial fraction expansion and Eq.(??), we obtain the following result

$$F = \frac{1}{x_1 - x_2} \left(\frac{b}{a} - 2x_1\right) \int_0^1 \frac{\ln x}{x - x_1} \, \mathrm{d}x + \frac{1}{x_1 - x_2} (2x_2 - \frac{b}{a}) \int_0^1 \frac{\ln x}{x - x_2} \, \mathrm{d}x$$

$$= \frac{1}{x_1 - x_2} \left\{ \left(\frac{b}{a} - 2x_1\right) \mathrm{Li}_2 \left[\frac{1}{x_+} - i\varepsilon \operatorname{sgn}(x_+)\right] - \left(\frac{b}{a} - 2x_2\right) \mathrm{Li}_2 \left[\frac{1}{x_-} + i\varepsilon \operatorname{sgn}(x_-)\right] \right\}.$$
(35)

with the function sgn(x) defined in Eq.(A3).

III. RESULTS AND DISCUSSIONS

We shall now apply the results obtained in Section 2 to calculate the integral left in Eq.(10). Comparing Eq.(10) and Eq.(11), we identify that

$$a = \frac{m_1^2}{\omega_2^2}, \qquad b = \frac{m_1^2 - \omega_1^2 + \omega_2^2}{\omega_2^2},$$
(36)

In the case $b^2 - 4a < 0$ which implies that $\lambda(m_1^2, \omega_1^2, \omega_2^2) < 0$, where $\lambda(x, y, z)$ is the well-known Källen function

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz,$$
(37)

By employing Eq.(32), yields the following explicit result

$$I = \frac{1}{m_1^2} \left\{ \frac{1}{2} \operatorname{Li}_2 \left(1 - \frac{\omega_1^2}{\omega_2^2} \right) - \left(\arcsin \frac{m_1^2 - \omega_1^2 + \omega_2^2}{2m_1 \omega_2} \right)^2 - 2 \arcsin \frac{m_1^2 - \omega_1^2 + \omega_2^2}{2m_1 \omega_2} \operatorname{arctan} \frac{m_1^2 + \omega_1^2 - \omega_2^2}{\sqrt{\lambda(m_1^2, \omega_1^2, \omega_2^2)}} + \frac{m_1^4 - (\omega_1^2 - \omega_2^2)^2}{8m_1^2 \omega_2^2} \left[\sum_{n=0}^{+\infty} \frac{2^n n!}{(n+1)(2n+1)!!} \left(\frac{(m_1^2 - \omega_1^2 + \omega_2^2)^2}{4m_1^2 \omega_1^2} \right)^n \right] \times {}_2F_1 \left(1, n+1; n+2; 1 - \frac{\omega_1^2}{\omega_2^2} \right) \right\}.$$
(38)

In order to apply the result in Eq.(38) correctly, the following comments are necessary. First, from the conditions of a and b declared in Eq.(11), to guarantee Eq.(38) holds true for the evaluation, in addition to $\lambda(m_1^2, \omega_1^2, \omega_2^2) < 0$, the masses must obey

$$m_1^2 > \omega_1^2 - \omega_2^2, \tag{39}$$

Second, since Eq.(38) is summed over hypergeometric functions, a crucial issue is that if the summation of the infinite series is convergent. Due to $\lambda(m_1^2, \omega_1^2, \omega_2^2) < 0$, it is obvious that

$$0 < \frac{(m_1^2 - \omega_1^2 + \omega_2^2)^2}{4m_1^2 \omega_1^2} < 1.$$
(40)

and the hypergeometric functions are always taking finite value, thus the summation is convergent. Finally, in considering the analytic property of the dilogarithm in Eq.(A2), a question is that if Eq.(38) can develop imaginary part. In other words, if the argument of the dilogarithm can be greater than 1. But it is impossible since both ω_1 and ω_2 are assumed to be real, thus $0 < 1 - (\omega_1^2/\omega_2^2) < 1$ is always satisfied, therefore there is no imaginary part can be developed. In the case $b^2 - 4a > 0$, this implies that $\lambda(m_1^2, \omega_1^2, \omega_2^2) > 0$, exploiting Eq.(35) we get

$$I = \frac{1}{m_1^2} \Big\{ \frac{1}{2} \operatorname{Li}_2(1 - \frac{\omega_1^2}{\omega_2^2}) - \operatorname{Li}_2[\frac{1}{x_+} - i\varepsilon \operatorname{sgn}(x_+)] - \operatorname{Li}_2[\frac{1}{x_-} + i\varepsilon \operatorname{sgn}(x_-)] \Big\},\tag{41}$$

where

$$x_{+} = \frac{(m_{1}^{2} - \omega_{1}^{2} + \omega_{2}^{2}) + \lambda^{1/2}(m_{1}^{2}, \omega_{1}^{2}, \omega_{2}^{2})}{2m_{1}^{2}},$$

$$x_{-} = \frac{(m_{1}^{2} - \omega_{1}^{2} + \omega_{2}^{2}) - \lambda^{1/2}(m_{1}^{2}, \omega_{1}^{2}, \omega_{2}^{2})}{2m_{1}^{2}},$$
(42)

Since $m_1^2 > \omega_1^2 - \omega_2^2$ as presented in Eq.(39), both x_+ and x_- are positive definite, thus Eq.(41) simplified to

$$I = \frac{1}{m_1^2} \Big[\frac{1}{2} \operatorname{Li}_2(1 - \frac{\omega_1^2}{\omega_2^2}) - \operatorname{Li}_2(\frac{1}{x_+} - i\varepsilon) - \operatorname{Li}_2(\frac{1}{x_-} + i\varepsilon) \Big].$$
(43)

In order to explore phenomenological implications of the results presented in Eq.(38) and Eq.(43), it is instructive to consider the special case that the coefficients in Eq.(11) are constrained by the following conditions

$$a = b > 0, \tag{44}$$

This is the integral indispensable in the evaluation of $H \to gg$ decay at one loop approximation. Supposing the mass of each propagator of the triangle is ω_1 , setting

$$a = b = \frac{m_1^2}{\omega_1^2},$$
(45)

Hence, Eq.(10) reduces to

$$I = \frac{1}{m_1^2} \int_0^1 \mathrm{d}x \frac{\ln(ax^2 - ax + 1 - i\varepsilon)}{x}, \qquad a = \frac{m_1^2}{\omega_1^2}$$
(46)

First consider the case 0 < a < 4, i.e., $0 < m_1 < 2\omega_1$. To get the correct results we must trace back to Eq.(12) and Eq.(13), other than taking advantage of Eq.(32) and simply setting a = b, otherwise it will make some mistakes. By exploiting Eq.(17), we immediately obtain

$$I = -\frac{a}{2m_1^2} \int_0^1 dz \int_0^1 dx \frac{1}{1+zax(x-1)}$$

= $-\frac{a}{2m_1^2} \int_0^1 dz \frac{4}{\sqrt{4az-a^2z^2}} \arctan \frac{az}{\sqrt{4az-a^2z^2}}$
= $-\frac{2}{m_1^2} \left(\arctan \frac{a}{\sqrt{4a-a^2}}\right)^2$, (47)

By using Eq.(19), we get the well-known function appeared in one loop evaluation of $H \to gg$

$$I = -\frac{2}{m_1^2} \left(\arcsin \frac{m_1}{2\omega_1} \right)^2, \qquad 0 < m_1 < 2\omega_1.$$
(48)

Next, we consider the case of a > 4, i.e., $m_1 > 2\omega_1$. From Eq.(35) we get

$$I = \frac{1}{m_1^2} \left[-\operatorname{Li}_2(\frac{1}{x_-} + i\varepsilon) - \operatorname{Li}_2(\frac{1}{x_+} - i\varepsilon) \right],\tag{49}$$

 x_+ and x_- are given by

$$x_{+} = \frac{1}{2} \left(1 + \sqrt{1 - \frac{4}{a}} \right), \qquad x_{-} = \frac{1}{2} \left(1 - \sqrt{1 - \frac{4}{a}} \right), \tag{50}$$

Since both x_+ and x_- are less than 1, in order to simplify the Eq.(49), defining

$$\alpha = \frac{1}{x_{-}} > 1, \quad \frac{1}{x_{+}} = \frac{1}{1 - x_{-}} = \frac{\alpha}{\alpha - 1} > 1,$$
(51)

Then Eq.(49) can be written as

$$I = \frac{1}{m_1^2} \Big[-\operatorname{Li}_2(\alpha + i\varepsilon) - \operatorname{Li}_2(\frac{\alpha}{\alpha - 1} - i\varepsilon) \Big],$$
(52)

Combining Eq.(A2) we obtain

$$I = \frac{1}{m_1^2} \left\{ -\left[\operatorname{Re}\operatorname{Li}_2(\alpha) + i\pi \ln \alpha\right] - \left[\operatorname{Re}\operatorname{Li}_2(\frac{\alpha}{\alpha - 1}) - i\pi \ln \frac{\alpha}{\alpha - 1}\right] \right.$$

$$= \frac{1}{m_1^2} \left\{ -\operatorname{Re}\left[\operatorname{Li}_2(\alpha) + \operatorname{Li}_2(\frac{\alpha}{\alpha - 1})\right] - i\pi \ln(\alpha - 1) \right\},$$
(53)

In considering the property of the dilogarithm

$$\operatorname{Re}\left[\operatorname{Li}_{2}(x) + \operatorname{Li}_{2}(\frac{x}{x-1})\right] = \frac{\pi^{2}}{2} - \frac{\ln^{2}(x-1)}{2}, \quad x > 1$$
(54)

which leads to

$$I = \frac{1}{m_1^2} \Big[-\frac{\pi^2}{2} + \frac{\ln^2(\alpha - 1)}{2} - i\pi \ln(\alpha - 1) \Big],$$
(55)

Substituting Eq.(51) into Eq.(55) and noticing that

$$\alpha - 1 = \frac{x_+}{x_-},\tag{56}$$

we get

$$I = \frac{1}{m_1^2} \left(-\frac{\pi^2}{2} + \frac{1}{2} \ln^2 \frac{x_+}{x_-} - i\pi \ln \frac{x_+}{x_-} \right), \tag{57}$$

Plugging Eq.(50) into Eq.(57) we arrive at the well-known function in the one loop evaluation of $H \rightarrow gg$ decay

$$I = -\frac{1}{2m_1^2} \left(\pi + i \ln \frac{1 + \sqrt{1 - \frac{4\omega_1^2}{m_1^2}}}{1 - \sqrt{1 - \frac{4\omega_1^2}{m_1^2}}} \right)^2. \quad m_1 > 2\omega_1$$
(58)

IV. SUMMARY

In this paper, the amplitude of the scalar one loop three-point diagram in which the three propagators are assigned two different masses is evaluated. A general type integral is extracted and analytic results are obtained. The conditions for correct applications of the results are discussed in details. If each propagator of the triangle taking the same mass, we find that the general results will simplify to the functions obtained in the lowest order evaluation of $H \rightarrow gg$ decay. The results and techniques presented in this paper can be applied also to decaying processes which are mediated by one loop triangle diagram.

Acknowledgements

The author thanks H. E. Haber for reminding the work in Ref.[20] and the updated version in his webpage.

Appendix A: The dilogarithm function and integral representation of Gauss hypergeometric function

In this section we list some necessary formula in our evaluation. The dilogarithm is defined as[22]

$$\operatorname{Li}_{2}(x) = \sum_{k=1}^{+\infty} \frac{x^{k}}{k^{2}} = -\int_{0}^{1} \frac{\ln(1-xt)}{t} \,\mathrm{d}t, \quad |x| < 1$$
(A1)

There is a branch cut from 1 to ∞ , for $\varepsilon \to 0$

$$\operatorname{Li}_{2}(x+i\varepsilon) = \operatorname{Re}\operatorname{Li}_{2}(x) + i\pi\operatorname{sgn}(\varepsilon)\Theta(x-1)\ln x, \qquad (A2)$$

where Θ is the step function, the sgn(x) is

$$\operatorname{sgn}(x) = \begin{cases} 1 & x > 0 \\ -1 & x < 0 \end{cases}$$

Another formula needed is the integral representation of the Gauss hypergeometric function [25, 26]

$${}_{2}F_{1}(\alpha,\beta;\gamma;z) = \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\gamma-\beta)} \int_{0}^{1} t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-zt)^{-\alpha} \mathrm{d}t, \tag{A3}$$

where

$$\operatorname{Re}(\gamma) > \operatorname{Re}(\beta) > 0, \quad |\operatorname{arg}(1-z)| < \pi.$$
(A4)

- [1] G. 't Hooft, M.Veltman, Nucl. Phys. B 153, 365(1779).
- [2] W. Beenakker, H. Kuijf, W. L. van Neerven et al., Phys. Rev. D 40, 54(1989).
- [3] G. Aad *et al.*(ATLAS), PhysLett. B **716**, 1(2012).
- [4] S. Chatrchyan *et al.*(CMS), B **716**, 30(2012).
- [5] J. F. Gunion, H. E. Haber, G. Kane, S. Dawson, *The Higgs Hunter's Guide* (Perseus Publishing, Cambridge, Massachusetts, 1990).
- [6] B. A. Kniehl, Phys. Rept. **240**, 211(1994).
- [7] A. Djouadi, Phys. Rept. 457, 1(2008), Phys. Rept. 459, 1(2008).
- [8] M. Spira, Prog. Part. Nucl. Phys. **95**, 98(2017).
- [9] M. Carena, C. Grojean, M. Kado et al, "Status of Higgs Boson Physics", R. L. Workman et al.(Particle Data Group), Prog. Theor. Exp. Phys. 2022, 083C01(2022).
- [10] M. Roth, A. Denner, Nucl. Phys. B **479**, 495(1996).
- [11] J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, Nucl. Phys. B 106, 292(1976).
- [12] T. G. Rizzo, Phys. Rev. D 22, 178(1980), Phys. Rev. D 22, 1824(1980).
- [13] A. I. Vainshtein, M. B. Voloshin, V. I. Zakharov et al, Sov. J. Nucl. Phys. 30, 711(1979).
- [14] L. B. Okun, Leptons and Quarks(North-Holland, Amsterdam, 1982).
- [15] J. F. Gunion and H. E. Haber, Nucl. Phys. B 278, 449(1986), Erratum: Nucl. Phys. B 402, 569(1993).
- [16] R. K. Ellis, I. Hinchliffe, M. Soldate et al, Nucl. Phys. B 297, 221(1988).
- [17] D. Huang, Y. Tang and Y-L. Wu, Commun. Theor. Phys. 57, 427(2012).
- [18] M. Shifman, A. Vainshtein, M. B. Voloshin, V. Zakharov, Phys. Rev. D 85, 013015(2012).
- [19] W. J. Marciano, C. Zhang and S. Willenbrock, Phys. Rev. D 85, 013002(2012).
- [20] H. E. Haber and D. Wyler, Nucl. Phys. B **323**, 257(1989).
- [21] H. B. Dwight, Tables of Integrals and Other Mathematical Data(The Macmillan Company, New York, 1957), Third edition.

- [22] L. Lewin, Polylogarithms and Associated Functions(North Holland, New York, 1981), Second Edition.
- [23] I. S. Gradshteyn I. M. Ryzhik, *Table of Integrals, Series, and Products* (Academic Press, London, 2014), Eighth Edition.
- [24] M. Abramowitz and I. Stegun, Handbook of Mathematical Functions with formulas, Graphs and Mathematical Tables(Dover Publications, New York, 1972).
- [25] A. Erdelyi, Higher Transcendental Functions, Vol.I (McGrill-Hall Book Company, New York, 1953).
- [26] Z. X. Wang, D. R. Guo, *Special Functions*(World Scientific, Singapore, 1989).