

A comprehensive study on the relation between the metal enrichment of ionised and atomic gas in star-forming galaxies

M. ARABSALMANI,^{1,2} L. GARRATT-SMITHSON,^{3,4} N. WIJERS,^{5,6} J. SCHAYE,⁵ A. BURKERT,^{7,8} C.D.P. LAGOS,^{3,4,9} E. LE FLOC'H,¹⁰
D. OBRESCHKOW,³ C. PEROUX,^{11,12} AND B. SCHNEIDER¹⁰

¹*Excellence Cluster ORIGINS, Boltzmannstraße 2, 85748 Garching, Germany*

²*Ludwig-Maximilians-Universität, Schellingstraße 4, 80799 München, Germany*

³*International Centre for Radio Astronomy Research (ICRAR), University of Western Australia, 35 Stirling Hwy, Crawley, WA 6009, Australia*

⁴*ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D)*

⁵*Leiden Observatory, Leiden University, Niels Bohrweg 2, NL-2333 CA Leiden, The Netherlands*

⁶*CIERA and Department of Physics and Astronomy, Northwestern University, 1800 Sherman Ave, Evanston, IL 60201, USA*

⁷*University Observatory Munich (USM), Scheinerstraße 1, 81679 München, Germany*

⁸*Max-Planck-Institut für extraterrestrische Physik (MPE), Giessenbachstr. 1, 85748 Garching, Germany*

⁹*Cosmic Dawn Center (DAWN)*

¹⁰*CEA, IRFU, DAP, AIM, Université Paris-Saclay, Université Paris Cité, Sorbonne Paris Cité, CNRS, 91191 Gif-sur-Yvette, France*

¹¹*European Southern Observatory, Karl-Schwarzschildstrasse 2, D-85748 Garching bei München, Germany*

¹²*Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388, Marseille, France*

ABSTRACT

We study the relation between the metallicities of ionised and atomic gas in star-forming galaxies at $z = 0 - 3$ using the EAGLE cosmological, hydrodynamical simulations. This is done by constructing a dense grid of sightlines through the simulated galaxies and obtaining the star formation rate- and HI column density-weighted metallicities, Z_{SFR} and Z_{HI} , for each sightline as proxies for the metallicities of ionised and atomic gas, respectively. We find $Z_{\text{SFR}} \gtrsim Z_{\text{HI}}$ for almost all sightlines, with their difference generally increasing with decreasing metallicity. The stellar masses of galaxies do not have a significant effect on this trend, but the positions of the sightlines with respect to the galaxy centres play an important role: the difference between the two metallicities decreases when moving towards the galaxy centres, and saturates to a minimum value in the central regions of galaxies, irrespective of redshift and stellar mass. This implies that the mixing of the two gas phases is most efficient in the central regions of galaxies where sightlines generally have high column densities of HI. However, a high HI column density alone does not guarantee a small difference between the two metallicities. In galaxy outskirts, the inefficiency of the mixing of star-forming gas with HI seems to dominate over the dilution of heavy elements in HI through mixing with the pristine gas. We find good agreement between the available observational data and the $Z_{\text{SFR}}-Z_{\text{HI}}$ relation predicted by the EAGLE simulations. Though, observed regions with a nuclear starburst mode of star formation appear not to follow the same relation.

Keywords: Hydrodynamical simulations, Interstellar medium, Metallicity, Galaxy chemical evolution

1. INTRODUCTION

The content of heavy elements in galaxies over cosmic time provides essential information on how galaxies form and evolve. The metal enrichment of galaxies is not only linked to their star formation history, but is also related to outflowing and infalling gas in galaxies, through which metals are lost or diluted with pristine gas (e.g., [Matteucci 2012](#); [Silk & Mamon 2012](#); [Naab & Ostriker 2017](#); [Dayal & Fer-](#)

[rara 2018](#)). Metallicity is therefore one of the key properties of galaxies, and its relation with other galaxy properties provides crucial information on the mechanisms involved in galaxy formation and evolution (e.g., [Tremonti et al. 2004](#); [Mannucci et al. 2009, 2010](#); [Bothwell et al. 2013](#); [Møller et al. 2013](#); [Zahid et al. 2014](#); [Yabe et al. 2015](#); [Arabsalmani et al. 2015, 2018](#)).

The metallicity of a galaxy is defined as the ratio between the abundance of the elements heavier than helium (metals) and that of hydrogen+helium in the galaxy. This is observationally obtained from the abundances of elements in the stellar populations or in the interstellar medium (ISM) of a galaxy. The latter measurements are derived separately for

the ionised and the atomic gas phases. The ionised gas metallicities are generally measured from the ratios of strong emission lines in the spectra of galaxies. Such measurements are based on calibrations which are only well-constrained at low redshifts (and only within a metallicity range) and may not hold at higher redshifts (e.g., see [Maiolino et al. 2008](#); [Steidel et al. 2014](#)). Moreover, it is shown that different calibrations give rise to metallicities that differ significantly for a given galaxy, even at low redshifts ([Kewley & Ellison 2008](#)). The atomic gas metallicity measurements are instead based on the ratio between the column densities of heavy elements and atomic hydrogen derived from ISM absorption lines against background continuum sources. These measurements provide more reliable estimates of metal enrichment up to very high redshifts ($z > 5$, [Sparre et al. 2014](#); [Hartoog et al. 2015](#)), and do not rely on any calibration (negligible ionisation corrections and also dust corrections are assumed for these measurements). However, they provide the metallicity measurement only in a narrow beam along the bright background source – typically the quasars (QSOs), or the afterglows of Gamma Ray Bursts (GRBs), unlike the case of ionised gas metallicities measured from emission-lines where the metallicity measured is representative of an entire galactic region. Note that these ratios measure the metal enrichment of the atomic gas, and not that of the combined neutral molecular and atomic gas.

Despite the crucial role of the metal enrichment of galaxies in our understanding of galaxy formation and evolution, measurements of the true metallicities of galaxies remain uncertain. It is not clear whether the metallicities derived from the stellar populations are / should be the same as those obtained for the gas. Even the metallicities derived for different gas phases (ionised and atomic) in a galaxy might yield different values. Understanding how these measurements relate to each other will not only provide essential information on the mixing of metals in different gas phases, but is also required for interpreting the measurements derived with different methods.

To date, there are only a handful of measurements for the metallicities of both ionised and atomic phases of gas in the same galaxies. These are mainly obtained for intervening galaxies along quasar sightlines, showing deviations between the two measurements (see e.g., [Christensen et al. 2014](#); [Friis et al. 2015](#); [Péroux et al. 2016](#); [Rahmani et al. 2016](#); [Hamanowicz et al. 2020](#)). In several cases the measurement are obtained for nearby galaxies (e.g., [Heckman et al. 2001](#); [Thuan et al. 2002, 2005](#); [Cannon et al. 2005](#); [Lebouteiller et al. 2009, 2013](#); [Hernandez et al. 2021](#)). Given the rarity of these measurements, cosmological hydrodynamical simulations (e.g., [Schaye et al. 2015](#); [Pillepich et al. 2018](#); [Maio et al. 2022](#)) offer a natural test-bed to explore the relationship between the metallicities of ionised and atomic gas

which can also be confronted to the few observational constraints.

In this paper we present a detailed and comprehensive analysis of the relation between the atomic and ionised gas metallicities in star-forming galaxies in the redshift range $0 \leq z \leq 3$ and investigate the dependence of this relation on various parameters such as the global properties of galaxies as well as the local properties of regions of interest. For this study we use the EAGLE (Evolution and Assembly of GaLaxies and their Environments) set of cosmological hydrodynamical simulations ([Schaye et al. 2015](#)). We describe the numerical methods in Section 2. The results and discussion are presented in Section 3. We summarise our findings in Section 4. We use a solar metallicity value of 0.0127 throughout this work. This is the default value used in CLOUDY v7.02 (last documented in [Ferland et al. \(1998\)](#)), with element abundances from [Grevesse & Sauval \(1998\)](#), [Allende Prieto et al. \(2001\)](#), [Holweber \(2001\)](#), and [Allende Prieto et al. \(2002\)](#).

2. NUMERICAL METHODS

2.1. Simulations

This paper uses the publicly available cosmological hydrodynamical simulation EAGLE RecalL025N0752 ([Schaye et al. 2015](#); [Crain et al. 2015a](#); [McAlpine et al. 2016](#); [The EAGLE team 2017a](#)). The correlations between metallicity and other galaxy properties – such as stellar mass, star formation rate (SFR), specific star formation rate, gas fraction – have been extensively studied using EAGLE (e.g., [Lagos et al. 2016](#); [De Rossi et al. 2017](#); [Trayford & Schaye 2019](#); [van Loon et al. 2021](#)), with authors finding a good agreement with observed trends down to stellar masses of $10^9 M_\odot$ using the EAGLE run RecalL0025N0752. This is the highest resolution EAGLE simulation and is consequently the simulation we use for this analysis. Particularly relevant to this work, EAGLE has been shown to reproduce both the observed radial metallicity gradients within the central part of galaxies (< 0.5 times the half-mass radius; [Collacchioni et al. 2020a](#)), along with $z \sim 0$ metallicity gradients within star-forming regions of disc galaxies (within observed scatter; [Tissera et al. 2019](#)), and the redshift evolution of the metallicity gradient within star-forming regions (although deviations in scatter were seen between the simulations and observations; [Tissera et al. 2022](#)). However, it is worth noting the recent paper by [Yates et al. \(2021\)](#), where it was highlighted that the cosmic metal density of the neutral gas phase at high redshift ($3 \lesssim z \lesssim 5$) is significantly lower in existing cosmological simulations (including EAGLE), than observations of Damped Lyman- α systems (systems with $N(\text{H I}) \geq 2 \times 10^{20} \text{ cm}^{-2}$) imply. This tension was seen across different cosmological simulations, hence it is not clear whether one simulation is favoured over another for this type of metallicity

study (note that the study presented in this paper is focused on $z \leq 3$). Our choice to use EAGLE for this analysis was further motivated by the fact the RecalL025N0752 run has been shown to reproduce the $z = 0$ global H_2 scaling relations (e.g., Lagos et al. 2015), alongside the HI scaling relations (e.g., Crain et al. 2017) and the HI column density distribution function (CDDF; e.g., Rahmati et al. 2015).

The simulations follow the evolution of a cubic volume of side-length 25 cMpc of the universe to $z = 0$, where c notes ‘comoving’. They use a dark matter particle mass of $1.21 \times 10^6 M_\odot$ and an (initial) gas particle mass of $2.26 \times 10^5 M_\odot$. These simulations use an updated version of the hybrid N-body Tree-PM/smooth particle hydrodynamics (SPH) code GADGET 3, presented in Springel (2005). The modifications to the numerical methods in GADGET – i.e. the inclusion of an artificial thermal conduction switch (Price 2008), a viscosity switch (Cullen & Dehnen 2010), a time step limiter (Durier & Dalla Vecchia 2012) and the pressure-entropy formalism described in Hopkins (2013) – are collectively known as “Anarchy” and are described in the appendix of Schaye et al. (2015) and in Schaller et al. (2015). The EAGLE simulations also model processes such as star formation and stellar/AGN feedback using sub-grid prescriptions, where the feedback efficiencies are calibrated (Crain et al. 2015b) to match the $z = 0$ galaxy stellar mass function (GSMF), the sizes of disk galaxies, and the observed black hole mass – stellar mass relation.

The star formation and metal enrichment/stellar-mass-loss sub-grid prescriptions are of particular importance to this work. Within the simulation, stellar populations are represented by star particles. These are formed using a metallicity-dependent density threshold designed to track the transition from the warm, atomic to the cold, molecular interstellar gas-phase (first described in Schaye 2004), alongside a pressure-dependent star formation rate that reproduces the observed Kennicutt-Schmidt star formation law (as described in Schaye & Dalla Vecchia 2008). Star particle formation is stochastic, with a probability linked to the star formation rate of a gas particle. Further, as described in Wiersma et al. (2009a), the stellar-mass-loss for each star particle is approximated using the metallicity-dependent lifetimes presented in Portinari et al. (1998) and the AGN/stellar wind/core-collapse supernovae yields given in Marigo (2001) and Portinari et al. (1998), on top of the mass-loss associated with Type Ia supernovae. EAGLE tracks 9 elements individually; H, He, C, N, O, Ne, Mg, Si and Fe (Wiersma et al. 2009b). These elements are distributed across the neighbours of each star particle using the SPH kernel weighting. For this paper we use the metallicities associated with each particle, as opposed to the SPH-smoothed version of the metallicities (those used in the cooling calculations of the simulation, though not when calculating the star formation rate of each particle (see

The EAGLE team 2017b). This is because in eq. 2 we are interpolating to a grid, and hence by smoothing the metallicities we would smooth twice which in return would artificially decrease the resolution. Further, when we do use the SPH-smoothed version of the particle metallicities we see an artificial reduction in scatter at high impact parameters/low densities, driven by the fact the metallicity is smoothed over larger distances in the outskirts of the galaxy/CGM. These differences highlight the constraints of the simulation – since the resolution of SPH simulations naturally follows mass, the CGM/IGM remain poorly resolved, hence the metallicity values of any HI in these regions should be interpreted with caution. For completeness, we reproduce all the plots presented in Section 3 using the SPH-smoothed version of the particle metallicities and present them in the Appendix.

2.2. The atomic gas phase

Similar to other cosmological-scale simulations (e.g., Dubois et al. 2014; Pillepich et al. 2018), EAGLE does not have the resolution to follow the cold gas phase of the galaxy; instead it imposes a density-dependent pressure floor, normalised to 8000 K at $n_{\text{H}} = 10^{-1} \text{ cm}^{-3}$. Therefore, the simulation needs to be post-processed in order to estimate the neutral (atomic and molecular) gas fraction of each gas particle. We do this using the method described in Figure 1 of Garratt-Smithson et al. (2021), based on previous works (e.g., Lagos et al. 2015; Bahé et al. 2016; Crain et al. 2017; Diemer et al. 2018). Note that we obtain both the atomic gas fraction and molecular gas fraction separately and use the former in order to estimate the metallicity of the atomic gas. We summarise the method in the following five steps.

(i) First, we calculate the total non-ionized hydrogen fraction, which combines molecular hydrogen and HI. We do this according to the prescription of Rahmati et al. (2013), assuming a Haardt & Madau (2012) photo-ionizing background. In this prescription, the photo-ionization rate is estimated using a density-dependent fitting function. The fitting function was calibrated using cosmological simulations that included radiative transfer calculations. This photo-ionization rate is then used to estimate the fraction of non-ionized hydrogen in each gas particle. In the next steps, we calculate the molecular hydrogen fraction in the non-ionized gas, which allows us to calculate the HI fraction.

(ii) From here, the molecular-dissociating UV-flux is estimated for each gas particle. The method we utilise here depends on whether the gas particle is star-forming or not. If it is, the flux is estimated using the ratio between the SFR density of the particle and that of the solar neighbourhood (as in Lagos et al. 2015). If it is non-star-forming, the particle uses a minimum value set by the Haardt & Madau (2012) UV background. Additionally, following the work by Diemer et al. (2018), later adapted for EAGLE in Garratt-Smithson




Figure 1. Example of the construction of artificial sightlines about two galaxies of stellar mass $2 \times 10^9 M_\odot$ (top row) and $9 \times 10^9 M_\odot$ (bottom row). The four columns from left to right show the maps of gas metallicity, the HI column density, the star formation rate surface density, and $\log_{10}(Z_{\text{SFR,med}}/Z_{\text{HI}})$ defined as δ (see Section 3), respectively. Each grid has a set pixel size of 2 kpc. The side length is determined by $2R_{200}$, though in the interest of comparison, only the inner 100 kpc for each galaxy is shown here.

et al. (2021), we assume 10 per cent of the UV photons emitted by star-forming particles escape and propagate through the surrounding medium (assumed to be optically thin). This provides an additional source of UV for non-star-forming particles.

(iii) We then calculate two theoretically-motivated density floors for the cold neutral medium; the first assumes a 2-phase ISM model (following [Wolfire et al. 2003](#)). The second is the minimum density of the cold neutral medium required to maintain pressure equilibrium (as described in [Krumholz 2013](#)).

(iv) Next, we take the maximum of these density floors and use this value, alongside the molecular-dissociating radiation field, to estimate the optical depth of dust grains (following [Krumholz et al. 2009](#); [Krumholz 2013](#)) – fundamental to the formation of molecular hydrogen.

(v) Lastly, we calculate the fraction of molecular hydrogen using both the UV-flux and the optical depth of dust, utilising the method described in [McKee & Krumholz \(2010\)](#). The atomic gas, HI, defined in this study excludes the molecular hydrogen as well as helium.

To illustrate our post-processing method, Figure 2 plots the gas phase diagram for all gas particles associated with 10 $z = 0$ systems taken from our sample. Here we use all

gas particles found within the virial radius of each system. This will therefore include the circumgalactic medium alongside the interstellar medium. The top plot is coloured by the HI mass fraction (f_{HI}), the middle plot shows the molecular hydrogen mass fraction (f_{H_2}), while the lower plot uses the ionised gas fraction (f_{ion}). Here we can see high f_{HI} and f_{H_2} values are associated with the low temperature (in particular, note the large values of f_{HI} and f_{H_2} at $T \sim 10^4$ K), high density regime, while the opposite is true for gas particles with high f_{ion} . In particular, galaxies sitting on the artificial pressure floor (seen in the bottom right of the phase diagram), are typically H_2 dominated.

2.3. Constructing artificial sightlines

We constructed artificial sightlines for the simulation box using an $N = 8000^2$ grid, with each grid cell representing a single sightline (each cell has a side length 3.125 ckpc). The box was then sliced into 5 in order to approximate typical velocity cuts used in observations (as in [Rahmati et al. 2015](#)). The metallicity of each pixel in our grid was calculated using a cloud-in-cell (CIC) algorithm. The CIC algorithm is a multi-linear interpolation scheme to evaluate fields on a regular grid based on irregularly placed particles. Here each particle contributes a metallicity weighted by the HI mass or




Figure 2. Gas phase diagram for the gas particles associated with 10 $z=0$ systems in our sample, with points colored by the HI mass fraction (f_{HI} ; upper plot), the molecular hydrogen mass fraction (f_{H_2} ; middle plot) and the ionised gas mass fraction (f_{ion} ; lower plot) respectively. Only gas particles at radii below the virial radius of each system were included. Note the large molecular hydrogen mass fractions as well as the large HI mass fractions at $T \sim 10^4$ K.

SFR that gas particle contributed to the cell, i.e.

$$Z_{\text{SFR}} = \frac{\sum_{i=1}^N \text{SFR}_{\text{cic},i} \times Z_i}{\sum_{i=1}^N \text{SFR}_{\text{cic},i}}, \quad (1)$$

where $\text{SFR}_{\text{cic},i}$ is the cloud-in-cell contribution of the i th particle to the SFR of the cell and Z_i is the metallicity of the i th particle. Similarly,

$$Z_{\text{HI}} = \frac{\sum_{i=1}^N \text{MHI}_{\text{cic},i} \times Z_i}{\sum_{i=1}^N \text{MHI}_{\text{cic},i}}, \quad (2)$$

where $\text{MHI}_{\text{cic},i}$ is the cloud-in-cell contribution of the i th particle to the total HI mass of the cell. The metallicities weighted by HI mass and star formation rate provide proxies

for the metallicities of atomic gas and ionised gas, respectively, thus we can compare the typical metallicity as measured using global cold gas in galaxies, with the metallicity of the star-forming regions. We mention again that throughout this paper, we use the metallicity of atomic gas, as the neutral gas metallicities obtained from observations in the literature refer to the metal enrichment of only atomic gas. Given that the EAGLE simulations reproduce the Kennicutt-Schmidt star formation law (as described in Schaye & Dalla Vecchia 2008), we expect the H_2 -weighted metallicities to be very similar to SFR-weighted metallicities. We investigate this and we indeed find this to be the case.

In order to investigate the relation between the HI-weighted metallicity and the SFR-weighted metallicity on a galaxy-by-galaxy level, we also took all central galaxies with a stellar mass of $> 10^8 M_\odot$ out of RecalL025N0752 and created a grid centered on each individual galaxy, using a side-length of $2R_{200}$ – the physical radius within which density is 200 times the critical density of the Universe, calculated for each halo identified via the friends-of-friends algorithm. Here we negate any projection-effects by explicitly rotating each system so that we are observing it perpendicularly to the angular momentum axis of the star particles. Further, the size of each pixel is set to 2 pkpc, independent of the value of R_{200} . This means there will be a smaller number of pixels for lower mass galaxies. As above, we then use this grid to interpolate Z_{HI} , Z_{SFR} , along with the HI column density $N(\text{HI})$ for each pixel. In observations, the impact parameter of a sightline, b , is defined as the projected distance between the sightline and the centre of its corresponding galaxy. We too calculate the impact parameter, b , using the centre of each pixel in the 2D $x-y$ plane (where the z direction is the rotation axis). Figure 1 shows an example of this process for two galaxies of stellar mass $2 \times 10^9 M_\odot$ (top row) and $9 \times 10^9 M_\odot$ (bottom row).

In the following section we present and discuss our findings using the sightlines constructed for individual galaxies. This is to ensure that there is no more than one galaxy per sightline, which is likely not the case for sightlines in the entire box (the sightlines constructed in the full box might correspond to more than one galaxy). We note that the results show the same trends no matter if we use all sightlines in the simulation box, or the sightlines produced using galaxies cut-out from the simulation.

3. RESULTS

3.1. $Z_{\text{HI}}-Z_{\text{SFR}}$ plane of EAGLE galaxies

The SFR- and HI- weighted metallicities (Z_{SFR} and Z_{HI} respectively) of all the sightlines, excluding those with zero SFR or zero HI, in EAGLE galaxies at $z=2$ are presented in Figure 3. The first notable finding is that for almost all the sightlines, the Z_{SFR} values are larger than the Z_{HI} val-




Figure 3. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plane for all the sightlines in EAGLE galaxies at $z = 2$. The medians of the Z_{SFR} values in each Z_{HI} bin are shown with pink diamonds. The shaded area shows the 1σ spread around the medians of the Z_{SFR} values. The $x=y$ line in the plane is presented by a dashed line.

ues. This implies that the star-forming gas in all regions of the galaxies is more enriched with metals than the HI gas located along the sightlines passing the same regions. It is also apparent that the majority of the sightlines have Z_{SFR} above ~ 0.1 solar metallicity. For the same sightlines the Z_{HI} values can be an order of magnitude lower. The saturation of the Z_{SFR} values could be a result of the local star formation enriching the surrounding gas to $Z_{\text{SFR}} \gtrsim 0.1 Z_{\odot}$.

To undertake a more quantitative investigation, we use the medians of the Z_{SFR} values, $Z_{\text{SFR,med}}$, within bins of Z_{HI} . The medians are marked with the pink diamonds in Figure 3, and the shaded area in the plot shows the 1σ region around the median values. In each Z_{HI} bin, we measure the difference between $\log_{10}(Z_{\text{SFR,med}}/Z_{\odot})$ and the central $\log_{10}(Z_{\text{HI}}/Z_{\odot})$ value of the bin, and refer to it as δ . δ clearly increases with a decrease in Z_{HI} and consequently a decrease in $Z_{\text{SFR,med}}$: for a $Z_{\text{SFR,med}}$ of Z_{\odot} , δ is about 0.2 dex, while its value increases to 1.2 dex for a $Z_{\text{SFR,med}}$ of $0.3 Z_{\odot}$. Equivalently, δ increases from 0.2 dex to 1.2 dex when Z_{HI} decreases from $0.5 Z_{\odot}$ to $0.02 Z_{\odot}$. The scatter in the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation also increases with decreasing metallicity. The 1σ (68%) spread around $Z_{\text{SFR,med}}$ varies from 0.4 dex to 0.7 dex between $Z_{\text{SFR,med}}$ values of Z_{\odot} and $0.3 Z_{\odot}$ (or equivalently between Z_{HI} values of $0.5 Z_{\odot}$ and $0.02 Z_{\odot}$).

The large differences between the metallicities of the two gas phases in the low metallicity regime indicates that the

low metallicities of the atomic gas could have a large contribution from dilution of heavy elements through mixing with the metal-poor, infalling pristine gas that is not yet enriched by heavy elements. We note that the metals in the EAGLE simulation remain fixed to their Lagrangian resolution element after being injected by a stellar population. Therefore, the mixing we refer to here is not an exchange of metals between resolution elements. Here, ‘mixing’ refers to the gas (resolution elements) moving in(to) and out of the galaxy, intermingling and transporting metals and pristine gas, or in other words, it refers to the effective distribution of metals in the galaxy through multiple feedback events. The dilution by inflowing pristine gas is particularly expected in the outskirts of galaxies, where the infall of gas from intergalactic medium reduces the average metal enrichment of the diffuse, HI without directly affecting the metallicity of the ionised gas (Collacchioni et al. 2020b; Wright et al. 2021). On the other hand, the decreased difference between the two metallicities in the high-metallicity regime could also imply that the mixing of neutral and ionised gas in metal-rich regions (such as the central regions of galaxies) is more efficient, leading to a more uniform metal enrichment in different gas phases. The enrichment of metals due to gas mixing in different phases, and the dilution of heavy elements in neutral gas by mixing with the pristine gas, are both likely to contribute in the difference between the metallicities of atomic and ionised gas. Although, the contributions of these two effects seem to depend on, directly or indirectly, the metallicity regime.

It is important to note that the sightlines with low metallicities in Figure 3 not only sample the (metal-poor) outskirts of galaxies (independent of the stellar mass regime), but also sample the entire extent of the low-mass galaxies, irrespective of the positions of the sightlines in galaxies with respect to the galaxy centres (given the mass-metallicity relation of star-forming galaxies; Lequeux et al. 1979; Tremonti et al. 2004; Zahid et al. 2014; Kudritzki et al. 2015). The large δ values in the low metallicity regime therefore raises the following questions: “is the mixing of gas in low-mass galaxies less efficient than in massive galaxies?”, “does the position of a sightline with respect to the centre of the galaxy have a significant effect on the mixing of gas?”, and “does the stellar mass of a galaxy affect the contribution of the pristine gas in diluting the metal enrichment of HI in the system and result in lower values of δ ?”.

To answer these questions, one needs to separately investigate the effects of the stellar mass and the impact parameter on the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation. Considering that lower-mass galaxies are smaller in size (see e.g., Shen et al. 2003; Graham & Worley 2008; Maltby et al. 2010; van der Wel et al. 2014, for the stellar-mass-size relation of galaxies), they are expected to host sightlines with generally smaller impact parameters. We therefore use the impact parameter of the sight-




Figure 4. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plane for the sightlines at $z = 2$ in three bins of stellar mass (rows) and four bins of impact parameter (columns). The colour of the points indicates the $N(\text{HI})$ value of each sightline, with the colour bar presented in the right side of each row. The grey lines show the $Z_{\text{SFR},\text{med}}$ values in each plot. The $x=y$ line in the plane is presented by a dashed line. Note that R_{200} varies between $\sim 40 - 90$ kpc for galaxies with stellar masses between 10^8 and $10^{10} M_{\odot}$.

lines, normalised by the R_{200} values of galaxies (varying between $\sim 40 - 90$ kpc for galaxies with stellar masses between 10^8 and $10^{10} M_{\odot}$) in our analysis. The large values of b/R_{200} therefore corresponds to the outskirts of galaxies, independent of the stellar mass or equivalently the size of a galaxy.

Figure 4 compares the effects of the stellar mass and the impact parameter on the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation at $z = 2$. Each row in the figure represents a stellar mass bin, and each column represents a specific range of the normalised impact parameter. The colouring of the points are according to the HI column densities along the sightlines. It is clear that at a given metallicity the sightlines with smaller normalised impact parameters are more concentrated around the $x=y$ line. In other words, the difference between the Z_{HI} and Z_{SFR} values reduces in the central regions of galaxies, irrespective of their stellar mass.

The stellar mass of galaxies does not seem to have a direct impact on the relation between Z_{HI} and Z_{SFR} . Of course galaxies with higher stellar mass would have larger sizes and naturally would accommodate more sightlines at larger impact parameters. Therefore the increase in δ and also the scatter of the relation at larger impact parameters (see the column for $0.1 \leq b/R_{200} < 0.3$) in galaxies with higher stellar mass is expected. At small impact parameters (see the column for $b/R_{200} < 0.05$), both δ and the scatter in the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation appears to decrease as the stellar mass of the host galaxies increases. But noting the smaller spread of HI column densities in massive galaxies in this regime of impact parameters, the smaller deviation between the two metallicities is likely not directly due to larger stellar mass, but is plausibly linked to the column density of HI. In fact, the smaller deviations between the two metallicities and the smaller scatter of the relation in low impact parameters in all stellar mass

bins could be a result of higher column densities – HI column densities are generally higher in the central regions of galaxies.

To investigate the effect of the $N(\text{HI})$ on δ and the scatter of the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation, we consider only the sightlines with high HI column densities – $N(\text{HI}) \geq 10^{20} \text{ cm}^{-2}$, the column density typically found within the optical disks of galaxies. We also restrict the sightlines to be within a distance of $b/R_{200} \leq 0.3$ from the galaxy centres. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plots for these sightlines at $z = 2$ are presented in the top panels of Figure 5. δ and the scatter in the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation remain quite large even at such high column densities. In fact, there is a significant fraction of sightlines with high HI column densities for which the difference between the two metallicities exceeds 0.5 dex.

On the other hand, when we consider the sightlines within the $0.02 R_{200}$ central regions of galaxies (irrespective of their HI column densities), δ and the overall scatter on the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation appear to be much smaller. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plot for these sightlines are shown in the bottom-right panel of Figure 5. It is clear from this Figure that these sightlines have large $N(\text{HI})$ values, as expected to be the case in the central regions of galaxies.

Note that the region within $0.02 R_{200}$ corresponds to the ~ 1 kpc, ~ 1.5 kpc, and ~ 2 kpc central regions of galaxies with stellar masses of $10^8 M_{\odot}$, $10^9 M_{\odot}$, and $10^{10} M_{\odot}$, respectively. The gravitational softening length of 0.35 kpc in the simulations thus allows us to explore the separation of metallicity between gas phases at spatial scales of $0.02 R_{200}$.

In order to quantitatively compare the role of $N(\text{HI})$ and the impact parameter on the difference between the two metallicities, we measure δ and the 1σ scatter of the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation for the sightlines with $N(\text{HI}) \gtrsim 10^{21} \text{ cm}^{-2}$ (shown in the top-right panel of Figure 5) and also the sightlines located within the $0.02 R_{200}$ central regions of galaxies (presented in the bottom-right panel of Figure 5). For the $Z_{\text{SFR,med}}$ values of $\gtrsim 0.3 Z_{\odot}$, δ remains around 0.2 dex for both the groups, and the 1σ scatter around δ remains around 0.1–0.2 dex for both the groups as well. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation in this metallicity regime therefore is very similar for sightlines with $N(\text{HI}) \gtrsim 10^{21} \text{ cm}^{-2}$ and sightlines in the $0.02 R_{200}$ central regions of galaxies. However, both δ and the 1σ scatter increase rapidly for the former group at $Z_{\text{SFR,med}} \lesssim 0.3 Z_{\odot}$.

We can therefore safely state that the location of a sightline being within the $0.02 R_{200}$ central regions of galaxies guarantees a small, systematic difference between Z_{HI} and Z_{SFR} with $\delta = 0.2$ dex. This finding implies that the mixing between star-forming gas and the HI gas is efficient in the central regions of galaxies (the mixing time scale is shorter than the typical gas depletion time). Note that this is not a direct consequence of a subgrid model for mixing since the simulations do not model subgrid diffusion. The efficiency of

mixing seems to reduce by moving outwards from the galaxy centres, as indicated by the large values of δ in large impact parameters. We find this to be independent of the stellar mass of the galaxies and hence the metallicity regime, as discussed before (see Figure 5).

On the other hand, a high HI column density of a sightline alone does not imply a small difference between the two metallicity measurements, and depending on the metallicity regime, δ could be as high as ~ 1.5 dex even for a sightline with $N(\text{HI}) \gtrsim 10^{21} \text{ cm}^{-2}$. Sightlines located at large distances from galaxy centres, even those with high HI column densities, are less enriched with heavy elements compared to the ionised gas in projected regions which are more metal enriched due to local star formation. We note that some of these sightlines could be associated with low-mass satellite galaxies, and some could be linked to the intergalactic medium, appearing at certain impact parameters due to projection effects. But given the significant number of high $N(\text{HI})$ sightlines, with large δ values, a large number of them should be associated with the outskirts of galaxies. Given the high $N(\text{HI})$ of these sightlines, they are less likely to be affected by mixing with pristine gas. Therefore, their large δ values should be mainly due to the inefficiency of the mixing of star-forming gas with HI. This implies that the inefficiency of the mixing of the two gas phases should be the dominant effect compared to the metal dilution in HI through mixing with pristine gas in the outskirts. Mitchell et al. (2020) and Wright et al. (2021) used the EAGLE simulations and showed that accreting gas in massive galaxies have more contributions from the recycled (hence more metal-rich) gas compared to low mass galaxies. Given that the stellar mass has an insignificant role on the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation, the findings of Mitchell et al. (2020) and Wright et al. (2021) support the dominant effect of the mixing of the two gas phases in the outskirts of galaxies.

We also investigate the redshift evolution of the $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation in the redshift range $z = 0 - 3$. The left panel of Figure 6 shows the $Z_{\text{SFR,med}}$ values and the 1σ scatter of the relation for all the sightlines at redshifts $z = 0, 1, 2$ and 3. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation appears to have a slight evolution with redshift: the $Z_{\text{SFR,med}}$ of sightlines at a fixed Z_{HI} slightly decreases with increasing redshift. This is consistent with the overall decrease in the average ionised gas metallicity in galaxies measured from observations. However, the relation between the two metallicities does not vary much with redshift for sightlines in the central regions of galaxies, as demonstrated in the right panel of Figure 6. The small, systematic difference between Z_{HI} and Z_{SFR} in the central regions of galaxies remains around 0.2 dex between $z = 0$ to $z = 3$. This suggests the efficiency of gas mixing in the central regions of galaxies to be very similar at $z = 0 - 3$.




Figure 5. The $Z_{\text{HI}}-Z_{\text{SFR}}$ plane for sightlines at $z = 2$ selected based on $N(\text{HI})$ and impact parameter. The $Z_{\text{SFR,med}}$ values are shown with black diamonds and the $x=y$ line is presented with the dashed lines. **Top panels:** Sightlines with HI column densities of $10^{20} \text{ cm}^{-2} \leq N(\text{HI}) < 10^{21} \text{ cm}^{-2}$ and $N(\text{HI}) \geq 10^{21} \text{ cm}^{-2}$ in the left and right panels respectively. All the presented sightlines are within the $0.3 R_{200}$ from galaxy centres. The colour bar represents the value for b/R_{200} . **Bottom panels:** Sightlines with $0.02 < b/R_{200} \leq 0.05$ in left and those within the $0.02 R_{200}$ central regions of galaxies in the right panel respectively. The colour bar represents the HI column density of the sightlines.

3.2. Comparison with observations

To compare our findings from simulations with the available observations, we use the metallicity measurements of both ionised and atomic phases of gas in the same galaxies reported in the literature. These measurements and the relevant references are provided in Table 1. The gathered observational data are for 22 intervening galaxies along quasar lines of sight, a GRB host galaxy, and 7 nearby galaxies.

For the nearby galaxies, the metallicity of HI gas is obtained from the ratio between the column densities of heavy elements (such as argon, nitrogen, and oxygen) and HI, derived from the imprinted absorption lines on the continuum emission from the bright UV targets (such as UV-bright massive stars) within these galaxies. The reported atomic metallicities of these galaxies in Table 1 are all based on abundances of oxygen. For all these galaxies but M83 the values of Z_{atomic} and Z_{ionised} are the average values over the entire galaxies. For galaxy M83, the reported metallicities of both




Figure 6. **Left:** The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ relation for all the sightlines of EAGLE galaxies at redshifts $z = 0, 1, 2$ and 3 . The $Z_{\text{SFR,med}}$ values are marked with diamonds and the 1σ scatter around the medians are shown with shaded areas. **Right:** The $Z_{\text{SFR,med}}$ values and the 1σ scatter around them for sightlines within the $0.02 R_{200}$ central regions of galaxies at four redshifts.

gas phases are derived in individual regions in the galaxy. The impact parameters of all these regions are provided in Table. 1.

The metallicities of HI in the QSO absorbers and the GRB host are derived based on the ratio between the column densities of heavy elements (such as Zn and S) and HI, obtained from ISM absorption lines imprinted on quasar/GRB spectrum. These metallicities are therefore measured for a narrow beam along the sightlines of the bright background quasars or GRBs. The ionised gas metallicities on the other hand are derived from the flux measurement of the bright nebular lines, originating from the star-forming regions in galaxies. These metallicities therefore provide an average value over the entire galaxies.

For a fair comparison between the measurements from observations and our findings from simulations, we require a representative Z_{SFR} value for each of the EAGLE galaxies which is measured over the entire star-forming disk of the galaxy, similar to the metallicity that is measured for ionised gas in galaxies based on observations. We obtain this value in two different ways: (i) we measure the mean SFR-weighted metallicity of all star-forming particles within the stellar half-mass radius of the galaxy, and (ii) we obtain the mean of the Z_{SFR} values for all the sightlines in a galaxy (within a side-length of $2R_{200}$, as discussed in Section 2.3). We find the two mean metallicities to be in very good agreement, and therefore use the latter definition for $Z_{\text{SFR,mean}}$ as the global Z_{SFR} in each of the EAGLE galaxies. We also use the mean values of the Z_{HI} , $Z_{\text{HI,mean}}$, by measuring the mean HI-weighted metallicity of all particles within a radial aperture of a galaxy.

All the systems in Table 1 have a range of HI column density between $\sim 10^{19} \text{ cm}^{-2}$ and 10^{22} cm^{-2} . The QSO-sightlines and all the regions in M83 have an impact parameter $\lesssim 60$ kpc. For the host galaxy of GRB 121024, the impact parameter of the GRB sightline in its host galaxy is not measured, but it is likely to be within the central few kpc of its host galaxy (see Friis et al. 2015). The median impact parameter measured for GRBs is reported to be $\sim 1 - 2$ kpc (see Arabsalmani et al. 2015; Lyman et al. 2017). For the nearby dwarfs, given their sizes, the continuum emission originates from regions within the $\lesssim 60$ kpc range. We therefore restrict the $Z_{\text{HI,mean}}$ measurements to HI column densities $10^{19} \text{ cm}^{-2} - 10^{22} \text{ cm}^{-2}$ and impact parameters of 0-60 kpc for a fair comparison with the observational measurements in hand. Figure 7 presents the metallicity measurements of all the systems from observations (a total number of 37 data points), along with the obtained metallicities of EAGLE galaxies in the $Z_{\text{HI,mean}}\text{--}Z_{\text{SFR,mean}}$ plane, at $z = 0 - 3$, similar to the redshifts of the observational points.

We find a good agreement between the metallicity measurements from observations and the prediction from EAGLE simulations. The four observational data points that stand out in the relation correspond to the four regions in the 200 pc nuclear region of M83. Hernandez et al. (2021) discusses that the efficient conversion of atomic to molecular gas and hence the depletion of the HI gas in the nuclear region of M83 is responsible for the relatively high atomic gas metallicity in that region. Our findings (see Fig. 4 and Fig. 5) clearly demonstrate that simply having low HI column densities in star forming regions (with normal mode of




Figure 7. The ionised and atomic gas metallicity measurements from observations, in comparison with the obtained metallicities of EAGLE galaxies. The quasar absorbers are shown with pink squares, and the host galaxy of GRB 121024 is presented with a purple square. Nearby dwarf galaxies are marked by black diamonds. The individual regions of M83 are shown by gray stars. The x values for these systems are measured in single sightline from the ratio between the column densities of heavy elements and HI, obtained from ISM absorption lines. The y values for these systems are obtained from the flux measurement of the bright nebular lines coming from the entire star-forming disk of the galaxy. The size of the squares are proportional to the redshift of the system. The blue points show the EAGLE galaxies at $z = 0 - 3$ in the $Z_{\text{HI,mean}} - Z_{\text{SFR,mean}}$ plane.

star formation) do not necessarily lead to $Z_{\text{SFR}} < Z_{\text{HI}}$. However, the presence of an ongoing starburst in the nuclear region of M83 (Dopita et al. 2010; Wofford et al. 2011) might be the cause for the significant difference between the metallicities with $Z_{\text{SFR}} < Z_{\text{HI}}$. This indicates that regions with a nuclear starburst mode of star formation might be following a different $Z_{\text{HI}} - Z_{\text{SFR}}$ relation. Note that the effects of enhanced radiation fields in nuclear starbursts is not included when simulating the EAGLE galaxies.

We also note that a few of the local dwarf galaxies presented in Fig. 7 have stellar masses $\sim 10^6 M_{\odot}$, below the lower stellar mass limit of the EAGLE galaxies ($M_{*} = 10^8 M_{\odot}$). These dwarf galaxies trace the lower metallicity end of the $Z_{\text{HI}} - Z_{\text{SFR}}$ relation predicted by the EAGLE galaxies.

4. DISCUSSION AND SUMMARY

The metal enrichment of different gas phases in galaxies can provide valuable information on fundamental phenomena in galaxy formation and evolution. We present a detail study of the relation between the metallicities of ionised and atomic gas in star-forming galaxies at $z = 0 - 3$ using the EAGLE simulations. For each of the simulated galaxies we construct several sight lines and for each sightline, we obtain the SFR- and N(HI)- weighted metallicities of gas, Z_{SFR} and Z_{HI} respectively, and use them as proxies for the metallicities of ionised and atomic gas in the sightline.

We find that $Z_{\text{SFR}} > Z_{\text{HI}}$ in almost all sightlines, with $\delta := \log_{10}(Z_{\text{SFR,med}}/Z_{\odot})$ exceeding a few dex in some sightlines, where $Z_{\text{SFR,med}}$ is the median of Z_{SFR} values in a Z_{HI} bin. In addition, δ increases with decreasing metallicity. We investigate whether stellar mass and hence the overall galaxy metallicity play a role in gas mixing and making the metal enrichment in different gas phases more uniform. Moreover we explore the effect of the impact parameter on δ , considering that galaxy outskirts are less enriched with heavy elements and hence the position of the sightlines with respect to the galaxy centre could have an effect on δ (e.g. through the dilution of heavy elements by pristine gas in outskirts). We find that irrespective of galaxy stellar mass, δ decreases by decreasing the impact parameter and reaches a minimum value of ~ 0.2 dex in the central regions of galaxies. This implies that mixing of star-forming gas and HI is most efficient in the central region of galaxies. The large values of δ in large impact parameters indicate that the efficiency of mixing decreases by moving outwards from the galaxy centre. This finding is independent of redshift, i.e. δ saturates to ~ 0.2 in the central $0.02 R_{200}$ of galaxies at $z = 0 - 3$.

Even though the low impact parameter sightlines are generally associated with high HI column densities, having a high N(HI) alone does not mean a small δ . Atomic gas located at large distances from galaxy centres (large impact parameters), even those with high HI column densities, are less enriched with heavy elements compared to the ionised gas in projected regions which are more metal enriched due to local star formation. In fact, we find a significant number of sightlines with $N(\text{HI}) \geq 10^{20} \text{ cm}^{-2}$ to be associated with large impact parameters, and large values of δ . Some of these sightlines could be associated with low-mass satellite galaxies, and some could be linked to the intergalactic medium, appearing at certain impact parameters due to projection effects. But a large number of them are associated with the outskirts of galaxies. This suggests that in the galaxy outskirts, the large δ values are mainly due to the inefficiency of mixing of star-forming gas with HI rather than dilution of heavy elements in HI through mixing with the pristine gas.

Mitchell et al. (2020) and Wright et al. (2021) used the EAGLE simulations and showed that accreting gas in massive galaxies have more contributions from the recycled (hence

Table 1. Systems with measured ionised and atomic gas metallicities, gathered from literature.

Name	z	$\log_{10}(N(\text{HI})/\text{cm}^{-2})$	b (kpc)	$Z_{\text{atomic}}/Z_{\odot}$	$Z_{\text{ionised}}/Z_{\odot}$	References
QSO-absorbers						
Q0302-223	1.009	20.36	25	-0.51 ± 0.12	0.04 ± 0.20	1,2
Q0452-1640	1.007	20.98	16	-0.96 ± 0.08	-0.46 ± 0.20	2,3
Q1009-0026	0.887	19.48	39	0.25 ± 0.06	0.04 ± 0.8	1,2
J1422-0001	0.91	20.40	12	-0.1 ± 0.4	0.04 ± 0.2	4
J1659+3735	0.1998	18.89	58	-1.12 ± 0.02	-0.21 ± 0.08	5
J2222-0946	2.354	20.65	6.3	-0.49 ± 0.05	-0.30 ± 0.13	2,3
HE2243-60	2.324	20.60	26	-0.72 ± 0.05	-0.32 ± 0.10	6
Q0235+164ID2	0.5243	21.70	13.2	-0.60 ± 0.41	-0.19 ± 0.15	7
Q0439-433	0.101	19.85	7.2	-0.20 ± 0.30	0.33 ± 0.14	7
Q0918+1636	2.5832	20.96	16.2	-0.12 ± 0.05	0.01 ± 0.20	7
Q2222-0946	2.354	20.65	6.3	-0.46 ± 0.07	-0.44 ± 0.19	7
J0238+1636	0.5253	21.70	7	-0.6 ± 0.4	-0.3 ± 0.2	8
J0304-2211	1.0095	20.36	25	-0.51 ± 0.12	0.04 ± 0.2	8
J0441-4313	0.1010	19.63	7	0.10 ± 0.15	0.50 ± 0.16	8
J0452-1640	1.007	20.98	16	-0.96 ± 0.10	-0.46 ± 0.20	8
J0918+1636	2.583	20.96	16	-0.12 ± 0.05	0.04 ± 0.2	8
J0958+0549	0.6546	20.54	20	-1.33 ± 0.23	-0.81 ± 0.06	8
J1009-0026	0.8864	19.48	39	0.25 ± 0.06	0.04 ± 0.8	8
J1138+0139	0.6126	21.25	12	-0.78 ± 0.16	-0.41 ± 0.18	8
J1204+0953	0.6390	21.04	10	-0.72 ± 0.16	-0.15 ± 0.14	8
J1436-0051	0.7390	20.08	50	-0.05 ± 0.12	0.33 ± 0.30	8
J1544+5912	0.0102	20.35	1	-0.50 ± 0.33	-0.4 ± 0.30	8
GRB host galaxies						
GRB 121024A	2.30	21.88		-0.6 ± 0.2	-0.1 ± 0.2	9
M83						
M83-3		18.94	0.2	1.19 ± 0.27	0.16 ± 0.15	10
M83-4		19.55	0.2	1.01 ± 0.23	0.17 ± 0.14	10
M83-6		20.72	3.1	0.12 ± 0.11	0.17 ± 0.20	10
M83-8		20.65	3.8	0.09 ± 0.09	0.24 ± 0.14	10
M83-9		20.56	5.1	0.08 ± 0.1	-0.05 ± 0.14	10
M83-12		20.98	3.3	-0.12 ± 0.07	0.15 ± 0.15	10
M83-14		20.43	3.2	0.18 ± 0.15	0.20 ± 0.14	10
M83-POS1		19.92	0.2	0.59 ± 0.09	0.31 ± 0.14	10
M83-POS2		18.91	0.2	0.69 ± 0.10	0.20 ± 0.14	10
Local dwarfs						
POX36		20.28		-1.38 ± 0.41	-0.61 ± 0.05	11
IZW18		21.34		-1.66 ± 0.15	-1.48 ± 0.06	12
SBS0335-052		21.86		-1.73 ± 0.55	-1.39 ± 0.01	13
IZw36		21.30		-1.76 ± 0.50	-0.89 ± 0.05	14
Mark59		20.85		-1.68 ± 0.3	-0.7 ± 0.003	15
NGC1705		20.18		-1.23 ± 0.22	-0.45 ± 0.07	16

Columns: (1) name of the system, (2) redshift, (3) HI column density, (4) impact parameter, (5) metallicity of atomic gas, (6) metallicity of ionised gas, (7) references.

References: (1) Péroux et al. (2011); (2) Augustin et al. (2018); (3) Péroux et al. (2012); (4) Bouché et al. (2016); (5) Kacprzak et al. (2014); (6) Bouché et al. (2013); (7) Christensen et al. (2014) and references therein; (8) Rahmani et al. (2016) and references therein; (9) Friis et al. (2015); (10) Hernandez et al. (2021); (11) Leboutteiller et al. (2009); (12) Leboutteiller et al. (2013); (13) Thuan et al. (2005); (14) Leboutteiller et al. (2004); (15) Thuan et al. (2002); (16) Heckman et al. (2001).

more metal-rich) gas compared to low mass galaxies. Given that the stellar mass has an insignificant role on the $Z_{\text{HI}}\text{-}Z_{\text{SFR}}$ relation, the findings of [Mitchell et al. \(2020\)](#) and [Wright et al. \(2021\)](#) support the dominant effect of the mixing of the two gas phases in the outskirts of galaxies.

We use the metallicity measurements of the ionised and atomic gas from observations, available in the literature, and compare them with the simulations. We find these observational measurements to be in a good agreement with the $Z_{\text{HI}}\text{-}Z_{\text{SFR}}$ relation of galaxies in the EAGLE simulations. There is however an indication that systems with starburst mode of star formation are likely to follow a different relation. Further analysis is required to investigate whether and how the presence of starburst can affect the metal enrichment of different gas phases.

Increasing the number of observational data points is essential for drawing conclusions on the relation between the metallicity of the atomic and ionised gas phases in galaxies. Although there are a very large number of intervening systems detected in quasar sightlines with accurate atomic gas metallicities, the presence of the bright background quasars make the ionised gas metallicity measurements quite challenging. Additionally, quasar sightlines typically trace the gas in the outskirts of the intervening systems, located at large distances ($\gtrsim 10$ kpc) from the star-forming regions where the ionised gas resides, making it even more challenging to measure the metallicity of the ionised gas in the same sightline. GRB sightlines on the other hand originate within the star-forming regions of their host galaxies. Therefore, the metallicities measured from the bright emission lines from the host galaxies are more likely to be representative of the metallicity of ionised gas in the GRBs sightlines. This provides a better chance to have a fair comparison between the metallicity of the ionised gas in GRBs sightlines (even without direct measurements) with the atomic gas metallicity obtained from imprinted absorption lines on GRBs spectra.

Moreover, GRBs are transients and hence they do not interfere with the observations of the bright emission lines from their host galaxies after they fade away. Observations with James Webb Space Telescope (JWST) will allow increasing the number of GRB hosts with metallicity measurements from both the methods ([Schady et al. 2021](#)). This, however, is limited by the number of detected GRB host galaxies.

Detailed studies of individual galaxies in sub-kpc scales, such as the study presented in [Hernandez et al. \(2021\)](#), are proposed to be carried out by combining the Integral Field Units (IFU) observations of galaxies with observations of bright background sources, to derive the metallicities of both the gas phases in similar regions of nearby galaxies (see [Kulkarni et al. 2020](#); [James et al. 2020](#)). Extending such studies to a large number of galaxies will allow us to test the findings presented in this paper, predicted by EAGLE, and will bring invaluable insights in our understanding of metal enrichment in galaxies.

ACKNOWLEDGMENTS

M.A. thanks Vianney Lebouteiller, Martin A. Zwaan, and Lise Christensen for helpful discussions. The presented study is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311. We acknowledge the Virgo Consortium for making their simulation data available. The EAGLE simulations were performed using the DiRAC-2 facility at Durham, managed by the ICC, and the PRACE facility Curie based in France at TGCC, CEA, Bruyères-le-Châtel. This work was supported by resources provided by the Pawsey Supercomputing Centre with funding from the Australian Government and the Government of Western Australia. D.O. is a recipient of an Australian Research Council Future Fellowship (FT190100083) funded by the Australian Government. Nastasha Wijers was supported by a CIERA Postdoctoral Fellowship.

REFERENCES

- Allende Prieto, C., Lambert, D. L., & Asplund, M. 2001, *ApJ*, 556, L63, doi: [10.1086/322874](#)
- . 2002, *ApJL*, 573, L137, doi: [10.1086/342095](#)
- Arabsalmani, M., Møller, P., Fynbo, J. P. U., et al. 2015, *MNRAS*, 446, 990, doi: [10.1093/mnras/stu2138](#)
- Arabsalmani, M., Møller, P., Perley, D. A., et al. 2018, *MNRAS*, 473, 3312, doi: [10.1093/mnras/stx2451](#)
- Augustin, R., Péroux, C., Møller, P., et al. 2018, *MNRAS*, 478, 3120, doi: [10.1093/mnras/sty1287](#)
- Bahé, Y. M., Crain, R. A., Kauffmann, G., et al. 2016, *MNRAS*, 456, 1115, doi: [10.1093/mnras/stv2674](#)
- Bothwell, M. S., Maiolino, R., Kennicutt, R., et al. 2013, *MNRAS*, 433, 1425, doi: [10.1093/mnras/stt817](#)
- Bouché, N., Murphy, M. T., Kacprzak, G. G., et al. 2013, *Science*, 341, 50, doi: [10.1126/science.1234209](#)
- Bouché, N., Finley, H., Schroetter, I., et al. 2016, *ApJ*, 820, 121, doi: [10.3847/0004-637X/820/2/121](#)
- Cannon, J. M., Skillman, E. D., Sembach, K. R., & Bomans, D. J. 2005, *ApJ*, 618, 247, doi: [10.1086/425897](#)
- Christensen, L., Møller, P., Fynbo, J. P. U., & Zafar, T. 2014, *MNRAS*, 445, 225, doi: [10.1093/mnras/stu1726](#)
- Collacchioni, F., Lagos, C. D. P., Mitchell, P. D., et al. 2020a, *MNRAS*, 495, 2827, doi: [10.1093/mnras/staa1334](#)
- . 2020b, *MNRAS*, 495, 2827, doi: [10.1093/mnras/staa1334](#)
- Crain, R. A., Schaye, J., Bower, R. G., et al. 2015a, *MNRAS*, 450, 1937, doi: [10.1093/mnras/stv725](#)

- . 2015b, MNRAS, 450, 1937, doi: [10.1093/mnras/stv725](https://doi.org/10.1093/mnras/stv725)
- Crain, R. A., Bahé, Y. M., Lagos, C. d. P., et al. 2017, MNRAS, 464, 4204, doi: [10.1093/mnras/stw2586](https://doi.org/10.1093/mnras/stw2586)
- Cullen, L., & Dehnen, W. 2010, MNRAS, 408, 669, doi: [10.1111/j.1365-2966.2010.17158.x](https://doi.org/10.1111/j.1365-2966.2010.17158.x)
- Dayal, P., & Ferrara, A. 2018, PhR, 780, 1, doi: [10.1016/j.physrep.2018.10.002](https://doi.org/10.1016/j.physrep.2018.10.002)
- De Rossi, M. E., Bower, R. G., Font, A. S., Schaye, J., & Theuns, T. 2017, MNRAS, 472, 3354, doi: [10.1093/mnras/stx2158](https://doi.org/10.1093/mnras/stx2158)
- Diemer, B., Stevens, A. R. H., Forbes, J. C., et al. 2018, ApJS, 238, 33, doi: [10.3847/1538-4365/aae387](https://doi.org/10.3847/1538-4365/aae387)
- Dopita, M. A., Blair, W. P., Long, K. S., et al. 2010, ApJ, 710, 964, doi: [10.1088/0004-637X/710/2/964](https://doi.org/10.1088/0004-637X/710/2/964)
- Dubois, Y., Pichon, C., Welker, C., et al. 2014, MNRAS, 444, 1453, doi: [10.1093/mnras/stu1227](https://doi.org/10.1093/mnras/stu1227)
- Durier, F., & Dalla Vecchia, C. 2012, MNRAS, 419, 465, doi: [10.1111/j.1365-2966.2011.19712.x](https://doi.org/10.1111/j.1365-2966.2011.19712.x)
- Ferland, G. J., Korista, K. T., Verner, D. A., et al. 1998, Publications of the Astronomical Society of the Pacific, 110, 761, doi: [10.1086/316190](https://doi.org/10.1086/316190)
- Friis, M., De Cia, A., Krühler, T., et al. 2015, MNRAS, 451, 167, doi: [10.1093/mnras/stv960](https://doi.org/10.1093/mnras/stv960)
- Garratt-Smithson, L., Power, C., Lagos, C. d. P., et al. 2021, MNRAS, 501, 4396, doi: [10.1093/mnras/staa3870](https://doi.org/10.1093/mnras/staa3870)
- Graham, A. W., & Worley, C. C. 2008, MNRAS, 388, 1708, doi: [10.1111/j.1365-2966.2008.13506.x](https://doi.org/10.1111/j.1365-2966.2008.13506.x)
- Grevesse, N., & Sauval, A. J. 1998, SSRv, 85, 161, doi: [10.1023/A:1005161325181](https://doi.org/10.1023/A:1005161325181)
- Haardt, F., & Madau, P. 2012, ApJ, 746, 125, doi: [10.1088/0004-637X/746/2/125](https://doi.org/10.1088/0004-637X/746/2/125)
- Hamanowicz, A., Péroux, C., Zwaan, M. A., et al. 2020, MNRAS, 492, 2347, doi: [10.1093/mnras/stz3590](https://doi.org/10.1093/mnras/stz3590)
- Hartoog, O. E., Malesani, D., Fynbo, J. P. U., et al. 2015, A&A, 580, A139, doi: [10.1051/0004-6361/201425001](https://doi.org/10.1051/0004-6361/201425001)
- Heckman, T. M., Sembach, K. R., Meurer, G. R., et al. 2001, ApJ, 554, 1021, doi: [10.1086/321422](https://doi.org/10.1086/321422)
- Hernandez, S., Aloisi, A., James, B. L., et al. 2021, ApJ, 908, 226, doi: [10.3847/1538-4357/abd6c4](https://doi.org/10.3847/1538-4357/abd6c4)
- Holweger, H. 2001, in American Institute of Physics Conference Series, Vol. 598, Joint SOHO/ACE workshop “Solar and Galactic Composition”, ed. R. F. Wimmer-Schweingruber, 23–30, doi: [10.1063/1.1433974](https://doi.org/10.1063/1.1433974)
- Hopkins, P. F. 2013, MNRAS, 428, 2840, doi: [10.1093/mnras/sts210](https://doi.org/10.1093/mnras/sts210)
- James, B. L., Aloisi, A., Emerick, A., Hernandez, S. S., & Kumari, N. 2020, Pinning down multi-phase mixing of metals within star-forming galaxies, HST Proposal
- Kacprzak, G. G., Martin, C. L., Bouché, N., et al. 2014, ApJL, 792, L12, doi: [10.1088/2041-8205/792/1/L12](https://doi.org/10.1088/2041-8205/792/1/L12)
- Kewley, L. J., & Ellison, S. L. 2008, ApJ, 681, 1183, doi: [10.1086/587500](https://doi.org/10.1086/587500)
- Krumholz, M. R. 2013, MNRAS, 436, 2747, doi: [10.1093/mnras/stt1780](https://doi.org/10.1093/mnras/stt1780)
- Krumholz, M. R., McKee, C. F., & Tumlinson, J. 2009, ApJ, 699, 850, doi: [10.1088/0004-637X/699/1/850](https://doi.org/10.1088/0004-637X/699/1/850)
- Kudritzki, R.-P., Ho, I. T., Schruha, A., et al. 2015, MNRAS, 450, 342, doi: [10.1093/mnras/stv522](https://doi.org/10.1093/mnras/stv522)
- Kulkarni, V. P., Bershad, M. A., Lundgren, B. F., et al. 2020, The Baryonic Content of Galaxies Mapped by MaNGA and Gas Flows Around Them, HST Proposal
- Lagos, C. d. P., Crain, R. A., Schaye, J., et al. 2015, MNRAS, 452, 3815, doi: [10.1093/mnras/stv1488](https://doi.org/10.1093/mnras/stv1488)
- Lagos, C. d. P., Theuns, T., Schaye, J., et al. 2016, MNRAS, 459, 2632, doi: [10.1093/mnras/stw717](https://doi.org/10.1093/mnras/stw717)
- Lebouteiller, V., Heap, S., Hubeny, I., & Kunth, D. 2013, A&A, 553, A16, doi: [10.1051/0004-6361/201220948](https://doi.org/10.1051/0004-6361/201220948)
- Lebouteiller, V., Kunth, D., Lequeux, J., et al. 2004, A&A, 415, 55, doi: [10.1051/0004-6361:20034592](https://doi.org/10.1051/0004-6361:20034592)
- Lebouteiller, V., Kunth, D., Thuan, T. X., & Désert, J. M. 2009, A&A, 494, 915, doi: [10.1051/0004-6361:200811089](https://doi.org/10.1051/0004-6361:200811089)
- Lequeux, J., Peimbert, M., Rayo, J. F., Serrano, A., & Torres-Peimbert, S. 1979, A&A, 80, 155
- Lyman, J. D., Levan, A. J., Tanvir, N. R., et al. 2017, MNRAS, 467, 1795, doi: [10.1093/mnras/stx220](https://doi.org/10.1093/mnras/stx220)
- Maio, U., Péroux, C., & Ciardi, B. 2022, A&A, 657, A47, doi: [10.1051/0004-6361/202142264](https://doi.org/10.1051/0004-6361/202142264)
- Maiolino, R., Nagao, T., Grazian, A., et al. 2008, A&A, 488, 463, doi: [10.1051/0004-6361:200809678](https://doi.org/10.1051/0004-6361:200809678)
- Maltby, D. T., Aragón-Salamanca, A., Gray, M. E., et al. 2010, MNRAS, 402, 282, doi: [10.1111/j.1365-2966.2009.15953.x](https://doi.org/10.1111/j.1365-2966.2009.15953.x)
- Mannucci, F., Cresci, G., Maiolino, R., Marconi, A., & Gnerucci, A. 2010, MNRAS, 408, 2115, doi: [10.1111/j.1365-2966.2010.17291.x](https://doi.org/10.1111/j.1365-2966.2010.17291.x)
- Mannucci, F., Cresci, G., Maiolino, R., et al. 2009, MNRAS, 398, 1915, doi: [10.1111/j.1365-2966.2009.15185.x](https://doi.org/10.1111/j.1365-2966.2009.15185.x)
- Marigo, P. 2001, A&A, 370, 194, doi: [10.1051/0004-6361:20000247](https://doi.org/10.1051/0004-6361:20000247)
- Matteucci, F. 2012, Chemical Evolution of Galaxies, doi: [10.1007/978-3-642-22491-1](https://doi.org/10.1007/978-3-642-22491-1)
- McAlpine, S., Helly, J. C., Schaller, M., et al. 2016, Astronomy and Computing, 15, 72, doi: [10.1016/j.ascom.2016.02.004](https://doi.org/10.1016/j.ascom.2016.02.004)
- McKee, C. F., & Krumholz, M. R. 2010, ApJ, 709, 308, doi: [10.1088/0004-637X/709/1/308](https://doi.org/10.1088/0004-637X/709/1/308)
- Mitchell, P. D., Schaye, J., & Bower, R. G. 2020, MNRAS, 497, 4495, doi: [10.1093/mnras/staa2252](https://doi.org/10.1093/mnras/staa2252)
- Møller, P., Fynbo, J. P. U., Ledoux, C., & Nilsson, K. K. 2013, MNRAS, 430, 2680, doi: [10.1093/mnras/stt067](https://doi.org/10.1093/mnras/stt067)
- Naab, T., & Ostriker, J. P. 2017, ARA&A, 55, 59, doi: [10.1146/annurev-astro-081913-040019](https://doi.org/10.1146/annurev-astro-081913-040019)

- Péroux, C., Bouché, N., Kulkarni, V. P., York, D. G., & Vladilo, G. 2011, *MNRAS*, 410, 2237, doi: [10.1111/j.1365-2966.2010.17598.x](https://doi.org/10.1111/j.1365-2966.2010.17598.x)
- . 2012, *MNRAS*, 419, 3060, doi: [10.1111/j.1365-2966.2011.19947.x](https://doi.org/10.1111/j.1365-2966.2011.19947.x)
- Péroux, C., Quiret, S., Rahmani, H., et al. 2016, *MNRAS*, 457, 903, doi: [10.1093/mnras/stw016](https://doi.org/10.1093/mnras/stw016)
- Pillepich, A., Springel, V., Nelson, D., et al. 2018, *MNRAS*, 473, 4077, doi: [10.1093/mnras/stx2656](https://doi.org/10.1093/mnras/stx2656)
- Portinari, L., Chiosi, C., & Bressan, A. 1998, *A&A*, 334, 505, <https://arxiv.org/abs/astro-ph/9711337>
- Price, D. J. 2008, *Journal of Computational Physics*, 227, 10040, doi: [10.1016/j.jcp.2008.08.011](https://doi.org/10.1016/j.jcp.2008.08.011)
- Rahmani, H., Péroux, C., Turnshek, D. A., et al. 2016, *MNRAS*, 463, 980, doi: [10.1093/mnras/stw1965](https://doi.org/10.1093/mnras/stw1965)
- Rahmati, A., Pawlik, A. H., Raičević, M., & Schaye, J. 2013, *MNRAS*, 430, 2427, doi: [10.1093/mnras/stt066](https://doi.org/10.1093/mnras/stt066)
- Rahmati, A., Schaye, J., Bower, R. G., et al. 2015, *MNRAS*, 452, 2034, doi: [10.1093/mnras/stv1414](https://doi.org/10.1093/mnras/stv1414)
- Schady, P., Tanvir, N. R., Arabsalmani, M., et al. 2021, Mapping emission and absorption line metallicities onto the same universal scale, JWST Proposal. Cycle 1
- Schaller, M., Dalla Vecchia, C., Schaye, J., et al. 2015, *MNRAS*, 454, 2277, doi: [10.1093/mnras/stv2169](https://doi.org/10.1093/mnras/stv2169)
- Schaye, J. 2004, *ApJ*, 609, 667, doi: [10.1086/421232](https://doi.org/10.1086/421232)
- Schaye, J., & Dalla Vecchia, C. 2008, *MNRAS*, 383, 1210, doi: [10.1111/j.1365-2966.2007.12639.x](https://doi.org/10.1111/j.1365-2966.2007.12639.x)
- Schaye, J., Crain, R. A., Bower, R. G., et al. 2015, *MNRAS*, 446, 521, doi: [10.1093/mnras/stu2058](https://doi.org/10.1093/mnras/stu2058)
- Shen, S., Mo, H. J., White, S. D. M., et al. 2003, *MNRAS*, 343, 978, doi: [10.1046/j.1365-8711.2003.06740.x](https://doi.org/10.1046/j.1365-8711.2003.06740.x)
- Silk, J., & Mamon, G. A. 2012, *Research in Astronomy and Astrophysics*, 12, 917, doi: [10.1088/1674-4527/12/8/004](https://doi.org/10.1088/1674-4527/12/8/004)
- Sparre, M., Hartoog, O. E., Krühler, T., et al. 2014, *ApJ*, 785, 150, doi: [10.1088/0004-637X/785/2/150](https://doi.org/10.1088/0004-637X/785/2/150)
- Springel, V. 2005, *MNRAS*, 364, 1105, doi: [10.1111/j.1365-2966.2005.09655.x](https://doi.org/10.1111/j.1365-2966.2005.09655.x)
- Steidel, C. C., Rudie, G. C., Strom, A. L., et al. 2014, *ApJ*, 795, 165, doi: [10.1088/0004-637X/795/2/165](https://doi.org/10.1088/0004-637X/795/2/165)
- The EAGLE team. 2017a, arXiv e-prints, arXiv:1706.09899, <https://arxiv.org/abs/1706.09899>
- . 2017b, arXiv e-prints, arXiv:1706.09899, <https://arxiv.org/abs/1706.09899>
- Thuan, T. X., Lecavelier des Etangs, A., & Izotov, Y. I. 2002, *ApJ*, 565, 941, doi: [10.1086/324684](https://doi.org/10.1086/324684)
- . 2005, *ApJ*, 621, 269, doi: [10.1086/427469](https://doi.org/10.1086/427469)
- Tissera, P. B., Rosas-Guevara, Y., Bower, R. G., et al. 2019, *MNRAS*, 482, 2208, doi: [10.1093/mnras/sty2817](https://doi.org/10.1093/mnras/sty2817)
- Tissera, P. B., Rosas-Guevara, Y., Sillero, E., et al. 2022, *MNRAS*, 511, 1667, doi: [10.1093/mnras/stab3644](https://doi.org/10.1093/mnras/stab3644)
- Trayford, J. W., & Schaye, J. 2019, *MNRAS*, 485, 5715, doi: [10.1093/mnras/stz757](https://doi.org/10.1093/mnras/stz757)
- Tremonti, C. A., Heckman, T. M., Kauffmann, G., et al. 2004, *ApJ*, 613, 898, doi: [10.1086/423264](https://doi.org/10.1086/423264)
- van der Wel, A., Franx, M., van Dokkum, P. G., et al. 2014, *ApJ*, 788, 28, doi: [10.1088/0004-637X/788/1/28](https://doi.org/10.1088/0004-637X/788/1/28)
- van Loon, M. L., Mitchell, P. D., & Schaye, J. 2021, *MNRAS*, 504, 4817, doi: [10.1093/mnras/stab1254](https://doi.org/10.1093/mnras/stab1254)
- Wiersma, R. P. C., Schaye, J., Theuns, T., Dalla Vecchia, C., & Tornatore, L. 2009a, *MNRAS*, 399, 574, doi: [10.1111/j.1365-2966.2009.15331.x](https://doi.org/10.1111/j.1365-2966.2009.15331.x)
- Wiersma, R. P. C., Schaye, J., Theuns, T., Dalla Vecchia, C., & Tornatore, L. 2009b, *MNRAS*, 399, 574, doi: [10.1111/j.1365-2966.2009.15331.x](https://doi.org/10.1111/j.1365-2966.2009.15331.x)
- Wofford, A., Leitherer, C., & Chandar, R. 2011, *ApJ*, 727, 100, doi: [10.1088/0004-637X/727/2/100](https://doi.org/10.1088/0004-637X/727/2/100)
- Wolfire, M. G., McKee, C. F., Hollenbach, D., & Tielens, A. G. G. M. 2003, *ApJ*, 587, 278, doi: [10.1086/368016](https://doi.org/10.1086/368016)
- Wright, R. J., Lagos, C. d. P., Power, C., & Correa, C. A. 2021, *MNRAS*, 504, 5702, doi: [10.1093/mnras/stab1057](https://doi.org/10.1093/mnras/stab1057)
- Yabe, K., Ohta, K., Akiyama, M., et al. 2015, *PASJ*, 67, 102, doi: [10.1093/pasj/psv079](https://doi.org/10.1093/pasj/psv079)
- Yates, R. M., Péroux, C., & Nelson, D. 2021, *MNRAS*, 508, 3535, doi: [10.1093/mnras/stab2837](https://doi.org/10.1093/mnras/stab2837)
- Zahid, H. J., Dima, G. I., Kudritzki, R.-P., et al. 2014, *ApJ*, 791, 130, doi: [10.1088/0004-637X/791/2/130](https://doi.org/10.1088/0004-637X/791/2/130)




Figure 8. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plane for all the sightlines in EAGLE galaxies at $z = 2$ with smoothed metallicities. The medians of the Z_{SFR} values in each Z_{HI} bin are shown with pink diamonds. The shaded area shows the 1σ spread around the medians of the Z_{SFR} values. The $x=y$ line in the plane is presented by a dashed line.

APPENDIX

In the main body of the paper, we have used the metallicities associated with each particle, as opposed to the SPH-smoothed version of the metallicities. Here we present all the plots presented in Section 3 using the SPH-smoothed version of the particle metallicities.




Figure 9. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plane with smoothed metallicities for the sightlines at $z = 2$ in three bins of stellar mass (rows) and four bins of impact parameter (columns). The colour of the points indicates the $N(\text{HI})$ value of each sightline, with the colour bar presented in the right side of each row. The grey lines show the $Z_{\text{SFR},\text{med}}$ values in each plot. The $x=y$ line in the plane is presented by a dashed line. Note that R_{200} varies between $\sim 40 - 90$ kpc for galaxies with stellar masses between 10^8 and $10^{10} M_{\odot}$.




Figure 10. The $Z_{\text{HI}}\text{--}Z_{\text{SFR}}$ plane with smoothed metallicities for sightlines at $z = 2$ selected based on $N(\text{HI})$ and impact parameter. The $Z_{\text{SFR,med}}$ values are shown with black diamonds and the $x=y$ line is presented with the dashed lines. **Top panels:** Sightlines with HI column densities of $10^{20} \text{ cm}^{-2} \leq N(\text{HI}) < 10^{21} \text{ cm}^{-2}$ and $N(\text{HI}) \geq 10^{21} \text{ cm}^{-2}$ in the left and right panels respectively. All the presented sightlines are within the $0.3 R_{200}$ from galaxy centres. The colour bar represents the value for b/R_{200} . **Bottom panels:** Sightlines with $0.02 < b/R_{200} \leq 0.05$ in left and those within the $0.02 R_{200}$ central regions of galaxies in the right panel respectively. The colour bar represents the HI column density of the sightlines.




Figure 11. Left: The $Z_{\text{HI}}\text{-}Z_{\text{SFR}}$ relation with smoothed metallicities for all the sightlines of EAGLE galaxies at redshifts $z = 0, 1, 2$ and 3 . The $Z_{\text{SFR,med}}$ values are marked with diamonds and the 1σ scatter around the medians are shown with shaded areas. **Right:** The $Z_{\text{SFR,med}}$ values of smoothed metallicities and the 1σ scatter around them for sightlines within the $0.02 R_{200}$ central regions of galaxies in four redshifts.




Figure 12. The ionised and atomic gas metallicity measurements from observations, in comparison with the obtained metallicities of EAGLE galaxies using smoothed metallicities. The quasar absorbers are shown with pink squares, and the host galaxy of GRB 121024 is presented with a purple square. Nearby dwarf galaxies are marked by black diamonds. The individual regions of M83 are shown by gray stars. The x values for these systems are measured in single sightline from the ratio between the column densities of heavy elements and HI, obtained from ISM absorption lines. The y values for these systems are obtained from the flux measurement of the bright nebular lines coming from the entire star-forming disk of the galaxy. The size of the squares are proportional to the redshift of the system. The blue points show the EAGLE galaxies at $z = 0 - 3$ in the $Z_{\text{HI,mean}}-Z_{\text{SFR,mean}}$ plane. The quasar absorbers are shown with pink squares and the host galaxy of GRB 121024 is presented with a purple square. The x values for these systems are measured in single sightline from the ratio between the column densities of heavy elements and HI, obtained from ISM absorption lines. The y values for these systems are obtained from the flux measurement of the bright nebular lines coming from the entire star-forming disk of the galaxy. The size of the squares are proportional to the redshift of the system. The blue points show the EAGLE galaxies at $z = 0 - 3$ in the $Z_{\text{HI,mean}}-Z_{\text{SFR,mean}}$ plane.