

# Three New Models of Layered Inhomogeneous Elliptical Galaxies

S. A. Gasanov

Sternberg Astronomical Institute, Lomonosov Moscow State University, Moscow, 119991  
Russia e-mail: gasanovsa57@gmail.com

**Abstract.** To solve some problems of celestial mechanics and astrophysics, three new models of an elliptical galaxy (EG) have been created, which are in good agreement with modern understanding of the structure of such galaxies. Based on these models, the total gravitational (potential) energy and rotational kinetic energy of an EG, as well as the velocity dispersion at a distance of its effective radius, are determined. A new method is proposed for determining the average values of the scale radius of an EG and the density at its center, as well as the average value of its key parameter density  $\beta$  and its value at a distance corresponding to the effective radius of the galaxy. The results obtained are applied to sixty EGs and presented in the form of tables for ten galaxies.

**Keywords:** elliptical galaxy, new models of elliptical galaxies, dynamic parameters, gravitational energy, rotational kinetic energy

## 1. INTRODUCTION

In [1, 2], the problem of the spatial motion of a passively gravitating body (a star or the center of mass of a globular cluster, GC) inside a rotating elliptical galaxy (EG) was considered. An EG was regarded as a two - layer ellipsoidal body: its luminous part is a triaxial ellipsoid, and the space between the boundaries of its luminous part and the halo is a homeoid filled with homogeneous dark matter.

In [1], the luminous part of an EG (LP EG) is considered a homogeneous triaxial ellipsoid, and, in [2], uses the so-called astrophysical law of density distribution. The EG model considered in [1] will be called model 1, and the model in [2], model 2. Within models 1 and 2, an analogue of the Jacobi integral was found and the region of possible motion of the star (or the mass center of the GC) was determined. The type and Lyapunov stability of the stationary solutions found libration points — were established, and zero velocity surfaces were constructed.

The problem of the spatial motion of a star inside (near) a globular cluster belonging to an inhomogeneous rotating EG was considered in [3]. The motion of a star near a GC was considered with allowance for the perturbations caused by the attraction of the EG, which, together with the halo, is a two-layer body [1]. The motion of a star near a GC takes place outside the luminous part of the EG, but inside the homeoid. The notion "near a GC" as a "sphere of action" (Hill's gravitational sphere) has been refined and concretized. In connection with the introduced concept of the sphere of action, two variants of the motion of a star were considered: inside and outside the sphere of action of a globular cluster, and the regions of the possibility of motion were determined. A quasi - integral and minimum - energy surfaces were found, which, under certain conditions, are transformed into an analog of the Jacobi integral and zero-velocity surfaces, respectively.

The results of [1–3] are applied to model elliptical galaxies with parameters that exactly coincide with the parameters of the elliptical galaxies NGC 4472 (M 49), NGC 4636, and NGC 4374 (M 84) and are presented in the form of figures and tables. When finding libration points

and analyzing their stability, exact expressions for the potentials of the LP EG and the homeoid rather than their series expansions are used.

In this paper, three new models of an EG are considered, in which a galaxy with a halo is considered a two-layer inhomogeneous ellipsoid of revolution: a spheroid. In this case, the outer and inner layers are assumed to be similar and concentric, and their centers coincide with the center of the EG. The LP EG is considered an inner layer and comprises an inhomogeneous ellipsoid of revolution (spheroid) with a homothetic (spheroidal) density distribution, or a layered inhomogeneous spheroid.

In the LP EG, baryon mass (BM) with an “astrophysical law” of density distribution prevails. The outer part is an inhomogeneous spherical layer with a spherical density distribution (model 3) or a spheroidal layer (conditionally, a homeoid) with a spheroidal density distribution (model 4). According to model 3, the outer layer and the halo of a galaxy are bounded by a sphere with a radius equal to the scale radius of the EG, and, according to model 4, they are bounded by a spheroidal surface with a semi-major axis equal to the scale radius of the galaxy. It is believed that the spherical layer and the conditional homeoid mainly consists of dark matter (DM) and, depending on its presence in the inner (central) regions of the EG, in models 3 and 4, two variants are considered. Variant (a), in which the main part of DM is outside the LP EG [4], and variant (b), in which the DM content in the inner regions of the EG is comparable to the BM content [5, 6]. In this case, the matching conditions for the potentials at the interface between the LP EG and the spherical layer or conditional homeoid are determined.

As correctly noted in [4], the nature of DM is unknown, and there is no clear understanding of its physical relationship with the observed astronomical objects. Nevertheless, its presence in galaxies is recognized and indirectly confirmed. In this work, three types of EG together with a halo are simulated, which cannot claim to be comprehensive of the DM problem as a whole.

According to model 5, an EG with (variant 1) or without (variant 2) a halo comprises a layered inhomogeneous spheroid consisting of BM and DM. In model 5, there is no interface between the LP EG and the conditional homeoid; therefore, the fulfillment of the matching conditions for the potentials is not considered. In models 3–5, the conditional boundaries (apparent sizes) of the LP EG are determined by the quantities  $D_{25}$  and  $R_{25}$  [7].

Beside the problem of the presence or absence of DM in the inner regions of an EG, there is another problem, namely, the true spatial orientation of the galaxy, which is unknown to us. Finding the true shape of such galaxies is of great importance for constructing a dynamic theory of equilibrium and the theory of the origin of EGs. As shown in [8], the true shape of an EG can be determined on the basis of two observational tests of 1) whether the rotation axis coincides with the apparent minor axis of the galaxy and 2) whether the isophotes of the galaxy are aligned. The use of these tests allows one to judge more confidently about the shape of EGs and indicates the existence of EGs both in the form of oblate or prolate spheroids and triaxial ellipsoids. If an EG has the shape of a prolate spheroid or a triaxial ellipsoid, then the rotation axis will, as a rule, be observed as not aligned with the apparent minor axis and the isophotes of the galaxy will be misaligned. If the galaxy has the shape of an oblate spheroid, then the projection of the minor axis onto the plane of the sky coincides with the direction of the rotation axis for any orientation of the galaxy relative to the observer and the isophote alignment is not disturbed.

In [9], a new method for solving the inverse geometric problem of recovering the shape of ellipsoidal bodies through their projection (limb) onto the plane of the sky was developed and used. Using this method, the semiaxes of the dwarf planet Haumea as a triaxial ellipsoid were determined. Thus, for each value of the photometric parameters of the dwarf planet, its shape and average density were determined, as well as the orientation relative to its ring and the orbits of satellites. It turned out that the planes of the planet's ring and the orbit of its satellite Hi'iaka do not coincide with the planet's equator plane and both satellites are in direct motion [9].

It is important to note that the method developed in [9] and applied to the dwarf planet Haumea cannot be used to recover the true shape of the EG as a triaxial ellipsoidal body or spheroid. In the present work, an EG is regarded as a layered inhomogeneous prolate ellipsoid of revolution (or a prolate spheroid for short). For a triaxial ellipsoid, the rotation axis does not coincide with the apparent minor axis and the isophotes will be misaligned. A more convenient or simpler variant in which an EG is an oblate spheroid, with its apparent axis coinciding with the axis of rotation and the isophote alignment not disturbed, is not considered.

Within the created models, a new method is proposed for determining the average values of the EG scale radius  $r_s$  and the densities in the center,  $\rho_0$ , and at the boundary of the galactic halo,  $\rho_s$ . The total gravitational (potential) energy  $W$  and rotational kinetic energy  $T_{rot}$  of an inhomogeneous EG, the velocity dispersion  $\sigma_{eff}$  at a distance of the effective radius of the galaxy  $R_{eff}$ , as well as the average value of the parameter  $\beta$  and its value  $\beta_{eff}$  corresponding to the effective radius of the galaxy, are determined according to these models.

Thus, it is currently not possible to accurately determine the true shape of an EG as a triaxial ellipsoid or even a spheroid. Therefore, we determine the values of the apparent major and minor semiaxes of the EG. Thus, the obtained values of the dynamic parameters listed above are approximate.

The problem of equilibrium and stability of dynamical systems appearing in these models is of particular interest and will be considered in a separate work.

## 2. DENSITY DISTRIBUTION LAWS

Assume that the luminous part of an EG (LP EG) is a layered inhomogeneous ellipsoid of revolution (prolate spheroid) with semiaxes  $a > b = c$ . As the density distribution law (LP EG profile), consisting of the BM, we take the "astrophysical" profile  $\rho(m)$  [10, 11] obtained by applying the Abel integral equation to the Hubble surface brightness profile  $I(m)$  [12]:

$$\rho(m) = \frac{\rho_0}{(1 + \beta m^2)^{3/2}}, \quad I(m) = \frac{I_0}{1 + \beta m^2}, \quad (1)$$

where  $\rho_0$  and  $I_0$  are the density at the center of the EG and the central surface brightness and the parameter  $\beta \gg 1$  for each EG is selected separately and is found by aligning the photometry data [10, 11]. In addition,  $m$  is a parameter of a family of similar and concentric spheroids

$$\frac{x^2}{a^2} + \frac{y^2 + z^2}{c^2} = m^2, \quad (a > b = c, \quad 0 \leq m \leq 1), \quad (2)$$

Here,  $m = 0$  corresponds to the center of the EG and  $m = 1$  corresponds to the spheroidal surface limiting the LP EG.

Next, we assume that an inhomogeneous spherical layer (model 3) is filled with DM with a density distribution law (profile) [13, 14]

$$\rho_S(r) = \frac{K r_s}{r} \left(1 + \frac{r}{r_s}\right)^{-2}, \quad \tilde{\rho}_S \equiv \rho_S(r_s) = \frac{K}{4} \quad (3)$$

Here,  $K$  is a normalizing coefficient, which has the dimension of density in the solar masses per cubic parsec,  $r_s$  is the scale radius of the EG, and  $\tilde{\rho}_S$  is the density of DM at the outer boundary of the layer. Formulas for calculating the coefficient  $K$  are given in [3, 13,14].

It should be noted that, in addition to profile (3), there are other laws of density distribution. In the book [15], a more general profile is given, from which, as special cases, the Dehnen [16, 17], Hernquist [18], Jaffe [19], and NFW [13] profiles are obtained. If an inhomogeneous spheroidal layer (model 4) filled with DM is considered, then profile (3) cannot be used. In this case, the density distribution law, which we call an analog of the NFW profile, is defined as

$$\rho_G(\mu) = \frac{K}{\xi \mu (1 + \xi \mu)^2}, \quad \left( \frac{r}{r_s} = \xi \mu, \quad \xi = \frac{\sqrt[3]{\tilde{a} \tilde{c}^2}}{r_s} < 1 \right), \quad (4)$$

which was proposed by B.P. Kondratyev. Here,  $\mu$  is the parameter of a family of similar and concentric spheroids with semiaxes  $\tilde{a}$  and  $\tilde{b} = \tilde{c}$ :

$$\mu^2 = \frac{x^2}{\tilde{a}^2} + \frac{y^2 + z^2}{\tilde{c}^2}, \quad (\tilde{a} = r_s > \tilde{b} = \tilde{c}) \quad (5)$$

In this case, families (2) and (5) are assumed to be similar and concentric spheroids. The value  $\mu^2 = 1$  corresponds to the outer boundary of the spheroidal layer, at which, according to expression (4), the density is equal to

$$\bar{\rho}_G \equiv \rho_G(1) = \frac{K}{\xi (1 + \xi)^2} \quad (6)$$

### 3. POTENTIAL ENERGY AND ROTATIONAL KINETIC ENERGY OF THE LUMINOUS PART OF AN ELLIPTICAL GALAXY

The total potential (gravitational) energy  $W(1)$  and rotational kinetic energy  $T(1)$  of a layered inhomogeneous spheroid with density and semiaxes  $a > b = c$  are determined by the formulas [10]

$$W(1) = -\pi G a c^2 J_0 \Psi(1), \quad T(1) = \frac{\pi G a c^2}{2} J_1 \Psi(1), \quad J_1 = J_0 - 3c^2 K_0 \quad (7)$$

respectively. Here,  $\Psi(1)$  is the value of the function

$$\Psi(m) = \int_0^{m^2} \rho(m) M(m) dm^2, \quad M(m) = 4\pi a c^2 \int_0^m m^2 \rho(m) dm \quad (8)$$

at  $m = 1$ , and

$$J_0 = \int_0^\infty \frac{du}{\Delta(u)}, \quad K_0 = \int_0^\infty \frac{du}{(c^2 + u)\Delta(u)}, \quad \Delta(u) = (c^2 + u)\sqrt{a^2 + u}, \quad (9)$$

or

$$J_0 = \frac{1}{\sqrt{a^2 - c^2}} \ln \frac{a + \sqrt{a^2 - c^2}}{a - \sqrt{a^2 - c^2}}, \quad K_0 = \frac{2a - c^2 J_0}{2c^2(a^2 - c^2)}, \quad J_1 = \frac{(2a^2 + c^2)J_0 - 6a}{2(a^2 - c^2)} \quad (10)$$

If the density  $\rho(m)$  of the LP EG is determined by astrophysical law (1), then, by virtue of (8), the mass  $M(m)$  of the intermediate ellipsoid and the function  $\Psi(m)$  will be

$$M(m) = 4\pi\rho_0 a c^2 g_1(m), \quad \Psi(m) = 4\pi\rho_0^2 a c^2 f_1(m) \quad (11)$$

Here

$$g_1(m) = \frac{1}{\beta\sqrt{\beta}} \left[ \ln \varphi_1(m) - \frac{h}{\sqrt{1+h^2}} \right], \quad \varphi_1(m) = h + \sqrt{1+h^2}, \quad h = \sqrt{\beta} m \quad (12)$$

$$f_1(m) = \frac{1}{\beta^2\sqrt{\beta}} \left[ \arctg h + \frac{h}{1+h^2} - \frac{2 \ln \varphi_1(m)}{\sqrt{1+h^2}} \right] \quad (13)$$

Then, by virtue of (7), we find the gravitational energy  $W_L = W(1)$  of the LP EG and its rotational kinetic energy  $T_L = T(1)$ , according to variant (a), in the form

$$a) \quad W_L = -2W_0 J_0, \quad T_L = W_0 J_1, \quad W_0 = 2\pi^2 G \rho_0^2 a^2 c^4 f_1(1) \quad (14)$$

where the function  $f_1(m)$  and coefficients  $J_0$  and  $J_1$  are defined above.

In variant (b), the gravitational energy  $\tilde{W}_L$  of the LP EG and its rotational kinetic energy  $\tilde{T}_L$  are determined differently; namely, in formulas (7) and (8), the function  $\Psi(m)$  is replaced by  $\tilde{\Psi}(m)$ . The latter is obtained from expression (8) by replacing the profile  $\rho(m)$  and mass  $M(m)$  by the general profile  $\rho(m) + \rho_G(m)$  and the total mass  $M(m) + \tilde{M}_G(m)$  of BM and DM. Then, the sought expressions  $\tilde{W}_L = \tilde{W}_L(1)$  and  $\tilde{T}_L = \tilde{T}_L(1)$  are represented as (b)

$$b) \quad \tilde{W}_L = -2\tilde{W}_0 J_0, \quad \tilde{T}_L = \tilde{W}_0 J_1, \quad \tilde{W}_0 = \frac{\pi G a c^2}{2} \tilde{\Psi}(1), \quad (15)$$

where

$$\begin{aligned} \tilde{\Psi}(m) &= \sum_{n=1}^4 \tilde{\psi}_n(m), \quad \tilde{\psi}_2(m) = \int_0^{m^2} \rho(m) \tilde{M}_G(m) dm^2, \\ \tilde{\psi}_3(m) &= \int_0^{m^2} \rho_G(m) M(m) dm^2, \quad \tilde{\psi}_4(m) = \int_0^{m^2} \rho_G(m) \tilde{M}_G(m) dm^2, \end{aligned}$$

Here,  $\tilde{\psi}_1(m) \equiv \Psi(m)$ , the function  $\Psi(m)$  is defined by equality (11) and is the mass  $\tilde{M}_G(m)$  of an inhomogeneous spheroid  $a > b = c$  consisting of DM with a profile  $\rho_G(m)$  and is calculated by the formula

$$\tilde{M}_G(m) = 4\pi a c^2 \int_0^m m^2 \rho_G(m) dm = 4\pi K \bar{a}^3 \left( \ln \bar{g} - \frac{\bar{g} - 1}{\bar{g}} \right), \quad \bar{g} = 1 + \bar{\xi} m, \quad \bar{\xi} = \frac{\sqrt[3]{a c^2}}{\bar{a}} \quad (16)$$

Here,  $\bar{a}$  is the scale of the galaxy. After calculating the functions  $\tilde{\psi}_k(m)$ , ( $k = 2, 3, 4$ ), we obtain for  $\tilde{\Psi}(m)$  the following expression:

$$\tilde{\Psi}(m) = 4\pi a c^2 \left[ \rho_0^2 f_1(m) + 2K\rho_0 f_2(m) + K^2 f_3(m) \right], \quad (17)$$

where the function  $f_1(m)$  is defined by equality (13),

$$f_2(m) = \frac{1}{\beta \bar{\xi}^3 \bar{g}} \left[ \frac{\bar{g} - 1}{\sqrt{1 + h^2}} - \frac{\bar{g} - 1}{h} \ln \varphi_1(m) + \frac{\bar{\xi} \bar{g}}{2\sqrt{\beta + \bar{\xi}^2}} \ln \frac{\varphi_2(m)}{\varphi_2(0)} - \frac{\bar{g}}{\sqrt{1 + h^2}} \ln \bar{g} \right], \quad (18)$$

$$f_3(m) = \frac{1}{\bar{\xi}^5} \left( 1 - \frac{1}{\bar{g}^2} - \frac{2 \ln \bar{g}}{\bar{g}} \right), \quad \varphi_2(m) = \frac{\beta m - \bar{\xi} + \sqrt{1 + h^2} \sqrt{\bar{\xi}^2 + \beta}}{-\beta m + \bar{\xi} + \sqrt{1 + h^2} \sqrt{\bar{\xi}^2 + \beta}}, \quad (19)$$

where the function  $\varphi_1(m)$  and parameter  $h$  are defined by equality (12) and  $\bar{g}$  and  $\bar{\xi}$ , by equality (16).

It is obvious that expression (15) for energies  $\tilde{W}_L$  and  $\tilde{T}_L$  coincides with expression (14) for energies  $W_L$  and  $T_L$  if the LP EG consists only of BM. Indeed, in this case  $\rho_G(m) = 0$ , or  $\tilde{\psi}_k(m) = 0$ , ( $k = 2, 3, 4$ ). Therefore,  $\tilde{\Psi}(m) = \Psi(m)$  and  $\tilde{W}_0 = W_0$ , or, which is the same,  $\tilde{W}_L = W_L$  and  $\tilde{T}_L = T_L$ .

It is important to note that the gravitational energy and rotational kinetic energy of the EG corresponding to variant (2) of model 5 are also determined by equality (15). In addition, from this equality, expressions for these energies corresponding to variant (1) of model 5 can be easily obtained by replacing  $a, b = c$  with  $\tilde{a}, \tilde{b} = \tilde{c}$  and  $\bar{a}$  with  $r_s$ .

#### 4. POTENTIAL ENERGIES AND ROTATIONAL KINETIC ENERGIES OF INHOMOGENEOUS SPHEROIDAL AND SPHERICAL LAYERS

First, let us determine the potential energy  $W_G$  and rotational kinetic energy  $T_G$  of an inhomogeneous spheroidal layer consisting of DM with a profile  $\rho_G(\mu)$  and mass  $M_G(\mu)$ . We assume that the inner and outer boundaries of this layer coincide with the spheroids with semiaxes  $\mu \tilde{a}$ ,  $\mu \tilde{b} = \mu \tilde{c}$  and  $\tilde{a}, \tilde{b} = \tilde{c}$ , respectively. Then, the energies  $W_G$  and  $T_G$  will be determined by the formulas [11]

$$W_G = -\pi G \tilde{a} \tilde{c}^2 J_0 \Psi_G(\mu), \quad T_G = \frac{\pi G \tilde{a} \tilde{c}^2}{2} J_1 \Psi_G(\mu), \quad (20)$$

where the coefficients  $J_0$  and  $J_1$  are given above and the function  $\Psi_G(\mu)$  is defined by the formula

$$\Psi_G(\mu) = \int_{\mu^2}^1 \rho_G(\mu) M_G(\mu) d\mu^2, \quad M_G(\mu) = 4\pi \tilde{a} \tilde{c}^2 \int_{\mu}^1 \mu^2 \rho_G(\mu) d\mu$$

After calculating the integrals for the mass of the homeoid  $M_G(\mu)$  and the function  $\Psi_G(\mu)$ , we obtain the following expressions:

$$M_G(\mu) = 4\pi K r_s^3 \left[ \ln \frac{1 + \xi}{1 + \xi \mu} - \frac{\xi(1 - \mu)}{(1 + \xi)(1 + \xi \mu)} \right], \quad \Psi_G(\mu) = \frac{8\pi K^2 r_s^3}{\xi^2} H_G(\mu), \quad (21)$$

where

$$H_G(\mu) = \frac{1}{1 + \xi\mu} \left[ \frac{\xi(1 - \mu)(2 + 3\xi + \xi\mu + 2\xi^2\mu)}{2(1 + \xi)^2(1 + \xi\mu)} - \ln \frac{1 + \xi}{1 + \xi\mu} \right], \quad \xi = \frac{\sqrt[3]{a\tilde{c}^2}}{r_s}$$

By virtue of (21), the energies  $W_G$  and  $T_G$  can be rewritten in a more compact form:

$$W_G = -2\tilde{W}_0 J_0 H_G(\mu), \quad T_G = \tilde{W}_0 J_1 H_G(\mu), \quad \tilde{W}_0 = 4\pi^2 G K^2 \xi r_s^6 \quad (22)$$

Now let us calculate the potential energy  $W_S(r)$  of an inhomogeneous spherical layer (model 3) with a profile  $\rho_S(r)$  defined by equality (3), using the formula from [11]:

$$W_S(r) = -2\pi \int_a^r u^2 \rho_S(u) U_S(u) du, \quad U_S(r) = \frac{4\pi G}{r} \int_a^r u^2 \rho_S(u) du + 4\pi G \int_r^{r_s} u \rho_S(u) du$$

Next, we will consider that the radius of the inner circle of this layer is equal to  $a$ : the semi-major axis of the midrange EG, and the radius of the outer circle is equal to the scale radius  $r_s$  of the EG. Then the internal potential  $U_S$  and mass  $M_S(r)$  of the intermediate spherical layer will be equal to

$$U_S(r) = \frac{4\pi G K r_s^3}{r} \left( \ln \frac{r + r_s}{a + r_s} - \frac{r}{2r_s} + \frac{a}{a + r_s} \right), \quad M_S(r) = 4\pi K r_s^3 \left( \ln \frac{r + r_s}{a + r_s} - \frac{r_s}{a + r_s} + \frac{r_s}{r + r_s} \right)$$

respectively, and, at  $r = r_s$ , we obtain the total mass of this layer.

Let us substitute the expressions for the potential  $U_S(r)$  of this layer and its density  $\rho_S(r)$  from (3) into the expression for  $W_S(r)$ . Then, for the total gravitational energy  $W_S = W_S(r_s)$  of an inhomogeneous spherical layer, we obtain

$$W_S = -8\pi^2 G K^2 r_s^5 \psi_1(r_s), \quad \psi_1(r_s) = \ln \frac{a + r_s}{2r_s} + \frac{(r_s - a)(5a + 3r_s)}{4(a + r_s)^2} \quad (23)$$

The rotational kinetic energy  $T_S(r)$  of a spherical layer can be represented as the difference between the energies  $\tilde{E}_1$  and  $\tilde{E}_2$  of two inhomogeneous balls of radii  $r_s$  and  $a$  with the same density  $\rho_S(r)$ , respectively. To determine the rotational energies  $\tilde{E}_1$  and  $\tilde{E}_2$ , we can use the formula

$$\tilde{E}_k = \frac{1}{2} \tilde{J}_k \Omega_k^2, \quad \Omega_k^2 = \frac{GM_k}{r_k^3}, \quad (k = 1, 2),$$

where  $M_1 = M_H$  and  $M_2 = M^*$  are the masses of the halo and the LP EG,  $r_1 = r_s$  and  $r_2 = a$  are the scale radius of the EG and the semi-major axis of its luminous part, and are the moments of inertia of inhomogeneous balls with density  $\rho_S(r)$  and radii  $r_s$  and relative to the axis of rotation. To determine  $\tilde{J}_1$  and  $\tilde{J}_2$ , we use the general formula for the moment of inertia:

$$J(r) = \frac{8\pi}{3} \int_0^r u^4 \rho_S(u) du = \frac{8\pi}{3} K r_s^3 \left[ 3r_s^2 \ln \frac{r + r_s}{r_s} + \frac{r(r^2 - 3rr_s - 6r_s^2)}{2(r + r_s)} \right],$$

setting  $r = r_s$  and  $r = a$  and find, respectively,  $J_1$  and  $J_2$  and, hence,  $\tilde{E}_1$  and  $\tilde{E}_2$ . Therefore, the sought for expression for the rotational energy  $T_S$  of a spherical layer is obtained in the form

$$T_S = \tilde{E}_1 - \tilde{E}_2, \quad \tilde{E}_1 = \frac{4\pi}{3} G M_H K r_s^2 (3 \ln 2 - 2), \quad (24)$$

$$\tilde{E}_2 = \frac{4\pi GM^*}{3a^3} K r_s^3 \left[ 3r_s^2 \ln \frac{a+r_s}{r_s} + \frac{a(a^2 - 3ar_s - 6r_s^2)}{2(a+r_s)} \right]$$

## 5. TOTAL POTENTIAL AND KINETIC ENERGY OF A LAYERED INHOMOGENEOUS ELLIPTICAL GALAXY

The total gravitational energy  $W$  and rotational kinetic energy  $T$  of a dynamical system according to variant (a) of models 3 and 4 will be presented in the form of sums  $W = W_1 + W_2 + W_3$  and  $T = T_1 + T_2$ , and, according to variant (b), in the form of sums  $\tilde{W} = \tilde{W}_1 + W_2 + \tilde{W}_3$  and  $T = \tilde{T}_1 + T_2$ , respectively. Here,  $W_1, T_1$  and  $\tilde{W}_1, \tilde{T}_1$  are the gravitational energies and rotational energies of the LP EG in variants (a) and (b), respectively. In addition,  $W_2$  and  $T_2$  are the gravitational energy and rotational energy of the inhomogeneous spherical (spheroidal) layer  $W_3$  and  $\tilde{W}_3$  are the mutual gravitational energies of the LP EG and this layer according to variants (a) and (b), respectively. According to variant (a) of both models, the energies  $W_1 = W_L$  and  $T_1 = T_L$  of the LP EG were determined by equalities (14), respectively. In this case, the energies  $W_2 = W_G$  and  $T_2 = T_G$  of the homeoid are determined by equality (22) and the energies  $W_2 = W_S$  and  $T_2 = T_S$  of the spherical layer are determined by equalities (23) and (24), respectively.

The energies  $\tilde{W}_1 = \tilde{W}_L$  and  $\tilde{T}_1 = \tilde{T}_L$  corresponding to variant (b) of models 3 and 4 are calculated differently, namely, by formula (15), the explicit form of which is given at the end of Section 5. Then, the gravitational energies  $W_3$  and  $\tilde{W}_3$  according to models 3 and 4 need to be calculated. First, we calculate them according to variants (a) and (b) of model 4 by the formula [11]:

$$a) \quad W_3 = -U_G(\nu)M(1), \quad b) \quad \tilde{W}_3 = -U_G(\nu) [M(1) + \tilde{M}_G(1)]$$

respectively. The total masses  $M(1)$  and  $\tilde{M}_G(1)$  of both BM and DM in the LP EG have been determined in Section 3, and the potential  $U_G(\nu)$  of an inner point of the homeoid cavity is equal to [11]

$$U_G(\nu) = \pi G \tilde{a} \tilde{c}^2 J_0 \int_{\nu^2}^1 \rho_G(m) dm^2 = \frac{2\pi G K \tilde{a} \tilde{c}^2 J_0}{\xi(1+\xi)} \frac{1-\nu}{1+\nu\xi} = \text{const.}$$

Here, an internal point of the homeoid interior means any passively gravitating body (e.g., a star) located in the LP EG. Next, setting  $\nu = \mu$  in the expression for the potential  $U_G(\nu)$ , the sought-for energies  $W_3$  and  $\tilde{W}_3$  can be represented as

$$a) \quad W_3 = -W_0 \rho_0 g_1(1), \quad b) \quad \tilde{W}_3 = -W_0 [\rho_0 g_1(1) + K g_2(1)], \quad (25)$$

where the function  $g_1(m)$  is defined by equality (12). In addition,

$$W_0 = \frac{8\pi^2 G K \tilde{a} \tilde{c}^2 a c^2 J_0}{\xi(1+\xi)} \frac{\mu(1-\mu)}{1+\mu\xi}, \quad g_2(m) = \frac{1}{\xi^3} \left[ \ln(1+\xi m) - \frac{\xi m}{1+\xi m} \right] \quad (26)$$

Now, let us calculate the mutual potential energies  $W_3$  and  $\tilde{W}_3$  according to variants (a) and (b) of model 3. By analogy with formula (25), we obtain

$$a) \quad W_3 = -U_S M(1), \quad b) \quad \tilde{W}_3 = -U_S [M(1) + \tilde{M}_G(1)],$$

where  $U_S$  is the potential at the inner point of the cavity of the spherical layer, equal to [20]

$$U_S = 4\pi G \int_a^{r_s} u \rho_S(u) du = 2\pi G K r_s^2 \frac{r_s - a}{r_s + a} = \text{const},$$

Then

$$a) W_3 = -W_0 \rho_0 g_1(1), \quad b) \widetilde{W}_3 = -W_0 [\rho_0 g_1(1) + K g_2(1)], \quad W_0 = 8\pi^2 G K r_s^2 a c^2 \frac{r_s - a}{r_s + a} \quad (27)$$

The function  $g_2(1)$  is defined above.

In conclusion, let us present a list of the total gravitational,  $W$ , and kinetic,  $T$ , energies calculated according to models 3, 4, and 5 in the following order.

Model 3: (a)  $W = W_L + W_S + W_3$  and  $T = T_L + T_S$ ; (b)  $\widetilde{W} = \widetilde{W}_L + W_S + \widetilde{W}_3$ , and  $\widetilde{T} = \widetilde{T}_L + T_S$ . The energies  $W_L$  and  $T_L$  are determined by equality (14);  $W_L$  and  $T_L$ , by formulas (23) and (24), respectively;  $\widetilde{W}_L$  and  $\widetilde{T}_L$ , by formula (15); and the energies  $W_3$  and  $\widetilde{W}_3$ , by equality (27).

Model 4: (a)  $W = W_L + W_G + W_3$  and  $T = T_L + T_G$ ; (b)  $\widetilde{W} = \widetilde{W}_L + W_G + \widetilde{W}_3$ , and  $\widetilde{T} = \widetilde{T}_L + T_G$ . The energies  $W_G$  and  $T_G$  are determined by formula (22), and the energies  $W_3$  and  $\widetilde{W}_3$ , by equality (25).

Model 5: (2)  $W = \widetilde{W}_L$  and  $T = \widetilde{T}_L$ . Here, the energies  $\widetilde{W}_L$  and  $\widetilde{T}_L$  are determined by formula (15). In this case, to obtain the necessary expressions corresponding to variant (1) of model 5, it is sufficient to replace the semiaxes  $a$  and  $b = c$  in expression (15) with  $\tilde{a}$  and  $\tilde{b} = \tilde{c}$ , respectively.

## 6. VELOCITY DISPERSION OF A LAYERED INHOMOGENEOUS ELLIPTICAL GALAXY

Let us consider the ratio of the total rotational energy  $T_{rot}$  of a layered inhomogeneous ellipsoid with semiaxes  $a > b > c$  and density  $\rho(m)$  to the absolute value of its total gravitational energy  $W$ . According to formula (7),  $W = W(1) = -\pi G a b c J_0 \Psi(1)$ ; for  $T_{rot} = T_1$ , we find the value of  $t_z$  [11]:

$$t_z = \frac{T_{rot}}{|W|} = \frac{1}{2} \left( 1 - \frac{3c^2 K_0}{J_0} \right) = \frac{J_1}{2J_0}, \quad (28)$$

which exactly coincides with the cognominal relation for classical homogeneous equilibrium figures (Maclaurin spheroids:  $a = b \geq c$ , or Jacobi ellipsoids:  $a \geq b \geq c$ ) and is a function of only the eccentricities of the layer [11].

It is important to note that the ratio  $t_z$  plays a key role in establishing equilibrium and stability of an axisymmetric dynamical system (e.g., homogeneous Maclaurin spheroids). A criterion for the stability of such a system, according to the Peebles–Ostriker hypothesis [21, 22], is the fulfillment of the inequality

$$t_z < t_{crit} \approx 0.14 \pm 0.03.$$

Next, the ratio  $t_z$  can be related with the rotational velocity  $v_{rot}(R)$  and velocity dispersion  $\sigma_s(R)$  observed in elliptical galaxies at a distance [11]:

$$\frac{v_{rot}(R)}{\sigma_s(R)} = \sqrt{\frac{t_z}{0.5 - t_z}}$$

The ratio  $v_{rot}(R)/\sigma_s(R)$  calculated by this formula should be compared with observations, which allows one to draw some conclusions about the dynamic state of this galaxy. In addition, calculating the linear rotation velocity  $v_{rot}(R) = \Omega R$ , we find the velocity spatial dispersion  $\sigma_s(R)$  as a function on the distance  $R$  from the center of the galaxy [11]:

$$\sigma_s(R) = \Omega R \cdot \sqrt{\frac{0.5 - t_z}{t_z}}, \quad \Omega = \sqrt{\frac{GM}{a^3}} \quad (29)$$

where  $\Omega$  is the angular rotation velocity of the galaxy. Since the centripetal and gravitational forces acting on a point located at a distance  $R$  from the center balance each other, we obtain:

$$\frac{V^2}{R} = \frac{GM}{R^2}, \quad V_{max} = \sqrt{\frac{GM}{a}} = \Omega a \quad (30)$$

Let us present other methods for determining the velocity spatial dispersion. Knowing the density distribution law  $\rho(r)$ , one can determine the velocity spatial dispersion  $\sigma_s(R)$  at a distance  $R$  from the center of the galaxy [23, 24]:

$$\sigma_s^2(R) = \frac{G}{\rho(R)} \int_R^\infty \frac{\rho(r) M(r)}{r^2} dr, \quad (31)$$

where the mass  $M(r)$  of the intermediate ball and the rotation velocity  $V_c(r)$  are determined by the expressions

$$M(r) = 4\pi \int_0^r u^2 \rho(u) du, \quad V_c(r) = \sqrt{\frac{GM(r)}{r}} \quad (32)$$

respectively.

Another expression for determining the velocity spatial dispersion  $\sigma_s(R)$  at a distance  $R$  from the center of a spherically symmetric galaxy is given in [25]:

$$\sigma_s^2(R) = \frac{2G}{I(R)} \int_R^\infty \frac{\sqrt{r^2 - R^2}}{r^2} \rho(r) M(r) dr, \quad I(R) = \frac{2}{\Upsilon} \int_R^\infty \frac{\rho(r) r dr}{\sqrt{r^2 - R^2}}, \quad (33)$$

where  $I(R)$  is the surface brightness [26] and  $\Upsilon$  is the mass–luminosity ratio.

It is important to note that, for an EG with a homothetic (astrophysical) density distribution (1), the velocity spatial dispersion  $\sigma_s(R)$  cannot be determined by formulas (31) and (33). To apply these formulas, we set

$$m = \frac{r}{q}, \quad m_{eff} = \frac{R_{eff}}{q}, \quad q = \sqrt[3]{ac^2}, \quad w = \frac{r}{q} \sqrt{\beta} = m \sqrt{\beta}, \quad W = \frac{R \sqrt{\beta}}{q}. \quad (34)$$

Then, expression (31), by virtue of expressions (1) and (15) for  $\rho(m)$  and  $M(m)$  takes the form

$$\sigma_s^2(R) = \frac{4\pi G \rho_0 \tilde{R}^3}{q \beta} H(R), \quad \tilde{R} = \sqrt{q^2 + R^2 \beta} \quad (35)$$

where

$$H(R) = \frac{4Q^4}{Q^4 - 1} \ln Q - 2 \ln(Q^2 + 1) + \frac{2Q^2}{(Q^2 + 1)^2}, \quad Q = \frac{R \sqrt{\beta} + \tilde{R}}{q}$$

In accordance with substitution (34), from (33), we find  $I(R)$  and  $\sigma_s^2(R)$

$$I(R) = \frac{2\rho_0 q^3}{\Upsilon \sqrt{\beta} \tilde{R}^2}, \quad \sigma_s^2(R) = \frac{4\pi G \rho_0 \tilde{R}^2 \Upsilon}{\beta} [K_1(R) + K_2(R)], \quad (36)$$

where

$$K_1(R) = \int_W^\infty \frac{\sqrt{w^2 - W^2}}{w^2(1+w^2)^{3/2}} \ln(w + \sqrt{1+w^2}) dw, \quad K_2(R) = -\frac{\pi (\tilde{R} - R\sqrt{\beta})^2}{4\tilde{R}^2}$$

Here, the integral  $K_1(R)$  cannot be calculated explicitly, but can be calculated numerically.

Formulas (35) and (36) of the velocity dispersion correspond to variant (a) of models 3 and 4. In this case, the LP SP does not contain DM; therefore, the total velocity dispersion is  $\sigma(R) \equiv \sigma_s(R)$ .

Now let us determine the velocity dispersion  $\sigma(R)$  according to variant (b) of models 3 and 4 and according to variants (1) and (2) of model 5. In this case, the LP SP contains DM and the velocity dispersion can be represented as the sum  $\sigma^2(R) = \sigma_s^2(R) + \sigma_d^2(R)$ , where  $\sigma_s^2(R)$  is the BM component of the velocity dispersion and  $\sigma_d^2(R)$  is the DM component. In this case, it suffices to calculate  $\sigma_d^2(R)$ , since  $\sigma_s^2(R)$  is defined by formulas (35) and (36). As for a formula similar to (29), it suffices to calculate the ratio  $t_z$ . According to variant (2) of model 5, this ratio will be equal to  $t_z = \tilde{T}_L/|\tilde{W}_L|$ . Here,  $\tilde{W}_L$  and  $\tilde{T}_L$  are determined by equality (15). This ratio is easily determined according to variant (1) of model 5 with the help of an appropriate substitution, which was mentioned above.

Let us proceed to calculating  $\sigma_d^2(R)$  first by formula(31)

$$\sigma_d^2(R) = \frac{G}{\tilde{q} \rho_G(\tilde{\mu})} \int_{\tilde{\mu}}^\infty \frac{\rho_G(\mu) M_G(\mu)}{\mu^2} d\mu, \quad \left( r = \tilde{q} \mu, \quad \tilde{\mu} = \frac{R}{\tilde{q}}, \quad \tilde{q} = \sqrt[3]{\tilde{a}c^2}, \quad \tilde{\mu}\xi = \frac{R}{r_s} \right)$$

Using expression (4) for the density  $\rho_G(\mu)$  and mass  $M_G(\mu)$  given in Section 4, we find

$$\sigma_d^2(R) = B_0 \left[ B_1(R) \ln \frac{R}{r_s} + B_2(R) \ln \frac{r_s + R}{r_s} - 6B_1(R) \text{dilog} \left( \frac{r_s + R}{r_s} \right) + B_3(R) \right], \quad (37)$$

where

$$B_1(R) = -\frac{R}{r_s^4} (r_s + R)^2, \quad B_2(R) = \frac{r_s + R}{r_s^4 R} (r_s^3 - 3Rr_s^2 + R^2r_s + 7R^3),$$

$$B_3(R) = \frac{1}{r_s^4} \left[ \pi^2 R (r_s + R)^2 - r_s^3 - 9Rr_s^2 - 7R^2r_s \right], \quad B_0 = \frac{2\pi G K r_s \tilde{q}^2}{\xi^2},$$

and

$$\text{dilog}(s) = \int_1^s \frac{\ln t}{1-t} dt, \quad \text{dilog}(1) = 0, \quad \text{dilog}(0) = \frac{\pi^2}{6}, \quad \text{dilog}\left(\frac{1}{2}\right) = \frac{\pi^2}{12} - \frac{\ln^2 2}{2}$$

Next, using the second formula (33), we determine the surface brightness  $I(R)$ :

$$I(R) = \frac{K \tilde{q} r_s^2}{\xi \Upsilon \tilde{R}^3} \left( -\tilde{R} + r_s \ln \frac{r_s + \tilde{R}}{r_s - \tilde{R}} \right), \quad \tilde{R} = \sqrt{r_s^2 - R^2}$$

Then, in the first formula (33), we consider the expressions for  $\rho_G(\mu)$  and masses  $M_G(\mu)$ . Then we determine the DM component of the velocity dispersion  $\sigma_d^2(R)$ :

$$\sigma_d^2(R) = \frac{2G}{I(R)} \int_{\bar{\mu}}^{\infty} \frac{\sqrt{\mu^2 - \bar{\mu}^2}}{\mu^2} \rho_G(\mu) M_G(\mu) d\mu = \psi_0(R) [\psi_1(R) - \xi \psi_2(R)], \quad (38)$$

where

$$\begin{aligned} \psi_0(R) &= \frac{8\pi G K r_s \Upsilon \bar{R}^3}{\tilde{q} \left( -\bar{R} + r_s \ln \frac{r_s + \bar{R}}{r_s - \bar{R}} \right)}, & \psi_1(R) &= \int_{\bar{\mu}}^{\infty} \frac{\sqrt{\mu^2 - \bar{\mu}^2}}{\mu^3} \frac{\ln(1 + \xi\mu)}{(1 + \xi\mu)^2} d\mu, \\ \psi_2(R) &= \frac{1}{4\bar{R}^3 r_s} \left[ (2r_s^4 - 9r_s^2 R^2 + 6R^4) \ln \frac{r_s + \bar{R}}{r_s - \bar{R}} - 2\bar{R} (5r_s^3 - 6R^2 r_s - 3\pi R \bar{R}^2) \right] \end{aligned} \quad (39)$$

The integral  $\psi_1(R)$  cannot be calculated analytically and is calculated numerically. The values of the velocity spatial dispersion  $\sigma(R)$  at a distance of the effective radius  $R_{eff}$  for ten EGs [27] are given in section Examples.

## 7. NEW DETERMINATION OF THE DENSITY VALUES IN THE CENTER OF EG, ITS SCALE RADIUS, AND PARAMETER $\beta_{eff}$

Assuming the value of the spatial velocity dispersion  $\sigma_{eff}$  at a distance of the effective radius  $R_{eff}$  to be known (see Section 6), we can determine the density  $\rho_0$  at the center of the EG. Using the astrophysical law of density distribution  $\rho(r)$  for the EG, according to formula (35), we obtain

$$\rho_0 = \frac{\sigma_s^2(R_{eff}) \beta_{eff} q}{4\pi G \sqrt{(q^2 + R_{eff}^2 \beta_{eff})^3} H(R_{eff})}, \quad \left( \rho_{crit} = \frac{3H^2}{8\pi G} \approx 1.37 \cdot 10^{-7} \right) \quad (40)$$

In the parentheses, the critical density value in the units  $M_{\odot}/pc^3$  is given.

If the velocity spatial dispersion  $\sigma_s(R)$  is determined by equality (36), then  $\rho_0$  is expressed as follows:

$$\rho_0 = \frac{\sigma_s^2(R_{eff}) \beta_{eff}}{4\pi G (q^2 + R_{eff}^2 \beta_{eff}) \Upsilon [K_1(R_{eff}) + K_2(R_{eff})]} \quad (41)$$

The values of  $\sigma_s(R_{eff})$ , as well as the density values  $\rho_0$  calculated from them at the center of the galaxy, are given in Section 8.

The value  $\beta_{eff}$  of the parameter  $\beta$  in equalities (40) and (41), corresponding to the effective radius, can be determined, e.g., from the Hubble law:

$$I_{eff} = \frac{I_0}{1 + \beta_{eff} m_{eff}^2}, \quad \beta_{eff} = \frac{1}{m_{eff}^2} \left( \frac{I_0}{I_{eff}} - 1 \right), \quad m_{eff} = \frac{R_{eff}}{\sqrt[3]{a c^2}}, \quad (42)$$

in which the central and effective surface brightness  $I_0$  and  $I_{eff}$  are assumed to be known. Another way to determine these parameters is related to the following formula:

$$L(m) = 4aqE(e) \int_0^m m I(m) dm = \frac{2qaE(e)I_0}{\beta} \ln(1 + \beta m^2), \quad e = \frac{\sqrt{a^2 - c^2}}{a},$$

in which the expression (1) for the surface brightness  $I(m)$  is taken into account. Here,  $L(m)$  is the total surface brightness of the intermediate ellipse with semiaxes  $ma$  and  $mc$  and  $E(e)$  is the complete elliptic integral of the second kind. Hence, it is easy to determine the total,  $L_T = L(m = 1)$ , and effective,  $L_{eff} = L(m_{eff})$ , luminosity:

$$L_T = 2qaE(e)I_0 \frac{\ln(1 + \beta_T)}{\beta_T}, \quad L_{eff} = 2qaE(e)I_0 \frac{\ln(1 + \beta_{eff}m_{eff}^2)}{\beta_{eff}},$$

where  $\beta_T$  is the value of the parameter  $\beta$  corresponding to  $m = 1$ . In addition, by definition,  $L_{eff} = L_T/2$ , i.e.,

$$\frac{\ln(1 + \beta_{eff}m_{eff}^2)}{\beta_{eff}} = \frac{\ln(1 + \beta_T)}{2\beta_T} = \frac{L_T}{4qaE(e)I_0}$$

Hence, assuming  $I_0$  and  $L_T$  to be known, we determine the parameters  $\beta_T$  and  $\beta_{eff}$ . In addition, the parameter  $\beta_T$  can be calculated differently. For example, if the stellar mass  $M^*$  of the galaxy, its density  $\rho_0$  at the center, and the semiaxes  $a, b = c$  are known, then, from formula (15), for the mass  $M(m)$ , we determine the value of the parameter  $\beta_T$  as a solution of the equation

$$\frac{g_1(\beta_T)}{\beta_T \sqrt{\beta_T}} = \frac{M^*}{4\pi\rho_0ac^2}, \quad M^* \equiv M(m = 1),$$

where the function  $g_1(\beta_T) \equiv g_1(m = 1)$  is defined by equality (12).

The scale radius  $r_s$  of the EG can be determined from the condition

$$M_h = M^* + M_G + M_S(r_s) = M_{bm} + M_S(r_s) = M_{bm} + M_{dm}$$

Here, the stellar mass  $M^*$ , the mass of the gas  $M_G$  (equal to the sum of the BM  $M_{bm}$ ), and the mass of the galactic halo  $M_h$  are assumed to be known. In this case, the total mass  $M_{dm} = M_S(r_s)$  of an inhomogeneous layer (model 3) consisting of DM is determined by equality (25) at  $r = r_s$ . In expression (25), the normalization coefficient  $K$  is replaced by its approximate value

$$K \approx \left(\frac{r_s}{10}\right)^{-2/3}, \quad (43)$$

where  $r_s$  is expressed in pc [13]. Then, assuming the semi-major axis  $a$  of the EG to be known, we represent the mass  $M_{dm} = M_S(r_s)$  as a function of only the scale radius and substitute it into the expression for  $M_h$ . This gives us an equation for determining  $r_s$ :

$$M_h - M_{bm} = B(r_s) r_s^{7/3}, \quad B(r_s) = 4\pi \sqrt[3]{100} \left[ \ln \frac{2r_s}{a + r_s} - \frac{r_s - a}{2(a + r_s)} \right] \quad (44)$$

The value of  $r_s$  determined from Eq. (44) corresponds to model 3.

In model 4, the value of  $r_s$  is found from the expression for  $M_h$  when  $M_{dm} = M_G(\mu)$ :

$$M_h - M_{bm} = A(r_s) r_s^{7/3}, \quad A(r_s) = 4\pi \sqrt[3]{100} \left[ \ln \frac{1 + \xi}{1 + \xi\mu} - \frac{\xi(1 - \mu)}{(1 + \xi)(1 + \xi\mu)} \right] \quad (45)$$

Here

$$\xi = \frac{\sqrt[3]{a\tilde{c}^2}}{r_s} = \frac{\sqrt[3]{ac^2}}{a}, \quad \mu = \frac{a}{r_s}$$

Now, assuming to be known  $r_s$  (in pc), we determine the normalization coefficient  $K$ , expressed in solar masses per cubic parsec, using formula (43). The values of the scale radii  $r_s$  thus found and the coefficient  $K$  according to formula (43) in models 3 and 4 for ten EGs are given in section Examples.

Thus, the formulas given in this section make it possible to determine the values of such key parameters of an EG as the density at the center of the galaxy and its scale radius  $r_s$ . The values of  $\rho_0$  and  $r_s$  of specific EGs presented in the database or in works by other authors are not given. In [28], the scale radius  $\rho_s$  of an EG is understood as the radius of its disk: the radial scale. In this case, the angular values of the minor,  $c$ , and major,  $a$ , axes are determined from observations. To determine these parameters, according to the above formulas, it is necessary to know the stellar (or baryon) mass and halo mass of the galaxy, its effective radius, and the velocity dispersion calculated at the distance of the effective radius of the EG. These data are available in the database and publications.

## 8. EXAMPLES

We took ten elliptical galaxies, for example, with parameters necessary for the calculations given in Tables 1–4.

The quantities  $D_{25}$  and  $R_{25}$ , which are measured in the photometric  $B$  band up to a distance with a limiting brightness of 25 magnitudes per square arcsecond, as well as the heliocentric distance  $D$ , are taken from the HyperLeda database (leda.univ-lyon1.fr).

Using the values of  $D_{25}$  and  $R_{25}$  given in [7], we determine the apparent values of the major and minor semi-axes  $a$  and  $c$  of the EG, which is regarded as an elongated spheroid with semi-axes  $a > b = c$ . For such galaxies and galaxies that have the shape of a triaxial ellipsoid, the rotation axis does not coincide with the apparent minor axis and the isophotes will be misaligned. The variant of an EG in the form of an oblate spheroid—a more convenient and simpler variant, for which the visible axis coincides with the axis of rotation and the isophote alignment is not violated—is not considered.

Table 1 presents the values of the heliocentric distance  $D$  (in Mpc), BM  $M_{bm}$ , DM mass  $M_{dm}$ , and halo mass  $M_h$ , scale radius  $r_s$  (in kpc), as well as the normalization coefficient  $K$  (in  $10^{-3}M_\odot/\text{pc}^3$ ), determined using models 3 and 4. The masses  $M_{bm}$ ,  $M_{dm}$ , and  $M_h$  are expressed in  $10^{12}$  solar masses.

Table 2 presents the values of the EG semi-major,  $a$ , and semi-minor,  $c$  axes (in kpc), parameters  $\beta_T$ ,  $m_{eff}$ , and  $\beta_{eff}$ , as well as the effective radius  $R_{eff}$  (in kpc) from [29, 30]. The parameters  $\beta_T$  and  $\beta_{eff}$  correspond to  $m = 1$  and  $m = m_{eff}$ , and the latter corresponds to  $R_{eff}$ .

The stellar masses  $M^*$ , BMs  $M_{bm}$ , and halo masses  $M_h$  of galaxies are given in [29–32]. For  $M_h$ , it is possible to use its approximate dependence on the stellar mass of the galaxy,  $M^*$ ,  $M_h = (56 + \Delta M) \cdot M^*$ , where  $\Delta M$  varies from  $\Delta M = -10$  to 16 [33]. In this case, the BMs  $M_{bm}$  of EGs can be determined from the estimates of the masses of neutral hydrogen  $M_{HI}$  and molecular,  $M_{H_2}$ , and central molecular,  $M_{H_{1c}}$ , gases given in [33, 34]. In addition, in [35], an approximate formula for determining the BM of an EG is given in the form  $M_{bm}/M_h \approx \Omega_b/\Omega_m \approx 0.17$ .

Table 3 presents the values of the central velocity dispersion  $\sigma_0$  and the spatial velocity dispersion  $\sigma(R_{eff})$  (in km/s) at the distance of the effective radius, as well as the density

at the center of the EG. Column 1 corresponds to the value of  $\sigma_s(R_{eff})$  given in [29], and column 2, to its value calculated by formula (29). Columns 3 and 4 in the first row give the values of  $\sigma_s(R_{eff})$  calculated by formulas (35) and (36), i.e., without considering the DM component. This corresponds to variant (a): the LP EG does not contain DM. The second row of these columns gives its values with allowance for the DM components:  $\sigma(R_{eff}) = \sqrt{\sigma_s^2(R_{eff}) + \sigma_d^2(R_{eff})}$ , where  $\sigma_d^2(R_{eff})$  is calculated using formulas (37) and (38). Columns 5 and 6 correspond to the values of  $\rho_0$ , and column 7, to the average  $\rho_0$  expressed in  $M_\odot/pc^3$ .

The discrepancies in the values of  $\sigma_s(R_{eff})$  are due to the fact that, in this work, the density is distributed in accordance with astrophysical law (6).

Table 4 presents the values of the total gravitational and rotational kinetic energies (in Joules) of ten elliptical galaxies calculated using models 3, 4, and 5. For model 5, the values of these quantities in variants (1) and (2) are given, and, for models 3 and 4, in two variants: (a) the main part of DM is contained outside the luminous part of the EG and (b) the fractions of dark and baryonic matter in the central regions of the galaxy are comparable.

**Table 1.** Values of the heliocentric distance  $D$  and baryon mass  $M_{bm}$  of EGs

NGC	$D, Mpc$	$M_{bm}$	Models	$M_{dm}$	$M_h$	$r_s$	$K$
4365 E3	23.3	0.6842	3	14.5814	15.2656	157.80	1.590
			4	14.5694	15.2536	176.00	1.478
4374 E1 (M 84)	18.5	0.7258	3	15.2556	15.9814	160.60	1.571
			4	15.2597	15.9855	168.80	1.520
4406 E3 (M 86)	16.8	0.8268	3	12.6832	13.5100	148.30	1.657
			4	12.6746	13.5014	165.50	1.540
4472 E2 (M 49)	17.1	1.1330	3	21.0525	22.1856	184.20	1.434
			4	21.0657	22.1988	197.0	1.371
4494 E2	16.6	0.1734	3	4.4278	4.6012	94.470	2.238
			4	4.4240	4.5974	102.70	2.117
4621 E4	14.9	0.2586	3	5.9330	6.1916	106.70	2.063
			4	5.9285	6.1871	124.60	1.861
4636 E2	14.3	0.4680	3	11.5722	12.0402	141.70	1.708
			4	11.5560	12.0239	152.90	1.623
4649 E2 (M 60)	17.3	0.9479	3	17.9113	18.8593	171.40	1.504
			4	17.9158	18.8637	181.80	1.446
4697 E4	11.4	0.2433	3	6.3532	6.5964	109.60	2.027
			4	6.3512	6.5944	130.50	1.804
7454 E2	23.2	0.0741	3	1.9067	1.9808	65.60	2.854
			4	1.9061	1.9802	72.02	2.681

The masses of dark matter,  $M_{dm}$ , and halo,  $M_h$ , as well as the scales radius  $r_s$  (in kpc) were calculated according to models 3 and 4. The masses are expressed in  $10^{12}M_{\odot}$ , and the normalization coefficient  $K$ , in  $10^{-3}M_{\odot}/pc^3$ .

**Table 2.** Values of the semi-major,  $a$ , and semi-minor,  $c$ , axes of EGs and the effective radius  $R_{eff}$

NGC	$a, kpc$	$c, kpc$	$\beta_T$	$m_{eff}$	$\beta_{eff}$	$R_{eff}, kpc$
4365 E3	20.896	15.490	1485.30	0.284	2086.13	6.775
4374 E1 (M 84)	19.947	17.373	2555.69	0.223	3333.71	5.492
4406 E3 (M 86)	18.114	13.428	1032.91	0.341	1547.86	10.140
4472 E2 (M 49)	22.166	18.437	554.84	0.447	915.70	8.661
4494 E2	11.556	9.179	2014.97	0.247	2706.64	3.764
4621 E4	11.114	7.343	790.19	0.397	1253.04	3.219
4636 E2	12.824	10.424	415.36	0.542	742.45	6.50
4649 E2 (M 60)	18.228	15.515	452.54	0.476	763.50	6.421
4697 E4	9.991	6.304	854.44	0.375	1324.18	3.922
7454 E2	6.889	5.348	913.44	0.346	1366.87	2.692

The quantities  $\beta_T$  and  $\beta_{eff}$  correspond to  $m = 1$  and  $m = m_{eff}$ .

**Table 3.** Values of the central,  $\sigma_0$ , and spatial,  $\sigma(R_{eff})$ , velocity dispersion at the distance of the effective radius and density  $\rho_0$  in the center of EGs

NGC	$\sigma_0, km/s$	$\sigma(R_{eff}), km/s$				$\rho_0, M_{\odot}/pc^3$		
		1	2	3	4	5	6	7
4365 E3	255.90	221.31	289.33	221.09 387.66	118.90 382.24	68.194	10.73	39.46
4374 E1 (M 84)	288.40	258.23	388.14	239.80 377.34	128.80 371.51	132.71	17.52	75.113
4406 E3 (M 86)	216.80	190.55	500.52	219.23 426.45	118.61 380.13	22.58	3.45	13.01380
4472 E2 (M 49)	288.30	250.04	527.89	255.82 442.19	137.85 419.12	50.844	7.89	29.366
4494 E2	261.80	224.39	248.31	152.10 266.16	81.82 265.60	98.34	21.42	59.878
4621 E4	223.90	197.69	188.99	195.54 293.71	105.02 298.20	254.57	37.97	146.27
4636 E2	199.50	181.55	627.01	195.39 358.66	105.58 349.67	48.12	5.18	26.650
4649 E2 (M 60)	314.80	267.92	519.45	262.99 410.56	141.57 394.82	105.36	13.22	59.289
4697 E4	180.70	169.43	263.52	183.70 301.03	98.93 307.87	129.73	22.31	76.017
7454 E2	231.70	223.36	240.49	118.60 210.25	63.90 209.71	113.05	18.10	65.572

Columns 1 and 2 correspond to the values of  $\sigma(R_{eff})$  given in [29] and calculated according to (29). Columns 3 and 4 in the first row give the values of  $\sigma_s(R_{eff})$  taking into account only the BM component, and in the second column, the values of  $\sigma(R_{eff})$  with allowance for the DM components. Columns 5 and 6 correspond to the values of  $\rho_0$ , and column 7 to the average value of  $\rho_0$ .

**Table 4.** Values of total gravitational energies  $W$  and rotational kinetic energies  $T$  of EGs, calculated according to models 3, 4, and 5

NGC	Models	$W$	$T$	NGC	Models	$W$	$T$
4365	3	a) $-1.0238$	$0.1727$	4621	3	a) $-0.2567$	$0.0419$
		b) $-1.0303$	$0.1728$			b) $-0.2574$	$0.0419$
	4	a) $-9.4667$	$0.3803$		4	a) $-2.0220$	$0.1124$
		b) $-9.4689$	$0.3803$			b) $-2.0223$	$0.1125$
	5	1) $-1951.72$	$78.516$		5	1) $-2372.6$	$132.12$
		2) $-0.0461$	$0.0018$			2) $-0.0139$	$0.0008$
4374	3	a) $-1.1024$	$0.1843$	4636	3	a) $-0.7105$	$0.1177$
		b) $-1.1089$	$0.1843$			b) $-0.7122$	$0.1178$
	4	a) $-12.644$	$0.2339$		4	a) $-7.8006$	$0.2168$
		b) $-1.1089$	$0.1843$			b) $-7.8012$	$0.2168$
	5	1) $-2466.2$	$45.666$		5	1) $-6223.4$	$173.13$
		2) $-0.0568$	$0.0011$			2) $-0.0258$	$0.0007$
4406	3	a) $-0.8899$	$0.1415$	4649	3	a) $-1.4781$	$0.2379$
		b) $-0.8946$	$0.1416$			b) $-1.4836$	$0.2379$
	4	a) $-7.7024$	$0.3093$		4	a) $-16.171$	$0.3492$
		b) $-7.7042$	$0.3093$			b) $-16.173$	$0.3492$
	5	1) $-3570.2$	$143.63$		5	1) $-897.39$	$193.99$
		2) $-0.0561$	$0.0023$			2) $-0.0909$	$0.0020$
4472	3	a) $-1.8781$	$0.3079$	4697	3	a) $-0.2809$	$0.0462$
		b) $-1.8881$	$0.3081$			b) $-0.2815$	$0.04621$
	4	a) $-19.874$	$0.4906$		4	a) $-2.1293$	$0.1315$
		b) $-19.877$	$0.4906$			b) $-2.1295$	$0.1315$
	5	1) $-5665.061$	$140.03$		5	1) $-4244.45$	$262.47$
		2) $-0.1022$	$0.00253$			2) $-0.0111$	$0.0007$
4494	3	a) $-0.1501$	$0.0261$	7454	3	a) $-0.0412$	$0.0068$
		b) $-0.1535$	$0.0262$			b) $-0.0414$	$0.0069$
	4	a) $-1.6082$	$0.0497$		4	a) $-0.4231$	$0.0144$
		b) $-1.6084$	$0.0497$			b) $-0.4232$	$0.0145$
	5	1) $-289.60$	$8.9556$		5	1) $-178.75$	$6.0820$
		2) $-0.0052$	$0.0002$			2) $-0.0014$	$0.00005$

For models 3 and 4, the values of  $W$  and  $T$  are given according to variants (a) and (b), and, for model 5, according to variants (1) and (2). The energies are expressed in units of  $10^{55} J$ .

## 9. CONCLUSIONS

Three new models of an EG, which are in good agreement with modern ideas about the structure of such galaxies, have been created for solving some problems of celestial mechanics and astrophysics. According to these models, an EG together with the halo is regarded as a two-layer inhomogeneous ellipsoid of revolution: a prolate spheroid. For a triaxial ellipsoid, the axis of rotation does not coincide with the apparent minor axis and the isophotes are misaligned. For an oblate spheroid, in which the apparent axis coincides with the axis of rotation and the isophote alignment is not violated, is not considered. In this case, the outer and inner layers are assumed to be similar and concentric and their centers coincide with the center of the EG. The LP EG is considered the inner layer and comprises an inhomogeneous prolate spheroid with a spheroidal density distribution. In the LP EG, a BM with an astrophysical law of density distribution prevails. The outer part comprises an inhomogeneous spherical layer with a spherical density distribution (model 3) or a spheroidal layer with a spheroidal density distribution (model 4). According to model 3, the outer layer and the halo of the galaxy are bounded by a sphere with a radius equal to the scale radius of the EG, and, according to model 4, they are bounded by a spheroidal surface with a semi-major axis equal to the scale radius of the galaxy. It is assumed that the spherical and spheroidal layers mainly consist of dark matter (DM) and, depending on its presence in the central regions of the EG, in models 3 and 4, two variants are considered: (a) the main part of DM is outside the LP EG and (b) the DM content in the inner regions of the EG is comparable to the BM content. In this case, the matching conditions for the potential at the interface between the LP SC and the spherical (spheroidal) layer are determined.

According to model 5, an EG with (variant 1) or without (variant 2) a halo comprises an inhomogeneous ellipsoid of revolution: a prolate spheroid consisting of BM and DM. In model 5, there is no interface between the LP SC and the homeoid; therefore, the fulfillment of the matching conditions for the potential is not considered.

Within the three new models of a layered inhomogeneous EG, the total gravitational (potential) energy, rotational kinetic energy, and velocity dispersion at a distance of the effective radius of the galaxy have been determined. A new method for determining the average values of the scale radius of an EG, the density at its center, and the value of the parameter corresponding to the effective radius of the galaxy is proposed.

The results obtained have been applied to sixty EGs and are presented in the form of tables for ten galaxies.

Analysis of the equilibrium and stability of a dynamic system created on the basis of models 3 and 4 will be carried out by the author separately.

## ACKNOWLEDGMENTS

I am grateful to professor B.P. Kondratyev for valuable advices and remarks.

## REFERENCES

- 1.S. A. Gasanov, *Astron. Rep.* 56, 469 (2012).
- 2.S. A. Gasanov, *Astron. Rep.* 58, 167 (2014).

- 3.S. A. Gasanov, *Astron. Rep.* 59, 238 (2015).
- 4.A. V. Zasov, A. S. Saburova, A. V. Khoperskov, and S.A. Khoperskov, *Phys. Usp.* 60, 3 (2017).
- 5.G. Bertin, R. P. Saglia, and M. Stiavelli, *Astrophys. J.* 384, 423 (1992).
- 6.M. Oguri, C. E. Rusu, and E. E. Falco, *Mon. Not. R. Astron. Soc.* 439, 2494 (2014).
- 7.G. de Vaucouleurs, A. de Vaucouleurs, H. Corwin, R.J. Buta, G. Paturel, and P. Fouque, *Third Reference Catalogue of Bright Galaxies* (Springer, New York, 1991), Vols. 2, 3.
- 8.B. P. Kondratyev and L. M. Ozernoi, *Sov. Astron. Lett.* 5, 37 (1979).
- 9.B. P. Kondratyev and V. S. Kornoukhov, *Mon. Not. R. Astron. Soc.* 478, 3159 (2018).
- 10.B. P. Kondrat'ev, *Cand. Sci. Dissertation* (Moscow, 1982).
- 11.B. P. Kondrat'ev, *Theory of Potential. New Methods and Problems with Solutions* (Mir, Moscow, 2007) [in Russian].
- 12.E. Hubble, *Astrophys. J.* 71, 231 (1930).
- 13.J. F. Navarro, C. S. Frenk, and S. D. M. White, *Astrophys. J.* 490, 493 (1997).
- 14.P. Cote, D. E. McLaughlin, J. G. Cohen, and J. P. Blakeslee, *Astrophys. J.* 591, 850 (2003).
- 15.J. Binney and S. Tremaine, *Galactic Dynamics* (Princeton Univ. Press, Princeton, 2008).
- 16.W. Dehnen, *Mon. Not. R. Astron. Soc.* 265, 250 (1993).
- 17.S. Tremaine, D. O. Richstone, Y.-I. Byun, A. Dressler, S. M. Faber, C. Grillmair, J. Kormendy, and T.R.Lauer, *Astron. J.* 107, 634 (1994).
- 18.L. Hernquist, *Astrophys. J.* 356, 359 (1990).
- 19.W. Jaffe, *Mon. Not. R. Astron. Soc.* 202, 995 (1983).
- 20.G. N. Duboshin, *Celestial Mechanics. Fundamental Problems and Methods* (Nauka, Moscow, 1968) [in Russian].
- 21.J. P. Ostriker and P. J. E. Peebles, *Astrophys. J.* 186, 467(1973).
- 22.V. L. Polyachenko and A. M. Fridman, *Physics of Gravitating Systems I: Equilibrium and Stability* (Nauka, Moscow, 1976; Springer, Berlin, 1984).
- 23.J. Binney, *Mon. Not. R. Astron. Soc.* 190, 873 (1980).
- 24.J. Binney, *Mon. Not. R. Astron. Soc.* 190, 421 (1980).
- 25.Ph. Prugniel and F. Simien, *Astron. Astrophys.* 321, 111 (1997).
- 26.B. Terzic and A. W. Graham, *arXiv: astro-ph/0506192* (2005).
- 27.R. L. Davies, E. M. Sadler, and R. F. Peletier, *Mon. Not. R. Astron. Soc.* 262, 650 (1993).
- 28.S. Samurovic, *Astron. Astrophys.* 470, A132 (2014).
- 29.M. Cappellari, E. Emsellem, D. Krajnovic, R. M. McDermid, et al., *Mon. Not. R. Astron. Soc.* 413, 813 (2011).
- 30.M. Cappellari, N. Scott, K. Alatalo, L. Blitz, et al., *Mon. Not. R. Astron. Soc.* 432, 1709 (2013).
- 31.L. R. Spitler, A. F. Dunkan, J. Strader, J. P. Brodie, and J. S. Gallagher, *Mon. Not. R. Astron. Soc.* 385, 361 (2008).
- 32.D. A. Forbes, L. Sinertu, G. Savorgnan, A. J. Romanowsky, C. Usher, and J. Brodie, *Mon. Not. R. Astron. Soc.* 464, 4611 (2017).
- 33.E. van Uitert, M. Cacciato, H. Hoekstra, M. Brower, et al., *Mon. Not. R. Astron. Soc.* 459, 3251 (2016).

34.D.-W. Kim and G. Fabbiano, *Astrophys. J.* 812, 127 (2015).

35.D. N. Spergel, R. Bean, O. Dore, M. R. Nolta, et al., *Astrophys. J. Suppl.* 170, 377 (2007).