

Resolving competition of charge-density wave and superconducting phases using the MPS+MF algorithm

Gunnar Bollmark,¹ Thomas Köhler,¹ and Adrian Kantian^{1,2}

¹*Department of Physics and Astronomy, Uppsala University, Box 516, S-751 20, Uppsala, Sweden*

²*SUPA, Institute of Photonics and Quantum Sciences,
Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom*

Materials with strong electronic correlations may exhibit a superconducting phase when tuning some parameters, but they almost always also have multiple other phases, typically insulating ones, that are in close competition with superconductivity. It is highly challenging to resolve this competition with quantitative numerics for the group of quasi-two-dimensional (Q2D) materials such as the cuprates. This is the case even for the simplified minimal models of these materials, the doped 2D Hubbard model with repulsive interactions, where clusters of sufficient size to determine the phase in the thermodynamic limit can be hard-to-impossible to treat in practice. The present work shows how quasi-one-dimensional (Q1D) systems, 2D and 3D arrays of weakly coupled 1D correlated electrons, are much more amenable to resolve the competition between superconducting and insulating orders on an equal footing using many-body numerics. Using the recently established matrix product state plus mean field (MPS+MF) approach for fermions [1], we demonstrate that large systems are readily reachable in these systems, and thus the thermodynamic regime by extrapolation. Focussing on basic model systems, 3D arrays of negative- U Hubbard chains with additional nearest-neighbour interaction V , we show that despite the MF component of the MPS+MF technique we can reproduce the expected coexistence of superconductivity and charge-density wave at $V = 0$ for density $n = 1$. We then show how we can tune away from coexistence by both tuning V and doping the system. This work thus paves the way to deploy two-channel MPS+MF theory on some highly demanding high- T_c superconducting systems, such as 3D arrays of repulsive- U doped Hubbard ladders, where we have recently characterized the properties of such arrays in single-channel MPS+MF calculations [1]. The present approach could thus conclusively show that this superconducting order would actually be obtained, by explicitly comparing superconductivity against its insulating competitors (magnetic orders already being eliminated by design in these systems).

I. INTRODUCTION

Understanding the macroscopic physics of strongly correlated electrons remains difficult for many systems. One of the main issues is the necessity of properly taking correlations between electrons into account, beyond effective single-particle descriptions. Problems where electronic correlations play a major role feature interesting and exotic physics ranging from unconventional superconductivity (USC) to spin-charge separation to quasi-one-dimensional (Q1D) systems [2–4].

A prominent example of the difficulty of accounting for correlations can be found in the two-dimensional (2D) Hubbard models with repulsive interactions, originally proposed to describe the physics of CuO_2 sheets in cuprate superconductors [5]. This class of superconducting materials attain superconductivity at large critical temperatures when doped away from unit density. Yet, trying to match the complex known phase diagram of, e.g., the cuprates with that of a doped 2D Hubbard model at strong electron-electron repulsion remains very difficult [2, 6–13]. This is due to the challenge of being able to treat a cluster of sufficient size to draw meaningful inferences about the thermodynamic limit of the system, which typically requires to resolve the competition of insulating and superconducting phases occurring for such systems [5]. For techniques based on quantum Monte Carlo (QMC), this is due to an apparently incurable sign-

problem, and for matrix-product state (MPS)-based ones on the superlinear scaling of entanglement with subsystem surface [14]. Just for one particular doped 2D Hubbard model (pure square lattice with only on-site repulsion) it has taken a sustained multi-group, multi-method effort over many years to accrue evidence that its ground state is more likely than not a stripe-order insulator instead of a d -wave superconductor [8, 9, 11, 12]. How these findings might change for any change in band structure, or for longer-range repulsion, or when coupling many 2D systems into a 3D array as required to represent an actual solid, is then another completely open problem.

Q1D many-body systems of fermions, which are 2D or 3D arrays of weakly coupled correlated one-dimensional (1D) fermions, offer a much better outlook for efficiently resolving the competition between competing ordered phases in unconventionally and high- T_c superconducting systems. Recently, an algorithm for solving such Q1D systems using a combination of MPS numerics and mean-field (MF) theory [1, 15, 16] has been developed. The MPS+MF algorithm used in these works either studies cases where no competition of phases could occur or the winning phase was surmised from effective field theory arguments available due to the 1D nature of the sub-units from which the 2D or 3D array is constituted [4]. However, these arguments are predicated entirely on the fastest diverging susceptibility of the 1D sub-units, which are then re-summed within a random




FIG. 1. A schematic appearance of: (a) the full Hamiltonian given in Eq. (1) and (b) the effectively one-dimensional mean-field Hamiltonian of Eq. (4).

phase or MF approximation to yield susceptibilities for the entire array. The reasoning is thus incomplete – in practice, different susceptibilities have different prefactors, which can also tip the balance towards a particular phase. Yet these prefactors are not known quantitatively from the effective field theory.

In the present work we introduce and test a multi-channel-MF version of the MPS+MF framework for correlated-fermion Q1D systems that can resolve the competition between superconductivity (SC) and charge-density wave (CDW) order, which are treated on an equal footing. This is done without pre-supposing any partic-

ular structure in the multiple ordering channels (other than the superconducting order being of singlet-type). This general approach allows the self-consistent flow to minimize the global (free) energy across multiple channels at once. This allows the expanded MPS+MF framework to also detect phase coexistence where this is physically possible.

In order to accomplish this a simple attractive fermion model composed of negative- U Hubbard chains weakly coupled together is studied. Such a system admits solution via other methods allowing us to determine whether the constructed algorithm performs in accordance with expectation. Thus, the paper is structured as follows: In Sec. II the model is presented and microscopic parameters defined. Continuing, the MF theory used to turn the Q1D system into an effective 1D one is derived in Sec. III. Additionally, the new channel of potential CDW ordering is presented and its behavior discussed. In Sec. IV the model is studied in both a strong and weak attractive regime. Coexistence is investigated by varying microscopic parameters and doping that affect the balance of orders. Finally, conclusions are given in Sec. V.

II. MODEL

In order to produce a MF framework capable of resolving competition of CDW and SC in Q1D systems we focus on a simple model amenable both to analytical methods and exact numerical methods:

$$\hat{H} = \hat{H}_0 - t_{\perp} \hat{H}_{\perp}. \quad (1)$$

The basic Hamiltonian is a set of disconnected 1D systems, described by \hat{H}_0 , arranged equidistantly from each other. These 1D systems are indexed by a set of 2D vectors $\{\mathbf{R}_n\}$. The disconnected Hamiltonian is given by (suppressing the \mathbf{R}_n on each operator)

$$\hat{H}_0 = \sum_{\{\mathbf{R}_n\}} \hat{H}_{1D}(\mathbf{R}_n) = \sum_{\{\mathbf{R}_n\}} \left\{ -t \sum_{i\sigma} \left[\hat{c}_{i+1,\sigma}^{\dagger} \hat{c}_{i,\sigma} + \text{h.c.} \right] - |U| \sum_i n_{i,\uparrow} \hat{n}_{i,\downarrow} + V \sum_i \hat{n}_i \hat{n}_{i+1} - \mu \sum_i \hat{n}_i \right\}, \quad (2)$$

and the perpendicularly connecting Hamiltonian is given by

$$\hat{H}_{\perp} = \frac{1}{2} \sum_i \sum_{\{\mathbf{R}_n\}} \sum_{\langle \mathbf{R}_n, \mathbf{R}_m \rangle} \hat{c}_{i,\sigma}^{\dagger}(\mathbf{R}_n) \hat{c}_{i,\sigma}(\mathbf{R}_m) + \text{h.c.} \quad (3)$$

Notably, tunneling only takes place between nearest-neighbor pairs $\langle \mathbf{R}_n, \mathbf{R}_m \rangle$. Dealing with a Hamiltonian of this form has been done using the MPS+MF algorithm [1, 15]. For strongly correlated fermions previous work using MPS+MF has assumed SC correlations dominate the ground state. However, studying systems with unit filling or adding the finite V in Eq. (2) this is no

longer necessarily true. This requires a formulation of MPS+MF with an additional MF channel for describing insulating phases.

III. CHARGE-DENSITY WAVES WITH MEAN-FIELD THEORY

In deriving the self-consistent equations for MPS+MF a perturbation theory is first applied to turn tunneling into effective interaction. Subsequently, a MF treatment is applied to this interaction: The usual Hartree-Fock and Bogoliubov contributions are taken into account produc-




FIG. 2. Figure showing the competition between SC and CDW phases at $t_{\perp} = 0.3t$ and $U = -10t$ for a system of length $L = 96$. The upper row shows a plot of local pairing amplitude over system size while the bottom row shows local density. In each figure the two final self-consistency iterations are plotted: When the plot is blue it lies on top of the red one. If both iterations are visible then they are dissimilar indicating a CDW phase.

ing a self-consistent effective 1D Hamiltonian [1]

$$\begin{aligned} \hat{H}_{\text{MF}} = & \hat{H}_{\text{1D}}(\mathbf{R}_n) - \sum_{i,k} \alpha_{i,k} \left(\hat{c}_{i,\uparrow} \hat{c}_{k,\downarrow} + \hat{c}_{k,\downarrow}^{\dagger} \hat{c}_{i,\uparrow}^{\dagger} \right) \\ & + \sum_{i,\sigma} \sum_{r=1}^{L-i} \beta_{i,r,\sigma} \left(\hat{c}_{i+r,\sigma}^{\dagger} \hat{c}_{i,\sigma} + \hat{c}_{i,\sigma}^{\dagger} \hat{c}_{i+r,\sigma} \right) \\ & + \sum_{i,\sigma} \mu_{i,\sigma} \hat{n}_{i,\sigma}, \end{aligned} \quad (4)$$

where

$$\alpha_{i,k} = \frac{2z_c t_{\perp}^2}{\Delta E_p} \langle \hat{c}_{i,\uparrow} \hat{c}_{k,\downarrow} \rangle, \quad (6)$$

$$\beta_{i,r,\sigma} = \frac{2z_c t_{\perp}^2}{\Delta E_p} \langle \hat{c}_{i+r,\sigma}^{\dagger} \hat{c}_{i,\sigma} \rangle \quad (7)$$

$$\mu_{i,\sigma} = \frac{z_c t_{\perp}^2}{\Delta E_p} (2 \langle \hat{n}_{i,\sigma} \rangle - 1), \quad (8)$$

where all amplitudes are assumed to be real. This requires the definition of pairing energy

$$\begin{aligned} \Delta E_p = 2E \left(N+1, S = \frac{1}{2} \right) - E(N+2, S=0) \\ - E(N, S=0), \end{aligned} \quad (9)$$

with $E(N, S)$ the spectrum of \hat{H}_{1D} indexed by particle number and spin. The coordination number z_c count the neighboring systems of a single 1D sub-system.

Notably, the MF terms Eqs. (6)-(8) are easily interpreted: Pairing lowers system energy whereas particle-hole terms carry a cost instead. Thus, pairing with

any separation of the constituents is generally favored whereas dispersion is disfavored. The final term is unique in that it provides a repulsive interaction, capable of driving charge order. At the same time, density is constant for a system which orders in the pairing operator and $\mu_{i,\sigma}$ would simply modify the chemical potential μ . In this work we expand the MF theory to include a fully site-dependent $\mu_{i,\sigma}$ -term, which can drive charge-ordering.

Solving such simplified systems is the subject of MPS+MF and is done by iteration of the MF amplitudes until they achieve self-consistency. While, in general the initial configuration of these amplitudes should not affect the final solution, in practice there are several choices that are more advantageous than others as expounded on in App. A.

At the basis of the MF Hamiltonian in Eq. (4) lies perturbation theory which produces an effective inter-chain interaction. It is notable that one form of interaction produced is a repulsive one which will tend to disfavor neighboring particles of like spin in the transverse direction. Thus, it would lead to a pinning behavior in these directions yielding a checkerboard appearance. Hence, we call such a phase CDW inspired by the 1D physics of \hat{H}_{1D} [4]. In order to solve this expanded set of amplitudes we must answer how such a term behaves under the MF theory used in MPS+MF.

A. Two-period in mean-fields

Performing MF theory on the inter-chain interaction changes the meaning of the term. While, in general, such

interaction would compare densities at different chains, the MF treatment assumes that all 1D sub-systems are the same, turning the transverse interaction into a density comparison in the same chain. This is undesirable since it gives rise to a local interaction between same-spin particles which would be prohibited by the Pauli principle. However, numerically solving for self-consistency requires iterative solution. Hence, the energy contribution to minimize at each iteration is really given by

$$V_{\perp} \propto (2 \langle \hat{n}_{i,\sigma} \rangle_{j-1} - 1) \langle \hat{n}_{i,\sigma} \rangle_j, \quad (10)$$

where $\langle \rangle_j$ indicates measurement at iteration j of the self-consistent calculation. Notably, since the interaction is repulsive the system will strive to minimize $\langle \hat{n}_{i,\sigma} \rangle_{j-1} \langle \hat{n}_{i,\sigma} \rangle_j$. In other words, a state ordering in this operator will in iteration j avoid high density in the regions of iteration $j-1$ which already had high density there. Hence a checkerboard-like pattern is achieved.

The cost of this is that the self-consistent iteration will at best converge to two different states, endlessly oscillating between the two so as to keep $\langle \hat{n}_{i,\sigma} \rangle_{j-1} \langle \hat{n}_{i,\sigma} \rangle_j$ small. When this occurs we name it a two-periodic behavior and surmise that it indicates that ordering in the CDW channel of the system is preferred. One concern of this interpretation is whether the Hamiltonian maintains tunneling equally strong to and from sites of large and small density. Since the MF amplitudes in Eq. (7) depend on this measurement it may be a concern whether hermiticity is maintained. We reason that this is not an issue case due to high-amplitude transitions always end up in low-amplitude states and vice versa (see App. B).

IV. RESULTS

In this work we are primarily interested in whether and how an MF theory based on the two-periodic MF amplitude in Sec. III can resolve physics expected from analytical methods in the same system. As such, we first study strongly interacting fermions at $U = -10t$ in a unit filled system

$$n = \frac{1}{L} \sum_{i=1}^L (\langle \hat{n}_{i,\uparrow} \rangle + \langle \hat{n}_{i,\downarrow} \rangle) = 1. \quad (11)$$

The pairing gap in such a system we compute to approximately $\Delta E_p \approx 6.5t$ yielding tightly bound pairs. This allows us to cut off the number of mean-field amplitudes (this is explained in more detail in [1])

$$\alpha_{i,k} = \begin{cases} \alpha_{i,k} & \text{if } |i-k| \leq R \\ 0 & \text{otherwise} \end{cases}, \quad (12)$$

$$\beta_{i,r,\sigma} = \begin{cases} \beta_{i,r,\sigma} & \text{if } |i-k| \leq R \\ 0 & \text{otherwise} \end{cases}, \quad (13)$$

where R is the cut-off point. With very tight binding such as for $U = -10t$ we choose this value at $R = 1$.

The great advantage of this system is that it is amenable to practically exact treatment which can provide useful standards of comparison to the developed method.

Once the method is established we attempt to develop a methodology in a lightly doped system at weaker interaction of $U = -2t$. In this case the pairing energy is somewhat affected by the other microscopic parameters and we find that it evaluates to $\Delta E_p \in [0.1t, 0.2t]$. Previous work on these systems indicate that a range of $R = 5$ should yield acceptable precision and as such we limit simulation to $R = 5$ when $U = -2t$. Interestingly, such a weakly attractive system lies close in similarity to previously studied lightly doped Hubbard ladders [1]. Therefore we also study the effects of doping in the weakly attractive case to see its effects on competition.

Notably, since the algorithm solving the effective 1D problem at each iteration is a MPS-based implementation of density-matrix renormalization group (DMRG) it is subject to the precision limitations of such an algorithm [17–20]. The two parameters which affect precision are system size and bond dimension which are considered in App. C. It is found that the precision is not sufficiently affected by these parameters to change the qualitative nature of the results. Thus, the sizes and bond dimensions which are found in App. C are presented in this section.

A. Unit filling competition

At unit density the CDW and SC phases can be shown to coexist analytically by virtue of a particle-hole transformation [21]. Specifically, there is symmetry between CDW and SC phases in such conditions meaning that if one occurs then the other is also present by symmetry. Any method aiming to resolve competition between these phases should be able to find such coexistence at unit density. In order to achieve this we focus on strongly attracting fermions. For such a case there is particularly good control as fermionic particles pair up more often than not leading to a bosonic description.

In order to resolve the competition we study the MF amplitudes corresponding to CDW and SC order respectively. For CDW order the local density should enter a two-periodic behavior when the system is minimizing energy using this channel. Conversely, a superconductor can be studied with the Bardeen-Cooper-Schrieffer (BCS) order parameter $\langle \hat{c}_{i,\uparrow} \hat{c}_{i,\downarrow} \rangle$. Whether the local density enters a two-periodic behavior or the BCS order parameter attains finite value will determine which order the ground state obtains.

Knowing what to look for, the nearest neighbor interaction V is varied in order to bias the system towards any given order: repulsion, $V > 0$, for CDW and attraction, $V < 0$ for SC. As can be seen in Fig. 2 the system's ground state changes notably over a small range of nearest neighbor interaction. The appearance can be roughly summarized as follows:

1. Weak attraction: The systems local density ap-

proaches a constant throughout the system and likewise the local pairing.

2. No interaction: Local density enters a two-periodic behavior while simultaneously local pairing remains in a single period and constant throughout the system.
3. Weak repulsion: Local pairing is now zero and local density has a strong two-period.

These results indicate that MPS+MF can be used to meaningfully study competition of CDW and SC phases in the model under study. Notably, the strongly interacting regime is amenable to other methods: particularly due to its likeness to a bosonic model where each doublon represent a single hard-core boson.

While it is important to keep the structure of the solution provided in Fig. 2 in mind it contains a lot more information than is necessary for determination of phases. The superconducting order parameter is approximately constant throughout the system inviting the definition of a system-wide order parameter

$$\langle \hat{c}\hat{c} \rangle = \frac{1}{i_l - i_f + 1} \sum_{i=i_f}^{i_l} \langle \hat{c}_{i,\uparrow} \hat{c}_{i,\downarrow} \rangle. \quad (14)$$

The inclusion of limits to the average is due to the open-boundary conditions (OBC) nature of DMRG: the edges act as impurities which disturb the bulk solution in which we are truly interested. In this manner we can eliminate edge effects by averaging over the center of the system.

Typically, a CDW phase would be identified by the long-range behavior of density-density correlations. However, when the term which drives CDW orders, the self-consistent solution provides two different ground states and self-consistent Hamiltonians. A linear combination of these ground states may solve the issue but requires in general that the states do not overlap: Something we cannot guarantee. Instead, it is clear that the two-periodic behavior can be diagnosed by a difference of local density at two different self-consistent iterations:

$$\delta n = \frac{1}{i_l - i_f + 1} \sum_{i=i_f}^{i_l} |\langle \hat{n}_i \rangle_j - \langle \hat{n}_i \rangle_{j-1}|. \quad (15)$$

Using these definitions a mapping of the order parameters at several different V and t_\perp can be easily displayed as shown in Fig. 3.

While being able to qualitatively replicate known results is useful as verification for the method, we can use the MPS+MF routine to move on to models of weakly interacting fermions. Moving outside the composite bosonic regime we target a weak attraction of $U = -2t$. Such a system has comparable pairing strength to weakly doped repulsive Hubbard ladders, making for an interesting study on the effects of competition in such systems.




FIG. 3. Order parameter of SC and CDW order from Eqs. (14) and (15), respectively. The simulation is performed for $U = -10t$, $n = 1$, $L = 160$ and varying t_\perp and V . The SC order uses red color and CDW order blue.




FIG. 4. Order parameter of SC and CDW order from Eqs. (14) and (15), respectively. The simulation is performed for $U = -2t$, $t_\perp = 0.05t$, $L = 192$ and varying n and V . The SC order uses red color and CDW order blue.

Notably, we find the qualitative picture remains: Figure 4 shows the competition when moving away from $V = 0$. The coexistence is found once again for $V = 0$ in the same figure.

B. Hole doping

In previous studies the MPS+MF method has been applied to systems which exhibit USC as a consequence of repulsively mediated electron pairing [1]. Specifically, lightly doped repulsive Hubbard ladders weakly coupled in a Q1D array are known to feature a quasi-long range order instability to SC but simultaneously to that of CDW. This competition is currently not possible to conclusively resolve with analytical methods and a more quantitative method would be the safest alternative.

In Fig. 4 we show an example of a weakly attracting system doped away from unit filling. Slightly doping the system to a filling of $n = 15/16$ sees the local pairing and superconductivity remain in the system while order parameter even increases. Interestingly, the local density

difference is clearly finite at $n = 1$ with a two-periodic behavior that disappears with doping. Furthermore, applying a repulsive potential to the system as when $n = 1$, thereby increasing the cost of pair condensation no longer destroys the superconductor.

V. CONCLUSIONS

In this work we have developed a modified version of the existing MPS+MF framework. Lifting assumptions of dominant ordering the competition between SC and CDW phases in fermionic systems with heuristic pairing is studied. This method carries the benefit of resolving competition in a MF sense without the use of Renormalization Group theory.

From the developed method it is apparent that the expectation of coexistence of CDW and SC phases at unit filling can be resolved. Including nearest neighbor interactions allows the biasing of the system towards either a SC or CDW phase. With weak bias to either attraction or repulsion we find the coexistence broken in the expected direction. Thus, we conclude that the method is able to resolve competition between these two phases at the very least on the MF level.

Utilizing the newly developed method we apply it to a more complicated case: Weakly interacting fermions doped away from unit filling. In this case, doping lifts the coexistence and favors a SC phase: Something that is also concluded by quasi-ordering arguments in bosonization calculations.

For future work, this method can be utilized in resolving competition between phases in more complex systems such as, e.g., arrays of doped repulsive Hubbard ladders: Conclusively answering the question of which phase such a system orders in.

ACKNOWLEDGMENTS

This work was supported by an ERC Starting Grant from the European Union’s Horizon 2020 research and innovation programme under grant agreement No. 758935; and the UK’s Engineering and Physical Sciences Research Council [grant number EP/W022982/1]. The computations were enabled by resources provided by the Swedish National Infrastructure for Computing (SNIC), as well as by compute time awarded by the UK’s Engineering and Physical Sciences Research Council on the ARCHER2, Peta4-Skylake and Cirrus compute clusters through the “Access to HPC” and “Scottish Academic Access” calls.

Appendix A: Initial values for self-consistent iteration

In order to establish the symmetry between CDW and SC phases for $n = 1$ care must be taken to what initial values are chosen for the MF amplitudes. In order to produce coexistence we choose

$$\alpha_{i,k}^{(\text{init})} = \begin{cases} \alpha_{i-k} & \text{if } i = k \\ 0 & \text{otherwise} \end{cases}, \quad (\text{A1})$$

which represents a strongly local pairing constant throughout the system. This drives towards a SC phase. Conversely, since coexistence occurs at $n = 1$ the periodicity of a CDW phase is 2, i.e., the density repeats every other site. Hence, the CDW MF is initialized with

$$\mu_{i\sigma}^{(\text{init})} = \begin{cases} \mu_{\sigma} & \text{if } i \text{ even} \\ -\mu_{\sigma} & \text{if } i \text{ odd} \end{cases}, \quad (\text{A2})$$

with the additional condition that $\mu_{\uparrow} = \mu_{\downarrow}$. This choice drives towards a CDW phase with period 2. Lastly the particle-hole are not deemed as leading drivers toward any order and are initialized at zero:

$$\beta_{i,r,\sigma}^{(\text{init})} = 0. \quad (\text{A3})$$

1. Doped CDW

When doping away from $n = 1$ the SC phase should remain equally simple: Constant local pairing throughout the system. Conversely, the CDW phase can longer be argued to have a simple period of 2 in the density profile. In fact, theory would expect each 1D system to obtain a period of $2/n$ [4]. However, light doping causes density in a discrete system to repeat only at a large number of sites.

One method, used in this work, to obtain reasonable initial values for CDW in a doped system is given by letting $\alpha_{i,k} = \beta_{i,r,\sigma} = 0$ at all iterations. With this restriction no pairing is possible and only the CDW MF is kept in the solution leaving no other ordering channel than a CDW. The resulting solution is kept as initial for a competition between CDW and SC phases.

Appendix B: Hermiticity of two-period solution

One potential concern for the two-periodic behavior is that the Hamiltonian strongly disrespects translational symmetry. Due to this property care must be taken that qualities such as hermiticity are preserved: Tunneling out of a high-density site into a low-density one could possibly have different amplitude than its conjugate. Such an observation is relevant for the two-periodic behavior associated with the occurrence of CDW. In order to resolve




FIG. 5. Comparison of measurements with $U = -10t$ and $n = 1$ at $L \in [128, 160]$ and bond dimension $\chi \in [100, 200]$ with various t_{\perp} and V .

this, consider the ground state of one of the two-period solutions

$$|\psi\rangle = |\phi\rangle + |\text{CDW}\rangle, \quad (\text{B1})$$

where the CDW part contains all two-periodic behavior and $|\phi\rangle$ single period effects. For unit filling the density is large at every other site and almost zero at the others. It makes intuitive sense that a tunneling event out of a high-density region is more probable than into one yet such events are conjugate.

Using $|\psi\rangle$ the measurement becomes

$$\begin{aligned} \langle\psi|\hat{c}_i^{\dagger}\hat{c}_j|\psi\rangle &= \langle\phi|\hat{c}_i^{\dagger}\hat{c}_j|\phi\rangle + \langle\text{CDW}|\hat{c}_i^{\dagger}\hat{c}_j|\text{CDW}\rangle \\ &+ \langle\phi|\hat{c}_i^{\dagger}\hat{c}_j|\text{CDW}\rangle + \langle\text{CDW}|\hat{c}_i^{\dagger}\hat{c}_j|\phi\rangle, \end{aligned} \quad (\text{B2})$$

and the conjugate process with $i \leftrightarrow j$. Performing the application and writing down amplitudes yields

$$\begin{aligned} \langle\psi|\hat{c}_i^{\dagger}\hat{c}_j|\psi\rangle &= \alpha_1 \underbrace{\langle\phi|\phi'\rangle}_{\beta_1} + \alpha_2 \underbrace{\langle\text{CDW}|\text{CDW}'\rangle}_{\beta_2} \\ &+ \alpha_3 \underbrace{\langle\phi|\text{CDW}'\rangle}_{\beta_3} + \alpha_4 \underbrace{\langle\text{CDW}|\phi'\rangle}_{\beta_4}, \end{aligned} \quad (\text{B3})$$

where α_i indicate the amplitudes of the surviving states after application of $\hat{c}_i^{\dagger}\hat{c}_j$. The β_i then indicate the amplitudes of the states which contain finite overlap with the primed states. Importantly, the primed states contain a single particle moved from j to i . Thus, any finite

β_i result from having states in the primed superpositions which are already inside the unprimed ones. However, if those states are already inside the unprimed superpositions then the conjugate operation $\hat{c}_j^{\dagger}\hat{c}_i$ generates the same states as those spared by application of $\hat{c}_i^{\dagger}\hat{c}_j$. Hence,

$$\begin{aligned} \langle\psi|\hat{c}_j^{\dagger}\hat{c}_i|\psi\rangle &= \beta_1^* \underbrace{\langle\phi|\phi''\rangle}_{\alpha_1^*} + \beta_2^* \underbrace{\langle\text{CDW}|\text{CDW}''\rangle}_{\alpha_2^*} \\ &+ \beta_3^* \underbrace{\langle\phi|\text{CDW}''\rangle}_{\alpha_3^*} + \beta_4^* \underbrace{\langle\text{CDW}|\phi''\rangle}_{\alpha_4^*}, \end{aligned} \quad (\text{B4})$$

which indicates that $\langle\hat{c}_i^{\dagger}\hat{c}_j\rangle = \langle\hat{c}_j^{\dagger}\hat{c}_i\rangle$ if the amplitudes are real.

Appendix C: Precision of simulation

While simulation is done with quasi-exact methods in combination with mean-field theory the former always carry a quantitative error. Since it is not the aim of this paper to precisely compute any property of the system under study the commonly applied linear extrapolation in truncation error is not utilized [22, 23]. Instead, a much more qualitative comparison is performed where the precision parameters are varied and we check for notable changes.

It is important to note that errors in the state due to insufficient bond dimension can be more than a question of precision. In fact, if bond dimension is taken low




FIG. 6. Comparison of measurements at $U = -2t$ and $t_{\perp} = 0.05t$ for $L \in [128, 192]$ and bond dimension $\chi \in [400, 600]$, varying n and V .

$U/t \backslash \chi$	100	200	400	600
-10.0	10^{-9}	10^{-13}	-	-
-2.0	10^{-4}	10^{-5}	10^{-7}	10^{-8}

TABLE I. Table showing the order of magnitude of truncation error ϵ_{ψ} for the various bond dimension χ used.

enough qualitative differences can appear as well. An example of such a difference is when an incorrect phase

for the ground state is realized due to insufficient bond dimension to capture the true ground state. In order to ward against this issue we show the order of magnitude for truncation errors associated with our simulations in table I [19]. In general, we find that given enough bond dimension the truncation error goes towards zero.

Additionally, since the DMRG routine is run in a finite size manner we also expect finite size effects. Assuming that very large sizes correctly capture the essential physics we compare measurements for different sizes and bond dimension in Figs. 5 and 6. Notably, we find that neither size nor bond dimension has a large qualitative impact on the measurements.

-
- [1] G. Bollmark, T. Köhler, L. Pizzino, Y. Yang, H. Shi, J. S. Hofmann, H. Shi, S. Zhang, T. Giamarchi, and A. Kantian, [Solving 2D and 3D lattice models of correlated fermions – combining matrix product states with mean field theory](#) (2022).
- [2] P. W. Anderson, *The Theory of Superconductivity in the High- T_c Cuprate Superconductors* (Princeton University Press, Princeton, NJ, 1997).
- [3] G. R. Stewart, *Unconventional superconductivity*, [Adv. Phys. **66**, 75 \(2017\)](#).
- [4] T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, New York, 2003).
- [5] D. P. Arovas, E. Berg, S. A. Kivelson, and S. Raghu, *The Hubbard Model*, [Annu. Rev. Condens. Matter Phys. **13**, 239 \(2022\)](#).
- [6] J. Orenstein, *Advances in the Physics of High-Temperature Superconductivity*, [Science **288**, 468 \(2000\)](#).
- [7] D. J. Scalapino, *A common thread: The pairing interaction for unconventional superconductors*, [Rev. Mod. Phys. **84**, 1383 \(2012\)](#).
- [8] J. P. F. LeBlanc, A. E. Antipov, F. Becca, I. W. Bulik, G. K. Chan, C.-M. Chung, Y. Deng, M. Ferrero, T. M. Henderson, C. A. Jiménez-Hoyos, E. Kozik, X.-W. Liu, A. J. Millis, N. V. Prokof'ev, M. Qin, G. E. Scuseria, H. Shi, B. V. Svistunov, L. F. Tocchio, I. S. Tupitsyn, S. R. White, S. Zhang, B.-X. Zheng, Z. Zhu, and E. Gull, *Solutions of the Two-Dimensional Hubbard Model: Benchmarks and Results from a Wide Range of*

- Numerical Algorithms*, *Phys. Rev. X* **5**, 41041 (2015).
- [9] B.-X. Zheng, C.-M. Chung, P. Corboz, G. Ehlers, M.-P. Qin, R. M. Noack, H. Shi, S. R. White, S. Zhang, and G. K.-L. Chan, *Stripe order in the underdoped region of the two-dimensional Hubbard model*, *Science* **358**, 1155 (2017).
- [10] A. Bohrdt, C. S. Chiu, G. Ji, M. Xu, D. Greif, M. Greiner, E. Demler, F. Grusdt, and M. Knap, *Classifying snapshots of the doped Hubbard model with machine learning*, *Nat. Phys.* **15**, 921 (2019).
- [11] M. Qin, C. M. Chung, H. Shi, E. Vitali, C. Hubig, U. Schollwöck, S. R. White, and S. Zhang, *Absence of Superconductivity in the Pure Two-Dimensional Hubbard Model*, *Phys. Rev. X* **10**, 031016 (2020).
- [12] A. Wietek, Y. Y. He, S. R. White, A. Georges, and E. M. Stoudenmire, *Stripes, Antiferromagnetism, and the Pseudogap in the Doped Hubbard Model at Finite Temperature*, *Phys. Rev. X* **11**, 31007 (2021).
- [13] A. Bohrdt, L. Homeier, C. Reinmoser, E. Demler, and F. Grusdt, *Exploration of doped quantum magnets with ultracold atoms*, *Ann. Phys.* **435**, 168651 (2021), arXiv: 2107.08043 Publisher: Elsevier Inc.
- [14] E. Stoudenmire and S. R. White, *Studying Two-Dimensional Systems with the Density Matrix Renormalization Group*, *Annu. Rev. Condens. Matter Phys.* **3**, 111 (2012), publisher: Annual Reviews.
- [15] G. Bollmark, N. Laflorencie, and A. Kantian, *Dimensional crossover and phase transitions in coupled chains: Density matrix renormalization group results*, *Phys. Rev. B* **102**, 195145 (2020).
- [16] S. Marten, G. Bollmark, T. Köhler, S. R. Manmana, and A. Kantian, *Transient superconductivity in three-dimensional Hubbard systems by combining matrix product states and self-consistent mean-field theory*, (2022), arXiv: 2207.09841v1.
- [17] S. R. White, *Density matrix formulation for quantum renormalization groups*, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [18] S. R. White, *Density-matrix algorithms for quantum renormalization groups*, *Phys. Rev. B* **48**, 10345 (1993).
- [19] U. Schollwöck, *The density-matrix renormalization group*, *Rev. Mod. Phys.* **77**, 259 (2005).
- [20] U. Schollwöck, *The density-matrix renormalization group in the age of matrix product states*, *Ann. Phys.* **326**, 96 (2011).
- [21] A. F. Ho, M. A. Cazalilla, and T. Giamarchi, *Quantum simulation of the Hubbard model: The attractive route*, *Phys. Rev. A* **79**, 33620 (2009).
- [22] S. R. White and D. A. Huse, *Numerical renormalization-group study of low-lying eigenstates of the antiferromagnetic $S=1$ Heisenberg chain*, *Phys. Rev. B* **48**, 3844 (1993).
- [23] Ö. Legeza and G. Fáth, *Accuracy of the density-matrix renormalization-group method*, *Phys. Rev. B* **53**, 14349 (1996).