quTARANG: A python GPE solver to study turbulence in quantum systems

Shawan Kumar Jha^{*,†,1}, Sachin Singh Rawat^{*,2}, Mahendra Kumar Verma², and Pankaj Kumar Mishra¹

¹Department of Physics, Indian Institute of Technology Guwahati, Guwahati 781039, India ²Department of Physics, Indian Institute of Technology Kanpur,

Kanpur 208016, India

Abstract

quTARANG is a Python-based general-purpose Gross-Pitaevskii Equation (GPE) solver. It can solve GPE in 1D, 2D and 3D and has the ability to run on both CPU and GPU. It has been developed to study turbulence in quantum systems, specifically in atomic Bose-Einstein condensates, and can be used to study different quantities, such as the varied spectra associated with quantum turbulence.

Summary

Turbulence is a phenomenon associated with the chaotic nature of flows in space and time solely arising due to the nonlinear nature of the interactions. Turbulence in classical fluids, as best characterised by the Navier-Stokes equation, remains unresolved to this day. The complex nonlinear interactions at various length and time scales make it a difficult problem to handle analytically as well as numerically. One recent approach to shed some light on this long-standing problem has been the study

^{*}These authors contributed equally to this work.

[†]Correspondence: shawankumar@iitg.ac.in

of turbulence in quantum fluids. The zero viscosity and quantized vortices of quantum fluid systems like Bose-Einstein Condensates (BECs) [1] distinguish it from its classical counterparts. BEC is a state of matter where Bose particles occupy the ground state upon cooling to a very low temperature and thus can be represented by a macroscopic wave function. One can model the dynamics of BECs using the mean-field Gross-Pitaevskii Equation (GPE) given by [2]

$$i\hbar\partial_t\psi(\vec{r},t) = -\frac{\hbar^2}{2m}\nabla^2\psi(\vec{r},t) + V(\vec{r},t)\psi(\vec{r},t) + NU_0|\psi(\vec{r},t)|^2\psi(\vec{r},t),$$
(1)

where $\psi(\vec{r}, t)$ is the macroscopic complex wave function, m is the atomic mass, $V(\vec{r}, t)$ is the trapping potential, N is the number of particles, $U_0 = (4\pi\hbar^2 a_s)/m$ is the nonlinear interaction parameter and a_s denotes the scattering length for the interaction of the atomic particles.

Our quantum simulator code quTARANG¹ is primarily designed for studying quantum turbulence in BECs by solving the GPE in laminar as well as in the turbulent regime.

Statement of Need

quTARANG is a robust and easy-to-use application that solves the GPE with Graphics Processing Unit (GPU). GPUs are specialized units primarily designed for image processing and for performing massive multigrid simulations at high speed. They have a large number of parallelizing units compared to CPUs, due to which they are being widely used to speed up code. There are no packages available in python that can solve turbulent GPE in 2D and 3D on both CPUs and GPUs. There exist, however, some software packages in other languages that can solve the GPE such as GPELab [3], Massively Parallel Trotter-Suzuki Solver [4], CUDA-enabled GPUE [5], a split-step Crank-Nicolson based Fortran code [2] and MPI-OpenMP enabled Gross-Pitaevskii Solver (GPS) [6].

Numerical Scheme and functionalities

We have chosen the frequency (ω^{-1}) as the time scale, oscillator length $a_0 = \sqrt{\hbar/m\omega}$ as the characteristic length scale and the harmonic oscillator ground state energy $\hbar\omega$ as the energy scale. With this formalism, the non-dimensional variables can be

 $^{^1} github.com/sachinrawat2207/quTARANG$




Figure 1: A comparison of the dynamical evolution of the root-mean-square size of the condensate in the x (σ_x), y (σ_y), and z (σ_z) direction obtained from our simulation (for 3D GPE) and those obtained by Bao et al. [7]. A perfect match is being obtained.

written as $t' = \omega t$, $\vec{r'} = \vec{r}/a_0$ and $\psi' = a_0^{3/2}\psi$. In what follows, we omit the prime (') from the variables. The non-dimensional form of the GPE is given by [2]

$$i\partial_t \psi(\vec{r},t) = -\frac{1}{2}\nabla^2 \psi(\vec{r},t) + V(\vec{r},t)\psi(\vec{r},t) + g|\psi(\vec{r},t)|^2\psi(\vec{r},t)$$
(2)

where g is the non-dimensional interaction parameter.

quTARANG uses a pseudo-spectral scheme, Time-splitting spectral (TSSP) method [7], to solve the dynamics of the GPE. The main advantage of using the TSSP scheme is that it is unconditionally stable and conserves the total particle number.

The ground state calculations in quTARANG are done by using TSSP with an imaginary time propagation method. In this method, one replaces t with -it. As we propagate in imaginary time, the eigenstates with higher energies begin to decay faster than the ground state as a result of which only the ground state survives. The wavefunction needs to be normalised at each time step in order to conserve total particle number.

quTARANG is equipped with various features, which include:

- 1. Ground state calculations for different potentials such as harmonic and anharmonic trap, optical lattice potential, time-dependent potential and stochastic potential.
- 2. Long-time dynamical evolution of different states using either CPUs or GPUs.
- 3. Computation of different quantities relevant to the study of turbulence phenomenon in BECs, such as components of kinetic energy (KE) and various spectra (compressible KE spectrum, incompressible KE spectrum and particle number spectrum).

Table 1: The condensate width (r_{rms}) and chemical potential (μ) obtained for the ground state using quTARANG. r_{rms}^* and μ^* are the corresponding values from [2] for comparison.

Dimension	g	r_{rms}	r_{rms}^{*}	μ	μ^*
	-2.5097	0.87771	0.87759	0.49987	0.49978
	0	1.00000	1.00000	1.00000	1.00000
	3.1371	1.10504	1.10513	1.42009	1.42005
2	12.5484	1.30656	1.30687	2.25609	2.25583
	62.742	1.78722	1.78817	4.61136	4.60982
	313.71	2.60122	2.60441	10.07639	10.06825
	627.42	3.07914	3.08453	14.20569	14.18922
	0	1	1	3.0000	3.0000
	18.81	1.3778	1.3249	4.3618	4.3611
	94.05	1.8222	1.7742	6.6824	6.6797
3	188.1	2.0881	2.0411	8.3718	8.3671
	940.5	2.8912	2.8424	14.9663	14.9487
	1881	3.3268	3.2758	19.5058	19.4751
	7524	4.3968	4.3408	33.5623	33.4677
	15048	5.0497	4.9922	44.1894	44.0234

Results

We have calculated the ground state and dynamics for given sets of initial conditions and compared them with the standard results. 1. Validation of ground state : The initial condition for 2D and 3D cases are given as:

2D:
$$\psi(\vec{r}, 0) = \left(\frac{1}{\pi}\right)^{1/2} e^{-\frac{(x^2+y^2)}{2}}, \quad V(\vec{r}) = \frac{1}{2}(x^2+y^2)$$

3D: $\psi(\vec{r}, 0) = \left(\frac{1}{\pi}\right)^{3/4} e^{-\frac{(x^2+y^2+z^2)}{2}}, \quad V(\vec{r}) = \frac{1}{2}(x^2+y^2+4z^2)$

The ground state obtained by quTARANG has been compared with that obtained by using the finite difference code of Muruganandam and Adhikari [2]. The results for harmonic potential well for 2D and 3D are in good agreement with Muruganandam and Adhikari [2] as shown in Table 1. The root-mean-square size (r_{rms}) of the condensate is defined as $r_{rms} = (\int r^2 |\psi(\vec{r}, t)|^2 dV)^{1/2}$.

2. Validation of dynamics : We have validated the dynamic evolution of a state by comparing our results with Bao et al. [7] for the following condition:

$$\psi(\vec{r},0) = \frac{(\gamma_y \gamma_z)^{1/4}}{\sqrt{(\pi\epsilon_1)^{3/4}}} e^{-\frac{(x^2 + \gamma_y y^2 + \gamma_z z^2)}{2\epsilon_1}}, \quad V(\vec{r},0) = \frac{1}{2}(x^2 + \gamma_y^2 y^2 + \gamma_z^2 z^2),$$

where $\gamma_y = 2.0$, $\gamma_z = 4.0$, $\epsilon_1 = 0.25$ and g = 0.1. Fig 1 shows the comparisons of the rms size of the condensate in x, y and z directions at different times calculated by using our code and those obtained by Bao et al. [7]. The results obtained from the **quTARANG** are in good agreement with the results obtained from Bao et al. for the same initial conditions.

References

- L. Madeira, A. Cidrim, M. Hemmerling, M. A. Caracanhas, F. E. A. dos Santos, and V. S. Bagnato, AVS Quantum Sci., 2(3), 035901 (2020).
- [2] P. Muruganandam and S. K. Adhikari, Computer Physics Communications, 180(10), 1888-1912 (2009).
- [3] X. Antoine and R. Duboscq, Computer Physics Communications, 185(11), 2969–2991 (2014).
- [4] P. Wittek and F. M. Cucchietti, Computer Physics Communications, 184(4), 1165–1171 (2013).
- [5] J. R. Schloss and L. J. O'Riordan, The Journal of Open Source Software, 3(32), 1037 (2018).

- [6] M. Kobayashi, P. Parnaudeau, F. Luddens, C. Lothodé, L. Danaila, M. Brachet and I. Danaila, Computer Physics Communications, 258, 107579 (2021).
- [7] W. Bao, D. Jacksch, and P. A. Markowich, J. Comp. Phys., 187(1), 318-342 (2003).