

Gravitational Baryogenesis: Problems and Possible Resolution

Presented at 6th International Conference on Particle Physics and Astrophysics (ICCPA-2022)

E. Arbuzova^{a,b}, A. Dolgov^{b,c}, K. Dutta^d, R. Rangarajan^e

January 23, 2023

^aDepartment of Higher Mathematics, Dubna State University, Universitetskaya st. 19,
Dubna, Moscow region 141983, Russia;

^bDepartment of Physics, Novosibirsk State University,
Pirogova st. 2, Novosibirsk, 630090 Russia

^cBogolyubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
Joliot-Curie st. 6, Dubna, Moscow region, 141980 Russia

^dDepartment of Physical Sciences, Indian Institute of Science Education and Research
Kolkata, Mohanpur 741246, India;

^eMathematical and Physical Sciences Division, School of Arts and Sciences, Ahmedabad
University, Navrangpura, Ahmedabad 380009, India.

Abstract

The coupling of baryonic current to the derivative of the curvature scalar, R , inherent to gravitational baryogenesis (GBG), leads to a fourth order differential equation of motion for R instead of the algebraic one of General Relativity (GR). The fourth-order differential equation is generically unstable. We consider a possible mechanism of stabilization of GBG by modification of gravity, introducing an R^2 -term into the canonical action of GR. It is shown that this mechanism allows for stabilization of GBG with bosonic and fermionic baryon currents. We have established the region of the model parameters leading to stabilization of R . Still, the standard cosmology would be noticeably modified.

1 Introduction

An excess of matter over antimatter in our Universe is crucial for our very existence and is well supported by various observations. The local Universe is clearly matter dominated. The amount of antimatter is very small and it can be explained as the result of high energy collisions in space. On the other hand, matter and antimatter seem to have similar properties, therefore we could expect a matter-antimatter symmetric universe. The existence of large regions of antimatter in our neighbourhood would produce high energy radiation created by matter-antimatter annihilation on the boundaries between matter and antimatter domains, which is not observed. A satisfactory model of our Universe should be able to explain the origin of the matter-antimatter asymmetry. Any initial asymmetry at inflation could not solve the problem of observed excess of matter over antimatter, because the energy density associated with the observed non-zero baryonic number density would not allow for sufficiently long inflation.

The term baryogenesis is used to indicate the generation of the excess of matter (baryons) over antimatter (antibaryons) or vice versa.

In 1967 Andrey Sakharov formulated three conditions today known as Sakharov Principles [1], necessary to produce a matter-antimatter asymmetry in the initially symmetric universe. These conditions include:

1. Non-conservation of baryonic number;
2. Breaking of symmetry between particles and antiparticles;
3. Deviation from thermal equilibrium.

However, not all of three Sakharov Principles are strictly necessary. For example, spontaneous baryogenesis (SBG) and gravitational baryogenesis (GBG) do not demand an explicit C and CP violation and can proceed in thermal equilibrium. Moreover, these mechanisms are usually most efficient in thermal equilibrium.

The statement that the cosmological baryon asymmetry can be created by spontaneous baryogenesis in thermal equilibrium was mentioned in the original paper by A. Cohen and D. Kaplan in 1987 [2] and in the subsequent papers by A. Cohen, D. Kaplan, and A. Nelson [3, 4] (for a review see [5–8]).

The term "spontaneous" is related to spontaneous breaking of underlying symmetry of the theory, which ensures the conservation of the total baryonic number in the unbroken phase. This symmetry is supposed to be spontaneously broken and in the broken phase the Lagrangian density acquires the term

$$\mathcal{L}_{SBG} = (\partial_\mu \theta) J_B^\mu, \quad (1)$$

where θ is a (pseudo) Goldstone field, and J_B^μ is the baryonic current of matter fields, which becomes non-conserved as a result of the symmetry breaking.

For a spatially homogeneous field, $\theta = \theta(t)$, the Lagrangian is reduced to a simple form

$$\mathcal{L}_{SBG} = \dot{\theta} n_B, \quad n_B \equiv J_B^0. \quad (2)$$

Here n_B is the baryonic number density, so it is tempting to identify $\dot{\theta}$ with the chemical potential, μ_B , of the corresponding system. However, such an identification is questionable [9, 10]. It depends upon the representation chosen for the fermionic fields and is heavily based on the assumption $\dot{\theta} \approx \text{const}$. In Ref. [9] the assumption $\dot{\theta} \approx \text{const}$ is relaxed.

Stimulated by spontaneous baryogenesis the idea of gravitational baryogenesis was put forward [11]. The scenario of SBG was modified by the introduction of the coupling of the baryonic current to the derivative of the curvature scalar R :

$$\mathcal{S}_{GBG} = -\frac{1}{M^2} \int d^4x \sqrt{-g} (\partial_\mu R) J_B^\mu, \quad (3)$$

where g is the determinant of the space-time metric tensor and the mass parameter M determines the energy scale of baryogenesis. There are a lot of articles on the subject, and a partial list of references is included in Refs. [12–16]. According to these papers, the GBG mechanism can successfully explain the magnitude of the cosmological baryon asymmetry of the universe.

However, it was argued in Refs. [17, 18], that the back reaction of the created non-zero baryonic density on the space-time curvature leads to strong instability of the cosmological evolution. In this paper we show that the problem of stability can be solved by adding to the Hilbert-Einstein action the quadratic in curvature term generated by quantum corrections [19, 20]. The underlying gravitational action has the form:

$$S_{Grav} = -\frac{M_{Pl}^2}{16\pi} \int d^4x \sqrt{-g} \left(R - \frac{R^2}{6M_R^2} \right), \quad (4)$$

where $M_{Pl} = 1.22 \cdot 10^{19}$ GeV is the Planck mass, and we use the metric signature $(+, -, -, -)$. As is known, the R^2 -term leads to excitation of the scalar degree of freedom, named scalaron, and M_R is the scalaron mass. In the very early universe the R^2 -term can generate inflation [21], and density perturbations. The amplitude of the observed density perturbations demands that $M_R = 3 \cdot 10^{13}$ GeV [22] if the scalaron is the inflaton. Otherwise $M_R > 3 \cdot 10^{13}$ GeV is allowed. Below we presume that the scalaron is the inflaton.

2 Instability problem of gravitational baryogenesis

The essential ingredient of the spontaneous baryogenesis is the coupling of the baryonic current the derivative of the curvature scalar $\partial_\mu R$ (3). Taken over canonical cosmological Friedmann-Lemaître-Robertson-Walker background, this interaction can successfully fulfil the task of generating the proper value of the baryon asymmetry of the universe.

However, any curvature dependent term in the Lagrangian of the theory would modify the equations of the General Relativity (GR). The modified GR equations have been analysed in Refs. [9, 18]. Since interaction (3) is not just linear in the curvature term multiplied by a constant, it leads to higher order equations describing evolution of gravitational fields. Higher order equations of motion are typically unstable with respect to small perturbations. According to the results of Refs. [9, 18], it indeed happens in the

frameworks of the SBG scenario and the characteristic time of the exponential instability is much shorter than the cosmological time. It creates serious problem for realisation of the SBG mechanism.

In this work we suggest to consider possible stabilisation of SBG and have proved that it can be realised but the resulting cosmological model suffers from too large value of R , much larger than that in the classical Friedmann cosmology. Possible ways to cure this shortcoming are mentioned.

3 Stabilisation of gravitational baryogenesis in modified gravity

3.1 Bosonic case.

Let us first consider the case when baryonic number is carried by a complex scalar field ϕ [17]. The total action has the form:

$$S_{tot}[\phi] = - \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{16\pi} \left(R - \frac{R^2}{6M_R^2} \right) + \frac{1}{M^2} (\partial_\mu R) J_{(\phi)}^\mu - g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi^* + U(\phi, \phi^*) \right] + S_{matt} \quad (5)$$

where $U(\phi, \phi^*)$ is the potential of field ϕ and S_{matt} is the matter action which does not include the field ϕ . In Eq. (5) $R(t)$ is the classical curvature field, while $\phi(\vec{x}, t)$ is the quantum operator of light scalar particles.

We assume that the potential $U(\phi, \phi^*)$ is not invariant with respect to phase transformation $\phi \rightarrow \exp(iq\beta)\phi$ and thus the corresponding current

$$J_{(\phi)}^\mu = iq g^{\mu\nu} (\phi^* \partial_\nu \phi - \phi \partial_\nu \phi^*) \quad (6)$$

is not conserved. Here q is the baryonic number of field ϕ . The non-conservation of the current is necessary for the proper performance of the model, otherwise S_{GBG} in Eq. (3) can be integrated away by parts.

Varying action (5) over $g^{\mu\nu}$ we come to the following equations:

$$\begin{aligned} & \frac{M_{Pl}^2}{16\pi} \left[R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R - \frac{1}{3M_R^2} \left(R_{\mu\nu} - \frac{1}{4} g_{\mu\nu} R + g_{\mu\nu} D^2 - D_\mu D_\nu \right) R \right] \\ & - \frac{1}{M^2} \left[(R_{\mu\nu} - (D_\mu D_\nu - g_{\mu\nu} D^2)) D_\alpha J_{(\phi)}^\alpha + \frac{1}{2} g_{\mu\nu} J_{(\phi)}^\alpha D_\alpha R - \frac{1}{2} (J_{(\phi)\nu} D_\mu R + J_{(\phi)\mu} D_\nu R) \right] \\ & - \frac{1}{2} (D_\mu \phi D_\nu \phi^* + D_\nu \phi D_\mu \phi^*) + \frac{1}{2} g_{\mu\nu} [D_\alpha \phi D^\alpha \phi^* - U(\phi)] - (D_\mu \phi)(D_\nu \phi^*) \\ & = \frac{1}{2} T_{\mu\nu}^{(matt)}, \end{aligned} \quad (7)$$

where D_μ is the covariant derivative in metric $g_{\mu\nu}$ (of course, for scalars $D_\mu = \partial_\mu$) and $T_{\mu\nu}^{(matt)}$ is the energy-momentum tensor of matter obtained from action S_{matt} .

Taking the trace of equation (7) with respect to μ and ν and changing sign we obtain:

$$\begin{aligned} \frac{M_{Pl}^2}{16\pi} \left(R + \frac{1}{M_R^2} D^2 R \right) + \frac{1}{M^2} [(R + 3D^2) D_\alpha J_{(\phi)}^\alpha + J_{(\phi)}^\alpha D_\alpha R] - D_\alpha \phi D^\alpha \phi^* + 2U(\phi) \\ = -\frac{1}{2} T^{(matt)} = 0, \end{aligned} \quad (8)$$

where $T^{(matt)} = g^{\mu\nu} T_{\mu\nu}^{(matt)}$ is the trace of the energy-momentum tensor of matter. For the usual relativistic matter $T^{(matt)} = 0$, while for scalar field ϕ the trace of the energy-momentum tensor is nonzero:

$$T_\mu^\mu(\phi) = -2D_\alpha \phi D^\alpha \phi^* + 4U(\phi). \quad (9)$$

The equation of motion for field ϕ is:

$$D^2 \phi + \frac{\partial U}{\partial \phi^*} = -\frac{iq}{M^2} (2D_\mu R D^\mu \phi + \phi D^2 R). \quad (10)$$

According to definition (6) and Eq. (10), the current divergence is:

$$D_\mu J^\mu = \frac{2q^2}{M^2} [D_\mu R (\phi^* D^\mu \phi + \phi D^\mu \phi^*) + |\phi|^2 D^2 R] + iq \left(\phi \frac{\partial U}{\partial \phi} - \phi^* \frac{\partial U}{\partial \phi^*} \right). \quad (11)$$

For homogeneous curvature scalar $R(t)$ in spatially flat FLRW-metric

$$ds^2 = dt^2 - a^2(t) d\mathbf{r}^2 \quad (12)$$

Eq. (8) is reduced to:

$$\begin{aligned} \frac{M_{Pl}^2}{16\pi} \left[R + \frac{1}{M_R^2} (\partial_t^2 + 3H\partial_t) R \right] + \frac{1}{M^2} [(R + 3\partial_t^2 + 9H\partial_t) D_\alpha J_{(\phi)}^\alpha + \dot{R} J_{(\phi)}^0] \\ + 2U(\phi) - (D_\alpha \phi)(D^\alpha \phi^*) = 0. \end{aligned} \quad (13)$$

where $J_{(\phi)}^0$ is the baryonic number density of the ϕ -field, $H = \dot{a}/a$ is the Hubble parameter, and the divergence of the current is given by the expression:

$$D_\alpha J_{(\phi)}^\alpha = \frac{2q^2}{M^2} [\dot{R} (\phi^* \dot{\phi} + \phi \dot{\phi}^*) + (\ddot{R} + 3H\dot{R}) \phi^* \phi] + iq \left(\phi \frac{\partial U}{\partial \phi} - \phi^* \frac{\partial U}{\partial \phi^*} \right). \quad (14)$$

As we see in what follows, the last two terms in Eq. (13) do not have an essential impact on the cosmological instability found in Ref. [17] and will be disregarded below.

Let us note that the statement of exponential instability of $R(t)$ [17] does not depend on the conservation or non-conservation of the current from the potential term $(\phi \partial U / \partial \phi - \phi^* \partial U / \partial \phi^*)$ in Eq. (14). However if the current from this term is conserved then the baryon asymmetry is not generated. On the other hand the term in square brackets in Eq. (14) does not lead to generation of the baryon asymmetry but leads to exponential instability of $R(t)$. Below we ignore the last term of Eq. (14).

Performing thermal averaging of the normal ordered bilinear products of field ϕ in the high temperature limit (see Appendix of Ref. [17]) in accordance with equations:

$$\langle \phi^* \phi \rangle = \frac{T^2}{12}, \quad \langle \phi^* \dot{\phi} + \dot{\phi}^* \phi \rangle = 0, \quad (15)$$

and using Eq. (14) we obtain the fourth order differential equation:

$$\begin{aligned} \frac{M_{Pl}^2}{16\pi} \left(R + \frac{1}{M_R^2} D^2 R \right) + \frac{q^2}{6M^4} (R + 3\partial_t^2 + 9H\partial_t) \left[(\ddot{R} + 3H\dot{R}) T^2 \right] + \frac{1}{M^2} \dot{R} \langle J_{(\phi)}^0 \rangle \\ = -2U(\phi) + (D_\alpha \phi)(D^\alpha \phi^*). \end{aligned} \quad (16)$$

Here $\langle J_{(\phi)}^0 \rangle$ is the thermal average value of the baryonic number density of ϕ , which is supposed to vanish initially, but created through the process of the gravitational baryogenesis. This term can be neglected because the baryon asymmetry is normally quite small. Even if it is not small it does not have considerable impact on the explosive rise of the curvature scalar. As we see in what follows the evolution of $R(t)$ proceeds much faster than the cosmological evolution, that is $\ddot{R}/\dot{R} \gg H$. Consequently, we neglect the terms proportional to R with respect to the terms proportional to the second derivative of R , \ddot{R} . We also consider the terms of the type HR as small w.r.t. to dR/dt . We can check that this presumption is true a posteriori with the obtained solution for $R(t)$.

Keeping only the dominant terms we simplify the above equation to:

$$\frac{d^4 R}{dt^4} + \frac{\kappa^4}{M_R^2} \frac{d^2 R}{dt^2} + \kappa^4 R = -T_\mu^\mu(\phi) \frac{M^4}{q^2 T^2}, \quad (17)$$

where

$$\kappa^4 = \frac{M_{Pl}^2 M^4}{8\pi q^2 T^2}. \quad (18)$$

While studying the instability of the solution we do not take into account the r.h.s. of Eq. (17) which does not depend upon R . Looking for the solution of Eq. (17) in the form $R = R_{in} \exp(\lambda t)$, we obtain the characteristic equation:

$$\lambda^4 + \frac{\kappa^4}{M_R^2} \lambda^2 + \kappa^4 = 0 \quad (19)$$

with the eigenvalues λ defined by the expression:

$$\lambda^2 = -\frac{\kappa^4}{2M_R^2} \pm \kappa^2 \sqrt{\frac{\kappa^4}{4M_R^4} - 1}. \quad (20)$$

There is no instability, if $\lambda^2 < 0$ and Eq. (17) has only oscillating solutions. It is realised, if $\kappa^4 > 4M_R^4$. Using the expression in Eq. (18) for κ^4 and taking $M_R = 3 \cdot 10^{13}$ GeV we find the stability condition:

$$M > 3 \cdot 10^4 \text{ GeV} \left(\frac{qT}{\text{GeV}} \right)^{1/2}, \quad (21)$$

which is fulfilled for all interesting values of M .

The value of λ depends upon the relation between κ and M_R . If $\kappa \sim M_R$ then the frequency of the oscillations of curvature is of the order of M_R and $|\lambda| \sim M_R$. If $\kappa \gg M_R$ then there are two possible solutions $|\lambda| \sim M_R$ and $|\lambda| \sim \kappa(\kappa/M_R) \gg M_R$. High frequency oscillations of R would lead to efficient gravitational particle production and, as a result, to damping of the oscillations.

3.2 Fermionic case

In this section we consider the case when baryonic number is carried by fermions. The gravitational part of the action has the form as in Eq. (4), while the fermionic part of the action is the same as in Refs. [10, 18]:

$$\begin{aligned} \mathcal{L}[Q, L] = & \frac{i}{2}(\bar{Q}\gamma^\mu\nabla_\mu Q - \nabla_\mu\bar{Q}\gamma^\mu Q) - m_Q\bar{Q}Q \\ & + \frac{i}{2}(\bar{L}\gamma^\mu\nabla_\mu L - \nabla_\mu\bar{L}\gamma^\mu L) - m_L\bar{L}L \\ & + \frac{g}{m_X^2}[(\bar{Q}Q^c)(\bar{Q}L) + (\bar{Q}^cQ)(\bar{L}Q)] + \frac{d}{M^2}(\partial_\mu R)J^\mu + \mathcal{L}_{matt}, \end{aligned} \quad (22)$$

where Q is the quark-like field with non-zero baryonic number B_Q , Q^c is the charged conjugated quark operator, L is another fermionic field (lepton), and ∇_μ is the covariant derivative of the Dirac fermions in tetrad formalism. The quark current is $J^\mu = B_Q\bar{Q}\gamma^\mu Q$ with γ^μ being the curved space gamma-matrices, and \mathcal{L}_{matt} describes all other forms of matter. The four-fermion interaction between quarks and leptons is introduced to ensure the necessary non-conservation of the baryon number with m_X being a constant parameter with dimension of mass and g being a dimensionless coupling constant. In the term, describing interaction of the baryonic current of fermions with the derivative of the curvature scalar, M is a constant parameter with dimension of mass and $d = \pm 1$ is dimensionless coupling constant which is introduced to allow for an arbitrary sign of the above expression.

Gravitational equations of motion with an account of R^2/M_R^2 -term in analogy with Eq. (7) take the form:

$$\begin{aligned} & \frac{M_{Pl}^2}{8\pi} \left[R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R - \frac{1}{3M_R^2} \left(R_{\mu\nu} - \frac{1}{4}g_{\mu\nu}R + g_{\mu\nu}D^2 - D_\mu D_\nu \right) R \right] \\ = & \frac{g_{\mu\nu}}{2} \frac{g}{m_X^2} [(\bar{Q}Q^c)(\bar{Q}L) + (\bar{Q}^cQ)(\bar{L}Q)] \\ & + \frac{i}{4} [\bar{Q}(\gamma_\mu\nabla_\nu + \gamma_\nu\nabla_\mu)Q - (\nabla_\nu\bar{Q}\gamma_\mu + \nabla_\mu\bar{Q}\gamma_\nu)Q] \\ & + \frac{i}{4} [\bar{L}(\gamma_\mu\nabla_\nu + \gamma_\nu\nabla_\mu)L - (\nabla_\nu\bar{L}\gamma_\mu + \nabla_\mu\bar{L}\gamma_\nu)L] \\ & - \frac{2d}{M^2} [R_{\mu\nu} + g_{\mu\nu}D^2 - D_\mu D_\nu] D_\alpha J^\alpha + \frac{d}{2M^2} (J_\mu\partial_\nu R + J_\nu\partial_\mu R) + T_{\mu\nu}^{matt}. \end{aligned} \quad (23)$$

Taking the trace of Eq. (23) with an account of fermion equations of motion we obtain:

$$-\frac{M_{Pl}^2}{8\pi} \left(R + \frac{1}{M_R^2} D^2 R \right) = m_Q \bar{Q} Q + m_L \bar{L} L + \frac{2g}{m_X^2} [(\bar{Q} Q^c)(\bar{Q} L) + (\bar{Q}^c Q)(\bar{L} Q)] - \frac{2d}{M^2} (R + 3D^2) D_\alpha J^\alpha + T_{matt}, \quad (24)$$

where T_{matt} is the trace of the energy momentum tensor of all other fields. In the early universe when various species are relativistic, we can take $T_{matt} = 0$. The average expectation value of the quark-lepton interaction term proportional to g is also small, so the contribution of all matter fields may be neglected and hence the only term which remains in the r.h.s. of Eq. (24) is that proportional to $D_\alpha J^\alpha$.

A higher order differential equation for R is obtained after we substitute the current divergence, $D_\alpha J^\alpha$, calculated from the kinetic equation in the external field R [18], into Eq. (24). For the spatially homogeneous case

$$D_\alpha J^\alpha = (\partial_t + 3H)n_B = I_B^{coll}, \quad (25)$$

where the collision integral, I_B^{coll} , in the lowest order of perturbation theory is equal to:

$$I_B^{coll} = -3B_q(2\pi)^4 \int d\nu_{q_1, q_2} d\nu_{\bar{q}_3, l_4} \delta^4(q_1 + q_2 - q_3 - l_4) [|A(q_1 + q_2 \rightarrow \bar{q}_3 + l_4)|^2 f_{q_1} f_{q_2} - |A(\bar{q}_3 + l_4 \rightarrow q_1 + q_2)|^2 f_{\bar{q}_3} f_{l_4}]. \quad (26)$$

Here $A(a \rightarrow b)$ is the amplitude of the transition from state a to state b , B_Q is the baryonic number of quark, f_a is the phase space distribution (the occupation number), and

$$d\nu_{q_1, q_2} = \frac{d^3 q_1}{2E_{q_1} (2\pi)^3} \frac{d^3 q_2}{2E_{q_2} (2\pi)^3}, \quad (27)$$

where $E_q = \sqrt{q^2 + m^2}$ is the energy of particle with three-momentum q and mass m . The element of phase space of final particles, $d\nu_{\bar{q}_3, l_4}$, is defined analogously.

We choose such representation of the quark operator, Q , for which the interaction of baryonic current with the derivative of the curvature scalar in Eq. (22) vanishes but reappears in the quark-lepton interaction term:

$$\frac{2g}{m_X^2} \left[e^{-3idB_Q R/M^2} (\bar{Q} Q^c)(\bar{Q} L) + e^{3idB_Q R/M^2} (\bar{Q}^c Q)(\bar{L} Q) \right]. \quad (28)$$

We make the simplifying assumption that the evolution of R can be approximately described by the law

$$R(t) \approx R(t_0) + (t - t_0) \dot{R}. \quad (29)$$

We assume that $\dot{R}(t)$ slowly changes at the characteristic time scale of the reactions, which contribute to the collision integral (26), and so we can approximately take $\dot{R} \approx const$.

According to the rules of quantum field theory the reaction probability is given by the square of the integral over space and time of the amplitude of the corresponding process. In

the case of time independent interaction it leads to the energy conservation, $\Sigma E_{in} = \Sigma E_{fin}$. If the interaction depends upon time the energy evidently is non-conserved and in our case, e.g. for the reaction $q_1 + q_2 \rightarrow \bar{q}_3 + l_4$, the energy balance has the form:

$$E(q_1) + E(q_2) = E(q_3) + E(l_4) + 3dB_Q\dot{R}/M^2. \quad (30)$$

In kinetic equilibrium the phase space distribution of fermions has the form

$$f = \frac{1}{e^{(E/T-\xi)} + 1} \approx e^{-E/T+\xi}, \quad (31)$$

where $\xi = \mu/T$ is the dimensionless chemical potential, different for quarks, ξ_q , and leptons, ξ_l . In thermal equilibrium case the condition of conservation of chemical potentials is fulfilled, that is $\Sigma \xi_{in} = \Sigma \xi_{fin}$. In particular it demands that chemical potentials of particles and antiparticles are equal by magnitude and have opposite signs: $\xi = -\bar{\xi}$, as follows e.g. from the consideration of particle-antiparticle annihilation into different numbers of photons. If energy is not conserved, due to time-dependent $R(t)$, the conservation of chemical potentials is also broken, as we see in what follows.

We assume that $\xi \ll 1$ and hence distribution (31) turns into:

$$f \approx e^{-E/T}(1 + \xi). \quad (32)$$

We also assume that $3dB_Q\dot{R}/(M^2T) \ll 1$ and correspondingly the balance of chemical potentials in equilibrium for the reactions $q_1 + q_2 \leftrightarrow \bar{q}_3 + l_4$ leads to:

$$3\xi_q - \xi_l - \frac{3dB_Q\dot{R}(t)}{M^2T} = 0. \quad (33)$$

Following Ref. [18], we express

$$n_B \approx \frac{g_s B_Q}{6} \xi_q T^3, \quad (34)$$

where g_s is the number of quark spin states. Since we are studying instability of R whose timescale is presumed to be much smaller than the expansion rate of the Universe, we approximate

$$D_\alpha J^\alpha \approx \dot{n}_B \approx \frac{g_s B_Q}{6} \dot{\xi}_q T^3 \quad (35)$$

$$\approx \frac{g_s B_Q}{6} \dot{\xi}_q^{eq} T^3, \quad (36)$$

ξ_q^{eq} is obtained from Eq. (33), using the conservation of the sum of baryonic and leptonic numbers which implies $\xi_l = -\xi_q/3$. Then

$$\xi_q^{eq} = \frac{9dB_Q\dot{R}(t)}{10M^2T}. \quad (37)$$

Substituting Eq. (37) in Eq. (36) and neglecting the \dot{T} -term, Eq. (24) gives the following fourth order differential equation for the curvature scalar:

$$\frac{d^4 R}{dt^4} + \frac{\kappa_f^4}{M_R^2} \frac{d^2 R}{dt^2} + \kappa_f^4 R = 0, \quad (38)$$

where

$$\kappa_f^4 = \frac{5M_{Pl}^2 M^4}{36\pi g_s B_Q^2 T^2}. \quad (39)$$

Once again, we consider terms containing R as small with respect to the terms containing \ddot{R} . The value of κ_f is only slightly numerically different from κ in Eq. (18) and has the same dependence upon the essential parameters, so the solutions of Eqs. (17) and (38) practically coincide.

4 Discussion

We have shown that discovered in Refs. [17, 18] exponential instability of the curvature scalar inherent to the mechanism of spontaneous baryogenesis can be successfully cured in modified gravity. The special form of gravity modification by introduction of R^2 -term into canonical Hilbert-Einstein action of General Relativity was explored as a workable mechanism.

However, the stabilized asymptotic value of R is extremely large and together with possibly successful baryogenesis would still strongly perturb canonical cosmology. Possible ways out of this problem could either be a more complicated model of $F(R)$ gravity or a proper account of particle production created by high frequency oscillations of $R(t)$. Both options open interesting possibilities for future research.

References

- [1] Sakharov A. D. Violation of CP Invariance, C asymmetry, and baryon asymmetry of the universe. *Pisma Zh. Eksp. Teor. Fiz.* **1967**, 5, 32–35.
- [2] Cohen A. G. and Kaplan D. B. Thermodynamic Generation of the Baryon Asymmetry. *Phys. Lett. B* **1987**, 199, 251–258.
- [3] Cohen A. G. and Kaplan D. B. Spontaneous Baryogenesis. *Nucl. Phys. B* **1988**, n308, 913–928.
- [4] Cohen A. G., Kaplan D. B and Nelson A. E. Spontaneous baryogenesis at the weak phase transition. *Phys. Lett. B* **1991**, 263, 86–92.
- [5] Dolgov A. D. NonGUT baryogenesis. *Phys. Rept.* **1992**, 222 (1992), 309–386.

- [6] Rubakov V. A. and Shaposhnikov M. E. Electroweak baryon number nonconservation in the early universe and in high-energy collisions. *Usp. Fiz. Nauk* **1996**, *166*, 493–537.
- [7] Riotto A. and Trodden M. Recent progress in baryogenesis. *Ann. Rev. Nucl. Part. Sci.* **1999**, *49*, 35–75.
- [8] Dolgov A.D. Baryogenesis, 30 years after. *Surveys in High Energy Physics* **1998** *13*, 83.
- [9] Arbuzova E. V., Dolgov A.D. and Novikov V. A. General properties and kinetics of spontaneous baryogenesis. *Phys. Rev. D* **2016**, *94*, no.12, 123501.
- [10] Dasgupta A., Jain R. K. and Rangarajan R., Effective chemical potential in spontaneous baryogenesis. *Phys. Rev. D* **2018** *98*, no.8, 083527.
- [11] Davoudiasl H., R. Kitano R., Kribs G. D., H. Murayama H., Steinhardt P. J. Gravitational baryogenesis. *Phys. Rev. Lett.* **2004**, *93*, 201301.
- [12] Lambiase G. and Scarpetta G. Baryogenesis in f(R): Theories of Gravity. *Phys. Rev. D* **2006**, *74*, 087504.
- [13] Sadjadi H. M. A Note on Gravitational Baryogenesis. *Phys. Rev. D* **2007**, *76*, 123507.
- [14] Lambiase G., Mohanty S. and Pizza L. Consequences of f(R)-theories of gravity on gravitational leptogenesis. *Gen. Rel. Grav.* **2013**, *45*, 1771–1785.
- [15] Fukushima M., Mizuno S. and Maeda K. i. Gravitational Baryogenesis after Anisotropic Inflation. *Phys. Rev. D* **2016** *93*, no.10, 103513.
- [16] Odintsov S. D. and Oikonomou V. K. Gauss–Bonnet gravitational baryogenesis. *Phys. Lett. B* **2016**, *760*, 259–262. doi:10.1016/j.physletb.2016.06.074 [arXiv:1607.00545 [gr-qc]].
- [17] Arbuzova E. V. and Dolgov A.D. Intrinsic problems of the gravitational baryogenesis. *Phys. Lett. B* **2017**, *769*, 171–175.
- [18] Arbuzova E. V. and Dolgov A.D. Instability of gravitational baryogenesis with fermions. *JCAP* **2017**, *06*, 001.
- [19] Ginzburg V. L., Kirzhnits D. A. and Lyubushin A. A. The role of quantum fluctuations of the gravitational fields in general relativity theory and cosmology. *Zh. Eksp. Teor. Fiz.* **1971**, *60*, 451–459.
- [20] Gurovich V. T. and Starobinsky A. A. Quantum effects and regular cosmological models. *Sov. Phys. JETP* **1979**, *50*, 844–852.
- [21] Starobinsky A. A. A New Type of Isotropic Cosmological Models Without Singularity. *Phys. Lett. B* **1980**, *91*, 99–102.

- [22] Gorbunov D. S. and Panin A. G. Free scalar dark matter candidates in R^2 -inflation: the light, the heavy and the superheavy. *Phys. Lett. B* **2012** 718, 15–20.