# THE HORNICH-HLAWKA FUNCTIONAL INEQUALITY FOR FUNCTIONS WITH POSITIVE DIFFERENCES

CONSTANTIN P. NICULESCU AND SUVRIT SRA

Version 2

ABSTRACT. We analyze the role played by *n*-convexity for the fulfillment of a series of linear functional inequalities that extend the Hornich-Hlawka functional inequality,  $f(x) + f(y) + f(z) + f(x + y + z) \ge f(x + y) + f(y + z) + f(z + x) + f(0)$ , including extensions to the case of positive operators.

#### 1. INTRODUCTION

Many noteworthy inequalities are related to the following problem:

**Problem 1.** Suppose that S is an abelian additive semigroup with neutral element 0, f a function defined on S and taking values in an ordered vector space E (or in its positive cone  $E_+$ ). For  $n \ge 2$ , find the linear inequalities relating

$$\sum_{i=1}^{n} f(x_i), \ \sum_{1 \le i < j \le n} f(x_i + x_j), \dots, \ f(\sum_{i=1}^{n} x_i)$$

for all  $x_1, \ldots, x_n \in \mathcal{S}$ .

Due to its many ramifications, this problem is still the subject of intense activity, and the present paper reports some new results in this direction, under the umbrella of the *Hornich-Hlawka functional inequality*. Specifically, we are interested in studying conditions under which a continuous function  $f : S \to \mathbb{R}_+$  satisfies

(1.1) 
$$f(x) + f(y) + f(z) + f(x+y+z) \ge f(x+y) + f(y+z) + f(z+x)$$

for all  $x, y, z \in S$ . When f is a real-valued function it is usual to replace (1.1) by

$$(1.2) \ f(x) + f(y) + f(z) + f(x + y + z) \ge f(x + y) + f(y + z) + f(z + x) + f(0)$$

A good start for understanding the Hornich-Hlawka functional inequality is provided by the following elementary (but powerful) inequality:

(1.3) 
$$|x| + |y| + |z| + |x + y + z| \ge |x + y| + |y + z| + |z + x|$$
 for all  $x, y, z \in \mathbb{R}$ .

As was noticed by Levi [17], every piecewise linear inequality like (1.3) remains true when the real variables x, y, z are replaced by arbitrary vectors  $\boldsymbol{x}, \boldsymbol{y}, \boldsymbol{z}$  in  $\mathbb{R}^N$  and the absolute value function is replaced by the Euclidean norm,

$$(1.4) \|x\| + \|y\| + \|z\| + \|x + y + z\| \ge \|x + y\| + \|y + z\| + \|z + x\|.$$

Date: January 19, 2023.

<sup>2000</sup> Mathematics Subject Classification. Primary 26B25; Secondary 26B35, 26D15, 26A48, 26A51.

*Key words and phrases.* Hornich-Hlawka functional inequality, completely monotone functions, function with positive differences, higher order convexity, positive (semi)definite matrix.

Inequality (1.4) is what is nowadays known as the *Hornich-Hlawka inequality*. See the paper of Hornich [14], which includes the marvelous argument of Hlawka, based on the triangle inequality and an identity (due to Fréchet [10]) which characterizes inner product spaces.

Using a standard technique, one can easily infer from (1.3) that the Hornich-Hlawka inequality (1.4) also works for all Lebesgue spaces  $L^1(\mu)$ , and so also for all spaces that can be embedded linearly and isometrically into an  $L^1(\mu)$ . The latter comment includes all Lebesgue spaces  $L^p(\mu)$  with  $p \in [1,2]$ )—see Lindenstrauss and Pełczyński [19].

In 1946, Popoviciu [32] proved that every continuous function  $f : \mathbb{R}_+ \to \mathbb{R}$  that vanishes at the origin and admits a nondecreasing derivative of second order on  $(0, \infty)$ , verifies the Hornich-Hlawka functional inequality (1.1). One can easily put Popoviciu's result in full generality by showing that actually all continuous 3-convex functions on  $\mathbb{R}_+$  taking values in an ordered Banach space verify inequality (1.2). This fact and its analogue in the case of continuous *n*-convex functions,

$$f(\sum_{i=1}^{n} x_i) - \sum_{1 \le i_1 < \dots < i_{n-1} \le n} f(x_{i_1} + \dots + x_{i_{n-1}}) + \sum_{1 \le i_1 < \dots < i_{n-2} \le n} f(x_{i_1} + \dots + x_{i_{n-2}}) - \dots + (-1)^{n-1} \sum_{i=1}^{n} f(x_i) \ge f(0),$$

will be the subject of Section 3.

Close to the above inequality is the characterization of the property of *n*-convexity via differences  $(\Delta_h f)(x) = f(x+h) - f(x)$ , rather than via divided differences as is usual. See Theorem 5, which expresses the identity of the class of continuous *n*-convex functions with the class of continuous functions having positive differences of order *n* in the sense that  $\Delta_{x_1}\Delta_{x_2}\cdots\Delta_{x_n}f(x) \geq 0$ . The connection with Popoviciu's inequality is evident when *n* is an odd integer because the condition  $\Delta_{x_1}\Delta_{x_2}\cdots\Delta_{x_n}f(x) \geq 0$ , simply means the introduction of a new variable in Popoviciu's inequality as follows:

$$\sum_{i=1}^{n} f(x_i + x) - \sum_{1 \le i < j \le n} f(x_i + x_j + x) + \sum_{1 \le i < j < k \le n} f(x_i + x_j + x_k + x) - \dots + (-1)^{n-1} f(x_1 + \dots + x_n + x) \ge f(x).$$

It is worth noticing that the functions  $f : \mathbb{R}_+ \to \mathbb{R}_+$  that have positive differences of any order are precisely the absolutely monotonic functions in the terminology of Bernstein [4]. Leaving the elegant framework of analysis on intervals one easily discovers that *n*-convexity and the property of having positive differences of order *n* are different concepts. This idea is detailed at the end of Section 3.

Two important classes of functions that mix a string of properties of *n*-convexity are those of completely monotone functions and of Bernstein functions. See Section 2 for their definitions and some examples. Sendov and Zitikis [36] prove that these functions verify inequalities of the form

(1.5) 
$$\sum_{i=1}^{n} f(x_i) - \sum_{1 \le i < j \le n} f(x_i + x_j) + \dots + (-1)^{n-1} f\left(\sum_{i=1}^{n} x_i\right) \ge 0,$$

for all  $x_1, \ldots, x_n$  in  $\mathbb{R}_+$  and  $n \ge 1$ . Their proof combines the classical integral representation theorems (respectively the Bernstein theorem and the Lévy–Khintchine representation theorem) with some probabilistic considerations. In Section 4 we extend this result as a double inequality that holds for completely monotone functions defined on cones. Combining this result with [35, Theorem 1.3 (a)], we then show that the function  $f(X) = (\det X)^{-\rho}$  (defined on  $N \times N$ -real symmetric positive definite matrices) also verifies the whole string of inequalities (1.5) if  $\rho \in \{0, 1/2, 1, 3/2, ...\} \cup [(N-1)/2, \infty).$ 

Section 5 considers functions defined on cones and having positive differences of a certain order n > 0. A surprising result is Theorem 9, which shows that the function det has positive differences of any order (though it is not completely monotonic). Probably the same happens for other immanants function (like the permanents), but we were able to prove only the positivity of differences of order 3. [TODO TODO].

For the reader's convenience, background on higher order convexity and the theory of ordered Banach spaces is summarized in Section 2.

## 2. Preliminaries

The study of higher order convexity was initiated by Hopf [13] and Popoviciu [29, 31], who defined it in terms of divided differences of a function. Assuming f a real-valued function defined on a real interval I, the divided differences of order  $0, 1, \ldots, n$  associated to a family  $x_0, x_1, \ldots, x_n$  of n+1 distinct points are respectively defined by the formulas

-

$$[x_0; f] = f(x_0)$$

$$[x_0, x_1; f] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$
...
$$[x_0, x_1, ..., x_n; f] = \frac{[x_1, x_2, ..., x_n; f] - [x_0, x_1, ..., x_{n-1}; f]}{x_n - x_0}$$

$$= \sum_{j=0}^n \frac{f(x_j)}{\prod_{k \neq j} (x_j - x_k)}.$$

Notice that all these divided differences are invariant to permutations of the points  $x_0, x_1, ..., x_n$ . As a consequence, we may always assume that  $x_0 < x_1 < \cdots < x_n$ .

A function f is called *n*-convex (respectively *n*-concave) if all divided differences  $[x_0, x_1, \ldots, x_n; f]$  are nonnegative (respectively nonpositive). In particular, 0-convex functions are precisely the nonnegative functions, 1-convex functions the nondecreasing ones, while 2-convex functions are simply the usual convex functions.

If f is n times differentiable, then a repeated application of Lagrange's mean value theorem yields the existence of a point  $\xi \in (\min_k x_k, \max_k x_k)$  such that

$$[x_0, x_1, ..., x_n; f] = \frac{f^{(n)}(\xi)}{n!}.$$

As a consequence, one obtains the following practical criterion of n-convexity.

**Lemma 1.** Every continuous function f defined on an interval I which is n times differentiable on the interior of I is n-convex provided that  $f^{(n)} \ge 0$ .

A big source of convex functions of higher order is provided by the Bernstein functions and the completely monotone functions. Recall that a function  $f:(0,\infty)\to$  $\mathbb{R}_+$  is a *Bernstein function* if it is infinitely differentiable and verifies the condition

$$(-1)^{n+1} f^{(n)}(x) \ge 0$$
 for all  $x > 0$  and  $n \ge 1$ ;

while, the function f is completely monotone if instead

$$(-1)^n f^{(n)}(x) \ge 0$$
 for all  $x > 0$  and  $n \ge 0$ .

By definition, a function  $f : [0, \infty) \to \mathbb{R}_+$  is a *Bernstein function* (respectively a *completely monotone* function) if it is continuous and its restriction of to  $(0, \infty)$ has the respective property.

Every Bernstein function is (2n + 1)-convex and every completely monotone function is 2n-convex for every  $n \ge 0$ .

If  $f: [0, \infty) \to \mathbb{R}_+$  is a Bernstein function then so is f - f(0); if f is a completely monotone function then f(0) - f is a Bernstein function. Some simple examples of Bernstein functions are

$$x/(x+1), \ 1-e^{-\alpha x}$$
 (for  $\alpha > 0$ ),  
 $\log(1+x), \ (x-1)/\log x \text{ and } x^{\alpha}$  (for  $0 < \alpha \le 1$ ).

A nice account of the two aforementioned classes of functions is offered by the authoritative monograph of Schilling, Song and Vondraček [34].

Besides the five examples mentioned above some other examples of 3-convex functions on  $\mathbb{R}_+$  are  $x^{\alpha}$  (for  $\alpha \in (0,1] \cup [2,\infty)$ ),  $-x^2 + \sqrt{x}$ ,  $-x \log x$ , sinh, cosh,  $-\log(\Gamma(x))$  etc.

The function  $1 - (x - 3) + \frac{(x-3)^3}{6}$  is continuous and 3-convex on  $\mathbb{R}_+$  but not *n*-convex for any  $n \in \{0, 1, 2\}$ .

The polynomials with positive coefficients and the exponential are *n*-convex for every  $n \ge 0$ .

All polynomials of degree less than or equal to 2 are both 3-convex and 3-concave.

The following approximation theorem due to Popoviciu [30] (see also [11, Theorem 1.3.1 (i), pg. 20]) allows us to reduce reasoning with *n*-convex functions to the case where they are also differentiable.

**Theorem 1** (Popoviciu's approximation theorem). If a continuous function  $f : [0,1] \to \mathbb{R}$  is k-convex, then so are the Bernstein polynomials associated to it,

$$B_n(f)(x) = \sum_{i=0}^n \binom{n}{i} x^i (1-x)^{n-i} f\left(\frac{i}{n}\right).$$

Moreover, by the well-known property of simultaneous uniform approximation of a function and its derivatives by Bernstein polynomials and their derivatives, it follows that  $B_n(f)$  and any derivative (of any order) of it, converge uniformly to f and to its derivatives, correspondingly.

Using a change of variable, one can easily see that the approximation theorem extends to functions defined on compact intervals [a, b] with a < b.

**Lemma 2.** (i) The composition of two continuous functions that are increasing, concave or 3-convex is a function of the same nature.

(ii) If  $f : \mathbb{R}_+ \to \mathbb{R}_+$  is a continuous 3-convex function which is also nondecreasing and concave, then the same properties hold for  $f^{\alpha}$  if  $\alpha \in (0, 1]$ .

*Proof.* According to Theorem 1, we may reduce the proof to the case where the involved functions are also of class  $C^3$ . In this case the proof can be completed by computing the sign of the derivatives of order 1, 2 and 3.

The n-convex functions taking values in an ordered Banach space can be introduced in the same manner as real-valued n-convex functions by using divided differences. We recall useful definitions below.

Recall that an ordered Banach space is any Banach space E endowed with the ordering  $\leq$  associated to a closed convex cone  $E_+$  via the formula

$$x \leq y$$
 if and only if  $y - x \in E_+$ 

such that

$$E = E_+ - E_+, \quad (-E_+) \cap E_+ = \{0\},\$$

and

$$0 \le x \le y$$
 in E implies  $||x|| \le ||y||$ .

The basic facts concerning the theory of ordered Banach spaces are made available by the book of Schaefer and Wolff [33]. See [25] for a short overview centered on two important particular cases:  $\mathbb{R}^n$ , the *n*-dimensional Euclidean space endowed with the coordinate-wise ordering, and  $\operatorname{Sym}(n, \mathbb{R})$  the ordered Banach space of all  $n \times n$  symmetric matrices with real coefficients endowed with the operator norm

$$|A|| = \sup_{\|x\| \le 1} |\langle Ax, x \rangle|,$$

and the Löwner (partial) ordering,

 $A \leq B$  if and only if  $\langle A\boldsymbol{x}, \boldsymbol{x} \rangle \leq \langle B\boldsymbol{x}, \boldsymbol{x} \rangle$  for all  $\boldsymbol{x} \in \mathbb{R}^n$ .

Here the operator norm can be replaced by any Schatten norm, in particular with the *Frobenius norm*,

$$||A||_F = \left(\sum_{i=1}^N \sum_{j=1}^N a_{ij}^2\right)^{1/2}$$

The Frobenius norm is associated to the trace inner product

$$\langle A, B \rangle = \operatorname{trace}(AB).$$

The positive cone of  $\mathbb{R}^n$  is the first orthant  $\mathbb{R}^n_+$ , while the positive cone of  $\operatorname{Sym}(n,\mathbb{R})$  is the set  $\operatorname{Sym}^+(n,\mathbb{R})$  consisting of all positive semi-definite matrices. We denote by  $\mathbb{R}^n_{++}$  and  $\operatorname{Sym}^{++}(n,\mathbb{R})$  respectively the interior of  $\mathbb{R}^n_+$  and  $\operatorname{Sym}^+(n,\mathbb{R})$ .

**Remark 1.** Much of the study of vector-valued convex functions can be reduced to that of real-valued functions. Indeed, in any ordered Banach space E, any inequality of the form  $u \leq v$  is equivalent to  $x^*(u) \leq x^*(v)$  for all  $x^* \in E^*_+$ .

As a consequence, a function  $f : I \to E$  is respectively nondecreasing, convex or n-convex if and only if  $x^* \circ f$  has this property whenever  $x^* \in E^*$  is a positive functional. For  $E = \mathbb{R}^n$ , this inequality reduces to the components of f.

Combining Remark 1 with Lemma 1 one obtains the following practical test of 3-convexity for vector-valued differentiable functions:

**Theorem 2.** Suppose that f is a continuous function defined on an interval I and taking values in an ordered Banach space E. If f is three times differentiable on the interior of I and  $f''' \ge 0$ , then f is a 3-convex function.

An example illustrating Theorem 2 is provided by the function

$$f : \mathbb{R}_+ \to \operatorname{Sym}(n, \mathbb{R}), \quad f(t) = -e^{-tA},$$

associated to a positive semi-definite matrix  $A \in \text{Sym}(n, \mathbb{R})$ . This function is of class  $C^{\infty}$  and its first three derivatives are given by the formulas

$$f'(t) = Ae^{-tA}, \quad f''(t) = -A^2e^{-tA}, \quad f'''(t) = A^3e^{-tA}.$$

Thus f is nondecreasing, concave and 3-convex (according to the ordering of  $\text{Sym}(n, \mathbb{R})$ ). The matrix  $A^3 e^{-tA}$  is positive semidefinite since the product of commuting positive semi-definite matrices is also positive semidefinite.

## 3. The functional inequality of Popoviciu

Popoviciu [32] published in 1946 a short note on a functional inequality that we restate here in a slightly more general form.

**Theorem 3.** Suppose that E is an ordered Banach space and  $f : [0, A] \to E$  is a continuous n-convex function  $(n \ge 1)$ . Then

$$f(\sum_{i=1}^{n} x_i) - \sum_{1 \le i_1 < \dots < i_{n-1} \le n} f(x_{i_1} + \dots + x_{i_{n-1}}) + \sum_{1 \le i_1 < \dots < i_{n-2} \le n} f(x_{i_1} + \dots + x_{i_{n-2}}) - \dots + (-1)^{n-1} \sum_{i=1}^{n} f(x_i) \ge (-1)^{n-1} f(0),$$
  
for all  $x_1, x_2, \dots, x_n \ge 0$  with  $\sum_{i=1}^{n} x_i \le A.$ 

Notice that this inequality can be reformulated as

$$\sum_{\varepsilon_1,\ldots,\varepsilon_n\in\{0,1\}} (-1)^{n-(\varepsilon_1+\cdots+\varepsilon_n)} f(\varepsilon_1 x_1+\cdots+\varepsilon_n x_n) \ge 0.$$

Popoviciu supplied the details only in the case n = 3, for functions f that vanish at the origin and admit a nondecreasing second order derivative.

The proof of Theorem 3 is by induction, starting with the following instance of Hardy-Littlewood-Pólya's majorization inequality (see [26, Theorem 4.1.3, pg. 186]):

**Lemma 3.** If  $g : [a,b] \to \mathbb{R}$  is a continuous convex function and c and d are two points in [a,b] such that a + b = c + d, then

$$g(c) + g(d) \le g(a) + g(b).$$

**Proof of Theorem 3.** The case n = 1 is trivial since 1-convexity is equivalent to the fact that f is nondecreasing. For n = 2 the inequality under attention reads as

$$f(x_1) + f(x_2) \le f(x_1 + x_2) + f(0),$$

which follows from Lemma 3. Suppose thus that the statement of Theorem 3 holds for all continuous *n*-convex functions and all  $x_1, x_2, ..., x_n \ge 0$  with  $\sum_{i=1}^n x_i \le A$ .

Let f be a continuous (n + 1)-convex function. According to Remark 1 we may assume that f is real-valued, while Popoviciu's approximation theorem (Theorem 1) allows us to restrict ourselves functions of class  $C^1$ . Then f' is continuous and nconvex and the same is true for the function  $\varphi(x) = f'(x_1 + x)$ . According to the induction hypothesis, if  $x_1, ..., x_n, x_{n+1} \ge 0$  and  $x_1 + \cdots + x_{n+1} \le A$ , we have

$$f'(\sum_{i=1}^{n+1} x_i) - \sum_{2 \le i_1 < \dots < i_{n-1} \le n+1} f'(x_1 + x_{i_1} + \dots + x_{i_{n-1}}) + \sum_{2 \le i_1 < \dots < i_{n-2} \le n+1} f'(x_1 + x_{i_1} + \dots + x_{i_{n-2}}) - \dots + (-1)^{n-1} \sum_{2 \le j \le n+1} f'(x_1 + x_j) \ge (-1)^{n-1} f'(x_1).$$

Similar inequalities occur by permuting the variables.

Consider  $x_2, x_3, \ldots, x_{n+1}$  fixed in [0, A] and  $x_1 \ge 0$  variable such that  $x_1 + \cdots + x_{n+1} \le A$ . The function F defined by the formula

$$F(x_1) = f(\sum_{i=1}^{n+1} x_i) - \sum_{1 \le i_1 < \dots < i_n \le n+1} f(x_{i_1} + \dots + x_{i_n}) + \dots + (-1)^{n-1} \sum_{1 \le i < j \le n+1} f(x_i + x_j) + (-1)^n \sum_{i=1}^{n+1} f(x_i) + (-1)^{n+1} f(0)$$

is differentiable and, according to the induction hypothesis,

$$F'(x_1) = f'\left(\sum_{i=1}^{n+1} x_i\right) - \sum_{\substack{2 \le i_1 < \dots < i_{n+1} \le n+1 \\ + (-1)^{n-1} \sum_{\substack{2 \le j \le n+1 \\ }} f'(x_1 + x_j) + (-1)^n f'(x_1) \ge 0.$$

Therefore F is a nondecreasing function, whence  $F(x_1) \ge F(0) = 0$ . In conclusion F is a nonnegative function and the proof is done.

It is worth noticing that Popoviciu's inequality can be turned into a characterization of *n*-convexity using difference operators.

The difference operators  $\Delta_h$  (of step size  $h \ge 0$ ) can be introduced in a large category of situations including the case of functions defined on *n*-dimensional intervals or on convex cones, ordered abelian semigroups, etc. They associate to each such function f, the function  $\Delta_h f$  defined by

$$\left(\Delta_h f\right)(x) = f(x+h) - f(x),$$

for all x and h such that the right-hand side formula makes sense.

Clearly, difference operators are linear and commute with each other,

$$\Delta_{h_1} \Delta_{h_2} = \Delta_{h_2} \Delta_{h_1}$$

They also verify the following property of invariance under translation:

$$\Delta_h \left( f \circ T_a \right) = \left( \Delta_h f \right) \circ T_a,$$

where  $T_a$  is the translation defined by the formula  $T_a(x) = x + a$ .

**Lemma 4.** If n is a positive integer, then the following formula holds:

$$\Delta_{h_1}\Delta_{h_2}\cdots\Delta_{h_n}f(x) = \sum_{\varepsilon_1,\dots,\varepsilon_n\in\{0,1\}} (-1)^{n-(\varepsilon_1+\dots+\varepsilon_n)} f(x+\varepsilon_1h_1+\dots+\varepsilon_nh_n)$$

The proof is immediate, by mathematical induction.

The property of convexity of a continuous function f defined on an interval I can be characterized via the difference operators as follows:

**Theorem 4.** A continuous function  $f : I \to \mathbb{R}$  is convex if and only if the following inequality holds,

(3.1) 
$$\Delta_a \Delta_b f(x) = f(a+b+x) - f(a+x) - f(b+x) + f(x) \ge 0$$

at all interior points  $x \in I$  and all  $a, b \ge 0$  for which  $x + a + b \in I$ .

In other words, for continuous functions defined on intervals, convexity is equivalent to the property of having positive differences of second order. *Proof.* The fact that convexity implies the inequality  $\Delta_a \Delta_b f \geq 0$  is a consequence of the Hardy-Littlewood-Pólya inequality of majorization. See Lemma 3. On the other hand, for u < v arbitrarily fixed in I, choosing a = b = (v - u)/2 and x = u, we infer from (3.1) that

$$\frac{f(u) + f(v)}{2} \ge f\left(\frac{u+v}{2}\right),$$

which is equivalent to convexity since f was assumed to be continuous. See [26, Theorem 1.1.8, pg. 5].

The following result extends Theorem 4 to the case of higher-order convexity and originates from an old paper of Boas and Widder [7].

**Theorem 5.** Suppose that  $f : [0, A] \to \mathbb{R}$  is a continuous function and  $n \ge 1$  is an integer. Then the following conditions are equivalent:

(i) f is n-convex;

(ii) f has positive differences of order n in the sense that

 $\Delta_{x_1}\Delta_{x_2}\cdots\Delta_{x_n}f(t)\geq 0$ 

for all points  $t, x_1, x_2, ..., x_n \ge 0$  such that  $t + x_1 + \cdots + x_n \le A$ .

The same works if the interval [0, A] is replaced by  $\mathbb{R}_+$  and  $\mathbb{R}$ .

*Proof.* The implication  $(i) \Longrightarrow (ii)$  follows from Theorem 3, when applied to the *n*-convex function g(x) = f(x+t) - f(t). An alternative proof is made available by the paper of Boas and Widder [7].

The converse implication is immediate and is the objective of [16, Theorem 15.3.1, pg. 430] in Kuczma's book. A very short proof of the implication  $(ii) \Longrightarrow (i)$  when n = 3 can be found in [2, Proposition 1].

**Corollary 1.** Suppose that E is an ordered Banach space and  $f : [0, A] \to E$  is a continuous function. Then f is n-convex if and only if it verifies the inequality

$$\Delta_{x_1}\Delta_{x_2}\cdots\Delta_{x_n}f(x) = \sum_{\varepsilon_1,\dots,\varepsilon_n\in\{0,1\}} \left(-1\right)^{n-(\varepsilon_1+\dots+\varepsilon_n)} f\left(x+\varepsilon_1x_1+\dots+\varepsilon_nx_n\right) \ge 0$$

for all points  $x, x_1, \ldots, x_n \in [0, A]$  such that  $x + x_1 + \cdots + x_n \leq A$ .

In particular, a continuous function  $f : [0, A] \to E$  is 3-convex if and only if it verifies the inequality

$$\begin{aligned} f\left(x+t\right) + f\left(y+t\right) + f\left(z+t\right) + f\left(x+y+z+t\right) \\ &\geq f\left(x+y+t\right) + f\left(y+z+t\right) + f\left(z+x+t\right) + f(t) \end{aligned}$$

for all points  $x, y, z, t \in [0, A]$  such that  $x + y + z + t \leq A$ .

Both Theorem 3 and Theorem 5 can be applied successfully to derive a number of useful inequalities satisfied by the Bernstein functions and the completely monotonic functions on  $[0, \infty)$ . We will come back to this matter in the next section.

In higher dimensions, the equivalence between usual convexity and the property of having positive differences of second order is no anymore valid. Some simple examples are indicated in what follows. **Example 1.** Consider the case of the infinitely differentiable function

$$f(x,y) = -2(xy)^{1/2}, \quad x,y \in (0,\infty).$$

This function is convex, its Hessian being the positive semidefinite matrix

$$H = \frac{1}{2} \begin{pmatrix} x^{-3/2}y^{1/2} & -x^{-1/2}y^{-1/2} \\ -x^{-1/2}y^{-1/2} & x^{1/2}y^{-3/2} \end{pmatrix}.$$

However,  $\Phi$  fails the inequality

$$\Delta_A \Delta_B f(X) \ge 0 \quad \text{for } A, B, X \in (0, \infty) \times (0, \infty);$$

for example, choose A = (1, 2), B = (2, 1) and X near the origin.

**Example 2.** The function

$$M: \mathbb{R}^2_+ \to \mathbb{R}, \quad M(x, y) = \min\{x, y\}$$

is continuous and concave. Besides it has positive differences of second order as

 $\min\{x + s + u, y + t + v\} - \min\{x + s, y + t\}$ 

$$-\min\{x+u, y+v\} + \min\{x, y\} \ge 0,$$

for all s, t, u, v > 0. The function M proves useful in statistics as the Fréchet-Hoeffding upper bound for joint distribution functions of random variables. See Nelsen [24].

Popoviciu [29, 31] introduced the concept of higher order convexity for functions of several variables using multiple divided differences. To gain some insight, let us consider the case of a function f = f(x, y) defined on a product  $I \times J$  of intervals, and let  $x_0, x_1, \ldots, x_m$  be distinct points in I, and  $y_0, y_1, \ldots, y_n$  be distinct points in J. The divided double differences are defined via the formula

(3.2) 
$$\begin{bmatrix} x_0, x_1, \dots, x_m \\ y_0, y_1, \dots, y_n \end{bmatrix} = [x_0, x_2, \dots, x_m; [y_0, y_1, \dots, y_n; f((x, \cdot)]] \\ = [y_0, y_1, \dots, y_n; [x_0, x_1, \dots, x_m; f((\cdot, y)]].$$

Notice that this formula is invariant under the permutation of variables  $x_k$  (and also under the permutation of the variables  $y_k$ ).

Drawing a parallel to the one dimensional case, Popoviciu [29, pg. 78] calls a function  $f: I \times J \to \mathbb{R}$  convex of order (m, n) if the divided differences

$$\begin{bmatrix} x_0, & x_1, & \dots & , & x_m \\ y_0, & y_1, & \dots & , & y_n \end{bmatrix}; f$$

are nonnegative for all distinct points  $x_0, x_1, ..., x_m \in I$  and  $y_0, y_1, ..., y_n \in J$ .

Needless to say, the study of this concept of convexity implies a formidable formalism, so little progress was made since the times of Popoviciu. The only one recent contribution is [12] that studies the cases m = n = 1 and m = n = 2.

### 4. The case of completely monotone functions on cones

The theory of completely monotone functions can be easily extended to the context of several variables using convex analysis. In what follows V denotes a finite-dimensional real vector space and C an open convex cone in V with closure  $\overline{C}$ . Its dual cone is  $C^* = \{y \in E^* : \langle y, x \rangle \ge 0 \text{ for all } x \in C\}$ . The points in  $C^*$  are linear functionals that are nonnegative on  $\overline{C}$ .

**Definition 1.** A function  $f : C \to \mathbb{R}_+$  is called completely monotone if f is  $C^{\infty}$  on C and, for all integers  $k \ge 1$  and all vectors  $v_1, ..., v_k \in C$ , we have

(4.1) 
$$(-1)^k D_{v_1} \cdots D_{v_k} f(x) \ge 0 \quad \text{for all } x \in \mathcal{C}.$$

Here  $D_v$  denotes the directional derivative along the vector v.

A function  $f : \overline{C} \to \mathbb{R}_+$  is called completely monotone if it is the continuous extension of a completely monotone function on C.

When  $\mathcal{C} = (0, \infty)^n$ , the condition (4.1) means that

$$(-1)^k \frac{\partial^k f}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_k}}(x) \ge 0$$

for all  $x \in (0,\infty)^n$  and all sets of indices  $1 \le i_1 \le i_2 \le \cdots \le i_k \le n$  of arbitrary length k.

As in the case of completely monotone functions of one real variable, these functions can be obtained as Laplace transforms of Borel measures on the dual cone.

**Theorem 6.** (Bernstein-Hausdorff-Widder-Choquet theorem). Let f be a nonnegative continuous function on the open convex cone C. Then f is completely monotone if and only if it is the Laplace transform of a unique Borel measure  $\mu$  supported on the dual cone  $C^*$ , that is,

$$f(x) = \int_{\mathcal{C}^*} e^{-\langle y, x \rangle} d\mu(y) \text{ for all } x \in \mathcal{C}.$$

When f admits a continuous extension to  $\overline{\mathcal{C}}$ , the last equality works for all  $x \in \overline{\mathcal{C}}$ .

For details, see Choquet [8].

**Remark 2.** Finding the positive Borel measure  $\mu$  that makes the formula of Theorem 6 working represents a practical way for checking the complete monotonicity of f. So is the case of Riesz kernels: If  $\alpha_1, \alpha_2, ..., \alpha_N > 0$ , then

$$x_1^{-\alpha_1}x_2^{-\alpha_2}\cdots x_N^{-\alpha_N} = \int_{\mathbb{R}^N_{++}} e^{-\langle \boldsymbol{y}, \boldsymbol{x} \rangle} \frac{x_1^{-\alpha_1}x_2^{-\alpha_2}\cdots x_N^{-\alpha_N}}{\Gamma(\alpha_1)\Gamma(\alpha_2)\cdots\Gamma(\alpha_N)} \mathrm{d}\boldsymbol{y}$$

for all  $\boldsymbol{x} \in \mathbb{R}^{N}_{++}$ . See [15, Proposition 2.7]. A more subtle case is that of inverse powers of the determinant

$$f(X) = (\det X)^{-\rho}, \quad X \in \operatorname{Sym}^{++}(N, \mathbb{R}),$$

for which Scott and Sokal [35] have shown that is completely monotone if and only if  $\rho \in \{0, 1/2, 1, 3/2, ..., (N-1)/2\} \cup ((N-1)/2, \infty)$ . See also [15, Theorem 4.1]. It is worth noticing that Siegel established in 1929 the formula

$$(\det A)^{-\rho} = \int_{\text{Sym}^{++}(N,\mathbb{R})} e^{-\operatorname{trace} AX} \frac{(\det X)^{\rho} \, dX}{\pi^{n(n-1)/4} \Gamma(\rho) \Gamma(\rho - 1/2) \cdots \Gamma(\rho - (n-1)/2))}$$

for all  $A \in \text{Sym}^{++}(N, \mathbb{R})$  and  $\rho \ge (N+1)/2$ . See [38, Hilfssatz 37, pg. 585].

We next extend (and improve) a result due to Sendov and Zitikis; see [36, Theorem 4.1, pg. 76].

**Theorem 7.** Every completely monotone function  $f : \mathcal{C} \to \mathbb{R}_+$  satisfies

(4.2) 
$$\sum_{i=1}^{n} f(x_i) - \sum_{1 \le i < j \le n} f(x_i + x_j) + \dots + (-1)^{n-1} f(\sum_{i=1}^{n} x_i) \ge 0,$$

for every  $x_1, \ldots, x_n \in C$  and  $n \geq 1$ . When f admits a continuous extension to  $\overline{C}$ , then then f satisfies the double inequality

$$(4.3) \ f(0) \ge \sum_{i=1}^{n} f(x_i) - \sum_{1 \le i < j \le n} f(x_i + x_j) + \dots + (-1)^{n-1} f(\sum_{i=1}^{n} x_i) \ge 0,$$
  
for every  $x_1, \dots, x_n \in \overline{\mathcal{C}}$  and  $n \ge 1$ .

For n = 3, the first conclusion of Theorem 7 reads as

$$\sum_{i=1}^{3} f(x_i) - \sum_{1 \le i < j \le 3} f(x_i + x_j) + f(\sum_{i=1}^{3} x_i) \ge 0,$$

which is nothing but a Hornich-Hlawka type inequality.

The proof of Theorem 7 needs the following auxiliary result.

Lemma 5. We have

$$P = \sum_{i=1}^{n} e^{-\alpha_i} - \sum_{1 \le i < j \le n} e^{-(\alpha_i + \alpha_j)} + \dots + (-1)^{n+1} e^{-\sum_{i=1}^{n} \alpha_i} \ge 0$$

and

$$Q = \sum_{i=1}^{n} (1 - e^{-\alpha_i}) - \sum_{1 \le i < j \le n} (1 - e^{-(\alpha_i + \alpha_j)}) + \dots + (-1)^{n+1} (1 - e^{-\sum_{i=1}^{n} \alpha_i}) \ge 0,$$

whenever  $\alpha_1, \ldots, \alpha_n \in \mathbb{R}_+$  and  $n \geq 2$ .

Proof. Indeed,

$$S = 1 - \prod_{i=1}^{n} (1 - e^{-\alpha_i})$$
 and  $Q = \sum_{i=1}^{n} (-1)^{k+1} \binom{n}{k} - S = 1 - S.$ 

**Proof of Theorem 7.** As per to Theorem 6, f admits the integral representation

$$f(x) = \int_{\mathcal{C}^*} e^{-\langle y, x \rangle} d\mu(y) \text{ for all } x \in \mathcal{C},$$

where  $\mu$  is a Borel measure on  $C^*$ . Then, taking into account to the first assertion of Lemma 5, we have the inequality

$$0 \leq \int_{\mathcal{C}^*} \sum_{i=1}^n e^{-\langle x_i, y \rangle} - \sum_{1 \leq i < j \leq n} e^{-\langle x_i + x_j, y \rangle} + \dots + (-1)^{n-1} e^{-\langle \sum_{i=1}^n x_i, y \rangle} d\mu(y) = \sum_{i=1}^n f(x_i) - \sum_{1 \leq i < j \leq n} f(x_i + x_j) + \dots + (-1)^{n-1} f(\sum_{i=1}^n x_i).$$

The case where f is defined on  $\overline{C}$  can be settled in the same manner, using both assertions of Lemma 5.

Combining Theorem 7 with the aforementioned result of Scott and Sokal (see Remark 2) one obtains the following result:

**Corollary 2.** If 
$$\rho \in \{0, 1/2, 1, 3/2, ...\} \cup [(N-1)/2, \infty)$$
, then

$$\sum_{i=1}^{n} \det^{-\rho}(A_i) - \sum_{1 \le i < j \le n} \det^{-\rho}(A_i + A_j) + \dots + (-1)^{n-1} \det^{-\rho}(\sum_{i=1}^{n} A_i) \ge 0,$$

for every  $A_1, \ldots, A_n \in \text{Sym}^{++}(N, \mathbb{R})$  and  $n \ge 1$ .

Given Corollary 2 one may wonder whether Theorem 7 also specializes to elementary symmetric polynomials. The situation here turns out to be more subtle, and a qualified answer follows from the discussion below. Recall that for any m = 0, 1, ..., N, the *m*-th elementary symmetric polynomial of  $\boldsymbol{x} \in \mathbb{R}^N$  is defined by the formula

$$E_{m,N}(\boldsymbol{x}) = \sum_{1 \le i_1 < \dots < i_m \le N} x_{i_1} \dots x_{i_m}.$$

Notice that det and the functions  $E_{m,N}$  are hyperbolic polynomials in the sense of Gårding. For details concerning this notion see [26, Section 4.6]. Kozhasov, Michalek and Sturmfels provided a constructive proof for the fact that any elementary symmetric polynomial  $E_{m,n}$  admits a real exponent  $\alpha' > 0$  such that  $E_{m,n}^{-\alpha}$  is completely monotone on  $\mathbb{R}_{++}^N$  for all  $\alpha \geq \alpha'$ —see [15, Theorem 6.4].

**Remark 3.** The restriction of  $e^{-x}$  to  $\mathbb{R}_+$  is a completely monotone function and thus it verifies the inequality

$$e^{-x_1} + e^{-x_2} + e^{-x_3} + e^{-(x_1+x_2+x_3)} \ge e^{-(x_1+x_2)} + e^{-(x_2+x_3)} + e^{-(x_3+x_1)}$$

However there  $x_1, x_2, x_3 > 0$  such that

$$e^{-x_1} + e^{-x_2} + e^{-x_3} + e^{-(x_1 + x_2 + x_3)} \not\geqslant e^{-(x_1 + x_2)} + e^{-(x_2 + x_3)} + e^{-(x_3 + x_1)} + e^0.$$

This shows that the inequality

$$\sum_{i=1}^{3} f(x_i) - \sum_{1 \le i < j \le 3} f(x_i + x_j) + f(\sum_{i=1}^{3} x_i) \ge f(0),$$

does not characterize the 3-convexity within the class of continuous functions.

The fact that the Bernstein functions  $f : \mathbb{R}_+ \to \mathbb{R}_+$  also verify the inequalities (4.2) follows from Lemma 5 and the Lévy-Khintchine theorem (Theorem 3.2, p. 15 of [34]), which asserts that each such function admits the integral representation

$$f(x) = a + bx + \int_0^\infty (1 - e^{-xt}) d\mu(t)$$

for some constants  $a, b \ge 0$  and a positive measure  $\mu$  on  $[0, \infty)$  such that

$$\int_0^\infty \min\left\{1,t\right\} \mathrm{d}\mu(t) < \infty.$$

As far as we know, no attempt was made to extend the theory of Bernstein functions to the framework of functions defined on cones.

### 5. Functions with positive differences on cones

The difference operators  $\Delta_h f: x \to \Delta_h f(x) = f(x+h) - f(x)$  are well defined in the case of functions f defined on a convex cone C. Such a function is said to be a function with positive differences of order  $n \ (n \ge 1)$  if

$$\Delta_{h_1}\Delta_{h_2}\cdots\Delta_{h_n}f(x) \ge 0 \text{ for all } x, h_1, h_2, \dots, h_n \in \mathcal{C}.$$

For convenience, we say that f has positive differences of order 0 if  $f \ge 0$ .

In the literature, the concept of *n*-absolute monotonicity is used with the meaning that the function under attention has *positive differences of order* k for all  $k \in \{0, 1, ..., n\}$ .

**Theorem 8.** If  $f : C \to F_+$  is a completely monotonic function, then f has positive differences of any even order k = 0, 2, 4, ...; under the same hypothesis, -f has positive differences of any odd order k = 1, 3, 5, ....

This follows easily from the characterization of monotonicity in terms of Gâteaux differentiability (as was established by Amann [1], Proposition 3.2, p. 184):

**Lemma 6.** Suppose that E and F are two ordered Banach spaces, C is a convex subset of E with nonempty interior int C and  $\Phi : C \to F$  is a function, continuous on C and Gâteaux differentiable on int C. Then  $\Phi$  is monotone nondecreasing on C if and only if

$$D\Phi(a)[v] = \frac{\Phi(a+tv) - \Phi(a)}{t} \ge 0$$

for all points  $a \in int \mathcal{C}$  and all vectors  $v \in E_+$ .

**Example 3.** Every function of the form

$$\Phi(oldsymbol{x}) = f\left(\langle oldsymbol{x}, oldsymbol{w}
angle
ight), \quad oldsymbol{x} \in \mathbb{R}^N_+,$$

associated to a continuous n-convex function  $f : \mathbb{R}_+ \to \mathbb{R}$  and a vector  $\boldsymbol{w} \in \mathbb{R}^N_+$  has positive differences of order n.

The proof is straightforward, as evident from the case n = 3. Indeed, in this case, for every  $\mathbf{x}, \mathbf{y}, \mathbf{z}, \mathbf{t} \in \mathbb{R}^n_+$  we have

$$\begin{split} \Phi\left(\boldsymbol{x}+\boldsymbol{t}\right) + \Phi\left(\boldsymbol{y}+\boldsymbol{t}\right) + \Phi\left(\boldsymbol{z}+\boldsymbol{t}\right) + \Phi\left(\boldsymbol{x}+\boldsymbol{y}+\boldsymbol{z}+\boldsymbol{t}\right) \\ &-\Phi\left(\boldsymbol{x}+\boldsymbol{y}+\boldsymbol{t}\right) - \Phi\left(\boldsymbol{y}+\boldsymbol{z}+\boldsymbol{t}\right) - \Phi\left(\boldsymbol{z}+\boldsymbol{x}+\boldsymbol{t}\right) - \Phi(\boldsymbol{t}) \\ &= f\left(\langle \boldsymbol{x}+\boldsymbol{t},\boldsymbol{v}\rangle\right) + f\left(\langle \boldsymbol{y}+\boldsymbol{t},\boldsymbol{v}\rangle\right) + f\left(\langle \boldsymbol{z}+\boldsymbol{t},\boldsymbol{v}\rangle\right) + f\left(\langle \boldsymbol{x}+\boldsymbol{y}+\boldsymbol{z}+\boldsymbol{t},\boldsymbol{v}\rangle\right) \\ &- f\left(\langle \boldsymbol{x}+\boldsymbol{y}+\boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{y}+\boldsymbol{z}+\boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{z}+\boldsymbol{x}+\boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{t},\boldsymbol{v}\rangle\right) \\ &= f\left(\langle \boldsymbol{x},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) + f\left(\langle \boldsymbol{y},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) + f\left(\langle \boldsymbol{z},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) \\ &+ f\left(\langle \boldsymbol{x}+\boldsymbol{y}+\boldsymbol{z},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{x}+\boldsymbol{y},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{y}+\boldsymbol{z},\boldsymbol{v}\rangle + \boldsymbol{t},\boldsymbol{v}\rangle\right) \\ &- f\left(\langle \boldsymbol{z}+\boldsymbol{x},\boldsymbol{v}\rangle + \langle \boldsymbol{t},\boldsymbol{v}\rangle\right) - f\left(\langle \boldsymbol{t},\boldsymbol{v}\rangle\right) \geq 0. \end{split}$$

A variant of this example is provided by the map

 $\Psi(A) = f\left(\operatorname{trace}(AW)\right), \quad A \in \operatorname{Sym}^+(n, \mathbb{R}),$ 

associated to a continuous n-convex function  $f : \mathbb{R}_+ \to \mathbb{R}$  and to an operator  $W \in \mathbb{R}^N_+$ . The ambient Hilbert space in this case is  $Sym(n, \mathbb{R})$ , endowed with the Frobenius norm and Löwner ordering.

Linear algebra offers many examples of functions that have positive differences of any order  $n \geq 0$ . So is the case of the determinant function det, restricted to  $\operatorname{Sym}^+(N,\mathbb{R})$  (though det it is not completely monotonic).

Clearly,  $det(X) \ge 0$  and since det is monotonic on the semidefinite matrices,

$$(\Delta_A \det)(X) = \det(X + A) - \det(X) \ge 0,$$

for all  $A, X \in \text{Sym}^+(N, \mathbb{R})$ . Also simple is the fact that

 $\left(\Delta_A \left(\Delta_B \det\right)\right)(X) = \det\left(A + B + X\right) - \det\left(A + X\right) - \det(B + X) + \det(X) \ge 0$ 

for all  $A, B, X \in \text{Sym}^+(N, \mathbb{R})$ . This inequality is mentioned in [42, Problem 36, pg. 215]. For X = 0 it reduces to the property of superaditivity of the function det,

$$\det(A+B) \ge \det(A) + \det(B).$$

Our next goal is to show the more challenging third-order inequality

 $\Delta_A \left( \Delta_B (\Delta_C \det) \right) \ge 0.$ 

We require some preparatory lemmas before stating our proof.

**Lemma 7.** Let  $A, B, C \in \text{Sym}^+(N, \mathbb{R})$ , and let  $e_k(X)$  denote the k-th elementary symmetric function of a matrix X ( $0 \le k \le N$ ). Then,

$$e_k(A) + e_k(B) + e_k(C) + e_k(A + B + C) \ge e_k(A + B) + e_k(B + C) + e_k(C + A).$$

*Proof.* Recall that for an  $N \times N$  matrix X, we have  $e_k(X) = \operatorname{tr}(\wedge^k X)$ , where  $\wedge$  denotes the anti-symmetric tensor product; moreover,  $\wedge^k(X) = P_k^*(\otimes^k X)P_k$  for a suitable projection matrix  $P_k$ —see e.g., [5, pg. 18] for these facts. Using  $P_k$  in Theorem 2.1 in [3], the claimed inequality for  $e_k$  follows.

**Lemma 8.** If  $A, B, C, X \in \text{Sym}^+(N, \mathbb{R})$ , then

$$det (A + X) + det (B + X) + det (C + X) + det (A + B + C + X)$$
  

$$\geq det (A + B + X) + det (B + C + X) + det (C + A + X) + det X.$$

For X = 0 we get the determinantal Hornich-Hlawka inequality

 $\det A + \det B + \det C + \det(A + B + C)$ 

 $\geq \det (A+B) + \det (B+C) + \det (C+A),$ 

first noticed by Lin [18], who provided a proof based on eigenvalue majorization.

**Proof.** Without loss of generality we may assume that X is invertible (for example, replace X by  $X + \varepsilon I$  if necessary). Then

$$\det(A+X) = \det(X) \det(X^{-1/2}AX^{-1/2}+I),$$

where I is the identity matrix; the inequality under attention is then equivalent to

$$det(A+I) + det(B+I) + det(C+I) + det(A+B+C+I)$$
  

$$\geq det(A+B+I) + det(B+C+I) + det(C+A+I) + 1.$$

Consider the function  $f(A) := \det(A + I) - 1$ . Then the above inequality becomes (5.1)  $f(A) + f(B) + f(C) + f(A + B + C) \ge f(A + B) + f(B + C) + f(C + A)$ . Now recall the well-known expansion

$$\det(A+I) = \sum_{k=0}^{N} e_k(A),$$

where  $e_k(\cdot)$  denotes the k-th elementary symmetric polynomial  $(e_0 = 1, e_1 = \text{tr} \dots, e_N = \text{det})$ —see [22, Theorem 7.1.2, pg. 197] Thus,  $f(A) = \sum_{k=1}^{N} e_k(A)$ , and inequality (5.1) becomes

$$\sum_{k=1}^{N} [e_k(A) + e_k(B) + e_k(C) + e_k(A + B + C)]$$
  

$$\geq \sum_{k=1}^{N} [e_k(A + B) + e_k(B + C) + e_k(C + A)].$$

The proof ends by applying Lemma 7 for each  $k \in \{1, ..., N\}$ .

A similar argument using [3, Corollary 3.4] yields that the function det has positive differences of any order.

Theorem 9. We have

$$\Delta_{A_1} (\Delta_{A_2}(...(\Delta_{A_n} \det)))(X)$$
  
=  $\sum_{k=0}^n (-1)^{k+1} \sum_{1 \le i_1 < i_2 < \dots < i_k \le n} \det(A_{i_1} + \dots + A_{i_k} + X) \ge 0,$ 

whenever  $A_1, \ldots, A_n, X \in \text{Sym}^+(N, \mathbb{R})$  and  $n \ge 1$ . In other words, the restriction of the det function to  $\text{Sym}^+(N, \mathbb{R})$  has positive differences of any order  $n \ge 0$ .

Lemma 8 can be extended to a larger class of matrix functions, that of immanants. The *immanant function*  $d_{\chi}^{G}$ , associated to a subgroup G of the symmetric group  $S_N$  of N letters and to an irreducible character  $\chi$  of G, is defined via the formula

$$d_{\chi}^{G}(A) = \sum_{\sigma \in G} \chi(\sigma) \prod_{i=1}^{N} a_{i \sigma(i)}, \ A \in \operatorname{Sym}^{+}(N, \mathbb{R}).$$

When  $G = \mathfrak{G}_m$  and  $\chi(\sigma) = \operatorname{sgn} \sigma$  we have  $d_{\chi}^G(A) = \det A$ , while for  $\chi(\sigma) \equiv 1$  we obtain the permanent of A.

The following two inequalities

$$\begin{aligned} & d_{\chi}^G \left(A + X\right) - d_{\chi}^G(X) \geq 0 \\ & d_{\chi}^G \left(A + B + X\right) - d_{\chi}^G \left(A + X\right) - d_{\chi}^G (B + X) + d_{\chi}^G(X) \geq 0 \end{aligned}$$

occur for all  $A, B, X \in \text{Sym}^+(N, \mathbb{R})$ . See respectively Merris [21], p. 228 and Paksoy, Turkmen and Zhang [27]. The fact that  $d_{\chi}^G$  has positive differences of third order on  $\text{Sym}^+(N, \mathbb{R})$  makes the objective of the following result:

**Theorem 10.** If  $A, B, C, X \in \text{Sym}^+(N, \mathbb{R})$ , then

$$\begin{split} d_{\chi}^{G}(A+X) + d_{\chi}^{G}(B+X) + d_{\chi}^{G}(C+X) + d_{\chi}^{G}(A+B+C+X) \\ &\geq d_{\chi}^{G}(A+B+X) + d_{\chi}^{G}(B+C+X) + d_{\chi}^{G}(C+A+X) + d_{\chi}^{G}(X). \end{split}$$

Using arguments from multilinear algebra one can show that there exists a matrix  $Z_{G,\chi}$  such that

(5.2) 
$$d_{\chi}^{G}(X) = Z_{G,\chi}^{*}\left(\otimes^{N} X\right) Z_{G,\chi}.$$

See [20], p. 126. The representation (5.2) turns the assertion of Theorem 10 into an immediate consequence of the following general result:

**Theorem 11.** (Operator Hornich-Hlawka inequality). If  $A, B, C, X \in \text{Sym}^+(N, \mathbb{R})$ and  $p \ge 1$  an integer, then

(5.3)

$$\otimes^{p}(A+X) + \otimes^{p}(B+X) + \otimes^{p}(C+X) + \otimes^{p}(A+B+C+X)$$
$$\geq \otimes^{p}(A+B+X) + \otimes^{p}(B+C+X) + \otimes^{p}(C+A+X) + \otimes^{p}X,$$

in the Löwner order. Here  $\otimes$  denotes the usual tensor product.

The proof of Theorem 11 follows the proof structure of [3, Theorem 2.1], but due to the additional X term in (5.3) it turns out to be more intricate and requires some preparation. We start by introducing the following convenient notation:

(5.4) 
$$X^j \equiv \otimes^j X = X^{\otimes j}, \text{ for } j \ge 1, \text{ and } X^0 = 1 \in \mathbb{N}.$$

The slight abuse of notation  $X^0 = 1$  will be helpful in simplifying the presentation.

**Lemma 9.** Let  $k, l \ge 0$  be integers and  $X, Y, Z, V \in \text{Sym}^+(N, \mathbb{R})$ . Then,

$$X^{k} \otimes V \otimes X^{l} + (X + Y + Z)^{k} \otimes V \otimes (X + Y + Z)^{l}$$
  

$$\geq (X + Y)^{k} \otimes V \otimes (X + Y)^{l} + (X + Z)^{k} \otimes V \otimes (X + Z)^{l},$$

in the sense of Löwner order.

*Proof.* The proof is by induction on k and l; we provide the argument for k, which holds essentially unchanged for l. This approach suffices since for a fixed but arbitrary  $l \ge 0$  we prove that the result holds for all  $k \ge 0$ ; similarly, for an arbitrarily fixed  $k \ge 0$  the result holds for all  $l \ge 0$ .

For the base case of induction, suppose k = 0. Then the inequality under attention reduces to the following one:

(5.5) 
$$V \otimes X^{l} + V \otimes (X + Y + Z)^{l} \ge V \otimes (X + Y)^{l} + V \otimes (X + Z)^{l}.$$

We know from [3, Theorem 2.1] that

(5.6) 
$$X^{l} + (X + Y + Z)^{l} \ge (X + Y)^{l} + (X + Z)^{l}.$$

Since tensor product preserves inequalities, taking tensor product with V on both sides of the inequality (5.6) one immediately get (5.5). Assume therefore that inequality in the the statement of Lemma 9 holds for a fixed l and some k > 0. Then, consider

$$(X + Y + Z)^{k+1} \otimes V \otimes (X + Y + Z)^{l}$$
  
=  $(X + Y + Z) \otimes [(X + Y + Z)^{k} \otimes V \otimes (X + Y + Z)^{l}]$   
 $\geq (X + Y + Z) \otimes [(X + Y)^{k} \otimes V \otimes (X + Y)^{l}$   
 $+ (X + Z)^{k} \otimes V \otimes (X + Z)^{l} - X^{k} \otimes V \otimes X^{l}],$   
=  $(X + Y)^{k+1} \otimes V \otimes (X + Y)^{l} + (X + Z)^{k+1} \otimes V \otimes (X + Z)^{l} - X^{k+1} \otimes V \otimes X^{l} + \mathcal{T},$ 

where the inequality follows from the induction hypothesis and the elementary monotonicity properties of the tensor product. It remains to show that the term

$$\mathcal{T} = Z \otimes (X+Y)^k \otimes V \otimes (X+Y)^l + Y \otimes (X+Z)^k \otimes V \otimes (X+Z)^l - (Y+Z) \otimes X^k \otimes V \otimes X^k \otimes X^k \otimes V \otimes X^k \otimes X$$

is a nonnegative operator. Since  $X, Y, Z \ge 0$ , it follows that  $X + Y \ge X$  and  $X + Z \ge X$ . Thus, the positive terms in  $\mathcal{T}$  attached to Y and Z are clearly bigger than the respective negative terms, whence  $\mathcal{T} \ge 0$ . Inductively, we can conclude that for fixed l, the inequality in the statement of Lemma 9 holds for all  $k \ge 0$ . Applying a similar argument for l, we conclude that this inequality works in full generality.

We are now in a position to detail the proof of Theorem 11.

**Proof of Theorem 11**. Using the auxiliary function

$$f_p(Z) = (Z + X)^p - X^p,$$

the inequality of interest (5.3) can be rewritten as

$$f_p(A) + f_p(B) + f_p(C) + f_p(A + B + C) \ge f_p(A + B) + f_p(B + C) + f_p(C + A).$$

Now introduce the function

$$g_p(Z) = f_p(Z) + f_p(B) + f_p(C) + f_p(Z + B + C) - f_p(Z + B) - f_p(Z + C) - f_p(B + C).$$

We will show that  $g_p$  is monotonic (as a map from  $\text{Sym}^+(N, \mathbb{R})$  into itself, under the Löwner order). Once this monotonicity is established we can conclude that

$$g_p(A) \ge g_p(0) = 0$$
 for every  $A \in \operatorname{Sym}^+(N, \mathbb{R})$ ,

a fact equivalent to the assertion of Theorem 11.

Monotonicity of  $g_p$  follows from Lemma 6 by considering its derivative. To that end, consider the (directional) derivative of the map  $\Phi(Z) = Z^p$ :

$$D(Z^p)[V] = V \otimes Z \otimes \cdots \otimes Z + Z \otimes V \cdots \otimes Z + \cdots + Z \otimes \cdots \otimes Z \otimes V$$
$$= \sum_{j=0}^{p-1} Z^j \otimes V \otimes Z^{p-1-j},$$

whenever  $Z \in \text{Sym}^{++}(N, \mathbb{R})$  and  $V \in \text{Sym}^{+}(N, \mathbb{R})$ . See [6, Eq. (2.13), pg. 44]. Indeed, applying this formula to  $f_p(Z) = (Z + X)^p - X^p$  we obtain

$$Df_p(Z)[V] = \sum_{j=0}^{p-1} (Z+V)^j \otimes V \otimes (Z+V)^{p-1-j},$$

which in turn leads to the identity

$$Dg_p(Z)[V] = Df_p(Z)[V] + Df_p(Z + B + C)[V] - Df_p(Z + B)[V] - Df_p(Z + C)[V]$$
  
=  $\sum_{j=0}^{p-1} \left[ (Z + V)^j \otimes V \otimes (Z + V)^{p-1-j} + (Z + B + C + V)^j \otimes V \otimes (Z + B + C + V)^{p-1-j} - (Z + B + V)^j \otimes V \otimes (Z + C + V)^{p-1-j} \right].$ 

But this sum evaluates to a positive quantity, which follows from Lemma 9 upon setting  $k \leftarrow j$ ,  $l \leftarrow p - 1 - j$  and  $x \leftarrow x + j$ . Now the proof is complete.

## 6. Further comments and extensions

6.1. Multivariable case for positive operators. Given  $A_1, ..., A_n, X \in \text{Sym}^+(N, \mathbb{R})$ and  $p \in \mathbb{N}$ , let us consider the matrices

(6.1) 
$$S_0^p = \otimes^p X, \quad S_k^p = \sum_{1 \le i_1 < \dots < i_k \le n} \otimes^p (A_{i_1} + \dots + A_{i_k} + X) \ (1 \le k \le N).$$

Then one can prove the following generalization of the operator Hornich-Hlawka inequality (Theorem 11):

(6.2)  $S_n^p + S_{n-2}^p + \dots \ge S_{n-1}^p + S_{n-3}^p + \dots ,$ 

Theorem 3.3 of [3] proves inequality (6.2) for the special case X = 0. One can prove the general case by a suitable monotonicity argument as in Section 5. The details are tedious so we omit them, leaving them as an exercise for the interested reader. 6.2. Two inequalities for determinants. Serre [37] noticed that  $det^{1/2}$  viewed as a function on  $Sym^+(2,\mathbb{R})$  verifies the opposite of the Hornich-Hlawka inequality,

(6.3) 
$$\det^{1/2} A + \det^{1/2} B + \det^{1/2} C + \det^{1/2} (A + B + C) \\ \leq \det^{1/2} (A + B) + \det^{1/2} (B + C) + \det^{1/2} (C + A).$$

While in [40], the second named author proved the following Minkowski-like inequality for det  $\frac{1}{n}$ , by viewing it as a function on Sym<sup>+</sup>( $n, \mathbb{R}$ ):

(6.4) 
$$\det^{\frac{1}{n}}(A+B)\det^{\frac{1}{n}}(A+C) \ge \det^{\frac{1}{n}}B\det^{\frac{1}{n}}C + \det^{\frac{1}{n}}A\det^{\frac{1}{n}}(A+B+C).$$

Inequality (6.4) is stronger the a usual log-supermodularity inequality for det, given the extra det  $\frac{1}{n} B \det \frac{1}{n} C$  term on its right hand side.

6.3. Vasic-Adamovic inequalities. We close the paper by briefly mentioning the related class of Vasic-Adamovic inequalities. Inspired by the work of D. M. Smiley and M. F. Smiley [39] on polygonal inequalities, P. M. Vasić and D. D. Adamović [41] found an inductive scheme for generating inequalities for any function that verifies the functional Hornich-Hlawka inequality. We state here a slightly modified version of their result [23, Theorem 2, pg. 528]:

**Theorem 12.** Let S be a commutative additive semigroup with 0 and G be an ordered abelian group (i.e., an abelian group with an order relation  $\leq$ ) such that

$$x, y, z \in \mathcal{G}$$
 and  $x \leq y$  implies  $x + z \leq y + z$ .

If  $\varphi : S \to G$  is a function such that for all  $x_1, x_2, x_3 \in S$ , we have

$$\sum_{1 \le i < j \le 3} \varphi \left( x_i + x_j \right) \le \sum_{k=1}^3 \varphi \left( x_k \right) + \varphi \left( \sum_{k=1}^3 x_k \right)$$

then for each pair  $\{k, n\}$  of integers with  $2 \le k < n$  we also have

$$\sum_{1 \le i_1 < \dots < i_k \le n} \varphi\left(\sum_{j=1}^k x_{i_j}\right) \le \binom{n-2}{k-1} \sum_{k=1}^n \varphi\left(x_k\right) + \binom{n-2}{k-2} \varphi\left(\sum_{k=1}^n x_k\right),$$

whenever  $x_1, ..., x_n \in \mathcal{S}$ .

This claim applies to every function f (defined on a convex cone C and taking values in an ordered Banach space E) that has positive differences of the third order. Indeed, in their case,

$$\begin{split} f\left(x+t\right) + f\left(y+t\right) + f\left(z+t\right) + f\left(x+y+z+t\right) \\ & \geq f\left(x+y+t\right) + f\left(y+z+t\right) + f\left(z+x+t\right) + f(t) \end{split}$$

for all points  $x, y, z, t \in C$  and performing the change of function  $\varphi(v) = f(v+t) - f(t)$  we obtain a function that verifies the Hornich-Hlawka inequality

$$\varphi\left(x\right) + \varphi\left(y\right) + \varphi\left(z\right) + \varphi\left(x + y + z\right) \ge \varphi\left(x + y\right) + \varphi\left(y + z\right) + \varphi\left(z + x\right).$$

#### References

- Amann H.: Multiple positive fixed points of asymptotically linear maps. J. Funct. Anal. 17, 174-213 (1974).
- [2] G. Bennett, Some forms of majorization, Houston J. Math. 36 (2010), 1037-1066.
- [3] Berndt, W., Sra, S.: Hlawka–Popoviciu inequalities on positive definite tensors. Linear Algebra Appl. 486, 317-327 (2015).
- [4] Bernstein, S.: Sur les fonctions absolument monotones, Acta Math. 52, 1–66 (1929).
- [5] Bhatia, R.: Matrix Analysis. Springer (1997).
- [6] Bhatia, R.: Positive Definite Matrices. Princeton University Press (2007).
- [7] Boas, R.P., Jr., Widder, D.V.: Functions with positive differences. Duke Math. J. 7, 496–503 (1940).
- [8] G. Choquet: Deux exemples classiques de représentation intégrale, Enseign. Math. 15, 63-75 (1969).
- [9] Del Moral, P., Niclas, A.: A Taylor expansion of the square root matrix function. J. Math. Anal. Appl. 465(1), 259-266 (2018).
- [10] Fréchet, M.: Sur la definition axiomatique d'une classe d'espaces vectoriels distanciés applicables vectoriellement sur l'espace de Hilbert. Ann. of Math. 36, 705–718 (1935).
- [11] Gal, S.G.: Shape Preserving Approximation by Real and Complex Polynomials. Birkhäuser, Boston (2008).
- [12] Gal, S.G., Niculescu C.P.: A new look at Popoviciu's concept of convexity for functions of two variables. J. Math. Anal. Appl. 479(1), 903-925 (2019)
- [13] Hopf, E.: Über die Zusammenhänge zwischen gewissen höheren Differenzenquotienten reeller Funktionen einer reellen Variablen und deren Differenzierbarkeitseigenschaften. Dissertation. Univ. Berlin (1926).
- [14] Hornich, H.: Eine Ungleichung für Vektorlängen. Math. Z. 48, 268–274 (1942)
- [15] Kozhasov K., Michalek M., Sturmfels B.: Positivity certificates via integral representations. volume II of London Math Society Lecture Notes Series, pages 84–114. Cambridge University Press (2022).
- [16] Kuczma, M.: An Introduction to the Theory of Functional Equations and Inequalities: Cauchy's Equation and Jensen's Inequality. Springer Science & Business Media (2009).
- [17] Levi, F.W.: Ein Reduktionsverfahren f
  ür lineare Vektorungleichungen. Archiv der Mathematik 2, 24-26 (1949).
- [18] Lin, M.: A determinantal inequality for positive semidefinite matrices. The Electronic Journal of Linear Algebra 27, 821-826 (2014)
- [19] J. Lindenstrauss, A. Pełczyński, Absolutely summing operators in  $\mathcal{L}_p$ -spaces and their applications, Studia Math. **29**, 275–326 (1968).
- [20] Marcus, M.: Finite Dimensional Multilinear Algebra, vol.I. Marcel Dekker (1973).
- [21] Merris, R.: Multilinear Algebra. Gordon & Breach. Amsterdam (1997).
- [22] Mirsky, L.: An introduction to linear algebra. Oxford University Press (1961).
- [23] Mitrinović, D.S., Pečarić, J.E., Fink, A.M.: Classical and new inequalities in analysis. Kluwer Academic Publishers, Dordrecht (1993).
- [24] Nelsen, R.B.: An introduction to copulas. Springer Science & Business Media (2006).
- [25] Niculescu, C.P., Olteanu, O.: From the Hahn-Banach extension theorem to the isotonicity of convex functions and the majorization theory. Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales (RACSAM), Serie A. Matemáticas 114(4), 1-19 (2020).
- [26] Niculescu, C.P., Persson, L.-E.: Convex Functions and their Applications. A Contemporary Approach, 2nd Ed., CMS Books in Mathematics Vol. 23, Springer-Verlag, New York (2018).
- [27] V. Paksoy, V., Turkmen, R., Zhang, F.: Inequalities of generalized matrix functions via tensor products. Electron. J. Linear Algebra, 27, 332–341 (2014).
- [28] Pecaric, J.E., Proschan, F., Tong, Y.L.: Convex Functions, Partial Orderings, and Statistical Applications. Mathematics in Science and Engineering vol. 187. Acad. Press., Inc., Boston (1992).
- [29] Popoviciu, T.: Sur l'approximation des fonctions convexes d'ordre supérieur. Mathematica (Cluj) 8, 1-85 (1934).
- [30] Popoviciu, T.: Sur l'approximation des fonctions convexes d'ordre supérieur. Mathematica (Cluj) 10, 49-54 (1935).
- [31] Popoviciu, T.: Les Fonctions Convexes. Hermann Cie. Editeurs, Paris (1944).

- [32] Popoviciu, T.: On some inequalities. Gaz. Mat., București 51, 81–85 (1946). (Romanian)
- [33] Schaefer, H.H., Wolff, M.P.: Topological Vector Spaces. Graduate Texts in Mathematics vol. 3, Springer Verlag, 1966.
- [34] Schilling, R.L., Song, R., Vondraček, Z.: Bernstein Functions. Theory and Applications. De Gruyter, Berlin (2010).
- [35] Scott, A.D., Sokal, A.D.: Complete monotonicity for inverse powers of some combinatorially defined polynomials. Acta Mathematica 213(2), 323-392 (2014).
- [36] Sendov, H.S., Zitikis, R.: The shape of the Borwein–Affleck–Girgensohn function generated by completely monotone and Bernstein functions. Journal of Optimization Theory and Applications 160(1), 67-89 (2014).
- [37] Serre, D.: The reverse Hlawka inequality in a Minkowski space. Comptes Rendus Mathematique 353(7), 629-633 (2015).
- [38] Siegel, C. L.: Uber die analytische Theorie der quadratischen Formen. Ann. of Math. 36, 527–606 (1935).
- [39] Smiley, D.M., Smiley, M.F.: The polygonal inequalities. Am. Math. Mon. 71, 755–760 (1964).
- [40] Sra, S. Reverse Minkowski (and related) determinant inequalities. MathOverflow. https://mathoverflow.net/q/251684 (version: 10/10/2016).
- [41] Vasić, P.M., Adamović D.D.: Sur un système infini d'inégalités fonctionnelles. Publ. Inst. Math. Nouv. Sér. 9 (23), 107–114 (1969).
- [42] Zhang, F.: Matrix Theory: Basic Results and Techniques, second edition. Springer, New York (2011).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRAIOVA, CRAIOVA 200585, ROMANIA *Email address*: constantin.p.niculescu@gmail.com

MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA *Email address:* suvrit@mit.edu