Probe of a Randall-Sundrum-like model from muon pair production at high energy muon collider

S.C. İnan^{*}

Department of Physics, Sivas Cumhuriyet University, 58140, Sivas, Turkey

and

A.V. Kisselev^{\dagger}

A.A. Logunov Institute for High Energy Physics, NRC "Kurchatov Institute", 142281, Protvino, Russian Federation

Abstract

We have examined inclusive $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\rm miss}$ and annihilation $\mu^+\mu^- \rightarrow \mu^+\mu^-$ processes at future high energy muon colliders in the framework of the Randall-Sundrum-like model with a small curvature of space-time. The collision energies of 3 TeV, 14 TeV and, 100 TeV are addressed. Both differential and total cross sections are calculated, and exclusion bounds on a 5-dimensional gravity scale are obtained depending on collision energy and integrated luminosity of the muon colliders.

1 Introduction

The Standard Model (SM) has been proven in a lot number of collider experiments. Nevertheless, we are still searching for solutions for many problems that SM cannot give a satisfactory solution. One of such problem is the so-called hierarchy problem which means the large energy gap between the

^{*}Electronic address: sceminan@cumhuriyet.edu.tr

[†]Electronic address: alexandre.kisselev@ihep.ru

electroweak scale and gravity scale. The most elegant answer to this phenomenon has been given in the framework of the Randall-Sundrum (RS) model [1] which is based on a 5D theory with one extra dimension compactified in an orbifold S_1/Z_2 . The main parameters of the RS model are a compactification radius r_c and AdS₅ curvature parameter k (hereinafter referred to as the *curvature* k). The model predicts Kaluza-Klein (KK) gravitons which are heavy resonances with masses around the TeV scale. The most stringent limits on KK graviton masses come from the LHC searches for heavy resonances. The experimental limits depend on a ratio $k/M_{\rm Pl}$, where $M_{\rm Pl}$ is the Planck mass. The CMS collaboration have excluded KK graviton masses below 2.3 to 4.0 TeV for the diphoton final state [2]. For the dilepton final state the CMS have excluded the RS graviton masses in the region 2.47-4.78 TeV [3]. The best lower limit of the ATLAS collaboration, 4.6 TeV, has been obtained in searching for the diphoton final state [4].

In papers [5, 6] the RS-like model with a small curvature of the 5dimensional space-time (RSSC model) has been proposed. In particular, a general solution for the warped metric has been obtained [7]. In contrast to the original RS model, the RSSC model has an almost continuous graviton mass spectrum which is similar to that of the ADD model [8]-[10], if $k \ll M_{\rm Pl}$. Thus, the above mentioned experimental bounds are not applied to the RS scenario with a small value of k. A probe of the RSSC model at the LHC can be found in [11, 12]. A detailed comparison of the RSSC model with the RS model is given in section 2.

In the present paper we intend to examine the RSSC model through the $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$ and $\mu^+\mu^- \rightarrow \mu^+\mu^-$ processes at a future muon collider. The idea of the muon collider was proposed by F. Tikhonin and G. Budker in the late 1960's [13, 14], and it was also discussed in the early 1980's [15, 16]. At present, a great physical potential of the muon collider for collisions of elementary particles at very high energies is being actively examined. Its advantage lies in the fact that muons can be accelerated in a ring without limitation from synchrotron radiation compared to linear or circular electron-positron colliders [17]-[22]. For instance, the muon collider may provide a determination of the electroweak couplings of the Higgs boson which is significantly better than what is considered attainable at other future colliders [23]-[29]. Interest in designing and building a muon collider is also based on its capability of probing the physics beyond the SM. In a number of recent papers searches for SUSY particles [30], WIMPs [31]-[33], and dark matter [34], vector boson fusion [35], leptoquarks [36], lepton flavor violation [37]-[39], vector-like leptons [40], heavy leptons [41, 42], and heavy neutrinos [43], top Yukawa couplings [44], multi-boson processes [45], and physics of the muon (g-2) [46]-[48] are presented. In our recent paper we have probed axion-like particles (ALPs) at high energy muon colliders [49]. In a number of papers anomalous quartic [50]-[52] and triple [53, 54] gauge couplings at the muon collider were studied. For more details on a spectacular opportunity of the muon collider in the direct exploration of the energy frontier, see [55].

Our goal is to obtain exclusion bounds on the 5-dimensional Planck scale M_5 which can be probed from two mentioned above processes at TeV and multi-TeV muon colliders. In the inclusive $\mu^+\mu^- \to \mu^+\mu^- + E_{\rm miss}$ scattering one pair of muons with large transverse momenta and large invariant mass is detected, while the other two scattered muons or produced neutrinos escape a detector. The gravity contribution comes from two subprocess, $V_1V_2 \to G \to \mu^+\mu^-$, with $V_{1,2} = \gamma, Z$, and $W^+W^- \to G \to \mu^+\mu^-$, where G denotes a KK graviton, and a summation over all KK gravitons is assumed. The other process we are interested in is the annihilation $\mu^+\mu^- \to \mu^+\mu^-$ scattering which has contributions from s- and t-channel graviton exchanges. Note that the processes can be easily distinguished experimentally from each other, since they have quite different distributions in invariant mass of the detected dimuon pair.

The paper is organized as follows. In the next section, the detailed description of the RSSC model is presented. In section 3 we examine the production of the muon pair accompanied by missing energy via vector boson fusion at the muon collider. The bounds on 5-dimensional Planck scale M_5 are obtained. In section 4 we study the exclusive dimuon production and calculate the values of M_5 which can be probed in this collision at the muon collider.

2 Model of warped extra dimension with small curvature

In this section, we describe the RSSC model in detail and compare it with the original RS model. The RS scenario with one extra dimension and two branes [1] was proposed as an alternative to the ADD scenario with large flat extra dimensions (EDs) [8]-[10]. It has the following background warped metric

$$ds^{2} = e^{-2\sigma(y)} \eta_{\mu\nu} \, dx^{\mu} \, dx^{\nu} - dy^{2} \,, \tag{1}$$

where $\eta_{\mu\nu}$ is the Minkowski tensor with the signature (+, -, -, -), y is an extra coordinate, and $\sigma(y)$ is the warp factor. The periodicity condition $y = y + 2\pi r_c$ is imposed, and the points (x_{μ}, y) and $(x_{\mu}, -y)$ are identified. Thus, we have a model of gravity in a slice of the AdS₅ space-time compactified to the orbifold S^1/Z_2 . This orbifold has two fixed points, y = 0, and $y = \pi r_c$. There are two branes located at these points (called Planck and TeV brane, respectively). The SM fields are confined to the TeV brane.

The classical action of the RS model is given by [1]

$$S = \int d^4x \int_{-\pi r_c}^{\pi r_c} dy \sqrt{G} \left(2\bar{M}_5^3 \mathcal{R} - \Lambda \right) + \int d^4x \sqrt{|g^{(1)}|} \left(\mathcal{L}_1 - \Lambda_1 \right) + \int d^4x \sqrt{|g^{(2)}|} \left(\mathcal{L}_2 - \Lambda_2 \right), \qquad (2)$$

where $G_{MN}(x, y)$ is the 5-dimensional metric, with $M, N = 0, 1, 2, 3, 4, \mu = 0, 1, 2, 3$. The quantities

$$g_{\mu\nu}^{(1)}(x) = G_{\mu\nu}(x, y=0) , \quad g_{\mu\nu}^{(2)}(x) = G_{\mu\nu}(x, y=\pi r_c)$$
 (3)

are induced metrics on the branes, \mathcal{L}_1 and \mathcal{L}_2 are brane Lagrangians, $G = \det(G_{MN})$, $g^{(i)} = \det(g^{(i)}_{\mu\nu})$. The parameter \overline{M}_5 is a *reduced* 5-dimensional Planck scale, $\overline{M}_5 = M_5/(2\pi)^{1/3}$. The parameter Λ is a 5-dimensional cosmological constant, and $\Lambda_{1,2}$ are brane tensions.

For the first time, the solution for $\sigma(y)$ in (1) has been obtained in [1],

$$\sigma_{\rm RS}(y) = k|y| , \qquad (4)$$

where k is a parameter with a dimension of mass. It defines the curvature of the 5-dimensional space-time. Later on in [7] the following general solution for $\sigma(y)$ was derived

$$\sigma(y) = \frac{kr_c}{2} \left[\left| \operatorname{Arccos}\left(\cos \frac{y}{r_c} \right) \right| - \left| \pi - \operatorname{Arccos}\left(\cos \frac{y}{r_c} \right) \right| \right] + \frac{\pi |k|r_c}{2} - C , \quad (5)$$

where $\operatorname{Arccos}(z)$ is a principal value of the multivalued inverse trigonometric function $\operatorname{arccos}(z)$, and C is a y-independent quantity.¹ The constant C

¹At the same time, C may depend on the parameter r_c , see below.

arises in (5) because Einstein–Hilbert's equations for $\sigma(y)$ contain only $\sigma'(y)$ and $\sigma''(y)$, where prime denotes a derivative with respect to the variable y, but not the function $\sigma(y)$ itself. Let us underline that solution (5) (i) obeys the orbifold symmetry $y \to -y$; (ii) makes the jumps of $\sigma'(y)$ on both branes; (iii) has explicit symmetry with respect to the branes. More details can be found in [7].

By taking C = 0 in (5), we get the RS model (4), while taking $C = \pi k r_c$ we come to the RS-like scenario with the small curvature of space-time (RSSC model, see [5]-[7]). In general, different values of C in the warp function (5) result in different spectra of the KK gravitons, see Appendix A for more details.

It is worth to remind the main features of the RSSC model in comparison with those of the RS model. The interactions of the Kaluza-Klein (KK) gravitons $h^{(n)}_{\mu\nu}$ with the SM fields on the TeV brane are given by the effective Lagrangian density

$$\mathcal{L}_{\rm int} = -\frac{1}{\bar{M}_{\rm Pl}} h_{\mu\nu}^{(0)}(x) T_{\alpha\beta}(x) \eta^{\mu\alpha} \eta^{\nu\beta} - \frac{1}{\Lambda_{\pi}} \sum_{n=1}^{\infty} h_{\mu\nu}^{(n)}(x) T_{\alpha\beta}(x) \eta^{\mu\alpha} \eta^{\nu\beta} , \quad (6)$$

were $\bar{M}_{\rm Pl} = M_{\rm Pl}/\sqrt{8\pi}$ is the reduced Planck mass, $T^{\mu\nu}(x)$ is the energymomentum tensor of the SM fields. The coupling constant is equal to

$$\Lambda_{\pi} = \bar{M}_5 \sqrt{\frac{\bar{M}_5}{k}} \,. \tag{7}$$

The hierarchy relation looks like

$$\bar{M}_{\rm Pl}^2 = \frac{\bar{M}_5^3}{k} \left[e^{2\pi k r_c} - 1 \right]. \tag{8}$$

To compare, in the original RS model the hierarchy relation is different, $\bar{M}_{\rm Pl}^2 = (\bar{M}_5^3/k) \left(1 - e^{-2\pi k r_c}\right)$, and $\Lambda_{\pi} = M_{\rm Pl} e^{-\pi k r_c}$. As for the Λ -term and brane tensions, they obey the fine-tuning relations [7]²

$$\Lambda = -24\bar{M}_{5}^{3}k^{2} , \quad \Lambda_{1} = -\Lambda_{2} = 12\bar{M}_{5}^{3}k .$$
(9)

In the RSSC model masses of the KK gravitons are proportional to the curvature k [5, 6],

$$m_n = x_n k , \quad n = 1, 2, \dots ,$$
 (10)

²In the RS model $\Lambda = -24\bar{M}_5^3k^2$, but $\Lambda_1 = -\Lambda_2 = 24\bar{M}_5^3k$ [1].

where x_n are zeros of the Bessel function $J_1(x)$. Should we take $k \ll \overline{M}_5 \sim 1$ TeV, the mass splitting Δm will be very small, $\Delta m \simeq \pi k$, and we come to an *almost continuous* mass spectrum, similar to the mass spectrum of the ADD model [8]. Nevertheless, even for rather small curvature k the RSSC model cannot be regarded as an IR modification of the ADD model with one ED, see Appendix B. On a contrary, in the RS model the gravitons are *heavy* resonances with masses above one-few TeV. For the first time heavy graviton searches were discussed in [56].

As was shown in a number of phenomenological papers on the RSSC model [11, 12], cross sections weakly depend on the parameter k, if $k \ll \overline{M}_5$. That is why, in what follows we will fix this parameter to be k = 1 GeV. Let us underline once more that the graviton mass spectra are quite different in the RSSC and RS models. It means that all experimental bounds and phenomenological predictions for parameters of the original RS model can not be applied to the RS-like scheme examined in the present paper.

3 Production of muon pair with missing energy in muon collisions

Let us study a KK graviton contribution to the inclusive dimuon production $\mu^{-}\mu^{+} \rightarrow \mu^{+}\mu^{-} + E_{\text{miss}}$ in the collision of muon beams. It is defined by gauge boson fusions, $V_1V_2 \rightarrow \mu^{+}\mu^{-}$ ($V_{1,2} = \gamma, Z$) and $W^+W^- \rightarrow \mu^{+}\mu^{-}$ as shown in Figs. 1 and 2.³ In the framework of the RSSC model a scattering amplitude is equal to $M_{\text{SM}} + M_{\text{KK}}$, where M_{SM} is the SM term, and M_{KK} is given by a sum of *s*-channel KK gravitons

$$M_{KK} = \frac{1}{2\Lambda_{\pi}^2} \sum_{n=1}^{\infty} \left[\bar{u}(p_1) \Gamma_2^{\mu\nu} v(p_2) \frac{B_{\mu\nu\alpha\beta}}{s - m_n^2 + i \, m_n \Gamma_n} \Gamma_1^{\alpha\beta\rho\sigma} e_\rho(k_1) e_\sigma(k_2) \right].$$
(11)

Here k_1, k_2 , and $e_{\rho}(k_1)$, $e_{\sigma}(k_2)$ are, respectively, the momenta and polarization vectors of the incoming bosons, and p_1, p_2 are momenta of the outgoing leptons. Γ_n is the total width of the graviton with the mass m_n . The coherent sum in (11) is over all massive KK modes. The Feynman rules for the KK graviton were derived in [57, 58] (see also [59]). In particular, the vector

³In what follows, $V_1(V_2)$ denotes γ or Z, while V without subscript means γ, Z or W.




Figure 1: The Feynman diagrams describing contribution of the KK graviton G to the collision of two neutral gauge bosons V_1, V_2 with outgoing charged leptons at the muon collider.




Figure 2: The Feynman diagrams describing contribution of the KK graviton G to the fusion of the W bosons, with two outgoing charged leptons at the muon collider.

boson-graviton vertex VVG has the form

$$\Gamma_1^{\alpha\beta\rho\sigma} = -\frac{i}{2} \left\{ \left[m_V^2 + (k_1 \cdot k_2) \right] C^{\alpha\beta\rho\sigma} + D^{\alpha\beta\rho\sigma} \right\}, \qquad (12)$$

where m_V is a mass of the gauge boson, and

$$C^{\alpha\beta\rho\sigma} = \eta^{\alpha\rho}\eta^{\beta\sigma} + \eta^{\alpha\sigma}\eta^{\beta\rho} - \eta^{\alpha\beta}\eta^{\rho\sigma} , \qquad (13)$$

$$D^{\alpha\beta\rho\sigma} = \eta^{\alpha\beta}k_{1}^{\sigma}k_{2}^{\rho} - (\eta^{\alpha\sigma}k_{1}^{\beta}k_{2}^{\rho} + \eta^{\alpha\rho}k_{1}^{\sigma}k_{2}^{\beta} - \eta^{\rho\sigma}k_{1}^{\alpha}k_{2}^{\beta}) - (\eta^{\beta\sigma}k_{1}^{\alpha}k_{2}^{\rho} + \eta^{\beta\rho}k_{1}^{\sigma}k_{2}^{\alpha} - \eta^{\rho\sigma}k_{1}^{\beta}k_{2}^{\alpha}) .$$
(14)

The KK graviton-lepton vertex Gl^+l^- is given by [57, 58]

$$\Gamma_2^{\mu\nu} = -\frac{i}{8} \left[\gamma^{\mu} (p_1^{\nu} - p_2^{\nu}) + \gamma^{\nu} (p_1^{\mu} - p_2^{\mu}) \right].$$
(15)

Finally, $B_{\mu\nu\alpha\beta}$ in (11) is a tensor part of the KK graviton propagator. Its explicit expression was also derived in [57, 58]. We can safely omit terms in $B_{\mu\nu\alpha\beta}$ which give zero contribution to eq. (11). Then we can write

$$B_{\mu\nu\alpha\beta} = \eta_{\mu\alpha}\eta_{\nu\beta} + \eta_{\mu\beta}\eta_{\nu\alpha} - \frac{2}{3}\eta_{\mu\nu}\eta_{\alpha\beta} .$$
 (16)

The s-channel contribution of the KK gravitons is equal to

$$S(s) = \frac{1}{\Lambda_{\pi}^2} \sum_{n=1}^{\infty} \frac{1}{s - m_n^2 + i \, m_n \Gamma_n} \,. \tag{17}$$

This series has been analytically calculated in [60]

$$\mathcal{S}(s) = -\frac{1}{4\bar{M}_5^3\sqrt{s}} \frac{\sin(2A) + i\sinh(2\varepsilon)}{\cos^2 A + \sinh^2 \varepsilon} , \qquad (18)$$

where

$$A = \frac{\sqrt{s}}{k} , \quad \varepsilon = 0.045 \left(\frac{\sqrt{s}}{\bar{M}_5}\right)^3.$$
 (19)

If the ratio \sqrt{s}/\bar{M}_5 is large enough, we get $\mathcal{S}(s) \simeq -i/(2\bar{M}_5^3\sqrt{s})$.

The squared amplitude of the subprocess $VV \rightarrow l^- l^+$ is a sum of three terms,

$$|M|^{2} = |M_{\rm SM}|^{2} + |M_{\rm KK}|^{2} + |M_{\rm int}|^{2} , \qquad (20)$$

where $M_{\rm SM}$ denotes the SM amplitude, while $M_{\rm KK}$ and $M_{\rm int}$ are the gravity and interference terms. In [59] the quantities $|M_{\rm SM}(\gamma\gamma \rightarrow l^-l^+)|^2$, $|M_{\rm KK}(\gamma\gamma \rightarrow l^-l^+)|^2$, and $|M_{\rm int}(\gamma\gamma \rightarrow l^-l^+)|^2$ were calculated for massless leptons. The results of our calculations of squared amplitudes $|M(VV \rightarrow l^-l^+)|^2$ for nonzero m_l and m_V are presented in Appendix C.

The virtual KK graviton production should lead to deviations from SM predictions in a magnitude of the cross section. The cross section of the

 $\mu^-\mu^+ \to \mu^+\mu^- + E_{\rm miss}$ scattering is defined by

$$d\sigma = \int_{\tau_{\min}}^{\tau_{\max}} d\tau \int_{x_{\min}}^{x_{\max}} \frac{dx}{x} \Bigg[\sum_{V_1, V_2 = \gamma, Z_T, Z_L} f_{V_1/\mu^+}(x, Q^2) f_{V_2/\mu^-}(\tau/x, Q^2) \, d\hat{\sigma}(V_1 V_2 \to \mu^+ \mu^-) \\ + \sum_{W_1, W_2 = W_T, W_L} f_{W_1^+/\mu^+}(x, Q^2) f_{W_2^-/\mu^-}(\tau/x, Q^2) \, d\hat{\sigma}(W_1^+ W_2^- \to \mu^+ \mu^-) \Bigg],$$
(21)

where

$$x_{\max} = 1 - \frac{m_{\mu}}{E_{\mu}}, \ \tau_{\max} = \left(1 - \frac{m_{\mu}}{E_{\mu}}\right)^2, \ x_{\min} = \tau/x_{\max}, \ \tau_{\min} = \frac{p_{\perp}^2}{E_{\mu}^2}, \quad (22)$$

and p_{\perp} is the transverse momenta of the outgoing photons. $f_{\gamma/\mu^{\pm}}(x, Q^2)$, $f_{Z_T/\mu^{\pm}}(x, Q^2)$, $f_{Z_L/\mu^{\pm}}(x, Q^2)$, $f_{W_T^{\pm}/\mu^{\pm}}(x, Q^2)$, and $f_{W_L^{\pm}/\mu^{\pm}}(x, Q^2)$ are unpolarized boson distributions inside unpolarized muon beams. In the leading order they are given by [61]

$$f_{\gamma/\mu^{\pm}}(x,Q^2) = \frac{\alpha}{2\pi} \frac{1 + (1-x)^2}{x} \ln \frac{Q^2}{m_{\mu}^2} , \qquad (23)$$

and [62, 63]

$$f_{Z_T/\mu^{\pm}}(x,Q^2) = \frac{\alpha_Z^{\pm}}{2\pi} \frac{1 + (1-x)^2}{x} \ln \frac{Q^2}{m_Z^2},$$

$$f_{Z_L/\mu^{\pm}}(x,Q^2) = \frac{\alpha_Z^{\pm}}{\pi} \frac{(1-x)}{x},$$

$$f_{W_T^{\pm}/\mu^{\pm}}(x,Q^2) = \frac{\alpha_W}{2\pi} \frac{1 + (1-x)^2}{x} \ln \frac{Q^2}{m_W^2},$$

$$f_{W_L^{\pm}/\mu^{\pm}}(x,Q^2) = \frac{\alpha_W}{\pi} \frac{(1-x)}{x},$$
(24)

where

$$\alpha_Z^{\pm} = \frac{\alpha}{(\cos\theta_W \sin\theta_W)^2} \left[(g_V^{\pm})^2 + (g_A^{\pm})^2 \right], \quad \alpha_W = \frac{\alpha}{4\sin\theta_W^2} , \qquad (25)$$

$$g_V^{\pm} = \pm 1/4 \mp \sin^2 \theta_W , \quad g_A^{\pm} = \pm 1/4 ,$$
 (26)

and m_{μ} is the muon mass. The variable x in eqs. (23), (24) is a ratio of the boson energy and energy of the incoming muon E_{μ} . Note that the Z and W bosons have different distributions for their transverse (T) and longitudinal (L) polarizations (24). Note also that the distribution of the massive bosons V (V = Z, W) is suppressed with respect to the photon distribution by the factor $\ln(Q^2/m_V^2)/\ln(Q^2/m_{\mu}^2)$. The mixed γ -Z and Z- γ terms are present in (21) along with the γ - γ and Z-Z terms.

The differential cross section of the $VV \rightarrow \mu^+\mu^-$ collision is a sum of helicity amplitudes squared,

$$\frac{d\hat{\sigma}}{d\Omega} = \frac{1}{64\pi^2 \hat{s}} \sum_{\lambda_1, \lambda_2, \lambda_3, \lambda_4} |M_{\lambda_1 \lambda_2 \lambda_3 \lambda_4}|^2,$$
(27)

where $\sqrt{\hat{s}} = 2E_{\mu}\sqrt{\tau}$ is an invariant energy of this collision, and $\lambda_{1,2}$ ($\lambda_{3,4}$) are boson (muon) helicities. In distributions (23), (24) we put $Q^2 = \hat{s}$.

The scattering angle for high energy muons in Fig. 1 is peaked near $\theta_{\mu} \approx 0.02^{\circ} - 1.2^{\circ}$ [64]. These very forward muons would most likely escape a muon detector away from colliding beams. Thus, only muons produced in the fusion of the neutral bosons will be detected. In the case of the W boson fusion, as in Fig. 2, the neutrino will escape detection.

For numerical analysis we apply the cut on the rapidity and transverse momenta of the detected muons, $|\eta| < 2.5$, $p_t > 50$ GeV. As was already mentioned above, we take k = 1 GeV. The results of our calculations of the differential cross sections for the $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$ scattering at the future muon collider are presented in Fig. 3. The predictions for three collision invariant energies of the muon collider are shown. As one can see, for each energy the cross sections rise as the invariant mass of the detected muons $m_{\mu^+\mu^-}$ grows, while the SM cross sections decrease more rapidly with an increase of $m_{\mu^+\mu^-}$. We have also calculated the differential cross sections via transverse momentum of the detected muons, see Fig. 4. They look like the cross sections in Fig. 3.

The total cross section as function of the minimal invariant mass of two detected muons $m_{\mu^+\mu^-,\min}$ is shown in Fig. 5. As one can see, it strongly depends on the gravity scale \bar{M}_5 . If \bar{M}_5 is of order one TeV, the cross section exceeds the SM one for all collision energies. For larger values of \bar{M}_5 the total cross section strongly dominates over the SM cross section at $\sqrt{s} = 14$ TeV and $\sqrt{s} = 100$ TeV.

All this enables us to derive the exclusion bounds on the 5-dimensional




Figure 3: The differential cross sections for the process $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$ via invariant mass of two detected muons at the muon collider. The left, middle and right panels correspond to the colliding energy of 3 TeV, 14 TeV, and 100 TeV, respectively. Three color curves (from the top down) correspond to different values of \bar{M}_5 . The SM cross sections (low curves) are also shown.

reduced Planck scale \overline{M}_5 . To derive them, we apply the following formula for the statistical significance SS [65]

$$SS = \sqrt{2[(S - B \ln(1 + S/B)]},$$
 (28)

where S is the number of signal events and B is the number of background (SM) events. We define the regions $SS \leq 1.645$ as the regions that can be excluded at the 95% C.L. To reduce the SM background, we additionally used the cuts $m_{\mu^-\mu^+} > 1$ TeV, $m_{\mu^-\mu^+} > 5$ TeV, and $m_{\mu^-\mu^+} > 50$ TeV for the collision energy of 3 TeV, 14 TeV, and 100 TeV, respectively. The results are shown in Fig. 6. Our best limits for $\sqrt{s} = 3$ TeV, 14 TeV and 100 TeV are, respectively, $\overline{M}_5 = 5.86$ TeV, 28.45 TeV and 213.5 TeV.




Figure 4: The differential cross sections for the process $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$ via transverse momentum of the detected muons at the muon collider.

4 Muon pair production in muon collision

In the previous section we assumed that in the $\mu^+\mu^- \to \mu^+\mu^- + E_{\text{miss}}$ process only two final muons are detected, while two scattered muons or neutrinos escape the detector. It means that the invariant mass of the outgoing muons can vary from one event to another. On the contrary, in the annihilation $\mu^-\mu^+ \to \mu^+\mu^-$ process the invariant mass of the final system $m_{\mu^+\mu^-}$ is fixed and close to the collision energy \sqrt{s} . Note that the muon collider has a low level of beamstrahlung and synchrotron radiation compared to linear or circular electron-positron colliders. As a result, an energy spread in the collision is significantly reduced, and it enables an improved energy resolution. That is why one can easily discriminate between two processes by measuring the invariant mass of the detected muon pair.

As in the previous case, a virtual production of the KK gravitons should give a contribution to the cross sections of the $\mu^-\mu^+ \rightarrow \mu^+\mu^-$ scattering. It is shown in Fig. 7. Our analytical expressions for amplitudes squared of this collision are collected in Appendix D. It is natural to present the cross section of the $\mu^-\mu^+ \rightarrow \mu^+\mu^-$ process as a function of transverse momentum




Figure 5: The total cross sections for the process $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$ via minimal invariant mass of two detected muons at the muon collider $m_{\mu^+\mu^-,\text{min}}$.

of the final muons p_t . We have calculated the differential cross sections for the $\mu^+\mu^- \rightarrow \mu^+\mu^-$ scattering at the muon collider, taking into account a contribution from the massive KK gravitons. The results of our calculations are given in Fig. 8 for three values of the collision energy \sqrt{s} and different values of the reduced 5-dimensional Planck scale \bar{M}_5 . As we can see, for $\sqrt{s} = 14$ TeV and $\sqrt{s} = 100$ TeV the cross section significantly dominates the SM one, especially for large p_t . The differential cross sections integrated in p_t from the minimal transverse momentum of the detected muons $p_{t,\min}$ are presented in Fig. 9.

As before, we aim at calculating exclusion bounds on M_5 which can be probed in the process $\mu^+\mu^- \rightarrow \mu^+\mu^-$ depending on the integrated luminosity of the future muon collider. We have used eq. (28) for the statistical significance. In doing so, the cuts $p_{t,\min} = 0.5$ TeV, $p_{t,\min} = 2.5$ TeV, and $p_{t,\min} = 25$ TeV were applied for the 3 TeV, 14 TeV, and 100 TeV center-of-




Figure 6: The exclusion bounds on the reduced fundamental gravity scale \overline{M}_5 via integrated luminosity of the muon collider for the process $\mu^+\mu^- \rightarrow \mu^+\mu^- + E_{\text{miss}}$. The left, middle and right panels correspond to the colliding energy of 3 TeV, 14 TeV, and 100 TeV.

mass energies, respectively. The exclusion regions are given in Fig. 10. We conclude that in the $\mu^+\mu^- \rightarrow \mu^+\mu^-$ process scales up to $\bar{M}_5 = 3.85$ TeV, 17.8 TeV and 126.3 TeV can be probed for $\sqrt{s} = 3$ TeV, 14 TeV, and 100 TeV, respectively. We see that these bounds on \bar{M}_5 are stronger than our constraints for the $\mu^-\mu^+ \rightarrow \mu^+\mu^- + E_{\rm miss}$ scattering (see section 3).

5 Conclusions

We have examined two collisions at future TeV and multi-TeV muon colliders in the Randall-Sundrum-like model with the small curvature (or RSSC model, for short) [5, 6]. It is the model with one ED and warped metric whose 5-dimensional space-time curvature k is about one GeV. The other main parameter of the RSSC model, the reduced 5-dimensional Planck scale \bar{M}_5 , is equal to (larger than) one TeV.

We have studied the $\mu^-\mu^+ \rightarrow \mu^+\mu^- + E_{\rm miss}$ scattering first. It goes via




Figure 7: The Feynman diagrams describing contribution of the KK graviton G to the $\mu^{-}\mu^{+} \rightarrow \mu^{+}\mu^{-}$ scattering.

the $VV \rightarrow \mu^+\mu^-$ collision, where V means one of the gauge bosons, γ, Z , or W. The squared amplitudes for the $VV \rightarrow l^+l^-$ collision, including the gravity, SM and, interference terms, have been analytically calculated for the first time for massive leptons l^{\pm} . Their explicit expressions are collected in Appendix C. The differential cross sections depending on the invariant mass of the detected muon pair $m_{\mu^+\mu^-}$ are calculated for three values of the reduced 5-dimensional Planck scale \bar{M}_5 for 3 TeV, 14 TeV and 100 TeV muon colliders. The total cross sections are presented as functions of the minimal value of $m_{\mu^+\mu^-}$. As a result, the exclusion bounds on the scale \bar{M}_5 are obtained. They are equal to $\bar{M}_5 = 5.86$ TeV, 28.45 TeV and 213.5 TeV for the collision energy of $\sqrt{s} = 3$ TeV, 14 TeV and 100 TeV, respectively.

The $\mu^-\mu^+ \to \mu^+\mu^-$ scattering has been also studied. As in the previous case, we have calculated the gravity, SM and interference squared amplitudes analytically, see Appendix D. It enabled us to estimate numerically the differential cross sections as functions of the transverse momenta of the outgoing muons. The total cross sections are also calculated. Finally, the exclusion bounds on the scale \bar{M}_5 , have been obtained. We have shown that the values of $\bar{M}_5 = 3.85$ TeV, 17.8 TeV and 126.3 TeV can be probed at 3 TeV, 14 TeV, and 100 TeV muon colliders. Remember that $\bar{M}_5 = M_5/(2\pi)^{1/3} \approx 0.54M_5$, where M_5 is the fundamental 5-dimensional gravity scale M_5 in the RSSC model. It means that the bounds on the scale M_5 are approximately twice stronger that the above mentioned limits for \bar{M}_5 .

Let us stress again that our bounds on M_5 should not be compared with the current experimental limits on the 5-dimensional gravity scale of the original RS model [1], since the mass spectra of the KK gravitons and, correspondingly, experimental signatures for the RSSC and RS models are quite




Figure 8: The differential cross sections for the process $\mu^+\mu^- \rightarrow \mu^+\mu^-$ via transverse momentum of the detected muons at the muon collider. The left, middle and right panels correspond to the colliding energy of 3 TeV, 14 TeV, and 100 TeV, respectively. Three color curves (from the top down) correspond to different values of \overline{M}_5 . The SM cross sections (low curves) are also shown.

different [5]-[7]. The comparative analysis of these models was presented in section 2.

Appendix A. Shift of warp function

Here we show that a shift of the warp function of the original RS model,

$$\sigma(y) \to \sigma(y) - C$$
, (A.1)

where C is a constant, results in a model with a quite different graviton spectrum. Note that (A.1) is equivalent to the rescaling of the 4D metric

$$g_{\mu\nu} \to g_{\mu\nu} e^{2C}. \tag{A.2}$$

The Einstein tensor $R_{\mu\nu} - (1/2)g_{\mu\nu}R$ is invariant under such a transformation. As for the energy-momentum tensor, it is scale-invariant only for massless




Figure 9: The total cross sections for the process $\mu^+\mu^- \to \mu^+\mu^-$ via minimal transverse momentum of the outgoing muons $p_{t,\min}$.

fields. As a simple example, consider the energy-momentum tensor of the massive scalar field,

$$T_{\mu\nu} = \partial_{\mu}\phi \,\partial_{\nu}\phi - \frac{1}{2} g_{\mu\nu} \left[g^{\alpha\beta} \partial_{\alpha}\phi \,\partial_{\beta}\phi - m^2 \phi^2 \right]. \tag{A.3}$$

It is not scale-invariant due to the third term in (A.3). In general, theories with *massive* fields are not invariant under transformation (A.2).

Now consider an effective 4-dimensional gravity action on the TeV brane (with radion term omitted) [66]

$$S_{\text{eff}} = \frac{1}{4} \sum_{n=0}^{\infty} \int d^4x \Big[\partial_{\mu} h_{\varrho\alpha}^{(n)}(x) \partial_{\nu} h_{\delta\lambda}^{(n)}(x) \eta^{\mu\nu} - m_n^2 h_{\varrho\alpha}^{(n)}(x) h_{\delta\lambda}^{(n)}(x) \Big] \eta^{\varrho\delta} \eta^{\alpha\lambda} \,. \tag{A.4}$$

The transformation (A.1) can be also regarded as a rescaling of four-dimensional coordinates

$$x^{\mu} = e^C x^{\prime \mu} , \qquad (A.5)$$

but without change of the metric [67]. Note that (A.5) is not a particular




Figure 10: The exclusion bounds on the reduced fundamental gravity scale \overline{M}_5 via integrated luminosity of the muon collider for the process $\mu^+\mu^- \rightarrow \mu^+\mu^-$. The left, middle and right panels correspond to the colliding energy of 3 TeV, 14 TeV, and 100 TeV.

case of general coordinate transformation in gravity, since the metric tensor $g_{\mu\nu}$ remains fixed.

The invariance of the action (A.4) under transformation (A.5) needs rescaling of the graviton fields and their mass: $h_{\mu\nu}^{(n)} = e^{-C} h_{\mu\nu}^{\prime(n)}$, $m_n = e^{-C} m'_n$. We see that the theory of massive KK gravitons is not scale-invariant. Only zero mode (massless graviton) remains unchanged. We conclude that the warp functions $\sigma(y)$ and $\sigma(y) - C$ correspond to two *nonequivalent* fourdimensional theories whose spectra of massive gravitons differ from each other [5, 7].

Appendix B. RSSC model versus ADD model with one dimension

We show that the AdS_5 space-time differs significantly from a 5-dimensional flat space-time with *one* large ED even for very small curvature parameter

k. Compare the hierarchy relations in both scenarios. The hierarchy relation for the ADD model with one ED of the size R_c looks like [8]-[10]

$$\bar{M}_{Pl}^2 = (2\pi R_c) \,\bar{M}_5^3,$$
 (B.1)

where \overline{M}_5 is the reduce 5-dimensional Planck scale in the ADD model. One can see that eq. (B.1) follows from eq. (8) in the limit

$$\pi kr \ll 1 , \qquad (B.2)$$

after replacement $r \to R_c$. However, it follows from (B.1) and (B.2) that then k should be unnaturally small,

$$k \ll \frac{\bar{M}_5^3}{\bar{M}_{\rm Pl}^2}$$
, (B.3)

even if \overline{M}_5 is a few TeV or tens of TeV. That is why, the RS-like model with the small (but not negligibly small) curvature k cannot be regarded as an IR modification of the ADD model, at least, for the parameters used in the present paper.⁴

Appendix C. Squared amplitudes for $VV \rightarrow l^-l^+$ scattering

Our calculations give the following analytical expressions for the squared amplitudes of the $\gamma\gamma \rightarrow l^-l^+$ collision in eq. (20)

$$|M_{\rm SM}|^2 = \frac{8e^4}{(t-m_l^2)^2(s+t-m_l^2)^2} [-34m_l^8 + m_l^6(60s+64t) - m_l^4(31s^2+52st+28t^2) + m_l^2s(s^2-2st-4t^2) - t(s+t)(s^2+2st+2t^2)], \qquad (C.1)$$

$$|M_{\rm KK}|^2 = -\frac{1}{8}|S(s)|^2[2m_l^8 - 8m_l^6t + m_l^4(s^2 + 4st + 12t^2) - 2m_l^2t(s + 2t)^2 + t(s+t)(s^2 + 2st + 2t^2)], \qquad (C.2)$$

⁴Let us remember that we study the case k = 1 GeV, $\overline{M}_5 = 1 \div 100$ TeV.

$$|M_{\rm int}|^2 = -\frac{e^2[S(s) + S^*(s)]}{2(t - m_l^2)(s + t - m_l^2)} [-2m_l^8 + m_l^6(3s + 4t) + m_l^4s(3s - 4t) - m_l^2(s^3 + 2s^2t + 3st^2 + 4t^3) + t(s + t)(s^2 + 2st + 2t^2)], \quad (C.3)$$

where s, t are Mandelstam variables, m_l is the lepton mass, and S(s) is defined in the text (17)-(19). If we take $m_l = 0$, we get known results obtained in [59].

For the $ZZ \rightarrow l^- l^+$ collision our calculations result in the following formulas

$$\begin{split} |M_{\rm SM}|^2 &= \frac{g_Z^4}{(t-m_l^2)^2(s+t-m_l^2-2m_Z^2)^2} \{-2m_l^8[-300\cos(2\theta_w)+184\cos(4\theta_w) \\ &- 68\cos(6\theta_w)+17\cos(8\theta_w)+195] \\ &+ 4m_l^6[-4m_Z^2(-94\cos(2\theta_w)+59\cos(4\theta_w)-24\cos(6\theta_w) \\ &+ 6\cos(8\theta_w)+56)+163s+196t-8(33s+37t)\cos(2\theta_w) \\ &+ 18(9s+10t)\cos(4\theta_w)+(15s+16t)(\cos(8\theta_w)-4\cos(6\theta_w))] \\ &+ m_l^4[-2m_Z^4(-164\cos(2\theta_w)+128\cos(4\theta_w) \\ &+ 23(-4\cos(6\theta_w)+\cos(8\theta_w)+3)) \\ &+ 8m_Z^2(72s+65t-4(32s+27t)\cos(2\theta_w)+(84s+68t)\cos(4\theta_w) \\ &+ (10s+7t)(\cos(8\theta_w)-4\cos(6\theta_w))) - 301s^2-436t^2-668st \\ &+ 4(125s^2+260st+156t^2)\cos(2\theta_w)-8(39s^2+78st+46t^2)\cos(4\theta_w) \\ &- (31s^2+52st+28t^2)(\cos(8\theta_w)-4\cos(6\theta_w))] \\ &+ m_l^2[7s^3+50s^2t+92st^2+s(s^2-2st-4t^2)(\cos(8\theta_w)-4\cos(6\theta_w)) \\ &+ 80t^3-4(3s^3+14s^2t+24st^2+24t^3)\cos(2\theta_w) \\ &+ 8(s+2t)(s^2+st+3t^2)\cos(4\theta_w)+2m_Z^2(79s^2+252st+112t^2) \\ &- 4(31s^2+108st+52t^2)\cos(2\theta_w)+8(9s^2+34st+17t^2)\cos(4\theta_w) \\ &+ (5s^2+28st+16t^2)(\cos(8\theta_w)-4\cos(6\theta_w))) \\ &+ 2m_Z^4(-263s-414t+(428s+696t)\cos(2\theta_w)-16(16s+27t)\cos(4\theta_w) \\ &- 21(s+2t)(\cos(8\theta_w)-4\cos(6\theta_w))) \\ &+ 2m_Z^6(-156\cos(2\theta_w)+96\cos(4\theta_w)-36\cos(6\theta_w)+9\cos(8\theta_w)+91)] \\ &- [-28\cos(2\theta_w)+16\cos(4\theta_w)-4\cos(6\theta_w)+\cos(8\theta_w)+19] \\ \times [4m_Z^8-4m_Z^6(s+3t)+m_Z^4(s^2+6st+14t^2)-2m_Z^2t(s+2t)^2 \\ &+ t(s+t)(s^2+2st+2t^2)]\}, \end{split}$$

$$|M_{\rm KK}|^2 = \frac{1}{288} |S(s)|^2 \{-72m_Z^8 + 6m_Z^6(-40m_l^2 + 9s + 48t) - 4m_Z^4[-2m_l^2(5s + 96t) + 136m_l^4 + 9t(7s + 12t)] + 3m_Z^2[-80m_l^6 + m_l^4(256t - 14s) - 4m_l^2(s^2 + 29st + 68t^2) + 9s^3 + 42s^2t + 114st^2 + 96t^3)] - 36[m_l^4 - 2m_l^2t + t(s + t)] \times (2m_l^4 - 4m_l^2t + s^2 + 2st + 2t^2)\},$$
(C.5)

$$\begin{split} |M_{\rm int}|^2 &= -\frac{g_Z^2[S(s) + S^*(s)]}{96(t - m_l^2)(s + t - m_l^2 - 2m_Z^2)} \{-12m_l^8[\cos(4\theta_w) - 2\cos(2\theta_w)] \\ &+ 2m_l^6[(16m_Z^2 + 3(3s + 4t)(\cos(4\theta_w) - 2\cos(2\theta_w)) + 6(s - 2t)] \\ &+ 2m_l^4[-m_Z^2((9s + 4t)(\cos(4\theta_w) - 2\cos(2\theta_w)) \\ &+ 24s + 60t) + 8m_Z^4(-2\cos(2\theta_w) + \cos(4\theta_w) + 4) + 6(3s^2 + st + 6t^2) \\ &+ 3s(3s - 4t)(\cos(4\theta_w) - 2\cos(2\theta_w))] \\ &+ m_l^2[4m_Z^2((2s^2 - 3st + 14t^2)(\cos(4\theta_w) - 2\cos(2\theta_w)) \\ &+ 2s^2 + 5st + 46t^2) + 4m_Z^4((s - 10t)(\cos(4\theta_w) - 2\cos(2\theta_w)) \\ &+ 8m_Z^6(-2\cos(2\theta_w) + \cos(4\theta_w) + 5) \\ &- 3(3s^3 + 10s^2t + 2(s^3 + 2s^2t + 3st^2 + 4t^3) \\ &\times (\cos(4\theta_w) - 2\cos(2\theta_w)) + 24st(s + t))] \\ &+ 6[-2\cos(2\theta_w) + \cos(4\theta_w) + 2][2m_Z^8 + m_Z^6(s - 8t) \\ &+ m_Z^4(-s^2 + 2st + 12t^2) - m_Z^2t(s^2 + 7st + 8t^2) \\ &+ t(s + t)(s^2 + 2st + 2t^2)]\} \,, \end{split}$$

where θ_w is the Weinberg angle, $g_Z = e/[\sin(\theta_w)\cos(\theta_w)]$, and m_Z is the mass of the Z boson. Because of conservation of helicity, in the massless limit $m_l = m_Z = 0$ s-channel graviton amplitudes squared (C.2) and (C.5) are proportional to the factor $t(s+t) = s(\sin\theta)^2/4$, where θ is the scattering angle.

The SM squared amplitude for the $\gamma Z \to l^+ l^-$ collision looks like

$$\begin{split} |M_{\rm SM}|^2 &= \frac{4g_Z^4}{(t-m_l^2)^2(s+t-m_l^2-m_Z^2)^2} \\ &\times \{-2m_l^8[184\cos(4\theta_w)+17\cos(8\theta_w)-300\cos(2\theta_w) \\ &- 68\cos(6\theta_w)+195] + 4m_l^6[-2(59\cos(4\theta_w)+6\cos(8\theta_w)-94\cos(2\theta_w) \\ &- 24\cos(6\theta_w)+56)m_Z^2+163s+196t-8(33s+37t)\cos(2\theta_w) \\ &+ 18(9s+10t)\cos(4\theta_w)+(15s+16t)(\cos(8\theta_w)-4\cos(6\theta_w))] \\ &+ m_l^4[(68\cos(2\theta_w)+44\cos(6\theta_w)-56\cos(4\theta_w)-11\cos(8\theta_w)-25)m_Z^4 \\ &+ 4[72s+65t-4(32s+27t)\cos(2\theta_w)+(84s+68t)\cos(4\theta_w) \\ &+ (10s+7t)(\cos(8\theta_w)-4\cos(6\theta_w))]m_Z^2-301s^2-436t^2-668st \\ &+ 4(125s^2+260ts+156t^2)\cos(2\theta_w)-8(3s^2+78st+46t^2)\cos(4\theta_w) \\ &- (31s^2+52st+28t^2)(\cos(8\theta_w)-4\cos(6\theta_w))] \\ &+ m_l^2[7s^3+50s^2t+92st^2+(s^2-2st-4t^2)(\cos(8\theta_w) \\ &- 4\cos(6\theta_w))s+80t^3+(79s^2+252st+112t^2 \\ &+ ((5(\cos(8\theta_w)-4\cos(6\theta_w)+11)+56\cos(4\theta_w)) \\ &- 92(\cos(2\theta_w)))m_Z^2-141s-226t+4(57s+94t)(\cos(2\theta_w)) \\ &- 8(17s+29t)(\cos(4\theta_w))-11(s+2t)(\cos(8\theta_w)-4\cos(6\theta_w)))m_Z^2 \\ &- 4(31s^2+108st+52t^2)\cos(2\theta_w)+8(9s^2+34st+17t^2)\cos(4\theta_w) \\ &+ (5s^2+28st+16t^2)(\cos(8\theta_w)-4\cos(6\theta_w)))m_Z^2 \\ &- 4(3s^3+14s^2t+24st^2+24t^3)\cos(2\theta_w) \\ &+ 8(s+2t)(s^2+st+3t^2)\cos(4\theta_w)] \\ &- t[16\cos(4\theta_w)+\cos(8\theta_w)-28\cos(2\theta_w)-4\cos(6\theta_w)+19] \\ &\times (s+t-m_Z^2)(s^2+2t^2+2st-2tm_Z^2+m_Z^4)\} . \end{split}$$

The contribution to the $\gamma Z \rightarrow l^+ l^-$ collision from the gravitons G is zero, since there is no γZG vertex. Note that in the limit $m_l = m_Z = 0$ all squared amplitudes depend on variables s and t(s + t) = tu, where u is the Mandelstam variable.

For the $W^+W^- \rightarrow l^-l^+$ collision we have

$$|M_{\rm SM}|^2 = -\frac{4g_e^2 g_w^2}{s^2 (s - m_Z^2)^2} [(2s - m_Z^2)^2 (\cos \theta_w)^2 (3sm_l^2 - 10tm_l^2 + 14m_l^2 m_W^2 + 5m_l^4 + 2sm_W^2 - 10tm_W^2 + 5m_W^4 + 4s^2 + 5st + 5t^2)] + \frac{4g_e g_w^3}{st(s - m_Z^2)} [(2s - m_Z^2) \cos \theta_w \times (sm_l^2 + tm_l^2 + m_l^2 m_W^2 - 2m_l^4 - sm_W^2 - 2tm_W^2 + m_W^4 + 3st + t^2)] - \frac{4g_w^4}{t^2} [m_l^4 - 2m_l^2 m_W^2 - 2tm_W^2 + m_W^4 + st + t^2], \qquad (C.8)$$

where $g_w = g_e / \sin \theta_w$, and a neutrino mass is taken to be zero.

The interference term looks like

$$|M_{\rm int}|^2 = [S(s) + S^*(s)] \Big[\frac{g_e g_w (2s - m_Z^2) \cos \theta_w}{2s(s - m_Z^2)^2} (-2m_l^2 - 2m_W^2 + s + 2t) \\ \times (sm_l^2 - 4tm_l^2 + 4m_l^2 m_W^2 + 2m_l^4 + sm_W^2 - 4tm_W^2 + 2m_W^4 + s^2 + 2st + 2t^2) \\ - \frac{g_w^2}{24t} (3s^2m_l^2 - 6sm_l^2m_W^2 + 3sm_l^4 + 10tm_l^2m_W^2 - 18tm_l^4 + 4m_l^4m_W^2 \\ - 10m_l^2m_W^4 + 12m_l^6 - 3s^2m_W^2 - 18stm_W^2 + 3sm_W^4 - 18t^2m_W^2 + 18tm_W^4) \\ - 6m_W^6 + 9s^2t + 15st^2 + 6t^3 \Big].$$
(C.9)

Finally, the squared amplitude $|M_{\rm KK}(W^+W^- \to l^-l^+)|^2$ is obtained from eq. (C.5) by the replacement $m_Z \to m_W$.

Appendix D. Squared amplitudes for $l^+l^- \rightarrow l^+l^-$ scattering

Here we present the result of our calculations of the squared amplitudes for the $l^+l^- \rightarrow l^+l^-$ process (both incoming and outgoing leptons have the same flavor). The SM squared amplitude has both the Z boson and photon contributions. The latter one is given by the formula

$$|M_{\rm SM}|^2 = \frac{16e^4}{s^2t^2} [m_l^4(5s^2 + 11st + 5t^2) + 4m_l^2(s+t)(s^2 + 5st + t^2) + (s^2 + st + t^2)^2], \qquad (D.1)$$

where m_l is the lepton mass. Note that the photon contribution to $|M_{\rm SM}|^2$ is dominant. That is why, we do not present (rather complicated) analytical expression for the Z boson contribution to $|M_{\rm SM}|^2$. The graviton squared amplitude is defined by KK graviton exchanges in s- and t-channels,

$$|M_{\rm KK}|^2 = \frac{1}{4608} \{ |S(s)|^2 F_1(s,t) + |S(t)|^2 F_1(t,s) + [S(s)S(t)^* + S(s)^*S(t)] F_2(s,t) \} .$$
(D.2)

Finally, the interference term of $|{\cal M}|^2$ is equal to (neglecting small Z boson contribution)

$$|M_{\rm int}|^2 = -\frac{e^2}{24st} \{ [S(s) + S(t)^*] F_3(s,t) + [S(s)^* + S(t)] F_3(t,s) \} .$$
(D.3)

Here S(s) is defined by eqs. (18), (19), and the following functions are introduced

$$F_1(s,t) = 6656m_l^8 + m_l^6(8576s + 10752t) + m_l^4(3440s^2 + 7296t^2 + 10752st) + m_l^2(360s^3 + 2376s^2t + 4320st^2 + 2304t^3) + 9s^4 + 90s^3t + 378s^2t^2 + 576st^3 + 288t^4 ,$$
(D.4)

$$F_{2}(s,t) = 7552m_{l}^{8} + 15168m_{l}^{6}(s+t) + m_{l}^{4}(7248(s^{2}+t^{2}) + 14968st) + m_{l}^{2}(1032(s^{3}+t^{3}) + 3690st(s+t)) + 36(s^{4}+t^{4}) + 225st(s+t) + 378s^{2}t^{2},$$
(D.5)

$$F_{3}(s,t) = m_{l}^{6}(576s + 512t) + m_{l}^{4}(552s^{2} + 448t^{2} + 1120st) + m_{l}^{2}(86s^{3} + 360s^{2}t + 432st^{2} + 144t^{3}) + 3s^{4} + 21s^{3}t + 45s^{2}t^{2} + 48st^{3} + 24t^{4},$$
(D.6)

where m_l is the lepton mass. Note that $F_2(s,t) = F_2(t,s)$.

References

 L. Randall and R. Sundrum, Large mass hierarchy from a small extra dimension, Phys. Rev. Lett. 83, 3370 (1999) [arXiv:hep-ph/9905221].

- [2] A.M. Sirunyan et al. (CMS Collaboration), Search for physics beyond the standard model in high-mass diphoton events from protonproton collisions at $\sqrt{s} = 13$ TeV, Phys. Rev. D **98**, 092001 (2018) [arXiv:1809.00327].
- [3] A.M. Sirunyan et al. (CMS Collaboration), Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV, JHEP 07, 7 (2021) [arXiv:2103.02708].
- [4] G. Aad et al. (ATLAS Collaboration), Search for resonances decaying into photon pairs in 139 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector Phys. Lett. 822 (2021) 136651 [arXiv:2102.13405].
- [5] G.F. Giudice, T. Plehn, and A. Strumia, Graviton collider effects in one and more large extra dimensions, Nucl. Phys. B 706, 455 (2005) [arXiv:hep-ph/0408320].
- [6] A.V. Kisselev and V.A. Petrov, Gravi-Reggeons and trans-Planckian scattering in models with one extra dimension, Phys. Rev. D 71, 124032 (2005) [arXiv:hep-ph/0504203].
- [7] A.V. Kisselev, Generalization of the Randall-Sundrum solution, Nucl. Phys. B 909, 218 (2016) [arXiv:1512.01091].
- [8] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, *The Hierarchy problem and new dimensions at a millimeter*, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315].
- [9] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phenomenology, astrophysics, and cosmology of theories with submillimeter dimensions and TeV scale quantum gravity*, Phys. Rev. D 59, 086004 (1999) [arXiv:hep-ph/9807344].
- [10] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, New dimensions at a millimeter to a fermi and superstrings at a TeV, Phys. Lett. B 436, 257 (1998) [arXiv:hep-ph/9804398].
- [11] A.V. Kisselev, RS model with a small curvature and two-photon production at the LHC, JHEP 09, 039 (2008) [arXiv:0804.3941].

- [12] A.V. Kisselev, Randall-Sundrum model with a small curvature and dimuon production at the LHC, JHEP 04, 025 (2013) [arXiv:1210.3238].
- [13] F.F. Tikhonin, On the effects of the clashing μ-meson beams (in Russian), JINR Report P2-4120, Dubna, 1968; On the effects at colliding mu-meson beams, arXiv:0805.3961.
- [14] G.I. Budker, Accelerators and colliding beams. In Proceedings of the 7th International Conference on High-Energy Accelerators. (HEACC 1969), 27 August – 2 September 1969, Yerevan, USSR. Vol. 1, pp. 33-39 (1970).
- [15] A.N. Skrinsky and V.V. Parkhomchuk, Cooling methods for beams of charged particles (in Russian), Sov. J. Part. Nucl. 12, 223 (1981) [Fiz. Elem. Chast. Atom. Yadra 12, 557 (1981)].
- [16] D. Neuffer, Principles and applications of muon cooling, in Proceedings of the 12th International Conference on High Energy Accelerators (HEACC 1983), Fermilab, Batavia, USA. 11-16 August 1983. Conf. Proc. C 830811, pp. 481-484 (1983).
- [17] A. Blondel, J.R. Ellis, and B. Autin, Prospective study of muon storage rings at CERN. In CERN Yellow Reports: Monographs. CERN: Geneva, Switzerland, 1999.
- [18] C.M. Ankenbrandt, M. Atac, B. Autin, V.I. Balbekov, and V.D. Barger, Status of muon collider research and development and future plans, Phys. Rev. ST Accel. Beams 2, 081001 (1999) [arXiv:physics/9901022].
- [19] J.P. Delahaye *et al.* (Muon Collider Working Group), *Muon colliders*, arXiv:1901.06150.
- [20] M. Boscolo, J.P. Delahaye, and M. Palmer, *The future prospects of muon colliders and neutrino factories*, Rev. Accel. Sci. Tech. **10**, 189 (2019) [arXiv:1808.01858].
- [21] D. Buttazzo, R. Franceschini, and A. Wulzer, Two paths towards precision at a very high energy lepton collider, JHEP 05, 219 (2021) [arXiv:2012.11555].
- [22] K.R. Long, D. Lucchesi, M.A. Palmer, N. Pastrone, D. Schulte, and V. Shiltsev, *Muon colliders to expand frontiers of particle physics*, Nature Phys. **17**, 289 (2021) [arXiv:2007.15684].

- [23] V. Barger, M.S. Berger, J.F. Gunion, and T. Han, *Higgs boson physics in the s-channel at* $\mu^+\mu^-$ *colliders*, Phys. Rep. **286**, 1 (1997) [arXiv:hep-ph/9602415].
- [24] M. Chiesa, Measuring the quartic Higgs self-coupling at a multi-TeV muon collider, JHEP 09, 098 (2020) [arXiv:2003.13628].
- [25] R. Franceschini and M. Greco, Higgs and BSM physics at the future muon collider, Symmetry 13, 851 (2021) [arXiv:2104.05770].
- [26] P. Bandyopadhyay and A. Costantini, Obscure Higgs boson at colliders, Phys. Rev. D 103, 015025 (2021) [arXiv:2010.02597].
- [27] T. Han, D. Liu, I. Low, and X. Wang, Electroweak couplings of the Higgs boson at a multi-TeV muon collider, Phys. Rev. D 103, 013002 (2021) [arXiv:2008.12204].
- [28] T. Han, S. Li, S. Su, W. Su, and Y. Wu, *Heavy Higgs bosons in 2HDM at a muon collider*, Phys. Rev. D 104, 055029 (2021) [arXiv:2102.08386].
- [29] A. Costantini, (New) physics at a multi-TeV μ collider, arXiv:2111.02507.
- [30] R. Capdevilla, F. Meloni, R. Simoniellod, and J. Zurita, Hunting wino and higgsino dark matter at the muon collider with disappearing tracks, JHEP 06, 133 (2021) [arXiv:2102.11292].
- [31] T. Han, Z. Liu, L.-T. Wang, and X. Wang, WIMPs at high energy muon colliders, Phys. Rev. D 103, 075004 (2021) [arXiv:2009.11287].
- [32] K. Black et al., Prospects for heavy WIMP dark matter searches at muon colliders, arXiv:2205.10404.
- [33] R. Franceschini and X. Zhao, Going all the way in the search for WIMP dark matter at the muon collider through precision measurements, aeXiv:2212.11900.
- [34] A. Jueid and S. Nasri, Lepton portal dark matter at muon colliders: Total rates and generic features for phenomenologically viable scenarios, arXiv:2301.12524.

- [35] A. Costantini et al., Vector boson fusion at multi-TeV muon colliders, JHEP 10, 080 (2020) [arXiv:2005.10289].
- [36] P. Asadi, R. Capdevilla, C. Cesarotti, and S. Homiller, Searching for leptoquarks at future muon colliders, JHEP 10, 182 (2021) [arXiv:2104.05720].
- [37] F. Bossi and P. Ciafaloni, Lepton flavor violation at muon-electron colliders, JHEP 10, 033 (2020) [arXiv:2003.03997].
- [38] S. Homiller, Q. Lu, and M. Reece, Complementary signals of lepton flavor violation at a high-energy muon collider, JHEP **07**, 036 (2022) [arXiv:2203.08825].
- [39] G. Haghighat, M.M. Najafabadi, Search for lepton-flavor-violating decays of the tau lepton at a future muon collider, arXiv:2204.04433.
- [40] Q. Guo, L. Gao, Y. Mao, and Q. Li, Search for vector-like leptons at a muon collider, arXiv:2204.01885.
- [41] P. Li, Z. Liu, and K.-F. Lyu, *Heavy neutral leptons at muon colliders*, arXiv:2301.07117.
- [42] K. Mekala, J. Reuter, and A.F. Zarnecki, *Optimal search reach for heavy* neutral leptons at a muon collider, arXiv:2301.02602.
- [43] I. Chakraborty, H. Roy, and T. Srivastava, Searches for heavy neutrinos at multi-TeV muon collider: a resonant leptogenesis perspective, arXiv:2206.07037.
- [44] M. Chen and D. Liu, Top Yukawa coupling at the muon collider, arXiv:2212.11067.
- [45] P. Bredt, W. Kilian, J. Reuter, and P. Stienemeier, NLO electroweak corrections to multi-boson processes at a muon collider, JHEP 12, 138 (2022) [arXiv:2208.09438].
- [46] R. Capdevilla, D. Curtin, Y. Kahn, and G. Krnjaic, Discovering the physics of (g – 2)_μ at future muon colliders, Phys. Rev. D 103, 075028 (2021) [arXiv:2006.16277].

- [47] W. Yin and M. Yamaguchi, Muon g 2 at multi-TeV muon collider, Phys. Rev. D 106, 033007 (2022) [arXiv:2012.03928].
- [48] J. Arakawa, A. Rajaraman, T. Sui, and T.M.P. Tait, *Probing muon* g-2 at a future muon collider, arXiv:2208.14464.
- [49] S.C. Inan, and A.V. Kisselev, Probe of axion-like particles in vector boson scattering at a muon collider, J. Phys. G: Nucl. Part. Phys. (2023) [arXiv:2207.03325].
- [50] B. Abbot *et al.*, Anomalous quartic gauge couplings at a muon collider, in Proceedings of the US Community Study on the Future of Particle Physics (Snowmass 2021), arXiv:2203.08135.
- [51] J.-C. Yang, X.-Y. Han, Z.-B. Qin, T. Li, and Y.-C. Guo, Measuring the anomalous quartic gauge couplings in the $W^+W^- \rightarrow W^+W^-$ process at muon collider using artificial neural networks, JHEP **09**, 074 (2022).
- [52] J.-C. Yang, Z.-B. Qing, X.-Y. Han, Y.-C. Guo, and T. Li, Tri-photon at muon collider: a new process to probe the anomalous quartic gauge couplings, JHEP 07, 053 (2022) [arXiv:2204.08195].
- [53] A. Senol, S. Spor, E. Gurkanli, V. Cetinkaya, H. Denizli, and M. Köksal, Model-independent study on the anomalous ZZγ and Zγγ couplings at the future muon collider, Eur. Phys. J. Plus 137, 1354 (2022).
- [54] S. Spor, Probe of the anomalous neutral triple gauge couplings in photoninduced collision at future muon colliders, arXiv:2207.11585.
- [55] K.M. Black et al., Muon collider forum report, arXiv:2209.01318.
- [56] H. Davoudiasl, J.L. Hewett, and T.G. Rizzo, Experimental probes of localized gravity: On and off the wall, Phys. Rev. D 63, 075004 (2001) [arXiv:hep-ph/0006041].
- [57] G.F. Giudice, R. Ratazzi, and J.D. Wells, Quantum gravity and extra dimensions at high-energy colliders, Nucl. Phys. B 544, 3 (1999) [arXiv:hep-ph/9811291].
- [58] T. Han, J.D. Lykken, and R.-J. Zhank, Kaluza-Klein states from large extra dimensions, Phys. Rev. D 59, 105006 (1999) [arXiv:hep-ph/9811350].

- [59] S. Atağ, S.C. İnan, and İ Şahin, Extra dimensions in photon-induced two lepton final states at the CERN-LHC, Phys. Rev. D 80, 075009 (2009) [arXiv:0904.2687].
- [60] A.V. Kisselev, Virtual gravitons and brane field scattering in the Randall-Sundrum model with a small curvature, Phys. Rev. D 73, 024007 (2006) [arXiv:hep-th/0507145].
- [61] V.M. Budnev, I.F. Ginzburg, G.V. Meledin, and V.G. Serbo, The two photon particle production mechanism. Physical problems. Applications. Equivalent photon approximation, Phys. Rep. 15, 181 (1975).
- [62] J. Lindfors, Luminosity functions for W^{\pm} and Z^{0} initiated processes, Z. Phys. C **35**, 355 (1987).
- [63] R. Ruiz, A. Costantini, F. Maltoni, and O. Mattelaer, The effective vector boson approximation in high-energy muon collisions, JHEP 06, 114 (2022) [arXiv:2111.02442].
- [64] S.C. Inan and A.V. Kisselev. Muon pair production via photon-induced scattering at the CLIC in models with extra dimensions, Nucl. Phys. B 956, 115047 (2020) [arXiv:1907.12824].
- [65] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, Eur. Phys. J. C 71, 1554 (2011) [Erratum ibid. 73, 2501 (2013)] [arXiv:1007.1727].
- [66] E.E. Boos, Yu.A. Kubyshin, M.N. Smolyakov, and I.P. Volobuev, Effective Lagrangians for physical degrees of freedom in the Randall-Sundrum model, Class. Quant. Grav. 19 (2002) 4591 [hep-th/0202009].
- [67] V.A. Rubakov, Larg and infinite extra dimensions, Phys. Usp. 44 (2001)
 871 [Usp. Fiz. Nauk 171, 913 (2001)] [hep-ph/0104152].