

NEAR-OPTIMAL RESTRICTION ESTIMATES FOR CANTOR SETS ON THE PARABOLA

DONGGEUN RYOU

ABSTRACT. For any $0 < \alpha < 1$, we construct Cantor sets on the parabola of Hausdorff dimension α such that they are Salem sets and each associated measure ν satisfies the estimate $\|\widehat{f d\nu}\|_{L^p(\mathbb{R}^2)} \leq C_p \|f\|_{L^2(\nu)}$ for all $p > 6/\alpha$ and for some constant $C_p > 0$ which may depend on p and ν . The range $p > 6/\alpha$ is optimal except for the endpoint. The proof is based on the work of Laba and Wang on restriction estimates for random Cantor sets and the work of Shmerkin and Suomala on Fourier decay of measures on random Cantor sets. They considered fractal subsets of \mathbb{R}^d , while we consider fractal subsets of the parabola.

1. INTRODUCTION

Given $f : \mathbb{R}^d \rightarrow \mathbb{C}$, the Fourier transform of f is defined by

$$\widehat{f}(\xi) = \int_{\mathbb{R}^d} f(x) e^{-2\pi i x \cdot \xi} dx \quad \forall \xi \in \mathbb{R}^d.$$

Let μ be a compactly supported measure on \mathbb{R}^d . If f is μ -integrable, $\widehat{f d\mu}$ is defined by

$$\widehat{f d\mu}(\xi) = \int f(x) e^{-2\pi i x \cdot \xi} d\mu(x) \quad \forall \xi \in \mathbb{R}^d.$$

We consider a Borel probability measure μ in \mathbb{R}^d and assume that it satisfies the estimate

$$\|\widehat{f d\mu}\|_{L^p(\mathbb{R}^d)} \leq C_p \|f\|_{L^2(\mu)} \quad \forall f \in L^2(\mu) \quad (1.1)$$

for some constant C_p which depends on p . It is easy to show that (1.1) holds when $p = \infty$. However, we can say more if we focus on more specific measures. One typical example is a surface carried measure on the sphere or on the paraboloid. Then, (1.1) holds when $p \geq 2(d+1)/(d-1)$ and the range of p is sharp. This is the result of the Stein-Tomas theorem (see for example [24]).

Let $B(x, r)$ be a closed ball of radius r centered at x . If we have assumptions on the Fourier decay of μ and the upper bound of $\mu(B(x, r))$, the following result generalizes the Stein-Tomas theorem.

Theorem 1.1. *Let μ be a Borel probability measure on \mathbb{R}^d . Assume that there exist $\alpha, \beta \in (0, d)$ and $C_1, C_2 \geq 0$ such that*

$$\mu(B(x, r)) \leq C_1 r^\alpha \quad \forall x \in \mathbb{R}^d, r > 0, \quad (1.2)$$

$$|\widehat{\mu}(\xi)| \leq C_2 (1 + |\xi|)^{-\beta/2} \quad \forall \xi \in \mathbb{R}^d. \quad (1.3)$$

Then, the estimate (1.1) holds when $p \geq (4d - 4\alpha + 2\beta)/\beta$ for some constant $C_p \geq 0$ which depends on p, C_1 and C_2 .

Date: November 17, 2023.

2020 Mathematics Subject Classification. 42B10 (primary) 28A80 (secondary).

Key words and phrases. Random Cantor sets, Restriction estimate, Salem set.

Mockenhaupt [21] and Mitsis [19] proved that (1.1) holds when $p > (4d - 4\alpha + 2\beta)/\beta$ and the endpoint result was proved by Bak and Seeger [1]. If μ is a surface carried measure on the paraboloid, $\alpha = \beta = d - 1$. Thus, Theorem 1.1 recovers the range $p \geq 2(d + 1)/(d - 1)$.

Define the *critical exponent* p_c of the measure μ by

$$p_c = \inf\{p : (1.1) \text{ holds}\}.$$

Theorem 1.1 means $p_c \leq (4d - 4\alpha + 2\beta)/\beta$. This upper bound of p_c is known to be optimal when $d - 1 < \beta \leq \alpha < d$ in the sense that there exists a measure μ such that it satisfies the assumptions of Theorem 1.1 and $p_c = (4d - 4\alpha + 2\beta)/\beta$. When $d = 1$, Hambrook and Łaba [10] constructed a measure such that (1.3) holds for any β arbitrarily close to but smaller than α and $p_c = (4 - 2\alpha)/\alpha$. Chen [6] extended this result to general α and β such that $0 \leq \beta \leq \alpha \leq 1$. In higher dimensions, it was done in [11] by Hambrook and Łaba.

However, there exist some measures μ such that $p_c < (4d - 4\alpha + 2\beta)/\beta$. In [10], Hambrook and Łaba showed that $p_c \geq 2d/\alpha_0$ if μ is supported on a set of Hausdorff dimension α_0 . Note that $0 \leq \alpha, \beta \leq \alpha_0$. If (1.2) and (1.3) hold for values arbitrarily close to α_0 , the support of μ is called a *Salem set*. If $0 < \alpha_0 < d$, we have

$$\frac{2d}{\alpha_0} < \frac{4d - 2\alpha_0}{\alpha_0} \leq \frac{4d - 4\alpha + 2\beta}{\beta}.$$

Thus, $2d/\alpha_0$ is smaller than $(4d - 4\alpha + 2\beta)/\beta$ even when μ is supported on a Salem set. Examples such that $p_c = 2d/\alpha_0$ were provided by Chen [5], Chen and Seeger [7], Shmerkin and Suomala [23] (see also [22]) and Łaba and Wang [15]. Therefore, the lower bound of p_c is optimal for all $0 < \alpha_0 < d$ in the sense that there exists a measure μ whose $p_c = 2d/\alpha_0$ for each α_0 .

In [5] and [7], Chen and Seeger constructed measures supported on a Salem set of Hausdorff dimension α_0 such that $p_c = 2d/\alpha_0$ where $\alpha_0 = d/k$ and k is an integer. They considered k -fold self-convolution of μ . Shmerkin and Suomala [23] constructed the example through random fractal measures. Their result covers when $d = 1$ and $1/2 < \alpha_0 < 1$. Łaba and Wang [15] constructed measures of Hausdorff dimension α_0 such that (1.3) holds for $\beta < \min(\alpha_0/2, 1)$ and $p_c = 2d/\alpha_0$. They used $\Lambda(p)$ -set and decoupling. Their construction works for all α_0 such that $0 < \alpha_0 < d$. However, in their result, the estimate (1.1) only holds when $p > 2d/\alpha_0$, while the results of Chen, Chen-Seeger, and Shmerkin-Suomala work even at the endpoint. Thus, it is still open whether, for any $0 < \alpha_0 < d$, there exists a measure μ such that its support has Hausdorff dimension α_0 and (1.1) holds for $p = 2d/\alpha_0$.

We also mention that all the results are related to a question raised by Mitsis [20] (see also Mattila [17]). If α is not an integer such that $0 < \alpha < d$, Mitsis asked if there is a compactly supported measure μ such that

$$C_1 r^\alpha \leq \mu(B(x, r)) \leq C_2 r^\alpha \quad \forall x \in \text{supp}(\mu), \forall r > 0$$

for some constants $C_1, C_2 > 0$ and

$$|\widehat{\mu}(\xi)| \leq |\xi|^{-\alpha/2} \quad \forall \xi \in \mathbb{R}^d.$$

A related result can also be found in [14]. However, there is no known result yet, including this paper, since we have $|\widehat{\mu}(\xi)| \leq C_{\alpha, \epsilon} |\xi|^{-\alpha/2 + \epsilon}$ at best where $C_{\alpha, \epsilon}$ is a constant depends on α and ϵ .

Throughout the paper, we denote by $X \lesssim Y$ when $X \leq CY$ for some constant $C > 0$ and we write $X \approx Y$ to denote that $X \lesssim Y$ and $Y \lesssim X$. If the constant C depends on a parameter such as ϵ , we write $X(\epsilon) \lesssim_\epsilon Y(\epsilon)$ instead of $X(\epsilon) \leq C(\epsilon)Y(\epsilon)$ where the constant $C(\epsilon)$ depends on ϵ . Depending on the context, $|\cdot|$ denotes the ℓ^2 norm of a vector in \mathbb{R}^d ,

the cardinality of a finite set or the d -dimensional Lebesgue measure of a subset in \mathbb{R}^d . We abbreviate $e^{2\pi i x}$ to $e(x)$.

Now, let us turn to our setting. Earlier works constructed fractal measures on \mathbb{R}^d , but we consider fractal measures on the parabola $\mathbb{P}^1 := \{(x, x^2) : 0 \leq x \leq 1\}$.

Theorem 1.2. *Let $0 < \alpha < 1$. There exists a Borel probability measure ν supported on a subset of \mathbb{P}^1 which satisfies the following.*

- (1) *The support of the measure ν has Hausdorff dimension α .*
- (2) *For any $0 < \epsilon < \alpha$, we have*

$$\nu(B(x, r)) \lesssim_{\alpha, \epsilon} r^{\alpha - \epsilon} \quad \forall x \in \mathbb{R}^2, \forall r > 0. \quad (1.4)$$

- (3) *For any $\epsilon > 0$,*

$$|\widehat{\nu}(\xi)| \lesssim_{\alpha, \epsilon} (1 + |\xi|)^{-\alpha/2 + \epsilon} \quad \forall \xi \in \mathbb{R}^2. \quad (1.5)$$

- (4) *For every $p > 6/\alpha$, we have the estimate*

$$\left\| \widehat{f d\nu} \right\|_{L^p(\mathbb{R}^2)} \lesssim_p \|f\|_{L^2(\nu)} \quad \forall f \in L^2(\nu). \quad (1.6)$$

Equivalently, for any $1 \leq q < 6/(6 - \alpha)$ we have

$$\left\| \widehat{f} \right\|_{L^q(\nu)} \lesssim_q \|f\|_{L^q(\mathbb{R}^2)} \quad \forall f \in L^q(\mathbb{R}^2).$$

Note that the support of ν is a Salem set and it is a subset of the parabola, which is also a Salem set. Theorem 1.2 is sharp except for the endpoint in the following sense.

Theorem 1.3. *Let $\mathbb{P}^{d-1} := \{(x, |x|^2) : x \in [0, 1]^{d-1}\}$ and $0 < \alpha < d - 1$. Assume that ν is a Borel probability measure supported on a subset of \mathbb{P}^{d-1} whose Hausdorff dimension is α . Assume that*

$$\nu(B(x, r)) \lesssim_{\alpha} r^{\alpha} \quad \forall x \in \mathbb{P}^{d-1}, \forall r > 0. \quad (1.7)$$

Then, the estimate

$$\left\| \widehat{f d\nu} \right\|_{L^p(\mathbb{R}^2)} \lesssim_{p, q} \|f\|_{L^q(\nu)} \quad \forall f \in L^q(\nu) \quad (1.8)$$

cannot hold if $p < q'(d + 1)/\alpha$ where $1/q + 1/q' = 1$.

When $d = 2$ and $q = 2$, Theorem 1.3 implies that $p_c \geq 6/\alpha$. Since ν is on the parabola, the lower bound of p_c increased from $4/\alpha$ to $6/\alpha$ compared with the result of Hambrook and Łaba [10]. Also, since $0 < \alpha < 1$, we have $6/\alpha < (8 - 2\alpha)/\alpha$. Thus, it is smaller than the upper bound of p_c obtained from Theorem 1.1.

1.1. Outline of the paper. The proof was inspired by the work of Łaba and Wang [15], but we cannot apply it directly. The main obstacle is that we cannot make use of dual cubes as in [15]. For example, they considered a Cantor set which can be covered by $R^{\alpha + \epsilon}$ number of cubes Q of side length R^{-1} for any $R > 0$. Let $\psi_Q = \psi(R(x - x_Q))$ where ψ is a compactly supported Schwartz function and x_Q is the center of Q . For any Q , $\widehat{\psi_Q}$ is essentially supported on a cube of side length $O(R)$ centered at the origin.

Now, let us consider a subset of the parabola as follows. First, consider a Cantor set on $[0, 1]$ which can be covered by $R^{\alpha + \epsilon}$ number of intervals I of length $R^{-1/2}$. Then, we can consider a parallelogram Ω_I of sides $O(R^{-1/2})$ and $O(R^{-1})$ which contains the subset of the parabola above I , i.e. $\{(x, x^2) : x \in I\} \subseteq \Omega_I$. Let ψ_{Ω_I} be a Schwartz function supported on Ω_I . Specifically,

$$\psi_{\Omega_I} = \psi(R^{1/2}(x_1 - x_I), R(x_2 - (2x_I + R^{-1/2})(x_1 - x_I) - x_I^2))$$

where x_I is the center of I . Then, $\widehat{\psi_{\Omega_I}}$ is essentially supported on a parallelogram of sides $O(R^{1/2})$ and $O(R)$. However, the directions of long sides of them are all different, so that they overlap in a small cube of side $O(R^{1/2})$, while the union of these parallelograms is a cube of side $O(R)$. Thus, we need new arguments in addition to the modifications of the proof in [15].

In Section 2, we construct a Cantor set on $[0, 1]$ by using $\Lambda(p)$ -sets and consider the associated measure which is supported on a subset of parabola above the Cantor set. Let ν be the measure constructed in Section 2. Then ν is supported on a set of Hausdorff dimension α and satisfies (1.4). In Section 3, we obtain the decoupling inequality for the support of ν . We use $\Lambda(p)$ -sets and the result of [4]. In Section 4, we obtain the local restriction estimate for ν , which is an analogue of Corollary 2 in [15]. However, we use the mixed norm interpolation [2] in addition to the argument in [15]. In Section 5, we obtain the Fourier decay of ν . We only need a negative exponent to derive (1.6), but it turned out that almost surely ν has the Fourier decay arbitrary close to the optimal one, as in (1.5). We use the arguments in [13] and [22] in our setting. The local restriction estimate and the Fourier decay lead to the global restriction estimate in Section 6, which is (1.6). Tao's epsilon removal lemma [27] is used as in [15], but we simplify the proof of Lemma 9 in [15] in order to avoid the problem we described above. Then, we prove that Theorem 1.2 happens almost surely if ν is a random measure constructed in Section 6. Lastly, we prove Theorem 1.3 in Section 7.

Acknowledgements. The author would like to thank his advisor Alex Iosevich for many discussions of this work and encouragement. The author would also like to thank Shaoming Guo, Zane Kun Li for helpful conversations and thank Kyle Hambrook for the reference [13]. The author is grateful to the anonymous referees for many valuable comments and suggestions.

2. THE CONSTRUCTION OF A CANTOR SET

Let $\{n_j\}_{j \in \mathbb{N} \setminus \{0\}}$ be a sequence of positive integers, and let $N_j = n_0 \cdots n_j$. We assume the following conditions on n_j .

$$n_0 = 1 \quad \text{and} \quad 2 \leq n_1 \leq n_2 \leq \cdots \leq n_j \leq \cdots, \quad (2.1)$$

$$\forall a > 0, a^j \lesssim_a N_j, \quad (2.2)$$

$$\forall \epsilon > 0 \text{ and } \forall j \in \mathbb{N}, n_{j+1} \lesssim_\epsilon N_j^\epsilon, \quad (2.3)$$

$$\exists B \text{ such that } \forall j \in \mathbb{N}, N_{2j}^{1/2} N_j^{-1} \leq B^j. \quad (2.4)$$

For example, if $n_j \approx j$, all conditions from (2.1) to (2.4) are satisfied by Stirling's formula (see [9, p. 98]). The condition (2.4) was not assumed in [15], but it is necessary in Sections 3 and 4. We need the following theorem in order to use $\Lambda(p)$ -sets.

Theorem 2.1 (Existence of $\Lambda(p)$ -sets). *Let $p > 2$. For every $N \in \mathbb{N}$, there exists a set $S \subset \{0, 1, 2, \dots, N-1\}$ such that $N^{2/p} \leq |S| < N^{2/p} + 1$ and*

$$\left\| \sum_{a \in S} c_a e(ax) \right\|_{L^p([0,1])} \leq C_p \left(\sum_{a \in S} |c_a|^2 \right)^{1/2} \quad \forall \{c_a\}_{a \in S} \in \ell^2 \quad (2.5)$$

for some constant $C_p > 0$ which depends only on p , but not on N .

Theorem 2.1 was first proved by Bourgain [3] and another proof was given by Talagrand [25], see also [26, Section 19.3] for simpler proof.

Let $0 < \alpha < 1$ and for each $j \in \mathbb{N}$, let $\{t_j\}_{\mathbb{N}}$ be a sequence of integers such that

$$n_j^{\alpha/2} \leq t_j < n_j^{\alpha/2} + 1.$$

Define

$$\Sigma_j := \{S \subset [0, n_j^{1/2}) \cap \mathbb{Z} : |S| = t_j \text{ and (2.5) holds for } p = 2/\alpha, N = n_j^{1/2}\}.$$

The set Σ_j is non-empty because of Theorem 2.1.

Let $E_0 = [0, 1]$ and $A_0 = \{0\}$. For each $a \in A_j$, choose a set $S_{j+1,a} \in \Sigma_{j+1}$ and let

$$A_{j+1,a} = n_{j+1}^{1/2}a + S_{j+1,a} \quad A_{j+1} = \cup_{a \in A_j} A_{j+1,a} \quad E_{j+1} = N_{j+1}^{-1/2}(A_{j+1} + [0, 1]).$$

The set A_j is a subset of $\{0, 1, 2, \dots, N_j^{1/2} - 1\}$ and $[0, 1] \supseteq E_1 \supseteq \dots \supseteq E_j \supseteq E_{j+1}$. Let $E_\infty := \cap_{j=1}^\infty E_j$. Similarly, let $P_j := \{(x_1, x_1^2) : x_1 \in E_j\}$ and $P_\infty := \cap_{j=1}^\infty P_j$.

For each $j \in \mathbb{N} \cup \{0\}$, we define

$$\mu_j := \frac{1}{|E_j|} \mathbf{1}_{E_j}(x_1), \quad x_1 \in \mathbb{R}.$$

We identify the function μ_j with the absolutely continuous measure $\mu_j dx_1$ so that

$$\int g d\mu_j := \int g \mu_j dx_1.$$

From μ_j , we define the measure ν_j as follows.

$$\int f d\nu_j := \int f(x_1, x_1^2) d\mu_j.$$

For simplicity, define $\|\mu_j\| := \mu_j(\mathbb{R})$ and $\|\nu_j\| := \nu_j(\mathbb{R}^2)$. The measures μ_j and ν_j converge weakly as $j \rightarrow \infty$ to probability measures μ and ν supported on E_∞ and P_∞ respectively.

Lemma 2.2 (Lemma 6, [15]). *The measure μ and the set E_∞ constructed above satisfy the following.*

- (1) *The set E_∞ has Hausdorff dimension α .*
- (2) *For any $0 < \epsilon < \alpha$, we have*

$$\mu(B(x, r)) \lesssim_{\alpha, \epsilon} r^{\alpha - \epsilon} \quad \forall x \in \mathbb{R}, \forall r > 0.$$

It is easy to show that the same is true for ν .

Lemma 2.3. *The measure ν and the set P_∞ constructed above satisfy the following.*


- (1) *The set P_∞ has Hausdorff dimension α .*
- (2) *For any $0 < \epsilon < \alpha$, we have*

$$\nu(B(x, r)) \lesssim_{\alpha, \epsilon} r^{\alpha - \epsilon} \quad \forall x \in \mathbb{R}^2, \forall r > 0.$$

3. DECOUPLING INEQUALITIES

In this section, we will derive the decoupling estimates for E_∞ and P_∞ . We will define decoupling constants and discrete restriction constants, and we will verify relations between them. Since we need to look into E_j in different scales, we define the following.

For $0 \leq i \leq j$ and $a \in A_i$, we define $E_{j,i,a} = E_j \cap N_i^{-1/2}(a + [0, 1])$, see Figure 1. If $i = 0$, since $A_0 = \{0\}$, note that $E_{j,i,a} = E_{j,0,0} = E_j$. Let $\mathcal{P}_j(E_{j,i,a})$ be the partition of

FIGURE 1. $E_{j,i,a}$

$E_{j,i,a}$ into intervals of length $N_j^{-1/2}$ and $\widehat{f}_I(\xi_1) := \widehat{f}_I(\xi_1)\mathbf{1}_I(\xi_1)$ for $I \in \mathcal{P}_j(E_{j,i,a})$ and $\xi_1 \in \mathbb{R}$. Sometimes, we will write $I(b) \in \mathcal{P}_j(E_{j,i,a})$ for $N_j^{-1/2}(b + [0, 1]) \subseteq E_{j,i,a}$ for some $b \in A_j$.

For $p \geq 2$, let $D_p(E_{j,i,a})$ denote the best constant such that

$$\left\| \sum_{I \in \mathcal{P}_j(E_{j,i,a})} f_I \right\|_{L^p(\mathbb{R})} \leq D_p(E_{j,i,a}) \left(\sum_{I \in \mathcal{P}_j(E_{j,i,a})} \|f_I\|_{L^p(\mathbb{R})}^2 \right)^{1/2} \quad (3.1)$$

for any function f such that $\text{supp } \widehat{f} \subseteq E_{j,i,a}$ and let $D_p(E_{j,i}) = \sup_{a \in A_i} D_p(E_{j,i,a})$.

If $I \in \mathcal{P}_j(E_{j,i,a})$, there exist a unique element $b \in A_j$ such that $I(b) \in \mathcal{P}_j(E_{j,i,a})$. Thus, we define

$$\begin{aligned} \Omega_I &= \{\xi \in \mathbb{R}^2 : bN_j^{-1/2} \leq \xi_1 \leq (b+1)N_j^{-1/2}, \\ &\quad |\xi_2 - (2b+1)N_j^{-1/2}(\xi_1 - bN_j^{-1/2}) - b^2N_j^{-1}| \leq N_j^{-1}\} \end{aligned}$$

and $P_{j,i,a} = \cup_{I \in \mathcal{P}_j(E_{j,i,a})} \Omega_I$ and $f_{\Omega_I} = \widehat{f}(\xi)\mathbf{1}_{\Omega_I}(\xi)$ for $\xi \in \mathbb{R}^2$. Let $D_p(P_{j,i,a})$ be the best constant such that

$$\left\| \sum_{I \in \mathcal{P}_j(E_{j,i,a})} f_{\Omega_I} \right\|_{L^p(\mathbb{R}^2)} \leq D_p(P_{j,i,a}) \left(\sum_{I \in \mathcal{P}_j(E_{j,i,a})} \|f_{\Omega_I}\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2} \quad (3.2)$$

for any function f such that $\text{supp } \widehat{f} \subseteq P_{j,i,a}$. Similarly, let $D_p(P_{j,i}) = \sup_{a \in A_i} D_p(P_{j,i,a})$.

We also need discrete versions of $D_p(E_{j,i})$ and $D_p(P_{j,i})$. For $a \in A_i$, let us define

$$A_{j,i,a} = A_j \cap (N_j N_i^{-1})^{1/2}(a + [0, 1]),$$

i.e., $A_{j,i,a}$ is the set of j -th level descendants of $a \in A_i$. Then, let $K_p(E_{j,i,a})$ be the best constant such that

$$\left\| \sum_{b \in A_{j,i,a}} c_b e(bx) \right\|_{L^p([0,1])} \leq K_p(E_{j,i,a}) \left(\sum_{b \in A_{j,i,a}} |c_b|^2 \right)^{1/2}$$

for any $\{c_b\}_{b \in A_{j,i,a}} \in \ell^2$ and $K_p(E_{j,i}) := \sup_{a \in A_i} K_p(E_{j,i,a})$. Similarly, let $K_p(P_{j,i,a})$ be the best constant such that

$$\left\| \sum_{b \in A_{j,i,a}} c_b e(bx_1 + b^2x_2) \right\|_{L^p([0,1]^2)} \leq K_p(P_{j,i,a}) \left(\sum_{b \in A_{j,i,a}} |c_b|^2 \right)^{1/2}$$

for any $\{c_b\}_{b \in A_{j,i,a}} \in \ell^2$ and $K_p(P_{j,i}) := \sup_{a \in A_i} K_p(P_{j,i,a})$.

In short, $D_p(E_{j,i,a})$ and $D_p(P_{j,i,a})$ are decoupling constants for $E_{j,i,a}$ and $P_{j,i,a}$ respectively and $K_p(E_{j,i,a})$ and $K_p(P_{j,i,a})$ are discrete restriction constants for $E_{j,i,a}$ and $P_{j,i,a}$ respectively.

If $i = 0$, we simply write $D_p(E_j) := D_p(E_{j,0})$, $D_p(P_j) := D_p(P_{j,0})$, $K_p(E_j) := K_p(E_{j,0})$ and $K_p(P_j) := K_p(P_{j,0})$.

We will show that $K_{6/\alpha}(P_j) \lesssim_\epsilon N_j^\epsilon$ through the following inequalities:

$$K_{6/\alpha}(P_j) \lesssim_\alpha D_{6/\alpha}(P_j) \lesssim_\epsilon N_j^\epsilon D_{2/\alpha}(E_j)$$

and

$$D_{2/\alpha}(E_j) \approx_\alpha K_{2/\alpha}(E_j) \lesssim_\epsilon \tilde{C}_{2/\alpha}^j$$

where \tilde{C}_p is a constant multiple of C_p in (2.5).

3.1. From $\Lambda(p)$ -sets to decoupling for E_j . We need the following lemmas to use $\Lambda(p)$ -sets in multiscale.

Lemma 3.1. *For $p \geq 2$, let S_1 be a subset of \mathbb{Z}^d . For $a \in S_1$ and $k \in \mathbb{N}$, let $S_{2,a}$ be subsets of $([0, k] \cap \mathbb{Z})^d$. Assume that the sets S_1 and $S_{2,a}$ satisfy*

$$\left\| \sum_{a \in S_1} c_a e(a \cdot x) \right\|_{L^p([0,1]^d)} \leq C_1 \left(\sum_{a \in S_1} |c_a|^2 \right)^{1/2} \quad \forall \{c_a\}_{a \in S_1} \in \ell^2 \quad (3.3)$$

and

$$\left\| \sum_{b \in S_{2,a}} c_b e(b \cdot x) \right\|_{L^p([0,1]^d)} \leq C_2 \left(\sum_{b \in S_{2,a}} |c_b|^2 \right)^{1/2} \quad \forall \{c_b\}_{b \in S_{2,a}} \in \ell^2, \forall a \in S_1. \quad (3.4)$$

Then, we have

$$\left\| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e((ka + b) \cdot x) \right\|_{L^p([0,1]^d)} \lesssim_p C_1 C_2 \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{1/2} \quad (3.5)$$

for all $\{c_{a,b}\}_{a \in S_1, b \in S_{2,a}} \in \ell^2$.

Lemma 3.2. *For $p \geq 2$ and for any $a \in A_i$, $D_p(E_{j,i,a}) \approx_p K_p(E_{j,i,a})$.*

Lemma 3.1 and 3.2 follow from the proofs of Lemma 4 and Proposition 1 in [15] and the duality of L^p played a key role. In Section 8, we provide alternative proofs of them which do not rely on duality.

Remark 3.3. Readers may want to compare Lemma 3.1 with Proposition 3.5 in [4]. They are similar but there is a trade-off. Lemma 3.1 can cover more general cases, because p does not need to be an even integer and there is no assumption on carryover. However, the implicit constant of the inequality is larger than Proposition 3.5 in [4] because the constant in (3.5) is not exactly $C_1 C_2$, but $C'_p C_1 C_2$ where $C'_p > 1$.

Lemma 3.4. *For $0 < \alpha < 1$, $D_{2/\alpha}(E_{j,i}) \lesssim_\alpha \tilde{C}_{2/\alpha}^{j-i}$ where the $\tilde{C}_{2/\alpha}$ is a constant multiple of $C_{2/\alpha}$ in (2.5).*

Proof. For any $a \in A_i$, note that

$$\sum_{b \in A_{j,i,a}} c_b e(bx) = \sum_{b_1 \in A_{j-1,i,a}} \sum_{b_2 \in S_{j,b_1}} c_{b_1,b_2} e(n_j^{1/2} b_1 + b_2).$$

For fixed $a \in A_i$, let C_1 be a constant such that

$$\left\| \sum_{b_1 \in A_{j-1,i,a}} c_{b_1} e(b_1 x) \right\|_{L^{2/\alpha}([0,1])} \leq C_1 \left(\sum_{b_1 \in A_{j-1,i,a}} |c_{b_1}|^2 \right)^{1/2} \quad \forall \{c_{b_1}\}_{b_1 \in A_{j-1,i,a}} \in \ell^2.$$

Combining Theorem 2.1 and Lemma 3.1, we get

$$\left\| \sum_{b \in A_{j,i,a}} c_b e(bx) \right\|_{L^{2/\alpha}([0,1])} \leq C_1 \tilde{C}_{2/\alpha} \left(\sum_{b \in A_{j,i,a}} |c_b|^2 \right)^{1/2} \quad \forall \{c_b\}_{b \in A_{j,i,a}} \in \ell^2.$$

where $\tilde{C}_{2/\alpha}$ is a constant multiple of $C_{2/\alpha}$.

Iterating Theorem 2.1 and Lemma 3.1, we get $K_{2/\alpha}(E_{j,i,a}) \lesssim \tilde{C}_{2/\alpha}^{j-i}$. By Lemma 3.2, we obtain

$$D_{2/\alpha}(E_{j,i,a}) \approx_\alpha K_{2/\alpha}(E_{j,i,a}) \lesssim \tilde{C}_{2/\alpha}^{j-i}.$$

for any $a \in A_i$. Taking the supremum over $a \in A_i$ finishes the proof. \square

3.2. From decoupling for E_j to decoupling for P_j . In this section, we write $N_{j,i} = N_j N_i^{-1}$. Also, note that $N_{j,0} = N_j$. We will prove the following decoupling estimate for P_j .

Proposition 3.5. *For $0 < \alpha < 1$, we have $D_{6/\alpha}(P_j) \lesssim_{\alpha,\epsilon} N_j^\epsilon$.*

It is well known that $K_p(P_j) \lesssim_p D_p(P_j)$ (see for example [8, Theorem 13.1]). Thus, we have $K_{6/\alpha}(P_j) \lesssim_{\alpha,\epsilon} N_j^\epsilon$. We will use it in the proof of Proposition 4.1.

In [4], they proved that the decoupling estimate for a Cantor set on the parabola can be derived from the decoupling estimate for a Cantor set on the line. We adapt their argument to our setting. The proofs of Lemma 3.6 and 3.7 are standard, but we included them for convenience.

Lemma 3.6 (Almost multiplicativity). *For $i, i_0, j \geq 0$ such that $i \geq i_0$, we have*

$$D_p(P_{j+i,i_0}) \leq D_p(P_{j+i,i}) D_p(P_{i,i_0}).$$

Proof. For arbitrary fixed $a \in A_{i_0}$ and $I_i(b) = N_i^{-1/2}(b + [0,1])$ where $b \in \mathbb{Z}$,

$$\sum_{I \in \mathcal{P}_{j+i}(E_{j+i,i_0,a})} f_{\Omega_I} = \sum_{I_i(b) \in \mathcal{P}_i(E_{i,i_0,a})} \sum_{I \in \mathcal{P}_{j+i}(E_{j+i,i,b})} f_{\Omega_I}.$$

For any function f such that $\text{supp } \widehat{f} \subseteq P_{j+i, i_0, a}$, we obtain

$$\begin{aligned}
& \left\| \sum_{I \in \mathcal{P}_{j+i}(E_{j+i, i_0, a})} f_{\Omega_I} \right\|_{L^p(\mathbb{R}^2)} \\
& \leq D_p(P_{i, i_0, a}) \left(\sum_{I_i(b) \in \mathcal{P}_i(E_{i, i_0, a})} \left\| \sum_{I \in \mathcal{P}_{j+i}(E_{j+i, i, b})} f_{\Omega_I} \right\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2} \\
& \leq D_p(P_{i, i_0, a}) \left(\sum_{I_i(b) \in \mathcal{P}_i(E_{i, i_0, a})} D_p(P_{j+i, i, b}) \sum_{I \in \mathcal{P}_{j+i}(E_{j+i, i, b})} \|f_{\Omega_I}\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2} \\
& \leq D_p(P_{j+i, i}) D_p(P_{i, i_0}) \left(\sum_{I_i(b) \in \mathcal{P}_i(E_{i, i_0, a})} \sum_{I \in \mathcal{P}_{j+i}(E_{j+i, i, b})} \|f_{\Omega_I}\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2}.
\end{aligned}$$

□

For $0 \leq k \leq i_1, i_2 \leq j$ and $i_0 \geq 0$, we define the bilinear constant $M_{p, i_0}(j, k, i_1, i_2, a)$ which is the smallest constant such that

$$\begin{aligned}
& \int_{\mathbb{R}^2} \left| \sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} f_{\Omega_{J_1}} \right|^p \left| \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} g_{\Omega_{J_2}} \right|^{2p} dx \\
& \leq M_{p, i_0}(j, k, i_1, i_2, a)^{3p} \times \\
& \quad \left(\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} \|f_{\Omega_{J_1}}\|_{L^{3p}(\mathbb{R}^2)}^2 \right)^{p/2} \left(\sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} \|g_{\Omega_{J_2}}\|_{L^{3p}(\mathbb{R}^2)}^2 \right)^p
\end{aligned}$$

for all $I_1 \in \mathcal{P}_{i_1+i_0}(E_{i_1+i_0, i_0, a})$ and $I_2 \in \mathcal{P}_{i_2+i_0}(E_{i_2+i_0, i_0, a})$ such that $d(I_1, I_2) \geq N_{k+i_0}^{-1/2}$ and for any f and g such that $\text{supp } \widehat{f} \subseteq P_{j+i_0, i_0, a}$ and $\text{supp } \widehat{g} \subseteq P_{j+i_0, i_0, a}$ respectively. Let $M_{p, i_0}(j, k, i_1, i_2) = \sup_{a \in A_{i_0}} M_{p, i_0}(j, k, i_1, i_2, a)$.

For any $a \in A_{i_0}$, note that

$$\begin{aligned}
M_{p, i_0}(j, k, i_1, i_2, a) & \leq D_{3p}(P_{j+i_0, i_1+i_0})^{1/3} D_{3p}(P_{j+i_0, i_2+i_0})^{2/3} \\
& \leq D_{3p}(P_{j+i_0, \min(i_1, i_2)+i_0}).
\end{aligned}$$

Thus, we get

$$M_{p, i_0}(j, k, i_1, i_2) \leq D_{3p}(P_{j+i_0, \min(i_1, i_2)+i_0}). \quad (3.6)$$

We will use it later in the proof.

Lemma 3.7 (Bilinear reduction). *If $i \leq j$ and $i_0 \geq 0$, then*

$$D_{3p}(P_{j+i_0, i_0}) \lesssim D_{3p}(P_{j+i_0, i+i_0}) + N_{i+i_0, i_0}^2 M_{p, i_0}(j, i, i, i).$$

Proof. For any $a \in A_{i_0}$ and for any f with $\text{supp } \hat{f} \subseteq P_{j+i_0, i_0, a}$, we have

$$\begin{aligned}
& \left\| \sum_{J \in \mathcal{P}_{j+i_0}(E_{j+i_0, i_0, a})} f_{\Omega_J} \right\|_{L^{3p}(\mathbb{R}^2)}^2 \\
&= \left\| \sum_{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a})} \left(\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} f_{\Omega_{J_1}} \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} \overline{f_{\Omega_{J_2}}} \right) \right\|_{L^{3p/2}(\mathbb{R}^2)} \\
&\leq \left\| \sum_{\substack{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a}) \\ d(I_1, I_2) \leq N_{i+i_0}^{-1/2}}} (\dots) \right\|_{L^{3p/2}(\mathbb{R}^2)} + \left\| \sum_{\substack{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a}) \\ d(I_1, I_2) \geq N_{i+i_0}^{-1/2}}} (\dots) \right\|_{L^{3p/2}(\mathbb{R}^2)}. \quad (3.7)
\end{aligned}$$

By the Cauchy-Schwarz inequality, the first term of (3.7) is bounded by

$$\begin{aligned}
&\leq \sum_{\substack{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a}) \\ d(I_1, I_2) \leq N_{i+i_0}^{-1/2}}} \left\| \sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} f_{\Omega_{J_1}} \right\|_{L^{3p}(\mathbb{R}^2)} \left\| \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} f_{\Omega_{J_2}} \right\|_{L^{3p}(\mathbb{R}^2)} \\
&\lesssim \sum_{I \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a})} \left\| \sum_{J \in \mathcal{P}_{j+i_0}(I \cap E_{j+i_0})} f_{\Omega_J} \right\|_{L^{3p}(\mathbb{R}^2)}^2 \\
&\leq \sum_{I(b) \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a})} D_{3p}(P_{j+i_0, i+i_0, b})^2 \sum_{J \in \mathcal{P}_{j+i_0}(I(b) \cap E_{j+i_0})} \|f_{\Omega_J}\|_{L^{3p}(\mathbb{R}^2)}^2 \\
&\leq D_{3p}(P_{j+i_0, i+i_0})^2 \sum_{J \in \mathcal{P}_{j+i_0}(E_{j+i_0, i_0, a})} \|f_{\Omega_J}\|_{L^{3p}(\mathbb{R}^2)}^2.
\end{aligned}$$

By the Cauchy-Schwarz inequality, $\int F^{3p/2} G^{3p/2} dx \leq (\int F^p G^{2p})^{1/2} (\int F^{2p} G^p)^{1/2}$ for any two nonnegative functions. Thus, the second term of (3.7) is bounded by

$$\begin{aligned}
&\leq N_{i+i_0, i_0}^2 \max_{\substack{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a}) \\ d(I_1, I_2) \geq N_i^{-1/2}}} \left\| \left(\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} f_{\Omega_{J_1}} \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} f_{\Omega_{J_2}} \right) \right\|_{L^{3p/2}(\mathbb{R}^2)} \\
&\leq N_{i+i_0, i_0}^2 M_{p, i_0}(j, i, i, i)^2 \max_{\substack{I_1, I_2 \in \mathcal{P}_{i+i_0}(E_{i+i_0, i_0, a}) \\ d(I_1, I_2) \geq N_i^{-1/2}}} \left(\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} \|f_{\Omega_{J_1}}\|_{L^{3p}(\mathbb{R}^2)}^2 \right)^{1/2} \\
&\quad \left(\sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} \|f_{\Omega_{J_2}}\|_{L^{3p}(\mathbb{R}^2)}^2 \right)^{1/2} \\
&\leq N_{i+i_0, i_0}^2 M_{p, i_0}(j, i, i, i)^2 \sum_{J \in \mathcal{P}_{j+i_0}(E_{j+i_0, i_0, a})} \|f_{\Omega_J}\|_{L^{3p}(\mathbb{R}^2)}^2.
\end{aligned}$$

□

Lemma 3.8 (Key estimate in [4]). *Let $p = 2/\alpha$ where $0 < \alpha < 1$. If $0 \leq k \leq i$, $2i \leq j$ and $i_0 \geq 0$, then*

$$M_{p,i_0}(j, k, i, i) \lesssim_p N_{k+i_0,i_0}^{1/6} (\tilde{C}_p B)^{i/3} B^{i_0/3} M_{p,i_0}(j, k, 2i, i). \quad (3.8)$$

If $0 \leq k \leq i$, $4i \leq j$ and $i_0 \geq 0$, then

$$M_{p,i_0}(j, k, i, 2i) \lesssim_p N_{k+i_0,i_0}^{1/3} (\tilde{C}_p B)^i B^{2i_0/3} M_{p,i_0}(j, k, 4i, 2i). \quad (3.9)$$

where B is the constant in (2.4) and \tilde{C}_p is the constant in Lemma 3.4.

Proof. We follow the proof of Lemma 2.4 in [4] with modifications. The condition (2.4) comes into play since n_k is nondecreasing. Let us prove (3.8) first. Fix an arbitrary $a \in A_{i_0}$. We only need to consider when f and g satisfy

$$\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} \|f_{\Omega_{J_1}}\|_{L^{3p}(\mathbb{R}^2)}^2 = \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} \|g_{\Omega_{J_2}}\|_{L^{3p}(\mathbb{R}^2)}^2 = 1 \quad (3.10)$$

where I_1 and I_2 are intervals in the definition of $M_{p,i_0}(j, k, i_1, i_2, a)$. Then, it suffices to show that,

$$\begin{aligned} \int_{\mathbb{R}^2} \left| \sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} f_{\Omega_{J_1}} \right|^p \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} g_{\Omega_{J_2}}^{2p} dx \\ \lesssim_p N_{k+i_0,i_0}^{p/2} (\tilde{C}_p B)^{ip} B^{i_0 p} M_{p,i_0}(j, k, 2i, i)^{3p}. \end{aligned} \quad (3.11)$$

Let $I_1 = N_{i+i_0}^{-1/2}(b_1 + [0, 1])$ and $I_2 = N_{i+i_0}^{-1/2}(b_2 + [0, 1])$ for some $b_1, b_2 \in A_{i+i_0}$. We can also assume that I_2 is to the left of I_1 so that $b_1 - b_2 \geq N_{i+i_0}^{1/2} N_{k+i_0}^{-1/2}$. Using the change of variables

$$\begin{aligned} x_1 &= x'_1 + b_2 N_{i+i_0}^{-1/2} \\ x_2 &= x'_2 + 2b_2 N_{i+i_0}^{-1/2} x'_1 + b_2^2 N_{i+i_0}^{-1}, \end{aligned}$$

we can reduce (3.11) to when $I_2 = [0, N_{i+i_0}^{-1/2}]$ and $I_1 = N_{i+i_0}^{-1/2}((b_1 - b_2) + [0, 1])$. For each $J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})$, the center of J is a distance $\geq N_{k+i_0}^{-1/2}$ away from the origin. Also, note that $I_1, I_2 \subseteq [0, N_{i_0}^{-1/2}]$.

Let


$$F_J := \sum_{J_1 \in \mathcal{P}_{j+i_0}(J \cap E_{j+i_0})} f_{\Omega_{J_1}}$$

and

$$G := \sum_{J_2 \in \mathcal{P}_{j+i_0}([0, N_{i+i_0}^{-1/2}] \cap E_{j+i_0})} g_{\Omega_{J_2}}.$$

Since f and g are normalized in (3.10), we have

$$\begin{aligned} \sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \|F_J G^2\|_{L^p(\mathbb{R}^2)}^2 \\ \leq M_{p,i_0}(j, k, 2i, i)^6 \sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \sum_{J_1 \in \mathcal{P}_{j+i_0}(J \cap E_{j+i_0})} \|f_{\Omega_{J_1}}\|_{L^p(\mathbb{R}^2)}^2 \\ \leq M_{p,i_0}(j, k, 2i, i)^6. \end{aligned}$$

FIGURE 2. The supports of \widehat{F}_J and \widehat{G}

Therefore, we have (3.11), if we prove

$$\left\| \sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} F_J G^2 \right\|_{L^p(\mathbb{R}^2)} \lesssim_p N_{k+i_0, i_0}^{1/2} (\tilde{C}_p B)^i B^{i_0} \left(\sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \|F_J G^2\|_{L^p(\mathbb{R}^2)}^2 \right)^{1/2}. \quad (3.12)$$

We claim that for any fixed $x_1 \in \mathbb{R}$,

$$\begin{aligned} & \int_{\mathbb{R}} \left| \sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} F_J(x_1, x_2) G(x_1, x_2)^2 \right|^p dx_2 \\ & \lesssim_p D_p(E_{2i+i_0, i+i_0})^p \left(\frac{N_{k+i_0}^{1/2} N_{2i+i_0}^{1/2}}{N_{i+i_0}} \right)^p \\ & \left[\sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \left(\int_{\mathbb{R}} |F_J(x_1, x_2) G(x_1, x_2)^2|^p dx_2 \right)^{2/p} \right]^{p/2}. \end{aligned} \quad (3.13)$$

Assuming the claim (3.13) and using Lemma 3.4, we obtain $D_p(E_{2i+i_0, i+i_0}) \lesssim_p \tilde{C}_p^i$ and by (2.4),

$$\frac{N_{2i+i_0}^{1/2} N_{i_0}^{1/2}}{N_{i+i_0}} \leq \frac{N_{2i+2i_0}^{1/2}}{N_{i+i_0}} \leq B^{i+i_0}. \quad (3.14)$$

Since $p \geq 2$, we apply Minkowski's inequality to (3.13) and obtain (3.12).

Now we prove the claim. Since $J \subseteq [0, N_{i_0}^{-1/2}]$, \widehat{F}_J is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_J^2 + O(N_{2i+i_0}^{-1/2} N_{i_0}^{-1/2} + N_{j+i_0}^{-1})\}$$

where γ_J is the center of J and since $J_2 \subseteq [0, N_{i+i_0}^{-1/2}]$, \widehat{G} is supported on an $O(N_{i+i_0}^{-1/2}) \times O(N_{i+i_0}^{-1} + N_{j+i_0}^{-1})$ rectangle centered at the origin, see Figure 2. Since $2i \leq j$, we have $N_{j+i_0}^{-1} \leq N_{i+i_0}^{-1}$ and also note that $N_{2i+i_0}^{-1/2} N_{i_0}^{-1/2} \leq N_{i+i_0}^{-1}$. Therefore, $\widehat{F_J G^2}$ is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_J^2 + O(N_{i+i_0}^{-1})\}.$$

The Fourier transform of $F_J G^2$ in y for fixed x is also supported on an interval of length $O(N_{i+i_0}^{-1})$ centered at γ_J^2 .

Let $c \approx N_{k+i_0, i_0}^{1/2} N_{2i+i_0}^{1/2} N_{i+i_0}^{-1}$ and cJ be the interval of length $c|J|$ which has the same center with J . Since I_1 is the interval of length $N_{i+i_0}^{-1/2}$, we have

$$\left\| \sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} f_{cJ} \right\|_{L^p(\mathbb{R})} \lesssim D_p(E_{2i+i_0, i+i_0}) \left(\frac{N_{k+i_0}^{1/2} N_{2i+i_0}^{1/2}}{N_{i+i_0}} \right) \left(\sum_{J \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \|f_{cJ}\|_{L^p(\mathbb{R})}^2 \right)^{1/2}. \quad (3.15)$$

Let γ_m be the left endpoint of I_1 and let us consider

$$T(x) = (2\gamma_m + N_{i+i_0}^{-1/2})(x - \gamma_m) + \gamma_m^2.$$

Note that $T(x)$ is an equation of a line which passes through (γ_m, γ_m^2) and $(\gamma_m + N_{i+i_0}^{-1/2}, (\gamma_m + N_{i+i_0}^{-1/2})^2)$. Since $\gamma_m \geq N_{k+i_0}^{-1/2}$ and $|\gamma_J - \gamma_m| \leq |I_1| = N_{i+i_0}^{-1/2}$,

$$\begin{aligned} T^{-1}(\gamma_J^2 + O(N_{i+i_0}^{-1})) &= \gamma_J + \frac{(\gamma_J - \gamma_m)(\gamma_J - \gamma_m - N_{i+i_0}^{-1/2})}{2\gamma_m + N_{i+i_0}^{-1/2}} + O(N_{k+i_0}^{1/2} N_{i+i_0}^{-1}) \\ &= \gamma_J + O(N_{k+i_0}^{1/2} N_{i+i_0}^{-1}). \end{aligned}$$

We use $|J| = N_{2i+i_0}^{-1/2}$ and obtain that $\gamma_J^2 + O(N_{i+i_0}^{-1}) \subseteq T(cJ)$ if c is sufficiently large while still comparable to $N_{k+i_0, i_0}^{1/2} N_{2i+i_0}^{1/2} N_{i+i_0}^{-1}$. The intervals cJ in (3.15) can be replaced by $T(cJ)$ and we obtain (3.13).

Now, we turn to the proof of (3.9). Similarly, we fix an arbitrary $a \in A_{i_0}$. We only need to consider f and g such that

$$\sum_{J_1 \in \mathcal{P}_{j+i_0}(I_1 \cap E_{j+i_0})} \|f_{\Omega_{J_1}}\|_{L^{3p}(\mathbb{R}^2)}^2 = \sum_{J_2 \in \mathcal{P}_{j+i_0}(I_2 \cap E_{j+i_0})} \|g_{\Omega_{J_2}}\|_{L^{3p}(\mathbb{R}^2)}^2 = 1.$$

We can assume that $I_2 := [0, N_{2i+i_0}^{-1/2}]$ is on the left of an interval I_1 of length $N_{i+i_0}^{-1/2}$ and $I_1, I_2 \subseteq [0, N_{i_0}^{-1/2}]$. For each $J \in \mathcal{P}_{4i+i_0}(I_1 \cap E_{4i+i_0})$, the center of J is a distance $\geq N_{k+i_0}^{-1/2}$ away from the origin. Let

$$F_J := \sum_{J_1 \in \mathcal{P}_{j+i_0}(J \cap E_{j+i_0})} f_{\Omega_{J_1}}$$

and

$$G := \sum_{J_2 \in \mathcal{P}_{j+i_0}([0, N_{2i+i_0}^{-1/2}] \cap E_{j+i_0})} g_{\Omega_{J_2}}.$$

By the same argument in the proof of (3.8), it suffices to prove that, for any fixed $x_1 \in \mathbb{R}$,

$$\begin{aligned} & \int_{\mathbb{R}} \left| \sum_{J \in \mathcal{P}_{4i+i_0}(I_1 \cap E_{4i+i_0})} F_J(x_1, x_2) G(x_1, x_2)^2 \right|^p dx_2 \\ & \lesssim_p D_p(E_{4i+i_0, 2i+i_0})^p D_p(E_{2i+i_0, i+i_0})^p \left(\frac{N_{k+i_0}^{1/2} N_{4i+i_0}^{1/2}}{N_{2i+i_0}} \right)^p \left(\frac{N_{k+i_0}^{1/2} N_{2i+i_0}^{1/2}}{N_{i+i_0}} \right)^p \\ & \quad \left[\sum_{J \in \mathcal{P}_{4i+i_0}(I_1 \cap E_{4i+i_0})} \left(\int_{\mathbb{R}} |F_J(x_1, x_2) G(x_1, x_2)^2|^p dx_2 \right)^{2/p} \right]^{p/2}, \end{aligned} \quad (3.16)$$

which corresponds to (3.13). By Lemma 3.4 and using (3.14) twice, we obtain (3.9).

We now prove (3.16). For $J_0 \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})$, let

$$F_{J_0} = \sum_{J \in \mathcal{P}_{4i+i_0}(J_0 \cap E_{4i+i_0})} F_J,$$

so that

$$\sum_{J \in \mathcal{P}_{4i+i_0}(I_1 \cap E_{4i+i_0})} F_J = \sum_{J_0 \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} F_{J_0}.$$

Since $J_0 \subseteq [0, N_{i_0}^{-1/2}]$, \widehat{F}_{J_0} is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_{J_0}^2 + O(N_{2i+i_0}^{-1/2} N_{i_0}^{-1/2} + N_{j+i_0}^{-1})\}$$

where γ_{J_0} is the center of J_0 and since $J_2 \subseteq [0, N_{2i+i_0}^{-1/2}]$, \widehat{G} is supported on an $O(N_{2i+i_0}^{-1/2}) \times O(N_{2i+i_0}^{-1} + N_{j+i_0}^{-1})$ rectangle. Then, $\widehat{F_{J_0} G^2}$ is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_{J_0}^2 + O(N_{i+i_0}^{-1})\}.$$

By the same argument in the proof of (3.13), we obtain that

$$\begin{aligned} & \int_{\mathbb{R}} \left| \sum_{J \in \mathcal{P}_{4i+i_0}(I_1 \cap E_{j+i_0})} F_J(x_1, x_2) G(x_1, x_2)^2 \right|^p dx_2 \\ & = \int_{\mathbb{R}} \left| \sum_{J_0 \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} F_{J_0}(x_1, x_2) G(x_1, x_2)^2 \right|^p dx_2 \\ & \lesssim_p D_p(E_{2i+i_0, i+i_0})^p \left(\frac{N_{k+i_0}^{1/2} N_{2i+i_0}^{1/2}}{N_{i+i_0}} \right)^p \\ & \quad \left[\sum_{J_0 \in \mathcal{P}_{2i+i_0}(I_1 \cap E_{2i+i_0})} \left(\int_{\mathbb{R}} |F_{J_0}(x_1, x_2) G(x_1, x_2)^2|^p dx_2 \right)^{2/p} \right]^{p/2}. \end{aligned} \quad (3.17)$$

Similarly, \widehat{F}_J is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_J^2 + O(N_{4i+i_0}^{-1/2} N_{i_0}^{-1/2} + N_{j+i_0}^{-1})\}$$

where γ_J is the center of J . Since $4i \leq j$, Fourier transform of $F_J G^2$ in y for fixed x is supported in the horizontal strip

$$\{(\xi_1, \xi_2) : \xi_2 = \gamma_J^2 + O(N_{2i+i_0}^{-1})\}.$$

We use the argument in the proof of (3.13) with i replaced by $2i$ and obtain

$$\begin{aligned} & \int_{\mathbb{R}} \left| \sum_{J \in \mathcal{P}_{4i+i_0}(J_0 \cap E_{4i+i_0})} F_J(x_1, x_2) G(x_1, x_2)^2 \right|^p dx_2 \\ & \lesssim_p D_p(E_{4i+i_0, 2i+i_0})^p \left(\frac{N_{k+i_0}^{1/2} N_{4i+i_0}^{1/2}}{N_{2i+i_0}} \right)^p \\ & \quad \left[\sum_{J \in \mathcal{P}_{4i+i_0}(J_0 \cap E_{4i+i_0})} \left(\int_{\mathbb{R}} |F_J(x_1, x_2) G(x_1, x_2)^2|^p dx_2 \right)^{2/p} \right]^{p/2}. \end{aligned} \quad (3.18)$$

Combining (3.17) and (3.18), we get (3.16). \square

Lemma 3.9. *Let $k \leq i_2 \leq i_1 \leq j$. Then,*

$$M_{p, i_0}(j, k, i_1, i_2) \leq M_{p, i_0}(j, k, i_2, i_1)^{1/2} D_{3p}(P_{j+i_0, i_2+i_0})^{1/2}.$$

Proof. As in [4], it follows from $\int F^p G^{2p} dx \leq (\int F^{2p} G^p)^{1/2} (\int G^{3p})^{1/2}$ where F and G are nonnegative functions and the definition of $M_{p, i_0}(j, k, i_1, i_2)$. \square

We now can prove Proposition 3.5.

Proof of Proposition 3.5. Let $p = 2/\alpha$ and assume that λ is the smallest exponent such that

$$D_{3p}(P_{j+i_0, i_0}) \lesssim_{p, \epsilon} N_{j+i_0}^\epsilon N_{j+i_0, i_0}^\lambda \quad (3.19)$$

for all $0 \leq i_0, j$ and for any $0 < \epsilon < 1$. Then, it suffices to show that $\lambda = 0$. We can run the iteration as in [4]. But, we should also consider that n_j is nondecreasing. There is a trivial estimate $D_{3p}(P_{j+i_0, i_0}) \lesssim N_{j+i_0, i_0}^{1/4}$, so we assume that $0 < \lambda \leq 1/4$ toward a contradiction.

By (2.2), for any $i_0 \geq 0$ we have

$$(\tilde{C}_p B)^{i/3} \lesssim_p N_i \leq N_{i+i_0, i_0}. \quad (3.20)$$

Combining Lemma 3.7, Lemma 3.8 and (3.20), we obtain that if $j \geq 2i$, then

$$\begin{aligned} D_{3p}(P_{j+i_0, i_0}) & \leq D_{3p}(P_{j+i_0, i+i_0}) + N_{i+i_0, i_0}^{O(1)} (\tilde{C}_p B)^{i/3} B^{i_0/3} M_{p, i_0}(j, i, 2i, i) \\ & \leq D_{3p}(P_{j+i_0, i+i_0}) + N_{i+i_0, i_0}^{O(1)} B^{i_0/3} M_{p, i_0}(j, i, 2i, i). \end{aligned} \quad (3.21)$$

Now we need an estimate for $M_{p, i_0}(j, i, 2i, i)$. By Lemma 3.8 and Lemma 3.9, for any positive integer a such that $1 \leq a \leq \frac{j}{4i}$, we obtain that

$$\begin{aligned} M_{p, i_0}(j, i, 2ai, ai) & \leq M_{p, i_0}(j, i, ai, 2ai)^{1/2} D_{3p}(P_{j+i_0, ai+i_0})^{1/2} \\ & \lesssim_p N_{i+i_0, i_0}^{O(1)} (\tilde{C}_p B)^{ai/2} B^{i_0/3} M_{p, i_0}(j, i, 4ai, 2ai)^{1/2} D_{3p}(P_{j+i_0, ai+i_0})^{1/2}. \end{aligned} \quad (3.22)$$

First, we assume that $j = 2^{k+1}i$ where k is a sufficiently integer to be determined later. By iterating (3.22) and using (3.19) with ϵ replaced by $\epsilon/2$, we have

$$\begin{aligned} M_{p, i_0}(j, i, 2i, i) & \lesssim_{p, \epsilon} N_{i+i_0, i_0}^{O(1)} (\tilde{C}_p B)^{ki/2} B^{2i_0/3} \\ & \quad N_{j+i_0}^{(\lambda+\epsilon/2)(1-1/2^k)} (N_{i+i_0} N_{2i+i_0}^{1/2} \cdots N_{2^{k-1}i+i_0}^{1/2^{k-1}})^{-\lambda/2} M_{p, i_0}(j, i, 2^{k+1}i, 2^k i)^{1/2^k}. \end{aligned}$$

We use (3.6) and (3.19) with ϵ replaced by $\epsilon/2$ and we get

$$M_{p, i_0}(j, i, 2^{k+1}i, 2^k i) \leq D_{3p}(P_{j+i_0, 2^k i+i_0}) \lesssim_{p, \epsilon} N_{j+i_0}^{\epsilon/2} N_{j+i_0, 2^k i+i_0}^\lambda.$$

Therefore, we arrive at

$$M_{p,i_0}(j, i, 2i, i) \lesssim_{p,\epsilon} N_{j+i_0}^{\epsilon/2} N_{j+i_0,i_0}^\lambda N_{i+i_0,i_0}^{O(1)} (\tilde{C}_p B)^{ki/2} B^{2i_0/3} \left(\frac{N_{i_0}}{N_{i+i_0}^{1/2} N_{2i+i_0}^{1/4} \cdots N_{2^{k-1}i+i_0}^{1/2^k} N_{2^k i+i_0}^{1/2^k}} \right)^\lambda.$$

Since n_j is nondecreasing, we obtain that for any $s \geq 0$,

$$\left(\frac{N_{i_0}}{N_{2^s i+i_0}} \right)^{1/2^s} \leq \frac{N_{i_0}}{N_{i+i_0}}.$$

Therefore, we have

$$\left(\frac{N_{i_0}}{N_{i+i_0}^{1/2} N_{2i+i_0}^{1/4} \cdots N_{2^{k-1}i+i_0}^{1/2^k} N_{2^k i+i_0}^{1/2^k}} \right) \leq N_{i+i_0,i_0}^{-(k+1)/2},$$

and it leads to

$$M_p(j, i, 2i, i) \lesssim_{p,\epsilon} N_{j+i_0}^{\epsilon/2} N_{j+i_0,i_0}^\lambda N_{i+i_0,i_0}^{O(1)-\lambda(k+1)/2} (\tilde{C}_p B)^{ki/2} B^{2i_0/3}.$$

Since $j = 2^{k+1}i$, $ki \leq j$ for any $k \geq 0$. By using (2.2), we have

$$(\tilde{C}_p B)^{ki/2} \leq (\tilde{C}_p B)^{j/2} \lesssim_{p,\epsilon} N_j^{\epsilon/4} \leq N_{j+i_0}^{\epsilon/4}. \quad (3.23)$$

Since $\lambda \leq 1/4$, $N_{j+i_0,i_0}^\lambda \leq N_{j+i_0,i+i_0}^\lambda N_{i+i_0,i_0}^{O(1)}$ and by (3.23), we get

$$M_p(j, i, 2i, i) \lesssim_{p,\epsilon} N_{j+i_0}^{3\epsilon/4} N_{j+i_0,i+i_0}^\lambda N_{i+i_0,i_0}^{O(1)-\lambda(k+1)/2} B^{2i_0/3}. \quad (3.24)$$

We choose k such that $O(1) - \lambda(k+1)/2 < 0$ and $k \approx \lambda^{-1}$, then it follows from (3.19) with ϵ replaced by $3\epsilon/4$, (3.21) and (3.24) that

$$\begin{aligned} D_{3p}(P_{j+i_0,i_0}) &\leq D_{3p}(P_{j+i_0,i+i_0}) + N_{i+i_0,i_0}^{O(1)} B^{i_0/3} M_{p,i_0}(j, i, 2i, i) \\ &\lesssim_{p,\epsilon} N_{j+i_0}^{3\epsilon/4} N_{j+i_0,i+i_0}^\lambda + N_{j+i_0}^{3\epsilon/4} N_{j+i_0,i+i_0}^\lambda N_{i+i_0,i_0}^{O(1)-\lambda(k+1)/2} B^{i_0} \\ &\lesssim_{p,\epsilon} N_{j+i_0}^{3\epsilon/4} N_{j+i_0,i+i_0}^\lambda B^{i_0}. \end{aligned}$$

Since $j = 2^{k+1}i$, we use (3.14) at all scales $i, 2i, \dots, 2^k i$ and obtain that

$$N_{i+i_0}^{-1} \leq B^{i+i_0} N_{2i+i_0}^{-1/2} N_{i_0}^{-1/2} \leq \dots \leq B^{i(k+1)+2i_0} N_{j+i_0}^{-1/2^{k+1}} N_{i_0}^{-1+1/2^{k+1}}.$$

Therefore, we have

$$D_{3p}(P_{j+i_0,i_0}) \lesssim_{p,\epsilon} N_{j+i_0}^{3\epsilon/4} (N_{j+i_0,i_0})^{\lambda(1-1/2^{k+1})} B^{i(k+1)\lambda} B^{i_0(2\lambda+1)}.$$

Since $\lambda \leq 1/4$ and $ki \leq j$, by using (2.2),

$$B^{i(k+1)\lambda} B^{i_0(2\lambda+1)} \leq B^{2(j+i_0)} \lesssim_\epsilon N_{j+i_0}^{\epsilon/4}.$$

Now, we obtain that

$$D_{3p}(P_{j+i_0,i_0}) \lesssim_{p,\epsilon} N_{j+i_0}^\epsilon N_{j+i_0,i_0}^{\lambda(1-1/2^{k+1})}. \quad (3.25)$$

If $2^{k+1}i \leq j \leq 2^{k+1}(i+1)$, it follows from Lemma 3.6 that

$$D_{3p}(P_{j+i_0,i_0}) \leq D_{3p}(P_{2^{k+1}i+i_0,i_0}) D_{3p}(P_{j+i_0,2^{k+1}i+i_0}).$$

By (3.25) with ϵ replaced by $\epsilon/2$ and the trivial estimate,

$$\begin{aligned} D_{3p}(P_{j+i_0, i_0}) &\lesssim_{p, \epsilon} N_{2^{k+1}i+i_0}^{\epsilon/2} N_{2^{k+1}i+i_0, i_0}^{\lambda(1-1/2^{k+1})} N_{j+i_0, 2^{k+1}i+i_0}^{1/4} \\ &\leq N_{j+i_0}^{\epsilon/2} N_{j+i_0, i_0}^{\lambda(1-1/2^{k+1})} N_{j+i_0, 2^{k+1}i+i_0}^{1/4}. \end{aligned}$$

By (2.3), there exists a constant $C_{\lambda, \epsilon}$ such that

$$n_{j+i_0} \leq n_{j+i_0+1} \leq C_{\lambda, \epsilon} N_{j+i_0}^{\epsilon/2^k}.$$

Since $k \approx \lambda^{-1}$, the constant $C_{\lambda, \epsilon}$ depends on λ . Then, we have

$$N_{j+i_0, 2^{k+1}i+i_0}^{1/4} \leq (n_{j+i_0})^{2^{k-1}} \leq C_{\lambda, \epsilon}^{2^{k-1}} N_{j+i_0}^{\epsilon/2} \lesssim_{\lambda, \epsilon} N_{j+i_0}^{\epsilon/2}.$$

Therefore, we obtain that for any $j, i_0 \geq 0$,

$$D_{3p}(P_{j+i_0, i_0}) \lesssim_{p, \epsilon, \lambda} N_{j+i_0}^{\epsilon} N_{j+i_0, i_0}^{\lambda(1-1/2^{k+1})}.$$

This contradicts the assumption that $\lambda > 0$ is the smallest exponent which satisfies (3.19). Therefore, $\lambda = 0$. \square

4. LOCAL RESTRICTION ESTIMATE

We denote $B^d(R)$ by a cube of side length R in \mathbb{R}^d centered at the origin.

Proposition 4.1. *Let $p = 6/\alpha$ where $0 < \alpha < 1$ and ν be the measure constructed in Section 2. Then, we have*

$$\left\| \widehat{f d\nu} \right\|_{L^p(B^2(R))} \lesssim_{p, \epsilon} R^\epsilon \|f\|_{L^2(\nu)} \quad \forall f \in L^2(\nu). \quad (4.1)$$

Equivalently, for q such that $1/q + 1/p = 1$, we have

$$\left\| \widehat{f} \right\|_{L^2(\nu)} \lesssim_{q, \epsilon} R^\epsilon \|f\|_{L^q(\mathbb{R}^2)} \quad (4.2)$$

for any $f \in L^q(\mathbb{R}^2)$ supported on $B^2(R)$.

Proposition 4.1 follows from the following lemma.

Lemma 4.2. *For $\ell \geq 2j$ and for every $a \in A_{2j}$, we define*

$$\widehat{f_{\Omega_a}}(\xi) = \widehat{f}(\xi) \sum_{b \in A_{\ell, 2j, a}} \mathbf{1}_{[0,1] \times [-1,1]}(N_\ell^{1/2} \xi_1 - b, N_\ell \xi_2 - (2b+1)N_\ell^{1/2} \xi_1 + b^2 + b).$$

For any $\sum_{a \in A_{2j}} f_{\Omega_a} \in L^2(\mathbb{R}^2)$, we have the estimate

$$\left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^p(B^2(N_j))} \lesssim_{\epsilon, p} K_p(P_{2j}) N_{2j}^{\epsilon + \frac{3}{2p} - \frac{\alpha}{4}} N_\ell^{-3/4} |A_\ell|^{1/2} \left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^2(\mathbb{R}^2)}. \quad (4.3)$$

Proof of Proposition 4.1. Let η be a Schwartz function on \mathbb{R} such that $|\eta| \geq 1$ on $[-1, 1]$ and $\widehat{\eta}$ is supported on $[0, 1]$. Let

$$\widehat{F}(\xi) := f(\xi_1, \xi_1^2) \sum_{a \in A_{2j}} \sum_{b \in A_{\ell, 2j, a}} \mathbf{1}_{[0,1]}(N_\ell^{1/2} \xi_1 - b) \widehat{\eta}_\ell(\xi_2 - \xi_1^2) |E_\ell|^{-1}$$

where $\eta_\ell(x) = \eta(x/N_\ell)$ for $x \in \mathbb{R}$. Note that

$$\sum_{a \in A_{2j}} \sum_{b \in A_{\ell, 2j, a}} \mathbf{1}_{[0,1]}(N_\ell^{1/2} \xi_1 - b) = \mathbf{1}_{E_\ell}(\xi_1).$$

If $x \in B^2(N_j)$, we have

$$|\widehat{fd\nu_\ell}(x)| \leq |\widehat{fd\nu_\ell}(x)\eta_\ell(-x_2)| = |F(-x)|.$$

Therefore, (4.3) implies that

$$\begin{aligned} \left\| \widehat{fd\nu_\ell} \right\|_{L^p(B^2(N_j))} &\lesssim \|F\|_{L^p(B^2(N_j))} \\ &\lesssim_\epsilon K_p(P_{2j}) N_{2j}^{\epsilon + \frac{3}{2p} - \frac{\alpha}{4}} N_\ell^{-3/4} |A_\ell|^{1/2} \|F\|_{L^2(\mathbb{R}^2)} \\ &= K_p(P_{2j}) N_{2j}^{\epsilon + \frac{3}{2p} - \frac{\alpha}{4}} N_\ell^{-3/4} |A_\ell|^{1/2} \left\| \widehat{F} \right\|_{L^2(\mathbb{R}^2)}. \end{aligned}$$

Since $|E_\ell| = |A_\ell| N_\ell^{-1/2}$, we get $\left\| \widehat{F} \right\|_{L^2(\mathbb{R}^2)} \lesssim N_\ell^{3/4} |A_\ell|^{-1/2} \|f\|_{L^2(\nu_\ell)}$.

Since $p = 6/\alpha$, it follows from Proposition 3.5 that

$$\begin{aligned} \left\| \widehat{fd\nu_\ell} \right\|_{L^p(B^2(N_j))} &\lesssim_\epsilon K_p(P_{2j}) N_{2j}^\epsilon \|f\|_{L^2(\nu_\ell)} \\ &\lesssim_\epsilon N_{2j}^{2\epsilon} \|f\|_{L^2(\nu_\ell)}. \end{aligned}$$

If $N_{j-1} \leq R \leq N_j$, conditions (2.2), (2.3) and (2.4) imply that

$$N_{2j} \leq B^{2j} N_j^2 = B^{2j} n_j^2 N_{j-1}^2 \lesssim_\epsilon N_{j-1}^{2+\epsilon}.$$

Therefore,

$$\begin{aligned} \left\| \widehat{fd\nu_\ell} \right\|_{L^p(B^2(R))} &\leq \left\| \widehat{fd\nu_\ell} \right\|_{L^p(B^2(N_j))} \\ &\lesssim N_{2j}^{2\epsilon} \|f\|_{L^2(\nu_\ell)} \\ &\lesssim_\epsilon N_{j-1}^{O(\epsilon)} \|f\|_{L^2(\nu_\ell)} \\ &\lesssim_\epsilon R^{O(\epsilon)} \|f\|_{L^2(\nu_\ell)}. \end{aligned}$$

When we take the limit $\ell \rightarrow \infty$, we get (4.1) and by duality, (4.2) follows. \square

Now let us turn to the proof of Lemma 4.2.

Proof of Lemma 4.2. Let η be a Schwartz function on \mathbb{R}^2 such that $|\eta| \geq 1$ on $[-1, 1]^2$ and $\widehat{\eta}$ is supported on $[0, 1]^2$ and write $\eta_{2j}(x) = \eta(x/N_{2j}^{1/2})$. Since $N_j \leq N_{2j}^{1/2}$, we have

$$\begin{aligned} \left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^p(B^2(N_j))} &\leq \left\| \sum_{a \in A_{2j}} f_{\Omega_a} \eta_{2j} \right\|_{L^p(\mathbb{R}^2)} \\ &= \sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \left| \int \sum_{a \in A_{2j}} f_{\Omega_a}(x) \eta_{2j}(x) g(x) dx \right| \\ &= \sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \left| \int \sum_{a \in A_{2j}} \widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi) \widehat{g}(\xi) d\xi \right| \end{aligned}$$

where $1/p + 1/q = 1$.

The function \widehat{f}_{Ω_a} is supported on a rectangle of dimensions $O(N_{2j}^{-1/2}) \times O(N_{2j}^{-1})$ centered at $((a + 1/2)N_{2j}^{-1/2}, (a + 1/2)^2 N_{2j}^{-1})$ and the direction that it is pointing depends on a . The function $\widehat{\eta}_{2j}$ is supported on a square with side length $O(N_{2j}^{-1/2})$ centered at the origin. Therefore, $\widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi)$ is supported on a square with side length $O(N_{2j}^{-1/2})$ centered at $(aN_{2j}^{-1/2}, a^2 N_{2j}^{-1})$.

For $a \in A_{2j}$ and $c = (c_1, c_2) \in \mathbb{Z}^2$, let us consider characteristic functions

$$\mathbf{1}_{Q_{a,c}}(\xi) = \mathbf{1}_{[0,1]^2}(N_{2j}^{1/2}\xi_1 - (a + c_1), N_{2j}^{1/2}\xi_2 - (a^2 N_{2j}^{-1/2} + c_2)).$$

Then, we have

$$\left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^p(B^2(N_j))} \leq \sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \sum_{|c| \lesssim 1} \left| \int \sum_{a \in A_{2j}} \widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi) \widehat{g}(\xi) \mathbf{1}_{Q_{a,c}}(\xi) d\xi \right|.$$

By letting

$$z_1 = N_{2j}^{1/2}\xi_1 - (a + c_1) \quad \text{and} \quad z_2 = N_{2j}^{1/2}\xi_2 - (a^2 N_{2j}^{-1/2} + c_2), \quad (4.4)$$

we obtain

$$\sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \sum_{|c| \lesssim 1} \left| \int_{[0,1]^2} \sum_{a \in A_{2j}} \widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(N_{2j}^{-1/2}(z_1 + a + c_1), N_{2j}^{-1/2}(z_2 + c_2) + a^2 N_{2j}^{-1}) \widehat{g}(N_{2j}^{-1/2}(z_1 + a + c_1), N_{2j}^{-1/2}(z_2 + c_2) + a^2 N_{2j}^{-1}) dz N_{2j}^{-1} \right|.$$

Let $a' = (a'_1, a'_2) \in \mathbb{Z}^2$ and $a' \cdot u = a'_1 u_1 + a'_2 u_2$. Then, we have

$$\sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} \int_{[0,1]^2} e(au_1 + a^2 u_2) e(-a' \cdot u) du = 1 \quad (4.5)$$

if and only if $a'_1 = a$ and $a'_2 = a^2$ for $|a| \lesssim N_{2j}^{1/2}$.

Let

$$F_c(u, z) := \sum_{a \in A_{2j}} \widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(N_{2j}^{-1/2}(z_1 + a + c_1), N_{2j}^{-1/2}(z_2 + c_2) + a^2 N_{2j}^{-1}) e(au_1 + a^2 u_2)$$

and

$$G_c(u, z) := \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} \widehat{g}(N_{2j}^{-1/2}(z_1 + a'_1 + c_1), N_{2j}^{-1/2}(z_2 + c_2) + a'_2 N_{2j}^{-1}) e(-a' \cdot u).$$

Also, we denote the mixed norm of f by

$$\|f\|_{L_u^{p_1} L_z^{p_2}([0,1]^2)} = \left[\int_{[0,1]^2} \left(\int_{[0,1]^2} |f(u, z)|^{p_1} du \right)^{p_2/p_1} dz \right]^{1/p_2}.$$

By (4.5) and Hölder's inequality, we have

$$\begin{aligned} \left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^p(B^2(N_j))} &\leq \sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \sum_{|c| \lesssim 1} \left| \int_{[0,1]^2} \int_{[0,1]^2} F_c(u, z) G_c(u, z) du dz \right| N_{2j}^{-1} \\ &\leq \sup_{\|g\|_{L^q(\mathbb{R}^2)} \leq 1} \sum_{|c| \lesssim 1} \|F_c(u, z)\|_{L_u^p L_z^2([0,1]^2)} \|G_c(u, z)\|_{L_u^q L_z^2([0,1]^2)} N_{2j}^{-1}. \end{aligned} \quad (4.6)$$

Since $F_c(u, z)$ is a Fourier series with respect to u variable, the definition of $K_p(P_{2j})$ and (4.4) implies that

$$\begin{aligned} \|F_c\|_{L_u^p L_z^2([0,1]^2)} &\leq K_p(P_{2j}) \|F_c\|_{L_u^2 L_z^2([0,1]^2)} \\ &\leq K_p(P_{2j}) N_{2j}^{1/2} \left(\int \sum_{a \in A_{2j}} |\widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi)|^2 d\xi \right)^{1/2}. \end{aligned} \quad (4.7)$$

In order to obtain an estimate for $\|G_c\|_{L_u^q L_z^2([0,1]^2)}$, we consider G_c as a linear operator $T : L^q(\mathbb{R}^2) \rightarrow L_u^q L_z^2([0,1]^2)$ acting on g , which is defined by

$$\begin{aligned} Tg(u, z) &= \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} \int g(x) e(N_{2j}^{-1/2} z_1 x_1 + N_{2j}^{-1/2} z_2 x_2) \\ &\quad e(N_{2j}^{-1/2} (a'_1 + c_1) x_1 + (a'_2 N_{2j}^{-1} + c_2 N_{2j}^{-1/2}) x_2) e(-a' \cdot u) dx. \end{aligned}$$

If we show that

$$\|Tg\|_{L_u^1 L_z^2([0,1]^2)} \lesssim_\epsilon N_{2j}^\epsilon \|g\|_{L^1(\mathbb{R}^2)} \quad \forall g \in L^1(\mathbb{R}^2) \quad (4.8)$$

and

$$\|Tg\|_{L_u^2 L_z^2([0,1]^2)} \lesssim N_{2j}^{3/4} \|g\|_{L^2(\mathbb{R}^2)} \quad \forall g \in L^2(\mathbb{R}^2), \quad (4.9)$$

then the mixed norm interpolation theorem (see [2, Theorem 2 in Section 7]) implies that

$$\|G_c(u, z)\|_{L_u^q L_z^2([0,1]^2)} = \|Tg\|_{L_u^q L_z^2([0,1]^2)} \lesssim_\epsilon N_{2j}^{\epsilon + \frac{3}{2}(1-\frac{1}{q})} \|g\|_{L^q(\mathbb{R}^2)} \quad \forall g \in L^q(\mathbb{R}^2). \quad (4.10)$$

When $q = 1$, we only need to consider the case $c = 0$, since the similar argument works for $c \neq 0$. We have

$$Tg(u, z) = \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} \int_{\mathbb{R}^2} g(N_{2j}^{1/2} x_1, N_{2j} x_2) e(x_1 z_1 + N_{2j}^{1/2} x_2 z_2) N_{2j}^{3/2} e(-a' \cdot (u - x)) dx.$$

Note that $\sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} e(-a' \cdot u)$ is a product of Dirichlet kernels, since

$$\sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} e(-a' \cdot u) = \sum_{|a'_1| \lesssim N_{2j}^{1/2}} e(-a'_1 u_1) \sum_{|a'_2| \lesssim N_{2j}} e(-a'_2 u_2).$$

Uniformly in x , we have

$$\left\| \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} e(-a' \cdot (u - x)) \right\|_{L^1([0,1]^2, du)} \lesssim \log(N_{2j}^{1/2}) \log(N_{2j}).$$

Therefore, we obtain that

$$\begin{aligned}
\|Tg\|_{L^1([0,1]^2, du)} &\lesssim \int_{\mathbb{R}^2} \int_{[0,1]^2} \left| g(N_{2j}^{1/2}x_1, N_{2j}x_2) N_{2j}^{3/2} \right| \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} e(-a' \cdot (x-u)) \Big| dx du \\
&\lesssim \log(N_{2j})^2 \int_{\mathbb{R}^2} \left| g(N_{2j}^{1/2}x_1, N_{2j}x_2) N_{2j}^{3/2} \right| dx \\
&\lesssim_\epsilon N_{2j}^\epsilon \|g\|_{L^1(\mathbb{R}^2)}.
\end{aligned} \tag{4.11}$$

Since the estimate (4.11) holds uniformly in z , we established (4.8).

When $q = 2$, since $Tg(u, z)$ is a Fourier series with respect to u variable, we use Plancherel's theorem and by (4.4), we have

$$\begin{aligned}
&\|Tg\|_{L_u^2 L_z^2([0,1]^2)} \\
&= \left(\int_{[0,1]^2} \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} |\widehat{g}(N_{2j}^{-1/2}(z_1 + a'_1 + c_1), N_{2j}^{-1/2}(z_2 + c_2) + a'_2 N_{2j}^{-1})|^2 dz \right)^{1/2} \\
&= N_{2j}^{1/2} \left(\int |\widehat{g}(\xi)|^2 \sum_{|a'_1| \lesssim N_{2j}^{1/2}, |a'_2| \lesssim N_{2j}} \mathbf{1}_{Q_{a',c}}(\xi) d\xi \right)^{1/2}
\end{aligned}$$

where

$$\mathbf{1}_{Q_{a',c}}(\xi) = \mathbf{1}_{[0,1]^2}(N_{2j}^{1/2}\xi_1 - (a'_1 + c_1), N_{2j}^{1/2}\xi_2 - (a'_2 N_{2j}^{-1/2} + c_2)).$$

The function $\mathbf{1}_{Q_{a',c}}(\xi)$ is supported on a square with side length $O(N_{2j}^{-1/2})$ centered at $((a'_1 + c_1)N_{2j}^{-1/2}, a'_2 N_{2j}^{-1} + c_2 N_{2j}^{-1/2})$. These squares overlap at most $O(N_{2j}^{1/2})$ times. Therefore, we have

$$\|Tg\|_{L_u^2 L_z^2([0,1]^2)} \lesssim N_{2j}^{3/4} \|\widehat{g}\|_{L^2(\mathbb{R}^2)} \leq N_{2j}^{3/4} \|g\|_{L^2(\mathbb{R}^2)}.$$

Therefore, we established (4.9). Combining (4.6), (4.7) and (4.10), we obtain that

$$\left\| \sum_{a \in A_{2j}} f_{\Omega_a} \right\|_{L^p(B^2(N_j))} \lesssim_\epsilon K_p(P_{2j}) N_{2j}^{\epsilon + \frac{3}{2p} - \frac{1}{2}} \left(\int \sum_{a \in A_{2j}} |\widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi)|^2 d\xi \right)^{1/2}. \tag{4.12}$$

We define

$$\mathbf{1}_{\ell, 2j, a}(\xi) := \sum_{b \in A_{\ell, 2j, a}} \mathbf{1}_{[0,1] \times [-1,1]}(N_\ell^{1/2}\xi_1 - b, N_\ell \xi_2 - (2b+1)N_\ell^{1/2}\xi_1 + b^2 + b)$$

so that $\widehat{f}_{\Omega_a}(\xi) = \widehat{f}_{\Omega_a} \mathbf{1}_{\ell, 2j, a}(\xi)$. By Hölder's inequality, we have

$$\begin{aligned}
|\widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi)|^2 &\leq (|\widehat{f}_{\Omega_a} \mathbf{1}_{\ell, 2j, a}| * |\widehat{\eta}_{2j}|(\xi))^2 \\
&\leq (|\widehat{f}_{\Omega_a}|^2 * |\widehat{\eta}_{2j}|(\xi))(\mathbf{1}_{\ell, 2j, a} * |\widehat{\eta}_{2j}|(\xi)).
\end{aligned}$$

Since $|\widehat{\eta}_{2j}| \lesssim N_{2j}$, for any fixed ξ , we have

$$(\mathbf{1}_{\ell, 2j, a} * |\widehat{\eta}_{2j}|(\xi)) \leq \|\widehat{\eta}_{2j}\|_{L^\infty(\mathbb{R}^2)} \|\mathbf{1}_{\ell, 2j, a}\|_{L^1(\mathbb{R}^2)} \lesssim N_{2j} N_\ell^{-3/2} \frac{|A_\ell|}{|A_{2j}|}.$$

Since $N_{2j}^{\alpha/2} \leq |A_{2j}|$ and $\|\widehat{\eta}_{2j}\|_{L^1(\mathbb{R}^2)} \lesssim 1$, we get

$$\begin{aligned}
\int \sum_{a \in A_{2j}} |\widehat{f}_{\Omega_a} * \widehat{\eta}_{2j}(\xi)|^2 d\xi &\lesssim N_{2j}^{1-\alpha/2} N_\ell^{-3/2} |A_\ell| \int \sum_{a \in A_{2j}} |\widehat{f}_{\Omega_a}|^2 * |\widehat{\eta}_{2j}|(\xi) d\xi \\
&\lesssim N_{2j}^{1-\alpha/2} N_\ell^{-3/2} |A_\ell| \sum_{a \in A_{2j}} \left\| \widehat{f}_{\Omega_a} \right\|_{L^2(\mathbb{R}^2)}^2 \\
&\lesssim N_{2j}^{1-\alpha/2} N_\ell^{-3/2} |A_\ell| \left\| \sum_{a \in A_{2j}} \widehat{f}_{\Omega_a} \right\|_{L^2(\mathbb{R}^2)}^2.
\end{aligned} \tag{4.13}$$

By (4.12) and (4.13), we obtain (4.3). \square

5. FOURIER DECAY

We abbreviate almost surely to a.s. When we say an inequality holds a.s., the corresponding implicit constant may depend on the measure ν , but the probability that the constant exist is 1. Let us consider a nondecreasing sequence $\{m_j\}_{j \in \mathbb{N}}$ such that $2 \leq m_j$. Let $M_j := m_j \cdots m_1$ and assume that for any $\epsilon > 0$ and $j \in \mathbb{N}$,

$$m_{j+1} \lesssim_\epsilon M_j^\epsilon. \tag{5.1}$$

Let \mathcal{I}_j be the collection of M_j^{-1} -intervals such that

$$\mathcal{I}_j = \{M_j^{-1}(k + [0, 1]) : 0 \leq k \leq M_j - 1\}.$$

We consider a sequence of random functions μ_j which satisfies the following conditions for some deterministic nondecreasing sequence $\{\beta_j\}_{j \in \mathbb{N}}$:

- $\mu_0 = \mathbf{1}_{[0,1]}$.
- $\mu_j = \beta_j \mathbf{1}_{E_j}$ where E_j is a union of intervals in \mathcal{I}_j .
- $\mathbb{E}(\mu_{j+1}(x) | E_j) = \mu_j(x)$ for all $x \in [0, 1]$.
- Conditioned on E_j , the sets $I_j \cap E_{j+1}$ are chosen independently for each $I_j \subseteq E_j$.

We will identify the functions μ_j with the measures $\mu_j dx$ and write $\|\mu_j\| := \mu_j([0, 1])$.

Remark 5.1. First three conditions are same with (M1)-(M3) in [22] and the last condition is weaker than (M4) in [22]. Since we only prove Fourier decay, it suffices to assume the weaker condition.

Let α_0 be a number such that

$$1 - \alpha_0 = \lim_{j \rightarrow \infty} \frac{\log \beta_j}{\log M_j}. \tag{5.2}$$

Note that (5.2) implies that there exist $j_0(\epsilon)$ for every $\epsilon > 0$ such that if $j \geq j_0(\epsilon)$, then

$$M_j^{1-\alpha_0-\epsilon} \leq \beta_j \leq M_j^{1-\alpha_0+\epsilon}. \tag{5.3}$$

Shmerkin and Suomala showed that μ_j converges weakly to a measure μ supported on $[0, 1]$ and the support of the measure μ is a Salem set a.s. so that $|\widehat{\mu}(\xi)| \lesssim_\sigma (1 + |\xi|)^{-\sigma/2}$ for any $\sigma < \alpha_0$, see [22, Theorem 4.2] and [23, Theorem 14.1].

Now, let ν_j be a measure defined as

$$\int f(x_1, x_2) d\nu_j := \int f(x_1, x_1^2) \mu_j(x_1) dx_1.$$

Similarly, ν_j converges weakly to a measure ν supported on \mathbb{P}^1 .

Proposition 5.2. *Suppose that μ_j is a sequence of random measures that satisfies the conditions above. For any $0 < \sigma < \alpha_0$, the limit measure ν satisfies the following inequality a.s.*

$$|\widehat{\nu}(\xi)| \lesssim_\sigma (1 + |\xi|)^{-\sigma/2} \quad \forall \xi \in \mathbb{R}^2. \quad (5.4)$$

Remark 5.3. For each $0 < \beta < 1$, Kaufman [13] constructed a Salem set of dimension β which is a subset of \mathbb{P}^1 . The proof of Proposition 5.2 is similar to [13] in that we used probabilistic argument and Van der Corput lemma. However, we showed that (5.4) is satisfied a.s. in more general settings.

One of the main ingredients of the proof is Hoeffding's inequality.

Theorem 5.4 (Hoeffding's inequality [12]). *Let X_1, \dots, X_n be independent real random variables such that $a_i \leq X_i \leq b_i$ and $S_n := X_1 + \dots + X_n$. For $t > 0$,*

$$\mathbb{P} \left(\left| S_n - \mathbb{E}(S_n) \right| > t \right) \leq 2 \exp \left(\frac{-2t^2}{\sum_{i=1}^n (b_i - a_i)^2} \right). \quad (5.5)$$

According to the size of $|\xi|$, we will use different arguments to obtain estimates for $|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)|$. If $|\xi| \leq M_{j+1}$, we use (5.5). If $|\xi| \geq m_j M_{j+1}^2$, we use van der Corput lemma. Lastly, if $M_{j+1} \leq |\xi| \leq m_{j+1} M_j^2$, we use both of them.

Lemma 5.5. *For fixed ξ such that $|\xi| \leq M_{j+1}$, we have the following tail bound. There exists a constant $C_1 > 0$ such that*

$$\mathbb{P} \left(|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-\sigma/2} \|\mu_j\|^{1/2} \middle| E_j \right) \lesssim \exp(-C_1 M_j^{1-\sigma} \beta_j \beta_{j+1}^{-2}). \quad (5.6)$$

Proof. The proof is similar to the proof of Theorem 4.2 in [22]. For $I_j \in \mathcal{I}_j$, we consider

$$X_{I_j} = \int_{I_j} \beta_{j+1} \mathbf{1}_{E_{j+1}} e(-x\xi_1 - x^2\xi_2) dx$$

for $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$. Then, $\sum_{I_j \in \mathcal{I}_j} X_{I_j} = \widehat{\nu}_{j+1}(\xi)$ and $\mathbb{E}(\mu_{j+1}(x)|E_j) = \mu_j(x)$ implies that

$$\begin{aligned} \mathbb{E} \left(\sum_{I_j \in \mathcal{I}_j} X_{I_j} | E_j \right) &= \mathbb{E}(\widehat{\nu}_{j+1}(\xi) | E_j) \\ &= \int e(-x\xi_1 - x^2\xi_2) \mathbb{E}(\mu_{j+1}(x) | E_j) dx \\ &= \int e(-x\xi_1 - x^2\xi_2) \mu_j(x) dx = \widehat{\nu}_j(\xi). \end{aligned}$$

Let $S_j := \sum_{I_j \in \mathcal{I}_j} X_{I_j}$, then

$$|S_j - \mathbb{E}(S_j)| = |\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)|$$

conditioned on E_j . Also, note that X_{I_j} are independent conditioned on E_j , since $I_j \cap E_{j+1}$ are chosen independently.

Next, we need to estimate $|X_{I_j}|$ for $I_j \subseteq E_j$. The random variables X_{I_j} are not real-valued. But by considering real and imaginary parts of X_{I_j} separately, we can use Hoeffding's inequality (5.5) with $\sum_{i=1}^n (b_i - a_i)^2$ replaced by the upper bound of $\sum_{I_j} |X_{I_j}|^2$.

We have

$$|X_{I_j}| \leq \beta_{j+1} M_j^{-1} \quad (5.7)$$

and the number of $I_j \subseteq E_j$ is $M_j \beta_j^{-1} \|\mu_j\|$. Thus, we obtain

$$\sum_{I_j \subseteq E_j} |X_{I_j}|^2 \leq \frac{\beta_{j+1}^2}{\beta_j M_j} \|\mu_j\|.$$

For $t = M_j^{-\sigma/2} \|\mu_j\|^{1/2}$, Hoeffding's inequality (5.5) implies (5.6). \square

Then, the unconditional version of Lemma 5.5 easily follows.

Corollary 5.6. *For fixed ξ , if $|\xi| \leq M_{j+1}$, then there exists a constant $C_1 > 0$ such that*

$$\mathbb{P} \left(|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-\sigma/2} \|\mu_j\|^{1/2} \right) \lesssim \exp(-C_1 M_j^{1-\sigma} \beta_j \beta_{j+1}^{-2}).$$

Proof. Note that Lemma 5.5 holds uniformly in E_j . Thus, the result can be immediately derived from the law of total probability. \square

For the rest of this section, we will use the following notations additionally. For fixed $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$ such that $\xi \neq 0$, consider $I_{j+1}(k) := M_{j+1}^{-1}(k + [0, 1]) \in \mathcal{I}_{j+1}$ and let $m(k)$ be the minimum of $|\xi_1 + 2x\xi_2|$ on $I_{j+1}(k)$. Also, if $\xi_2 \neq 0$, let $m(k_0)$ be the smallest among $m(k)$ and x_0 be a point in $[0, 1]$ such that $m(k_0) = |\xi_1 + 2x_0\xi_2|$. If there are two k 's such that $m(k)$ is the smallest, k_0 can be either of them.

Lemma 5.7. *For any $|\xi| \geq m_{j+1} M_j^2$, we have*

$$|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \lesssim_\epsilon M_{j+1}^{1-\alpha_0+\epsilon} |\xi|^{-1/2}. \quad (5.8)$$

Proof. By Van der Corput lemma (see for example [24, Proposition 2.2, Chapter 8]), we get

$$\left| \int_{I_{j+1}(k)} e(-x\xi_1 - x^2\xi_2) dx \right| \lesssim m(k)^{-1} \quad \text{if } m(k) \neq 0. \quad (5.9)$$

If $|\xi_1| \geq 10|\xi_2|$, then $|\xi_1 + 2x\xi_2| \gtrsim |\xi|$. Therefore,

$$\begin{aligned} |\widehat{\nu}_{j+1}(\xi)| &\leq \sum_{I_{j+1}(k) \subseteq E_{j+1}} \left| \int_{I_{j+1}(k)} \beta_{j+1} e(-x\xi_1 - x^2\xi_2) dx \right| \\ &\lesssim \beta_{j+1} M_{j+1} |\xi|^{-1} \\ &\leq \beta_{j+1} m_{j+1}^{1/2} |\xi|^{-1/2}. \end{aligned} \quad (5.10)$$

In the last inequality, we used that $|\xi| \geq m_{j+1} M_j^2$.

If $|\xi_1| \leq 10|\xi_2|$, we consider $I_{j+1}(k_0 + p)$ for $p = -1, 0$ or 1 . The interval $I_{j+1}(k_0 + p)$ can be divided into I_1 and I_2 where $I_1 = \{x : |x - x_0| \leq |\xi|^{-1/2}\}$ and $I_2 = \{x : |x - x_0| \geq |\xi|^{-1/2}\}$. For $x \in I_2$, since $|\xi_1 + 2x\xi_2|$ attains its minimum at x_0 ,

$$\begin{aligned} |\xi_1 + 2x\xi_2| &= |(\xi_1 + 2x_0\xi_2) + 2(x - x_0)\xi_2| \\ &\geq 2|x - x_0||\xi_2| \gtrsim |\xi|^{1/2}. \end{aligned}$$

Therefore, (5.9) implies that

$$\begin{aligned} \left| \int_{I_{j+1}(k_0+p)} \beta_{j+1} e(-x\xi_1 - x^2\xi_2) dx \right| &\lesssim \beta_{j+1} |I_1| + \beta_{j+1} |\xi|^{-1/2} \\ &\lesssim \beta_{j+1} |\xi|^{-1/2}. \end{aligned} \quad (5.11)$$

If $p \neq -1, 0, 1$, then $m(k_0 + p) \geq 2|x - x_0||\xi| \gtrsim pM_{j+1}^{-1}|\xi|$. Using (5.9), we obtain

$$\left| \int_{I_{j+1}(k_0+p)} \beta_{j+1} e(-x\xi_1 - x^2\xi_2) dx \right| \lesssim \frac{\beta_{j+1}M_{j+1}}{p|\xi|}. \quad (5.12)$$

Combining (5.11) and (5.12), we get

$$\begin{aligned} |\widehat{\nu_{j+1}}(\xi)| &\leq \sum_{I_{j+1}(k) \subseteq E_{j+1}} \left| \int_{I_{j+1}(k)} \beta_{j+1} e(-x\xi_1 - x^2\xi_2) dx \right| \\ &\lesssim \beta_{j+1}|\xi|^{-1/2} + \sum_{p=1}^{M_{j+1}} \frac{\beta_{j+1}M_{j+1}}{p|\xi|} \\ &\lesssim \beta_{j+1}m_{j+1}^{1/2} \log(M_{j+1})|\xi|^{-1/2}. \end{aligned}$$

We considered when $|\xi_1| \geq 10|\xi_2|$ and $|\xi_1| \leq 10|\xi_2|$. In either cases, (5.1) and (5.3) imply that

$$\begin{aligned} |\widehat{\nu_{j+1}}(\xi)| &\lesssim \beta_{j+1}m_{j+1}^{1/2} \log(M_{j+1})|\xi|^{-1/2} \\ &\lesssim_\epsilon M_{j+1}^{1-\alpha_0+\epsilon} |\xi|^{-1/2} \end{aligned}$$

for any ξ . The same argument works for $\widehat{\nu}_j$ with $j+1$ replaced by j . Thus, we have $|\widehat{\nu}_j(\xi)| \lesssim_\epsilon M_j^{1-\alpha_0+\epsilon} |\xi|^{-1/2}$. Since $M_j \leq M_{j+1}$, we get (5.8). \square

Lemma 5.8. *For fixed ξ , if $M_{j+1} \leq |\xi| \leq m_{j+1}M_j^2$, let k be an integer such that $0 \leq k \leq j-1$ and*

$$M_{j+1}M_k \leq |\xi| \leq M_{j+1}M_{k+1}.$$

For such k , the following tail bounds hold.

If $|\xi_1| \geq 10|\xi_2|$, there exists a constant $C_2 > 0$ such that

$$\mathbb{P} \left(|\widehat{\nu_{j+1}}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0-\epsilon)/2} \left(\frac{M_j}{|\xi|} \right)^{1/2} \|\mu_j\|^{1/2} \middle| E_j \right) \lesssim \exp(-C_2 M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2} m_{j+1}^{-1}) \quad (5.13)$$

If $|\xi_1| \leq 10|\xi_2|$, there exists a constant $C_3 > 0$ such that

$$\begin{aligned} \mathbb{P} \left(|\widehat{\nu_{j+1}}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0-\epsilon)/2} \left[\sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)) \right]^{1/2} \middle| E_j \right) \\ \lesssim \exp(-C_3 M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2}) \end{aligned} \quad (5.14)$$

where $I_i(x_0)$ is an interval of length M_i^{-1} centered at x_0 .

Proof. If $|\xi_1| \geq 10|\xi_2|$, by the argument similar to (5.10), we have $|X_{I_j}| \lesssim \beta_{j+1}m_{j+1}|\xi|^{-1}$ and the number of interval I_j in E_j is $\beta_j^{-1}M_j \|\mu_j\|$. Since $M_{j+1} \leq |\xi|$, we have

$$\begin{aligned} \sum_{I_j \subseteq E_j} |X_{I_j}|^2 &\lesssim \beta_{j+1}^2 \beta_j^{-1} m_{j+1} M_{j+1} |\xi|^{-2} \|\mu_j\| \\ &\leq \beta_{j+1}^2 \beta_j^{-1} m_{j+1} |\xi|^{-1} \|\mu_j\|. \end{aligned}$$

For $t = M_j^{(1-\alpha_0+\epsilon)/2} |\xi|^{-1/2} \|\mu_j\|^{1/2}$, Hoeffding's inequality (5.5) implies (5.13).

If $|\xi_1| \leq 10|\xi_2|$, we consider $I_k(x_0)$ first. If $I_j \subseteq I_k(x_0)$, we use (5.7). Since the number of I_j in $I_k(x_0)$ is $\beta_j^{-1}M_j\mu_j(I_k(x_0))$, we obtain that

$$\sum_{I_j \subseteq I_k(x_0)} |X_{I_j}|^2 \leq \beta_{j+1}^2 \beta_j^{-1} M_j^{-1} \mu_j(I_k(x_0)). \quad (5.15)$$

If $I \subseteq I_i(x_0)$ but $I \not\subseteq I_{i+1}(x_0)$ for $0 \leq i \leq k-1$, then $|x - x_0| \geq M_{i+1}^{-1} - M_j^{-1} \gtrsim M_{i+1}^{-1}$ for any $x \in I$. Therefore,

$$|\xi_1 + 2x\xi_2| \gtrsim |x - x_0||\xi| \gtrsim M_{i+1}^{-1}|\xi|.$$

We use (5.9) again and obtain that $|X_{I_j}| \lesssim \beta_{j+1}m_{j+1}M_{i+1}|\xi|^{-1}$. Since the number of I_j in $I_i(x_0)$ is $\lesssim \beta_j^{-1}M_j\mu_j(I_i(x_0))$ and $M_{j+1}|\xi|^{-1} \leq M_k^{-1}$, we get

$$\begin{aligned} \sum_{I_j \subseteq I_i(x_0)} |X_{I_j}|^2 &\lesssim \frac{\beta_{j+1}^2}{\beta_j M_j} \frac{M_{j+1}^2}{|\xi|^2} M_{i+1}^2 \mu_j(I_i(x_0)) \\ &\leq \frac{\beta_{j+1}^2}{\beta_j M_j} \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)). \end{aligned} \quad (5.16)$$

Combining (5.15) and (5.16), we get

$$\sum_{I_j \subseteq E_j} |X_{I_j}|^2 \lesssim \frac{\beta_{j+1}^2}{\beta_j M_j} \sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)).$$

For

$$t = M_j^{-(\alpha_0 - \epsilon)/2} \left(\sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)) \right)^{1/2},$$

Hoeffding's inequality (5.5) implies (5.14). \square

We also have the following unconditional version of Lemma 5.8.

Corollary 5.9. *For fixed ξ , if $M_{j+1} \leq |\xi| \leq m_{j+1}M_j^2$, for k defined in Lemma 5.8, we have the following tail bounds.*

If $|\xi_1| \geq 10|\xi_2|$, there exists a constant $C_2 > 0$ such that

$$\mathbb{P} \left(|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0 - \epsilon)/2} \left(\frac{M_j}{|\xi|} \right)^{1/2} \|\mu_j\|^{1/2} \right) \lesssim \exp(-C_2 M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2} m_{j+1}^{-1})$$

If $|\xi_1| \leq 10|\xi_2|$, there exists a constant $C_3 > 0$ such that

$$\begin{aligned} \mathbb{P} \left(|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0 - \epsilon)/2} \left[\sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)) \right]^{1/2} \right) \\ \lesssim \exp(-C_3 M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2}) \end{aligned}$$

where $I_i(x_0)$ is an interval of length M_i^{-1} centered at x_0 .

Proof. Note that Lemma 5.8 holds uniformly in E_j . Thus, the result immediately follows from the law of total probability. \square

In order to prove Proposition 5.2, we also need an estimate for $\mu_j(I_i(x_0))$.

Lemma 5.10. *We have the following estimate a.s.*

$$\mu_j(I_i(x_0)) \lesssim_\epsilon M_j^\epsilon \beta_i M_i^{-1}$$

where $j \geq i$ and the implicit constant is uniform in i, j and x_0 .

Proof. Note that $I_i(x_0) \subseteq I_i(k-1) \cup I_i(k) \cup I_i(k+1)$ for some $0 \leq k \leq M_i - 1$. Thus, it suffices to show that for any $j \geq i$,

$$\mu_j(I_i) \lesssim_\epsilon M_j^\epsilon \beta_i M_i^{-1} \quad \text{a.s.} \quad (5.17)$$

where the implicit constant is uniform in i, j and $I_i \in \mathcal{I}_i$.

For fixed $I_i \in \mathcal{I}_i$, let $X_{I_j} = \mu_{j+1}(I_j \cap I_i)$. Then, $\sum_{I_j \in \mathcal{I}_j} X_{I_j} = \mu_{j+1}(I_i)$ and

$$\mathbb{E}\left(\sum_{I_j \in \mathcal{I}_j} X_{I_j} | E_j\right) = \sum_{I_j \in \mathcal{I}_j} \mathbb{E}(\mu_{j+1}(I_j \cap I_i) | E_j) = \sum_{I_j \in \mathcal{I}_j} \mu_j(I_j \cap I_i) = \mu_j(I_i).$$

Also, X_{I_j} are independent conditioned on E_j , since $I_j \cap E_{j+1}$ are chosen independently.

We have $|X_{I_j}| \leq \beta_{j+1} M_j^{-1}$ and the number of I_j such that $I_j \subseteq I_i$ is $\beta_j^{-1} M_j \mu_j(I_i)$. Note that $I_j \subseteq I_i$ if and only if $X_{I_j} > 0$. Therefore, we have

$$\sum_{I_j \in \mathcal{I}_j} |X_{I_j}|^2 \leq \frac{\beta_{j+1}^2}{\beta_j M_j} \mu_j(I_i).$$

For $t = M_j^{\epsilon/4} M_j^{-1/2} \beta_j^{1/2} \mu_j(I_i)^{1/2}$, Hoeffding's inequality (5.5) implies that there exist a constant $C > 0$ such that

$$\mathbb{P}(|\mu_{j+1}(I_i) - \mu_j(I_i)| \geq t | E_j) \lesssim \exp(-C \beta_j^2 \beta_{j+1}^{-2} M_j^{\epsilon/2}).$$

Therefore, we have

$$\mathbb{P}(|\mu_{j+1}(I_i) - \mu_j(I_i)| \geq t \text{ for some } I_i \in \mathcal{I}_i) \lesssim M_i \exp(-C \beta_j^2 \beta_{j+1}^{-2} M_j^{\epsilon/2}). \quad (5.18)$$

By (5.3),

$$\sum_{i=1}^{\infty} \sum_{j=i}^{\infty} M_i \exp(-C \beta_j^2 \beta_{j+1}^{-2} M_j^{\epsilon/2}) \lesssim_\epsilon \sum_{i=1}^{\infty} M_i \exp(-C M_i^{\epsilon/4}) < \infty.$$

Borel-Cantelli lemma and (5.18) imply that there exist i_0 a.s. such that if $j \geq i \geq i_0$, then for any $I_i \in \mathcal{I}_i$,

$$\mu_{j+1}(I_i) \leq \mu_j(I_i) + M_j^{\epsilon/4} M_j^{-1/2} \beta_j^{1/2} \mu_j(I_i)^{1/2}. \quad (5.19)$$

Using (5.3) again, we can choose $i_1(\epsilon)$ such that if $i \geq i_1(\epsilon)$, then

$$M_i^{1-\alpha_0-\epsilon/4} \leq \beta_i \leq M_i^{1-\alpha_0+\epsilon/4}.$$

Now, we claim that if $i \geq \max(i_0, i_1(\epsilon))$, then there exists a constant $C' \geq 1$ such that for any $j \geq i$ and $I_i \in \mathcal{I}_i$

$$\mu_j(I_i) \leq C' j^2 \beta_i M_i^{-1} M_j^{\epsilon(1-2^{-j})}. \quad (5.20)$$

If $j = i$, then $\mu_j(I_i) \lesssim \beta_i M_i^{-1}$ uniformly in i and $I_i \in \mathcal{I}_i$. If (5.20) holds for $j = k$, (5.19) implies that

$$\mu_{k+1}(I_i) \leq C' k^2 \beta_i M_i^{-1} M_k^{\epsilon(1-2^{-k})} + (C')^{1/2} k \beta_i M_i^{-1} M_k^{\frac{\epsilon}{2}(1-2^{-k})} M_k^{\epsilon/4} \left(\frac{\beta_k M_k^{-1}}{\beta_i M_i^{-1}} \right)^{1/2}.$$

By the choice of $\epsilon_1(\epsilon)$,

$$\frac{\beta_k M_k^{-1}}{\beta_i M_i^{-1}} \leq \frac{M_k^{-\alpha_0 + \epsilon/4}}{M_i^{-\alpha_0 - \epsilon/4}} \leq (M_k M_i)^{\epsilon/4} \leq M_k^{\epsilon/2}.$$

Thus, we obtain that for any $I_i \in \mathcal{I}_i$,

$$\mu_{k+1}(I_i) \leq C'(k^2 + k)\beta_i M_i^{-1} M_k^{\epsilon(1-2^{-k-1})} \leq C'(k+1)^2 \beta_i M_i^{-1} M_{k+1}^{\epsilon(1-2^{-k-1})}.$$

Hence, we proved (5.20). The estimate (5.17) follows when $j \geq i \geq \max(i_0, i_1(\epsilon))$.

To consider the case $i \leq \max(i_0, i_1(\epsilon))$, we first show that $\|\mu_j\|$ is a.s. bounded. We have $\mathbb{E}(\|\mu_{j+1}\| | E_j) = \|\mu_j\|$ for all E_j and the law of total probability implies that $\mathbb{E}(\|\mu_{j+1}\|) = \mathbb{E}(\|\mu_j\|)$. By iterating the equality, we get $\mathbb{E}(\|\mu_j\|) = \mathbb{E}(\|\mu_0\|) = 1$ for all $j \in \mathbb{N} \cup \{0\}$. By Martingale convergence theorem (see for example [9, Theorem 4.2.11]), as $j \rightarrow \infty$, $\|\mu_j\|$ converges to $\|\mu\|$ a.s. with $\mathbb{E}(\|\mu\|) = 1$. Therefore, $\|\mu_j\|$ is a.s. bounded.

Therefore, if $i \leq \max(i_0, i_1(\epsilon))$, for any $j \geq i$ and $I_i \in \mathcal{I}_i$ we have

$$\mu_j(I_i) \leq \|\mu_j\| \lesssim_{\epsilon} \min_{i \leq \max(i_0, i_1(\epsilon))} (\beta_i M_i^{-1}) \leq M_j^{\epsilon} \beta_i M_i^{-1} \quad \text{a.s.}$$

Thus, we get (5.17). \square

Proof of Proposition 5.2. As in [22] and [23], we use Lemma 9A4 in [28]. Then, it suffices to consider only $\xi \in \mathbb{Z}^2$. Let Ω_j be the event that there exists $\xi \in \mathbb{Z}^2$ such that one of the following happens.

(1) If $|\xi| \leq M_{j+1}$, then

$$|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-\sigma/2} \|\mu_j\|^{1/2}.$$

(2) If $M_{j+1} \leq |\xi| \leq m_{j+1} M_j^2$ and $|\xi_1| \geq 10|\xi_2|$, then

$$|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0 - \epsilon)/2} \left(\frac{M_j}{|\xi|} \right)^{1/2} \|\mu_j\|^{1/2}.$$

(3) If $M_{j+1} \leq |\xi| \leq m_{j+1} M_j^2$ and $|\xi_1| \leq 10|\xi_2|$, then

$$|\widehat{\nu}_{j+1}(\xi) - \widehat{\nu}_j(\xi)| \geq M_j^{-(\alpha_0 - \epsilon)/2} \left[\sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)) \right]^{1/2}$$

for k such that $0 \leq k \leq j-1$ and $M_{j+1} M_k \leq |\xi| \leq M_{j+1} M_{k+1}$.

Since we only consider $\xi \in \mathbb{Z}^2$ such that $|\xi| \leq m_{j+1} M_j^2$, Corollary 5.6 and 5.9 imply that

$$\mathbb{P}(\Omega_j) \lesssim m_{j+1}^2 M_j^4 \exp(-C M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2} m_{j+1}^{-1}) \quad (5.21)$$

for some constant $C > 0$. For sufficiently large j and sufficiently small $\epsilon > 0$, it follows from (5.1) and (5.3) that

$$M_j^{1-\alpha_0+\epsilon} \beta_j \beta_{j+1}^{-2} m_{j+1}^{-1} \gtrsim_{\epsilon} M_j^{\epsilon/2}$$

Therefore, $\sum_{j=1}^{\infty} \mathbb{P}(\Omega_j) < \infty$.

Using Lemma 5.10, (5.3) and that $M_{j+1}M_k \leq |\xi| \leq M_{j+1}M_k$, a.s. we have

$$\begin{aligned} \sum_{i=0}^k \left(\frac{M_{i+1}}{M_k} \right)^2 \mu_j(I_i(x_0)) &\lesssim_\epsilon \sum_{i=0}^k M_j^\epsilon \beta_i M_i m_{i+1}^2 M_k^{-2} \\ &\lesssim_\epsilon M_j^\epsilon \beta_k m_{k+1}^2 M_k^{-1} \lesssim_\epsilon M_j^\epsilon M_k^\epsilon \left(\frac{M_{j+1}}{|\xi|} \right)^{\alpha_0 - \epsilon} \\ &\lesssim M_j^{2\epsilon} \left(\frac{M_j}{|\xi|} \right)^{\alpha_0 - \epsilon}. \end{aligned} \quad (5.22)$$

We apply Borel-Cantelli lemma to Ω_j , Since $\|\mu_j\|$ is a.s. bounded, Lemma 5.7 and (5.22) imply that a.s. there exists j_0 such that for $j \geq j_0$,

$$\begin{aligned} |\hat{\nu}_{j+1}(\xi) - \hat{\nu}_j(\xi)| &\lesssim M_j^{-\sigma/2} && \text{if } |\xi| \leq M_{j+1}, \\ &\lesssim_\epsilon M_j^{2\epsilon} |\xi|^{(-\alpha_0 + \epsilon)/2} && \text{if } M_{j+1} \leq |\xi| \leq m_{j+1} M_j^2, \\ &\lesssim_\epsilon M_{j+1}^{1-\alpha_0 + \epsilon} |\xi|^{-1/2} && \text{if } |\xi| \geq m_{j+1} M_j^2. \end{aligned}$$

When $M_{j+1} \leq |\xi| \leq m_{j+1} M_j^2$ and $|\xi_1| \geq 10|\xi_2|$, we used that $M_j/|\xi| \leq (M_j/|\xi|)^{\alpha_0 - \epsilon}$ for $\epsilon < \alpha_0$.

Let j_1, j_2 be numbers such that $M_{j_2} \leq |\xi| \leq M_{j_2+1}$ and $m_{j_1+1} M_{j_1}^2 \leq |\xi| \leq m_{j_1+2} M_{j_1+1}^2$. By telescoping, we obtain for $k \geq j_0$,

$$\begin{aligned} |\hat{\nu}_k(\xi) - \hat{\nu}_{j_0}(\xi)| &\lesssim_\epsilon \sum_{j_0 \leq j \leq j_1} M_{j+1}^{1-\alpha_0 + \epsilon} |\xi|^{-1/2} + \sum_{\max(j_0, j_1) < j < j_2} M_j^{2\epsilon} |\xi|^{(-\alpha_0 + \epsilon)/2} \\ &\quad + \sum_{\max(j_0, j_2) \leq j \leq k} M_j^{-\sigma/2}. \end{aligned}$$

We drop the first sum if $j_1 < j_0$ and we drop the first and the second sum if $j_1, j_2 \leq j_0$. Since $\sigma < \alpha_0$, for sufficiently small $\epsilon > 0$, we get

$$\begin{aligned} |\hat{\nu}_k(\xi) - \hat{\nu}_{j_0}(\xi)| &\lesssim_\epsilon M_{j_1+1}^{1-\alpha_0 + \epsilon} |\xi|^{-1/2} + M_{j_2-1}^{2\epsilon} |\xi|^{(-\alpha_0 + \epsilon)/2} + M_{j_2}^{-\sigma/2} \\ &\lesssim_\epsilon |\xi|^{-\sigma/2 + \epsilon}. \end{aligned}$$

Since $|\hat{\nu}_{j_0}(\xi)| \lesssim_{j_0} |\xi|^{-1/2}$, by letting $k \rightarrow \infty$, we obtain (5.4). \square

6. GLOBAL RESTRICTION ESTIMATE

We need to randomize the choice of $S_{j,a}$ in order to get the global restriction estimate.

Proposition 6.1. *Let $\{n_j\}_{j \in \mathbb{N}}$ be a sequence of positive numbers which satisfies (2.1), (2.2), (2.3) and (2.4). Let $\{\mu_j\}_{j \in \mathbb{N} \cup \{0\}}$ be a sequence of random measures on $[0, 1]$ which satisfies the following.*

- (1) μ_0, μ_1, \dots are constructed through the process in Section 2.
- (2) For each $j \in \mathbb{N}$ and for each $a \in A_j$, the set $S_{j,a}$ are chosen randomly and independently from Σ_j with probability distribution such that

$$\mathbb{E}(\mu_j(x) | E_{j-1}) = \mu_{j-1}(x).$$

For the corresponding sequence of random measures $\{\nu_j\}_{j \in \mathbb{N} \cup \{0\}}$, its limiting measure ν satisfies all conclusions of Theorem 1.2 a.s.

To prove Proposition 6.1, we need the next lemma to combine it with Tao's epsilon removal argument.

Lemma 6.2. *Let $p = 6/\alpha$ where $0 < \alpha < 1$ and assume that ν is the measure constructed in Section 2 and satisfies (1.5). For sufficiently large $R > 0$, suppose that $\{Q_i\}_{i=1}^M$ is a sparse collection of R -cubes in \mathbb{R}^2 , which means that their centers x_1, \dots, x_M are $R^B M^B$ -separated from each other for some sufficiently large constant B which will be determined later. For any function f supported on $\cup_{i=1}^M Q_i$, we have*

$$\left\| \widehat{f} \right\|_{L^2(d\nu)} \lesssim_{q,\epsilon} R^\epsilon \|f\|_{L^q(\mathbb{R}^2)} \quad (6.1)$$

where $1/p + 1/q = 1$.

Proof. Let $f = \sum_{i=1}^M f \mathbf{1}_{Q_i}$ where $\mathbf{1}_{Q_i}$ is a characteristic function on Q_i .

$$\begin{aligned} \left\| \widehat{f} \right\|_{L^2(d\nu)}^2 &= \int \left| \sum_{i=1}^M \widehat{f \mathbf{1}_{Q_i}} \right|^2 d\nu \\ &= \int \sum_{i=1}^M |\widehat{f \mathbf{1}_{Q_i}}|^2 d\nu + \int \sum_{i \neq j} \widehat{f \mathbf{1}_{Q_i}} \overline{\widehat{f \mathbf{1}_{Q_j}}} d\nu. \end{aligned}$$

It follows from Proposition 4.1 that

$$\begin{aligned} \sum_{i=1}^M \int |\widehat{f \mathbf{1}_{Q_i}}|^2 d\nu &\lesssim_{q,\epsilon} R^\epsilon \sum_{i=1}^M \|f \mathbf{1}_{Q_i}\|_{L^q(\mathbb{R}^2)}^2 \\ &\leq R^\epsilon \left(\sum_{i=1}^M \|f \mathbf{1}_{Q_i}\|_{L^q(\mathbb{R}^2)}^q \right)^{2/q} = R^\epsilon \|f\|_{L^q(\mathbb{R}^2)}^2. \end{aligned} \quad (6.2)$$

In the last inequality, we used that $1 \leq q \leq 2$.

For the second term, Young's convolution inequality implies that

$$\begin{aligned} \int \sum_{i \neq j} \widehat{f \mathbf{1}_{Q_i}} \overline{\widehat{f \mathbf{1}_{Q_j}}} d\nu &= \sum_{i \neq j} \iint f \mathbf{1}_{Q_i}(x) \overline{f \mathbf{1}_{Q_j}(y)} \widehat{d\nu}(y-x) dx dy \\ &\leq \sum_{i \neq j} \|f \mathbf{1}_{Q_i}\|_{L^q(\mathbb{R}^2)} \|f \mathbf{1}_{Q_j}\|_{L^q(\mathbb{R}^2)} \left\| \widehat{d\nu} \mathbf{1}_{i,j} \right\|_{L^r(\mathbb{R}^2)} \end{aligned} \quad (6.3)$$

where $\frac{2}{q} + \frac{1}{r} = 2$, so that $r = \frac{p}{2} \geq 1$ and

$$\mathbf{1}_{i,j} = \mathbf{1}_{B^2(2R)}(x - (x_j - x_i))$$

where $B^2(2R)$ is a ball of radius $2R$ centered at the origin. Since x_i and x_j are separated at least by $R^B M^B$, (1.5) implies that

$$\left\| \widehat{d\nu} \mathbf{1}_{i,j} \right\|_{L^r(\mathbb{R}^2)} \lesssim_\epsilon (R^B M^B)^{-\alpha/2+\epsilon} R^{2/r}. \quad (6.4)$$

We choose B such that

$$B \geq \frac{8}{\alpha}$$

and let $\epsilon = \alpha/4$ in (6.4). Since $r \geq 1$ we have,

$$\left\| \widehat{d\nu} \mathbf{1}_{i,j} \right\|_{L^r(\mathbb{R}^2)} \lesssim (RM)^{-B\alpha/4} R^2 \leq M^{-2}. \quad (6.5)$$

By (6.3) and (6.5), we obtain that

$$\begin{aligned}
\int \sum_{i \neq j} \widehat{f \mathbf{1}_{Q_i}} \overline{\widehat{f \mathbf{1}_{Q_j}}} d\nu &\lesssim M^{-2} \sum_{i \neq j} \|f \mathbf{1}_{Q_i}\|_{L^q(\mathbb{R}^2)} \|f \mathbf{1}_{Q_j}\|_{L^q(\mathbb{R}^2)} \\
&\leq M^{-1} \left(\sum_{i=1}^M \|f \mathbf{1}_{Q_i}\|_{L^q(\mathbb{R}^2)}^2 \right)^{1/2} \left(\sum_{j=1}^M \|f \mathbf{1}_{Q_j}\|_{L^q(\mathbb{R}^2)}^2 \right)^{1/2} \\
&\leq M^{-1} \|f\|_{L^q(\mathbb{R}^2)}^2.
\end{aligned} \tag{6.6}$$

In the last inequality, we used again that $1 \leq q \leq 2$. Combining (6.2) and (6.6), we obtain (6.1). \square

Let us turn to the proof of Proposition 6.1.

Proof of Proposition 6.1. The inequality (1.4) follows from Lemma 2.3. If we let $n_k^{1/2} = m_k$, $N_k^{1/2} = M_k$, $\beta_k = |E_k|^{-1}$ and $\alpha_0 = \alpha$, then $\{\mu_k\}_{k \in \mathbb{N} \cup \{0\}}$ satisfies all conditions of Proposition 5.2. Thus, $\widehat{\nu}$ satisfies (1.5) a.s. Now, we can use Tao's epsilon removal argument to prove (1.6). Lemma 6.2 replaces Lemma 3.2 in [27] and rest of the proof (1.6) is same as in [27, Theorem 1.2]. \square

7. PROOF OF THEOREM 1.3

The proof is based on Knapp-type example.

Proof. Since ν is supported on \mathbb{P}^{d-1} , there exists a measure μ on $[0, 1]^{d-1}$ such that

$$\int f(x) d\nu = \int f(x', |x'|^2) d\mu$$

for all ν -measurable function f on \mathbb{R}^d where $x' \in \mathbb{R}^{d-1}$. The assumption (1.7) implies that

$$\mu(B(x', r)) \lesssim_\alpha r^\alpha \tag{7.1}$$

for all $x' \in \mathbb{R}^{d-1}$ and $\forall r > 0$.

We claim that for any $\epsilon > 0$, there exist sequences $\{a_i\}_{i \in \mathbb{N}}$ and $\{r_i\}_{i \in \mathbb{N}}$ such that $\lim_{i \rightarrow \infty} r_i = 0$ and

$$r_i^{\alpha+\epsilon} < \mu(B(a_i, r_i)). \tag{7.2}$$

The measure μ is supported on a set of Hausdorff dimension α in \mathbb{R}^{d-1} . By Frostman's lemma (see for example [18, Theorem 2.7]), α is the supremum of s such that $\mu(B(x', r)) \leq r^s$ holds for all $x' \in \mathbb{R}^{d-1}$ and $r > 0$. Towards a contradiction, for given $\epsilon > 0$, we assume that $r_0 > 0$ is the smallest such that $\mu(B(x', r_0)) > r^{\alpha+\epsilon}$ for some $x' \in \mathbb{R}^{d-1}$. Let $\mu' = r_0^{\alpha+\epsilon} \mu$. Since ν is a probability measure, $\mu([0, 1]^{d-1}) = 1$. If $r \geq r_0$, then $\mu'(B(x, r)) \leq r_0^{\alpha+\epsilon} \mu([0, 1]^{d-1}) \leq r^{\alpha+\epsilon}$. If $r < r_0$, then $\mu'(B(x, r)) \leq \mu(B(x, r)) \leq r^{\alpha+\epsilon}$. Since μ' is also supported on a set of Hausdorff dimension α , it contradicts Frostman's lemma. Therefore, such r_0 does not exist and we established the claim (7.2).

Assume that the estimate (1.8) holds for some p and q and let $h_i(x') = \mathbf{1}_{[0, 1]^{d-1}}(r_i^{-1}(x' - a_i))$. Then, we have

$$\left(\int \left| \int h_i(x') e(-x' \cdot \xi' - |x'|^2 \xi_d) d\mu(x') \right|^p d\xi \right)^{1/p} \lesssim_{p,q} \left(\int |h_i(x')|^q d\mu \right)^{1/q} \tag{7.3}$$

where $\xi' \in \mathbb{R}^{d-1}$ and $\xi_d \in \mathbb{R}$. By (7.1), we obtain that

$$\left(\int |h_i(x')|^q d\mu \right)^{1/q} \lesssim_\alpha r_i^{\alpha/q}. \quad (7.4)$$

By change of variables $\xi' = r_i^{-1}\zeta'$ and $\xi_d = r_i^{-2}\zeta_d$, left-hand side of (7.3) becomes

$$r_i^{-(d+1)/p} \left(\int \left| \int h_i(x') e(-r_i^{-1}(x' - a_i) \cdot (\zeta' + 2\zeta_d r_i^{-1} a_i) - r_i^{-2}\zeta_d |x' - a_i|^2) d\mu(x') \right|^p d\zeta \right)^{1/p}. \quad (7.5)$$

By considering when $|\zeta' + 2\zeta_d r_i^{-1} a_i| \leq 1/100$ and $|\zeta_d| \leq 1/100$, (7.2) and (7.5) implies that

$$\begin{aligned} & \left(\int \left| \int h_i(x') e(-x' \cdot \xi' - |x'|^2 \xi_d) d\mu(x') \right|^p d\xi \right)^{1/p} \\ & \gtrsim r_i^{-(d+1)/p} \left| \int \mathbf{1}_{[0,1]^{d-1}}(r_i^{-1}(x' - a_i)) d\mu(x') \right| \gtrsim_{\alpha, \epsilon} r_i^{-(d+1)/p + \alpha + \epsilon}. \end{aligned} \quad (7.6)$$

We combine (7.3), (7.4) and (7.6), then for any r_i , we get

$$r_i^{-(d+1)/p + \alpha + \epsilon} \lesssim_{\alpha, \epsilon, p, q} r_i^{\alpha/q}.$$

Therefore,

$$\frac{\alpha}{q} \leq -\frac{d+1}{p} + \alpha + \epsilon.$$

Since ϵ is arbitrary, $p \geq q'(d+1)/\alpha$ if the estimate (1.8) holds. \square

8. APPENDIX

Proof of Lemma 3.1. First, assume that S_1 is a subset of $[-k_0, k_0]^d \cap \mathbb{Z}^d$. Let ψ be a non-negative smooth function supported on $[0, 1]^d$ such that $\psi \geq 1$ on $[1/4, 3/4]^d$. For $x \in [-k, k]^d$,

$$1 \leq \sum_{|n| \lesssim k} |\psi(x - n)|^p + \sum_{|n'| \lesssim 1} \sum_{|n| \lesssim k} |\psi(x - n - n'/2)|^p$$

where $n, n' \in \mathbb{Z}^d$, but $n' \neq 0$. Thus, we have

$$\begin{aligned} & \left\| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e((ka + b) \cdot x) \right\|_{L^p([0,1]^d)}^p = \int_{[0,k]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e\left(\left(a + \frac{b}{k}\right) \cdot x\right) \right|^p k^{-d} dx \\ & \leq \sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e(a \cdot x) e\left(\frac{b}{k} \cdot (x + n)\right) \psi(x) \right|^p dx \\ & + \sum_{|n'| \lesssim 1} \sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e(a \cdot (x + n'/2)) e\left(\frac{b}{k} \cdot (x + n + n'/2)\right) \psi(x) \right|^p dx. \end{aligned}$$

We will show that the first term is bounded by a constant multiple of

$$C_1 C_2 \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{p/2}. \quad (8.1)$$

By the same argument, the second term will be also bounded by a constant multiple of (8.1).

Since ψ is supported on $[0, 1]^d$, we have

$$\psi(x)e(b \cdot x/k) = \sum_{m \in \mathbb{Z}^d} \widehat{\psi}_b(m)e(m \cdot x)$$

where $\widehat{\psi}_b(m)$ is the m -th Fourier coefficient of $\psi(x)e(b \cdot x/k)$. Since $\sum_m |\widehat{\psi}_b(m)| \lesssim 1$ and S_1 is a subset of $[-k_0, k_0]^d$, we obtain

$$\begin{aligned} & \left(\sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e(a \cdot x) e\left(\frac{b}{k} \cdot (x+n)\right) \psi(x) \right|^p dx \right)^{1/p} \\ & \leq \sum_{m \in \mathbb{Z}^d} \left(\sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) e((a+m) \cdot x) \right|^p dx \right)^{1/p} \\ & \leq \sum_{m \in \mathbb{Z}^d} \left(\sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \left(\sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right) e(a \cdot x) \right|^p dx \right)^{1/p} \\ & \leq C_1 \sum_{m \in \mathbb{Z}^d} \left(\sum_{|n| \lesssim k} k^{-d} \left(\sum_{a \in S_1} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right|^2 \right)^{p/2} \right)^{1/p}. \end{aligned} \quad (8.2)$$

In the last inequality, we used (3.3).

First, we consider when $|m| \leq 10$. Since $p \geq 2$ and $|\psi| \lesssim 1$, we get

$$\begin{aligned} & \left(\sum_{|n| \lesssim k} k^{-d} \left(\sum_{a \in S_1} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right|^2 \right)^{p/2} \right)^{1/p} \\ & \leq \left(\sum_{a \in S_1} \left(\sum_{|n| \lesssim k} k^{-d} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \int_{[0,1]^d} \psi(x) e\left(\left(\frac{b}{k} - m\right) \cdot x\right) dx \right|^p \right)^{2/p} \right)^{1/2} \\ & \lesssim \left(\sum_{a \in S_1} \left(k^{-d} \sum_{|n| \lesssim k} \int_{[0,1]^d} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b}{k} \cdot (x+n)\right) \right|^p dx \right)^{2/p} \right)^{1/2} \\ & \lesssim \left(\sum_{a \in S_1} \left\| \sum_{b \in S_{2,a}} c_{a,b} e(b \cdot x) \right\|_{L^p([0,1]^d)}^2 \right)^{1/2} \leq C_2 \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{1/2}. \end{aligned} \quad (8.3)$$

In the case when $|m| \geq 10$, since ψ is supported on $[0, 1]^d$ and decreases rapidly, we can use the principle of stationary phase. Let $(b/k - m)_i$ denote the i -th coordinate of $(b/k - m)$ and denote $\nabla_{b,k,m}$ by

$$\nabla_{b,k,m} = \sum_{i=1}^d \left| \frac{b}{k} - m \right|^{-2} \left(\frac{b}{k} - m \right)_i \frac{\partial}{\partial x_i}.$$

Note that

$$\widehat{\psi}_b(m) = \int_{[0,1]^d} (\nabla_{b,k,m}^2 \psi(x)) e\left(\left(\frac{b}{k} - m\right) x\right) dx.$$

If we let

$$\tilde{c}_{a,b,i,j,m} = c_{a,b} \left| \frac{b}{k} - m \right|^{-4} \left(\frac{b}{k} - m \right)_i \left(\frac{b}{k} - m \right)_j,$$

it is well defined since $|b| \leq k$ and $|m| \geq 10$. Then we obtain

$$\left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right| \leq \sum_{i,j=1}^d \int_{[0,1]^d} \left| \sum_{b \in S_{2,a}} \tilde{c}_{a,b,i,j,m} e\left(\frac{b}{k} \cdot (x+n)\right) \partial_{ij} \psi(x) \right| dx.$$

Since $|\partial_{i,j} \psi| \lesssim 1$ uniformly in i and j , we obtain that

$$\begin{aligned} & \left(\sum_{|n| \lesssim k} k^{-d} \left(\sum_{a \in S_1} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right|^2 \right)^{p/2} \right)^{1/p} \\ & \lesssim \sum_{i,j=1}^d \left(\sum_{a \in S_1} \left(k^{-d} \sum_{|n| \lesssim k} \int_{[0,1]^d} \left| \sum_{b \in S_{2,a}} \tilde{c}_{a,b,i,j,m} e\left(\frac{b}{k} \cdot (x+n)\right) \right|^p dx \right)^{2/p} \right)^{1/2} \\ & \lesssim \sum_{i,j=1}^d \left(\sum_{a \in S_1} \left\| \sum_{b \in S_{2,a}} \tilde{c}_{a,b,i,j,m} e(b \cdot x) \right\|_{L^p([0,1]^d)}^2 \right)^{1/2} \\ & \leq C_2 \sum_{i,j=1}^d \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |\tilde{c}_{a,b,i,j,m}|^2 \right)^{1/2} \lesssim_d C_2 |m|^{-2} \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{ab}|^2 \right)^{1/2}. \end{aligned}$$

In the last inequality, we used that $|b/k - m| \approx |m|$ since $|m| \geq 10$. Therefore,

$$\begin{aligned} & \sum_{|m| \geq 10} \left(\sum_{|n| \lesssim k} k^{-d} \left(\sum_{a \in S_1} \left| \sum_{b \in S_{2,a}} c_{a,b} e\left(\frac{b \cdot n}{k}\right) \widehat{\psi}_b(m) \right|^2 \right)^{p/2} \right)^{1/p} \\ & \lesssim C_2 \sum_{|m| \geq 10} |m|^{-2} \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{1/2} \\ & \lesssim C_2 \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{1/2}. \end{aligned} \tag{8.4}$$

By (8.2), (8.3) and (8.4), it follows that

$$\begin{aligned} & \left(\sum_{|n| \lesssim k} k^{-d} \int_{[0,1]^d} \left| \sum_{a \in S_1} \sum_{b \in S_{2,a}} c_{a,b} e(a \cdot x) e\left(\frac{b}{k} \cdot (x+n)\right) \psi(x) \right|^p dx \right)^{1/p} \\ & \lesssim C_1 C_2 \left(\sum_{a \in S_1} \sum_{b \in S_{2,a}} |c_{a,b}|^2 \right)^{1/2}. \end{aligned}$$

We proved Lemma 3.1 when S_1 is a subset of $[-k_0, k_0]^d \cap \mathbb{Z}^d$. The estimate holds for arbitrary k_0 . Thus, letting $k_0 \rightarrow \infty$ finishes the proof. \square

Proof of Lemma 3.2. The inequality $K_p(E_j) \lesssim_p D_p(E_j)$ is well known (see for example [8, Theorem 13.1]). Thus, it suffices to prove the converse. Let R be a positive large number and S be a subset of $[-R, R]^d \cap \mathbb{Z}^d$ such that

$$\left\| \sum_{a \in S} c_a e(a \cdot x) \right\|_{L^p([0,1]^d)} \leq C \left(\sum_{a \in S} |c_a|^2 \right)^{1/2} \quad (8.5)$$

and let

$$\hat{f}_a(\xi) = \hat{f}(\xi) \mathbf{1}_{[0,1]^d}(R\xi - a).$$

Let us consider a Schwartz function η such that $|\eta| \geq 1$ on $[-1, 1]^d$ and $\hat{\eta}$ is supported on $[0, 1]^d$. Recall that $B^d(R)$ is a cube of side length R in \mathbb{R}^d centered at the origin. We will show that for any f such that $f = \sum_{a \in S} f_a$,

$$\|f\|_{L^p(B^d(R))} \lesssim C \left(\sum_{a \in S} \|f_a\|_{L^p(\eta_R)}^2 \right)^{1/2} \quad (8.6)$$

where $\eta_R(x) = \eta(x/R)$ and C is the constant in (8.5). Then (8.6) implies that

$$\|f\|_{L^p(\mathbb{R}^d)} \lesssim C \left(\sum_{a \in S} \|f_a\|_{L^p(\mathbb{R}^d)}^2 \right)^{1/2}. \quad (8.7)$$

For the proof of (8.7) from (8.6), see [8, Proposition 9.15] and [16, Chapter 4.1]. Thus, Lemma 3.2 easily follows.

Now we prove (8.6). We have

$$\begin{aligned} \|f\|_{L^p(B^d(R))} & \leq \|f \eta_R\|_{L^p(B^d(R))} \\ & = \left\| \sum_{a \in S} \int \hat{f}_a * \hat{\eta}_R(\xi) e(-x \cdot \xi) d\xi \right\|_{L^p(B^d(R))}. \end{aligned} \quad (8.8)$$

The function $\hat{f}_a * \hat{\eta}_R$ is supported on a cube of side length $O(R^{-1})$ centered at a/R . Therefore, by change of variables $\xi = R^{-1}\zeta$ and $R^{-1}x = z$, (8.8) is

$$\begin{aligned} & \leq R^{-d+d/p} \sum_{|c| \lesssim 1} \int_{[0,1]^d} \left\| \sum_{a \in S} \hat{f}_a * \hat{\eta}_R(R^{-1}(\zeta + a + c)) e(-z \cdot (\zeta + a + c)) \right\|_{L^p(B^d(1))} d\zeta \\ & \lesssim R^{-d+d/p} \sum_{|c| \lesssim 1} \int_{[0,1]^d} \left\| \sum_{a \in S} \hat{f}_a * \hat{\eta}_R(R^{-1}(\zeta + a + c)) e(-z \cdot a) \right\|_{L^p(B^d(1))} d\zeta. \end{aligned}$$

By (8.5), we obtain that

$$\begin{aligned}
\|f\|_{L^p(B^d(R))} &\lesssim CR^{-d+d/p} \sum_{|c|\lesssim 1} \int_{[0,1]^d} \left(\sum_{a \in S} |\widehat{f}_a * \widehat{\eta}_R(R^{-1}(\zeta + a + c))|^2 \right)^{1/2} d\zeta \\
&\lesssim CR^{-d+d/p} \sum_{|c|\lesssim 1} \left(\sum_{a \in S} \int_{[0,1]^d} |\widehat{f}_a * \widehat{\eta}_R(R^{-1}(\zeta + a + c))|^2 d\zeta \right)^{1/2} \\
&\lesssim CR^{-d/2+d/p} \left(\sum_{a \in S} \left\| \widehat{f}_a * \widehat{\eta}_R(\xi) \right\|_{L^2(\mathbb{R}^d)}^2 \right)^{1/2} \\
&\leq CR^{-d/2+d/p} \left(\sum_{a \in S} \|f_a \eta_R\|_{L^2(\mathbb{R}^d)}^2 \right)^{1/2} \\
&\leq CR^{-d/2+d/p} \left(\sum_{a \in S} \|f_a\|_{L^p(\eta_R)}^2 \|\eta_R\|_{L^1(\mathbb{R}^d)}^{1-2/p} \right)^{1/2}.
\end{aligned}$$

Since $\|\eta_R\|_{L^1} \approx R^d$, this completes the proof of (8.6). \square

REFERENCES

- [1] Jong-Guk Bak and Andreas Seeger. Extensions of the Stein-Tomas theorem. *Math. Res. Lett.*, 18(4):767–781, 2011.
- [2] A. Benedek and R. Panzone. The space L^p , with mixed norm. *Duke Math. J.*, 28:301–324, 1961.
- [3] J. Bourgain. Bounded orthogonal systems and the $\Lambda(p)$ -set problem. *Acta Math.*, 162(3-4):227–245, 1989.
- [4] Alan Chang, Jaume de Dios Pont, Rachel Greenfeld, Asgar Janneshan, Zane Kun Li, and José Madrid. Decoupling for fractal subsets of the parabola. *Math. Z.*, 301(2):1851–1879, 2022.
- [5] Xianghong Chen. A Fourier restriction theorem based on convolution powers. *Proc. Amer. Math. Soc.*, 142(11):3897–3901, 2014.
- [6] Xianghong Chen. Sets of Salem type and sharpness of the L^2 -Fourier restriction theorem. *Trans. Amer. Math. Soc.*, 368(3):1959–1977, 2016.
- [7] Xianghong Chen and Andreas Seeger. Convolution powers of Salem measures with applications. *Canad. J. Math.*, 69(2):284–320, 2017.
- [8] Ciprian Demeter. *Fourier restriction, decoupling, and applications*, volume 184 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2020.
- [9] Rick Durrett. *Probability—theory and examples*, volume 49 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2019. Fifth edition of [MR1068527].
- [10] Kyle Hambrook and Izabella Łaba. On the sharpness of Mockenhaupt’s restriction theorem. *Geom. Funct. Anal.*, 23(4):1262–1277, 2013.
- [11] Kyle Hambrook and Izabella Łaba. Sharpness of the Mockenhaupt-Mitsis-Bak-Seeger restriction theorem in higher dimensions. *Bull. Lond. Math. Soc.*, 48(5):757–770, 2016.
- [12] Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *J. Amer. Statist. Assoc.*, 58:13–30, 1963.
- [13] R. Kaufman. Random measures on planar curves. *Ark. Mat.*, 14(2):245–250, 1976.
- [14] Thomas William Körner. Hausdorff and Fourier dimension. *Studia Math.*, 206(1):37–50, 2011.
- [15] Izabella Łaba and Hong Wang. Decoupling and near-optimal restriction estimates for Cantor sets. *Int. Math. Res. Not. IMRN*, 2018(9):2944–2966, 2018.
- [16] Zane Kun Li. *Decoupling for the parabola and connections to efficient congruencing*. University of California, Los Angeles, 2019.
- [17] Pertti Mattila. Hausdorff dimension, projections, and the Fourier transform. *Publ. Mat.*, 48(1):3–48, 2004.

- [18] Pertti Mattila. *Fourier analysis and Hausdorff dimension*, volume 150 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2015.
- [19] Themis Mitsis. A Stein-Tomas restriction theorem for general measures. *Publ. Math. Debrecen*, 60(1-2):89–99, 2002.
- [20] Themis Mitsis. A Stein-Tomas restriction theorem for general measures. *Publ. Math. Debrecen*, 60(1-2):89–99, 2002.
- [21] G. Mockenhaupt. Salem sets and restriction properties of Fourier transforms. *Geom. Funct. Anal.*, 10(6):1579–1587, 2000.
- [22] Pablo Shmerkin and Ville Suomala. A class of random Cantor measures, with applications. In *Recent developments in fractals and related fields*, Trends Math., pages 233–260. Birkhäuser/Springer, Cham, 2017.
- [23] Pablo Shmerkin and Ville Suomala. Spatially independent martingales, intersections, and applications. *Mem. Amer. Math. Soc.*, 251(1195):v+102, 2018.
- [24] Elias M. Stein and Rami Shakarchi. *Functional analysis*, volume 4 of *Princeton Lectures in Analysis*. Princeton University Press, Princeton, NJ, 2011. Introduction to further topics in analysis.
- [25] Michel Talagrand. Sections of smooth convex bodies via majorizing measures. *Acta Math.*, 175(2):273–300, 1995.
- [26] Michel Talagrand. Upper and lower bounds for stochastic processes. decomposition theorems. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, 60, 2021.
- [27] Terence Tao. The Bochner-Riesz conjecture implies the restriction conjecture. *Duke Math. J.*, 96(2):363–375, 1999.
- [28] Thomas H. Wolff. *Lectures on harmonic analysis*, volume 29 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003. With a foreword by Charles Fefferman and a preface by Izabella Laba, Edited by Laba and Carol Shubin.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY, USA
 Email address: `dryou@ur.rochester.edu`