

OpenMP Fortran programs for solving the time-dependent dipolar Gross-Pitaevskii equation

Luis E. Young-S.^a, Paulsamy Muruganandam^b, Antun Balaž^c, Sadhan K. Adhikari^{d,*}

^a*Grupo de Modelado Computacional, Facultad de Ciencias Exactas y Naturales, Universidad de Cartagena, 130014 Cartagena, Bolivar, Colombia*

^b*Department of Physics, Bharathidasan University, Palkalaiperur Campus, Tiruchirappalli 620024, Tamilnadu, India*

^c*Institute of Physics Belgrade, University of Belgrade, Pregrevica 118, 11080 Belgrade, Serbia*

^d*Instituto de Física Teórica, UNESP – Universidade Estadual Paulista, 01.140-70 São Paulo, São Paulo, Brazil*

Abstract

In this paper we present Open Multi-Processing (OpenMP) Fortran 90/95 versions of previously published numerical programs for solving the dipolar Gross-Pitaevskii (GP) equation including the contact interaction in one, two and three spatial dimensions. The atoms are considered to be polarized along the z axis and we consider different cases, e.g., stationary and non-stationary solutions of the GP equation for a dipolar Bose-Einstein condensate (BEC) in one dimension (along x and z axes), two dimensions (in x - y and x - z planes), and three dimensions. The algorithm used is the split-step semi-implicit Crank-Nicolson scheme for imaginary- and real-time propagation to obtain stationary states and BEC dynamics, respectively, as in the previous version [R. Kishor Kumar et al., Comput. Phys. Commun. 195, 117 (2015)]. These OpenMP versions have significantly reduced execution time in multicore processors.

Keywords: Dipolar Bose-Einstein condensate; Contact and dipolar interaction; Gross-Pitaevskii equation; Split-step Crank-Nicolson scheme; Fortran programs; Partial differential equation

New version program summary

Program title: DBEC-GP-OMP, a program package containing programs `imag3d-th.f90`, `real3d-th.f90`, `imag2dXY-th.f90`, `real2dXY-th.f90`, `imag2dXZ-th.f90`, `real2dXZ-th.f90`, `imag1dX-th.f90`, `real1dX-th.f90`, `imag1dZ-th.f90`, `real1dZ-th.f90`, with `fft3.f03` and `fftw3.mod`.

CPC Library link to program files:

Program obtainable from: CPC Program Library, Queens University, Belfast, N. Ireland.

Licensing provisions: Apache License 2.0

Programming language: Open Multi-Processing (OpenMP) Fortran 90/95. The program is tested with the GNU, Intel, and Oracle (former Sun) compilers.

Journal Reference of previous version: Comput. Phys. Commun. 195 (2015) 117. <https://doi.org/10.1016/j.cpc.2015.03.024>

Does the new version supersede the previous version?: Yes

Nature of problem: The present Open OpenMP Fortran 90/95 programs solve the time-dependent nonlinear partial differential Gross-Pitaevskii (GP) equation for a trapped dipolar Bose-Einstein condensate (BEC) in one (1D), two (2D), and three (3D) spatial dimensions.

Solution method: We employ the split-step Crank-Nicolson scheme to discretize the time-dependent GP equation in space and time. The discretized equation is then solved by imaginary- or real-time propagation, employing adequately small space and time steps, to yield the solution of stationary and non-stationary problems, respectively.

Reason for new version: Previously published Fortran programs [1] for solving the GP equation for a BEC have become useful

*Corresponding author.

Email addresses: `lyoung@unicartagena.edu.co` (Luis E. Young-S.), `anand@bdu.ac.in` (Paulsamy Muruganandam), `antun@ipb.ac.rs` (Antun Balaž), `sk.adhikari@unesp.br` (Sadhan K. Adhikari)

tools. These programs have been translated to the C programming language [2] and later extended to the more complex scenario of dipolar atoms [3], spinor condensates [4], and rotating condensates [5]. Now virtually all computers have multi-core processors and some have motherboards with more than one physical computer processing unit (CPU), which may increase the number of available CPU cores on a single computer to several tens. The Fortran [6] and C [7] programs for a nondipolar BEC have been adopted to be very fast on such multi-core modern computers. The C programs for a dipolar BEC have been adopted to multicore processors to yield OpenMP, OpenMP/MPI, and CUDA/MPI versions [8, 9].

The available Fortran 90/95 and C programs for the solution of the GP equation for a dipolar BEC in 1D, 2D, and 3D are enjoying widespread use [1, 2, 3]. Later, OpenMP/MPI, and CUDA/MPI [8, 9] versions of these C programs were published, which highly reduce the CPU times in multicore processors. Although, Fortran 90/95 is a popular programming language, the previous Fortran 90/95 programs for a dipolar BEC [3], lacking an efficient parallelization, suffer from a drawback of requiring large CPU time. In this paper we present the OpenMP version of the Fortran 90/95 programs for the solution of the GP equation for a dipolar BEC in 1D, 2D, and 3D, ideal for an efficient execution in multicore processors, which highly reduce the CPU time.

Summary of revisions:

The program package DBEC-GP-OMP contains 3D programs `imag3d-th.f90`, `real3d-th.f90`, 2D programs `imag2dXY-th.f90`, `real2dXY-th.f90`, `imag2dXZ-th.f90`, `real2dXZ-th.f90`, and 1D programs `imag1dX-th.f90`, `real1dX-th.f90`, `imag1dZ-th.f90`, `real1dZ-th.f90`, in the directory `src`, as well as the files `makefile`, `readme.txt`, and `readme-fftw.txt`. The 3D and 2D programs are OpenMP versions leading to a significant reduction in execution time in a multicore processor. However, in 1D the supplied programs are autparallel versions as there is no further reduction in the execution time in the OpenMP versions. The `makefile` allows an automated compilation of the programs using different supported compilers (GNU, Intel, Sun Oracle) by a simple `make` command, as in Ref. [3]. However, the 1D and 2D dipolar programs in previous publications [3, 8, 9] erroneously used the 3D nonlinearities, in place of corresponding 1D and 2D nonlinearities, in numerical calculations. The corresponding arXiv archive versions do not have these errors. The file `readme.txt` contains instructions on how to compile and run the programs. The file `readme-fftw.txt` provides instructions for installing the fast Fourier transformation (FFT) routine in Linux/Unix operating systems unless this routine is preinstalled. All input parameters are placed in the beginning of each program in MODULEs `GPE_DATA` and `COMM_DATA`. The directory output contains examples of matching outputs of imaginary- and real-time propagation programs in sub-directories with a generic name, e.g. `imag3d-th`, `real2dXY-th`, etc. The reader is advised to consult Ref. [3] for details. The imaginary-time programs implement the calculation starting from an initial analytic Gaussian wave function using a non-zero value of the parameter `NSTP`, whereas the real-time programs use the converged solution of the imaginary-time programs as the initial state of calculation employing `NSTP = 0`. The FFT algorithm works faster and more efficiently when we take the number of space grid points – N in 1D and NX, NY, NZ in 2D and 3D – in powers of 2, e.g. 2^n with n an integer, and should be so chosen for reducing the execution time. The operational scheme for running the codes is identical to that in Ref. [3].

Acknowledgments

L.E.Y.-S. acknowledges the financial support by the Vicerrectoria de Investigaciones, Universidad de Cartagena, Colombia through Project No. 019-2021. P.M. acknowledges the financial support by the Council of Scientific and Industrial Research (CSIR), India under Grant No. 03(1422)/18/EMR-II, and Science and Engineering Research Board (SERB), India under Grant No. CRG/2019/004059. A.B. acknowledges funding provided by the Institute of Physics Belgrade, through a grant by the Ministry of Education, Science, and Technological Development of the Republic of Serbia. S.K.A. acknowledges support by the CNPq, Brazil through grant 301324/2019-0.

References

- [1] P. Muruganandam, S. K. Adhikari, Fortran programs for the time-dependent Gross-Pitaevskii equation in a fully anisotropic trap, *Comput. Phys. Commun.* 180 (2009) 1888; arXiv:0904.3131.
- [2] D. Vudragović, I. Vidanović, A. Balaž, P. Muruganandam, S. K. Adhikari, C programs for solving the time-dependent Gross-Pitaevskii equation in a fully anisotropic trap, *Comput. Phys. Commun.* 183 (2012) 2021; arXiv:1206.1361.
- [3] R. Kishor Kumar, L.E. Young-S., D. Vudragović, A. Balaž, P. Muruganandam, S.K. Adhikari, Fortran and C programs for the time-dependent dipolar Gross-Pitaevskii equation in an anisotropic trap, *Comput. Phys. Commun.* 195 (2015) 117; arXiv:1506.03283.
- [4] R. Ravisankar, D. Vudragović, P. Muruganandam, A. Balaz, S. K. Adhikari, Spin-1 spin-orbit- and Rabi-coupled Bose–Einstein condensate solver, *Comput. Phys. Commun.* 259 (2021) 107657; arXiv:2009.13507.
- [5] R. K. Kumar, V. Lončar, P. Muruganandam, S. K. Adhikari, A. Balaž, C and Fortran OpenMP programs for rotating Bose–Einstein condensates, *Comput. Phys. Commun.* 240 (2019) 74; arXiv:1906.06327.
- [6] L. E. Young-S., P. Muruganandam, S. K. Adhikari, V. Loncar, D. Vudragović, Antun Balaž, OpenMP GNU and Intel Fortran programs for solving the time-dependent Gross-Pitaevskii equation, *Comput. Phys. Commun.* 220 (2017) 503; arXiv:1709.04423.

- [7] B. Satarić, V. Slavnić, A. Belić, A. Balaž, P. Muruganandam, and S. K. Adhikari, Hybrid OpenMP/MPI programs for solving the time-dependent Gross-Pitaevskii equation in a fully anisotropic trap, *Comput. Phys. Commun.* 200 (2016) 411; arXiv:1601.04641.
- [8] V. Lončar, L.E. Young-S., S. Škrbić, P. Muruganandam, S.K. Adhikari, A. Balaž, OpenMP, OpenMP/MPI, and CUDA/MPI C programs for solving the time-dependent dipolar Gross-Pitaevskii equation, *Comput. Phys. Commun.* 209 (2016) 190; arXiv:1610.05329.
- [9] V. Lončar, A. Balaž, A. Bogojević, S. Škrbić, P. Muruganandam, S.K. Adhikari, CUDA programs for solving the time-dependent dipolar Gross-Pitaevskii equation in an anisotropic trap, *Comput. Phys. Commun.* 200 (2016) 406; arXiv:1601.04640.