CENTRAL SUPPORT FOR TRIANGULATED CATEGORIES

HENNING KRAUSE

ABSTRACT. For any essentially small triangulated category the centre of its lattice of thick subcategories is introduced; it is a spatial frame and yields a notion of central support. A relative version of this centre recovers the support theory for tensor triangulated categories and provides a universal notion of cohomological support. Along the way we establish Mayer–Vietoris sequences for pairs of commuting subcategories.

1. INTRODUCTION

Triangulated categories are algebraic desiderata as they were introduced in order to deal with cohomologies arising in geometric or topological settings. But over the last years triangulated categories have been turned into geometric objects, thanks to the notion of support [1, 7, 9, 17, 19]. In this work we pursue this direction¹ and study in which way the geometry of an essentially small triangulated category \mathcal{T} is captured by its lattice of thick subcategories, and more precisely by its centre. Our first result (Corollary 3.8) is meant to illustrate this because many applications of geometric flavour involve Mayer–Vietoris sequences [4, 7, 21, 23].

Theorem 1. For a pair of thick subcategories $\mathcal{U}, \mathcal{V} \subseteq \mathcal{T}$ the following are equivalent.

- The pair U, V is commuting, that is, any morphism between objects of U and V factors through an object of U ∧ V.
- (2) For each pair of objects $X, Y \in T$ there exists a long exact sequence
- $\cdots \to \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(X,Y) \to \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(X,Y) \oplus \operatorname{Hom}_{\mathcal{T}/\mathcal{V}}(X,Y) \to$

$$\rightarrow \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\vee\mathcal{V})}(X,Y) \rightarrow \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(X,\Sigma Y) \rightarrow \cdots$$

The notion of support for objects of a triangulated category \mathcal{T} amounts to additional geometric structure. It requires a topological space and assigns to an object of \mathcal{T} a subset of this space. For instance, the space could be given by the set of prime tensor ideals provided that \mathcal{T} admits a symmetric tensor product [1], or by the set of prime ideals of a graded commutative ring acting centrally on \mathcal{T} [7].

An essential feature of any notion of support is a parameterisation of thick subcategories by subsets of a topological space. We take as input a lattice T of thick subcategories. Then its *centre*

 $Z(T) := \{ \mathcal{U} \in T \mid \mathcal{U} \text{ and } \mathcal{V} \text{ commute for all } \mathcal{V} \in T \}$

provides such a space because Z(T) is a *spatial frame*; this means it identifies with the lattice of open sets of a topological space. This space may be used to define a support for any object such that the central subcategories are parameterised by the open subsets.

Date: January 25, 2023.

²⁰²⁰ Mathematics Subject Classification. 18G80.

Key words and phrases. Centre, distributive lattice, Mayer–Vietoris sequence, support, thick subcategory, triangulated category.

¹We follow the slogan 'geometry is algebra is geometry', which is an adaptation of Ringel's 'algebra is geometry is algebra' in [22].

We summarise and state our main result (Theorem 5.2) for the lattice of thick subcategories Thick \mathcal{T} of an essentially small triangulated category \mathcal{T} .

Theorem 2. Let $T \subseteq$ Thick \mathfrak{T} be a sublattice which is closed under arbitrary joins. Then its centre $Z(T) \subseteq T$ is also a sublattice and closed under arbitrary joins. Moreover, Z(T) is a spatial frame and for any pair $\mathfrak{U}, \mathfrak{V}$ in T there is a Mayer-Vietoris sequence provided at least one of \mathfrak{U} and \mathfrak{V} is central.

A consequence is that T is a distributive lattice when Z(T) = T, but the converse is not necessarily true. For example, Z(T) = T for T the lattice of thick tensor ideals when \mathcal{T} is a rigid tensor triangulated category (Example 5.3). Thus we provide foundations for *triangular geometry* in the spirit of tensor triangular geometry [3, 17, 19], with additional structure on \mathcal{T} given by a distinguished sublattice $T \subseteq$ Thick \mathcal{T} .

Our discussion of central thick subcategories is inspired by recent work of Gratz and Stevenson [12], and in particular its presentation at the Abel Symposium 2022 in Ålesund. A crucial observation in [12] is the relevance of *distributivity* for the lattice of thick subcategories. In this work we provide evidence for a close connection between distributivity and *commutativity*, responding in particular to the quest for 'interesting distributive sublattices of Thick \mathcal{T} ' in [12, §8].

This paper is organised as follows. In §2 we briefly discuss our main tool: the category of cohomological functors of a triangulated category which identifies with its ind-completion. In §3 we introduce central subcategories and show that they form a spatial frame. Along the way we establish a couple of Mayer–Vietoris sequences for any pair of commuting subcategories. In §4 we provide many examples, including triangulated categories with additional structure given by a tensor product or a central ring action. This motivates a relative version of the lattice of central subcategories which is discussed in the final §5.

Acknowledgements. It is a pleasure to thank Greg Stevenson for inspiring discussions related to this work. Much thanks also to Dave Benson for providing specific examples from modular representation theory, and to Kent Vashaw for help with the non-commutative tensor products in this paper.

Part of this work was done during the Trimester Program 'Spectral Methods in Algebra, Geometry, and Topology' at the Hausdorff Institute in Bonn. It is a pleasure to thank for hospitality and for funding by the Deutsche Forschungsgemeinschaft (DFG) under Excellence Strategy EXC-2047/1-390685813.

2. The category of cohomological functors

Triangulated categories were introduced in order to deal with cohomologies arising in geometric or topological settings. If we wish to treat triangulated categories as geometric objects in their own right, it seems natural to study them by cohomological methods, using the category of cohomological functors. In this section we collect some basic and well-known facts.

Cohomological functors. Let \mathcal{T} be an essentially small triangulated category with suspension $\Sigma: \mathcal{T} \xrightarrow{\sim} \mathcal{T}$. Recall that a functor $\mathcal{T}^{\text{op}} \to \text{Ab}$ into the category of abelian groups is *cohomological* if it takes exact triangles to exact sequences. We denote by $\text{Coh}\mathcal{T}$ the category of cohomological functors. Morphisms in $\text{Coh}\mathcal{T}$ are natural transformation and the Yoneda functor $\mathcal{T} \to \text{Coh}\mathcal{T}$ sending $X \in \mathcal{T}$ to

$$H_X = \operatorname{Hom}_{\mathfrak{T}}(-, X)$$

is fully faithful. The suspension Σ extends to a functor $\operatorname{Coh} \mathfrak{T} \xrightarrow{\sim} \operatorname{Coh} \mathfrak{T}$ by taking F in $\operatorname{Coh} \mathfrak{T}$ to $F \circ \Sigma^{-1}$; we denote this again by Σ .

It is convenient to view Coh T as a full subcategory of the category Mod T of all additive functors $T^{op} \rightarrow Ab$. Note that (co)limits in Mod T are computed pointwise.

For E and F in Mod \mathfrak{T} we write $\operatorname{Hom}(E, F)$ for the set of morphisms from E to F. Thus $\operatorname{Hom}(H_X, F) \cong F(X)$ for X in \mathfrak{T} , by Yoneda's lemma.

Any additive functor $F\colon \mathbb{T}^{\mathrm{op}}\to \mathrm{Ab}$ can be written canonically as a colimit of representable functors

$$\operatorname{colim}_{H_X \to F} H_X \xrightarrow{\sim} F$$

where the colimit is taken over the slice category \mathfrak{T}/F ; see [13, Proposition 3.4]. Objects in \mathfrak{T}/F are morphisms $H_X \to F$ where X runs through the objects of \mathfrak{T} . A morphism in \mathfrak{T}/F from $H_X \xrightarrow{\phi} F$ to $H_{X'} \xrightarrow{\phi'} F$ is a morphism $\alpha \colon X \to X'$ in \mathfrak{T} such that $\phi'H_\alpha = \phi$.

A theorem of Lazard says that a module is flat if and only if it is a filtered colimit of finitely generated free modules; this has been generalised to functor categories by Oberst and Röhrl. The following lemma shows that cohomological and flat functors agree; this is well-known, for instance from [20, Lemma 2.1].

Lemma 2.1. The cohomological functors $\mathbb{T}^{\text{op}} \to \text{Ab}$ are precisely the filtered colimits of representable functors (in the category of additive functors $\mathbb{T}^{\text{op}} \to \text{Ab}$). In particular, the category $\text{Coh} \mathbb{T}$ has filtered colimits.

Exact functors. We say that a sequence of morphisms in $\operatorname{Coh} \mathcal{T}$ is *exact* provided that evaluation at each object in \mathcal{T} yields an exact sequence in Ab. This provides an exact structure in the sense of Quillen on the category $\operatorname{Coh} \mathcal{T}$.

Let \mathcal{T} and \mathcal{U} be essentially small triangulated categories. A functor $P: \operatorname{Coh} \mathcal{T} \to \operatorname{Coh} \mathcal{U}$ is said to be *exact* if it takes exact sequences to exact sequences and if there is a natural isomorphism $P \circ \Sigma \xrightarrow{\sim} \Sigma \circ P$.

An exact functor $f: \mathfrak{T} \to \mathfrak{U}$ induces a pair of functors

$$f^*\colon \operatorname{Coh} \mathfrak{T} \longrightarrow \operatorname{Coh} \mathfrak{U} \quad \text{and} \quad f_*\colon \operatorname{Coh} \mathfrak{U} \longrightarrow \operatorname{Coh} \mathfrak{T}$$

where $f^*(F) = \operatorname{colim}_{H_X \to F} H_{f(X)}$ and $f_*(G) = G \circ f$. We recall some basic facts.

Lemma 2.2 ([8, Lemma 2.4]). Let $f: \mathbb{T} \to \mathcal{U}$ be an exact functor between essentially small triangulated categories.

- (1) The functor f^* is a left adjoint of f_* .
- (2) The functors f^* and f_* are exact and preserve filtered colimits.
- (3) If f is fully faithful, then f^* is fully faithful and $id \xrightarrow{\sim} f_* \circ f^*$.

(4) If f is a quotient functor, then f_* is fully faithful and $f^* \circ f_* \xrightarrow{\sim} id$. \Box

Let $S \subseteq T$ be a triangulated subcategory. With $i: S \to T$ and $q: T \to T/S$ the canonical functors, set

$$\Gamma_{\mathbb{S}} = i^* \circ i_*$$
 and $L_{\mathbb{S}} = q_* \circ q^*$

These are exact functors on Coh \mathcal{T} . We may identify Coh \mathcal{S} and Coh \mathcal{T}/\mathcal{S} with full subcategories of Coh \mathcal{T} . Then we have for each $F \in \text{Coh }\mathcal{T}$

$$F \in \operatorname{Coh} S \iff \Gamma_S F \xrightarrow{\sim} F \iff L_S F = 0$$

and

$$F \in \operatorname{Coh} \mathfrak{T}/\mathfrak{S} \quad \Longleftrightarrow \quad F \xrightarrow{\sim} L_{\mathfrak{S}}F \quad \Longleftrightarrow \quad \Gamma_{\mathfrak{S}}F = 0.$$

The functors $\Gamma_{\mathbb{S}}$ and $L_{\mathbb{S}}$ fit into a long exact sequence. To explain this we note that for any $F \in \operatorname{Coh} \mathfrak{T}$ one has

$$\Gamma_{\mathfrak{S}}F = \operatorname{colim}_{H_{\mathfrak{S}} \to F} \operatorname{Hom}_{\mathfrak{T}}(-, S)$$

where the colimit is taken over the slice category S/F, which is filtered because S is a triangulated subcategory. On the other hand, the definition of the morphisms in the quotient category T/S gives for any $X \in T$

$$L_{\mathcal{S}}H_X = \operatorname{Hom}_{\mathcal{T}/\mathcal{S}}(-, X) = \operatorname{colim}_{X \to Y} \operatorname{Hom}_{\mathcal{T}}(-, Y)$$

where $X \to Y$ runs through all morphisms with cone in S. Combining this we obtain the following.

Proposition 2.3 ([8, Proposition 2.10]). For a triangulated subcategory $S \subseteq T$ there is the following long exact sequence of functors $\operatorname{Coh} T \to \operatorname{Coh} T$.

$$(2.1) \qquad \cdots \longrightarrow \Sigma^{n-1}L_{\mathbb{S}} \longrightarrow \Sigma^n \Gamma_{\mathbb{S}} \longrightarrow \Sigma^n \longrightarrow \Sigma^n L_{\mathbb{S}} \longrightarrow \Sigma^{n+1}\Gamma_{\mathbb{S}} \longrightarrow \cdots \qquad \Box$$

The sequence (2.1) is functorial with respect to S. Thus each inclusion $S_1 \subseteq S_2$ of triangulated subcategories of \mathcal{T} induces a morphism between the corresponding long exact sequences. An immediate consequence is the following.

Lemma 2.4 ([8, Lemma 2.16]). Let $S_1 \subseteq S_2 \subseteq T$ be triangulated subcategories and let (Γ_1, L_1) and (Γ_2, L_2) be the corresponding pairs of (co)localisation functors on Coh T. The morphisms in (2.1) induce isomorphisms

$$\Gamma_1\Gamma_2 \cong \Gamma_1 \cong \Gamma_2\Gamma_1, \quad L_1L_2 \cong L_2 \cong L_2L_1, \quad \Gamma_1L_2 = 0 = L_2\Gamma_1, \quad \Gamma_2L_1 \cong L_1\Gamma_2.$$

The rest of this paper makes excessive use of the category of functors

$$\operatorname{Coh} \mathfrak{T} \longrightarrow \operatorname{Coh} \mathfrak{T}$$

that are exact and perserve filtered colimits. We consider them up to isomorphism, which means that we identify functors F and G when there is a natural isomorphism $F \xrightarrow{\sim} G$. Let us denote by End \mathcal{T} the isomorphism classes of objects. Then we have the following relevant structures:

- (1) an associative multiplication End $T \times \text{End } T \to \text{End } T$ with an identity (given by the composition of functors),
- (2) an associative sum $\operatorname{End} \mathfrak{T} \times \operatorname{End} \mathfrak{T} \to \operatorname{End} \mathfrak{T}$ with an identity (given by the direct sum of functors),
- (3) a suspension Σ : End $\mathcal{T} \xrightarrow{\sim}$ End \mathcal{T} , and
- (4) a notion of exact sequence $F \to G \to H$ with connecting morphism $H \to \Sigma F$, providing a long exact sequence

$$\cdots \longrightarrow \Sigma^{n-1}H \longrightarrow \Sigma^n F \longrightarrow \Sigma^n G \longrightarrow \Sigma^n H \longrightarrow \Sigma^{n+1}F \longrightarrow \cdots$$

3. The lattice of central subcategories

Throughout we keep fixed an essentially small triangulated category \mathcal{T} . Let Thick \mathcal{T} denote the lattice of thick subcategories of \mathcal{T} . For \mathcal{U}, \mathcal{V} in Thick \mathcal{T} we write $\mathcal{U} \wedge \mathcal{V}$ for their meet (i.e. the intersection) and $\mathcal{U} \vee \mathcal{V}$ for their join (i.e the smallest thick subcategory containing \mathcal{U} and \mathcal{V}). For a class of objects $\mathcal{X} \subseteq \mathcal{T}$ let thick \mathcal{X} denote the smallest thick subcategory containing \mathcal{X} .

Central subcategories. Given a pair \mathcal{U}, \mathcal{V} of thick subcategories we consider the property that morphisms between objects from \mathcal{U} and \mathcal{V} factor through objects in $\mathcal{U} \wedge \mathcal{V}$. This condition appears already in [26, II.2.3], though the equivalent conditions (2)–(4) in the lemma below seem to be new.

Lemma 3.1. For a pair $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ the following are equivalent.

- (1) Every morphism $\mathcal{U} \ni U \to V \in \mathcal{V}$ factors through an object in $\mathcal{U} \land \mathcal{V}$.
- (2) The canonical morphism $\Gamma_{\mathcal{U}\wedge\mathcal{V}}\to\Gamma_{\mathcal{U}}\Gamma_{\mathcal{V}}$ is an isomorphism.
- (3) The canonical morphism $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \to L_{\mathcal{U}}\Gamma_{\mathcal{V}}$ is an isomorphism.
- (4) The canonical morphism $\Gamma_{\mathfrak{U}}L_{\mathfrak{U}\wedge \mathfrak{V}} \to \Gamma_{\mathfrak{U}}L_{\mathfrak{V}}$ is an isomorphism.
- (5) The canonical morphism $\operatorname{Hom}_{\mathcal{T}/(\mathfrak{U}\wedge \mathcal{V})}(-,V) \to \operatorname{Hom}_{\mathcal{T}/\mathfrak{U}}(-,V)$ is an isomorphism for all $V \in \mathcal{V}$.

Proof. When Coh \mathcal{U} is viewed as a full subcategory of Coh \mathcal{T} , then the objects in Coh \mathcal{U} are precisely the filtered colimits of objects H_U with $U \in \mathcal{U}$. From this it follows that $F \in \text{Coh } \mathcal{T}$ belongs to Coh \mathcal{U} if and only if every morphism $H_X \to F$ with $X \in \mathcal{T}$ factors through H_U for some $U \in \mathcal{U}$.

(1) \Leftrightarrow (2) The above general observation shows that the first two conditions are equivalent to the property that each object of the form $\Gamma_{\mathcal{U}}\Gamma_{\mathcal{V}}F$ belongs to Coh $\mathcal{U}\wedge\mathcal{V}$.

(2) \Leftrightarrow (3) Composition of $\Gamma_{\mathcal{V}}$ with the exact sequences (2.1) given by \mathcal{U} and $\mathcal{U} \wedge \mathcal{V}$ yield the following diagram.

The five lemma implies that α is an isomorphism if and only if β is an isomorphism.

(2) \Leftrightarrow (4) Composition of $\Gamma_{\mathcal{U}}$ with the exact sequences (2.1) given by \mathcal{V} and $\mathcal{U} \wedge \mathcal{V}$ yield the following diagram.

$$\cdots \to \Sigma^{-1} \Gamma_{\mathcal{U}} L_{\mathcal{U} \wedge \mathcal{V}} \to \Gamma_{\mathcal{U} \wedge \mathcal{V}} \to \Gamma_{\mathcal{U}} \to \Gamma_{\mathcal{U}} L_{\mathcal{U} \wedge \mathcal{V}} \to \Sigma \Gamma_{\mathcal{U} \wedge \mathcal{V}} \to \cdots$$

$$\downarrow^{\Sigma^{-1} \beta'} \qquad \downarrow^{\alpha} \qquad \downarrow^{\mathrm{id}} \qquad \downarrow^{\beta'} \qquad \downarrow^{\Sigma \alpha}$$

$$\cdots \to \Sigma^{-1} \Gamma_{\mathcal{U}} L_{\mathcal{V}} \to \Gamma_{\mathcal{U}} \Gamma_{\mathcal{V}} \to \Gamma_{\mathcal{U}} \to \Gamma_{\mathcal{U}} L_{\mathcal{V}} \to \Sigma \Gamma_{\mathcal{U} \wedge \mathcal{V}} \to \cdots$$

The five lemma implies that α is an isomorphism if and only if β' is an isomorphism. (3) \Leftrightarrow (5) For $V \in \mathcal{V}$ the inclusion $\mathcal{U} \wedge \mathcal{V} \subseteq \mathcal{U}$ induces the morphism

 $\operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,V)\cong L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}H_V\longrightarrow L_{\mathcal{U}}\Gamma_{\mathcal{V}}H_V\cong \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(-,V).$

As any functor of the form $\Gamma_{\mathcal{V}}F$ is a filtered colimit of representable functors H_V , this morphism is an isomorphism for all V if and only if $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \xrightarrow{\sim} L_{\mathcal{U}}\Gamma_{\mathcal{V}}$. \Box

We note that the equivalent conditions of the preceding lemma imply the *second Noether isomorphisms*, i.e. the canonical functors

$$\mathcal{V}/(\mathcal{U}\wedge\mathcal{V})\longrightarrow(\mathcal{U}\vee\mathcal{V})/\mathcal{U}$$
 and $\mathcal{U}/(\mathcal{U}\wedge\mathcal{V})\longrightarrow(\mathcal{U}\vee\mathcal{V})/\mathcal{V}$

are equivalences up to direct summands. This is clear for the first one and follows for the other by passing to the opposite category \mathcal{T}^{op} .

Definition 3.2. A pair of elements \mathcal{U}, \mathcal{V} in Thick \mathcal{T} is called *commuting* if we have canonical isomorphisms

$$\Gamma_{\mathfrak{U}}\Gamma_{\mathcal{V}} \xleftarrow{\sim} \Gamma_{\mathfrak{U}\wedge\mathcal{V}} \xrightarrow{\sim} \Gamma_{\mathcal{V}}\Gamma_{\mathfrak{U}}.$$

An equivalent condition is that a morphism between objects of \mathcal{U} and \mathcal{V} (in either direction) factors through an object of $\mathcal{U} \wedge \mathcal{V}$.

An element \mathcal{U} in Thick \mathcal{T} is called *central* if it commutes with all \mathcal{V} in Thick \mathcal{T} .

We denote by $Z(\text{Thick } \mathcal{T})$ the set of central thick subcategories of \mathcal{T} ; it is partially ordered by inclusion. In the following we study the basic properties of $Z(\text{Thick } \mathcal{T})$. In particular we will see that it is a sublattice of Thick \mathcal{T} .

We continue with a sequence of technical lemmas which establish further identities for exact functors $\operatorname{Coh} \mathcal{T} \to \operatorname{Coh} \mathcal{T}$ given by a pair of thick subcategories. The next one shows that commutativity is equivalent to an excision property.

Lemma 3.3. For a pair \mathcal{U}, \mathcal{V} in Thick \mathcal{T} the following are equivalent.

- (1) The pair \mathcal{U}, \mathcal{V} is commuting.
- (2) The canonical morphism

 $\psi\colon L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{U}\vee\mathcal{V}}\longrightarrow L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}\oplus L_{\mathcal{V}}\Gamma_{\mathcal{U}\vee\mathcal{V}}$

is an isomorphism.

(3) The canonical morphism

$$\psi_X \colon \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,X) \longrightarrow \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(-,X) \oplus \operatorname{Hom}_{\mathcal{T}/\mathcal{V}}(-,X)$$

is an isomorphism for all $X \in \mathcal{U} \lor \mathcal{V}$.

(4) The canonical morphism $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \to L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}$ is an isomorphism.

Proof. (1) \Rightarrow (2) The objects $X \in \mathcal{T}$ such that

$$L_{\mathcal{U}\wedge\mathcal{V}}H_X\longrightarrow L_{\mathcal{U}}H_X\oplus L_{\mathcal{V}}H_X$$

is an isomorphism form a thick subcategory $\mathfrak{X} \subseteq \mathfrak{T}$. From Lemma 3.1 it follows that $\mathfrak{U}, \mathfrak{V} \subseteq \mathfrak{X}$. Thus ψ is an isomorphism, as any functor of the form $\Gamma_{\mathfrak{U}\vee\mathfrak{V}}F$ is a filtered colimit of representable functors H_X with $X \in \mathfrak{U} \vee \mathfrak{V}$.

(2) \Rightarrow (3) We have $\psi H_X = \psi_X$ for each $X \in \mathcal{U} \vee \mathcal{V}$.

(3) \Rightarrow (1) For $X \in \mathcal{V}$ the morphism ψ_X specialises to $\operatorname{Hom}_{\mathcal{T}/(\mathcal{U} \wedge \mathcal{V})}(-, X) \rightarrow \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(-, X)$, and similar for $X \in \mathcal{U}$. Thus \mathcal{U} and \mathcal{V} are commuting, again by Lemma 3.1.

(1) \Rightarrow (4) The objects $X \in \mathcal{T}$ such that $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}H_X \rightarrow L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}H_X$ is an isomorphism form a thick subcategory $\mathcal{X} \subseteq \mathcal{T}$. From Lemma 3.1 it follows that $\mathcal{U}, \mathcal{V} \subseteq \mathcal{X}$. Thus the composite

$$L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}\xleftarrow{\sim} L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}\vee\mathcal{V}}\to L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}$$

is an isomorphism, as any functor of the form $\Gamma_{\mathcal{U}\vee\mathcal{V}}F$ is a filtered colimit of representable functors H_X with $X \in \mathcal{U}\vee\mathcal{V}$.

(4) \Rightarrow (1) Suppose $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \xrightarrow{\sim} L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}$. Composing with $\Gamma_{\mathcal{V}}$ yields $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \xrightarrow{\sim} L_{\mathcal{U}}\Gamma_{\mathcal{V}}$, and therefore $\Gamma_{\mathcal{U}\wedge\mathcal{V}} \xrightarrow{\sim} \Gamma_{\mathcal{U}}\Gamma_{\mathcal{V}}$ by Lemma 3.1. Composing with $\Gamma_{\mathcal{U}}$ yields $L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}} = 0$, and therefore

$$\Gamma_{\mathcal{U}\wedge\mathcal{V}} \xleftarrow{\sim} \Gamma_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}} \xrightarrow{\sim} \Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}}.$$

Thus the pair \mathcal{U}, \mathcal{V} is commuting.

Lemma 3.4. Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ be commuting. Then the following holds.

- (1) The canonical morphism $L_{\mathcal{U}}L_{\mathcal{V}} \to L_{\mathcal{U}\vee\mathcal{V}}$ is an isomorphism.
- (2) The canonical morphism $L_{\mathcal{U}}\Gamma_{\mathcal{V}} \to L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}$ is an isomorphism.
- (3) The canonical morphism $\Gamma_{\mathcal{U}}L_{\mathcal{V}} \to \Gamma_{\mathcal{U}\vee\mathcal{V}}L_{\mathcal{V}}$ is an isomorphism.

Proof. Consider the exact sequences (2.1) given by \mathcal{V} and $\mathcal{U} \lor \mathcal{V}$. We compose with $L_{\mathcal{U}}$ and obtain the following diagram.

Now consider the composite

$$L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}}\xrightarrow{\beta}L_{\mathcal{U}}\Gamma_{\mathcal{V}}\xrightarrow{\gamma}L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}}.$$

Then β is an isomorphism by Lemma 3.1, and $\gamma\beta$ is an isomorphisms by Lemma 3.3. Thus γ is an isomorphism, and it follows from the five lemma that δ is an isomorphism. A similar argument yields the last isomorphism.

Lemma 3.5. Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ be commuting. Then we have canonical isomorphisms

$$L_{\mathfrak{U}}L_{\mathcal{V}} \xrightarrow{\sim} L_{\mathfrak{U}\vee\mathcal{V}} \xleftarrow{\sim} L_{\mathcal{V}}L_{\mathfrak{U}}$$

Also, we have canonical isomorphisms

 $\Gamma_{\mathcal{V}}L_{\mathcal{U}} \xleftarrow{\sim} \Gamma_{\mathcal{V}}L_{\mathcal{U}\wedge\mathcal{V}} \cong L_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{V}} \xrightarrow{\sim} L_{\mathcal{U}}\Gamma_{\mathcal{V}}$

6

and

$$\Gamma_{\mathcal{V}}L_{\mathcal{U}} \xrightarrow{\sim} \Gamma_{\mathcal{U}\vee\mathcal{V}}L_{\mathcal{U}} \cong L_{\mathcal{U}}\Gamma_{\mathcal{U}\vee\mathcal{V}} \xleftarrow{\sim} L_{\mathcal{U}}\Gamma_{\mathcal{V}}.$$

Proof. The first and third sequence of isomorphisms follow from Lemma 3.4 and the second sequence from Lemma 3.1. $\hfill \Box$

The following lemma says that meet and join preserve central elements.

Lemma 3.6. Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ and suppose that \mathcal{U} is central.

- (1) The element $\mathcal{U} \wedge \mathcal{V}$ is central in Thick \mathcal{V} .
- (2) The element $(\mathcal{U} \vee \mathcal{V})/\mathcal{V}$ is central in Thick \mathcal{T}/\mathcal{V} .

Proof. (1) Let $\mathfrak{X} \in \text{Thick } \mathcal{V}$. Then any morphism between objects of $\mathfrak{U} \wedge \mathcal{V}$ and \mathfrak{X} (in either direction) factors through an object of $\mathfrak{U} \wedge \mathfrak{X} = (\mathfrak{U} \wedge \mathcal{V}) \wedge \mathfrak{X}$.

(2) Observe that for any $\mathfrak{X} \in \text{Thick } \mathfrak{T}$ containing \mathfrak{V} the functor $\Gamma_{\mathfrak{X}}$ restricts to a functor $\text{Coh } \mathfrak{T}/\mathfrak{V} \to \text{Coh } \mathfrak{T}/\mathfrak{V}$ where it identifies with $\Gamma_{\mathfrak{X}/\mathfrak{V}}$, since $\Gamma_{\mathfrak{X}}L_{\mathfrak{V}} \cong L_{\mathfrak{V}}\Gamma_{\mathfrak{X}}$. We obtain using Lemma 3.5

$$\Gamma_{\mathfrak{X}}\Gamma_{\mathfrak{U}\vee\mathfrak{V}}L_{\mathfrak{V}}\cong\Gamma_{\mathfrak{X}}\Gamma_{\mathfrak{U}}L_{\mathfrak{V}}\cong\Gamma_{\mathfrak{U}}\Gamma_{\mathfrak{X}}L_{\mathfrak{V}}\cong\Gamma_{\mathfrak{U}}L_{\mathfrak{V}}\Gamma_{\mathfrak{X}}\cong\Gamma_{\mathfrak{U}\vee\mathfrak{V}}L_{\mathfrak{V}}\Gamma_{\mathfrak{X}}\cong\Gamma_{\mathfrak{U}\vee\mathfrak{V}}\Gamma_{\mathfrak{X}}L_{\mathfrak{V}},$$

and therefore $\Gamma_{\mathcal{X}}\Gamma_{\mathcal{U}\vee\mathcal{V}}\cong\Gamma_{\mathcal{U}\vee\mathcal{V}}\Gamma_{\mathcal{X}}$ when restricted to $\operatorname{Coh} \mathfrak{T}/\mathcal{V}$.

Mayer–Vietoris sequences. A pair of commuting subcategories gives rise to a pair of long exact sequences. It turns out that commutativity is actually equivalent to having such Mayer–Vietoris sequences.

Proposition 3.7. Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ be commuting. Then there are canonical exact sequences

$$(\lambda_{\mathcal{U},\mathcal{V}}) \quad \cdots \longrightarrow \Sigma^{-1} L_{\mathcal{U}\vee\mathcal{V}} \longrightarrow L_{\mathcal{U}\wedge\mathcal{V}} \xrightarrow{(1)} L_{\mathcal{U}} \oplus L_{\mathcal{V}} \longrightarrow L_{\mathcal{U}\vee\mathcal{V}} \longrightarrow \Sigma L_{\mathcal{U}\wedge\mathcal{V}} \longrightarrow \cdots$$

$$(\gamma_{\mathfrak{U},\mathfrak{V}}) \quad \cdots \longrightarrow \Sigma^{-1} \varGamma_{\mathfrak{U} \wedge \mathfrak{V}} \longrightarrow \varGamma_{\mathfrak{U} \wedge \mathfrak{V}} \stackrel{(2)}{\longrightarrow} \varGamma_{\mathfrak{U}} \oplus \varGamma_{\mathfrak{V}} \longrightarrow \varGamma_{\mathfrak{U} \wedge \mathfrak{V}} \longrightarrow \Sigma \varGamma_{\mathfrak{U} \wedge \mathfrak{V}} \longrightarrow \cdots$$

where (1) and (2) are induced by the inclusions $\mathcal{U} \land \mathcal{V} \subseteq \mathcal{U}$ and $\mathcal{U} \land \mathcal{V} \subseteq \mathcal{V}$.

Proof. The exact sequence (2.1) given by \mathcal{U} and composed with $L_{\mathcal{U}\wedge\mathcal{V}}\to L_{\mathcal{V}}$ yields a morphisms of exact sequences.

Here we use the identification $L_{\mathcal{U}}L_{\mathcal{V}} = L_{\mathcal{U}\vee\mathcal{V}}$ from Lemma 3.4, and the isomorphisms follow from Lemma 3.1. Inverting them yields the connecting morphism $L_{\mathcal{U}\vee\mathcal{V}} \rightarrow \Sigma L_{\mathcal{U}\wedge\mathcal{V}}$, and then a standard argument turns the diagram into an exact sequence of the form $\lambda_{\mathcal{U},\mathcal{V}}$; cf. [24, Lemma 10.7.4]. The proof for the second sequence is analogous.

Corollary 3.8. For a pair \mathcal{U}, \mathcal{V} in Thick \mathcal{T} the following are equivalent.

- (1) The pair \mathcal{U}, \mathcal{V} is commuting.
- (2) There is an exact Mayer-Vietoris sequence $\lambda_{\mathcal{U},\mathcal{V}}$.
- (3) For each object $X \in \mathcal{T}$ there is a canonical exact sequence

 $\cdots \to \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,X) \to \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(-,X) \oplus \operatorname{Hom}_{\mathcal{T}/\mathcal{V}}(-,X) \to$ $\to \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,X) \to \operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,\Sigma X) \to \cdots .$

Proof. $(1) \Rightarrow (2)$ This follows from Proposition 3.7.

(2) \Rightarrow (3) For $X \in \mathcal{T}$ apply the Mayer–Vietoris sequence $\lambda_{\mathcal{U},\mathcal{V}}$ to H_X .

(3) \Rightarrow (1) For $X \in \mathcal{U} \vee \mathcal{V}$ the sequence $\lambda_{\mathcal{U},\mathcal{V}}H_X$ yields an isomorphism

$$\operatorname{Hom}_{\mathcal{T}/(\mathcal{U}\wedge\mathcal{V})}(-,X) \xrightarrow{\sim} \operatorname{Hom}_{\mathcal{T}/\mathcal{U}}(-,X) \oplus \operatorname{Hom}_{\mathcal{T}/\mathcal{V}}(-,X).$$

Thus the pair \mathcal{U}, \mathcal{V} is commuting by Lemma 3.3.

This result reflects the classical situation given by a pair of subspaces $A, B \subseteq X = A \cup B$ of a topological space X. In that case the Mayer–Vietoris sequence for a homology theory is equivalent to an excision property which is symmetric in A and B; cf. [24, 10.7]. The triangulated analogue of this is the isomorphism $L_{U \wedge V} \Gamma_V \xrightarrow{\sim} L_U \Gamma_{U \vee V}$ from Lemma 3.3. Related (though not equivalent) is the second Noether isomorphism $\mathcal{V}/(\mathcal{U} \wedge \mathcal{V}) \xrightarrow{\sim} (\mathcal{U} \wedge \mathcal{V})/\mathcal{U}$.

There is a plethora of Mayer–Vietoris sequences for triangulated categories in the literature [4, 7, 21, 23], including interesting applications. We recover some of them and note that all are based (implicitly) on a pair of commuting thick subcategories. A typical example is given by subcategories \mathcal{U}, \mathcal{V} satisfying $\operatorname{Hom}_{\mathcal{T}}(U, V) = 0$ and $\operatorname{Hom}_{\mathcal{T}}(V, U) = 0$ for all $U \in \mathcal{U}$ and $V \in \mathcal{V}$.

Distributivity. Recall that a lattice is *distributive* if for all elements x, y, z there is an equality

 $(x \wedge z) \lor (y \wedge z) = (x \lor y) \land z.$

An equivalent condition is that for all elements x, y, z there is an equality

 $(x \lor z) \land (y \lor z) = (x \land y) \lor z.$

The lattice Thick \mathcal{T} is rarely distributive. However, we get distributivity when we restrict to central elements in this lattice. The proof requires another technical lemma.

Lemma 3.9. Let $\mathcal{U}, \mathcal{V}, \mathcal{W} \in \text{Thick } \mathcal{T}$ and suppose $\mathcal{W} \subseteq \mathcal{U} \land \mathcal{V}$. If $\Gamma_{\mathcal{W}} \xrightarrow{\sim} \Gamma_{\mathcal{U}} \Gamma_{\mathcal{V}}$ then $\mathcal{W} = \mathcal{U} \land \mathcal{V}$.

Proof. Let $X \in \mathcal{U} \land \mathcal{V}$. Then $\Gamma_{\mathcal{W}} H_X \cong \Gamma_{\mathcal{U}} \Gamma_{\mathcal{V}} H_X \cong H_X$. Thus $H_X \in \operatorname{Coh} \mathcal{W}$ and therefore $X \in \mathcal{W}$.

Proposition 3.10. Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ be central. Then $\mathcal{U} \lor \mathcal{V}$ and $\mathcal{U} \land \mathcal{V}$ are central. Moreover, we have for all $\mathcal{W} \in \text{Thick } \mathcal{T}$

$$(\mathcal{U} \land \mathcal{W}) \lor (\mathcal{V} \land \mathcal{W}) = (\mathcal{U} \lor \mathcal{V}) \land \mathcal{W}$$

and

$$(\mathcal{U} \lor \mathcal{W}) \land (\mathcal{V} \lor \mathcal{W}) = (\mathcal{U} \land \mathcal{V}) \lor \mathcal{W}$$

Proof. It is clear that $\mathcal{U} \wedge \mathcal{V}$ is central since

$$\Gamma_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{W}}\cong\Gamma_{\mathcal{U}}\Gamma_{\mathcal{V}}\Gamma_{\mathcal{W}}\cong\Gamma_{\mathcal{W}}\Gamma_{\mathcal{U}}\Gamma_{\mathcal{V}}\cong\Gamma_{\mathcal{W}}\Gamma_{\mathcal{U}\wedge\mathcal{V}}.$$

Now we show that $\mathcal{U} \vee \mathcal{V}$ is central and combine this with the proof of the first identity. Consider the following Mayer–Vietoris sequences, where the middle one uses that $\mathcal{U} \wedge \mathcal{W}$ and $\mathcal{V} \wedge \mathcal{W}$ are commuting by Lemma 3.6.

$$\cdots \longrightarrow \Gamma_{\mathcal{U}\wedge\mathcal{V}}\Gamma_{\mathcal{W}} \longrightarrow \Gamma_{\mathcal{U}}\Gamma_{\mathcal{W}} \oplus \Gamma_{\mathcal{V}}\Gamma_{\mathcal{W}} \longrightarrow \Gamma_{\mathcal{U}\vee\mathcal{V}}\Gamma_{\mathcal{W}} \longrightarrow \cdots$$

$$\uparrow^{2} \qquad \uparrow^{2} \qquad \downarrow^{2} \qquad \downarrow^{$$

8

The vertical isomorphisms are clear since \mathcal{U} and \mathcal{V} central. The inclusion

$$(\mathcal{U} \land \mathcal{W}) \lor (\mathcal{V} \land \mathcal{W}) \subseteq (\mathcal{U} \lor \mathcal{V}) \land \mathcal{W}$$

is automatic. Thus we can apply Lemma 3.9, and then the five lemma yields the assertion for $\mathcal{U} \vee \mathcal{V}$. We conclude that the central elements form a sublattice of Thick \mathcal{T} which is distributive.

We extend Proposition 3.10 and establish the same assertions for arbitrary joins, using some standard arguments.

Proposition 3.11. Let $(\mathfrak{U}_i)_{i \in I}$ be a family of central elements in Thick \mathfrak{T} . Then $\bigvee_{i \in I} \mathfrak{U}_i$ is central and

$$\bigvee_{i\in I} (\mathfrak{U}_i \wedge \mathfrak{V}) = \big(\bigvee_{i\in I} \mathfrak{U}_i\big) \wedge \mathfrak{V}$$

for all $\mathcal{V} \in \operatorname{Thick} \mathfrak{T}$.

Proof. Set $\mathcal{U} = \bigvee_{i \in I} \mathcal{U}_i$. The element $\mathcal{U}_J = \bigvee_{i \in J} \mathcal{U}_i$ is central for all finite $J \subseteq I$ by Proposition 3.10. Now choose objects $U \in \mathcal{U}$ and $V \in \mathcal{V}$. Then $U \in \mathcal{U}_J$ for some finite $J \subseteq I$ and therefore any morphism between U and V factors through an object in $\mathcal{U}_J \land \mathcal{V} \subseteq \mathcal{U} \land \mathcal{V}$. Thus \mathcal{U} is central.

The inclusion $\bigvee_{i \in I} (\mathcal{U}_i \wedge \mathcal{V}) \subseteq \mathcal{U} \wedge \mathcal{V}$ is clear. For the other inclusion we use again Proposition 3.10. In fact any object $X \in \mathcal{U} \wedge \mathcal{V}$ belongs to $\mathcal{U}_J \wedge V = \bigvee_{i \in J} (\mathcal{U}_i \wedge \mathcal{V})$ for some finite $J \subseteq I$, and therefore X belongs to $\bigvee_{i \in I} (\mathcal{U}_i \wedge \mathcal{V})$. \Box

The frame of central subcategories. Recall that a lattice is a *frame* if it is complete (i.e. arbitrary joins exist) and the following *infinite distributivity* holds: for all elements $(x_i)_{i \in I}$ and y there is an equality

$$\bigvee_{i \in I} (x_i \wedge y) = \big(\bigvee_{i \in I} x_i\big) \wedge y.$$

A morphism of frames is a morphism of lattices that preserves finite meets and arbitrary joins.

To a frame F corresponds its space $\operatorname{Pt} F$ of points. A *point* is by definition a frame morphisms $p: F \to \{0, 1\}$ and the open sets are of the form

$$U(a) := \{ p \in \Pr F \mid p(a) = 1 \} \qquad (a \in F).$$

To a topological space X we can associate its frame of open subsets ΩX , and this yields an adjoint pair of functors

$$\mathbf{Frm} \xrightarrow{\mathrm{Pt}} \mathbf{Top}^{\mathrm{op}}.$$

A frame F is spatial if the assignment $a \mapsto U(a)$ gives an isomorphism $F \xrightarrow{\sim} \Omega \operatorname{Pt} F$. It is convenient to identify the points in $\operatorname{Pt} F$ with the *prime elements* of F via the assignment

$$\operatorname{Pt} F \ni p \longmapsto \bigvee_{p(a)=0} a \in F.$$

The central thick subcategories form a sublattice

$$Z(\operatorname{Thick} \mathfrak{T}) \subseteq \operatorname{Thick} \mathfrak{T}$$

by Proposition 3.10. This lattice is in fact a spatial frame.

Corollary 3.12. The central thick subcategories form a spatial frame, and therefore the lattice of central thick subcategories identifies with the lattice of open subsets of a sober topological space. *Proof.* The infinite distributivity follows from Proposition 3.11. An application of Zorn's lemma produces sufficiently many prime elements such that the frame is spatial; see the proof of Theorem 4.7 in [12]. \Box

The assignment $\mathfrak{T} \mapsto Z(\operatorname{Thick} \mathfrak{T})$ is functorial in the following sense.

Corollary 3.13. For any triangulated subcategory $S \subseteq T$ the canonical functors $S \to T$ and $T \to T/S$ induce frame morphisms

$$Z(\operatorname{Thick} \mathfrak{T}) \longrightarrow Z(\operatorname{Thick} \mathfrak{S}), \qquad \mathfrak{U} \mapsto \mathfrak{U} \wedge \mathfrak{S}$$

and

$$Z(\operatorname{Thick} \mathfrak{T}) \longrightarrow Z(\operatorname{Thick} \mathfrak{T}/\mathfrak{S}), \qquad \mathfrak{U} \mapsto (\mathfrak{U} \vee \mathfrak{S})/\mathfrak{S}$$

Proof. The maps are well defined by Lemma 3.6. They preserve finite meets and arbitrary joins by Propositions 3.10 and 3.11.

Central support. For an essentially small triangulated category \mathcal{T} let

$$\operatorname{Spc}_{\operatorname{cent}} \mathfrak{T} := \operatorname{Pt} Z(\operatorname{Thick} \mathfrak{T})$$

denote the space associated with the frame of central thick subcategories of \mathfrak{T} . For any object $X \in \mathfrak{T}$ its *central support* is the set of prime elements \mathfrak{P} in $Z(\text{Thick }\mathfrak{T})$ not containing X, so

(3.1)
$$\operatorname{supp}_{\operatorname{cent}} X := \{ \mathcal{P} \in \operatorname{Spc}_{\operatorname{cent}} \mathcal{T} \mid X \notin \mathcal{P} \}.$$

For a class of objects $\mathfrak{X} \subseteq \mathfrak{T}$ we set

$$\operatorname{supp}_{\operatorname{cent}} \mathfrak{X} := \bigcup_{X \in \mathfrak{X}} \operatorname{supp}_{\operatorname{cent}} X,$$

which equals the set of primes \mathcal{P} such that $\mathcal{X} \not\subseteq \mathcal{P}$.

- **Corollary 3.14.** (1) The assignment $\mathcal{U} \mapsto \operatorname{supp}_{\operatorname{cent}} \mathcal{U}$ induces an isomorphism between $Z(\operatorname{Thick} \mathcal{T})$ and the frame of open subsets of $\operatorname{Spc}_{\operatorname{cent}} \mathcal{T}$.
 - (2) For any pair \mathcal{U}, \mathcal{V} in Thick \mathcal{T} there is a Mayer-Vietoris sequence $\lambda_{\mathcal{U},\mathcal{V}}$ provided at least one of \mathcal{U} and \mathcal{V} is central.

Proof. This is a reformulation of Corollaries 3.8 and 3.12.

4. Examples

Having introduced for any triangulated category the lattice of central thick subcategories, it is now time to look at examples.

Tensor triangulated categories. Let $\mathcal{T} = (\mathcal{T}, \otimes, \mathbb{1})$ be a tensor triangulated category, i.e. a triangulated category equipped with a symmetric monoidal structure which is exact in each variable. Then Coh \mathcal{T} inherits a symmetric monoidal structure which is exact and preserves filtered colimit in each variable by setting $H_X \otimes H_Y = H_{X \otimes Y}$ for all X, Y in \mathcal{T} . The category \mathcal{T} is called *rigid* if there is an exact functor $D: \mathcal{T}^{\text{op}} \to \mathcal{T}$ and a natural isomorphism

$$\operatorname{Hom}_{\mathfrak{T}}(X \otimes Y, Z) \cong \operatorname{Hom}_{\mathfrak{T}}(Y, DX \otimes Z)$$

for all $X, Y, Z \in \mathcal{T}$. Note that in this case $X \xrightarrow{\sim} D^2 X$ for all $X \in \mathcal{T}$.

Lemma 4.1. Let \mathcal{T} be a rigid tensor triangulated category and $\mathcal{U} \subseteq \mathcal{T}$ a thick tensor ideal. Then Coh \mathcal{U} and Coh \mathcal{T}/\mathcal{U} are tensor ideals of Coh \mathcal{T} .

Proof. We begin with Coh \mathcal{U} . Let $E, F \in \text{Coh }\mathcal{T}$ written as filtered colimits $E = \text{colim}_i H_{X_i}$ and $F = \text{colim}_j H_{X_j}$. If $F \in \text{Coh }\mathcal{U}$, then we may assume $X_j \in \mathcal{U}$ for all j. Thus $H_{X_i} \otimes F \cong \text{colim}_j H_{X_i \otimes X_j}$ belongs to Coh \mathcal{U} for all i because \mathcal{U} is a tensor ideal, and therefore $E \otimes F \cong \text{colim}_i (H_{X_i} \otimes F)$ is in Coh \mathcal{U} .

Now consider $\operatorname{Coh} \mathcal{T}/\mathcal{U}$, where rigidity is used as follows. Let $F \in \operatorname{Coh} \mathcal{T}/\mathcal{U}$ and $U, X \in \mathcal{T}$. Then we have

$$(H_X \otimes F)(U) \cong \operatorname{Hom}(H_U, H_X \otimes F) \cong \operatorname{Hom}(H_{DX \otimes U}, F) \cong F(DX \otimes U) = 0$$

when $U \in \mathcal{U}$. For an arbitrary $E = \operatorname{colim}_i H_{X_i}$ in $\operatorname{Coh} \mathcal{T}$ we conclude that $E \otimes F$ lies in $\operatorname{Coh} \mathcal{T}/\mathcal{U}$, since

$$(E \otimes F)(U) \cong \operatorname{colim}_{i}(H_{X_i} \otimes F)(U) = 0 \quad \text{for} \quad U \in \mathfrak{U}.$$

Proposition 4.2. Let T be a rigid tensor triangulated category. Then any pair of thick tensor ideals is commuting.

Proof. We provide two proofs; the second one is due to Kent Vashaw [25] and does not use the commutativity of the tensor product.

Let $\mathcal{U}, \mathcal{V} \in \text{Thick } \mathcal{T}$ be tensor ideals. From Lemma 4.1 we know that $\text{Coh } \mathcal{U}$ and $\text{Coh } \mathcal{T}/\mathcal{U}$ are tensor ideals of $\text{Coh } \mathcal{T}$. From this we deduce isomorphisms

$$\Gamma_{\mathfrak{U}} \cong (\Gamma_{\mathfrak{U}}H_{\mathfrak{1}}) \otimes -$$
 and $L_{\mathfrak{U}} \cong (L_{\mathfrak{U}}H_{\mathfrak{1}}) \otimes -$

as follows. Consider for $F \in \operatorname{Coh} \mathcal{T}$ the exact sequence

$$\cdots \longrightarrow (\Gamma_{\mathfrak{U}}H_{\mathbb{1}}) \otimes F \longrightarrow F \longrightarrow (L_{\mathfrak{U}}H_{\mathbb{1}}) \otimes F \longrightarrow \cdots$$

and apply $\Gamma_{\mathcal{U}}$. We have

$$\Gamma_{\mathfrak{U}}((\Gamma_{\mathfrak{U}}H_1)\otimes F)\cong (\Gamma_{\mathfrak{U}}H_1)\otimes F$$
 and $\Gamma_{\mathfrak{U}}((L_{\mathfrak{U}}H_1)\otimes F)=0$

because

$$(\Gamma_{\mathcal{U}}H_1) \otimes F \in \operatorname{Coh}\mathcal{U}$$
 and $(L_{\mathcal{U}}H_1) \otimes F \in \operatorname{Coh}\mathcal{T}/\mathcal{U}$

by the first part of this proof. Thus $(\Gamma_{\mathfrak{U}}H_1) \otimes F \xrightarrow{\sim} \Gamma_{\mathfrak{U}}F$, and the proof of the isomorphism $(L_{\mathfrak{U}}H_1) \otimes F \xrightarrow{\sim} L_{\mathfrak{U}}F$ is analogous. Now the commutativity $\Gamma_{\mathfrak{U}}\Gamma_{\mathcal{V}} \cong \Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}}$ follows.

The second proof is more direct. Given $U \in \mathcal{U}$ and $V \in \mathcal{V}$, any morphism $U \to V$ admits a factorisation

$$U \to U \otimes DU \otimes U \to V \otimes DU \otimes U \to V$$

with $V \otimes DU \otimes U$ in $\mathcal{U} \wedge \mathcal{V}$. By symmetry the same holds for a morphism $V \to U$. Thus the pair \mathcal{U}, \mathcal{V} is commuting.

Remark 4.3. (1) In Example 4.13 one finds a pair of thick tensor ideals which is not commuting.

(2) The second proof shows that the proposition remains true when the tensor product is not commutative [25].

(3) For a predecessor of the proposition, see Proposition 6.5 in [10].

The following immediate consequence explains why in several interesting examples all thick subcategories are central. This includes the category of perfect complexes over a commutative ring or the stable homotopy category of finite spectra.

Corollary 4.4. If a tensor triangulated category is generated by its tensor identity then all thick subcategories are central.

Proof. The assumption means $\mathcal{T} = \text{thick } \mathbb{1}$ and this implies that all thick subcategories are tensor ideals. In particular, $\text{Coh } \mathcal{U}$ and $\text{Coh } \mathcal{T}/\mathcal{U}$ are tensor ideals of $\text{Coh } \mathcal{T}$ for all thick $\mathcal{U} \subseteq \mathcal{T}$. Now apply the argument from Proposition 4.2.

Remark 4.5. If one assumes rigidity the corollary remains true when the tensor product is not commutative, using the second proof of Proposition 4.2.

Central actions. Let R be a graded commutative ring. We consider only graded R-modules and Spec R denotes the set of homogeneous prime ideals of R with the Zariski topology. We fix a triangulated category \mathfrak{T} with an R-linear action via a homomorphism of graded rings $R \to Z^*(\mathfrak{T})$ into the graded centre of \mathfrak{T} ; cf. [7, 11]. This means that for all objects X, Y in \mathfrak{T}

$$\operatorname{Hom}_{\mathcal{T}}^*(X,Y) := \bigoplus_{n \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{T}}(X,\Sigma^n Y)$$

is a graded R-module. Also, for $F \in \operatorname{Coh} \mathfrak{T}$ and $X \in \mathfrak{T}$ the graded abelian group

$$F^*(X) := \bigoplus_{n \in \mathbb{Z}} F(\Sigma^{-n}X)$$

is an *R*-module.

Let Φ be a multiplicatively closed set of homogeneous elements in R. For an R-module M set

$$M[\Phi^{-1}] := M \otimes_R R[\Phi^{-1}]$$

and note that $M[\Phi^{-1}] = 0$ if and only if $M_{\mathfrak{p}} = 0$ for all $\mathfrak{p} \in \operatorname{Spec} R$ such that $\mathfrak{p} \cap \Phi = \emptyset$. We define a triangulated category $\mathfrak{T}[\Phi^{-1}]$ as follows. The objects are the same as in \mathfrak{T} . For the morphisms set

$$\operatorname{Hom}_{\mathfrak{T}\left[\Phi^{-1}\right]}^{*}(X,Y) := \operatorname{Hom}_{\mathfrak{T}}^{*}(X,Y)[\Phi^{-1}]$$

The exact triangles in $\mathcal{T}[\Phi^{-1}]$ are given by the ones isomorphic to images of exact triangles in \mathcal{T} under the canonical functor $\mathcal{T} \to \mathcal{T}[\Phi^{-1}]$. The kernel of this functor is the thick subcategory

$$\mathfrak{T}_{\Phi} := \{ X \in \mathfrak{T} \mid H_X^*[\Phi^{-1}] = 0 \}$$

and from [8, Lemma 3.5] we have a triangle equivalence

(4.1)
$$\mathfrak{T}/\mathfrak{T}_{\Phi} \xrightarrow{\sim} \mathfrak{T}[\Phi^{-1}].$$

A special case of the above construction arises for each $\mathfrak{p} \in \operatorname{Spec} R$. We denote by $\mathfrak{T}_{\mathfrak{p}}$ the triangulated category given by localising the morphisms at \mathfrak{p} , so

$$\operatorname{Hom}_{\mathcal{T}_{\mathfrak{n}}}^{*}(X,Y) := \operatorname{Hom}_{\mathcal{T}}^{*}(X,Y)_{\mathfrak{p}}$$

The notation $X \mapsto X_{\mathfrak{p}}$ is used for the canonical functor $\mathfrak{T} \to \mathfrak{T}_{\mathfrak{p}}$.

Lemma 4.6. A thick subcategory of the form \mathbb{T}_{Φ} is central.


Proof. To simplify notation we set $\Gamma_{\Phi} := \Gamma_{\mathcal{T}_{\Phi}}$ and $L_{\Phi} := L_{\mathcal{T}_{\Phi}}$. Note that

$$(L_{\Phi}F)^* \cong F^*[\Phi^{-1}]$$

for each F in Coh T by (4.1). Now fix a thick subcategory $\mathcal{U} \subseteq \mathcal{T}$. We have canonical isomorphisms

$$L_{\Phi}\Gamma_{\mathfrak{U}} \xleftarrow{\sim} \Gamma_{\mathfrak{U}}L_{\Phi}\Gamma_{\mathfrak{U}} \xrightarrow{\sim} \Gamma_{\mathfrak{U}}L_{\Phi}$$

since L_{Φ} restricts to a functor $\operatorname{Coh} \mathcal{U} \to \operatorname{Coh} \mathcal{U}$. It remains to consider the following commutative diagram with exact rows.



Then the five lemma yields the isomorphism $\Gamma_{\Phi}\Gamma_{\mathcal{U}} \cong \Gamma_{\mathcal{U}}\Gamma_{\Phi}$. Thus \mathfrak{T}_{Φ} is central. \Box

For an ideal \mathfrak{a} of R that is generated by finitely many homogeneous elements set

 $V(\mathfrak{a}) := \{\mathfrak{p} \in \operatorname{Spec} R \mid \mathfrak{a} \subseteq \mathfrak{p}\}$

and consider the thick subcategory

$$\mathfrak{I}_{V(\mathfrak{a})} := \{ X \in \mathfrak{T} \mid X_{\mathfrak{p}} = 0 \text{ for } \mathfrak{p} \in (\operatorname{Spec} R) \setminus V(\mathfrak{a}) \}.$$

Lemma 4.7. A thick subcategory of the form $\mathcal{T}_{V(\mathfrak{a})}$ is central.

Proof. Choose homogeneous generators r_1, \ldots, r_n of \mathfrak{a} and set $\Phi_i = \{r_i^p \mid p \ge 1\}$ for $1 \le i \le n$. Then $\mathcal{T}_{V(r_i)} = \mathcal{T}_{\Phi_i}$ is central for each *i* by Lemma 4.6. We have $V(\mathfrak{a}) = V(r_1) \cap \cdots \cap V(r_n)$ and therefore

$$\mathcal{T}_{V(\mathfrak{a})} = \mathcal{T}_{V(r_1)} \wedge \dots \wedge \mathcal{T}_{V(r_n)}$$

is central by Proposition 3.10.

The sets $V(\mathfrak{a})$ given by finitely generated ideals form a distributive lattice with

$$V(\mathfrak{a}) \cap V(\mathfrak{b}) = V(\mathfrak{a} + \mathfrak{b})$$
 and $V(\mathfrak{a}) \cup V(\mathfrak{b}) = V(\mathfrak{a}\mathfrak{b}).$

Lemma 4.8. For $U = V(\mathfrak{a})$ and $V = V(\mathfrak{b})$ we have

$$\mathfrak{T}_{U\cap V} = \mathfrak{T}_U \land \mathfrak{T}_V \qquad and \qquad \mathfrak{T}_{U\cup V} = \mathfrak{T}_U \lor \mathfrak{T}_V.$$

Proof. The first equality is clear. For the second equality choose homogeneous generators r_1, \ldots, r_m of \mathfrak{a} and s_1, \ldots, s_n of \mathfrak{b} such that

$$V(\mathfrak{a}) = V(r_1) \cap \cdots \cap V(r_m)$$
 and $V(\mathfrak{b}) = V(s_1) \cap \cdots \cap V(s_n)$.

For a homogeneous element $r \in R$ of degree |r| and an object $X \in \mathcal{T}$ we denote by $X/\!\!/r$ the cone of $X \xrightarrow{r} \Sigma^{|r|} X$ and have

$$\mathcal{T}_{V(r)} = \operatorname{thick}(\{X / \!\!/ r \mid X \in \mathcal{T}\})$$

by [8, Proposition 3.10]. For homogeneous elements $r, s \in R$ an application of the octahedral axiom shows that $X/\!\!/rs$ is an extension of $X/\!\!/r$ and $\Sigma^{|r|}X/\!\!/s$. Thus

$$\mathfrak{T}_{V(r)\cup V(s)} = \mathfrak{T}_{V(rs)} = \mathfrak{T}_{V(r)} \vee \mathfrak{T}_{V(s)}.$$

Using the distributivity in $Z(\text{Thick } \mathcal{T})$ from Proposition 3.10 we obtain

$$\mathfrak{I}_{V(\mathfrak{a})} \vee \mathfrak{I}_{V(\mathfrak{b})} = \bigwedge_{i,j} (\mathfrak{I}_{V(r_i)} \vee \mathfrak{I}_{V(s_j)}) = \bigwedge_{i,j} \mathfrak{I}_{V(r_i s_j)} = \mathfrak{I}_{V(\mathfrak{a}\mathfrak{b})} = \mathfrak{I}_{V(\mathfrak{a}) \cup V(\mathfrak{b})}$$

since

$$V(\mathfrak{ab}) = \bigcap_{i,j} V(r_i s_j).$$

Recall that a set $V \subseteq \operatorname{Spec} R$ has a Zariski open and quasi-compact complement if and only if $V = V(\mathfrak{a})$ for an ideal \mathfrak{a} of R that is generated by finitely many homogeneous elements. A subset $V \subseteq \operatorname{Spec} R$ is called *Thomason* if it can be written as $V = \bigcup_i V_i$ such that each $(\operatorname{Spec} R) \setminus V_i$ is Zariski open and quasi-compact. The Thomason subsets are precisely the open subsets for the *Hochster dual topology* on $\operatorname{Spec} R$ [14]. We write $\operatorname{Spec}^{\vee} R$ to denote the spectrum of prime ideals with this dual topology.

Given a frame F, one calls $x \in F$ finite or compact if $x \leq \bigvee_{i \in I} y_i$ implies $x \leq \bigvee_{i \in J} y_i$ for some finite subset $J \subseteq I$. A frame is coherent when every element can be written as a join of finite elements and the finite elements form a sublattice. In this case we write F^c for the sublattice of finite elements and observe that for any frame F' a lattice morphism $F^c \to F'$ extends uniquely to a frame morphism $F \to F'$.

The following proposition identifies the thick subcategories of \mathcal{T} which are given by a central ring action; they are central elements in Thick \mathcal{T} .

Proposition 4.9. The assignment

$$V\longmapsto \mathfrak{T}_V:=\bigvee_{V(\mathfrak{a})\subset V}\mathfrak{T}_{V(\mathfrak{a})}$$

induces a frame morphism from the open subsets of $\operatorname{Spec}^{\vee} R$ to the central thick subcategories of \mathfrak{T} .

Proof. The open sets of $\operatorname{Spec}^{\vee} R$ form a coherent frame and the finite elements are precisely the sets of the form $V(\mathfrak{a})$. Then the assertion follows from the preceding discussion, once we observe that a thick subcategory of the form $\mathcal{T}_{V(\mathfrak{a})}$ is central by Lemma 4.7 and that the map $V(\mathfrak{a}) \mapsto \mathcal{T}_{V(\mathfrak{a})}$ is a lattice morphism by Lemma 4.8. \Box

It would be interesting to know when for a given triangulated category all central subcategories are given by a central ring action. In algebraic examples this seems to be more likely, while it seems not to be true for the stable homotopy category of finite spectra.

Noetherian cohomology. Let R be a graded commutative ring and \mathcal{T} a triangulated category with an R-linear action via a homomorphism of graded rings $R \to Z^*(\mathcal{T})$. We assume that R is noetherian and that $\operatorname{Hom}^*_{\mathcal{T}}(X,Y)$ is a noetherian R-module for all objects X, Y in \mathcal{T} .

The *cohomological support* of an object X is by definition

$$\operatorname{supp}_R X := \{ \mathfrak{p} \in \operatorname{Spec} R \mid X_{\mathfrak{p}} \neq 0 \}.$$

Lemma 4.10. We have $\operatorname{supp}_R X = V(\mathfrak{a})$ where \mathfrak{a} denotes the kernel of the ring homomorphism $R \to \operatorname{End}^*_{\mathfrak{T}}(X)$.

Proof. Observe that $X_{\mathfrak{p}} = 0$ if and only $(H_X^*)_{\mathfrak{p}} = 0$. Then the equality follows from the fact that for each $Y \in \mathcal{T}$ the *R*-action on $H_X^*(Y) = \operatorname{Hom}_{\mathcal{T}}^*(Y, X)$ factors through $\operatorname{End}_{\mathcal{T}}^*(X)$, and that for any finitely generated *R*-module *M*

$$\operatorname{supp}_{R} M := \{ \mathfrak{p} \in \operatorname{Spec} R \mid M_{\mathfrak{p}} \neq 0 \} = V(\operatorname{Ann}_{R} M). \qquad \Box$$

Lemma 4.11. For an object $X \in \mathfrak{T}$ and a Thomason set $V \subseteq \operatorname{Spec} R$ we have

$$\operatorname{supp}_{R} X \subseteq V \iff X \in \mathfrak{T}_{V}.$$

Proof. From the preceding lemma we have an ideal \mathfrak{a} such that $\operatorname{supp}_R X = V(\mathfrak{a})$. Then $\operatorname{supp}_R X \subseteq V$ implies $X \in \mathfrak{T}_{V(\mathfrak{a})} \subseteq \mathfrak{T}_{\mathcal{V}}$. On the other hand, $X \in \mathfrak{T}_V$ implies $X \in \mathfrak{T}_{V(\mathfrak{b})}$ for some $V(\mathfrak{b}) \subseteq V$, and therefore $\operatorname{supp}_R X \subseteq V(\mathfrak{b}) \subseteq V$. \Box

Hereditary algebras. For a finite dimensional hereditary algebra we consider the category of perfect complexes. Thick subcategories that are generated by exceptional sequences have been classified in this case via non-crossing partitions [15, 16]. The following lemma shows that they are usually not central.

Lemma 4.12. Let $\mathcal{U} \subseteq \mathcal{T}$ be a thick subcategory such that the inclusion admits a left and a right adjoint. If \mathcal{U} is central, then ${}^{\perp}\mathcal{U} = \mathcal{U}^{\perp}$.

Proof. We have

$$\operatorname{Ker} \Gamma_{\perp_{\mathcal{U}}} = \operatorname{Coh} \mathcal{U} \quad \text{and} \quad \operatorname{Ker} \Gamma_{\mathcal{U}} = \operatorname{Coh} \mathcal{U}^{\perp}.$$

Thus $\Gamma_{\perp \mathcal{U}} \Gamma_{\mathcal{U}} = 0$, while $\Gamma_{\mathcal{U}} \Gamma_{\perp \mathcal{U}} \neq 0$ when $^{\perp}\mathcal{U} \not\subseteq \mathcal{U}^{\perp}$. On the other hand, we have Ker $L_{\mathcal{U}} = \operatorname{Coh} \mathcal{U}$. Thus $L_{\mathcal{U}} L_{\perp \mathcal{U}} = 0$, and $L_{\mathcal{U}} L_{\perp \mathcal{U}} \neq 0$ when $^{\perp}\mathcal{U} \not\supseteq \mathcal{U}^{\perp}$. \Box

Example 4.13. We consider the path algebra $A = k(\circ \rightarrow \circ)$ over a field k and the bounded derived category $\mathcal{T} = \mathbf{D}^{b} \pmod{A}$ of finitely generated A-modules, which identifies with the category of perfect complexes over A. The Hasse diagram of the lattice of thick subcategories is the following.



For any proper thick subcategory $\mathcal{U} \subseteq \mathcal{T}$ we have

Thick
$$\mathcal{T} = \{0, \mathcal{U}, {}^{\perp}\mathcal{U}, \mathcal{U}^{\perp}, \mathcal{T}\}.$$

In particular, $Z(\text{Thick } \mathfrak{T}) = \{0, \mathfrak{T}\}$ by Lemma 4.12.

The category of A-modules admits a symmetric monoidal structure given by the pointwise tensor product over k. All thick subcategories are tensor ideals, except the one generated by the tensor unit. This provides a pair of thick tensor ideals \mathcal{U}, \mathcal{V} such that $\Gamma_{\mathcal{U}} \Gamma_{\mathcal{V}} \ncong \Gamma_{\mathcal{V}} \Gamma_{\mathcal{U}}$, because we have non-zero morphisms between \mathcal{U} and \mathcal{V} in one direction but $\mathcal{U} \wedge \mathcal{V} = 0$.

The above example can be generalised as follows.

Example 4.14. Let R be a commutative ring and consider the category of perfect complexes $\mathcal{T} = \operatorname{Perf} T_2(R)$ over the matrix ring $T_2(R) = \begin{bmatrix} R & R \\ 0 & R \end{bmatrix}$. This category admits an obvious R-action which provides an embedding

Thick Perf $R \hookrightarrow Z(\operatorname{Thick} \mathfrak{T}) \subsetneq \operatorname{Thick} \mathfrak{T}$.

Example 4.15. We consider the category of coherent sheaves on the projective line \mathbb{P}^1 over a fixed field k and its derived category $\mathcal{T} = \mathbf{D}^b(\operatorname{coh} \mathbb{P}^1)$; it is equivalent to the category of perfect complexes over the path algebra $k(\circ \rightrightarrows \circ)$.

We recall from [18] the description of Thick $\mathbb T.$ There is a lattice isomorphism

{thick subcategories of $\mathbf{D}^{b}(\operatorname{coh} \mathbb{P}^{1})$ generated by vector bundles} $\xrightarrow{\sim} \mathbb{Z}$

given by thick $\mathcal{O}(n) \mapsto n$, where \mathbb{Z} denotes the lattice given by the following Hasse diagram:



We have

 $(\operatorname{thick} \mathcal{O}(n))^{\perp} = \operatorname{thick} \mathcal{O}(n-1) \text{ and } ^{\perp}(\operatorname{thick} \mathcal{O}(n)) = \operatorname{thick} \mathcal{O}(n+1).$

Thus a subcategory of the form thick O(n) is not central by Lemma 4.12. On the other hand, we have a lattice isomorphism

{thick tensor ideals of $\mathbf{D}^b(\operatorname{coh} \mathbb{P}^1)$ } $\xrightarrow{\sim}$ {Thomason subsets of \mathbb{P}^1 }

given by $\mathcal{U} \mapsto \bigcup_{X \in \mathcal{U}} \operatorname{supp} X$. This yields an isomorphism²

Thick $\mathbf{D}^b(\operatorname{coh} \mathbb{P}^1) \xrightarrow{\sim} \{ \text{Thomason subsets of } \mathbb{P}^1 \} \amalg \mathbb{Z}.$

Now let $\mathcal{U} \subseteq \mathcal{T}$ be a proper thick tensor ideal and $\mathcal{V} = \text{thick } \mathcal{O}(n)$. Then $\mathcal{U} \land \mathcal{V} = 0$ but $\Gamma_{\mathcal{V}}\Gamma_{\mathcal{U}} \neq 0$, because we have $\text{Hom}(\mathcal{O}(n), X) \neq 0$ for any non-zero torsion sheaf X in \mathcal{U} . Thus \mathcal{U} is not central. We conclude that $Z(\text{Thick }\mathcal{T}) = \{0, \mathcal{T}\}.$

²Let L', L'' be a pair of lattices with smallest elements 0', 0'' and greatest elements 1', 1''. Then $L' \amalg L''$ denotes the new lattice which is obtained from the disjoint union $L' \cup L''$ (viewed as sum of posets) by identifying 0' = 0'' and 1' = 1''.

Group algebras. Let G be a finite group and k a field. We consider the stable category $\mathcal{T} = \operatorname{stmod} kG$ of finitely generated modules over the group algebra kG. This is a rigid tensor triangulated category and the tensor ideals are precisely the thick subcategories given by the central action of the group cohomology [6].

Example 4.16 (Benson). In general there are central thick subcategories which are not tensor ideal. Let $B_0 = e(kG)$ denote the principal block of kG which is given by a central idempotent $e \in kG$. We write R := Z(kG) for the centre and set $V := V((1-e)R) \subseteq \text{Spec } R$. Then stmod $B_0 = \mathcal{T}_V$ is central, and this is a tensor ideal if and only if $B_0 = kG$. For example, we have thick $k = \text{stmod } B_0 \neq \mathcal{T}$ for $G = A_7$ and char k = 2.

Example 4.17 (Benson). Let $G = \mathbb{Z}/3 \times S_3$ and char k = 3. Then thick k is not central. There are two simple modules, denoted by k and ε (for sign representation). There is a uniserial module $X = [k, \varepsilon, k]$ supported in the nucleus [5] and a non-zero morphism $k \to X$. However, the intersection of the thick subcategories thick k and thick X is zero.

It seems interesting to find out more about $Z(\operatorname{stmod} kG)$ and its associated space. When $Z(\operatorname{stmod} kG)$ equals the lattice of thick tensor ideals, then the corresponding space is given by the projective variety of the group cohomology ring $H^*(G, k)$, but the general case requires a bigger space and is not clear. For instance, are all central subcategories given by the central action of a graded commutative ring?

5. The relative perspective

Let \mathcal{T} be an essentially small triangulated category. The examples suggest to get back to our Definition 3.2, introducing the following relative version of the lattice of central thick subcategories.

Definition 5.1. Let $T \subseteq \text{Thick } \mathcal{T}$ be a sublattice which is closed under finite meets and arbitrary joins, containing in particular 0 and \mathcal{T} . Then $\mathcal{U} \in T$ is called *central relative to* T if \mathcal{U} commutes with all \mathcal{V} in T. An equivalent condition is that a morphism between objects of \mathcal{U} and any $\mathcal{V} \in T$ (in either direction) factors through an object of $\mathcal{U} \wedge \mathcal{V}$. We set

 $Z(T) := \{ \mathcal{U} \in T \mid \mathcal{U} \text{ is central relative to } T \}.$

Theorem 5.2. Let $T \subseteq \text{Thick } \mathcal{T}$ be a sublattice which is closed under arbitrary joins. Then its centre $Z(T) \subseteq T$ is also a sublattice and closed under arbitrary joins. Moreover, Z(T) is a spatial frame and for any pair \mathcal{U}, \mathcal{V} in T the Mayer-Vietoris sequence $\lambda_{\mathcal{U},\mathcal{V}}$ exists provided at least one of \mathcal{U} and \mathcal{V} is central.

Proof. We have already seen the proof for $T = \text{Thick } \mathcal{T}$. The arguments given in Propositions 3.10 and 3.11 as well as in Corollary 3.12 carry over without change. For the Mayer–Vietoris sequence, see Corollary 3.8.

A consequence is that T is a distributive lattice when Z(T) = T. The converse is not necessarily true; cf. Example 4.13.

Example 5.3. Let \mathcal{T} be a rigid tensor triangulated category (not necessarily commutative) and $T \subseteq \text{Thick } \mathcal{T}$ the sublattice of all thick tensor ideals. Then Z(T) = T by Proposition 4.2. In particular, the space associated with this frame

$$\operatorname{Spc}_{\otimes} \mathfrak{T} := \operatorname{Pt} T$$

coincides (up to Hochster duality) with Balmer's space $\operatorname{Spc} \mathcal{T} [1, 17]$ and its non-commutative analogue [12, 19].

Example 5.4. Consider the graded centre $Z^*(\mathcal{T})$ of \mathcal{T} and set

$$T := \{ \mathfrak{T}_V \mid V \subseteq \operatorname{Spec} Z^*(\mathfrak{T}) \text{ is Thomason} \}.$$

Then $T \subseteq Z(\text{Thick } \mathcal{T})$ is a frame by Proposition 4.9 and in particular Z(T) = T. One may call the associated space

$$\operatorname{Spc}_{\operatorname{coh}} \mathfrak{T} := \operatorname{Pt} T$$

the cohomological spectrum of \mathcal{T} ; it is universal in the sense that any central action of a graded commutative ring R via a homomorphism $R \to Z^*(\mathcal{T})$ induces a continuous map $\operatorname{Spc}_{\operatorname{coh}} \mathcal{T} \to \operatorname{Spec}^{\vee} R$. In particular, the open subsets of $\operatorname{Spc}_{\operatorname{coh}} \mathcal{T}$ parameterise the thick subcategories of \mathcal{T} that are given by the central action of any graded commutative ring.

For a rigid tensor triangulated category \mathcal{T} there is an interesting relation between $\operatorname{Spc}_{\otimes} \mathcal{T}$ and $\operatorname{Spc}_{\operatorname{coh}} \mathcal{T}$; for a discussion of this we refer to [2].

For any choice of $T \subseteq$ Thick \mathcal{T} there is an obvious notion of *central support* relative to T for objects in \mathcal{T} , which is the analogue of the central support defined via (3.1) and parameterises the elements of Z(T); cf. Corollary 3.14. In particular, one may formulate a universal property of central support which takes T as input. The details are left to the interested reader.

References

- P. Balmer, The spectrum of prime ideals in tensor triangulated categories, J. Reine Angew. Math. 588 (2005), 149–168.
- [2] P. Balmer, Spectra, spectra, spectra—tensor triangular spectra versus Zariski spectra of endomorphism rings, Algebr. Geom. Topol. 10 (2010), no. 3, 1521–1563.
- [3] P. Balmer, Tensor triangular geometry, in Proceedings of the International Congress of Mathematicians. Volume II, 85–112, Hindustan Book Agency, New Delhi, 2010.
- [4] P. Balmer and G. Favi, Gluing techniques in triangular geometry, Q. J. Math. 58 (2007), no. 4, 415–441.
- [5] D. J. Benson, Cohomology of modules in the principal block of a finite group, New York J. Math. 1 (1994/95), 196–205.
- [6] D. J. Benson, J. F. Carlson and J. Rickard, Thick subcategories of the stable module category, Fund. Math. 153 (1997), no. 1, 59–80.
- [7] D. Benson, S. B. Iyengar and H. Krause, Local cohomology and support for triangulated categories, Ann. Sci. Éc. Norm. Supér. (4) 41 (2008), no. 4, 573–619.
- [8] D. Benson, S. B. Iyengar and H. Krause, A local-global principle for small triangulated categories, Math. Proc. Cambridge Philos. Soc. 158 (2015), no. 3, 451–476.
- [9] A.B. Buan, H. Krause, and Ø. Solberg, Support varieties-an ideal approach, Homology, Homotopy Appl., 9 (2007), 45–74.
- [10] A. B. Buan, H. Krause, N. Snashall, and Ø. Solberg, Support varieties—an axiomatic approach, Math. Z. 295 (2020), no. 1-2, 395–426.
- R.-O. Buchweitz and H. Flenner, Global Hochschild (co-)homology of singular spaces, Adv. Math. 217 (2008), no. 1, 205–242.
- [12] S. Gratz and G. Stevenson, Approximating triangulated categories by spaces, arXiv:2205.13356.
- [13] A. Grothendieck and J. L. Verdier, Préfaisceaux, in SGA 4, Théorie des Topos et Cohomologie Etale des Schémas, Tome 1. Théorie des Topos, Lect. Notes in Math., vol. 269, Springer, Heidelberg, 1972-1973, pp. 1-184.
- [14] M. Hochster, Prime ideal structure in commutative rings, Trans. Amer. Math. Soc. 142 (1969), 43–60.
- [15] A. Hubery and H. Krause, A categorification of non-crossing partitions, J. Eur. Math. Soc. (JEMS) 18 (2016), no. 10, 2273–2313.
- [16] C. Ingalls and H. Thomas, Noncrossing partitions and representations of quivers, Compos. Math. 145 (2009), no. 6, 1533–1562.
- [17] J. Kock and W. Pitsch, Hochster duality in derived categories and point-free reconstruction of schemes, Trans. Amer. Math. Soc. 369 (2017), no. 1, 223–261.
- [18] H. Krause and G. Stevenson, The derived category of the projective line, in Spectral structures and topological methods in mathematics, 275–297, EMS Ser. Congr. Rep. EMS Publ. House, Zürich, 2019.

- [19] D. K. Nakano, K. B. Vashaw and M. T. Yakimov, Noncommutative tensor triangular geometry, Amer. J. Math. 144 (2022), no. 6, 1681–1724.
- [20] A. Neeman, The Brown representability theorem and phantomless triangulated categories, J. Algebra 151 (1992), no. 1, 118–155.
- [21] J. Rickard, Idempotent modules in the stable category, J. London Math. Soc. (2) 56 (1997), no. 1, 149–170.
- [22] C. M. Ringel, Algebra at the turn of the century, Lecture at the National Conference on Algebra VII, Beijing Normal University, October 1999, Southeast Asian Bull. Math. 25 (2001), no. 1, 147–160.
- [23] R. W. Thomason, The classification of triangulated subcategories, Compositio Math. 105 (1997), no. 1, 1–27.
- [24] T. tom Dieck, Algebraic topology, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2008.
- [25] K. B. Vashaw, Private communication.
- [26] J.-L. Verdier, Des catégories dérivées des catégories abéliennes, Astérisque, 239, Société Mathématique de France, 1996.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, D-33501 BIELEFELD, GERMANY *Email address*: hkrause@math.uni-bielefeld.de