Variations on the Goroff-Sagnotti operator.

Enrique Álvarez, Jesús Anero and Eduardo Velasco-Aja.

Departamento de Física Teórica and Instituto de Física Teórica, IFT-UAM/CSIC, Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain

E-mail: enrique.alvarez@uam.es, jesusanero@gmail.com and eduardo.velasco@uam.es

Abstract

The effect of modifying General Relativity with the addition of some higher dimensional operators, generalizations of the Goroff-Sagnotti operator, is discussed. We determine in particular, the general solution of the classical equations of motion, assuming it to be spherically symmetric, not necessarily static. Even in the non-spherically symmetric case, we present a necessary condition for an algebraically generic spacetime to solve the corresponding equations of motion. Some examples of an application of said condition are explicitly worked out.

Contents

1	Prolegomena. 1.1 The operator $\mathcal{O}_2 \equiv W^{\mu\nu}{}_{\alpha\beta}W^{\alpha\beta}{}_{\mu\nu}$	2 3 4
2	Solving the trace equation.	5
3	Spherically symmetric spacetimes (S3).	7
4	Conclusions.	7
5	Acknowledgements.	9
A	The S3 Weyl tensor.	10

1 Prolegomena.

In [2] we have explored some consequences of the incorporation of the Goroff-Sagnotti counterterm to the Einstein-Hilbert action. Let us remind the reader that the Goroff-Sagnotti [10] counterterm is the first nontrivial renormalization of Einstein-Hilbert's lagrangian. It first appears at the two-loop order because general relativity is one-loop finite, as shown in [17]. To be specific,

$$\mathcal{O}_{GS} \equiv W^{\mu\nu}_{\ \rho\sigma} W^{\rho\sigma}_{\ \gamma\delta} W^{\gamma\delta}_{\ \mu\nu},\tag{1}$$

where $W^{\mu\nu}{}_{\rho\sigma}$ is Weyl's tensor¹. The structure of this operator is suggestive of a simple generalization. The purpose of the present letter is to study these more general operators, to wit

$$\mathcal{O}_p \equiv \operatorname{tr} W^p \equiv W_{\mu_1 \nu_1 \rho_1 \sigma_1} \dots W_{\mu_p \nu_p \rho_p \sigma_p} I^{\vec{\mu} \vec{\nu} \vec{\rho} \vec{\sigma}}, \qquad (2)$$

where $I^{\vec{\mu}\vec{\nu}\vec{\rho}\vec{\sigma}}$ is the product of 2p inverse metric tensors, for example

$$I^{\vec{\mu}\vec{\nu}\vec{\rho}\vec{\sigma}} \equiv g^{\mu_1\mu_2}\dots g^{\mu_{p-1}\mu_p} g^{\nu_1\nu_2}\dots g^{\nu_{p-1}\nu_p} g^{\rho_1\rho_2}\dots g^{\rho_{p-1}\rho_p} g^{\sigma_1\sigma_2}\dots g^{\sigma_{p-1}\sigma_p}, \tag{3}$$

although more complicated contractions are possible, they are all similar in character.

The main property of Weyl's tensor is that it is inert² under the so-called Weyl rescalings (a gauge transformation) of the spacetime metric

$$g_{\mu\nu} \to \Omega(x)^2 g_{\mu\nu},$$
 (4)

$$W^{\mu}{}_{\nu\rho\sigma} \to W^{\mu}{}_{\nu\rho\sigma}.$$
 (5)

 $^1\mathrm{Let}$ us recall the definition of Weyl's tensor as

$$W_{\alpha\beta\gamma\delta} \equiv R_{\alpha\beta\gamma\delta} - \frac{1}{n-2} \left\{ g_{\alpha\gamma} R_{\beta\delta} - g_{\alpha\delta} R_{\beta\gamma} - g_{\beta\gamma} R_{\delta\alpha} + g_{\beta\delta} R_{\gamma\alpha} \right\} + \frac{1}{(n-1)(n-2)} R \left\{ g_{\alpha\gamma} g_{\delta\beta} - g_{\alpha\delta} g_{\gamma\beta} \right\}$$

²Bear in mind the position of the indices in eq. (5).

This means that $S_p \equiv \int d(vol) \mathcal{O}_p$ (where $d(vol) \equiv \sqrt{|g|} d^n x$), is conformally invariant whenever n = 2p.

It would seem that, at least for compact spacetimes, the corresponding action S_p should coincide with the integral of Branson's Q-curvature [6]

$$S_Q \equiv \int d(vol)Q,\tag{6}$$

whose variation is Fefferman and Graham's [9] obstruction tensor (a generalization of sorts of Bach's tensor [4] to arbitrary even dimension). But in fact, Alexakis' theorem [1] proving Deser and Schwimmer's [7] conjecture guarantees that Branson's curvature is a linear combination of conformal invariants constructed out of Riemann tensor and its covariant derivatives plus total derivatives and Euler's density (the pfaffian).

$$Q = Weyl + Pfaff + \nabla_{\mu}j^{\mu}.$$
(7)

Unfortunately, starting in n = 6 [3, 5] there are other conformally invariant operators which are not Weyl polynomials. This means that starting on this dimension, our former conjecture ceases to hold.

The operators \mathcal{O}_p are, however, the only invariants not involving any derivative of Riemann's tensor.

The main interest of those operators from our point of view is that they are natural generalizations of the Goroff-Sagnotti counterterm [10]. Some of the consequences of incorporating this operator in the classical action have been studied in a previous paper [2]. The main purpose of this letter is to study generalizations of the type $p \ge 3$.

Also, in the preceding work, we concentrated mainly on the spherically symmetric case; here we attempt to make more general statements. We shall examine in detail a particular equation (the trace equation) which is a necessary (and in many cases sufficient) condition for solving the equations of motion (EM).

1.1 The operator $\mathcal{O}_2 \equiv W^{\mu\nu}{}_{\alpha\beta}W^{\alpha\beta}{}_{\mu\nu}$.

Let us now consider the four-dimensional conformal action

$$S_{Weyl^2} \equiv \int d^4x \sqrt{|g|} W_{\mu\nu\rho\sigma} W^{\mu\nu\rho\sigma}.$$
 (8)

In n = 4 dimensions, using Lanczos' identity for the four-dimensional Weyl tensor [11],

$$W_{\mu\alpha\beta\lambda}W_{\nu}^{\ \alpha\beta\lambda} = \frac{1}{4}g_{\mu\nu}W^{\rho\sigma}_{\ \alpha\beta}W^{\alpha\beta}_{\ \rho\sigma},\tag{9}$$

the EM are exactly [4] Bach's tensor

$$\frac{\delta S_{Weyl^2}}{\delta g^{\mu\nu}} = \sqrt{|g|} \left\{ \nabla^\lambda \nabla^\tau W_{\lambda\mu\tau\nu} + \frac{1}{2} W_{\lambda\mu\tau\nu} R^{\lambda\tau} \right\} = \sqrt{|g|} B_{\mu\nu} = 0, \tag{10}$$

which is traceless;

$$B^{\lambda}_{\ \lambda} = 0. \tag{11}$$

Were we to consider this same action in arbitrary dimensions the EM are,

$$\frac{4R^{\alpha\beta}W_{\mu\alpha\nu\beta}}{n-2} - \frac{1}{2}g_{\mu\nu}W_{\alpha\beta\gamma\epsilon}W^{\alpha\beta\gamma\epsilon} + 2W_{\mu}^{\ \alpha\beta\gamma}W_{\nu\alpha\beta\gamma} + 4\nabla_{(\alpha}\nabla_{\beta)}W_{\mu}^{\ \alpha}{}_{\nu}^{\ \beta} = 0, \qquad (12)$$

whose trace reads,

$$(2 - \frac{1}{2}n)W_{\alpha\beta\gamma\epsilon}W^{\alpha\beta\gamma\epsilon} = 0.$$
 (13)

We shall see later on the importance of this trace equation.

1.2 The Goroff-Sagnotti operator $\mathcal{O}_3 \equiv W_{\mu\nu\alpha\beta}W^{\alpha\beta\rho\sigma}W_{\rho\sigma}{}^{\mu\nu}$.

The action for \mathcal{O}_{GS} is

$$S_{GS} \equiv \int d^4x \sqrt{|g|} W_{\mu\nu\alpha\beta} W^{\alpha\beta\rho\sigma} W_{\rho\sigma}^{\ \mu\nu}.$$
 (14)

It is instructive to work out the EM in arbitrary dimensions. They read

$$V_{\mu\nu}^{(1)} = \frac{1}{2} g_{\mu\nu} \mathcal{O}_{GS} + \frac{6}{(n-2)} \Biggl[\frac{1}{n-1} \left(R W_{\mu}^{\ \alpha\beta\lambda} W_{\nu\alpha\beta\lambda} - R_{\mu\nu} W^2 \right) - R^{\alpha\beta} W_{\mu\alpha\lambda\tau} W^{\lambda\tau}_{\ \nu\beta} - 2\nabla^{\tau} \nabla_{\mu} (W_{\nu}^{\ \alpha\beta\lambda} W_{\tau\alpha\beta\lambda}) + g_{\mu\nu} \nabla^{\rho} \nabla^{\sigma} (W_{\rho}^{\ \alpha\beta\lambda} W_{\sigma\alpha\beta\lambda}) - (n-2) \nabla^{\rho} \nabla^{\sigma} (W_{\mu\rho\alpha\beta} W^{\alpha\beta}_{\ \nu\sigma}) + \left(\frac{(\nabla_{\mu} \nabla_{\nu} - g_{\mu\nu} \Box)}{n-1} W^2 + \Box (W_{\mu}^{\ \alpha\beta\lambda} W_{\nu\alpha\beta\lambda}) \Biggr) + R_{\mu}^{\ \tau} W_{\nu}^{\ \alpha\beta\lambda} W_{\tau\alpha\beta\lambda} \Biggr] - 3W_{\mu}^{\ \alpha\beta\lambda} W_{\nu\alpha\rho\sigma} W^{\rho\sigma}_{\ \beta\lambda}.$$
(15)

The trace of eq. (15) corresponds to,

$$V^{(1)} = \frac{n-6}{2} W_{\mu\nu\alpha\beta} W^{\alpha\beta\rho\sigma} W_{\rho\sigma}^{\ \mu\nu}.$$
 (16)

We see that in any dimension different from the conformal one (n = 6) a *necessary* condition for the EM to be satisfied is that

$$\mathcal{O}_3 := W_{\mu\nu\alpha\beta} W^{\alpha\beta\rho\sigma} W_{\rho\sigma}^{\ \mu\nu} = 0.$$
(17)

For the other contraction (which is independent of the former for $n \ge 6$), the story is quite similar

$$S_{Weyl^3} \equiv \int d^4x \sqrt{-g} W^{\alpha}_{\gamma}{}^{\beta}_{\epsilon} W^{\gamma}_{\rho}{}^{\epsilon}_{\sigma} W^{\rho}_{\alpha}{}^{\sigma}_{\beta}, \qquad (18)$$

varying eq. (18) yields the EM,

$$V_{\mu\nu}^{(2)} = -\frac{3}{n-2} \left[\frac{1}{n-1} \left(2RW_{\mu\alpha\beta\lambda} W_{\nu}^{\ \beta\alpha\lambda} + R_{\mu\nu} W_{\alpha\beta\rho\sigma} W^{\alpha\rho\beta\sigma} + (\nabla_{\mu} \nabla_{\nu} - g_{\mu\nu} \Box) (W_{\alpha\beta\rho\sigma} W^{\alpha\rho\beta\sigma}) - 2\nabla^{\lambda} \nabla_{\mu} (W_{\nu\alpha\beta\tau} W_{\lambda}^{\ \beta\alpha\tau}) + g_{\mu\nu} \nabla^{\rho} \nabla^{\sigma} (W_{\rho\alpha\beta\tau} W_{\sigma}^{\ \beta\alpha\tau}) + \Box (W_{\mu\alpha\beta\tau} W_{\nu}^{\ \beta\alpha\tau}) - 2R^{\rho\sigma} \left(W_{\mu}^{\ \alpha\beta} W_{\nu\beta\sigma\alpha} - W_{\mu}^{\ \alpha\beta} W_{\rho\alpha\sigma\beta} \right) \right] + 3R_{\mu}^{\ \alpha\lambda\beta} W_{\nu}^{\ \rho\sigma} W_{\alpha\rho\beta\sigma} - 6W_{\mu}^{\ \alpha\lambda\beta} W_{\nu}^{\ \rho\sigma} W_{\alpha\rho\beta\sigma}) + \frac{1}{2}g_{\mu\nu} W_{\gamma}^{\ \alpha\beta} W_{\rho}^{\ \sigma\sigma} W_{\alpha\beta}^{\ \sigma\sigma} - 3\nabla^{\rho} \nabla^{\sigma} (W_{\mu\nu}^{\ \alpha\beta} W_{\rho\beta\sigma\alpha}) + 3\nabla^{\rho} \nabla^{\sigma} (W_{\mu\nu}^{\ \alpha\beta} W_{\nu\beta\sigma\alpha}),$$
(19)

as in the previous case, the trace of the EM now corresponds to

$$V^{(2)} = \frac{n-6}{2} W^{\alpha}{}_{\gamma}{}^{\beta}{}_{\epsilon} W^{\gamma}{}_{\rho}{}^{\epsilon}{}_{\sigma} W^{\rho}{}_{\alpha}{}_{\beta}, \qquad (20)$$

with a similar necessary condition for the trace. We now see how this generalizes for $\mathcal{O}_{p\geq 3}$.

2 Solving the trace equation.

Let us consider the action

$$S_p \equiv \int d(vol)\mathcal{O}_p \equiv \int d^n x \mathcal{B}_p,\tag{21}$$

in arbitrary dimension n. Notice that we have defined $\mathcal{B}_p \equiv \sqrt{|g|} \mathcal{O}_p$

Consider a linearized Weyl rescaling

$$\delta_W g_{\mu\nu} = \omega(x) g_{\mu\nu}. \tag{22}$$

Then

$$\delta_W S_p = \int d^n x \delta_W \mathcal{B}_p = \int d^n x (n-2p) \omega \mathcal{B}_P = \int d^n x \frac{\delta S_p}{\delta g_{\mu\nu}} \omega g^{\mu\nu}.$$
 (23)

It follows that

$$g^{\mu\nu}\frac{\delta S_p}{\delta g_{\mu\nu}} = (n-2p)\mathcal{B}_P,\tag{24}$$

which conveys the fact that the action is conformal³ in dimension n = 2p.

This is the motivation for studying in some detail the trace equation (24), which will not always imply the vanishing of Weyl's tensor.

Let us represent, following $[12]^4$ and [15, 18], the 10 independent components of Weyl's tensor as a six-dimensional block matrix

$$W^{\mu\nu}{}_{\rho\sigma} = \begin{pmatrix} E & -B \\ B & E \end{pmatrix}, \tag{25}$$

where both B and E are three-dimensional, traceless, symmetric and real matrices. Clearly⁵,

$$W^{2} = N \begin{pmatrix} E^{2} - B^{2} & -(EB + BE) \\ BE + EB & E^{2} - B^{2} \end{pmatrix} \to \operatorname{tr} W^{2} = N \operatorname{tr} (E^{2} - B^{2}).$$
(26)

Similarly,

$$W^{3} = N \begin{pmatrix} E^{3} - EB^{2} - B^{2}E - BEB & B^{3} - BE^{2} - E^{2}B - BE^{2} - EBE \\ BE^{2} - B^{3} + EBE + E^{2}B & E^{3} - EB^{2} - B^{2}E - BEB \end{pmatrix},$$
(27)

 $^{^{3}}$ We stress that the utility of the trace equation lies on its simplicity which is directly tied to conformal invariance.

 $^{^4\}mathrm{In}$ fact, Matte's work precedes Petrov's by one year, although he did not achieve a full classification of Weyl's tensor.

⁵Here N takes care of index undercounting and its exact value is unmaterial for our purposes.

where now, using the cyclic property of the trace,

$$\mathcal{O}_{GS} = \operatorname{tr} W^3 = N \operatorname{tr} \left(E^3 - 3EB^2 \right).$$
⁽²⁸⁾

Given the fact that Schwarzschild spacetime, (as in fact all static, spherically symmetric spacetimes) have purely electric Weyl tensors, it can be easily shown that for them,

$$\mathcal{O}_{GS} = 0 \Longrightarrow W_{\mu\nu\rho\sigma} = 0. \tag{29}$$

For purely magnetic spacetimes, $\mathcal{O}_{GS} = 0$ is given. Unfortunately, there are not many known purely magnetic exact solutions. There is a theorem stated in [13] claiming that there are no purely magnetic Type D vacuum solutions of Einstein's equations.

Starting with \mathcal{O}_4 , the cyclic property of the trace is not enough to get a simple result. To be more specific, we need some explicit results [15, 16] on Petrov's classification of spacetimes. They are frequently expressed in terms of the properties of the matrix

$$Q \equiv E + iB. \tag{30}$$

The algebraically general spacetime is dubbed Petrov Type I. For these spacetimes, the general criterion is

$$(Q - \lambda_1)(Q - \lambda_2)(Q - \lambda_3) = 0, \qquad (31)$$

(where $\sum \lambda_i = 0$; and $\lambda_i \equiv e_i + ib_i$). The electric and magnetic Weyl tensors obey

$$E = \begin{pmatrix} e_1 & 0 & 0\\ 0 & e_2 & 0\\ 0 & 0 & e_3 \end{pmatrix}, \quad \text{and} \quad B = \begin{pmatrix} b_1 & 0 & 0\\ 0 & b_2 & 0\\ 0 & 0 & b_3 \end{pmatrix}$$
(32)

with $\sum e_i = \sum b_i = 0$. It is plain that in this case, [E, B] = 0 so the trace is given by

$$\operatorname{tr} W^{3} = 2 N \sum_{i} e_{i} (e_{i}^{2} - 3b_{i}^{2}).$$
(33)

Type D is a subtype that corresponds to $\lambda_1 = \lambda_2 = -2\lambda_3$ in eq. (31). This particular Petrov Type is important for us because all spherically symmetric solutions [16] are either Type D or else Type O, (which means that the spacetime is conformally flat).

The general result for the all-important Types I and D is obtained from the above discussion;

$$\operatorname{tr} W^{2m} = 2 \sum_{k=0}^{m} (-1)^k \binom{2m}{2k} \operatorname{tr} \left(E^{2(m-k)} B^{2k} \right), \qquad (34)$$

$$\operatorname{tr} W^{2m+1} = 2 \sum_{k=0}^{m} (-1)^k \binom{2m+1}{2k} \operatorname{tr} \left(E^{2(m-k)+1} B^{2k} \right).$$
(35)

This gives immediately the conditions for the vanishing of the trace in terms of products of the electric and magnetic eigenvalues $\{e_i, b_i\}$.

3 Spherically symmetric spacetimes (S3).

Let us now particularize to the case of a spherically symmetric spacetime S3 with metric tensor,

$$ds^{2} = B(t,r)dt^{2} - A(t,r)dr^{2} - r^{2} d\Omega_{2}^{2}.$$
(36)

In [2] we were mostly concerned with $\operatorname{tr} W^3$ but we also made the observation that for this particular S3 spacetimes,

$$\mathcal{O}_4 := W^{\alpha\beta}_{\ \gamma\delta} W^{\gamma\delta}_{\ \kappa\zeta} W^{\kappa\zeta}_{\ \lambda\sigma} W^{\lambda\sigma}_{\ \alpha\beta} \,, \tag{37}$$

is proportional to the lagrangian of Conformal Gravity (CG) in eq. (8).

As a matter of fact, [2, 8] Weyl's tensor for an S3 depends on a unique combination of A(t, r), B(t, r) and their derivatives. Using the notation of appendix A, it so happens that

$$\sqrt{-g}\mathcal{O}_2 = \frac{\sqrt{r^4 A B \sin^2(\theta)}}{12r^4 A^4 B^4} G^2,$$
(38)

and,

$$\sqrt{-g}\mathcal{O}_4 = \frac{\sqrt{r^4 A B \sin^2(\theta)}}{576 r^8 A^8 B^8} G^4.$$
(39)

However, we stressed that CG was different from any other S_p action in which it contained solutions not present for the $p \neq 2$ cases. It is important to understand how the p = 4fails to capture such solutions even when this similarity is present.

This simple relationship between both lagrangians does not carry over to the corresponding solutions. As shown in [2], the solutions for CG contain those of S_4 but not conversely. Of course, for CG the trace equation (11) does not yield any restriction on the conformal case n = 4. On the other hand, for the EM for S_4 , the trace condition applies. Therefore, the only way in which a metric can be a solution to both CG and quartic gravity is to satisfy, in addition

$$W^{2} = W^{\alpha\beta}_{\ \gamma\delta} W^{\gamma\delta}_{\ \alpha\beta} = 0 \Rightarrow G = 0 \Rightarrow W^{\alpha}_{\ \beta\gamma\delta} = 0, \tag{40}$$

i.e., it has to be conformally-flat. In fact for any $p \neq 2$, S3 solutions are determined by eq. (40).

4 Conclusions.

Let us finish with a couple of examples that depart from spherical symmetry and thus from the theorem proven in [2].

• First we consider the spacetime of p-p waves with line element [16],

$$ds^{2} = -2\mathrm{du}^{2}H(x, y, u) + 2\mathrm{dudv} - \left(\mathrm{dx}^{2} + \mathrm{dy}^{2}\right).$$
(41)

To solve Einstein's equations in vacuum, H(x, y, u) has to satisfy

$$R_{\mu\nu} = 0 \quad \Leftrightarrow \quad \partial_x^2 H(x, y, u) + \partial_y^2 H(x, y, u) = 0.$$
(42)

The above-introduced analysis yields electric and magnetic components of the Weyl tensor corresponding to,

$$E = \frac{1}{2H(x, y, u) + 2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \partial_x^2 H(x, y, u) & \partial_x \partial_y H(x, y, u) \\ 0 & \partial_x \partial_y H(x, y, u) & -\partial_x^2 H(x, y, u) \end{pmatrix},$$
(43)

and,

$$B = \frac{1}{2H(x, y, u) + 2} \begin{pmatrix} 0 & 0 & 0\\ 0 & -\partial_x \partial_y H(x, y, u) & \partial_x^2 H(x, y, u)\\ 0 & \partial_x^2 H(x, y, u) & \partial_x \partial_y H(x, y, u) \end{pmatrix}.$$
 (44)

In eqs. (43) and (44), eq. (42) has been used. Therefore, we see that p-p waves are everywhere of Petrov Type N. This implies,

$$Q^2 = 0, (45)$$

which translates into the trace condition being satisfied for them, i.e.,

$$\operatorname{tr} W^3 \propto \operatorname{tr} Q^3 = 0. \tag{46}$$

Thus, p-p waves comprise an example of a spacetime satisfying the necessary condition to be a solution to the EM. Here, direct computation allows us to see that this spacetime also satisfies the Goroff-Sagnotti EM, eq. (15).

In [2] it was proven that no spherically symmetric spacetime can simultaneously solve Einstein's and Goroff-Sagnotti's EM. It is interesting to remark that p-p waves are the only example of a solution to both EM we have been able to identify to this date.

• As a second non-spherically symmetric nor static example, we consider Kerr-Newmann spacetime, whose line element, in Boyer-Lindquist coordinates [14] reads,

$$ds^{2} = \frac{\Delta}{\rho^{2}} \left(dt - a \sin^{2} \theta \, d\phi \right)^{2} - \frac{\sin^{2} \theta}{\rho^{2}} \Big[\left(r^{2} + a^{2} \right) \, d\phi - a \, dt \Big]^{2} - \frac{\rho^{2}}{\Delta} dr^{2} - \rho^{2} \, d\theta^{2}, \quad (47)$$

where,

$$\Delta := r^2 - rr_s + a^2 + r_q^2, \qquad \text{and} \qquad \rho := a^2 \cos^2(\theta) + r^2.$$
(48)

As usual, a is related to the angular momentum, r_s is the Schwarzschild radius, related to the mass and r_q is related to the electric charge ⁶ of the solution.

In this case, the electric and magnetic matrices are,

$$E = e(r,\theta) \begin{pmatrix} 1 & 0 & 0\\ 0 & -\frac{1}{2} & 0\\ 0 & 0 & -\frac{1}{2} \end{pmatrix}, \text{ with } e(r,\theta) = \frac{4r_s r \left(3a^2 - 2r^2 + 3a^2 \cos(2\theta)\right)}{\left(a^2 + 2r^2 + a^2 \cos(2\theta)\right)^3},$$
(49)

$$B = b(r,\theta) \begin{pmatrix} 1 & 0 & 0\\ 0 & -\frac{1}{2} & 0\\ 0 & 0 & -\frac{1}{2} \end{pmatrix}, \text{ with } b(r,\theta) = \frac{4r_s a \cos(\theta) \left(a^2 - 6r^2 + a^2 \cos(2\theta)\right)}{\left(a^2 + 2r^2 + a^2 \cos(2\theta)\right)^3}.$$
 (50)

⁶We set $r_q = 0$ such that the line element in eq. (47) is a solution to the vacuum Einstein's EM.

This corresponds to a Petrov Type D spacetime, following the above discussion,

$$trW^{3} = \frac{3}{4}e(r,\theta)(e(r,\theta)^{2} - 3b(r,\theta)^{2}) = 0.$$
(51)

The trace in eq. (51) is nonvanishing⁷ for arbitrary r values. We conclude that an uncharged Kerr spacetime with mass cannot solve the EM for the Goroff-Sagnotti counterterm. This we believe to be an important result. Not only Schwarzschild but also its most important stationary extension, namely Kerr (widely believed to be the general endpoint of gravitational collapse), is incompatible with the Goroff-Sagnotti lagrangian deformation.

Let us finish with a quick summary of this work. In this letter, we have studied the effects of including generalizations of the Goroff-Sagnotti operator on Einstein-Hilbert's action for GR. Specifically, we have focused on the equations of motion and obtained one necessary condition for their solutions.

In detail, the operators we have studied are built of p contractions of the Weyl tensor, symbolically,

$$\int d^n x \,\sqrt{|g|} \mathcal{O}_p := \int d^n x \,\sqrt{|g|} \mathrm{tr} \,W^p.$$
(52)

These are conformally invariant in dimension n = 2p. Using the so-called trace equation and the Petrov classification of Weyl's tensor, we have found a necessary condition for the solutions of the EM in the general case.

In the more simple instance of spherically symmetric spacetimes, which motivated this study originally [2], we are able to find the general solution of the EM. This is because, in this case, the trace condition becomes a necessary and sufficient condition for a spacetime to solve for the complete EM.

Examples for the *less trivial* p-p waves and Kerr-spacetime are explicitly worked out. p-p waves are found to be a solution to the EM of GR with the p = 3 operator included.

5 Acknowledgements.

We have enjoyed fruitful discussions with Prof. Senovilla. We acknowledge partial financial support by the Spanish MINECO through the Centro de Excelencia Severo Ochoa Program under Grant CEX2020-001007-S funded by MCIN/AEI/10.13039/501100011033 All authors acknowledge the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No 860881-HIDDeN and also by Grant PID2019-108892RB-I00 funded by MCIN/AEI/ 10.13039/501100011033 and by "ERDF A way of making Europe". This project has received funds/support from the European Union's Horizon Europe programme uncer Marie Sklodowska-Curie Actions-Staff Exchanges (SE) grant agreement No 101086085-ASYMMETRY.

⁷Unless $r_s = 0$.

A The S3 Weyl tensor.

Direct computation of the Weyl tensor for S3 spacetimes yields

$$W^{\mu\nu}_{\ \rho\sigma} = \frac{G}{12A^2B^2} C^{\mu\nu}_{\ \rho\sigma}, \tag{A.1}$$

In particular,

$$G(A, A', B, B', \cdots) := A \left(r^2 \left(B'^2 - \dot{A}\dot{B} \right) + 2rB \left(r\ddot{A} + B' - rB'' \right) - 4B^2 \right) + rB \left(A' \left(rB' - 2B \right) - r\dot{A}^2 \right) + 4A^2B^2,$$
(A.2)

where $\dot{f} = \partial_t f(r, t)$ and $f' = \partial_r f(r, t)$. $C^{\mu\nu}_{\ \rho\sigma}$ is a tensor independent of the metric. In detail;

This Weyl tensor is purely electric, with

$$E = \frac{G}{12A^2B^2} \begin{pmatrix} -1 & 0 & 0\\ 0 & \frac{1}{2} & 0\\ 0 & 0 & \frac{1}{2} \end{pmatrix}.$$
 (A.4)

References

- S. Alexakis, "On the decomposition of global conformal invariants I". Ann. of Math. (2) 170 (2009), no. 3, 1241–1306. arXiv:0509571.
- [2] E. Álvarez, J. Anero and E. Velasco-Aja, "The fate of horizons under quantum corrections," [arXiv:2207.12721 [hep-th]].
 E. Alvarez, J. Anero and R. Santos-Garcia, "Structural stability of spherical horizons," [arXiv:2006.02463 [hep-th]].
- [3] E. Alvarez, J. Conde and L. Hernandez, "The Dirichlet obstruction in AdS / CFT," Int. J. Mod. Phys. D 12, 1415-1430 (2003) doi:10.1142/S0218271803003839
 [arXiv:hep-th/0303164 [hep-th]].

- [4] Rudolf Bach, "Zur Weylschen Relativitätstheorie und der Weylschen Erweiterung des Krümmungstensorbegriffs", Mathematische Zeitschrift, 9 (1921) pp. 110
- [5] L. Bonora, P. Pasti and M. Bregola, "Weyl Cocycles," Class. Quant. Grav. 3, 635 (1986) doi:10.1088/0264-9381/3/4/018
- [6] T. Branson and A. R. Gover, "Conformally invariant operators, differential forms, cohomology and a generalisation of Q-curvature," [arXiv:math/0309085 [math.DG]].
- S. Deser and A Schwimmer, "Geometric classification of conformal anomalies in arbitrary dimensions," Phys. Lett. B 309, 279-284 (1993) doi:10.1016/0370-2693(93)90934-A [arXiv:hep-th/9302047 [hep-th]].
- [8] S. Deser and J. Franklin, "Birkhoff for Lovelock redux," Class. Quant. Grav. 22 (2005), L103-L106 doi:10.1088/0264-9381/22/16/L03 [arXiv:gr-qc/0506014 [gr-qc]].
 S. Deser and A. V. Ryzhov, "Curvature invariants of static spherically symmetric geometries," Class. Quant. Grav. 22, 3315-3324 (2005) doi:10.1088/0264-9381/22/16/012 [arXiv:gr-qc/0505039 [gr-qc]].
 S. Deser and B. Tekin, "Shortcuts to high symmetry solutions in gravitational theories," Class. Quant. Grav. 20 (2003), 4877-4884 doi:10.1088/0264-9381/20/22/011 [arXiv:gr-qc/0306114 [gr-qc]].
- [9] C. Fefferman and C. R. Graham, "The ambient metric," Ann. Math. Stud. 178, 1-128 (2011) [arXiv:0710.0919 [math.DG]].
- [10] M. H. Goroff and A. Sagnotti, "The Ultraviolet Behavior of Einstein Gravity," Nucl. Phys. B 266, 709-736 (1986) doi:10.1016/0550-3213(86)90193-8
- C. Lanczos, "A Remarkable property of the Riemann-Christoffel tensor in four dimensions," Annals Math. 39 (1938), 842-850 doi:10.2307/1968467 Lovelock, D. (1970). Dimensionally dependent identities. Mathematical Proceedings of the Cambridge Philosophical Society, 68(2), 345-350. doi:10.1017/S0305004100046144
- [12] A.Matte, "Sur de nouvelles solutions oscillatoires des equations de la gravitation" Canadian J of Math 5,1 (1953)
- [13] C B G McIntosh, R Arianrhodt, S T Wadet and C Hoenselaers, "Electric and magnetic Weyl tensors: classification and analysis", 1994 Class. Quantum Grav. 11 1555
- [14] Barrett O'Neill, "The geometry of Kerr black holes" (Dover, NY, 2014)
- [15] FAE Pirani, "Invariant formulation of gravitational radiation theory," Phys. Rev. 105, 1089-1099 (1957) doi:10.1103/PhysRev.105.1089
- [16] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers and E. Herlt, "Exact solutions of Einstein's field equations," Cambridge Univ. Press, 2003, ISBN 978-0-521-46702-5, 978-0-511-05917-9 doi:10.1017/CBO9780511535185
- [17] G. 't Hooft and M. J. G. Veltman, "One loop divergencies in the theory of gravitation," Ann. Inst. H. Poincare Phys. Theor. A 20 (1974), 69-94

[18] J. A. Wheeler, "Gravitational and Electromagnetic Wave Flux Compared and Contrasted," Phys. Rev. D 16, 3384-3389 (1977) doi:10.1103/PhysRevD.16.3384