arXiv:2301.10867v2 [gr-qc] 9 Feb 2023

Resolution of cosmological singularity in Hořava-Lifshitz cosmology

Ewa Czuchry^{1,*}

¹National Centre for Nuclear Research, Pasteura 7, Warsaw, Poland (Dated: February 13, 2023)

The standard ACDM model despite its agreement with observational data still has some issues unaddressed, lie the problem of initial singularity. Solving that problem usually requires modifications of general relativity. However, there appeared the Hořava-Lifshitz (HL) theory of gravity, in which equations governing cosmological evolution include a new term scaling similarly as dark radiation term in the Friedmann equations, enabling a bounce of the universe instead of initial singularity. This review describes past works on a stability of such a bounce in different formulations of HL theory, initial detailed balance scenario and further projectable versions containing higher than quadratic term to the original action.

^{*} ewa.czuchry@ncbj.gov.pl

I. INTRODUCTION

Classical General Relativity (GR) apart its simple beauty and symmetry is also strongly confirmed in several experimental tests. However, it does not explain many issues like dark matter, spacetime singularities including the initial one in cosmology, and the ones inside the black holes. In order to answer these issues there has been many attempts to modify GR both on the classical and quantum level. Specifically, the quantisation of GR cosmology was supposed to resolve the initial singularity problem.

Attempts to quantize gravity could be divided into two categories. One way was to assume the classical theory of gravity and quantise it in various manners, with the first attempts performed via the the covariant quantum gravity. In that classical approach one repeats the method successful in quantising electrodynamics, namely considering the path integral of the Hilbert-Einstein action and then calculates the perturbation of the metric around a background one. The obtained equations unlike in electrodynamics are non-renormalizable in higher energies. The canonical quantum gravity considers ADM (3 + 1)-decomposition of the spacetime and quantisation of the constraints obtained from Hamiltonian. Other attempts included sophisticated theories like string theory and loop quantum gravity. These theories manage to solve some problems (such as a cosmological singularity [1]) but there are difficult to be phenomenologically tested [2, 3]. There are also attempts for resolving an initial singularity problem by combination of canonical and coherent state quantisation like the one in our paper [4], however at this moment they are difficult to be validated by observations data.

Although there is still no full theory of quantum gravity developed it is supposed to manifest beyond a characteristic energy scale for quantum gravity $E_{Pl} = \sqrt{\hbar c^5/G}$ built in terms of the speed of light c, the gravitational constant Gand the Planck's constant \hbar . Therefore, there is the second research direction which aims to construct a modified version of GR with an improved UV behavior. General relativity after many tests performed seems to be consistent with all current observations. This makes it a very good IR limit of potential quantum gravity model. There has been made some proposals for UV-completions of general relativity in the past [5, 6]. They have one thing in common, namely the existence of some cutoff energy scale beyond which quantum effects could be detected, specially for a cutoff energy in the range of TeV. The widely discussed recent proposal is Hořava gravity, which is a proposal of a UV complete theory of gravity. It seems to be renormalizable at high energies, which makes it a candidate for a quantum gravity model [7, 8]. The action of this theory contains additional higher order spatial derivatives and therefore the theory loses the full diffeomorphism invariance, keeping the (1+3) foliation preserving diffeomorphism. Moreover, there is an UV fixed point in this gravity model where there is an anisotropic Lifshitz scaling between time and space. Therefore, the resulting theory is called Hořava-Lifshitz (HL) gravity.

Significant work has been done on this theory where different aspects and properties were examined [9–22]. Many studies were devoted to cosmological solutions [12, 19, 23], braneworlds and dark radiation [12, 21]. Hořava-Lifshitz cosmology obtained a novel feature enabling the existence of bounce instead of initial singularity predicted by classical GR. There has been also other research focused on finding specific solutions, including black holes, and their properties, and many works devoted to phenomenological aspects both astrophysical and concerning dark matter.

Derivation of Hořava-Lifshitz cosmology [12, 19, 23] made via varying action written Friedmann-Robertson-Walker space-time metrics resulted in equations analogous to the standard Friedman ones. These equations contain a new term which scales similarly as dark radiation [12, 19, 21], i.e. $\sim 1/a^4$ (where *a* is a scale factor), and provides a negative contribution to the energy density. This feature enables obtaining non-singular cosmological evolution, resolving the initial singularity problem [14, 21, 24]. Such a possibility not only results in avoiding the initial singularity but may have other consequences for potential histories of the Universe like scenario of contraction from the infinite size connected by a bounce to the expansion to infinite size again, or eternal cycles of the similar scenario.

Despite many promises made by this modified theory of gravity it seems that it contains instabilities and pathologies in different formulations (see e.g. [22, 25–27]). The original Hořava formulation suffers from among many problems: the existence of ghost instabilities and strong coupling at IR [10, 28], the appearance of a term that violates parity [22], very large value and negative sign of cosmological constant [29, 30], issues with power counting renormalisation of the propagation of the scalar mode [13, 31]. Some of those problems might be solved by performing an analytic continuation of the parameters of the theory [20].

In the original Hořava formulation it is assumed via the so called detailed balance condition that a potential part of the action is derived from the so-called superpotential, which limits the big number of its terms and corresponding independent couplings. Another imposed condition is the demand of projectability, used in a standard cosmology. It requires that lapse function N depends only on time N = N(t). It might seem that this condition is too strict but on the other hand it seems that non-projectable version of Hořava gravity results in serious strong coupling problem ([28]) and does not possess a valid GR limit at IR. However, some authors [29, 32] claim the opposite, proposing adding additional terms to the superpotential (not to the action thus still keeping detailed balance or eventually softly breaking it) and relaxing projectability. Nonetheless, subsequent works demonstrated that it caused problems with the scalar mode power-counting renormalizability. One of the simplest models with the detailed balance condition relaxed is the Sotiriou-Visser-Weinfurtner (SVW) generalisation [22]. This version of HL gravity assumes a gravitational action containing terms not only quadratic in curvature, but also cubic ones, what was suggested already in [12, 19]. This model still maintains the projectability condition. Generalised Friedmann equations obtained from varying such an action contain not only a dark radiation term $\sim 1/a^4$ but also terms scaling as $\sim 1/a^6$ term. These new terms, although negligible at large values of a, become dominating at small ones and might modify or cancel bounce solutions. Specifically, as it has the opposite sign than the $1/a^4$ term, it may compensate the dark radiation term at small scales and result in singular solutions. Similar scenario arrives in the HL gravity with the softly broken detailed balance condition and negative spacial curvature [33].

Nonetheless, the issue of the initial singularity still remains one of the key questions of early Universe cosmology. The possibility that it might be avoided in a modified gravity and replaced by a bounce is a very promising feature. In this review we are going to present result of the research [14, 15] performed via phase portrait techniques, on occurrence and stability of the bounce in two simplest formulation of HL cosmology: original one with imposed detailed balance condition and SVW formulation relaxing this condition. As additional terms in analogs of Friedman equations are proportional the curvature parameter $K = \{-1, 0, 1\}$ only non-flat cosmologies with $K = \pm 1$ allow the existence of a bounce and existence of non-singular solutions.

In [14] matter sector was described in terms of a scalar field with a potential given by a quadratic power of that field. More general approach and easier for fitting with observational data is the hydrodynamical approach used in [15] where matter sector is described in terms of density ρ and pressure p. In the latter work it was assumed that wproviding the relation between density and pressure in the equation of state, is constant, which is at some level an idealisation and simplification. At the moment we do not have the history of the HL universe constructed in a similar way as in the standard Λ CDM model, where we have phases and epochs containing different matter or radiation sectors. Therefore, as we still have limited understanding on the physical aspects of the theory and its parameters, current research rather describes different analytical possibilities, not some exact physical solutions.

This paper is organised as follows: We first give a brief overview of HL cosmology in both scenarios under consideration in Section II. In Section III the possibility of bounce in both formulations is discussed. Section IV contains derivation and description of the phase portraits of the HL cosmology with imposed condition of detailed balance, while in the section V this condition is released. Section VI contains summary of results on possibility of a bounce in HL cosmology. In Section VII we discuss limitation of the underlying theory.

II. HOŘAVA-LIFSHITZ COSMOLOGY

The main obstacle in quantising gravity is that general relativity in its classical formulation is non-renormalisable. This might be visualised by expanding some quantity \mathcal{F} with respect to the gravitational constant [27] as follows:

$$\mathcal{F} = \sum_{n=0}^{\infty} a_n \left(G_N E^2 \right)^n.$$
⁽¹⁾

Here E is the energy of the system, a_n denotes a numerical coefficient and G_N is the gravitational coupling constant. Therefore, $E^2 \ge G^{-1}$ and the expansion above diverges. Consequently, as demonstrated, general relativity is not perturbatively renormalisable in the high energy regimes.

There has been many researches pointing out that the ultraviolet behaviour of general relativity might be improved by including higher-order derivatives in the standard gravitational metric. The latter is the Einstein-Hilbert action:

$$S = \int d^4x \sqrt{g}R,\tag{2}$$

where d^4x denotes volume element of the space-time, g is its metric matrix' determinant and R is a scalar curvature. Including higher order terms of the derivatives of the metric provides a following action:

$$S = \int d^4x \sqrt{g} (R + f(R_{\mu\nu}R^{\mu\nu})).$$
 (3)

The additional terms, containing different derivatives of R, $R_{\mu\nu}$ etc. change the graviton propagator from $1/k^2$ into $1/(k^2 - G_N k^4)$ [7, 8]. The propagator part proportional to k^{-4} cancels the ultraviolet divergence. However, the resulting theory has time derivatives of $\mathcal{O} > 2$ and therefore non-unitary. Moreover, it possesses a spin-2 ghost with a non-zero mass [27] and derived form that action field equations are of the fourth order.

The novel idea of Hořava [7] was to construct a higher-order theory of gravity breaking the Lorentz invariance in the ultraviolet. In his theory only the *spatial* derivatives are of $\mathcal{O} > 2$ which evaded the ghost. However, it is demanded that any theory of gravity theory should be consistent with all current experiments which have not detected any significant violation of Lorentz invariance. Thus it is necessary to restore the Lorentz invariance in the infrared limit. In order to overcome this problem Hořava proposed an anisotropic scaling of space and time in high UV energies, which is known as Lifshitz scaling. In a 4-dimensional spacetime this scaling takes the form:

$$t \to b^{-z}t, \, x^i \to b^{-1}x^i, \tag{4}$$

here i = 1, 2, 3 and z is a critical exponent. Lorentz invariance is restored when z = 1, but the power-counting renormalisability demands $z \ge 3$ [27], usually z = 3 is assumed. Therefore, the resulting theory is called Hořava-Lifshitz (HL) gravity. Lorentz symmetry is here broken down to transformations $t \to \xi_0(t), x^i \to \xi^i(t, x^k)$, preserving the spatial diffeomorphisms unlike full space time diffeomorphisms invariance of GR. Thus such a theory acquires a symmetry preserving a space-time foliation [7, 27], where on each constant time hypersurface there are allowed arbitrary changes of the spatial coordinates.

Preservation of a space-time foliation and anisotropic scaling between time and space and time introduces the ADM (1+3) decomposition of the spacetime. The standard ADM metrics in a preferred foliation and with (-+++) signature is following:

$$ds^{2} = -N^{2}dt^{2} + g_{ij}(dx^{i} + N^{i}dt)(dx^{j} + N^{j}dt).$$
(5)

The dynamics is now described in terms of the lapse function N, the shift vector N^i , and the spatial metric g_{ij} (i, j = 1, 2, 3). The most general action for such theory can be written as:

$$S = \int d^3x dt \ N \sqrt{g} \left[K^{ij} K_{ij} - \lambda K^2 - \mathcal{V}(g_{ij}) \right].$$
(6)

Here as usually g denotes the determinant of the spatial metric g_{ij} , λ is a dimensionless running coupling constant, \mathcal{V} is a potential term and K is a trace of the extrinsic curvature of the spatial 3-dimensional hypersurface K_{ij} :

$$K_{ij} = \frac{1}{2N} \left(\dot{g}_{ij} - \nabla_i N_j - \nabla_j N_i \right). \tag{7}$$

An overdot denotes a derivative with respect to the time coordinate t. The trace of K_{ij} is K. The potential \mathcal{V} is invariant only under three-dimensional diffeomorphisms [25] and depends only on the spatial metric and its (spatial) derivatives. Thus it contains only operators constructed from the spatial metric g_{ij} and of dimension 4 and 6.

A. Detailed Balance

As the action (6) is very complicated Hořava [7, 26, 29] proposed to impose additional condition, the so-called detailed balance. It assumes that the \mathcal{V} could be derived from a superpotential W [7, 26, 29]:

$$\mathcal{V} = E^{ij} \mathcal{G}_{ijkl} E^{kl}, \quad E^{ij} = \frac{1}{\sqrt{g}} \frac{\delta W}{\delta g_{ij}}, \tag{8}$$

and

$$\mathcal{G}^{ijkl} = \frac{1}{2} \left(g^{ik} g^{jl} + g^{il} g^{jk} \right) - \lambda g^{ij} g^{kl}.$$
(9)

By carrying out an analytic continuation (e.g. [20]) of two constant parameters ω and μ we obtain he action for Hořava-Lifshitz gravity in the detailed balance condition [26] and reads as

$$S_{db} = \int dt \, d^3x \sqrt{g} N \Biggl[\frac{2}{\kappa^2} \left(K_{ij} K^{ij} - \lambda K^2 \right) + \frac{\kappa^2}{2\omega^4} C_{ij} C^{ij} - \frac{\kappa^2 \mu}{2\omega^2} \frac{\epsilon^{ijk}}{\sqrt{g}} R_{il} \nabla_j R_k^l + \frac{\kappa^2 \mu^2}{8} R_{ij} R^{ij} + \frac{\kappa^2 \mu^2}{8(1 - 3\lambda)} \left(\frac{1 - 4\lambda}{4} R^2 + \Lambda R - 3\Lambda^2 \right) \Biggr],$$
(10)

where C^{ij} is the Cotton tensor:

$$C^{ij} = \epsilon^{ikl} \nabla_k \left(R^j_{\ l} - \frac{1}{4} R \delta^j_l \right), \tag{11}$$

 ϵ^{ikl} denotes the totally antisymmetric tensor. The parameters κ, ω , and μ arriving in the theory have mass dimension respectively -1, 0, and 1. The analytic continuation mentioned above reads as $\mu \mapsto i\mu$ and $\omega^2 \mapsto -i\omega^2$ and it enables obtaining the positive values of the cosmological constant Λ as predicted by current observational results in the low energy regime.

It is expected that action (10) reduces to the Einstein-Hilbert one in the IR limit of the theory. This is possible if the speed of light c and gravitational constant G correspond to HL parameters as follows:

$$G = \frac{\kappa^2}{32\pi c}, \quad c = \frac{\kappa^4 \mu^2 \Lambda}{8(3\lambda - 1)^2}.$$
(12)

The coupling constant λ present in the action (10) is dimensionless. It runs with energy and flows to the three infrared (IR) fixed points ([7]): $\lambda = 1/3$, $\lambda = 1$ or $\lambda = \infty$. However, some of those values seem unphysical, in the region $1 > \lambda > 1/3$ there appear ghost instabilities in the IR limit of the theory [34]. The attempt to solve this problem [20] resulted in instabilities re-emerging at the other energy region, in UV. Thus the most physically interesting case is the regime $\lambda \ge 1$ that allows for a possible flow towards GR, where $\lambda = 1$. Region $\lambda \le 1/3$ on the other hand is disconnected from $\lambda = 1$, therefore cannot be included in realistic physical considerations.

In order to obtain a cosmological model it is necessary to populate the universe with matter (and radiation). The simplest method would be to model the matter sector by assuming it is described by a scalar field φ with a quadratic potential $V(\varphi) = \frac{1}{2}m^2\varphi^2$. However, a more realistic approach is to apply a hydrodynamic approximation where matter is described by two quantities p and ρ , which are respectively pressure and energy density and fullil the continuity equation $\dot{\rho} + 3H(\rho + p) = 0$.

To derive equations of HL cosmology one uses the projectability condition N = N(t) [7] and the spatial part of the metrics being the standard FLRW line element: $g_{ij} = a^2(t)\gamma_{ij}$, $N_i = 0$, where γ_{ij} denotes a maximally symmetric metric with constant curvature:

$$\gamma_{ij} \mathrm{d}x^i \mathrm{d}x^j = \frac{\mathrm{d}r^2}{1 - Kr^2} + r^2 (\mathrm{d}\theta^2 + \sin^2\theta \mathrm{d}\varphi^2),\tag{13}$$

values $K = \{-1, 0, 1\}$ correspond respectively to closed, flat, and open Universe. This background metric implies that

$$C_{ij} = 0, \qquad R_{ij} = \frac{2K}{a^2} g_{ij}, \qquad K_{ij} = \frac{H}{N} g_{ij}, \qquad (14)$$

where $H \equiv \dot{a}/a$ denotes the Hubble parameter.

On this background the gravitational action (10) take the following form :

$$S_{\rm FRW} = \int dt \, d^3x \, N a^3 \, \left\{ \frac{3(1-3\lambda)}{2\kappa^2} \frac{H^2}{N^2} + \frac{3\kappa^2 \mu^2 \Lambda}{4(1-3\lambda)} \left(\frac{K}{a^2} - \frac{\Lambda}{3}\right) - \frac{\kappa^2 \mu^2}{8(1-3\lambda)} \frac{K^2}{a^4} \right\}.$$
 (15)

In order to obtain equations of motion on a cosmological background one needs to vary the action (15) with respect to N and a. Only after that and the lapse can be set to one: N = 1 and terms with density ρ and pressure p are added. This procedure provides the analogs to the Friedmann equations for the projectable Hořava-Lifshitz cosmology with imposed the detailed-balance condition:

$$H^{2} = \frac{\kappa^{2} \rho}{6(3\lambda - 1)} \pm \frac{\kappa^{4} \mu^{2}}{8(3\lambda - 1)^{2}} \left(\frac{K\Lambda}{a^{2}} - \frac{\Lambda^{2}}{2} - \frac{K^{2}}{2a^{4}}\right),$$
(16)

$$\dot{H} = -\frac{\kappa^2(\rho+p)}{4(3\lambda-1)} \mp \frac{\kappa^4\mu^2}{8(3\lambda-1)^2} \left(\frac{K\Lambda}{a^2} + \frac{K^2}{4a^4}\right),\tag{17}$$

together with the continuity equation:

$$\dot{\rho} + 3H(\rho + p) = 0.$$
 (18)

In the equations above there are two signs before the terms with Λ , namely the upper one corresponds the $\Lambda < 0$ case, the lower one describes the analytic continuation $\mu \mapsto i\mu$ providing a positive Λ .

Some terms in the above equations which scale as a^{-4} are similar to the dark energy expressions therefore parameters: energy density ρ_{de} and pressure density p_{de} are interpreted as dark energy parameters:

$$\rho_{de}|_{db} := \frac{3\kappa^2 \mu^2 K^2}{8(3\lambda - 1)a^4} + \frac{3\kappa^2 \mu^2 \Lambda^2}{8(3\lambda - 1)},\tag{19}$$

$$p_{de}|_{db} := \frac{\kappa^2 \mu^2 K^2}{8(3\lambda - 1)a^4} - \frac{3\kappa^2 \mu^2 \Lambda^2}{8(3\lambda - 1)}.$$
(20)

We require that eqs (16) and (17) coincide with the standard Friedmann equations. Thus, we can identify the following:

$$c = \frac{\kappa^2 \mu}{4} \sqrt{\frac{\Lambda}{1-3\lambda}}, \quad G = \frac{\kappa^2 c}{32\pi}, \quad \Lambda_E = -\frac{3\kappa^4 \mu^2}{3\lambda - 1} \frac{\Lambda^2}{32} = \frac{3c^2}{2}\Lambda, \tag{21}$$

respectively, as well as $\mu^2 \Lambda = 1/32\pi^2 G^2$ and $\lambda = 1$ (which is an IR fixed point). We demand a real value of the speed of light *c*, therefore the cosmological constant Λ has to be negative for $\lambda > 1/3$ and positive for $\lambda < 1/3$. In order to obtain a positive cosmological constant Λ , as suggested by observations, it is necessary to perform in (10) an analytic complex continuation of constant parameters μ and ω as follows $\mu \mapsto i\mu$ and $\omega^2 \mapsto -i\omega^2$. On the level of equations for Hořava-Lifshitz cosmology varying λ -parameter in the range $[1, \infty)$ results in the running of the speed of light, but does not change the structure of the equations (16) and (17).

When we substitute the equation of state $p = w\rho$ and the above expressions linking physical constants and HL parameters to (16) and (17) we obtain the following equations:

$$H^{2} = \frac{2}{3\lambda - 1} \left[\frac{\rho}{3} \pm \left(\frac{\Lambda_{E}}{3} - \frac{K}{a^{2}} + \frac{3}{4\Lambda_{E}} \frac{K^{2}}{a^{4}} \right) \right]$$
(22)

$$\dot{H} = \frac{2}{3\lambda - 1} \left[-\frac{(1+w)}{2} \rho \pm \left(\frac{K}{a^2} - \frac{3}{2\Lambda_E} \frac{K^2}{a^4} \right) \right].$$
(23)

B. Beyond detailed balance

The gravitational action (10) contains terms up to quadratic in the curvature. However, a more general renormalizable theory could also contain cubic terms and there is not *a priori* reason to keep only quadratic terms ([12, 19, 27]). Thus Sotiriou, Visser and Weinfurtner ([26]) built a projectable theory as the original Hořava theory, but without imposing the detailed balance condition in the action.

This formulation led to Friedmann equations with an additional term $\sim 1/a^6$, moreover with additional and uncoupled coefficients:

$$H^{2} = \frac{2}{(3\lambda - 1)} \left(\frac{\rho}{3} + \sigma_{1} + \sigma_{2} \frac{K}{a^{2}} + \sigma_{3} \frac{K^{2}}{a^{4}} + \sigma_{4} \frac{K}{a^{6}} \right),$$
(24)

$$\dot{H} = \frac{2}{(3\lambda - 1)} \left(-\frac{p}{2} - \frac{\rho}{2} - \sigma_2 \frac{K}{a^2} - 2\sigma_3 \frac{K^2}{a^4} - 3\sigma_4 \frac{K}{a^6} \right).$$
(25)

In order to coincide with the Friedmann equations in the IR limit $\lambda = 1$ and for large a, when terms proportional to $1/a^4$ and to $1/a^6$ become negligibly small, one has to set $\sigma_1 = \Lambda_E/3$ and $\sigma_2 = -1$. However, values of constants σ_3 , σ_4 are at this stage arbitrary. This way we obtain the following equations:

$$H^{2} = \frac{2}{(3\lambda - 1)} \left(\frac{\rho}{3} + \frac{\Lambda_{E}}{3} - \frac{K}{a^{2}} + \sigma_{3} \frac{K^{2}}{a^{4}} + \sigma_{4} \frac{K}{a^{6}} \right),$$
(26)

$$\dot{H} = \frac{2}{(3\lambda - 1)} \left(-\frac{\rho(1 + w)}{2} + \frac{K}{a^2} - 2\sigma_3 \frac{K^2}{a^4} - 3\sigma_4 \frac{K}{a^6} \right),\tag{27}$$

We can observe new terms in the above analogs of Friedmann equations, proportional to $1/a^6$. They mimic stiff matter, such that $\rho = p$ (w = 1) which scales similarly $\rho_{\text{stiff}} \sim 1/a^6$). These terms are negligibly small at large scales, but may play a significant role at small values of a scale parameter, thus changing the dynamics of the Universe around initial singularity or a bounce.

III. EXISTENCE OF BOUNCE

Hořava-Lifshitz cosmological equations contain additional terms proportional a^{-4} (DB) and to a^{-6} (BDB) that introduce the possibility of a cosmological bounce, namely a scenario in which contraction of the universe stops and reverse to expansion (or in the opposite direction). In a DB scenario, from the form of eq. (16) it follows that it is possible that H = 0. When this condition is fulfilled at some monet of time the realisation of the bounce is possible (but not necessary, for that we also need $\dot{H} \neq 0$. In the case $\lambda = 1$ [12], the bounce may happen in non-empty Universe equipped with matter, at the critical time t_* , $a = a_*$, when the critical energy density reaches the following value:

$$\rho = \rho_* = \frac{12}{\kappa^2} \left(\frac{K}{a_*^2} + \frac{\Lambda_E}{3} + \frac{\kappa^4 \mu^2}{64} \frac{K^2}{a_*^4} \right), \tag{28}$$

This value is determined by the values of couplings κ and μ .

Additionally, a continuity equation implies that at the bounce $\dot{H} > 0$. Therefore, when the condition H = 0 is fulfilled we also have the sufficient condition for existence of a bounce $\dot{H} \neq 0$. As $\dot{H} > 0$ is is only possible a transition from a contracting to an expanding phase, but not the reverse. Moreover, there is another condition for a realisation bounce [24] that requires that $(\frac{p}{12} - p) > 0$ and the energy density of regular matter scales less fast than dark matter terms.

Near the bounce so for small a the dominating terms in the Friedmann equations (22) and (23) are the terms scaling as a^{-4} , while others terms become insignificant. Particularly, H^2 , \dot{H} and ρ scale as $a^{-3(1+w)}$, where w is a constant parameter in the equation of state $p = w\rho$. Subsequently, if $w > -\frac{1}{3}$ the density term dominates over the curvature term $\sim 1/a^2$.

In the BDB scenario bounce might happed at the critical density:

$$\rho_* = -\Lambda_E + 3\frac{K}{a_*^2} - 3\frac{\sigma_3 K^2}{a_*^4} - \frac{3\sigma_4 K}{a_*^6}.$$
(29)

For flat universe and positive cosmological constant bounce is not positive as resulting critical density becomes negative.

IV. BOUNCE STABILITY IN THE DETAILED BALANCE FORMULATION

We are mainly interested in the possibility of appearing of a bounce which could be given by dynamics of variables a and H. From eq. (22) we might determine ρ and then insert its formula into (23). This way we obtain two systems, one containing the formula for density and its derivative via the continuity equation (22), but still dependent on a and H. The second system is independent and consist of two equations describing the evolution of a and H.

Specifically, eq. (22) provided a following expression for ρ :

$$\rho = \frac{3(3\lambda - 1)}{2} H^2 \mp \left(\Lambda_E - 3\frac{K}{a^2} + \frac{9}{4\Lambda_E}\frac{K^2}{a^4}\right).$$
(30)

This expression substituted in (23) results in

$$\dot{H} = \frac{\pm 1}{3\lambda - 1} \left[(1+w)\Lambda_E - (3w+1)\frac{K}{a^2} + \frac{3(3w-1)}{4\Lambda_E}\frac{K^2}{a^4} \right] - \frac{3}{2}(1+w)H^2.$$
(31)

Adding the the definition of the Hubble parameter:

$$\dot{a} = aH,\tag{32}$$

we have a two dimensional dynamical system.

The set of equations (31-32) is difficult to solve analytically. However, we are interested non in detailed solutions but in the qualitative analysis. In purpose of that we use the method of the phase portraits, where we search for critical points and analyse their character. These points are locations where the derivatives of all the dynamic variables, In our case when the r.h.s. of (31-32), vanish. What we obtain are the only points where phase trajectories could start, end or intersect. Moreover, they can also appear in infinity. In this case a suitable coordinate transformation, the so called Poincaré projection, is used that projects the complete phase space onto a compact region. The nature of these points, both finite and infinite, is given by the properties of the Jacobian matrix of the linearized equations at those points. All that information provides a qualitative analysis of the dynamical system.

The method of finding critical points consists of setting all right-hand-sides of dynamical equations to zero, thus finding points where derivative of dynamical variables vanish. In case of two equations (31)-(32) the corresponding solutions are two following P_1 and P_2 in phase-space (a, H):

$$P_1: a^2 = \frac{3K}{2\Lambda_E}, \ H = 0, \tag{33}$$

$$P_2: a^2 = \frac{(3w-1)K}{(1+w)2\Lambda_E}, \ H = 0.$$
(34)

These two points exist when the values of a obtained via square root of the expression on the right hand side of the above equations are real and nonnegative. Thus the point P_1 exists if $K/\Lambda_E > 0$, if we assume a positive cosmological constant therefore only for K > 0. Point P_2 exists when w > 1/3 and $K/\Lambda_E > 0$ or w < 1/3 and $K/\Lambda_E < 0$. Thus we have two critical points existing if the parameter of state w > 1/3. Moreover, they are both finite, unless w = -1, when P_2 blows to infinity. As mentioned above, due to $\dot{H} > 0$ both points represent a bouncing solution.

In order to complete the analysis, the stability properties of the critical points needed. They are determined by the eigenvalues of the Jacobian A of the system (31)-(32). Eigenvalues of A with non-zero real parts different from zero point to hyperbolic points. They include sources (unstable) with positive real parts, saddle for real parts of opposite sign and sinks (stable) corresponding to negative real parts. Critical points at which all the eigenvalues have real parts different from zero are called hyperbolic. Among them one can distinguish sources (unstable) with positive real parts, saddles with real parts of different sign and sinks (stable) for negative real parts. If at least one eigenvalue has a real part equal to zero it is then called a non-hyperbolic critical point. For such points it is not possible to obtain conclusive information about the stability from the Jacobian matrix and other tools like e.g. numerical simulation [35] should be then used.

In the case of (31)-(32) the eigenvalues of A at P_1 are both imaginary and it is a center for all the values of the parameters. The character of P_2 is more complicated and depend on the values of Λ_E , K and w. Thus P_2 is a center when $K/\Lambda_E < 0$ and $-1 \le w < 1/3$, with a special subcase for w = -1 that being so it becomes a linear center, so a center with only one eigenvector. Otherwise it becomes a saddle, thus without a bouncing possibility.

To have a full picture of the dynamics of the Universe also the information about critical points that occurring at infinity is necessary. For this purpose the so called Poincaré projection [36, 37] is used. It projects the whole infinite phase space (a, H) onto a compact region. Specifically, we introduce the new coordinates (\tilde{a}, \tilde{H}) which written in polar coordinates $r, \phi: \tilde{a} = r \cos \phi$ and $\tilde{H} = r \sin \phi$. Moreover:

$$a = \frac{r}{1 - r} \cos \phi, \tag{35}$$

$$H = \frac{r}{1-r}\sin\phi,\tag{36}$$

It is also necessary to rescale the time parameter t, which take infinite values, introducing the new time parameter T in a similar way, i.e. dT = dt/(1 - r). In such coordinates the phase space is now compactified to a sphere of radius one and its interior. Here infinity corresponds to r = 1. We have to keep in mind that a scale factor a may take only nonnegative values, so actually a semi-sphere.

This procedure provides the dynamical equations in terms of r, ϕ and their derivatives with respect to new time T. At the surface of the sphere, so limit r = 1 there are 3 solutions $P_3 = (1,0)$, $P_4 = (1, \pi/2)$, $P_6 = (1, -\pi/2)$ written in polar coordinates (r, ϕ) . These critical points are now hyperbolic unless w = -1, resulting in P_4 and P_6 being respectively a repelling and an attracting node. For w = -1 points P_4 and P_6 are non-hyperbolic and numerical simulations provide that they are saddles and also ends of a separatrice.

The numerical phase portraits are presented at fig. 1, which contains the deformed phase space, scaled to fit on the compactified sphere. We observe that bounce scenarios are only possible when one of the critical points P_1 and P_2 exist and is a center. Then we have closed orbits around them and the Universe might go through eternal cycles of expansion and collapse, connected by a bounce of a finite size, expansion etc. However, the point P_1 describes less physical bouncing solution with the density $\rho = 0$. More interesting case is when P_2 is a center, as a density ρ is non-zero at that point. The special case of w = -1 provides a third bounce scenario is around the linear center P_2 located now at ∞ . In this case the universe begins in a static infinite state as H = 0 $a = \infty$, then contracts to a finite size and rebounces to a static infinite universe.




FIG. 1: Projected phase space of HL universe [15] in DB condition. On the left a case with $K\Lambda_E > 0$ and w > 1/3, on the right $K/\Lambda_E < 0$ and -1 < w < 1/3.

V. BOUNCE STABILITY IN THE BEYOND DETAILED BALANCE FORMULATION

In the Sotiriou, Visser and Weinfurtner formulation the generalised Friedmann equations (26)-(27) contain additional terms $\sim 1/a^6$ and uncoupled coefficients.

Solving eq. (26) for ρ provides:

$$\rho = 3\frac{(3\lambda - 1)}{2}H^2 - \Lambda_E - 3\frac{K}{a^2} - 3\frac{\sigma_3 K^2}{a^4} - \frac{3\sigma_4 K}{a^6}.$$
(37)

Substituting this expression on ρ into ((27)) and using the equation of state $p = w\rho$ results in

$$\dot{H} = \frac{2}{3\lambda - 1} \left(\frac{\Lambda_E(1+w)}{2} - \frac{K(1+3w)}{2a^2} + \frac{\sigma_3(-1+3w)K^2}{2a^4} + \frac{3\sigma_4(1+w)K}{2a^6} \right) - \frac{3(1+w)}{2}H^2.$$
(38)

As in the DB case, supplementing the above equation with the definition of the Hubble parameter provides the two dimensional dynamical system.

Again we search for critical points where \dot{a} and H these points fulfil H = 0 and obtain the following condition:

$$\Lambda_E(1+w)a^6 - K(1+3w)a^4 + \sigma_3(-1+3w)K^2a^2 + 3\sigma_4(-1+w)K = 0$$
(39)

It is a bicubic equation, which in general possesses quite complicated solutions but might be simplified in two special cases. Namely when w = -1 describing the equation of state of the cosmological constant, and in case of radiation described by w = 1/3. Besides these two cases critical points of the system (32) and (38) have following coordinates $(a_x, 0)$. Here a_x^2 is a root of the cubic equation:

$$\Lambda_E(1+w)x^3 - K(1+3w)x^2 + \sigma_3(-1+3w)K^2x + 3\sigma_4(-1+w)K = 0.$$
(40)

Such an equation might have zero, one, two or three real solutions depending on the sign of its discriminant. Moreover, if they exist they are either always stable or always unstable depending on the sign of $K/(3\lambda - 1)$. Their character depends on the values of a_x , Λ_E , σ_3 and σ_4 . The most significant feature of oscillating (and bouncing) solutions in the SVW formulation is the existence of two centres, with a saddle between them (three finite critical points) for some values of parameters. In a more realistic situation, that includes dynamical change of state parameter, it would be possible to go from one oscillating bouncing solution to another.

In order to study stability properties of infinite critical points one again has to perform the Poincarè transformation. It leads to the similar results as in detailed balance scenario. Points at infinity are transformed to the sphere r = 1. Two points at $\phi = \pi/2$ and at $-\pi/2$ are respectively repelling and attracting node, respectively. The point at $\phi = 0$ is non-hyperbolic.

Figure 2. shows the example of the phase space of system with three finite critical points. Here points S_1 and S_3 are centres and a point S_2 is a saddle.




FIG. 2: Projected phase space of the HL universe in beyond detailed balance formulation with 3 critical points existing[15].

VI. DISCUSSION

This paper reviews the research performed the cosmological bounce in different formulations of projectable versions of Hořava-Lifshitz gravity, with and without detailed balance condition. The analogs of the Friedmann equations in both this models contain a term scaling as $1/a^4$ and similar to dark radiation. That additional term enables that fthe Hubble parameter might be H = 0 at some moment of time. This is a necessary condition for the realisation of the bounce while an additional condition $\dot{H} \neq 0$ makes it sufficient. In the Sotiriou, Visser and Weinfurtner model there is an additional term $1/a^6$ in the analogs of Friedmann equations. This term is of arbitrary sign, so it can enhance the possibility of a bounce or cancel it.

The biggest difference between the detailed balance theory and its breaking arrives for the small values of a scale parameter a as the SVW gravity term $1/a^6$ plays role only for the small values of a and becomes insignificant for the bigger ones. This difference is visible in phase portraits of both theories and number of potential bouncing solutions. In the original Hořava formulation there exists one bouncing solutions for all values of parameters but it corresponds to density $\rho = 0$. For non zero $\rho = 0$ there might be a bouncing solution if $K/\Lambda_E < 0$ and $-1 \le w < 1/3$, for other values of parameters a bounce is not possible.

The SVW HL cosmology is a bit more complicated as there are additional terms in the analogs of Friedmann equations. There exist bouncing solutions for some values of parameters of the theory, however a range of parameters that lead only to singular solutions is wider than in the detailed balance scenario. One very interesting special case includes two centres, with a saddle between them (corresponding to three finite critical points). If one takes into account dynamical change of state parameter, which is much more realistic scenario, it might be possible to go from one oscillating solution to another bouncing solution. The problem is that the existence of such solutions depends on the values of coupling constants σ_3 and σ_4 and their physical interpretation still remains an open question.

Moreover, in both these formulations, bouncing non-singular solutions exist only in case of a non-flat universe $K \neq 0$. Otherwise the bouncing solutions become singular.

VII. CONCLUSIONS

The obtained cosmological results presented here are promising and suggest there is a possibility to replace the initial cosmological singularity of GR by finite bouncing solutions. However, one must also consider that there are many problems and contradicting statements in the different formulations and extensions of HL-type theories. Aside from that aspect there are also observational bounds on the existence of the Hořava-Lifshitz gravity and the values of its constants and parameters.

At present, HL-type theories, including the original one and its extensions, are not yet ruled out by observational

data. However, there now are tight bounds on some parameters of the theory [38] from the binary neutron star merger GW170817 [39]. Therefore, it is possible that further observational data might either rule out some specific scenarios or the whole model. It is also possible that some agreement with observations could provide a better justification for additional theoretical research as it is still hoped that HL gravity could offer a promising cosmological scenario without initial singularity and solve some shortcomings of classical GR, like non-renormalisability and thus problems with quantisation.

There are several observational bounds on different regions of the Hořava-Lifshitz framework, *e.g.* using data from binary pulsars [40, 41], using general cosmological data [16] and also bounds in the context of dark energy [42]. In the context of dark matter and dark energy there are also bounds on generally Lorentz violation [9, 43]. There is also quite recent research performed in the effective field theory formalism [18] of the extension HL gravity [25]. However, this analysis is reduced to a flat background spacetime, which limits the overall number of parameters.

In our papers [44, 45] we have placed new bounds on parameters of Hořava-Lifshitz cosmology, in its projectable version with and without imposing detailed balance condition. We found very interesting results on spatial curvature. Namely the original HL model is well fitted with a positive non-zero spatial curvature with accuracy to more than 3σ , whereas when we relaxed the detailed balance condition we obtained again positive non-zero spatial curvature at 1σ accuracy. As this analysis also included BAO's, therefore there is needed further investigation of the curvature parameter which could possibly finally exclude some of the HL models. Anyway those results seem to be fascinating in view of future observation and also somehow demonstrate why an analysis limited to zero spatial curvature is somehow limited. Still non-singular bouncing solutions in HL universe appear only for non-zero spatial curvature, so these two topics are related.

We have to take into account that most obtained bounds on the parameters of the HL cosmology are similar to those in Λ CDM model, except the non-zero curvature parameter. Of course, the Λ CDM model has still fewer parameters and from this point of view should be preferred; it also fits the data well. However, one has to bear in mind also the theoretical aspects of Hořava gravity which make it a good candidate for an ultra violet complete theory of gravity. There also several implication like the possible resolution of the initial cosmological singularity, so there are still many reasons to keep investigating this model and its extensions.

- [1] M. Bojowald and G. Date, Phys. Rev. Lett. 92, 071302 (2004).
- [2] F. Quevedo, (2016), arXiv:1612.01569 [hep-th].
- [3] F. Girelli, F. Hinterleitner, and S. Major, SIGMA 8, 098 (2012), arXiv:1210.1485.
- [4] H. Bergeron, E. Czuchry, J.-P. Gazeau, P. Małkiewicz, and W. Piechocki, Phys. Rev. D 92, 124018 (2015), arXiv:1501.07871 [gr-qc].
- [5] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, Phys. Lett. B 429, 263 (1998), arXiv:hep-ph/9803315.
- [6] G. Dvali, Fortsch. Phys. 58, 528 (2010), arXiv:0706.2050 [hep-th].
- [7] P. Hořava, Phys. Rev. D79, 084008 (2009), arXiv:0901.3775 [hep-th].
- [8] P. Hořava and C. M. Melby-Thompson, Phys. Rev. D 82, 064027 (2010), arXiv:1007.2410 [hep-th].
- [9] B. Audren, D. Blas, M. M. Ivanov, J. Lesgourgues, and S. Sibiryakov, JCAP 1503, 016 (2015), arXiv:1410.6514 [astro-ph.CO].
- [10] D. Blas, O. Pujolas, and S. Sibiryakov, JHEP **10**, 029 (2009), arXiv:0906.3046 [hep-th].
- [11] D. Blas and H. Sanctuary, Phys. Rev. D84, 064004 (2011), arXiv:1105.5149 [gr-qc].
- [12] G. Calcagni, JHEP 09, 112 (2009), arXiv:0904.0829 [hep-th].
- [13] M. Colombo, A. E. Gümrükçüoğlu, and T. P. Sotiriou, Phys. Rev. D92, 064037 (2015), arXiv:1503.07544 [hep-th].
- [14] E. Czuchry, Class. Quant. Grav. 28, 085011 (2011), arXiv:0911.3891 [hep-th].
- [15] E. Czuchry, Class. Quant. Grav. 28, 125013 (2011), arXiv:1008.3410 [hep-th].
- [16] S. Dutta and E. N. Saridakis, JCAP 1001, 013 (2010), arXiv:0911.1435 [hep-th].
- [17] S. Dutta and E. N. Saridakis, JCAP 1005, 013 (2010), arXiv:1002.3373 [hep-th].
- [18] N. Frusciante, M. Raveri, D. Vernieri, B. Hu, and A. Silvestri, Phys. Dark Univ. 13, 7 (2016), arXiv:1508.01787 [astroph.CO].
- [19] E. Kiritsis and G. Kofinas, Nucl. Phys. B821, 467 (2009), arXiv:0904.1334 [hep-th].
- [20] H. Lü, J. Mei, and C. N. Pope, Phys. Rev. Lett. 103, 091301 (2009), arXiv:0904.1595 [hep-th].
- [21] E. N. Saridakis, Eur. Phys. J. C67, 229 (2010), arXiv:0905.3532 [hep-th].
- [22] T. P. Sotiriou, M. Visser, and S. Weinfurtner, JHEP 10, 033 (2009), arXiv:0905.2798 [hep-th].
- [23] S. Mukohyama, Class. Quant. Grav. 27, 223101 (2010), arXiv:1007.5199 [hep-th].
- [24] R. Brandenberger, Physical Review D 80 (2009), 10.1103/physrevd.80.043516.
- [25] D. Blas, O. Pujolas, and S. Sibiryakov, Phys. Rev. Lett. 104, 181302 (2010), arXiv:0909.3525 [hep-th].
- [26] T. P. Sotiriou, Proceedings, 14th Conference on Recent developments in gravity (NEB 14): Ioannina, Greece, June 8-11, 2010, J. Phys. Conf. Ser. 283, 012034 (2011), arXiv:1010.3218 [hep-th].
- [27] A. Wang, Int. J. Mod. Phys. **D26**, 1730014 (2017), arXiv:1701.06087 [gr-qc].

- [28] C. Charmousis, G. Niz, A. Padilla, and P. M. Saffin, JHEP 08, 070 (2009), arXiv:0905.2579 [hep-th].
- [29] D. Vernieri and T. P. Sotiriou, Phys. Rev. D85, 064003 (2012), arXiv:1112.3385 [hep-th].
- [30] C. Appignani, R. Casadio, and S. Shankaranarayanan, JCAP 04, 006 (2010), arXiv:0907.3121 [hep-th].
- [31] D. Vernieri, Phys. Rev. D 91, 124029 (2015), arXiv:1502.06607 [hep-th].
- [32] M. Pospelov and Y. Shang, Phys. Rev. D 85, 105001 (2012), arXiv:1010.5249 [hep-th].
- [33] E. J. Son and W. Kim, Mod. Phys. Lett. A 26, 719 (2011), arXiv:1007.5371 [gr-qc].
- [34] C. Bogdanos and E. N. Saridakis, Class. Quant. Grav. 27, 075005 (2010), arXiv:0907.1636 [hep-th].
- [35] D. K. Arrowsmith, C. M. Place, C. Place, et al., An introduction to dynamical systems (Cambridge university press, 1990).
- [36] G. N. Felder, A. Frolov, and L. Kofman, Classical and Quantum Gravity 19, 2983 (2002).
- [37] G. Felder, A. Frolov, L. Kofman, and A. Linde, Physical Review D 66, 023507 (2002).
- [38] A. Emir Gümrükçüoğlu, M. Saravani, and T. P. Sotiriou, Phys. Rev. D 97, 024032, arXiv:1711.08845 [gr-qc].
- [39] B. P. Abbott et al. (LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL), Astrophys. J. Lett. 848, L13 (2017), arXiv:1710.05834 [astro-ph.HE].
- [40] K. Yagi, D. Blas, E. Barausse, and N. Yunes, Phys. Rev. D 89, 084067 (2014), [Erratum: Phys.Rev.D 90, 069902 (2014), Erratum: Phys.Rev.D 90, 069901 (2014)], arXiv:1311.7144 [gr-qc].
- [41] K. Yagi, D. Blas, N. Yunes, and E. Barausse, Phys. Rev. Lett. **112**, 161101 (2014), arXiv:1307.6219 [gr-qc].
- [42] M.-i. Park, JCAP **01**, 001 (2010), arXiv:0906.4275 [hep-th].
- [43] B. Audren, D. Blas, J. Lesgourgues, and S. Sibiryakov, JCAP 08, 039 (2013), arXiv:1305.0009 [astro-ph.CO].
- [44] N. A. Nilsson and E. Czuchry, Phys. Dark Univ., 100253 (2018), [Phys. Dark Univ.C23,100253(2019)], arXiv:1803.03615 [gr-qc].
- [45] N. A. Nilsson and E. Czuchry, In preparation.