

Proof to count bound state nodes in supersymmetric quantum mechanics

A.Aynbund^{1,*} and V.V.Kiselev^{1,2,†}

¹*Landau Phystech-School, Moscow Institute of Physics and Technology,
Russia, 141701, Moscow Region, Dolgoprudny, Institutsky 9*

²*State Research Center of the Russian Federation “Institute for High
Energy Physics” of National Research Centre “Kurchatov Institute”,
Russia, 142281, Moscow Region, Protvino, Nauki 1*

A normalizable static supersymmetric bound ground state annihilated by the super-generators has got zero number of internal nodes in the framework of one-dimensional supersymmetric quantum mechanics. The super-generator transformations between excited super-partner bound states as combined with the standard technique of Wronskian provides an elegant and self-sufficient way to derive the equality of internal nodes amount to the number of consequent excitation.

I. INTRODUCTION

Counting an amount of internal nodes in wave functions of bound states is known as an oscillation theorem. The theorem relates a sequence number of excited state in the order of energy enlarge to number of its nodes. An ordinary proof, say, in classical book by F. Berezin and M. Shubin [1] is a set of theorems, lemmas, consequences and prepositions in Sturm–Liouville theory of differential equations. Such the way is rigorous and it looks like just straightforwardly mathematical, while physical explanations merely give some qualitative arguments but not enough to be improvements (see, for instance, [2]). In occasion there is a method based on coherent quantum structure, that provides an elegant and physically clear way to derive the oscillation theorem by means of supersymmetry in one dimensional quantum mechanics. This method is built on a construction introduced by M. Crum [3] and

*Electronic address: aynbund.asya@phystech.edu

†Electronic address: Valery.Kiselev@ihep.ru; kiselev.vv@mipt.ru

incorporated to the supersymmetry by E. Witten [4] as reviewed in [5–7].

In Section II basic notions and constructions are described to give the most important properties of supersymmetric Hamiltonian. Section III deals with a ground state, and we derive that the ground state has got no internal nodes. Section IV is devoted to a step to the level next to the ground state to prove that the number of nodes increases to the unit exactly. In Conclusion we summarize the results.

II. MACHINERY OF SUPERSYMMETRIC QUANTUM MECHANICS

The theory under study is one-dimensional stationary Schrödinger equation. Suppose that Ψ_0 is a ground state wave function normalised in a Hilbert space of quantum states for a Hamiltonian \hat{H}_1 . As it is well known, function Ψ_0 can be set to be real since it has got nodes at borders. Let the Hamiltonian have got several excitations of the ground state, and the excited levels are counted in the order of energy enlargement as $\Psi_k^{(1)}$. Following M. Crum [3] and M. Darboux [8], one constructs a partner Hamiltonian \hat{H}_2 , that possesses a set of bound states wave functions $\Psi_k^{(2)}$.

Define two linear differential operators A^- and A^+ , which are hermitian conjugated each to other:

$$A^- = -\frac{d}{dx} + \frac{\Psi'_0}{\Psi_0}, \quad A^+ = \frac{d}{dx} + \frac{\Psi'_0}{\Psi_0}.$$

Introduce operators of super-generators Q and $\bar{Q} \equiv Q^\dagger$ by E.Witten [4]

$$Q = \frac{\hbar}{\sqrt{2m}} \begin{pmatrix} 0 & 0 \\ A^- & 0 \end{pmatrix} \text{ and } \bar{Q} = \frac{\hbar}{\sqrt{2m}} \begin{pmatrix} 0 & A^+ \\ 0 & 0 \end{pmatrix} \text{ acting on } \begin{pmatrix} \Psi^{(1)} \\ \Psi^{(2)} \end{pmatrix}$$

where the column refers to wave functions of super-partner Hamiltonians \hat{H}_1 and \hat{H}_2 , respectively.

A supersymmetric Hamiltonian \mathcal{H} is constructed as

$$\mathcal{H} = \{Q, \bar{Q}\}.$$

Consider properties of Hamiltonian \mathcal{H} and supercharge operators. So, the super-generators are nilpotent, i. e. $Q^2 = \bar{Q}^2 = 0$ and they commute with Hamiltonian

$$[Q, \mathcal{H}] = 0.$$

Then the definition of super-partner Hamiltonians in terms of matrix \mathcal{H} reads off

$$\mathcal{H} = \frac{\hbar^2}{2m} \begin{pmatrix} A^+A^- & 0 \\ 0 & A^-A^+ \end{pmatrix} = \begin{pmatrix} \hat{H}_1 - E_0 & 0 \\ 0 & \hat{H}_2 - E_0 \end{pmatrix}.$$

Two Schrödinger equations under study are

$$\hat{H}_1 \Psi_k^{(1)} = E_k \Psi_k^{(1)}, \quad \hat{H}_2 \Psi_k^{(2)} = E_k \Psi_k^{(2)},$$

where the eigenvalues of Hamiltonians coincide. Indeed, according to the connection between \hat{H}_1 and operators A^- and A^+ the action of A^+A^- to the wave function $\Psi_k^{(1)}$ gives

$$\frac{\hbar^2}{2m} A^+A^- \Psi_k^{(1)} = (\hat{H}_1 - E_0) \Psi_k^{(1)} = (E_k - E_0) \Psi_k^{(1)}.$$

Acting by A^- results in the eigen state equation

$$A^-A^+(A^- \Psi_k^{(1)}) = (E_k - E_0)(A^- \Psi_k^{(1)}) \quad \Rightarrow \quad (\hat{H}_2 - E_0)(A^- \Psi_k^{(1)}) = (E_k - E_0)(A^- \Psi_k^{(1)}),$$

determining the wave function $\Psi_k^{(2)} \stackrel{\text{def}}{=} A^- \Psi_k^{(1)}$ as the bound state solution for Hamiltonian \hat{H}_2 . Note that the superscripts of wave functions point to means the super-partner Hamiltonians, while the subscript denotes the number of excitation with the same energy E_k for both Hamiltonians.

III. GROUND STATE IN THE DISCRETE SPECTRUM

The ground state wave function Ψ_0 satisfies the equation of ground state:

$$A^- \Psi_0 = 0.$$

Hence, any other function satisfying the same equation is a normalised solution for the ground state and it is proportional to Ψ_0 , since the bound state energies are not degenerate.

Let us show that $A^-|\Psi_0| = 0$. Therefore, $\Psi_0 = \text{const} \cdot |\Psi_0|$, and generically one can put the constant to be positive, while $\Psi_0 > 0$ at any internal point except borders. It means that Ψ_0 is nodeless.

Indeed,

$$\tilde{\Psi} \stackrel{\text{def}}{=} |\Psi_0| = \Psi_0 \text{sign}(\Psi_0),$$

while the action by A^- straightforwardly gives

$$\begin{aligned}
A^- \tilde{\Psi} &= \left(-\frac{d}{dx} + \frac{\Psi'_0}{\Psi_0} \right) \Psi_0 \text{sign}(\Psi_0) = -\frac{d}{dx} \left(\Psi_0 \text{sign}(\Psi_0) \right) + \Psi'_0 \text{sign}(\Psi_0) = \\
&= \cancel{-\text{sign}(\Psi_0)\Psi_0} - \Psi_0 \frac{d}{dx} \left(\text{sign}(\Psi_0) \right) + \cancel{\Psi'_0 \text{sign}(\Psi_0)} = -\Psi_0 \frac{d}{dx} \left(\text{sign}(\Psi_0) \right) = \\
&= -\sum_j \Psi_0(x_j) 2 \delta(x_j) \text{sign}(\Psi'_0(x_j)) \equiv 0, \quad (1)
\end{aligned}$$

where x_j are positions of possible nodes of Ψ_0 , if exist, while $\Psi'_0(x_j) \neq 0$ because

$$\Psi'(x_j) = \Psi(x_j) = 0$$

would lead to $\Psi(x) \equiv 0$ as a solution of Schrödinger equation.

Thus, the super-generators provide us with the short proof of statement that the ground state wave function Ψ_0 is nodeless, so we consider it to be positive anywhere except border points.

IV. COUNTING THE NODES

As we plot in Fig. 1, there are two sets of states for the super-partner Hamiltonians with energy E_1 . We show that one can transform one column of states to another by making use the linear operator A^- for moving from left to right (for example, if one acts on Ψ_0 , we will get 0) and the operator A^+ for moving back. These operators conserve the energy level.




FIG. 1: The ground state and the excited level of Hamiltonian \hat{H}_1 (left), the ground state of Hamiltonian \hat{H}_2 (right) with energies E_0 and E_1 , respectively.




FIG. 2: Node of the excited state and the direction of the derivative in it.

The figure points that the ground states for both super-partner Hamiltonians are nodeless, while more energy means that the wave function has more nodes as well known fact. So, with this explicit scheme for the super-partner system we are going to prove that the first excitation of ground state for \hat{H}_1 has got exactly single internal node, while further excitations have got numbers of nodes equal to its excitation numbers.

In previous section we proved so-called *base case* of mathematical induction method. The next point is to prove the induction step. Acting on the first excited wave function of Hamiltonian \hat{H}_2 and by revealing the operator by definition we get

$$A^+\Psi_1^{(2)} = \Psi_1^{(2)} + \frac{\Psi_0'}{\Psi_0}\Psi_1^{(2)} = \frac{(\Psi_0\Psi_1^{(2)})'}{\Psi_0} = \langle \text{or} \rangle = A^+(A^-\Psi_1^{(1)}) = (E_1 - E_0)\Psi_1^{(1)}.$$

Then, multiplying by Ψ_0 one can see the derivative of Wronskian for Ψ_0 and $\Psi_1^{(1)}$:

$$(\Psi_0\Psi_1^{(2)})' = (E_1 - E_0)\Psi_1^{(1)}\Psi_0 = -W'_{\Psi_0\Psi_1^{(1)}} \Rightarrow -W_{\Psi_0\Psi_1^{(1)}} = \Psi_0\Psi_1^{(2)}.$$

Therefore, we come to the following expression:

$$\Psi_0'\Psi_1^{(1)} - \Psi_0\Psi_1'^{(1)} = \Psi_0\Psi_1^{(2)}.$$

Consider the first term on the left side. We can see that as E_1 is greater than E_0 there are strictly more nodes for the function of excited level. Hence, there are at least one node which we will denote x_1 . Rewriting the previous equation at the node we get

$$\Psi_0(x_1)\Psi_1^{(2)}(x_1) = -\Psi_0(x_1)\Psi_1'^{(1)}(x_1),$$

where all of $\Psi_0(x_1)$ and $\Psi_1^{(2)}(x_1)$ are positive by construction for the ground states (remember, that $\Psi_1^{(2)}$ is the ground state for Hamiltonian \hat{H}^2).

Therefore, $\Psi_1'^{(1)}(x_1) < 0$, as it is pictured in Fig. 2, for any node x_1 . It means that in all of nodes the derivative has to be negative. Taking into account the continuity of wave

function, we have to conclude that there is no more than one change of sign $\Psi_1^{(1)}$, and the wave function of the first excitation $\Psi_1^{(1)}$ has got the single internal node, exactly.

Further steps of mathematical induction are quite transparent.

V. CONCLUSION

In the present paper we have presented the new original modern proof for counting the internal nodes of bound states wave functions instead of Sturm–Liouville theory by making use the elegant mathematical formalism of Supersymmetric Quantum Mechanics. It could be incorporated into textbooks on modern quantum mechanics in order to provide some acquaintance with up-to-date supersymmetric concepts and mathematical methods essential in nowadays science.

-
- [1] F. A. Berezin, M. Shubin, *The Schrödinger Equation*. (Springer Science & Business Media, 2012).
 - [2] M. Moriconi, American Journal of Physics **75(3)**, 284 (2007) [arXiv:quant-ph/0702260].
 - [3] M. M. Crum, Quart. J. Math. Oxford Ser. 2 **6**, 1216 (1955) [arXiv:physics/9908019].
 - [4] E. Witten, Nucl. Phys. B **188**, 513 (1981).
 - [5] C. V. Sukumar, Phys. A: Math. Gen. **18**, 2917 (1985).
 - [6] L. E. Gendenshtein, I. V Krive, Soviet Physics Uspekhi. **28**, 645 (1985).
 - [7] F. Cooper, F. Khare, U. Sukhatme, Physics Reports **25**, 267 (1995) [arXiv:hep-th/9405029].
 - [8] M. G. Darboux, Comptes Rendus Acad. Sci. Paris **94**, 1456 (1882) [arXiv:physics/9908003].