

TRACE METHODS FOR COHOCHSCHILD HOMOLOGY

SARAH KLANDERMAN AND MAXIMILIEN PÉROUX

ABSTRACT. Hochschild homology is a classical invariant of rings that plays an important role because of its connection to algebraic K -theory via the Dennis trace. At level zero, the Dennis trace is induced by the Hattori–Stallings trace. In this paper, we introduce new algebraic K -theories of coalgebras and obtain coalgebraic refinements of the Hattori–Stallings trace that connect these algebraic K -theories to coHochschild homology—the invariant analogous to Hochschild homology but for coalgebras. We employ bicategorical methods of Ponto to show that coHochschild homology is a shadow. Consequently, we obtain that coHochschild homology is Morita–Takeuchi invariant.

CONTENTS

1. Introduction	1
2. Coalgebraic variations of the Hattori–Stallings trace	8
3. Bicategories of bicomodules	16
4. CoHochschild homology as a shadow	20
5. Dualizable bicomodules	25
6. Bicategorical traces	29
7. Morita–Takeuchi invariance	31
Appendix A. Homotopy theory of connective bicomodules	33
References	40

1. INTRODUCTION

Trace methods provide an essential resource for accessing computations in algebraic K -theory. Such computations are crucial as they can shed light on problems across different fields of mathematics, including algebra, topology, and number theory. Algebraic K -theory is constructed using tools from stable homotopy theory, and consequently is challenging to compute. The first example of a trace map is the *Dennis trace* $K_*(R) \rightarrow \mathrm{HH}_*(R)$ which relates K -theory to a more computable invariant called *Hochschild homology* (HH). Bökstedt defined a closer approximation to K -theory by introducing a topological refinement of HH called *Topological Hochschild homology* (THH). The resulting Dennis trace $K(R) \rightarrow \mathrm{THH}(R)$ and its refinements have been utilized to great effect in computations, for example [BHM93, Dun97, NS18].

CoHochschild homology (coHH) is the invariant of coalgebras analogous to Hochschild homology that was first introduced by Doi [Doi81]. Based on previous work by Hess–Parent–Scott in chain complexes [HPS09], Hess–Shipley extended Doi’s definition to the setting of stable homotopy theory and introduce *topological coHochschild homology* (coTHH) as the spectral refinement of coHH in [HS21]. Given a connected space X , there is an equivalence of spectra

$$\mathrm{THH}(\Sigma_+^\infty \Omega X) \simeq \Sigma_+^\infty \mathcal{L}X,$$

where $\mathcal{L}X$ is the free loop space on X , see [Lod98, 7.3.11] or [NS18, IV.3.3]. Building on work of Malkiewich [Mal17, 2.22], Hess–Shipley show that further connectivity conditions on X gives an equivalence of spectra

$$\mathrm{coTHH}(\Sigma_+^\infty X) \simeq \Sigma_+^\infty \mathcal{L}X,$$

2020 *Mathematics Subject Classification*. Primary: 16E40, 16T15, 18N10, 19A99, 55P43, 55U15. Secondary: 16D90, 18F30, 18M70, 18N60, 57T30.

Key words and phrases. coalgebra, comodule, coHochschild homology, trace, bicategory, K -theory, Morita–Takeuchi equivalence.

see [HS21, 3.7]. These equivalences of spectra have led to new computations of the homology of the free loop space [BGS22], using a *coBökstedt spectral sequence* for coTHH constructed in [BGH⁺18]. This spectral sequence is analogous to the Bökstedt spectral sequence for THH defined by Angeltveit–Rognes [AR05].

Given the prominent role played by trace methods in computations of algebraic K -theory, it is natural to extend trace methods to coTHH . Combining the equivalences above with the Dennis trace, Hess–Shipley obtained a new trace to coTHH [HS21] for X simply connected:

$$K(\Sigma_+^\infty \Omega X) \longrightarrow \mathrm{THH}(\Sigma_+^\infty \Omega X) \simeq \mathrm{coTHH}(\Sigma_+^\infty X).$$

Additionally, work of the second author and Bayındır [BP23, 4.4] shows that applying Spanier–Whitehead duality on the Dennis trace determines a pairing $K(C^\vee) \wedge \mathrm{coTHH}(C) \longrightarrow \mathbb{S}$, that can be reformulated as

$$K(C^\vee) \longrightarrow \mathrm{coTHH}(C)^\vee, \quad (1.1)$$

where C^\vee is the Spanier–Whitehead dual of a coalgebra spectrum C with some finiteness conditions.

However, both of the coTHH traces mentioned above are not internal to the coalgebraic setting; they are directly induced by combining the usual Dennis trace with an identification of coTHH with THH . Moreover, there has been no clear candidate of an algebraic K -theory for coalgebras without using a duality on coalgebras into rings. In this paper, as a first step towards trace methods in the coalgebraic setting, we introduce new algebraic K -theories for coalgebras and construct analogues of the Dennis trace on $\mathrm{coHochschild}$ homology at level zero. Our results lay the foundational framework for building higher level traces on $\mathrm{coHochschild}$ homology as well as spectral refinements to coTHH .

Review of trace methods. If A is a connective ring spectrum, the Dennis trace $K(A) \rightarrow \mathrm{THH}(A)$ at level zero is precisely a homomorphism $K_0(\pi_0(A)) \rightarrow \mathrm{HH}_0(\pi_0(A))$. Therefore, it is induced by a homomorphism $K_0(R) \rightarrow \mathrm{HH}_0(R)$ for a (discrete) ring R . In order to obtain trace maps in the coalgebraic setting, we must first determine what happens at level zero for $\mathrm{coHochschild}$ homology. The homomorphism $K_0(R) \rightarrow \mathrm{HH}_0(R)$ is induced by the usual trace on square matrices in linear algebra. Essential properties of the trace are that it is additive and cyclic:

$$\mathrm{tr}(A + B) = \mathrm{tr}(A) + \mathrm{tr}(B), \quad \mathrm{tr}(AB) = \mathrm{tr}(BA),$$

and these properties make the trace easily computable. Abstracting these properties, the definition of the trace can be extended from matrices to any endomorphism on a dualizable object in a symmetric monoidal category [DP80, PS14]. A more subtle situation arises in non-symmetric contexts. For instance, the relative tensor product of bimodules over a non-commutative ring R does not define a symmetric monoidal structure. Nevertheless, in [Hat65, Sta65], Hattori and Stallings define the trace of an R -linear endomorphism on a finitely generated projective right R -module M as a homomorphism $\mathrm{End}_R(M) \rightarrow R/[R, R]$, called the *Hattori–Stallings trace*. The group $R/[R, R] = \mathrm{HH}_0(R)$ is the universal home of any trace map $\mathcal{M}_n(R) \rightarrow A$ valued in an abelian group A . The Hattori–Stallings trace extends the universal trace $R \rightarrow R/[R, R]$ to endomorphisms of finitely generated and projective R -modules, defining the induced homomorphism $K_0(R) \rightarrow \mathrm{HH}_0(R)$ by evaluating the Hattori–Stallings trace of the identity on such modules.

Ponto generalized this approach by introducing the notion of traces in *shadowed bicategories* [Pon10]. Essentially, a bicategory is a generalization of a monoidal category in which we allow a change of base ring, and a shadow is a choice of a symmetry in this monoidal structure. A prototypical example is the bicategory of rings and bimodules, with horizontal composition given by the (underived) relative tensor product and with shadow the zeroth Hochschild homology HH_0 . The key observation is that given (not-necessarily commutative) rings R and S , an (R, S) -bimodule M and a (S, R) -bimodule N , the relative tensor products $M \otimes_S N$ and $N \otimes_R M$ are *not* isomorphic. In fact, they do not even have the same algebraic structure, as the former is an (R, R) -bimodule while the latter is an (S, S) -bimodule. However, the two bimodules become equivalent after applying (relative) Hochschild homology:

$$\mathrm{HH}_0(R, M \otimes_S N) \cong \mathrm{HH}_0(S, N \otimes_R M).$$

Considering instead the derived tensor product shows that higher levels of Hochschild homology HH_* are also shadows, and replacing rings by ring spectra shows that THH is a shadow. In fact, THH is the universal cyclic invariant, and the universal shadow in a sense made precise in [HR21a]. Every shadowed bicategory comes equipped with a notion of a bicategorical trace that is cyclic, and the bicategorical trace on dualizable bimodules recovers the Hattori–Stallings trace in the underived case.

Results. We dualize the construction of trace maps in the coalgebraic setting, using the framework of bicategorical shadows. In doing so, we are confronted with several obstacles. For instance, given coalgebras C and D , the *cotensor product* $M \square_D N$ of a (C, D) -bicomodule M with a (D, C) -bicomodule N need not to be a (C, C) -bicomodule. Even worse, the cotensor product is not even associative in general when considering coalgebras over \mathbb{Z} [GP87, IV.2.5]. To remedy this, we assume that \mathbb{k} is a ring of global dimension zero, as this guarantees we have a nice associative relative cotensor product on bicomodules.

Theorem 1.2 (Theorem 4.5). *Let \mathbb{k} be a commutative ring with global dimension zero. The zeroth co-Hochschild homology coHH_0 defines a shadow on the bicategory of \mathbb{k} -coalgebras and bicomodules with their relative cotensor products.*

In particular, given \mathbb{k} -coalgebras C and D , a (C, D) -bicomodule M and a (D, C) -bicomodule N , there is a natural isomorphism of \mathbb{k} -modules:

$$\mathrm{coHH}_0(M \square_D N, C) \cong \mathrm{coHH}_0(N \square_C M, D).$$

Given a right dualizable (C, D) -bicomodule M and an endomorphism $f: M \rightarrow M$ of (C, D) -bicomodules, the fact that coHH_0 is a shadow induces a \mathbb{k} -linear homomorphism:

$$\mathrm{tr}_C^D(f): \mathrm{coHH}_0(C) \longrightarrow \mathrm{coHH}_0(D), \quad (1.3)$$

that is cyclic (Proposition 6.2). Considering either $C = \mathbb{k}$ or $D = \mathbb{k}$ leads to two particularly interesting traces.

Case 1: $D = \mathbb{k}$ in (1.3). In this case, if M is a finitely cogenerated and injective left C -comodule (see Definition 2.8), then it is right dualizable as a (C, \mathbb{k}) -bicomodule. Given a C -colinear endomorphism f of M , the assignment $f \mapsto \mathrm{tr}_C^{\mathbb{k}}(f)$ determines a \mathbb{k} -linear homomorphism that we call the *Hattori–Stallings cotrace*:

$${}_C\mathrm{End}(M) \rightarrow \mathrm{Hom}_{\mathbb{k}}(\mathrm{coHH}_0(C), \mathbb{k}). \quad (1.4)$$

We give a precise formula in Definition 2.13, and we subsequently denote $\mathrm{cotr}(f) = \mathrm{tr}_C^{\mathbb{k}}(f)$. The Hattori–Stallings cotrace is an extension of a universal cotrace $\mathrm{coHH}_0(C) \hookrightarrow C$ considered by Quillen in [Qui88, Qui89], see Remark 2.15.

The class of finitely cogenerated and injective left C -comodules is the dual analogue of the class of finitely generated and projective right R -modules considered when computing the algebraic K -theory $K(R)$ of a ring R . Therefore, this analogy leads us to introduce the following construction.

Definition 1.5. Let \mathbb{k} be a commutative ring with global dimension zero and let C be a \mathbb{k} -coalgebra. Let ${}_C\mathrm{CoMod}^{\mathrm{fc}, \mathrm{inj}}$ denote the category of finitely cogenerated and injective left C -comodules, viewed as a Waldhausen category whose cofibrations are monomorphisms and whose weak equivalences are isomorphisms. Let $K^c(C)$ denote the resulting algebraic K -theory spectrum associated to the Waldhausen category ${}_C\mathrm{CoMod}^{\mathrm{fc}, \mathrm{inj}}$, as in [Wal85] or [Wei13, IV.8.5].

In this paper, we obtain several properties for $K^c(C)$. First, if C is a cocommutative coalgebra, then $K^c(C)$ is a ring spectrum (Proposition 2.20). Second, if C is a conilpotent coaugmented coalgebra, then $K_0^c(C) \cong \mathbb{Z}$ (Proposition 2.19). This result applies in particular to the polynomial ring $\mathbb{k}[x]$ or the group ring $\mathbb{k}\Gamma$ of a group Γ seen as Hopf algebras.

Just as the Hattori–Stallings trace $\mathrm{End}_R(M) \rightarrow \mathrm{HH}_0(R)$ led to a homomorphism $K_0(R) \rightarrow \mathrm{HH}_0(R)$ of abelian groups by considering the trace on the identity, the Hattori–Stallings cotrace ${}_C\mathrm{End}(M) \rightarrow \mathrm{coHH}_0(C)^*$ on the identity (i.e., the *corank*) induces the following map.

Theorem 1.6 (Theorem 2.16). *Let \mathbb{k} be a commutative ring with global dimension zero, and let C be a \mathbb{k} -coalgebra. The Hattori–Stallings corank induces a homomorphism of abelian groups:*

$$K_0^c(C) \longrightarrow \mathrm{coHH}^0(C),$$

where $\mathrm{coHH}^0(C) = \mathrm{Hom}_{\mathbb{k}}(\mathrm{coHH}_0(C), \mathbb{k})$.

The cotrace $K_0^c(C) \rightarrow \mathrm{coHH}^0(C)$ is analogous to the Dennis trace $K_0(C^*) \rightarrow \mathrm{coHH}^0(C)$ obtained when C is finitely generated as \mathbb{k} -modules, a particular case of (1.1) from [BP23]. We view Theorem 1.6 as a first step in constructing the Dennis trace entirely in the coalgebraic setting, with no finiteness assumption on C .

Case 2: $C = \mathbb{k}$ in (1.3). In this case, relabelling D as C , if M is a right C -comodule that is finitely generated as a \mathbb{k} -module, then it is right dualizable as a (\mathbb{k}, C) -bicomodule. Given a C -colinear endomorphism f on M , the assignment $f \mapsto \mathrm{tr}_{\mathbb{k}}^C(f)(1)$ determines a \mathbb{k} -linear homomorphism that we call the *colinear Hattori–Stallings trace*:

$$\mathrm{End}_C(M) \longrightarrow \mathrm{coHH}_0(C). \quad (1.7)$$

In Section 2.3, we show that this map is a coalgebraic refinement of the twisted Hattori–Stallings trace on a \mathbb{k} -linear homomorphism $M \rightarrow M \otimes C$ and give a precise formula in Definition 2.23.

The class of right C -comodules that are finitely generated as \mathbb{k} -modules is similar to classes that have been considered before in the literature. For instance, given a space X , its A -theory $A(X)$ is defined to be the K -theory of the Waldhausen category of homotopy finite retractive spaces over X with cofibrations and weak equivalences created in spaces [Wal85]. For any generalized reduced homology \mathcal{E}_* , let $A(X; \mathcal{E}_*)$ be the Waldhausen K -theory of the same category with the same cofibrations, but replacing the usual weak equivalences of spaces by \mathcal{E}_* -equivalences. Hess–Shipley show in [HS16, 1.3] that $A(X; \mathcal{E}_*)$ is equivalent to the K -theory of the Waldhausen category $(\mathrm{CoMod}_{X_+}^{\mathrm{hf}})_{\mathcal{E}_*}$ of homotopically finite X_+ -comodules in pointed spaces, with cofibrations all monomorphisms and weak equivalences \mathcal{E}_* -equivalences. Moreover, in [Pér24a, 1.2], for $\mathcal{E} = H\mathbb{Q}_*$ and X a simply connected space, the second author showed that $A(X; H\mathbb{Q}_*) \simeq K(\mathrm{CoMod}_{C_*(X, \mathbb{Q})}^{\mathrm{perf}})$, where $\mathrm{CoMod}_{C_*(X, \mathbb{Q})}^{\mathrm{perf}}$ is the Waldhausen category of perfect chain complexes that are comodules over the singular chain complex $C_*(X, \mathbb{Q})$, with cofibrations all monomorphisms and weak equivalences quasi-isomorphisms. These examples suggest the following similar construction in our context.

Definition 1.8. Let \mathbb{k} be a commutative ring with global dimension zero, and let C be a \mathbb{k} -coalgebra. Let $\mathrm{CoMod}_C^{\mathrm{fg}}$ denote the category of right C -comodules that are finitely generated (and automatically projective) as \mathbb{k} -modules. It is a Waldhausen category whose cofibrations are monomorphisms and whose weak equivalences are isomorphisms. Let $K(\mathrm{CoMod}_C^{\mathrm{fg}})$ denote the resulting algebraic K -theory spectrum.

We show in Proposition 2.29 that this algebraic K -theory is similar to the G -theory of rings. A particular example of interest is the group ring $\mathbb{k}\Gamma$ for Γ a finite group, as $G(\mathbb{k}\Gamma)$ is the K -theory of representations of Γ in the category of \mathbb{k} -modules, also known as Swan theory [Swa70].

If X is a simply connected space, then we obtain equivalences $K(\Sigma_+^\infty \Omega X) \simeq A(X) \simeq K((\mathrm{CoMod}_{X_+}^{\mathrm{hf}})_{H\mathbb{Z}_*})$ by [HS16, 1.4]. Combining with $\mathrm{THH}(\Sigma_+^\infty \Omega X) \simeq \mathrm{coTHH}(\Sigma_+^\infty X)$ from [HS21], the Dennis trace can be reformulated as

$$K\left((\mathrm{CoMod}_{X_+}^{\mathrm{hf}})_{H\mathbb{Z}_*}\right) \longrightarrow \mathrm{coTHH}(\Sigma_+^\infty X).$$

The colinear Hattori–Stallings trace $\mathrm{End}_C(M) \rightarrow \mathrm{coHH}_0(C)$ on the identity (i.e., the rank) allows us to extend the trace above on their π_0 -group, removing the connectivity assumptions.

Theorem 1.9 (Theorem 2.27). *Let \mathbb{k} be a commutative ring with global dimension zero, and let C be a \mathbb{k} -coalgebra. The colinear Hattori–Stallings rank induces a homomorphism of abelian groups:*

$$K_0\left(\mathrm{CoMod}_C^{\mathrm{fg}}\right) \rightarrow \mathrm{coHH}_0(C).$$

In this paper, we also extend our shadow framework of Theorem 1.2 to the derived setting. Our approach dualizes the arguments which show Hochschild homology is a shadow [Pon10], using an adaptation of the so-called Dennis–Waldhausen Morita argument. However, we are again faced with challenges that we now discuss.

First, obtaining model structures on categories of comodules and coalgebras is a challenging problem [BHK⁺15, HKRS17, GKR20]. Even in the case when model structures do exist, they may not be well-behaved. For instance, given a nice enough monoidal model category, its associated module and algebra categories represent modules and algebras that are associative and unital up to higher homotopy, see for instance [Lur17, 4.1.8.4, 4.3.3.17]. However, the dual result for comodules and coalgebras may not even be true [PS19, Pér22b]. Nevertheless, the second author showed that there is a model structure on connective comodules over simply connected differential graded \mathbb{k} -coalgebras up to quasi-isomorphism representing homotopy coherent connective comodules in $H\mathbb{k}$ -spectra [Pér24b].

Second, constructing a derived version of the cotensor product is a difficult problem, but is necessary to obtain a shadow structure on derived bicomodules. The derived tensor product can be modeled by a two-sided

bar construction, and so for bicomodules, we would want a two-sided cobar construction to model a *derived cotensor* product. As we point out in Remark 3.11, topologists and algebraists have a different meaning for the cobar construction, and issues emerge in both constructions. The topological definition of cobar is well-behaved homotopically but usually does not preserve comodule structures, whereas the algebraic definition of cobar does preserve these structures but is not invariant under quasi-isomorphisms. To circumvent these issues, one could restrict to the context of simply connected differential graded coalgebras, which is the setting for Koszul duality, and mimic some of the arguments in algebra, as the second author does in [Pér24a, Pér24b]. However, there are still subtleties in this setting, such as totalization not commuting with the tensor product, which we resolve in this paper.

Theorem 1.10 (Theorem 4.9). *Let \mathbb{k} be a commutative ring with global dimension zero. CoHochschild homology coHH defines a shadow on the bicategory of simply connected differential graded \mathbb{k} -coalgebras and bicomodules with their derived relative cotensor product.*

One of the defining properties of K -theory and Hochschild homology is invariance under Morita equivalences — rings that have equivalent (possibly derived) categories of left modules. The dual notion for coalgebras and comodules is called *Morita–Takeuchi equivalence*. Despite the pathological nature of model categories of comodules discussed above, we are able to show that $\mathrm{coHochschild}$ homology is invariant under Morita–Takeuchi equivalence in the derived context using our shadow framework.

Theorem 1.11 (Theorem 7.4). *Let C and D be homotopically Morita–Takeuchi equivalent simply connected differential graded \mathbb{k} -coalgebras. Then there is an isomorphism:*

$$\mathrm{coHH}(C) \cong \mathrm{coHH}(D).$$

Vista. The present paper is the first step of a program to extend the Dennis trace $K(R) \rightarrow \mathrm{THH}(R)$ to the coalgebraic setting. Our results propose natural candidates for the algebraic K -theory of a coalgebra. The Hess–Shipley identification $K(\Sigma_+^\infty \Omega X) \simeq K((\mathrm{CoMod}_{X_+}^{\mathrm{hf}})_{H\mathbb{Z}_*})$ suggests that these new K -theories for coalgebras “deloop” the usual algebraic K -theories of rings. In fact, given any ring spectrum R , its bar construction BR is a coalgebra and modules over R are equivalent to certain comodules over BR under a Koszul duality [Heu20]. Therefore, obtaining computational tools to determine the algebraic K -theories of a coalgebra can shed new light on the usual K -theory of rings.

In forthcoming work with Agarwal and Mehrle, the second author intends to extend the traces of Theorems 1.6 and 1.9 to higher levels

$$K_n^c(C) \longrightarrow \mathrm{coHH}^n(C), \quad K_n\left(\mathrm{CoMod}_C^{\mathrm{fg}}\right) \rightarrow \mathrm{coHH}_n(C).$$

These higher traces are built by extending our notion of cotraces $V \rightarrow \mathcal{M}_n^c(C)$ on *co-matrices* introduced in this paper.

Additionally, in forthcoming work with Brazelton, Calle, Chan, and Keenan, the second author intends to extend current methods in this paper to the bicategory of bicomodules over the coalgebra spectrum $\Sigma_+^\infty X$, with a derived cotensor product obtained entirely through ∞ -categorical methods. In particular, it is expected to obtain a trace $K(\mathrm{CoMod}_{\Sigma_+^\infty X}^{\mathrm{hf}}) \rightarrow \mathrm{coTHH}(\Sigma_+^\infty X)$ that is a delooping of the Dennis trace $K(\Sigma_+^\infty \Omega X) \rightarrow \mathrm{THH}(\Sigma_+^\infty \Omega X)$, extending Hess–Shipley results. Moreover, after applying the linearization $H\mathbb{k} \wedge -: \mathrm{CoMod}_{\Sigma_+^\infty X} \rightarrow \mathrm{CoMod}_{H\mathbb{k} \wedge X_+}$, we expect to recover the same shadow structure as Theorem 1.10 when X is simply connected.

Organization. We finish this section with notations and definitions that are used throughout this paper. In section 2, we construct the Hattori–Stallings cotrace and the colinear Hattori–Stallings trace using only algebraic methods and motivation. This section does not require any knowledge of bicategories, shadows, or homotopy theory. In order to be a self-contained paper, we will recall all necessary bicategorical definitions in later sections. We introduce the different bicategories of bicomodules in Section 3. Our key result, that $\mathrm{coHochschild}$ homology defines a bicategorical shadow, is proved in Section 4. We then explore the notion of duality in bicomodules in Section 5, which has been under-documented in the literature thus far. In Section 6, we provide a bicategorical description of the traces that appear in Section 2. In Section 7, we show how Morita–Takeuchi equivalences are part of the bicategorical structures. In Appendix A, we carefully detail how to represent the derived cotensor product of bicomodules and explore what fibrant bicomodules are.

Acknowledgements. We thank Jonathan Campbell and Cary Malkiewich for discussions that sparked some of the initial ideas of this paper, as well as Teena Gerhardt and Gabriel Angelini-Knoll. We also thank Haldun Özgür Bayındır, Thomas Brazelton, Maxine Calle, David Chan, Andres Mejia, Svetlana Makarova, Mona Merling, and Manuel Rivera for helpful and enlightening conversations. We also thank Maxine Calle and David Chan for their valuable feedback on earlier versions of this paper.

Notation. We begin by establishing notation that we use throughout this paper.

- (1) The letter \mathbb{k} shall always denote a commutative ring with global dimension zero, i.e., a finite product of fields, or said differently, a commutative semisimple ring. Every \mathbb{k} -module is projective and injective.
- (2) Let $\text{Mod}_{\mathbb{k}}$ denote the category of \mathbb{k} -modules. We write the relative tensor product $\otimes_{\mathbb{k}}$ as \otimes .
- (3) Given \mathbb{k} -modules M and C , and \mathbb{k} -linear homomorphisms $\Delta: C \rightarrow C \otimes C$ and $\rho: M \rightarrow M \otimes C$, we may employ the *Sweedler notation* in coalgebraic contexts:

$$\Delta(c) = \sum_{i=1}^n c_{(1)i} \otimes c_{(2)i} =: \sum_{(c)} c_{(1)} \otimes c_{(2)}, \text{ and } \rho(m) = \sum_{i=1}^n m_{(0)i} \otimes m_{(1)i} =: \sum_{(m)} m_{(0)} \otimes m_{(1)}.$$

- (4) Let $\text{Ch}_{\mathbb{k}}^{\geq 0}$ be the category of non-negative chain complexes of \mathbb{k} -modules (graded homologically). The category is endowed with a symmetric monoidal structure. The tensor product of two chain complexes X and Y is defined by

$$(X \otimes Y)_n = \bigoplus_{i+j=n} X_i \otimes_{\mathbb{k}} Y_j,$$

with differential given on homogeneous elements by

$$d(x \otimes y) = dx \otimes y + (-1)^{|x|} x \otimes dy.$$

We denote the tensor simply as \otimes . The monoidal unit is denoted \mathbb{k} , which is the chain complex \mathbb{k} concentrated in degree zero.

- (5) We write $\text{Hom}_{\mathbb{k}}(-, -)$ for the internal hom for both $\text{Mod}_{\mathbb{k}}$ and $\text{Ch}_{\mathbb{k}}^{\geq 0}$. We write $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$, the linear dual.
- (6) Given any symmetric monoidal category (\mathcal{C}, \otimes) , we write $\tau: X \otimes Y \xrightarrow{\cong} Y \otimes X$, the natural symmetric isomorphism.

General definitions. We recall here the categorical definitions of coalgebras and comodules, and their related concepts.

Definition 1.12. Let $(\mathcal{C}, \otimes, \mathbb{I})$ be a symmetric monoidal category. A *coalgebra* (C, Δ, ε) in \mathcal{C} consists of an object C in \mathcal{C} together with a *coassociative* comultiplication $\Delta: C \rightarrow C \otimes C$, such that the following diagram commutes:

$$\begin{array}{ccc} C & \xrightarrow{\Delta} & C \otimes C \\ \Delta \downarrow & & \downarrow \text{id}_C \otimes \Delta \\ C \otimes C & \xrightarrow{\Delta \otimes \text{id}_C} & C \otimes C \otimes C, \end{array}$$

and that admits a *counit* morphism $\varepsilon: C \rightarrow \mathbb{I}$ such that we have the following commutative diagram:

$$\begin{array}{ccccc} C \otimes C & \xrightarrow{\text{id}_C \otimes \varepsilon} & C \otimes \mathbb{I} \cong C & \cong \mathbb{I} \otimes C & \xleftarrow{\varepsilon \otimes \text{id}_C} C \otimes C \\ & \searrow \Delta & \parallel C & \nearrow \Delta & \\ & & & & \end{array}$$

The coalgebra is *cocommutative* if the following diagram commutes:

$$\begin{array}{ccc} C \otimes C & \xrightarrow[\cong]{\tau} & C \otimes C \\ \Delta \swarrow & & \searrow \Delta \\ & C & \end{array}$$

Given a coalgebra (C, Δ, ε) , we can define its opposite coalgebra C^{op} to be the coalgebra $(C, \tau \circ \Delta, \varepsilon)$. If C is cocommutative, then $C = C^{\text{op}}$.

Definition 1.13. Let $(\mathbf{C}, \otimes, \mathbb{I})$ be a symmetric monoidal category, and let (C, Δ, ε) be a coalgebra in \mathbf{C} . A *right C -comodule* (M, ρ) is an object M in \mathbf{C} together with a *coassociative* and *counital* right coaction morphism $\rho : M \rightarrow M \otimes C$ in \mathbf{C} , i.e., the following diagrams commute:

$$\begin{array}{ccc} M & \xrightarrow{\rho} & M \otimes C \\ \rho \downarrow & & \downarrow \rho \otimes \text{id}_C \\ M \otimes C & \xrightarrow{\text{id}_M \otimes \Delta} & M \otimes C \otimes C, \end{array} \quad \begin{array}{ccc} M & \xrightarrow{\rho} & M \otimes C \\ & \searrow & \downarrow \text{id}_M \otimes \varepsilon \\ & & M \otimes \mathbb{I} \\ & & \downarrow \cong \\ & & M. \end{array}$$

We can similarly define a *left C -comodule* (M, λ) . In fact, notice that a left C -comodule is a right C^{op} -comodule. Given two coalgebras C and D in \mathbf{C} , a (C, D) -bicomodule (M, λ, ρ) is a left C -comodule (M, λ) and a right D -comodule (M, ρ) such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{\rho} & M \otimes D \\ \lambda \downarrow & & \downarrow \lambda \otimes \text{id}_D \\ C \otimes M & \xrightarrow{\text{id}_C \otimes \rho} & C \otimes M \otimes D. \end{array}$$

A morphism $(M, \rho_M) \rightarrow (N, \rho_N)$ of right C -comodules is a morphism $f : M \rightarrow N$ in \mathbf{C} such that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \rho_M \downarrow & & \downarrow \rho_N \\ M \otimes C & \xrightarrow{f \otimes \text{id}_C} & N \otimes C. \end{array}$$

This morphism will be referred as a *(right) C -colinear homomorphism*. A (C, D) -bilinear homomorphism is a morphism that is both a left C -colinear map and a right D -colinear homomorphism. The notation $\text{CoMod}_C(\mathbf{C})$ will denote the category of right C -comodules in \mathbf{C} . Similarly, we use ${}_C\text{CoMod}(\mathbf{C})$ to denote the category of left C -comodules in \mathbf{C} , and ${}_C\text{CoMod}_D(\mathbf{C})$ for the category of (C, D) -bicomodules. Notice that we have isomorphisms of categories ${}_C\text{CoMod}_{\mathbb{I}}(\mathbf{C}) \cong {}_C\text{CoMod}(\mathbf{C})$ and ${}_{\mathbb{I}}\text{CoMod}_C(\mathbf{C}) \cong \text{CoMod}_C(\mathbf{C})$.

Definition 1.14. We say a coalgebra (C, Δ, ε) is *flat* in a symmetric monoidal category $(\mathbf{C}, \otimes, \mathbb{I})$ if the induced functor $C \otimes - : \mathbf{C} \rightarrow \mathbf{C}$ preserves equalizers when they exist.

Remark 1.15. Given a symmetric monoidal category $(\mathbf{C}, \otimes, \mathbb{I})$, and coalgebras C and D in \mathbf{C} , note that we obtain an isomorphism of categories

$${}_C\text{CoMod}_D(\mathbf{C}) \cong \text{CoMod}_{C^{\text{op}} \otimes D}(\mathbf{C}).$$

In particular, given a (C, D) -bicomodule (M, λ, ρ) , we obtain a right $(C^{\text{op}} \otimes D)$ -comodule via

$$M \xrightarrow{\rho} M \otimes D \xrightarrow{\lambda \otimes \text{id}_D} C \otimes M \otimes D \xrightarrow{\tau \otimes \text{id}_D} M \otimes C \otimes D.$$

Definition 1.16. Let $(\mathbf{C}, \otimes, \mathbb{I})$ be a symmetric monoidal category, let C, D , and E be coalgebras in \mathbf{C} , and let (M, λ_M, ρ_M) be a (D, C) -bicomodule and (N, λ_N, ρ_N) be a (C, E) -bicomodule. Define the *cotensor product* $M \square_C N$ to be the following equalizer in \mathbf{C} :

$$M \square_C N \longrightarrow M \otimes N \rightrightarrows M \otimes C \otimes N.$$

$\begin{array}{c} \xrightarrow{\rho_M \otimes \text{id}_N} \\ \xleftarrow{\text{id}_M \otimes \lambda_N} \end{array}$

If D and E are flat coalgebras, then $M \square_C N$ is a (D, E) -bicomodule via

$$\begin{array}{ccccc}
M \square_C N & \longrightarrow & M \otimes N & \xrightleftharpoons[\text{id}_M \otimes \lambda_N]{\rho_M \otimes \text{id}_N} & M \otimes C \otimes N. \\
\downarrow \exists! & & \downarrow \lambda_X \otimes \rho_Y & & \downarrow \lambda_M \otimes \text{id}_C \otimes \rho_N \\
D \otimes (M \square_C N) \otimes E & \longrightarrow & D \otimes M \otimes N \otimes E & \xrightleftharpoons[\text{id}_D \otimes \text{id}_M \otimes \lambda_N \otimes \text{id}_E]{\text{id}_D \otimes \rho_M \otimes \text{id}_N \otimes \text{id}_E} & D \otimes M \otimes C \otimes N \otimes E.
\end{array}$$

Definition 1.17. Given a symmetric monoidal category $(\mathbf{C}, \otimes, \mathbb{I})$, we say two coalgebras C and D in \mathbf{C} are *Morita–Takeuchi equivalent* if the categories ${}_C \text{CoMod}(\mathbf{C})$ and ${}_D \text{CoMod}(\mathbf{C})$ are equivalent.

Definition 1.18. Given a closed symmetric monoidal category $(\mathbf{C}, \otimes, \mathbb{I}, [-, -])$, and a coalgebra C , we can provide an enrichment of $\text{CoMod}_C(\mathbf{C})$ as follows. Given right C -comodules M and N , define $\text{Hom}_C(M, N)$ as the equalizer in \mathbf{C}

$$\text{Hom}_C(M, N) \longrightarrow [M, N] \rightrightarrows [M, N \otimes C],$$

where the first parallel map is induced by the coaction $N \rightarrow N \otimes C$ while the second is defined by forgetful-cofree adjointness. We denote $\text{End}_C(M) = \text{Hom}_C(M, M)$. We can similarly define ${}_C \text{Hom}(-, -)$ and ${}_C \text{End}(-)$ for left C -comodules.

Definition 1.19. Let \mathbf{A} be an abelian symmetric monoidal category with enough injective objects. Suppose C is a flat coalgebra in \mathbf{A} . Then ${}_C \text{CoMod}(\mathbf{A})$ also has enough injective objects. If M is a right C -comodule in \mathbf{A} , then the functor $M \square_C - : {}_C \text{CoMod}(\mathbf{A}) \rightarrow \mathbf{A}$ is left exact between abelian categories with enough injectives and thus we can right derive it. We denote by $\text{CoTor}_C^i(M, -)$ the i^{th} right derived functor. Similarly, if N is a left C -comodule, we can right derive the functor $- \square_C N : \text{CoMod}_C(\mathbf{A}) \rightarrow \mathbf{A}$ and obtain functors $\text{CoTor}_C^i(-, N)$. Per usual, we can check that $\text{CoTor}_C^i(M, N)$ is unambiguous.

2. COALGEBRAIC VARIATIONS OF THE HATTORI–STALLINGS TRACE

We introduce the Hattori–Stallings cotrace (Definition 2.13) and the colinear Hattori–Stallings trace (Definition 2.23) using only (co)algebraic methods. We obtain trace maps on K -theories: Theorem 2.16 and Theorem 2.27. We first begin by briefly recalling how the Hattori–Stallings trace is built so that the reader sees the analogies between the algebraic setting and coalgebraic setting.

2.1. The Hattori–Stallings trace. We recall [Hat65, Sta65]. Let R be a ring, not necessarily commutative, and let A be an abelian group. Denote the ring of square matrices of size n with coefficients in R by $\mathcal{M}_n(R)$.

Definition 2.1. A *trace function on R with values in A* is a collection \mathbb{Z} -linear homomorphisms denoted $T_n : \mathcal{M}_n(R) \rightarrow A$ for each $n \geq 1$, such that $T_n(MN) = T_m(NM)$, where we chose M to be a $n \times m$ -matrix, and N a $m \times n$ -matrix.

Each map $T_n : \mathcal{M}_n(R) \rightarrow A$ is entirely determined by the map $T_1 : R = \mathcal{M}_1(R) \rightarrow A$, [Sta65, 1.4, 1.5] as $T_n(M) = \sum_{i=1}^n T_1(m_{ii})$, where $M = [m_{ij}]$. Therefore we can rephrase a *trace* on R with values in A to be a \mathbb{Z} -linear homomorphism $T : R \rightarrow A$ such that $T(rs) = T(sr)$, for all $r, s \in R$, i.e., the following diagram commutes:

$$R \otimes_{\mathbb{Z}} R \xrightleftharpoons[\text{m} \circ \tau]{\text{m}} R \xrightarrow{T} A,$$

where m denotes the multiplication on R .

If we let $[R, R] = \{rs - sr \mid r, s \in R\}$ be the subgroup of commutators, then we obtain a universal trace $\text{tr} : R \rightarrow R/[R, R]$ by the quotient homomorphism, and every trace $T : R \rightarrow A$ is uniquely determined by a \mathbb{Z} -linear homomorphism $R/[R, R] \rightarrow A$:

$$\begin{array}{ccc}
R & \xrightarrow{T} & A. \\
\text{tr} \downarrow & \nearrow \exists! & \\
R/[R, R] & &
\end{array}$$

Given M a finitely generated free right R -module, then an R -linear endomorphism $f: M \rightarrow M$ is represented by a square matrix on R , and thus any trace $T: R \rightarrow A$ defines a trace $T_M: \text{End}_R(M) \rightarrow A$ as we have the isomorphism of ring $\text{End}_R(M) \cong \mathcal{M}_n(R)$, where n is the rank of M .

More generally, if M is a finitely generated projective right R -module, then there exists another projective module N such that $M \oplus N$ is a free R -module. Therefore if we extend an R -linear endomorphism $f: M \rightarrow M$ to the endomorphism $f+0: M \oplus N \rightarrow M \oplus N$, we again obtain a square matrix, and thus any trace $T: R \rightarrow A$ defines a trace $T_M: \text{End}_R(M) \rightarrow A$.

Definition 2.2. Let R be a ring, and M be a finitely generated projective right R -module. The *Hattori–Stallings trace* on M is the trace induced by the universal trace $\text{tr}: R \rightarrow R/[R, R]$ on the endomorphisms of M :

$$\text{tr}_M: \text{End}_R(M) \longrightarrow R/[R, R].$$

The *Hattori–Stallings rank* of M is then the trace of the identity endomorphism on M . The abelian group $R/[R, R]$ is isomorphic to $\text{HH}_0(R) = R \otimes_{R \otimes R^{\text{op}}} R$, the zeroth *Hochschild homology* of R .

The definition above depends on the choice of generators of M and a section $R^{\oplus n} \rightarrow M$. However, the Hattori–Stallings trace can be defined coordinate-free as follows. Given a finitely generated projective right R -module M , we have an isomorphism

$$M \otimes_R \text{Hom}_R(M, R) \xrightarrow{\cong} \text{Hom}_R(M, M) = \text{End}_R(M),$$

and thus the Hattori–Stallings trace is the composition

$$\text{End}_R(M) \xleftarrow{\cong} M \otimes_R \text{Hom}_R(M, R) \xrightarrow{\text{ev}} R \xrightarrow{\text{tr}} \text{HH}_0(R).$$

Let $P(R)$ be the set of isomorphism classes of finitely generated projective right R -modules. It is a commutative monoid with respect to direct sum. Any trace $T: R \rightarrow A$ defines an additive homomorphism with its rank:

$$\begin{aligned} P(R) &\longrightarrow A \\ [M] &\longrightarrow T_M(\text{id}_M). \end{aligned}$$

Let $K_0(R)$ be the group completion of $P(R)$. Then the additive map factors through a \mathbb{Z} -linear homomorphism

$$T: K_0(R) \longrightarrow A.$$

Choosing T to be the universal trace $\text{tr}: R \rightarrow \text{HH}_0(R)$ defines the *Hattori–Stallings rank* $K_0(R) \rightarrow \text{HH}_0(R)$, which is the zeroth level of the Dennis trace $K_*(R) \rightarrow \text{HH}_*(R)$.

2.2. The Hattori–Stallings cotrace. We wish to dualize the approach of Hattori and Stallings in the coalgebraic context. Just as the group $\text{HH}_0(R)$ is the universal home for traces, we introduce a group $\text{coHH}_0(C)$ as the universal home for so-called cotraces. Throughout the rest of the section, let (C, Δ, ε) be a \mathbb{k} -coalgebra, not necessarily cocommutative.

Definition 2.3. Let V be a \mathbb{k} -module. A \mathbb{k} -linear homomorphism $T: V \rightarrow C$ is said to be a *cotrace on C from V* if it is *cocyclic* in the sense that the following diagram commutes:

$$V \xrightarrow{T} C \xrightarrow[\tau \circ \Delta]{\Delta} C \otimes C.$$

In other words, using the Sweedler notation $\Delta(c) = \sum_{(c)} c_{(1)} \otimes c_{(2)}$, the \mathbb{k} -linear homomorphism $T: V \rightarrow C$ is a cotrace if for all $v \in V$:

$$\sum_{(T(v))} T(v)_{(1)} \otimes T(v)_{(2)} = \sum_{(T(v))} T(v)_{(2)} \otimes T(v)_{(1)}.$$

Definition 2.4. The *cocommutator* submodule $\langle\langle C \rangle\rangle$ is the kernel in \mathbb{k} -modules:

$$\langle\langle C \rangle\rangle \xrightarrow{\text{cotr}} C \xrightarrow[\tau \circ \Delta]{\Delta} C \otimes C.$$

The *universal cotrace* on C is the resulting cotrace $\text{cotr}: \langle\langle C \rangle\rangle \hookrightarrow C$. Every cotrace $T: V \rightarrow C$ is uniquely determined by a \mathbb{k} -linear homomorphism $V \rightarrow \langle\langle C \rangle\rangle$:

$$\begin{array}{ccc} & & \langle\langle C \rangle\rangle \\ & \nearrow \exists! & \downarrow \text{cotr} \\ V & \xrightarrow{T} & C \end{array}$$

Notice that $\langle\langle C \rangle\rangle$ is isomorphic to the zeroth coHochschild homology $\text{coHH}_0(C) := C \square_{C \otimes C^{\text{op}}} C$. We recall the definition of $\text{coHH}_q(C)$ more generally in Definition 4.3.

Recall that any trace $T: R \rightarrow A$ can be extended on square matrices $T_n: \mathcal{M}_n(R) \rightarrow A$ by assigning $T_n(M) = \sum_{i=1}^n T(m_{ii})$, where $M = [m_{ij}]$. We wish to do similarly for cotraces.

Definition 2.5. Given $n, m \geq 1$, define the (n, m) -comatrices $\mathcal{M}_{n \times m}^c(\mathbb{k})$ to be usual \mathbb{k} -module $\mathcal{M}_{n \times m}(\mathbb{k})$ of matrices of size $n \times m$ with coefficients in \mathbb{k} , but endowed with the following two operations. Let $\{E_{ij}\}$ be the canonical basis of $\mathcal{M}_{n \times m}(\mathbb{k})$ composed of the elementary matrices. Define for all $r \geq 1$ the \mathbb{k} -linear homomorphism:

$$\begin{aligned} \Delta: \mathcal{M}_{n \times m}^c(\mathbb{k}) &\longrightarrow \mathcal{M}_{n \times r}^c(\mathbb{k}) \otimes \mathcal{M}_{r \times m}^c(\mathbb{k}) \\ E_{ij} &\longmapsto \sum_{\ell=1}^r E_{i\ell} \otimes E_{\ell j}, \end{aligned}$$

and the \mathbb{k} -linear homomorphism:

$$\begin{aligned} \varepsilon: \mathcal{M}_{n \times n}^c(\mathbb{k}) &\longrightarrow \mathbb{k} \\ E_{ij} &\longmapsto \delta_{ij}, \end{aligned}$$

where δ_{ij} is the Kronecker symbol. The maps Δ and ε are coassociative and counital in the sense that the comatrices $\mathcal{M}_{n \times m}^c(\mathbb{k})$ constitute the morphisms of a category $\text{Mat}^c(\mathbb{k})$ enriched in $\text{Mod}_{\mathbb{k}}^{\text{op}}$, the opposite category of \mathbb{k} -modules. In fact, the underlying category of $\text{Mat}^c(\mathbb{k})$ is anti-equivalent to the usual category $\text{Mat}(\mathbb{k})$ of matrices. Denote by $\mathcal{M}_{n \times m}^c(C) = C \otimes \mathcal{M}_{n \times m}^c(\mathbb{k})$ the (n, m) -comatrices with values in C . Combining with the comultiplication and counit of C , the \mathbb{k} -modules $\mathcal{M}_{n \times m}^c(C)$ extend the previous operations to $\mathcal{M}_{n \times m}^c(C) \rightarrow \mathcal{M}_{n \times r}^c(C) \otimes \mathcal{M}_{r \times m}^c(C)$ and $\mathcal{M}_{n \times m}^c(C) \rightarrow \mathbb{k}$, which are again coassociative and counital. When $n = m$, we denote $\mathcal{M}_n^c(C)$ the square comatrices: these are \mathbb{k} -coalgebras and $\mathcal{M}_n^c(\mathbb{k})$ is anti-equivalent to the usual matrix ring $\mathcal{M}_n(\mathbb{k})$. Moreover, by [BW03, 12.20], the coalgebras C and $\mathcal{M}_n^c(C)$ are Morita–Takeuchi equivalent for all $n \geq 1$ (Definition 1.17).

Given a cotrace $T: V \rightarrow C$, it can be extended to a \mathbb{k} -linear homomorphism $T_n: V \rightarrow \mathcal{M}_n^c(C)$ by assigning $v \mapsto T(v) \otimes I_n$, where I_n is the identity matrix of size n . These homomorphisms $T_n: V \rightarrow \mathcal{M}_n^c(C)$ are cocyclic in the sense that the following diagram commutes:

$$\begin{array}{ccc} & \mathcal{M}_n^c(C) & \xrightarrow{\Delta} \mathcal{M}_{n \times m}^c(C) \otimes \mathcal{M}_{m \times n}^c(C) \\ & \nearrow T_n & \downarrow \tau \\ V & & \\ & \searrow T_m & \downarrow \\ & \mathcal{M}_m^c(C) & \xrightarrow{\Delta} \mathcal{M}_{m \times n}^c(C) \otimes \mathcal{M}_{n \times m}^c(C). \end{array} \tag{2.6}$$

The commutative diagram captures the dual property $T_n(MN) = T_m(NM)$ of traces.

Lemma 2.7. A cotrace $T: V \rightarrow C$ is uniquely determined by a collection of \mathbb{k} -linear homomorphisms $T_n: V \rightarrow \mathcal{M}_n^c(C)$ satisfying (2.6) for which $T_n(v) = T(v) \otimes I_n$ for all $v \in V$ and $n \geq 1$.

We now wish extend further the cotraces just as a trace $T: R \rightarrow A$ can be extended to $T_M: \text{End}_R(M) \rightarrow A$ for M a finitely generated and projective right R -module.

Definition 2.8. We say a left C -comodule M is *finitely cogenerated* if there is a C -colinear monomorphism $M \hookrightarrow C^{\oplus n}$ for some $n \geq 0$.

A left C -comodule M is *injective* in the usual abelian category sense: any C -colinear monomorphisms $M \hookrightarrow N$ splits. It is immediate to obtain the following result.

Lemma 2.9. *For any finitely cogenerated and injective left C -comodule M , there is another finitely cogenerated and injective left C -comodule N such that $M \oplus N \cong C^{\oplus n}$ for some $n \geq 0$.*

Finitely cogenerated and injective left C -comodules are the dualizable comodules in a sense that we make precise in Section 5. From Definition 5.2 and Example 5.4, given any finitely cogenerated left C -comodule M , the functor $M \otimes -: \text{Mod}_{\mathbb{k}} \rightarrow {}_C\text{CoMod}$ admits a left adjoint $h_C(M, -): {}_C\text{CoMod} \rightarrow \text{Mod}_{\mathbb{k}}$ called the *cohom* functor. In other words, given any left C -comodule N , the cohom $h_C(M, N)$ is the universal \mathbb{k} -module providing a natural \mathbb{k} -linear isomorphism

$$\text{Mod}_{\mathbb{k}}(h_C(M, N), V) \cong {}_C\text{CoMod}(N, M \otimes V),$$

for any \mathbb{k} -module V . In particular, for $V = \mathbb{k}$, we obtain the isomorphism $h_C(M, N)^* \cong {}_C\text{Hom}(N, M)$. Given any left C -comodule N , there is a C -colinear homomorphism $\eta: N \rightarrow M \otimes h_C(M, N)$ called the *coevaluation* on N with M , adjoint of the identity on $h_C(M, N)$.

Definition 2.10 ([BW03, 12.9]). Let M be a finitely cogenerated left C -comodule. The *coendomorphism coalgebra* on M denoted $e_C(M)$ is the \mathbb{k} -module $h_C(M, M)$ endowed with the comultiplication:

$$h_C(M, M) \longrightarrow h_C(M, M) \otimes h_C(M, M),$$

which is the adjoint of the C -colinear homomorphism:

$$M \xrightarrow{\eta} M \otimes h_C(M, M) \xrightarrow{\eta \otimes 1} M \otimes h_C(M, M) \otimes h_C(M, M).$$

The counit $h_C(M, M) \rightarrow \mathbb{k}$ is the adjoint of the identity of the C -colinear homomorphism $M \xrightarrow{\cong} M \otimes \mathbb{k}$.

We have an isomorphism of \mathbb{k} -coalgebras $e_C(C^{\oplus n}) \cong \mathcal{M}_n^c(C)$, [BW03, 12.20]. More generally, we obtain the identification $h_C(C^{\oplus n}, C^{\oplus m}) \cong \mathcal{M}_{n \times m}^c(C)$. The linear dual $e_C(M)^*$ is anti-equivalent to the \mathbb{k} -algebra ${}_C\text{End}(M)$, see [BW03, 12.9].

Lemma 2.11. *Given a finitely cogenerated and injective left C -comodule M , there exists a \mathbb{k} -linear homomorphism $\mathcal{M}_n(C) \rightarrow e_C(M)$ that becomes precisely the following monomorphism after taking the linear dual:*

$$\begin{aligned} {}_C\text{End}(M) &\longrightarrow {}_C\text{End}(C^{\oplus n}) \\ (M \xrightarrow{f} M) &\longmapsto (M \oplus N \xrightarrow{f \oplus 0} M \oplus N), \end{aligned}$$

for some choice of N such that $M \oplus N \cong C^{\oplus n}$.

Proof. As M is finitely cogenerated, there exists $n \geq 0$ such that $M \hookrightarrow C^{\oplus n}$. Since M is injective, we can choose a C -colinear section $C^{\oplus n} \rightarrow M$. Define $h_C(C^{\oplus n}, C^{\oplus n}) = \mathcal{M}_n(C) \rightarrow e_C(M) = h_C(M, M)$ as the adjoint the composition:

$$C^{\oplus n} \longrightarrow M \otimes h_C(M, C^{\oplus n}) \longrightarrow C^{\oplus n} \otimes h_C(M, C^{\oplus n}) \longrightarrow C^{\oplus n} \otimes h_C(M, M).$$

The first map is induced by coevaluation on $C^{\oplus n}$, the second map is induced by $M \hookrightarrow C^{\oplus n}$, and the third map is induced by applying $h_C(M, -)$ on the section $C^{\oplus n} \rightarrow M$. \square

Therefore, given a finitely cogenerated and injective left C -comodule M , any cotrace $T: V \rightarrow C$ extends to $T_M: V \rightarrow e_C(M)$ as the composite:

$$V \xrightarrow{T_n} \mathcal{M}_n(C) \longrightarrow e_C(M),$$

for some $n \geq 1$. Applying the linear dual yields a \mathbb{k} -linear homomorphism

$$T_M: {}_C\text{End}(M) \cong e_C(M)^* \longrightarrow V^*.$$

As $C^{\oplus n}$ is the cofree C -comodule on $\mathbb{k}^{\oplus n}$, we obtain the identification:

$${}_C\text{End}(C^{\oplus n}) \cong \text{Hom}_{\mathbb{k}}(C^{\oplus n}, \mathbb{k}^{\oplus n}) \cong \text{Hom}_{\mathbb{k}}(C, \mathcal{M}_n(\mathbb{k})),$$

which sends the extended endomorphism $f \oplus 0$ on $C^{\oplus n}$ of a C -colinear endomorphism $f: M \rightarrow M$ to a \mathbb{k} -linear homomorphism $C \rightarrow \mathcal{M}_n(\mathbb{k})$ that we denote $c \mapsto f_c$. Unwinding the definitions, we obtain the following formula for $T_M: {}_C\text{End}(M) \rightarrow V^*$,

$$T_M(f)(v) = \sum_{(T(v))} \varepsilon(T(v)_{(1)}) \text{tr}\left(f_{T(v)_{(2)}}\right), \quad (2.12)$$

for any $v \in V$ and C -colinear endomorphism $f: M \rightarrow M$. Here $\varepsilon: C \rightarrow \mathbb{k}$ denotes the counit, and the comultiplication $C \rightarrow C \otimes C$ is denoted $c \mapsto \sum_{(c)} c_{(1)} \otimes c_{(2)}$ using Sweedler notation.

Definition 2.13. Let M be a finitely cogenerated and injective left C -comodule. The *Hattori–Stallings cotrace on M* is the cotrace induced by universal cotrace $\text{cotr}: \text{coHH}_0(C) \hookrightarrow C$ on the endomorphisms of M :

$$\text{cotr}_M: {}_C\text{End}(M) \rightarrow \text{coHH}_0(C)^*.$$

Explicitly, it is given by the formula, for all $c \in \text{coHH}_0(C)$:

$$\text{cotr}_M(f)(c) = \sum_{(c)} \varepsilon(c_{(1)}) \text{tr}(f_{c_{(2)}}).$$

The *Hattori–Stallings corank on M* is the cotrace on the identity.

Remark 2.14. A priori, the definition of the Hattori–Stallings cotrace may depend on the choice of the section $C^{\oplus n} \rightarrow M$, but we shall see in Proposition 6.4 that the cotrace is a bicategorical trace in a shadow framework and thus has a coordinate-free formulation, just as the usual Hattori–Stallings trace.

Remark 2.15. In [Qui88], Quillen already established $\text{coHH}_0(C)$ as the universal home for cotrace maps with universal cotrace $\text{coHH}_0(C) \hookrightarrow C$ as we do here. He identified $\text{coHH}_0(C)$ with cyclic homology HC_* in certain cases. In [Qui89], Quillen utilized the universal cotrace to extend trace maps on a ring R to traces $K_0(A) \rightarrow \text{HC}_{2n}(A)$, where A is a quotient algebra of R . However, Quillen did not extend cotraces $V \rightarrow C$ to cotraces $V \rightarrow \mathcal{M}_n^c(C)$ nor $V \rightarrow e_C(M)$. Our Hattori–Stallings cotrace is an extension of the universal cotrace $\text{coHH}_0(C) \hookrightarrow C$.

Let $\text{I}(C)$ be the isomorphism class of finitely cogenerated injective left C -comodules. It is a commutative monoid with respect to the direct sum. Any cotrace $T: V \rightarrow C$, defines an additive homomorphism with its rank:

$$\begin{aligned} \text{I}(C) &\longrightarrow V^* \\ [M] &\longmapsto T_M(\text{id}_M). \end{aligned}$$

Denote $K_0^c(C)$ the group completion of $\text{I}(C)$. Since the abelian category ${}_C\text{CoMod}^{\text{fc}, \text{inj}}$ is split exact, then $K_0^c(C)$ is indeed equivalent to the π_0 -group of the algebraic K -theory spectrum of Definition 1.5. Therefore, the above additive map factors through a \mathbb{Z} -linear homomorphism:

$$T: K_0^c(C) \longrightarrow V^*.$$

Choosing T to be the universal cotrace $\text{cotr}: \text{coHH}_0(C) \hookrightarrow C$ yields the following result.

Theorem 2.16. *Let C be a \mathbb{k} -coalgebra. The Hattori–Stallings corank defines a homomorphism of abelian groups:*

$$K_0^c(C) \longrightarrow \text{coHH}_0(C)^*.$$

Example 2.17. If $C = \mathbb{k}$, then $K^c(\mathbb{k})$ is equivalent to the algebraic K -theory $K(\mathbb{k})$ of \mathbb{k} as a ring, and thus $K_0^c(\mathbb{k}) \cong \mathbb{Z}^{\times n}$ as \mathbb{k} is a finite product of fields, and $\text{coHH}_0(C)^* = \mathbb{k}$, and the corank $\mathbb{Z}^n \rightarrow \mathbb{k}$ is induced by the unique ring homomorphism from \mathbb{Z} on each field in the direct summand of \mathbb{k} .

Remark 2.18 (Eilenberg Swindle). Just as in the case for rings, we needed to restrict to finitely cogenerated comodules, as otherwise $K_0^c(C) = 0$. Indeed, let $\mathbb{k}^{\oplus \infty}$ be the \mathbb{k} -vector space with countably infinite basis.

Let $C^{\oplus\infty} = C \otimes \mathbb{k}^{\oplus\infty}$ be an infinitely cogenerated cofree left C -comodule. Then we obtain isomorphisms of left C -comodules

$$\begin{aligned} C^{\oplus\infty} &= C \otimes \mathbb{k}^{\oplus\infty} \\ &\cong C \otimes (\mathbb{k}^{\oplus n} \oplus \mathbb{k}^{\oplus n} \oplus \dots) \\ &\cong (C \otimes \mathbb{k}^{\oplus n}) \oplus (C \otimes \mathbb{k}^{\oplus n}) \oplus \dots \\ &\cong C^{\oplus n} \oplus C^{\oplus n} \oplus \dots. \end{aligned}$$

Therefore if $M \oplus N \cong C^{\oplus n}$, then we obtain

$$\begin{aligned} M \oplus C^{\oplus\infty} &\cong M \oplus (N \oplus M) \oplus (N \oplus M) \oplus \dots \\ &\cong (M \oplus N) \oplus (M \oplus N) \oplus \dots \\ &\cong C^{\oplus\infty}. \end{aligned}$$

Hence, in $K_0^c(C)$ we would get $[M] = 0$.

Proposition 2.19. *If C is a conilpotent coaugmented \mathbb{k} -coalgebra, then $K_0^c(C) \cong \mathbb{Z}$.*

Proof. By [Pos23, 2.6], injective C -comodules are all cofree when C is a conilpotent coaugmented coalgebra. Thus $I(C) \cong \mathbb{N}$, and the result follows. \square

Proposition 2.20. *If C is a cocommutative \mathbb{k} -coalgebra, then $K_0^c(C)$ is a commutative ring. In fact, if C is cocommutative, then $K^c(C)$ is a ring spectrum.*

Proof. If C is cocommutative, then the cotensor product forms a symmetric monoidal structure on C -comodules. The cotensor product of injective C -comodules is injective [Doi81, Proposition 1]. The cotensor product of finitely cogenerated C -comodules is finitely cogenerated. Therefore $K_0^c(C)$ is a commutative ring via cotensor product of C -comodules. As the cotensor product preserves monomorphisms in each variable, then the Waldhausen category ${}_C\text{CoMod}^{\text{fc}, \text{inj}}$ is symmetric monoidal and thus $K^c(C)$ is a ring spectrum [BM11, 2.8]. \square

2.3. The colinear Hattori–Stallings trace. Another approach is to obtain a coalgebraic refinement of the Hattori–Stallings trace as we now explain. We can extend the notion of a trace $T: R \rightarrow A$ to a trace on any (R, R) -bimodule P as a \mathbb{Z} -linear homomorphism $T: P \rightarrow A$ such that $T(rx) = T(xr)$ for all $r \in R$, $x \in P$. A universal trace is then given by $\text{tr}: P \rightarrow \text{HH}_0(R, P) := R \otimes_{R \otimes R^{\text{op}}} P$. If M is a right R -module, then $\text{Hom}_R(M, R) \otimes_{\mathbb{Z}} M$ is an (R, R) -bimodule.

Definition 2.21 ([Pon10, 2.3.4]). Let M be a finitely generated projective right R -module, and P an (R, R) -bimodule. The *twisted Hattori–Stallings trace* $\text{Hom}_R(M, M \otimes_R P) \rightarrow \text{HH}_0(R, P)$ is defined on a right R -linear homomorphism $f: M \rightarrow M \otimes_R P$, referred to as an *endomorphism on M twisted by P* , as the image of f under the composition

$$\text{Hom}_R(M, M \otimes_R P) \xleftarrow{\cong} (M \otimes_R P) \otimes_R \text{Hom}_R(M, R) \xrightarrow{\text{ev}} P \xrightarrow{\text{tr}} \text{HH}_0(R, P).$$

Given a \mathbb{k} -coalgebra C , we now choose $R = \mathbb{k}$ and $P = C$ as a (\mathbb{k}, \mathbb{k}) -bimodule, then the twisted Hattori–Stallings trace becomes the \mathbb{k} -linear homomorphism $\text{tr}: \text{Hom}_{\mathbb{k}}(M, M \otimes C) \rightarrow \text{HH}_0(\mathbb{k}, C) = C$. If we require M to be a right C -comodule, finitely generated as a \mathbb{k} -module, with coaction $\rho: M \rightarrow M \otimes C$, we define $\text{tr}^C: \text{End}_C(M) \rightarrow C$ by $\text{tr}^C(f) = \text{tr}(\rho \circ f)$.

Proposition 2.22. *The \mathbb{k} -linear homomorphism $\text{tr}^C: \text{End}_C(M) \rightarrow C$ is a cotrace. In particular, it factors uniquely through the universal cotrace:*

$$\begin{array}{ccc} & & \text{coHH}_0(C) \\ & \nearrow \exists! \text{tr}^C & \downarrow \text{cotr} \\ \text{End}_C(M) & \xrightarrow{\text{tr}^C} & C. \end{array}$$

13

Definition 2.23. Given a right C -comodule M that is finitely generated as a \mathbb{k} -module, we call the induced \mathbb{k} -linear homomorphism $\text{tr}^C: \text{End}_C(M) \rightarrow \text{coHH}_0(C)$ the *colinear Hattori–Stallings trace on M* . Explicitly, given a basis (e_1, \dots, e_n) of M , and a C -colinear homomorphism $f: M \rightarrow M$, we have the formula:

$$\text{tr}^C(f) = \sum_{i=1}^n \sum_{(e_i)} e_i^*(f(e_{i(0)})) e_{i(1)}$$

where we denoted $m \mapsto \sum_{(m)} m_{(0)} \otimes m_{(1)}$ the coaction $M \rightarrow M \otimes C$. The *colinear Hattori–Stallings rank on M* is the colinear trace on the identity.

Proof of Proposition 2.22. Denote $\Delta: C \rightarrow C \otimes C$ the comultiplication on C . We need to show that for all colinear endomorphisms f , we have $\Delta(\text{tr}^C(f)) = \tau(\Delta(\text{tr}^C(f)))$. As f is colinear and M is a right C -comodule, we obtain the commutative diagram:

$$\begin{array}{ccccc} M & \xrightarrow{f} & M & \xrightarrow{\rho} & M \otimes C \\ \rho \downarrow & & \downarrow \rho & & \downarrow 1 \otimes \Delta \\ M \otimes C & \xrightarrow{f \otimes 1} & M \otimes C & \xrightarrow{\rho \otimes 1} & M \otimes C^{\otimes 2} \end{array} \quad (2.24)$$

As M is a right C -comodule, its linear dual M^* is a left C -comodule with coaction:

$$\begin{aligned} \lambda: M^* &\longrightarrow \text{Hom}_k(M, C) \xleftarrow{\cong} C \otimes M^* \\ \alpha &\longmapsto (M \rightarrow M \otimes C \xrightarrow{\alpha} C) \end{aligned}$$

The coactions are compatible on $M^* \otimes M$ in the sense that the following diagram commutes:

$$\begin{array}{ccccc} M^* \otimes M & \xrightarrow{\lambda \otimes 1} & C \otimes M^* \otimes M & & \\ 1 \otimes \rho \downarrow & & \downarrow 1 \otimes \text{ev} & \searrow \lambda \otimes 1 \otimes 1 & \\ M^* \otimes M \otimes C & \xrightarrow{\text{ev} \otimes 1} & C & & C^{\otimes 2} \otimes M^* \otimes M \\ & \searrow 1 \otimes 1 \otimes \rho & & & \downarrow 1 \otimes 1 \otimes \text{ev} \\ & & M^* \otimes M \otimes C^{\otimes 2} & \xrightarrow{\text{ev} \otimes 1 \otimes 1} & C^{\otimes 2}. \end{array} \quad (2.25)$$

Using the usual convention for Sweedler notation, and (2.24) and (2.25), we obtain

$$\begin{aligned} \Delta(\text{tr}^C(f)) &= \sum_i \sum_{(e_i)} e_i^*(f(e_{i(0)})) e_{i(1)} \otimes e_{i(2)} \\ &= \sum_i \sum_{(f(e_i))} e_i^*(f(e_i)_{(0)}) f(e_i)_{(1)} \otimes f(e_i)_{(2)} \\ &= \sum_i \sum_{(e_i^*)} (e_i^*)_{(0)} (f(e_i)) (e_i^*)_{(2)} \otimes (e_i^*)_{(1)} \\ &= \sum_i \sum_{(e_i)} e_i^*(f(e_i)_{(0)}) e_{i(2)} \otimes e_{i(1)} \\ &= \tau(\Delta(\text{tr}^C(f))). \end{aligned} \quad \square$$

Let $M(C)$ be the isomorphism class of right C -comodules that are finitely generated as \mathbb{k} -modules. By [Wei13, II.9.1.3, IV.8.4], the π_0 -group of the algebraic K -theory $K(\text{CoMod}_C^{\text{fg}})$ of Definition 1.8 is isomorphic to the free abelian group on $M(C)$ under the relation $[M] = [M'] + [M'']$ whenever $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is an exact sequence in $\text{CoMod}_C^{\text{fg}}$. The colinear Hattori–Stallings rank defines an additive homomorphism $M(C) \rightarrow \text{coHH}_0(C)$. We wish to show it factors through $K_0(\text{CoMod}_C^{\text{fg}})$.

Lemma 2.26 (Additivity). *Given a morphism of short exact sequences in $\text{CoMod}_C^{\text{fg}}$:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' \longrightarrow 0 \\ & & \downarrow f' & & \downarrow f & & \downarrow f'' \\ 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' \longrightarrow 0 \end{array}$$

we obtain $\text{tr}^C(f) = \text{tr}^C(f') + \text{tr}^C(f'')$ in $\text{coHH}_0(C)$.

Proof. This follows from $\text{tr}^C(f) = \text{tr}(\rho \circ f)$ and the fact that the usual twisted trace is additive. In more details, we have $M \cong M' \oplus M''$ as \mathbb{k} -modules. Denote the coactions as follows:

$$\begin{array}{lll} M \rightarrow M \otimes C & M' \rightarrow M' \otimes C & M'' \rightarrow M'' \otimes C \\ m \mapsto \sum_{(m)} m_{(0)}^M \otimes m_{(1)}^M & m \mapsto \sum_{(m)} m_{(0)}^{M'} \otimes m_{(1)}^{M'} & m \mapsto \sum_{(m)} m_{(0)}^{M''} \otimes m_{(1)}^{M''}. \end{array}$$

Choose a basis $(e_1, \dots, e_n, e_{n+1}, \dots, e_{n+m})$ of M such that (e_1, \dots, e_n) is a basis of M' and $(e_{n+1}, \dots, e_{n+m})$ is a basis of M'' . As f, f' and f'' are colinear, we obtain for all $1 \leq i \leq n$ and $n+1 \leq j \leq n+m$:

$$\sum_{(e_i)} f'(e_{i(0)}^{M'}) \otimes e_{i(1)}^{M'} = \sum_{(e_i)} f(e_{i(0)}^M) \otimes e_{i(1)}^M \quad \sum_{(e_i)} f''(e_{i(0)}^{M''}) \otimes e_{i(1)}^{M''} = \sum_{(e_i)} f(e_{i(0)}^M) \otimes e_{i(1)}^M.$$

Then:

$$\begin{aligned} \text{tr}^C(f) &= \sum_{i=1}^{n+m} \sum_{(e_i)} e_i^*(f(e_{i(0)}^M)) e_{i(1)}^M \\ &= \sum_{i=1}^n \sum_{(e_i)} e_i^*(f(e_{i(0)}^M)) e_{i(1)}^M + \sum_{i=n+1}^{n+m} \sum_{(e_i)} e_i^*(f(e_{i(0)}^M)) e_{i(1)}^M \\ &= \sum_{i=1}^n \sum_{(e_i)} e_i^*(f'(e_{i(0)}^{M'})) e_{i(1)}^{M'} + \sum_{i=n+1}^{n+m} \sum_{(e_i)} e_i^*(f''(e_{i(0)}^{M''})) e_{i(1)}^{M''} \\ &= \text{tr}^C(f') + \text{tr}^C(f''). \end{aligned} \quad \square$$

Theorem 2.27. *Let C be a \mathbb{k} -coalgebra. The colinear Hattori–Stallings rank defines a homomorphism of abelian groups:*

$$K_0(\text{CoMod}_C^{\text{fg}}) \longrightarrow \text{coHH}_0(C).$$

Proof. Considering trace on identity yields a homomorphism $\mathbb{Z}[\text{M}(C)] \rightarrow \text{coHH}_0(C)$. Since the colinear trace tr^C is additive, then the previous homomorphism must factor through $K_0(\text{CoMod}_C^{\text{fg}})$. \square

Since the cotensor product of finitely generated comodules is again finitely generated, we obtain a similar result as Proposition 2.20.

Proposition 2.28. *If C is a cocommutative \mathbb{k} -coalgebra, then $K(\text{CoMod}_C^{\text{fg}})$ is a ring spectrum.*

Any right C -comodule M can be given a left C^* -module with the action given by the coaction followed by evaluation:

$$C^* \otimes M \rightarrow C^* \otimes M \otimes C \cong C^* \otimes C \otimes M \rightarrow \mathbb{k} \otimes M \cong M.$$

This defines a fully faithful functor $\text{CoMod}_C \hookrightarrow {}_{C^*}\text{Mod}$ that is an equivalence of categories if and only if C is finitely generated as a \mathbb{k} -module [BW03, 4.7]. In that case, the ring C^* is also finitely generated as a \mathbb{k} -module, and a left C^* -module is finitely generated over C^* if and only if it is finitely generated over \mathbb{k} . This results into an equivalence $\text{CoMod}_C^{\text{fg}} \simeq {}_{C^*}\text{Mod}^{\text{fg}}$ between right C -comodules and left C^* -modules that are finitely generated as \mathbb{k} -modules. Considering ${}_{C^*}\text{Mod}^{\text{fg}}$ as a Waldhausen category with cofibrations all monomorphisms, we obtain $K({}_{C^*}\text{Mod}^{\text{fg}}) \simeq G(C^*)$, the G -theory of the ring C^* . The Cartan map $K(C^*) \rightarrow G(C^*)$ combined with the colinear Hattori–Stallings rank defines a homomorphism $K_0(C^*) \rightarrow \text{coHH}_0(C) \cong \text{HH}_0(C^*)^* \cong \text{HH}_0(C^*)$, and this is the usual Hattori–Stallings rank $K_0(C^*) \rightarrow \text{HH}_0(C^*)$ on the ring C^* . We record our observation

as follows choosing $C^* = R$, then R is a \mathbb{k} -algebra that is finitely generated as a \mathbb{k} -module, i.e. an Artinian \mathbb{k} -algebra.

Proposition 2.29. *Let R be an Artinian \mathbb{k} -algebra. There is an equivalence of algebraic K -theory spectra:*

$$G(R) \simeq K \left(\text{CoMod}_{R^*}^{\text{fg}} \right),$$

and the combination of the Cartan maps $K(R) \rightarrow G(R)$ and colinear Hattori–Stallings rank on R^* recovers the Dennis trace $K_0(R) \rightarrow \text{coHH}_0(R^*) \cong \text{HH}_0(R)$.

Remark 2.30. If R is a commutative Artinian \mathbb{k} -algebra, then $R = R_1 \times \cdots \times R_n$ is a finite product of Artinian local rings R_i [Lan02, X.7.7.(ii)]. By [Wei13, V.4.2.1], when \mathbb{k} is a field, we get $G(R_i) \simeq K(\mathbb{k})$. Therefore, for R a commutative Artinian algebra over a field \mathbb{k} , we obtain:

$$K \left(\text{CoMod}_{R^*}^{\text{fg}} \right) \simeq \prod_{i=1}^n K(\mathbb{k}).$$

3. BICATEGORIES OF BICOMODULES

In this section, we introduce the bicategories of bicomodules in Definition 3.5 and Definition 3.13. For convenience, we recall the definition of a bicategory, further details of which may be found in [Pon10, PS13, CP19].

Definition 3.1. A *bicategory* \mathcal{B} consists of following.

- A collection of objects, $ob(\mathcal{B})$, called *0-cells*. We write $C \in \mathcal{B}$ instead of $C \in ob(\mathcal{B})$.
- Categories $\mathcal{B}(C, D)$ for each pair of objects $C, D \in \mathcal{B}$. The objects in these categories are referred to as *1-cells* and the morphisms as *2-cells*. The composition is referred to as *vertical composition*.
- Unit functors $U_C \in \mathcal{B}(C, C)$ for all $C \in \mathcal{B}$.
- *Horizontal composition* functors for $C, D, E \in \mathcal{B}$

$$\odot: \mathcal{B}(C, D) \times \mathcal{B}(D, E) \rightarrow \mathcal{B}(C, E),$$

which are not required to be strictly associative or unital.

- Natural isomorphisms for $M \in \mathcal{B}(C, D)$, $N \in \mathcal{B}(D, E)$, $P \in \mathcal{B}(E, F)$, and $Q \in \mathcal{B}(F, G)$ given $C, D, E, F, G \in \mathcal{B}$:

$$a: (M \odot N) \odot P \xrightarrow{\cong} M \odot (N \odot P),$$

$$\ell: U_C \odot M \xrightarrow{\cong} M, \quad r: M \odot U_D \xrightarrow{\cong} M,$$

which satisfy the triangle identity

$$\begin{array}{ccc} (M \odot U_D) \odot N & \xrightarrow{a} & M \odot (U_D \odot N) \\ & \searrow r \odot \text{id}_N & \swarrow \text{id}_M \odot \ell \\ & M \odot N, & \end{array}$$

and the pentagon identity

$$\begin{array}{ccccc} & & (M \odot N) \odot (P \odot Q) & & \\ & \nearrow a & & \searrow a & \\ ((M \odot N) \odot P) \odot Q & & & & M \odot (N \odot (P \odot Q)) \\ \downarrow a \odot \text{id}_Q & & & & \uparrow \text{id}_M \odot a \\ (M \odot (N \odot P)) \odot Q & \xrightarrow{a} & & & M \odot ((N \odot P) \odot Q). \end{array}$$

A bicategory with one object is precisely a monoidal category. The bicategory of categories in which 0-cells are categories, 1-cells are functors, and 2-cells are natural transformations is a classical example. We recall other examples below that will be motivating in our context.

Example 3.2. We can define a bicategory whose 0-cells are rings and whose 1- and 2-cells come from the category of (R, S) -bimodules for rings R and S . In this case the horizontal composition is given by the tensor product of bimodules. The derived version of this bicategory has 0-cells which are rings and 1- and 2-cells from the derived category of (R, S) -bimodules, where horizontal composition is given by the derived tensor product $\otimes^{\mathbb{L}}$. Recall that if M is a right S -module, and N is a left S -module, then $M \otimes_S^{\mathbb{L}} N$ is equivalent to the two-sided bar construction, $\text{Bar}(M, S, N)$.

Example 3.3. When the 0-cells are given by ring spectra and the 1- and 2-cells come from the homotopy category of (R, S) -bimodules for ring spectra R and S , then the horizontal composition is given by the derived smash product of spectra. Note that as in the previous example, $\text{Bar}(M, S, N) \simeq M \wedge_S^{\mathbb{L}} N$, see [EKMM97, IV.7.5], and so we may also consider this horizontal composition as the two-sided bar construction.

In this paper, we will introduce two bicategories formed by bicomodules instead of bimodules. Their horizontal compositions will be given by the (derived) cotensor product.

Proposition 3.4. *Let $(\mathcal{C}, \otimes, \mathbb{I})$ be a symmetric monoidal category, and let C be a flat coalgebra in \mathcal{C} . Then $({}_C\text{CoMod}_{\mathcal{C}}(\mathcal{C}), \square_C, C)$ is a monoidal category. Further, it is symmetric monoidal if C is cocommutative.*

Definition 3.5. Define CoMod to be the bicategory whose 0-cells are coalgebras in $\text{Mod}_{\mathbb{k}}$, and whose 1-cells, 2-cells, and vertical compositions are given by the category ${}_C\text{CoMod}_D(\text{Mod}_{\mathbb{k}})$. The unit U_C is the (C, C) -bicomodule C , and horizontal composition is given by the cotensor product:

$$\odot : \mathcal{B}(C, D) \times \mathcal{B}(D, E) \rightarrow \mathcal{B}(C, E)$$

$$(M, N) \mapsto M \odot N := M \square_D N.$$

For (C, D) -bicomodule M , (D, E) -bicomodule N , and (E, F) -bicomodule P , the natural isomorphisms

$$a : (M \odot N) \odot P \xrightarrow{\cong} M \odot (N \odot P), \quad \ell : U_C \odot M \xrightarrow{\cong} M, \quad r : M \odot U_D \xrightarrow{\cong} M,$$

follow as in [Doi81] from the natural isomorphisms

$$(M \square_D N) \square_E P \cong M \square_D (N \square_E P), \quad C \square_C M \cong M, \quad M \square_D D \cong M.$$

More specifically, these maps are given by

$$\begin{aligned} a : (M \square_D N) \square_E P &\xrightarrow{\cong} M \square_D (N \square_E P) & \ell : C \square_C M &\xrightarrow{\cong} M & r : M \square_D D &\xrightarrow{\cong} M \\ (m \otimes n) \otimes p &\longmapsto m \otimes (n \otimes p) & c \otimes m &\longmapsto \varepsilon_C(c)m & m \otimes d &\longmapsto \varepsilon_D(d)m. \end{aligned}$$

We now consider the derived case. We first need a homotopy theory of bicomodules. Let $\mathbf{M} = \text{Ch}_{\mathbb{k}}^{\geq 0}$ be the category of non-negative chain complexes over \mathbb{k} . There is a model structure on $\text{Ch}_{\mathbb{k}}^{\geq 0}$ in which weak equivalences are quasi-isomorphisms, cofibrations are monomorphisms, and fibrations are positive levelwise epimorphisms, see [Hov99]. It is a combinatorial symmetric monoidal model category with its usual tensor product, and a simplicial model category via the Dold-Kan correspondence. In addition, every object is cofibrant and fibrant.

Proposition 3.6. *Let C and D be dg-coalgebras over \mathbb{k} . There exists a simplicial combinatorial model structure on ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ in which*

- *the weak equivalences \mathbf{W} are the morphisms of (C, D) -bicomodules that are quasi-isomorphisms in $\text{Ch}_{\mathbb{k}}^{\geq 0}$;*
- *the cofibrations are the morphisms of (C, D) -bicomodules that are monomorphisms in $\text{Ch}_{\mathbb{k}}^{\geq 0}$.*

In particular, every object is cofibrant.

Proof. The model structure is left-induced in the sense of [HKRS17] via the forgetful-cofree adjunction

$${}_C\text{CoMod}_D \left(\text{Ch}_{\mathbb{k}}^{\geq 0} \right) \xrightleftharpoons[C \otimes - \otimes D]{U} \text{Ch}_{\mathbb{k}}^{\geq 0}.$$

Indeed, this follows from [Pér24b, 2.12] and [HKRS17, 6.3.7] by Remark 1.15. \square

The above result is true more generally for unbounded chain complexes over any ring. However, it is not known if this forms an algebraic model for homotopy coherent comodules in this generality.

Definition 3.7. A (connective) *dg-coalgebra over \mathbb{k}* is a coalgebra C in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. We say it is *simply connected* if $C_0 \cong \mathbb{k}$ and $C_1 = 0$.

When C and D are simply connected, we show the model structure from the previous proposition defines the correct homotopy type, see Theorem A.4. Fibrant objects in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ are retracts of Postnikov towers, see more details in Appendix A.

In order to define horizontal composition for the bicategory in the derived setting, we now need to derive the cotensor product. However, it is more subtle than the usual tensor product of modules because the cotensor product is neither a left nor a right adjoint in general. Nevertheless, one can right derive the cotensor product using methods of [Pér24b, Pér24a]. We leave the details in Appendix A, but essentially, we show that the cotensor product of fibrant bicomodules is again fibrant (Proposition A.10) and that the cotensor preserves weak equivalences on fibrant objects (Proposition A.17). Moreover, just as the derived tensor product can be interpreted as a two-sided bar construction, the derived cotensor product can be interpreted as a two-sided cobar construction, as we shall now explain.

Definition 3.8. Let C and D be simply connected dg-coalgebras over \mathbb{k} , with comultiplication and counit of C given by $\Delta: C \rightarrow C \otimes C$ and $\varepsilon: C \rightarrow \mathbb{k}$ respectively. Let M be a (D, C) -bicomodule in $\text{Ch}_{\mathbb{k}}^{\geq 0}$, and let $\rho: M \rightarrow M \otimes C$ denote its right coaction over C . The *two-sided cosimplicial cobar construction* $\Omega^\bullet(M, C, C)$ of M is the cosimplicial object in ${}_D\text{CoMod}_C(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$M \otimes C \rightrightarrows M \otimes C \otimes C \rightrightarrows \cdots,$$

defined as follows.

- For all $n \geq 0$, $\Omega^n(M, C, C) = M \otimes C^{\otimes n+1}$.
- The zeroth coface map $d^0: \Omega^n(M, C, C) \rightarrow \Omega^{n+1}(M, C, C)$ is given by $d^0 = \rho \otimes \text{id}_{C^{n+1}}$.
- For $1 \leq i \leq n+1$, the i^{th} coface map $d^i: \Omega^n(M, C, C) \rightarrow \Omega^{n+1}(M, C, C)$ is given by

$$d^i = \text{id}_M \otimes \text{id}_{C^{\otimes i-1}} \otimes \Delta \otimes \text{id}_{C^{\otimes n+1-i}}.$$

- For all $0 \leq j \leq n$, the j^{th} codegeneracy map $s^j: \Omega^{n+1}(M, C, C) \rightarrow \Omega^n(M, C, C)$ is given by

$$s^j = \text{id}_M \otimes \text{id}_{C^{\otimes j}} \otimes \varepsilon \otimes \text{id}_{C^{\otimes n+1-j}}.$$

Since ${}_D\text{CoMod}_C(\text{Ch}_{\mathbb{k}}^{\geq 0})$ is a simplicial model category, homotopy limits over cosimplicial diagrams are computed as in [Hir03, 18.1.8]. We denote the homotopy limit of the cosimplicial diagram $\Omega^\bullet(M, C, C)$ in ${}_D\text{CoMod}_C(\text{Ch}_{\mathbb{k}}^{\geq 0})$ by $\Omega(M, C, C)$, and we say it is the *two-sided cobar resolution of M* . By [Pér24b, 2.5], we have $M \simeq \Omega(M, C, C)$ as a (D, C) -bicomodule if M is fibrant as a left D -comodule. Notice that each object in the cosimplicial diagram $\Omega^\bullet(M, C, C)$ is a fibrant right C -comodule by Lemma A.8. Thus $\Omega(M, C, C)$ is a fibrant (D, C) -bicomodule by [Hir03, 18.5.2] if M is a fibrant left D -comodule.

Definition 3.9. Let C , D and E be simply connected dg-coalgebras over \mathbb{k} . Let M be a (D, C) -bicomodule and N be a (C, E) -bicomodule. We define the *two-sided cosimplicial cobar construction of M and N* to be the cosimplicial object $\Omega^\bullet(M, C, N)$ in ${}_D\text{CoMod}_E(\text{Ch}_{\mathbb{k}}^{\geq 0})$ given by $\Omega^\bullet(M, C, C) \square_C N$. We write $\Omega(M, C, N)$ for the homotopy limit of $\Omega^\bullet(M, C, N)$ in ${}_D\text{CoMod}_E(\text{Ch}_{\mathbb{k}}^{\geq 0})$. As noted in [Pér24b, 2.5], it is equivalent to compute the homotopy limit in $\text{Ch}_{\mathbb{k}}^{\geq 0}$.

Proposition 3.10. Let C , D and E be simply connected dg-coalgebras over \mathbb{k} . Let M be a (D, C) -bicomodule and N be a fibrant (C, E) -bicomodule. Then we obtain an equivalence $\Omega(M, C, C) \square_C N \simeq \Omega(M, C, N)$ as (D, E) -bicomodules.

Proof. This argument is similar to discussion in [Pér24b, 4.2] (the cocommutative requirement there was not needed). \square

Remark 3.11. By the Dold-Kan correspondence, since C is a simply connected dg-coalgebra, the two-sided cosimplicial cobar resolution $\Omega(M, C, N)$ from Definition 3.8 is quasi-isomorphic to $\underline{\Omega}(M, C, N)$, the *conormalized cobar resolution of M and N over C* , which we now define. We first establish the following notation conventions.

- Given V a graded \mathbb{k} -module, we define

$$T(V) = \bigoplus_{n \geq 0} V^{\otimes n}.$$

Elements in the summands are denoted $v_1 | \cdots | v_n$, where $v_i \in V$.

- Let s^{-1} denote the desuspension functor on graded \mathbb{k} -modules where, for $V = \bigoplus_{i \in \mathbb{Z}} V_i$, we define $(s^{-1}V)_i = V_{i+1}$. Given a homogeneous element v in V , we write $s^{-1}v$ for the corresponding element in $s^{-1}V$.
- We denote the kernel of the counit $\varepsilon: C \rightarrow \mathbb{k}$ by \underline{C} , often referred to as the coideal of C .

The conormalized cobar resolution of C is the chain complex $\underline{\Omega}C := (T(s^{-1}\underline{C}), d_\Omega)$ where, if d denotes the differential on C , then

$$\begin{aligned} d_\Omega(s^{-1}c_1 | \cdots | s^{-1}c_n) &= \sum_{j=1}^n \pm s^{-1}c_1 | \cdots | s^{-1}(dc_j) | \cdots | s^{-1}c_n \\ &\quad + \sum_{j=1}^n \sum_{(c_j)} \pm s^{-1}c_1 | \cdots | s^{-1}c_{j(1)} | s^{-1}c_{j(2)} | \cdots | s^{-1}c_n, \end{aligned}$$

where $\Delta(c_j) = \sum_{(c_j)} c_{j(1)} \otimes c_{j(2)}$ denotes the (reduced) comultiplication of \underline{C} on the element c_j using the Sweedler notation. To determine the signs above, one needs to apply the Koszul rule. More generally, we define the conormalized cobar resolution of M and N over C to be the chain complex $\underline{\Omega}(M, C, N) := (M \otimes T(s^{-1}\underline{C}) \otimes N, \delta)$, where up to Koszul sign, the differential δ is defined as

$$\begin{aligned} \delta(m \otimes s^{-1}c_1 | \cdots | s^{-1}c_n \otimes n) &= dm \otimes s^{-1}c_1 | \cdots | s^{-1}c_n \otimes n \\ &\quad \pm \sum_{(m)} m_{(0)} \otimes s^{-1}m_{(1)} | s^{-1}c_1 | \cdots | s^{-1}c_n \otimes n \\ &\quad \pm m \otimes d_\Omega(s^{-1}c_1 | \cdots | s^{-1}c_n) \otimes n \\ &\quad \pm m \otimes s^{-1}c_1 | \cdots | s^{-1}c_n \otimes dn \\ &\quad \pm \sum_{(n)} m \otimes s^{-1}c_1 | \cdots | s^{-1}c_n | s^{-1}n_{(1)} \otimes n_{(0)}, \end{aligned}$$

where d denotes either the differential on M or N , and $m \mapsto \sum_{(m)} m_{(0)} \otimes m_{(1)}$ denotes the coaction of $M \rightarrow M \otimes C$ applied to an element m , and $n \mapsto \sum_{(n)} n_{(1)} \otimes n_{(0)}$ denotes the coaction of $N \rightarrow C \otimes N$ applied to an element n .

By the Dold-Kan correspondence, there is an isomorphism of categories between cosimplicial objects and non-positive cochain complexes of bicomodules given by the conormalization functor

$$N^*: \left({}_D\text{CoMod}_E(\text{Ch}_{\mathbb{k}}^{\geq 0}) \right)^\Delta \xrightarrow{\cong} \text{coCh}^{\leq 0} \left({}_D\text{CoMod}_E(\text{Ch}_{\mathbb{k}}^{\geq 0}) \right).$$

Therefore, we denote $N^*(\Omega^\bullet(M, C, N))$ by $\underline{\Omega}^\bullet(M, C, N)$. It is a double complex, and one can show that its total complex is precisely $\underline{\Omega}(M, C, N)$. A word of warning however: a double complex in general has two possible totalizations, one given using coproducts and one using products (see [Wei94, 1.2.6]). The product-total complex of $\underline{\Omega}^\bullet(M, C, N)$ is quasi-isomorphic to the homotopy limit of $\Omega^\bullet(M, C, N)$, i.e., $\Omega(M, C, N)$ (see [Bun12, 4.23] for instance). On the other hand, the coproduct-total complex of $\underline{\Omega}^\bullet(M, C, N)$ is $\underline{\Omega}(M, C, N)$. Of course, if the double complex is bounded, these are equal. For instance, when C is simply-connected, then $\underline{\Omega}^{-q}(M, C, N)_p = 0$ for $0 \leq p \leq 2q - 1$, and is therefore bounded. Thus when C is simply-connected, we obtain a quasi-isomorphism

$$\Omega(M, C, N) \simeq \underline{\Omega}(M, C, N).$$

But in general, if C is not simply-connected, no such claim can be made.

Remark 3.12. Let C be a simply-connected coalgebra in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. Let M and N be left and right C -comodules respectively. As noted in [Rav86, A1.2.12], for all $i \geq 0$ we have an isomorphism

$$\text{CoTor}_C^i(M, N) \cong H^i(\underline{\Omega}^\bullet(M, C, N)).$$

Therefore, on the homotopy category of comodules, we have a derived cotensor product, which we denote by $\hat{\square}$. In fact, $M\hat{\square}_C N$ is quasi-isomorphic to the two-sided cobar resolution $\Omega(M, C, N)$, see Corollary A.18. The derived cotensor preserves any homotopy coherent coactions, see [Pér24b].

Having established the necessary definitions for each of the components, we now define the appropriate bicategory for the derived bicomodule setting.

Definition 3.13. Define \mathcal{DCoMod} to be the bicategory whose 0-cells are simply connected dg-coalgebras over \mathbb{k} , and whose 1-cells, 2-cells, and vertical compositions are given by the homotopy category of the model category ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$. The unit U_C is the fibrant (C, C) -bicomodule C , and horizontal composition is given by the derived cotensor product of bicomodules. Given a fibrant (C, D) -bicomodule M and a fibrant (D, E) -bicomodule N , their horizontal composition $M \odot N$ is the fibrant (C, E) -bicomodule

$$M \square_D N \simeq M \hat{\square}_D N \simeq \Omega(M, D, N).$$

If P is a fibrant (E, F) -bicomodule, then we define the natural isomorphisms:

$$a: (M \square_D N) \square_E P \xrightarrow{\cong} M \square_D (N \square_E P),$$

$$\ell: C \square_C M \xrightarrow{\cong} M, \quad r: M \square_D D \xrightarrow{\cong} M,$$

as follows. The isomorphism a follows from the natural isomorphism $(M \otimes N) \otimes P \cong M \otimes (N \otimes P)$, and therefore automatically respects the pentagon identity. The natural isomorphisms ℓ and r are induced by the counits $C \rightarrow \mathbb{k}$ and $D \rightarrow \mathbb{k}$ respectively, and the triangle identities follow for the cotensor products since they hold for the tensor product.

Since we shall need it in next section, we provide an explicit definition of a , ℓ , and r from above, where we instead use the cobar construction as a model for the derived cotensor product. The associative equivalence

$$a: \Omega(\Omega(M, D, N), E, P) \xrightarrow{\cong} \Omega(M, D, \Omega(N, E, P))$$

is induced by an isomorphism of cosimplicial objects $\Omega^\bullet(\Omega^\bullet(M, D, N), E, P) \cong \Omega^\bullet(M, D, \Omega^\bullet(N, E, P))$, using Corollary A.21 from Appendix A. Indeed, the (i, j) -spot corresponds to the (j, i) -spot up to isomorphism

$$(M \otimes D^{\otimes j} \otimes N) \otimes E^{\otimes i} \otimes P \xrightarrow{\cong} M \otimes D^{\otimes j} \otimes (N \otimes E^{\otimes i} \otimes P)$$

from the usual associative isomorphism. The equivalences $\ell: \Omega(C, C, M) \xrightarrow{\cong} M$ and $r: \Omega(M, D, D) \xrightarrow{\cong} M$ are defined as in Definition 3.8, and are induced by the counits. For instance, the map $C^{\otimes n+1} \otimes M \rightarrow M$ is induced by repeatedly applying the counit on C .

4. COHOCHSCHILD HOMOLOGY AS A SHADOW

Here we show our main results, namely that coHochschild homology provides a shadow structure on the previously defined bicategories of bicomodules.

Definition 4.1 ([Pon10, PS13]). A *shadow functor* for a bicategory \mathcal{B} consists of functors

$$\langle\langle - \rangle\rangle_C: \mathcal{B}(C, C) \longrightarrow \mathbf{T}$$

for every $C \in \mathcal{B}$ and some fixed category \mathbf{T} equipped with a natural isomorphism for $M \in \mathcal{B}(C, D)$, $N \in \mathcal{B}(D, C)$:

$$\theta: \langle\langle M \odot N \rangle\rangle_C \xrightarrow{\cong} \langle\langle N \odot M \rangle\rangle_D.$$

For $P \in \mathcal{B}(C, C)$, these functors must satisfy the following commutative diagrams

$$\begin{array}{ccccc} \langle\langle (M \odot N) \odot P \rangle\rangle_C & \xrightarrow{\theta} & \langle\langle P \odot (M \odot N) \rangle\rangle_C & \xrightarrow{\langle\langle a \rangle\rangle} & \langle\langle (P \odot M) \odot N \rangle\rangle_C \\ \langle\langle a \rangle\rangle \downarrow & & & & \uparrow \theta \\ \langle\langle M \odot (N \odot P) \rangle\rangle_C & \xrightarrow{\theta} & \langle\langle (N \odot P) \odot M \rangle\rangle_D & \xrightarrow{\langle\langle a \rangle\rangle} & \langle\langle N \odot (P \odot M) \rangle\rangle_D \end{array}$$

$$\begin{array}{ccccc}
\langle\langle P \odot U_C \rangle\rangle_C & \xrightarrow{\theta} & \langle\langle U_C \odot P \rangle\rangle_C & \longrightarrow & \langle\langle P \odot U_C \rangle\rangle_C \\
& \searrow \langle\langle r \rangle\rangle & \downarrow \langle\langle \ell \rangle\rangle & \swarrow \langle\langle r \rangle\rangle & \\
& & \langle\langle P \rangle\rangle_C & &
\end{array}$$

In this case, we say \mathcal{B} is a *shadowed bicategory*, and we write $(\mathcal{B}, \langle\langle - \rangle\rangle)$ for the bicategory and its shadow.

Example 4.2. We can now consider shadows for the bicategories that we introduced in Examples 3.2 and 3.3. For a ring R , the 0^{th} Hochschild homology, $\mathrm{HH}_0(R, -)$, is a shadow on the bicategory with 1- and 2-cells from the category of (R, R) -bimodules to the category of abelian groups Ab :

$$\begin{aligned}
\langle\langle - \rangle\rangle_R: {}_R\mathrm{Mod}_R &\rightarrow \mathrm{Ab} \\
M &\mapsto R \otimes_{R \otimes R^{\mathrm{op}}} M \cong \mathrm{HH}_0(R, M).
\end{aligned}$$

More generally, a Dennis-Waldhausen Morita argument shows that Hochschild homology, $\mathrm{HH}(R, -)$, is a shadow in the derived setting [Wal79]. Further, [BM12] shows that topological Hochschild homology, $\mathrm{THH}(R, -)$, is a shadow to the homotopy category of spectra:

$$\begin{aligned}
\langle\langle - \rangle\rangle_R: \mathrm{Ho}({}_R\mathrm{Mod}_R) &\rightarrow \mathrm{Ho}(\mathrm{Sp}) \\
M &\mapsto \mathrm{THH}(R, M).
\end{aligned}$$

Since (topological) Hochschild homology provides a bicategorical shadow for the setting of modules, we want to consider the analogue of this construction for the context of comodules. Work of Doi defines coHochschild homology, denoted coHH , as an invariant of coalgebras analogous to Hochschild homology.

Definition 4.3 ([Doi81]). For a commutative ring R , a coassociative, counital R -coalgebra C , and a (C, C) -bicomodule M , build the cochain complex $\mathcal{H}(M, C)$:

$$\cdots \longleftarrow M \otimes_R C \otimes_R C \longleftarrow M \otimes_R C \longleftarrow M \longleftarrow 0,$$

as follows. Let $\mathcal{H}^r(M, C) = M \otimes_R C^{\otimes r}$ for $r \geq 0$ with coboundary map $\delta^r: \mathcal{H}^r(M, C) \rightarrow \mathcal{H}^{r+1}(M, C)$ defined by

$$\delta^r = \sum_{i=0}^{r+1} (-1)^i d_i,$$

for d_i given by

$$d_i = \begin{cases} \rho \otimes \mathrm{id}_C^{\otimes r} & i = 0 \\ \mathrm{id}_M \otimes \mathrm{id}_C^{\otimes i-1} \otimes \Delta \otimes \mathrm{id}_C^{\otimes (r-i)} & 1 \leq i \leq r \\ \tilde{t} \circ (\lambda \otimes \mathrm{id}_C^{\otimes r}) & i = r+1, \end{cases}$$

where $\rho: M \rightarrow M \otimes_R C$ denotes the right coaction, $\lambda: M \rightarrow C \otimes_R M$ denotes the left coaction, and \tilde{t} is the map that twists the first factor to the last. Then the q^{th} -coHochschild homology of C with coefficients in M is given by the cohomology of the cochain complex

$$\mathrm{coHH}_q(M, C) := H^q(\mathcal{H}(M, C)).$$

Remark 4.4. We can see that $\mathcal{H}(M, C) = \underline{\Omega}(M, C^e, C)$ as in Remark 3.11, where $C^e = C \otimes_R C^{\mathrm{op}}$, and in particular we get that $\mathrm{coHH}_q(M, C) \cong \mathrm{CoTor}_{C^e}^q(M, C)$, see Definition 1.19 and Remark 3.12. In particular $\mathrm{coHH}_0(M, C) \cong M \square_{C^e} C$.

Theorem 4.5. The 0^{th} coHochschild homology coHH_0 is a shadow on the bicategory CoMod . That is, it gives a family of functors

$$\begin{aligned}
\mathrm{coHH}_0(-, C): {}_C\mathrm{CoMod}_C &\rightarrow \mathrm{Mod}_{\mathbb{k}} \\
M &\mapsto \mathrm{coHH}_0(M, C)
\end{aligned}$$

that satisfy the required shadow properties.

Proof. Let C and D be coalgebras over \mathbb{k} . Given M a (C, D) -bicomodule and N a (D, C) -bicomodule, we need to show that we have an isomorphism

$$\theta: \text{coHH}_0(M \square_D N, C) \longrightarrow \text{coHH}_0(N \square_C M, D).$$

Using Sweedler notation, we denote the coactions on M as follows:

$$\begin{aligned} M &\longrightarrow C \otimes M, & M &\longrightarrow M \otimes D \\ m &\longmapsto \sum_{(m^C)} m_{(1)}^C \otimes m_{(0)}^C, & m &\longmapsto \sum_{(m^D)} m_{(0)}^D \otimes m_{(1)}^D, \end{aligned}$$

and the coactions on N as follows:

$$\begin{aligned} N &\longrightarrow D \otimes N, & N &\longrightarrow N \otimes C \\ n &\longmapsto \sum_{(n^D)} n_{(1)}^D \otimes n_{(0)}^D, & n &\longmapsto \sum_{(n^C)} n_{(0)}^C \otimes n_{(1)}^C. \end{aligned}$$

Recall that $\text{coHH}_0(M \square_D N, C)$ is defined as the kernel of

$$\delta^0: M \square_D N \rightarrow (M \square_D N) \otimes C.$$

The desired isomorphism will be induced by

$$\begin{aligned} \tau: M \otimes N &\longrightarrow N \otimes M \\ m \otimes n &\longmapsto n \otimes m. \end{aligned}$$

We need to verify that if we restrict τ to $\text{coHH}_0(M \square_D N, C)$, we indeed corestrict to $\text{coHH}_0(N \square_C M, D)$. In other words, we need to verify that if $m \otimes n \in \text{coHH}_0(M \square_D N, C)$, then $n \otimes m \in \text{coHH}_0(N \square_C M, D)$. Notice that because $m \otimes n \in \text{coHH}_0(M \square_D N, C)$, it follows that

$$\begin{aligned} (1) \quad &\sum_{(m^D)} m_{(0)}^D \otimes m_{(1)}^D \otimes n = \sum_{(n^D)} m \otimes n_{(1)}^D \otimes n_{(0)}^D, \text{ since } m \otimes n \in M \square_D N; \\ (2) \quad &\sum_{(n^C)} m \otimes n_{(0)}^C \otimes n_{(1)}^C = \sum_{(m^C)} m_{(0)}^C \otimes n \otimes m_{(1)}^C, \text{ since } m \otimes n \in \ker(\delta^0). \end{aligned}$$

To see that $n \otimes m \in \text{coHH}_0(N \square_C M, D)$, notice that $n \otimes m \in N \square_C M$ follows from (2) above, while $n \otimes m \in \ker(\delta^0)$ follows from (1) above. Therefore we have obtained the desired homomorphism, θ . An analogous argument defines its inverse, and thus θ is an isomorphism.

Next we must show that for a (C, C) -bicomodule P , the following diagram is commutative:

$$\begin{array}{ccccc} \text{coHH}_0((M \square_D N) \square_C P, C) & \xrightarrow{\theta} & \text{coHH}_0(P \square_C (M \square_D N), C) & \xrightarrow{\langle\langle a \rangle\rangle} & \text{coHH}_0((P \square_C M) \square_D N, C) \\ \langle\langle a \rangle\rangle \downarrow & & & & \uparrow \theta \\ \text{coHH}_0(M \square_D (N \square_C P), C) & \xrightarrow{\theta} & \text{coHH}_0((N \square_C P) \square_C M, D) & \xrightarrow{\langle\langle a \rangle\rangle} & \text{coHH}_0(N \square_C (P \square_C M), D) \end{array}$$

We check its commutativity directly by applying the definition of θ given above and a from Definition 3.5:

$$\begin{array}{ccccc} (m \otimes n) \otimes p & \longmapsto & p \otimes (m \otimes n) & \longmapsto & (p \otimes m) \otimes n. \\ \downarrow & & & & \uparrow \\ m \otimes (n \otimes p) & \longmapsto & (n \otimes p) \otimes m & \longmapsto & n \otimes (p \otimes m) \end{array}$$

Similarly, we need to check the commutativity of the diagram below:

$$\begin{array}{ccccc} \text{coHH}_0(P \square_C C, C) & \xrightarrow{\theta} & \text{coHH}_0(C \square_C P, C) & \xrightarrow{\theta} & \text{coHH}_0(P \square_C C, C) \\ & \searrow \langle\langle r \rangle\rangle & \downarrow \langle\langle \ell \rangle\rangle & \swarrow \langle\langle r \rangle\rangle & \\ & & \text{coHH}_0(P, C) & & \end{array}$$

This follows again by applying the definitions of θ , r , and ℓ (see Definition 3.5):

$$\begin{array}{ccccc} p \otimes c & \xrightarrow{\quad} & c \otimes p & \xrightarrow{\quad} & p \otimes c \\ & \searrow & \downarrow & \swarrow & \\ & & \varepsilon(c)p & & \end{array}$$

This proves that the 0^{th} coHochschild homology is a shadow in this bicategorical setting. \square

Having established that coHH_0 is a bicategorical shadow on CoMod , we now consider the derived setting. Recall that $\underline{\Omega}$ of Remark 3.11 is not invariant under quasi-isomorphisms in general. Therefore particular care is required in order to show that coHH is a shadow in the derived setting. To do so, we must instead consider coalgebras in chain complexes that are simply connected. Extending the definitions of [HS21] and [BGH⁺18], the first author introduced in [Kla22, 2.8] the notion of coHochschild homology with coefficients for any model category with a symmetric monoidal structure.

Definition 4.6. Let $(\mathbf{M}, \otimes, \mathbb{I})$ be a symmetric monoidal category with a model structure, and let $C \in \mathbf{M}$ be a coalgebra with coassociative comultiplication $\Delta : C \rightarrow C \otimes C$ and counit $\varepsilon : C \rightarrow \mathbb{I}$. Further, let M be a (C, C) -bicomodule with left and right coactions $\lambda : M \rightarrow C \otimes M$ and $\rho : M \rightarrow M \otimes C$ respectively. Define $\text{coHH}^{\mathbf{M}}(M, C)^{\bullet}$ to be the cosimplicial object with r -simplices $\text{coHH}^{\mathbf{M}}(M, C)^r = M \otimes C^{\otimes r}$, with coface maps

$$d_i = \begin{cases} \rho \otimes \text{id}_C^{\otimes r} & i = 0 \\ \text{id}_M \otimes \text{id}_C^{\otimes i-1} \otimes \Delta \otimes \text{id}_C^{\otimes (r-i)} & 1 \leq i \leq r \\ \tilde{t} \circ (\lambda \otimes \text{id}_C^{\otimes r}) & i = r+1, \end{cases}$$

where \tilde{t} is the map that twists the first factor to the last, and with codegeneracy maps $s_i : M \otimes C^{\otimes (r+1)} \rightarrow M \otimes C^{\otimes r}$, for $0 \leq i \leq r$,

$$s_i = \text{id}_M \otimes \text{id}_C^{\otimes i} \otimes \varepsilon \otimes \text{id}_C^{\otimes r-i}.$$

This gives a cosimplicial object of the form

$$\cdots \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M \otimes C \otimes C \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M \otimes C \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M.$$

The *coHochschild homology* in \mathbf{M} of the coalgebra C with coefficients in M is then defined by

$$\text{coHH}^{\mathbf{M}}(M, C) = \text{holim}_{\Delta} (\text{coHH}^{\mathbf{M}}(M, C)^{\bullet}).$$

Remark 4.7. In [HR21b], the authors give an ∞ -categorical definition of topological Hochschild homology with coefficients, extending the approach of [NS18]. In [BP23], a dual approach of [NS18] for topological coHochschild homology was given. Mimicking this approach, we can define a relative cyclic cobar construction $\text{CoBar}_{\mathcal{C}}^{\bullet}(-, -) : \text{CoAlg}_{\mathcal{AB}}(\mathcal{C}) \rightarrow \text{Fun}(\mathcal{N}(\Delta), \mathcal{C})$ on a symmetric monoidal ∞ -category \mathcal{C} , where $\text{CoAlg}_{\mathcal{AB}}(\mathcal{C})$ denotes the ∞ -category of pairs (M, C) consisting of an \mathbb{E}_1 -coalgebra C and a bicomodule M over C . Essentially, the construction $\text{CoBar}_{\mathcal{C}}^{\bullet}(C, M)$ is making precise the diagram in \mathcal{C} :

$$\cdots \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M \otimes C \otimes C \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M \otimes C \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} M.$$

Denote $\text{coHH}^{\mathcal{C}}(M, C)$ the totalization in \mathcal{C} of the cosimplicial object obtained by the cyclic cobar construction $\text{CoBar}_{\mathcal{C}}^{\bullet}(M, C)$. Per usual, we denote $\text{coHH}^{\mathcal{C}}(C, C)$ by $\text{coHH}^{\mathcal{C}}(C)$. If \mathbf{M} is a combinatorial symmetric monoidal model category with class of weak equivalences denoted by \mathbf{W} , then one could also obtain a symmetric monoidal ∞ -category $\mathcal{N}(\mathbf{M}_c)[\mathbf{W}^{-1}]$ obtained by Dwyer–Kan localization. Then one could consider $\text{coHH}^{\mathcal{N}(\mathbf{M}_c)[\mathbf{W}^{-1}]}(C, M)$. Suppose both C and M are cofibrant in \mathbf{M} . Then, similar to [BP23, 2.13] we obtain a weak equivalence

$$\text{coHH}^{\mathbf{M}}(M, C) \simeq \text{coHH}^{\mathcal{N}(\mathbf{M}_c)[\mathbf{W}^{-1}]}(M, C).$$

Example 4.8. Our main example of interest is $\mathbf{M} = \text{Ch}_{\mathbb{k}}^{\geq 0}$. In this case, when C is simply connected, the definition of coHochschild homology coincides with [HPS09], see also Remark 3.11 and [BGH⁺18, 2.11].

In what follows, we write $\text{coHH}^{\text{Ch}_{\mathbb{k}}^{\geq 0}}$ simply as coHH . We show in Proposition A.23 that $\text{coHH}(M, C) \simeq \Omega(M, C^e, C)$ for C simply connected and M a fibrant (C, C) -bicomodule.

Theorem 4.9. *CoHochschild homology defines a shadow on the bicategory of derived bicomodules over simply connected coalgebras $\mathcal{D}\text{CoMod}$.*

Proof. Let C and D be simply connected coalgebras in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. Given M a fibrant (C, D) -bicomodule, and N a fibrant (D, C) -bicomodule, we need to show that we have a quasi-isomorphism:

$$\theta: \text{coHH}(M \hat{\square}_D N, C) \longrightarrow \text{coHH}(N \hat{\square}_C M, D).$$

As M and N are fibrant, the derived cotensor $M \hat{\square}_D N$ is modeled by $M \square_D N$ or $\Omega(M, D, N)$. We choose the cobar construction as a model: it automatically gives us the desired quasi-isomorphism at the cost of some combinatorics. To provide this equivalence, we apply Corollary A.22 from Appendix A. In particular, we claim that θ is induced by an isomorphism of bicosimplicial objects:

$$\theta: \text{coHH}(\Omega^\bullet(M, D, N), C)^\bullet \xrightarrow{\cong} \text{coHH}(\Omega^\bullet(N, C, M), D)^\bullet.$$

This is the dual to the Dennis-Waldhausen Morita argument. Indeed, we define θ by the usual shuffling isomorphism in the bicosimplicial map below, in which the rows correspond to the two-sided cobar construction, while the columns correspond to the coHochschild complex. For ease of understanding, the diagrams below have been color-coded to illustrate this isomorphism via shuffling.

$$\begin{array}{ccccc}
M \otimes N & \rightrightarrows & M \otimes N \otimes C & \rightrightarrows & M \otimes N \otimes C^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
M \otimes D \otimes N & \rightrightarrows & M \otimes D \otimes N \otimes C & \rightrightarrows & M \otimes D \otimes N \otimes C^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
M \otimes D^{\otimes 2} \otimes N & \rightrightarrows & M \otimes D^{\otimes 2} \otimes N \otimes C & \rightrightarrows & M \otimes D^{\otimes 2} \otimes N \otimes C^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
\vdots & & \vdots & & \vdots & & \\
& & \theta & & & & \\
& & \downarrow & & & & \\
N \otimes M & \rightrightarrows & N \otimes M \otimes D & \rightrightarrows & N \otimes M \otimes D^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
N \otimes C \otimes M & \rightrightarrows & N \otimes C \otimes M \otimes D & \rightrightarrows & N \otimes C \otimes M \otimes D^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
N \otimes C^{\otimes 2} \otimes M & \rightrightarrows & N \otimes C^{\otimes 2} \otimes M \otimes D & \rightrightarrows & N \otimes C^{\otimes 2} \otimes M \otimes D^{\otimes 2} & \rightrightarrows & \dots \\
\Downarrow & & \Downarrow & & \Downarrow & & \\
\vdots & & \vdots & & \vdots & &
\end{array}$$

The (i, j) -spot in $\text{coHH}(\Omega^\bullet(M, D, N), C)^\bullet$ is isomorphic to the (j, i) -spot in $\text{coHH}(\Omega^\bullet(N, C, M), D)^\bullet$:

$$\theta: M \otimes D^{\otimes i} \otimes N \otimes C^{\otimes j} \xrightarrow{\cong} N \otimes C^{\otimes j} \otimes M \otimes D^{\otimes i},$$

which incorporates a Koszul sign. Moreover, the map θ is compatible with the cofaces and codegeneracies. By [Hir03, 18.5.3], we obtain the equivalence

$$\theta: \text{coHH}(\Omega(M, D, N), C) \xrightarrow{\cong} \text{coHH}(\Omega(N, C, M), D),$$

using the model of homotopy limit as in [Hir03, 18.1.8], providing the desired quasi-isomorphism θ .

Given a fibrant (C, C) -bicomodule P , we next must show that the following diagram commutes:

$$\begin{array}{ccccc}
\mathrm{coHH}(\Omega(M, D, N), C, P), C & \xrightarrow{\theta} & \mathrm{coHH}(\Omega(P, C, \Omega(M, D, N)), C) & \xrightarrow{\langle\langle a \rangle\rangle} & \mathrm{coHH}(\Omega(P, C, M), D, N), C \\
\downarrow \langle\langle a \rangle\rangle & & & & \uparrow \theta \\
\mathrm{coHH}(\Omega(M, D, \Omega(N, C, P)), C) & \xrightarrow{\theta} & \mathrm{coHH}(\Omega(N, C, P), C, M), D & \xrightarrow{\langle\langle a \rangle\rangle} & \mathrm{coHH}(\Omega(N, C, \Omega(P, C, M)), D).
\end{array}$$

This follows a similar argument as above. We need to consider an isomorphism of “tri-cosimplicial” isomorphisms in which we keep track of the swapping of the grading:

$$\begin{array}{ccccc}
((M \otimes D^{\otimes k} \otimes N) \otimes C^{\otimes j} \otimes P) \otimes C^{\otimes i} & \xrightarrow{\theta} & (P \otimes C^{\otimes i} \otimes (M \otimes D^{\otimes k} \otimes N)) \otimes C^{\otimes j} & \xrightarrow{\langle\langle a \rangle\rangle} & ((P \otimes C^{\otimes i} \otimes M) \otimes D^{\otimes k} \otimes N) \otimes C^{\otimes j}. \\
\downarrow \langle\langle a \rangle\rangle & & & & \uparrow \theta \\
(M \otimes D^{\otimes k} \otimes (N \otimes C^{\otimes j} \otimes P)) \otimes C^{\otimes i} & \xrightarrow{\theta} & ((N \otimes C^{\otimes j} \otimes P) \otimes C^{\otimes i} \otimes M) \otimes D^{\otimes k} & \xrightarrow{\langle\langle a \rangle\rangle} & (N \otimes C^{\otimes j} \otimes (P \otimes C^{\otimes i} \otimes M)) \otimes D^{\otimes k}
\end{array}$$

We then need to check if the following diagram commutes:

$$\begin{array}{ccccc}
\mathrm{coHH}(\Omega(P, C, C), C) & \xrightarrow{\theta} & \mathrm{coHH}(\Omega(C, C, P), C) & \xrightarrow{\theta} & \mathrm{coHH}(\Omega(P, C, C), C) \\
& \searrow \langle\langle r \rangle\rangle & \downarrow \langle\langle \ell \rangle\rangle & \swarrow \langle\langle r \rangle\rangle & \\
& & \mathrm{coHH}(P, C) & &
\end{array}$$

We will prove the left triangle is commutative, but the argument for the right triangle will follow analogously. We apply the definition of θ , ℓ , and r on the bicosimplicial objects:

$$\begin{array}{ccc}
(P \otimes C^{\otimes \bullet} \otimes C) \otimes C^{\otimes *} & \xrightarrow{\theta} & (C \otimes C^{\otimes *} \otimes P) \otimes C^{\otimes \bullet} \\
\downarrow \langle\langle r \rangle\rangle & & \downarrow \langle\langle \ell \rangle\rangle \\
P \otimes C^{\otimes *} & \xrightarrow{\cong} & P \otimes C^{\otimes \bullet}.
\end{array}$$

Here we kept track of the different gradings by \bullet and $*$. Therefore, once we apply the homotopy limit, we obtain the desired result. \square

5. DUALIZABLE BICOMODULES

As traces are always defined on dualizable objects, we investigate the notion of duality in our bicategories of bicomodules.

Definition 5.1. Let \mathcal{B} be a bicategory. A 1-cell $M \in \mathcal{B}(C, D)$ is *right dualizable* if there is another 1-cell M^* in $\mathcal{B}(D, C)$, with 2-cells $\eta: U_C \rightarrow M \odot M^*$ and $\varepsilon: M^* \odot M \rightarrow U_D$, called *coevaluation* and *evaluation* respectively, such that the compositions in $\mathcal{B}(C, D)$ and $\mathcal{B}(D, C)$ respectively are the identity 2-cells:

$$\begin{aligned}
M &\cong U_C \odot M \xrightarrow{\eta \odot \mathrm{id}} (M \odot M^*) \odot M \cong M \odot (M^* \odot M) \xrightarrow{\mathrm{id} \odot \varepsilon} M \odot U_D \cong M, \\
M^* &\cong M^* \odot U_C \xrightarrow{\mathrm{id} \odot \eta} M^* \odot (M \odot M^*) \cong (M^* \odot M) \odot M^* \xrightarrow{\varepsilon \odot \mathrm{id}} U_D \odot M^* \cong M^*.
\end{aligned}$$

We call M^* the *right dual* of M and say that (M, M^*) form a *dual pair* in \mathcal{B} .

For instance, in the bicategory of bimodules, an (R, S) -bimodule M is right dualizable if and only if it is finitely generated and projective as a right S -module, and its dual is given by its linear dual $\mathrm{Hom}_S(M, S)$, see [PS13, 6.1]. We shall obtain a very similar result for bicomodules with the subtlety that our bicategory is not “closed” (see [Pon10, 4.1.4]), and thus we cannot recognize dualizable objects (as in [Pon10, 4.3.3]) since we are not provided with an internal hom. Our bicategories will almost be “co-closed” thanks to the introduction of a cohom functor. We shall make use of the notion of “quasi-finite” comodules, as introduced by Takeuchi in [Tak77]. More modern reviews, in more general settings, can be found in [BW03] and [Al-02].

Let ${}_C\text{CoMod}$ be the category of left C -comodules. If M is a left C -comodule and V is a \mathbb{k} -module, then $M \otimes V$ is a left C -comodule. This defines a functor:

$$\begin{aligned} M \otimes -: \text{Mod}_{\mathbb{k}} &\longrightarrow {}_C\text{CoMod} \\ V &\longmapsto M \otimes V. \end{aligned}$$

Definition 5.2. We say a left C -comodule M is *quasi-finite* if the functor $M \otimes -: \text{Mod}_{\mathbb{k}} \rightarrow {}_C\text{CoMod}$ is a right adjoint. In this case, we denote its left adjoint by $h_C(M, -): {}_C\text{CoMod} \rightarrow \text{Mod}_{\mathbb{k}}$, and refer to it as the *cohom* functor. In other words, given any left C -comodule N , the cohom $h_C(M, N)$ is the universal \mathbb{k} -module providing a natural \mathbb{k} -linear isomorphism

$$\text{Mod}_{\mathbb{k}}(h_C(M, N), V) \cong {}_C\text{CoMod}(N, M \otimes V), \quad (5.3)$$

for any \mathbb{k} -module V . In particular, for $V = \mathbb{k}$, we obtain the isomorphism

$$h_C(M, N)^* \cong {}_C\text{Hom}(N, M).$$

Example 5.4. An example of a quasi-finite left C -comodule M is given by cofree comodules $M = C \otimes V$, where V is a finitely generated \mathbb{k} -module. Since we have

$$\begin{aligned} \text{Mod}_{\mathbb{k}}(h_C(C \otimes V, N), W) &\cong {}_C\text{CoMod}(N, C \otimes V \otimes W) \\ &\cong \text{Mod}_{\mathbb{k}}(N, V \otimes W) \\ &\cong \text{Mod}_{\mathbb{k}}(V^* \otimes N, W), \end{aligned}$$

for any \mathbb{k} -module W , we obtain $h_C(C \otimes V, N) \cong V^* \otimes N$. Choosing $V = \mathbb{k}^{\oplus n}$ so that $M = C^{\oplus n}$, then $h_C(C^{\oplus n}, N) \cong N^{\oplus n}$. More generally, any finitely cogenerated comodule (Definition 2.8) is quasi-finite.

The unit of the adjunction (5.3) provides the \mathbb{k} -linear coevaluation

$$\eta: N \longrightarrow M \otimes h_C(M, N).$$

If D is another coalgebra, and N is a (C, D) -bicomodule, then $h_C(M, N)$ is a right D -comodule, and the map η is a morphism of (C, D) -bicomodules, see [Tak77, 1.7, 1.19]. Therefore, for any quasi-finite left C -comodule M , we obtain that $h_C(M, C)$ is a right C -comodule, and we have a (C, C) -bicomodule homomorphism $\eta: C \rightarrow M \otimes h_C(M, C)$.

Let T be a left C -comodule, M a quasi-finite left C -comodule, and N a (C, C) -bicomodule. Denote by $\partial: h_C(M, N \square_C T) \rightarrow h_C(M, N) \square_C T$ the adjoint to the left C -colinear homomorphism

$$\eta \square 1: N \square_C T \longrightarrow (M \otimes h_C(M, N)) \square_C T \cong M \otimes (h_C(M, N) \square_C T).$$

It is noted that, in [Tak77, 1.14], the \mathbb{k} -linear homomorphism ∂ above is an isomorphism whenever M is an injective quasi-finite left C -comodule. If we choose $N = C$ and $T = M$, combining the isomorphism with the map $h_C(M, M) \rightarrow \mathbb{k}$ of Definition 2.10, we obtain the evaluation

$$\varepsilon: h_C(M, C) \square_C M \xleftarrow[\cong]{\partial} h_C(M, M) \longrightarrow \mathbb{k}.$$

Example 5.5. Consider the bicategory CoMod of Definition 3.5. A comodule M in ${}_C\text{CoMod}_{\mathbb{k}}$ is right dualizable if it is quasi-finite and injective as a left C -comodule. The dual of M is the right C -comodule $h_C(M, C)$, together with the coevaluation $\eta: C \rightarrow M \otimes h_C(M, C)$ and evaluation $\varepsilon: h_C(M, C) \square_C M \rightarrow \mathbb{k}$ defined above. The desired triangle identities follow from the adjunction of the cohom functor (5.3).

Example 5.6. In Example 5.5, choose $M = C \otimes V$, where V is a dualizable \mathbb{k} -module (i.e. finitely generated). We obtain $M^* = h_C(C \otimes V, C) \cong V^* \otimes C$. We can then explicitly describe the coevaluation and evaluation. Let us denote the comultiplication and counit of C by $\Delta_C: C \rightarrow C \otimes C$ and $\varepsilon_C: C \rightarrow \mathbb{k}$ respectively. Similarly, denote the coevaluation and evaluation of the dualizable \mathbb{k} -module V by $\eta_V: \mathbb{k} \rightarrow V \otimes V^*$ and $\varepsilon_V: V^* \otimes V \rightarrow \mathbb{k}$ respectively. Then the coevaluation $\eta: C \rightarrow M \otimes M^* \cong (C \otimes V) \otimes (V^* \otimes C)$ is the composite

$$C \xrightarrow{\Delta_C} C \otimes C \cong C \otimes \mathbb{k} \otimes C \xrightarrow{\text{id} \otimes \eta_V \otimes \text{id}} C \otimes (V \otimes V^*) \otimes C.$$

Explicitly, if we pick (e_1, \dots, e_n) for a basis of V , and (e_1^*, \dots, e_n^*) for the dual basis of V^* , and write $\Delta_C(c) = \sum_{(c)} c_{(1)} \otimes c_{(2)}$, we obtain the formula

$$\eta(c) = \sum_{i=1}^n \sum_{(c)} c_{(1)} \otimes e_i \otimes e_i^* \otimes c_{(2)}.$$

The evaluation $\varepsilon: M^* \square_C M \cong V^* \otimes C \otimes V \rightarrow \mathbb{k}$ is then the composite

$$V^* \otimes C \otimes V \xrightarrow{\text{id} \otimes \varepsilon_C \otimes \text{id}} V^* \otimes \mathbb{k} \otimes V \cong V^* \otimes V \xrightarrow{\varepsilon_V} \mathbb{k}.$$

Explicitly, we obtain the formula for $f \in V^*, c \in C, v \in V$:

$$\varepsilon(f \otimes c \otimes v) = \varepsilon_C(c) f(v).$$

Choosing in particular $V = \mathbb{k}^{\oplus n}$ so that $M = C^{\oplus n}$ and $M^* \cong C^{\oplus n}$ and $V \otimes V^* \cong \mathcal{M}_n(\mathbb{k})$, we get $C^{\oplus n} \otimes C^{\oplus n} \cong \mathcal{M}_n^c(C) \otimes C$ and the coevaluation can be formulated as the (C, C) -bilinear morphism:

$$\begin{aligned} \eta: C &\longrightarrow \mathcal{M}_n^c(C) \otimes C \\ c &\longmapsto \sum_{(c)} c_{(1)} \otimes I_n \otimes c_{(2)}. \end{aligned}$$

Notice that $M^* \square_C M \cong \mathcal{M}_n^c(C)$ as \mathbb{k} -modules, and thus the evaluation can be formulated as a \mathbb{k} -linear homomorphism:

$$\begin{aligned} \varepsilon: \mathcal{M}_n^c(C) &\longrightarrow \mathbb{k} \\ c \otimes f &\longmapsto \varepsilon_C(c) \text{tr}(f). \end{aligned}$$

Example 5.7. We now generalize Example 5.5. A comodule M in ${}_C\text{CoMod}_D$ is right dualizable whenever it is quasi-finite and injective as a left C -comodule. Define $M^* = h_C(M, C)$, which is a (D, C) -bicomodule. Indeed, by [Tak77, 1.8], for any (C, D) -bicomodule M such that M is quasi-finite as a left C -comodule, and any left C -comodule N , we have that $h_C(M, N)$ is a left D -comodule with coaction $h_C(M, N) \rightarrow D \otimes h_C(M, N)$ given by the adjoint to the left C -colinear map

$$N \xrightarrow{\eta} M \otimes h_C(M, N) \xrightarrow{\rho \otimes \text{id}} M \otimes D \otimes h_C(M, N).$$

Moreover, by [Tak77, 1.7], the C -bilinear map $\eta: C \rightarrow M \otimes h_C(M, C)$ factors through the (C, C) -bicomodule $M \square_D h_C(M, C)$, and the induced map $\eta: C \rightarrow M \square_D h_C(M, C)$ remains C -bilinear by [Tak77, 1.9]. Therefore this defines the desired C -bilinear coevaluation $\eta: C \rightarrow M \square_D M^*$. In fact, by [Tak77, 1.10], the cohom functor $h_C(M, -): {}_C\text{CoMod} \rightarrow \text{Mod}_{\mathbb{k}}$ can be promoted to a functor $h_C(M, -): {}_C\text{CoMod} \rightarrow {}_D\text{CoMod}$ whenever the quasi-finite left C -comodule M is endowed with a right D -coaction, and it is the left adjoint of the functor

$$\begin{aligned} {}_D\text{CoMod} &\longrightarrow {}_C\text{CoMod} \\ L &\longrightarrow M \square_D L, \end{aligned}$$

i.e., we obtain an equivalence

$${}_D\text{CoMod}(h_C(M, N), L) \cong {}_C\text{CoMod}(N, M \square_D L) \quad (5.8)$$

for any left C -comodule N and left D -comodule L . In particular, if we choose $M = N$ and $L = D$, the adjoint of the identity map on M provides a left D -colinear map $h_C(M, M) \rightarrow D$, which is in fact D -bilinear by [Tak77, 1.11]. Moreover, by [Tak77, 1.15], the map

$$h_C(M, M) \cong h_C(M, C \square_C M) \xrightarrow{\partial} h_C(M, C) \square_C M,$$

is D -bilinear. When M is injective as a left C -comodule, the map ∂ is an isomorphism. Therefore, the desired evaluation $\varepsilon: M^* \square_C M \rightarrow D$ is given by composing the previous D -bilinear maps

$$h_C(M, C) \square_C M \xrightarrow[\cong]{\partial^{-1}} h_C(M, C \square_C M) \cong h_C(M, M) \longrightarrow D.$$

Just as in Example 5.5, the triangle identities follow from the adjunction of the cohom functor (5.8).

Example 5.9. A particular case of Example 5.7 is when $C = \mathbb{k}$. A bicomodule M in ${}_{\mathbb{k}}\text{CoMod}_D$ is right dualizable when M is dualizable as a \mathbb{k} -module (i.e., finitely generated), and its right dual is $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$, the usual linear dual. This appeared in the proof of Proposition 2.22. Explicitly, its left D -coaction is given by:

$$\begin{aligned} \lambda: M^* &\longrightarrow \text{Hom}_{\mathbb{k}}(M, D) \cong D \otimes M^* \\ \left(M \xrightarrow{\alpha} \mathbb{k}\right) &\longmapsto \left(M \xrightarrow{\rho} M \otimes D \xrightarrow{\alpha \otimes \text{id}} D\right), \end{aligned}$$

where $\rho: M \rightarrow M \otimes D$ is the right D -coaction of M . By [BW03, 10.11], we have an isomorphism of \mathbb{k} -modules $M \square_D M^* \cong \text{Hom}_D(M, M)$. Therefore, we can interpret the coevaluation $\eta: \mathbb{k} \rightarrow M \square_D M^*$ as:

$$\begin{aligned} \mathbb{k} &\longrightarrow \text{Hom}_D(M, M) \\ 1 &\longmapsto \text{id}_M. \end{aligned}$$

Notice that the following diagram commutes:

$$\begin{array}{ccc} M^* \otimes M & \xrightarrow{\lambda \otimes \text{id}} & \text{Hom}_{\mathbb{k}}(M, D) \otimes M \\ \text{id} \otimes \rho \downarrow & & \downarrow \\ M^* \otimes M \otimes D & \longrightarrow & D, \end{array}$$

where the unlabeled maps are defined by evaluating. This defines the D -bilinear evaluation homomorphism $\varepsilon: M^* \otimes M \rightarrow D$:

$$\varepsilon(\alpha \otimes m) = \sum_{(m)} \alpha(m_{(0)}) \otimes m_{(1)}$$

where we denoted $\rho(m) = \sum_{(m)} m_{(0)} \otimes m_{(1)}$ for all $m \in M$.

We can further generalize the examples above to the differential graded case. Arguments for quasi-finite comodules are entirely categorical (see [Al-02]). However, the notion has not been documented before in this context, and therefore we introduce them here.

Definition 5.10. Let C be a simply connected dg-coalgebra over \mathbb{k} . A left dg- C -comodule M is *quasi-finite* if the functor

$$\begin{aligned} \text{Ch}_{\mathbb{k}}^{\geq 0} &\longrightarrow {}_C\text{CoMod} \\ V &\longmapsto M \otimes V \end{aligned}$$

admits a left adjoint, denoted $h_C(M, -): {}_C\text{CoMod} \rightarrow \text{Ch}_{\mathbb{k}}^{\geq 0}$, called the *cohom functor*. In other words, we obtain an equivalence

$$\text{Ch}_{\mathbb{k}}^{\geq 0}(h_C(M, N), V) \cong {}_C\text{CoMod}(N, M \otimes V),$$

for any left dg- C -comodule N and \mathbb{k} -chain complex V .

Example 5.11. If V is a perfect \mathbb{k} -chain complex, then just as in Example 5.4, we have that $C \otimes V$ is quasi-finite, and $h_C(C \otimes V, N) \cong V^* \otimes N$. If $C = \mathbb{k}$, then a left dg C -comodule is just a chain complex, and quasi-finite comodules are precisely the perfect chain complexes.

If M is also endowed with a right D -coaction, then $h_C(M, N)$ is a right D -comodule and we obtain the adjunction

$${}_C\text{CoMod} \begin{array}{c} \xrightarrow{h_C(M, -)} \\ \perp \\ \xleftarrow{M \square_D -} \end{array} {}_D\text{CoMod}.$$

Therefore, just as in the discrete case, the right dual of a (C, D) -bicomodule M should be the (D, C) -bicomodule $h_C(M, C)$. Notice, even if M is a fibrant (C, D) -bicomodule, there is no reason to expect

$h_C(M, C)$ is fibrant as a (D, C) -bicomodule. However, this is not needed. Indeed, by Proposition A.17, if M is a fibrant (C, D) -bicomodule, then

$$\begin{aligned} M \square_D h_C(M, C) &\simeq M \square_D \Omega(D, D, h_C(M, C)) \\ &\simeq \Omega(M, D, h_C(M, C)) \\ &\simeq M \hat{\square}_D h_C(M, C). \end{aligned}$$

Similarly, we obtain $h_C(M, C) \square_C M \simeq h_C(M, C) \hat{\square}_C M$.

Just as in the discrete case, the C -bilinear map $\eta: C \rightarrow M \square_D h_C(M, C)$ from the unit of the cohom adjunction induces the desired coevaluation on the dual pair $(M, h_C(M, C))$.

We now describe how to obtain the evaluation. Recall that a \mathbb{k} -chain complex V is perfect if and only if the natural map $V \otimes \text{Hom}_{\mathbb{k}}(V, \mathbb{k}) \rightarrow \text{Hom}_{\mathbb{k}}(V, V)$ is an isomorphism. Similarly, as in Example 5.5, for any quasi-finite left C -comodule M , we have a natural D -bilinear map $\partial: h_C(M, M) \rightarrow h_C(M, C) \square_C M$. We saw in the discrete case that if M is injective and quasi-finite, then ∂ is an isomorphism.

Definition 5.12. A fibrant left C -comodule M is said to be *coperfect* if it is quasi-finite and the induced map $\partial: h_C(M, M) \rightarrow h_C(M, C) \square_C M$ is an isomorphism.

Given a coperfect (C, D) -bicomodule M , define the evaluation as

$$\varepsilon: h_C(M, C) \square_C M \xrightarrow[\cong]{\partial^{-1}} h_C(M, M) \longrightarrow D.$$

Combining our arguments above, we obtain the desired triangle identities by adjunction of the cohom functor, and thus the following result.

Proposition 5.13. *Let C and D be simply connected dg-coalgebras. A (fibrant) (C, D) -bicomodule M is right dualizable if it is coperfect as a left C -comodule. Its right dual is given by $h_C(M, C)$.*

Example 5.14. Suppose $D = \mathbb{k}$, and $M = C \otimes V$, where V is a perfect chain complex. Then M is quasi-finite as a left C -comodule and $M^* = h_C(M, C) \cong V^* \otimes C$. Moreover, M is coperfect because

$$V^* \otimes M \cong h_C(C \otimes V, M) \rightarrow h_C(C \otimes V, C) \square_C M \cong V^* \otimes M$$

is an isomorphism.

Example 5.15. If $C = \mathbb{k}$, then if M is a right D -comodule such that M is a perfect chain complex, then M is dualizable and $M^* = \text{Hom}_{\mathbb{k}}(M, \mathbb{k})$.

6. BICATEGORICAL TRACES

Every shadowed bicategory defines a notion of traces on its dualizable objects. This extends the notion of traces on symmetric monoidal categories, as reviewed in [PS14]. We show the Hattori–Stallings cotrace ${}_C \text{End}(M) \rightarrow \text{coHH}_0(C)^*$ (Definition 2.13) and the colinear Hattori–Stallings trace $\text{End}_C(M) \rightarrow \text{coHH}_0(C)$ (Definition 2.23) are bicategorical traces on the shadowed bicategory $(\text{CoMod}, \text{coHH}_0)$. We first recall the general definition of a bicategorical trace.

Definition 6.1. [PS13] Let $(\mathcal{B}, \langle \langle - \rangle \rangle)$ be a shadowed bicategory. Let $M \in \mathcal{B}(C, D)$ be a right dualizable 1-cell. The *trace* of a 2-cell $f: M \rightarrow M$, denoted $\text{tr}^{\mathcal{B}}(f)$, is the composite

$$\langle \langle U_C \rangle \rangle_C \xrightarrow{\langle \langle \eta \rangle \rangle} \langle \langle M \odot M^* \rangle \rangle_C \xrightarrow{\langle \langle f \odot \text{id} \rangle \rangle} \langle \langle M \odot M^* \rangle \rangle_C \xrightarrow[\cong]{\theta} \langle \langle M^* \odot M \rangle \rangle_D \xrightarrow{\langle \langle \varepsilon \rangle \rangle} \langle \langle U_D \rangle \rangle_D.$$

More generally, given 1-cells $P \in \mathcal{B}(D, D)$ and $Q \in \mathcal{B}(C, C)$, the trace of a 2-cell $f: Q \odot M \rightarrow M \odot P$ is the composite

$$\langle \langle Q \rangle \rangle_C \xrightarrow{\langle \langle \text{id} \odot \eta \rangle \rangle} \langle \langle Q \odot M \odot M^* \rangle \rangle_C \xrightarrow{\langle \langle f \odot \text{id} \rangle \rangle} \langle \langle M \odot P \odot M^* \rangle \rangle_C \xrightarrow[\cong]{\theta} \langle \langle M^* \odot M \rangle \rangle_D \xrightarrow{\langle \langle \varepsilon \odot \text{id} \rangle \rangle} \langle \langle P \rangle \rangle_D.$$

The *Euler characteristic* of M , denoted $\chi(M)$, is the trace of its identity 2-cell $\text{tr}^{\mathcal{B}}(\text{id}_M)$.

Bicategorical traces enjoy many useful properties that are recorded in [PS13, Section 7]. Notably, we obtain the following cyclicity property.

Proposition 6.2 ([PS13, 7.3]). *If M and N are right dualizable 1-cells in a shadowed bicategory \mathcal{B} and if $f: M \rightarrow N$ and $g: N \rightarrow M$ are 2-cells in \mathcal{B} , then $\mathrm{tr}^{\mathcal{B}}(f \circ g) = \mathrm{tr}^{\mathcal{B}}(g \circ f)$.*

Example 6.3 ([Pon10, 4.2.2]). Let R be a ring and let M be a finitely generated projective right R -module. Let f be an R -linear endomorphism on M . The Hattori–Stallings trace (Definition 2.2) can be regarded as the image of 1 in the bicategorical trace $\mathrm{tr}(f): \mathbb{Z} \rightarrow \mathrm{HH}_0(R)$:

$$\mathbb{Z} \xrightarrow{\eta} M \otimes_R \mathrm{Hom}_R(M, R) \xrightarrow{f \otimes 1} M \otimes_R \mathrm{Hom}_R(M, R) \cong \mathrm{HH}_0(R, \mathrm{Hom}_R(M, R) \otimes_{\mathbb{Z}} M) \xrightarrow{\langle\langle \varepsilon \rangle\rangle} \mathrm{HH}_0(R).$$

Given \mathbb{k} -coalgebras C and D , and M a right dualizable (C, D) -bicomodule with dual M^* , for any (C, D) -bilinear homomorphism $f: M \rightarrow M$, we obtain a trace $\mathrm{tr}_C^D(f): \mathrm{coHH}_0(C) \rightarrow \mathrm{coHH}_0(D)$ defined as the composition:

$$\mathrm{coHH}_0(C) \xrightarrow{\langle\langle \eta \rangle\rangle} \mathrm{coHH}_0(M \square_D M^*, C) \xrightarrow{\langle\langle f \square 1 \rangle\rangle} \mathrm{coHH}_0(M \square_D M^*, C) \cong \mathrm{coHH}_0(M^* \square_C M, D) \xrightarrow{\langle\langle \varepsilon \rangle\rangle} \mathrm{coHH}_0(D).$$

We show how the bicategorical trace recovers the Hattori–Stallings cotrace of Definition 2.13. Recall from Example 5.5 that a finitely cogenerated and injective left C -comodule is a right dualizable (C, \mathbb{k}) -bicomodule.

Proposition 6.4. *Let C be a \mathbb{k} -coalgebra. Let M be a finitely cogenerated and injective left C -comodule. The Hattori–Stallings cotrace ${}_C\mathrm{End}(M) \rightarrow \mathrm{coHH}_0(C)^*$ sends a C -colinear homomorphism $f: M \rightarrow M$ to the bicategorical trace $\mathrm{tr}_C^{\mathbb{k}}(f)$:*

$$\mathrm{coHH}_0(C) \xrightarrow{\langle\langle \eta \rangle\rangle} \mathrm{coHH}_0(M \otimes h_C(M, C), C) \xrightarrow{\langle\langle f \otimes 1 \rangle\rangle} \mathrm{coHH}_0(M \otimes h_C(M, C), C) \cong h_C(M, C) \square_C M \xrightarrow{\varepsilon} \mathbb{k}.$$

Proof. Suppose first $M = C^{\oplus n}$. Then the bicategorical trace $\mathrm{tr}_C^{\mathbb{k}}(f)$ can be re-written as the composition:

$$\mathrm{coHH}_0(C) \xrightarrow{\langle\langle \eta \rangle\rangle} \mathrm{coHH}_0(\mathcal{M}_n^c(C) \otimes C, C) \xrightarrow{\langle\langle f \otimes 1 \rangle\rangle} \mathrm{coHH}_0(\mathcal{M}_n^c(C) \otimes C, C) \cong \mathcal{M}_n^c(C) \xrightarrow{\varepsilon} \mathbb{k}.$$

Recall given $f: M \rightarrow M$, we denoted the corresponding \mathbb{k} -linear homomorphism $C \rightarrow \mathcal{M}_n(\mathbb{k})$ by $c \mapsto f_c$.

Given $c \in \mathrm{coHH}_0(C) \subseteq C$, following the formulas of Example 5.6, we obtain that the image of c under the composition is given by:

$$c \mapsto \sum_{(c)} c_{(1)} \otimes I_n \otimes c_{(2)} \mapsto \sum_{(c)} c_{(1)} \otimes f_{c_{(2)}} \otimes c_{(3)} \mapsto \sum_{(c)} c_{(1)} \otimes f_{c_{(2)}} \mapsto \sum_{(c)} \varepsilon(c_{(1)}) \mathrm{tr}(f_{c_{(2)}}).$$

Therefore $\mathrm{tr}_C^{\mathbb{k}}(f)(c) = \mathrm{cotr}_M(f)(c)$ for $M = C^{\oplus n}$. We can conclude for a general finitely cogenerated and injective comodule M by applying Lemma 2.11. \square

Similarly, we can show the bicategorical trace recovers the colinear Hattori–Stallings trace of Definition 2.23. From Example 5.9, we know that right C -comodules that are finitely generated as \mathbb{k} -modules are right dualizable (\mathbb{k}, C) -bicomodules.

Proposition 6.5. *Let C be a \mathbb{k} -coalgebra. Let M be a right C -comodule that is finitely generated as a \mathbb{k} -module. The colinear Hattori–Stallings trace $\mathrm{End}_C(M) \rightarrow \mathrm{coHH}_0(C)$ sends a C -colinear homomorphism $f: M \rightarrow M$ to the image of 1 in \mathbb{k} in the bicategorical trace $\mathrm{tr}_C^C(f)$:*

$$\mathbb{k} \xrightarrow{\eta} M \square_C M^* \xrightarrow{f \square 1} M \square_C M^* \cong \mathrm{coHH}_0(M^* \otimes M, C) \xrightarrow{\langle\langle \varepsilon \rangle\rangle} \mathrm{coHH}_0(C).$$

Proof. We denote the right coaction $M \rightarrow M \otimes C$ by $m \mapsto \sum_{(m)} m_{(0)} \otimes m_{(1)}$. Choosing a basis (e_1, \dots, e_n) of M , and applying the formulas from Example 5.9, we obtain that the image of 1 in the composition is given by:

$$1 \mapsto \sum_{i=1}^n e_i \otimes e_i^* \mapsto \sum_{i=1}^n f(e_i) \otimes e_i^* \mapsto \sum_{i=1}^n e_i^* \otimes f(e_i) \mapsto \sum_{i=1}^n \sum_{(f(e_i))} e_i^*(f(e_i)_{(0)}) \otimes f(e_i)_{(1)}.$$

As f is C -colinear, we obtain in particular:

$$\sum_{(f(e_i))} f(e_i)_{(0)} \otimes f(e_i)_{(1)} = \sum_{(e_i)} f(e_i)_{(0)} \otimes e_{i(1)}.$$

Therefore, we can conclude $\mathrm{tr}_C^C(f)(1) = \mathrm{tr}^C(f)$ as in Definition 2.23. \square

Remark 6.6. In recent independent work of Justin Barhite [Bar23], Hochschild cohomology HH^0 is shown to be a so-called *coshadow* on the bicategory of bimodules over \mathbb{k} -algebras, and there are associated *bicategorical cotraces*. If C is a \mathbb{k} -coalgebra that is finitely generated as \mathbb{k} -module, then the bicategory of bimodules over C^* is intimately related to the bicategory of bicomodules over C since $M \square_C N \cong {}_{C^*}\mathrm{Hom}_{C^*}(C^*, M \otimes N)$ by [BW03, 10.10]. This induces an equivalence $\mathrm{coHH}_0(N \otimes M, C) \cong \mathrm{HH}^0(C^*, M \otimes N)$. From our perspective however, our cotrace comes from a shadow and not a coshadow, and we do not assume that our bicategory is closed. One can build a functor $(\mathrm{CoMod}^{\mathrm{fg}})^{\mathrm{co-op}} \rightarrow \mathrm{Mod}$ from the co-opposite bicategory (inverting both 1 and 2 cells) of finitely generated bicomodules to the bicategory of bimodules, and we expect that it sends the shadow coHH_0 to the coshadow HH^0 . Nonetheless, it does not seem that the language of coshadows and bicategorical cotraces is the appropriate vocabulary for the Hattori–Stallings cotrace, nor the colinear Hattori–Stallings trace.

Example 6.7. We can generalize the Hattori–Stallings cotrace for chain complexes. Let C be a simply connected dg-coalgebra over \mathbb{k} , let M be a coperfect left C -comodule, and let f be an C -colinear endomorphism on M . Then we obtain the cotrace $\mathrm{coHH}(C) \rightarrow \mathbb{k}$. Since the homomorphism is in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$, and \mathbb{k} is concentrated in degree zero, then the above trace is entirely determined by $\mathrm{coHH}_0(C)$. Unfortunately, as C is simply connected, $\mathrm{coHH}_0(C) = \mathbb{k}$, and this cotrace is simply the alternating sum of the usual trace of f on each degree. Similarly, we can also generalize the colinear Hattori–Stallings trace for chain complexes. Letting V denote a perfect chain complex with a right C -comodule structure, given a C -colinear endomorphism $f: V \rightarrow V$, its colinear trace defines a chain homomorphism $\mathbb{k} \rightarrow \mathrm{coHH}(C)$. It is non-trivial only in degree zero, but as C is simply connected, we get that the colinear trace is the alternating sum of the trace of f on each degree.

Example 6.8. Let X be a simply connected CW-complex, and let Y be a finite CW-complex. Let $C_*(-; \mathbb{k}) = C_*(-)$ denote the singular chain complex over \mathbb{k} associated to a space. Recall that $C_*(X)$ is a dg-coalgebra over \mathbb{k} using Alexander–Whitney formula and the diagonal $X \rightarrow X \times X$. Then $C_*(X \times Y) \cong C_*(X) \otimes C_*(Y)$ is a coperfect $C_*(X)$ -comodule. Let $f: Y \rightarrow Y$ be an endomorphism. This defines an endomorphism on $X \times Y$ that is the identity on X , and thus an endomorphism \hat{f} on the comodule $C_*(X \times Y)$. Therefore the cotrace of \hat{f} is a map $\mathrm{coHH}(C_*(X)) \rightarrow \mathbb{k}$ in which its image is the usual Lefschetz number of f .

Example 6.9. Let X be a simply connected CW-complex, and let Y be a finite CW-complex with a continuous map $\rho: Y \rightarrow X$. Then $C_*(Y)$ is a comodule over $C_*(X)$. Let f be an endomorphism over Y . Then the colinear trace induced on coHochschild homology $\mathbb{k} \rightarrow \mathrm{coHH}(C_*(X))$ is the usual alternating sum of the traces of $C_*(Y) \xrightarrow{f} C_*(Y) \xrightarrow{\rho} C_*(X)$.

7. MORITA–TAKEUCHI INVARIANCE

In bicategories, Morita equivalence is the natural notion of an equivalence in a bicategory. It extends the usual notion of equivalence of categories and Morita equivalence between rings.

Definition 7.1 ([CP19, 4.1, 4.2]). Let \mathcal{B} be a bicategory, let $M \in \mathcal{B}(C, D)$ be a right dualizable 1-cell, and denote its right dual by M^* . We say the dual pair (M, M^*) is a *Morita equivalence* in \mathcal{B} if the coevaluation $\eta: U_C \rightarrow M \odot M^*$ and evaluation $\varepsilon: M^* \odot M \rightarrow D$ maps are isomorphisms in $\mathcal{B}(C, C)$ and $\mathcal{B}(D, D)$ respectively. If such a pair exists, we say C and D are *Morita equivalent*.

Proposition 7.2 ([CP19, 4.5]). Let $(\mathcal{B}, \langle\langle - \rangle\rangle)$ be a shadowed bicategory. Let $M \in \mathcal{B}(C, D)$ be a right dualizable 1-cell and denote its right dual by M^* . If (M, M^*) is a Morita equivalence, then the Euler characteristic $\chi(M)$ is an isomorphism, with inverse $\chi(M^*)$. In particular, if C and D are Morita equivalent, then

$$\langle\langle U_C \rangle\rangle_C \cong \langle\langle U_D \rangle\rangle_D.$$

Example 7.3. Consider the bicategory CoMod of bicomodules over \mathbb{k} as in Definition 3.5. Then in [Tak77, 2.3], a Morita equivalence is referred to as a *set of equivalence data*. It is then shown that a Morita equivalence in this bicategory recovers the notion of Morita–Takeuchi equivalence in \mathbb{k} -modules (Definition 1.17). Therefore, if C and D are Morita–Takeuchi equivalent, then $\mathrm{coHH}_0(C) \cong \mathrm{coHH}_0(D)$ by Proposition 7.2. This recovers the results of [FS98] at level zero. By [Tak77, 3.5], if M is a left C -comodule that is a

quasi-finite injective cogenerator, and $D := e_C(M)$, the coalgebra of coendomorphisms, then C and D are Morita–Takeuchi equivalent, and M and M^* form a Morita equivalence in the bicategory \mathcal{CoMod} .

Consider the bicategory of derived bicomodules \mathcal{DCoMod} as in Definition 3.13. We say two simply connected coalgebras C and D in $\mathcal{Ch}_{\mathbb{k}}^{\geq 0}$ are *homotopically Morita–Takeuchi equivalent* if they are Morita equivalent in the bicategory \mathcal{DCoMod} . A major consequence of Theorem 4.9 together with Proposition 7.2 is the following.

Theorem 7.4. *Let C and D be homotopically Morita–Takeuchi equivalent simply connected differential graded \mathbb{k} -coalgebras. Then we obtain an isomorphism:*

$$\mathrm{coHH}(C) \cong \mathrm{coHH}(D).$$

Proof. This is a combination of Theorem 4.9 together with Proposition 7.2. \square

This extends the results of [HS21], where we suspect that the simply-connected condition was forgotten to be mentioned.

Proposition 7.5. *Suppose (M, M^*) form a dual pair in \mathcal{DCoMod} . Then we obtain a Quillen adjunction*

$$M^* \square_C - : {}_C \mathcal{CoMod}(\mathcal{Ch}_{\mathbb{k}}^{\geq 0}) \xrightarrow{\perp} {}_D \mathcal{CoMod}(\mathcal{Ch}_{\mathbb{k}}^{\geq 0}) : M \square_D -.$$

Moreover, it is a Quillen equivalence if and only if (M, M^) is a homotopical Morita–Takeuchi equivalence.*

Proof. Since M and M^* are dual to each other, the maps $\eta : C \rightarrow M \square_D M^*$ and $\varepsilon : M^* \square_C M \rightarrow D$ induce the adjunction. For instance, given a left C -comodule P , a left D -comodule Q , and a D -colinear map $\alpha : M^* \square_C P \rightarrow Q$, we obtain a C -colinear map $P \rightarrow M \square_D Q$ via the composite

$$P \cong C \square_C P \xrightarrow{\eta \square \mathrm{id}} (M \square_D M^*) \square_C P \cong M \square_D (M^* \square_C P) \xrightarrow{\mathrm{id} \square \alpha} M \square_D Q.$$

The axioms on η and ε guarantee that the procedure gives a correspondence:

$${}_D \mathcal{CoMod}(\mathcal{Ch}_{\mathbb{k}}^{\geq 0}) \left(M^* \square_C P, Q \right) \cong {}_C \mathcal{CoMod}(\mathcal{Ch}_{\mathbb{k}}^{\geq 0}) \left(P, M \square_D Q \right).$$

Since the cotensor product is left exact, then the functor $M^* \square_D -$ preserves monomorphisms, i.e., cofibrations. Since M^* is a fibrant right C -comodule, then $M^* \square_C -$ preserves quasi-isomorphisms, i.e., weak equivalences (by Proposition A.17). Therefore we obtain that it is a Quillen adjunction.

Suppose (M, M^*) is a homotopical Morita–Takeuchi equivalence and $P \rightarrow P'$ is a C -colinear map such that the induced map

$$M^* \square_C P \xrightarrow{\sim} M^* \square_C P'$$

is a quasi-isomorphism. Then if we apply $M \square_D -$, since M is a fibrant D -comodule, we obtain the quasi-isomorphism by Proposition A.17:

$$\underbrace{(M \square_D M^*) \square_C P}_{\simeq P} \cong M \square_D (M^* \square_C P) \xrightarrow{\sim} M \square_D (M^* \square_C P') \cong \underbrace{(M \square_D M^*) \square_C P'}_{\simeq P'}.$$

Therefore $P \rightarrow P'$ is a quasi-isomorphism, and thus $M^* \square_C -$ reflects quasi-isomorphism. Moreover, for any fibrant left D -comodule Q , we have a quasi-isomorphism

$$Q \cong D \square_D Q \simeq M^* \square_C M \square_D Q \rightarrow Q.$$

Thus by [Hov99, 1.3.16], we obtain that the adjunction is a Quillen equivalence.

Conversely, if we supposed the adjunction to be a Quillen equivalence, then if we apply the adjoints on C and D , we recover that the coevaluation and evaluation $C \rightarrow M \square_D M^*$ and $M^* \square_C M \rightarrow D$ are quasi-isomorphisms by again applying [Hov99, 1.3.16]. \square

Example 7.6. Given a map of simply connected coalgebras $f : C \rightarrow D$, we recover the expected result that if f is a quasi-isomorphism, then C and D are homotopically Morita–Takeuchi equivalent. In more details, recall that (C, C) form a dual pair of bicomodules over C , and $h_C(C, C) \cong C$. Moreover, C can be regarded as a left D -comodule via f :

$$C \longrightarrow C \otimes C \xrightarrow{f \otimes \mathrm{id}} D \otimes C.$$

In fact, any left C -comodule P can be regarded as a left D -comodule this way, we shall denote it by $f^*(C)$. Notice that $f^*(C) \cong C \square_C P$ as a left D -comodule. We obtain a Quillen adjunction:

$$f^*: {}_C\mathrm{CoMod}(\mathrm{Ch}_{\mathbb{k}}^{\geq 0}) \xrightleftharpoons[\perp]{} {}_D\mathrm{CoMod}(\mathrm{Ch}_{\mathbb{k}}^{\geq 0}): C \square_D -,$$

which is a Quillen equivalence if and only if f is a quasi-isomorphism. This recovers the usual change of coalgebras, see [Pér24b, 4.10].

APPENDIX A. HOMOTOPY THEORY OF CONNECTIVE BICOMODULES

The goal of the appendix is to show that the homotopy theory of bicomodules is endowed with a derived cotensor product which provides a homotopy coherent monoidal structure.

Let \mathcal{C}^{\otimes} be a symmetric monoidal ∞ -category, as in [Lur17, 2.0.0.7]. Subsequently, we only refer to its underlying ∞ -category, \mathcal{C} , as in [Lur17, 2.1.2.20]. By an \mathbb{E}_1 -algebra in \mathcal{C} , we mean an associative algebra in the sense of [Lur17, 4.1.1.6]. Given \mathbb{E}_1 -algebras A and B in \mathcal{C} , denote the ∞ -category of (A, B) -bimodule objects in \mathcal{C} by ${}_A\mathrm{Mod}_B(\mathcal{C})$, as in [Lur17, Section 4.3]. From [Pér22a, 2.1], define an \mathbb{E}_1 -coalgebra in \mathcal{C} to be an \mathbb{E}_1 -algebra in the opposite category $\mathcal{C}^{\mathrm{op}}$. Similarly as in [Lur17, 4.1.1.7], given an \mathbb{E}_1 -coalgebra C in \mathcal{C} , we can define C^{op} , the *opposite* \mathbb{E}_1 -coalgebra of C .

Definition A.1. Let \mathcal{C} be a symmetric monoidal ∞ -category. Let C and D be \mathbb{E}_1 -coalgebras in \mathcal{C} . A (C, D) -bicomodule object in \mathcal{C} is a (C, D) -bimodule object in $\mathcal{C}^{\mathrm{op}}$. The ∞ -category of (C, D) -bicomodules in \mathcal{C} is defined as

$${}_C\mathrm{CoMod}_D(\mathcal{C}) := ({}_C\mathrm{Mod}_D(\mathcal{C}^{\mathrm{op}}))^{\mathrm{op}}.$$

We define the ∞ -categories of left C -comodules ${}_C\mathrm{CoMod}(\mathcal{C})$ and right D -comodules $\mathrm{CoMod}_D(\mathcal{C})$ similarly.

Remark A.2. Just as in Remark 1.15, given \mathbb{E}_1 -coalgebras C and D in a symmetric monoidal ∞ -category \mathcal{C} , we obtain an equivalence of ∞ -categories

$${}_C\mathrm{CoMod}_D(\mathcal{C}) \simeq \mathrm{CoMod}_{C^{\mathrm{op}} \otimes D}(\mathcal{C}).$$

This follows from [Lur17, 4.6.3.11] applied to the opposite category. Notice that the statement requires the monoidal product of \mathcal{C} to commute with totalizations. However, as noted above [Lur17, 4.6.3.3], this requirement is not essential and remains true for any symmetric monoidal ∞ -category \mathcal{C} .

Remark A.3. Let \mathbf{C} be a symmetric monoidal category. Let \mathbf{C}^{\otimes} be its operator category as in [Lur17, 2.0.0.1]. Then its nerve $\mathcal{N}(\mathbf{C}^{\otimes})$ is a symmetric monoidal ∞ -category whose underlying ∞ -category is $\mathcal{N}(\mathbf{C})$, see [Lur17, 2.1.2.21]. Let C be a coalgebra in \mathbf{C} . It can be regarded as an \mathbb{A}_{∞} -coalgebra in $\mathcal{N}(\mathbf{C})$, see [Pér22a, 2.3]. If D is also a coalgebra in \mathbf{C} , then we obtain an equivalence of ∞ -categories

$$\mathcal{N}({}_C\mathrm{CoMod}_D(\mathbf{C})) \simeq {}_C\mathrm{CoMod}_D(\mathcal{N}(\mathbf{C})).$$

Let \mathbf{M} be a symmetric monoidal model category as in [Hov99, 4.2.6], with a class of weak equivalence denoted \mathbf{W} . We assume every object is cofibrant. Its Dwyer-Kan localization is the ∞ -category denoted $\mathcal{N}(\mathbf{M})[\mathbf{W}^{-1}]$ following [Lur17, 1.3.4.15] whose homotopy category is the homotopy category of \mathbf{M} . It is endowed with a symmetric monoidal structure via the derived tensor product, see [Lur17, 4.1.7.6]. Suppose now that the model structure on \mathbf{M} is combinatorial. Given algebras A and B in \mathbf{M} , the category of bimodules ${}_A\mathrm{Mod}_B(\mathbf{M})$ is endowed with a model structure whose class of weak equivalences, denoted \mathbf{W}' , are the morphisms of bimodules which are weak equivalences regarded as in \mathbf{M} . By [Lur17, 4.3.3.17], we obtain an equivalence of ∞ -categories

$$\mathcal{N}({}_A\mathrm{Mod}_B(\mathbf{M}))[\mathbf{W}'^{-1}] \simeq {}_A\mathrm{Mod}_B(\mathcal{N}(\mathbf{M})[\mathbf{W}^{-1}]).$$

The same statement for bicomodules is challenging if we insist on considering a combinatorial monoidal model category (and not “co-combinatorial”). Following [Pér24b], we show in Theorem A.4 when the equivalence above does hold for bicomodules. We denote the Dwyer-Kan localization of $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$ by $\mathcal{D}^{\geq 0}(\mathbb{k})$; it is equivalent to the symmetric monoidal ∞ -category of connective $H\mathbb{k}$ -modules in spectra.

Theorem A.4. *Let C and D be simply connected dg-coalgebras over \mathbb{k} . Then the natural functor*

$$\mathcal{N}\left({}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})\right)[W^{-1}] \longrightarrow {}_C\mathrm{CoMod}_D(\mathcal{D}^{\geq 0}(\mathbb{k}))$$

is an equivalence of ∞ -categories.

Proof. The result was proved in [Pér24b, 3.1] for right comodules. We can deduce the result for bicomodules using Remarks 1.15 and A.2. In more details, if C and D are simply connected, then so is $C \otimes D$:

$$(C \otimes D)_0 = C_0 \otimes D_0 \cong \mathbb{k} \otimes \mathbb{k} \cong \mathbb{k}, \quad (C \otimes D)_1 = (C_1 \otimes D_0) \oplus (C_0 \otimes D_1) = 0.$$

Therefore we can apply [Pér24b, 3.1] to the model category of right $(C^{\mathrm{op}} \otimes D)$ -comodules. In particular, we obtain the following diagram of ∞ -categories:

$$\begin{array}{ccc} \mathcal{N}\left({}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})\right)[W^{-1}] & \xrightarrow{\simeq} & \mathcal{N}\left(\mathrm{CoMod}_{C^{\mathrm{op}} \otimes D}(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})\right)[W'^{-1}] \\ \downarrow & & \downarrow \simeq \\ {}_C\mathrm{CoMod}_D(\mathcal{D}^{\geq 0}(\mathbb{k})) & \xrightarrow{\simeq} & \mathrm{CoMod}_{C^{\mathrm{op}} \otimes D}(\mathcal{D}^{\geq 0}(\mathbb{k})). \end{array}$$

Here, we let W' denote the class of right $(C^{\mathrm{op}} \otimes D)$ -comodule morphisms that are quasi-isomorphisms in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$. By [Pér24b, 3.1], the right vertical map on the diagram above is an equivalence of ∞ -categories. By Remark 1.15, the top horizontal map is an equivalence. By Remark A.2, the bottom horizontal map is an equivalence. Thus the left vertical map is an equivalence of ∞ -categories. \square

We now describe fibrant objects in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ using Postnikov towers. We say a tower $\{M(n)\}$ in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ *stabilizes in each degree* if for all $n \geq 0$, and all $0 \leq i \leq n$, the maps in the tower induce isomorphisms of \mathbb{k} -modules

$$M(n+1)_i \cong M(n+2)_i \cong M(n+3)_i \cong \dots$$

Although in general non-finite limits in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ do not correspond to the underlying limit in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$, limits of towers that stabilizes in each degree in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ do correspond to their underlying limits, see [Pér24a, 4.15]. Furthermore, usually the functor $M \square_{C-} : {}_C\mathrm{CoMod}(\mathrm{Ch}_{\mathbb{k}}^{\geq 0}) \rightarrow \mathrm{Ch}_{\mathbb{k}}^{\geq 0}$ does not preserve non-finite limits, but it does for towers that stabilize in each degree, see [Pér24a, 4.21].

Proposition A.5 ([Pér24a, 4.17]). *Let C and D be simply connected dg-coalgebras over \mathbb{k} . Let M be a (C, D) -bicomodule in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$. There exists a tower $\{M(n)\}$ in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ that stabilizes in each degree defined as follows:*

- $M(0) = 0$;
- $M(1) = C \otimes M \otimes D$;
- if the (C, D) -bicomodule $M(n)$ is defined together with a cofibration $M \hookrightarrow M(n)$ that induces an isomorphism $H_i(M) \cong H_i(M(n))$ for $0 \leq i \leq n$, then $M(n+1)$ and the cofibration $M \hookrightarrow M(n+1)$ are defined by the pullback (both in ${}_C\mathrm{CoMod}_D(\mathrm{Ch}_{\mathbb{k}}^{\geq 0})$ and in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$):

$$\begin{array}{ccc} M & \xrightarrow{\quad 0 \quad} & C \otimes P \otimes D \\ \downarrow & \searrow \text{dashed} & \downarrow \\ M(n+1) & \longrightarrow & C \otimes P \otimes D \\ \downarrow & \lrcorner & \downarrow \\ M(n) & \longrightarrow & C \otimes Q \otimes D, \end{array}$$

where the right vertical map is the (C, D) -cofree map induced by an epimorphism $P \rightarrow Q$ in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$. Moreover $\mathrm{holim}_n M(n) \simeq \lim_n M(n) \cong M$, and the (homotopy) limits are determined in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$.

We now show that fibrant objects in the category of bicomodules behave well with respect to the cotensor product.

Consider the general case of a symmetric monoidal category, $(\mathcal{C}, \otimes, \mathbb{I})$, that is abelian, and consider flat coalgebras C and D in \mathcal{C} . Then the category of bicomodules ${}_C\text{CoMod}_D(\mathcal{C})$ remains abelian, and the forgetful functor $U: {}_C\text{CoMod}_D(\mathcal{C}) \rightarrow \mathcal{C}$ preserves and reflects exact sequences, kernels, cokernels, monomorphisms, and epimorphisms.

Lemma A.6 ([Pér24b, 4.5]). *Let C and D be dg-coalgebras over \mathbb{k} . Let $f: M \rightarrow N$ be an epimorphism in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$, and let F be its kernel. Then f is a fibration in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ if and only if F is fibrant in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$.*

Lemma A.7. *Let C and D be simply connected dg-coalgebras over \mathbb{k} . If M is a fibrant (C, D) -bicomodule, then M is also a fibrant left C -comodule and a fibrant right D -comodule.*

Proof. Since M is a fibrant (C, D) -bicomodule, it is a retract of the limit \tilde{X} of its Postnikov tower $\{M(n)\}$. Since for all chain complexes V , the bicomodule $C \otimes V \otimes D$ is fibrant both as a left C -comodule and right D -comodule, then by Lemma A.6, we can conclude the desired result. \square

Lemma A.8. *Let V be in $\text{Ch}_{\mathbb{k}}^{\geq 0}$, let C and D be simply connected dg-coalgebras over \mathbb{k} , and let N be a fibrant right D -comodule. Then $(C \otimes V) \otimes N$ is a fibrant (C, D) -bicomodule.*

Proof. We consider $\{N(m)\}$, the Postnikov tower in $\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ of N . As N is a retract of the limit $\tilde{N} = \lim_m N(m)$, then it is enough to show that $C \otimes V \otimes \tilde{N}$ is a fibrant (C, D) -bicomodule. Therefore we need to prove that $\{(C \otimes V) \otimes N(m)\}$ is a fibrant tower of (C, D) -bicomodules. For $m = 0$, we get $(C \otimes V) \otimes N(0) = 0$, which is fibrant. For $m = 1$, we get $(C \otimes V) \otimes N(1) = (C \otimes V) \otimes N \otimes D$, which is a cofree (C, D) -bicomodule and is thus a fibrant (C, D) -bicomodule. Suppose $m \geq 1$, then we obtain the pullback in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$\begin{array}{ccc} (C \otimes V) \otimes N(m+1) & \longrightarrow & (C \otimes V) \otimes P \otimes D \\ \downarrow & \lrcorner & \downarrow \\ (C \otimes V) \otimes N(m) & \longrightarrow & (C \otimes V) \otimes Q \otimes D. \end{array}$$

Since the right vertical map is a fibration in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$, then so is the left vertical map. Thus $\{(C \otimes V) \otimes Y(m)\}$ is a fibrant tower of (C, D) -bicomodules. \square

Lemma A.9. *Let C and D be simply connected dg-coalgebras over \mathbb{k} . If M is a fibrant left C -comodule and N is a fibrant right D -comodule, then $M \otimes N$ is a fibrant (C, D) -bicomodule.*

Proof. Let $\{M(n)\}$ be the Postnikov tower in ${}_C\text{CoMod}(\text{Ch}_{\mathbb{k}}^{\geq 0})$ of X . As M is fibrant, it is a retract of $\tilde{X} = \lim_n M(n)$. Therefore it is sufficient to show that $\tilde{M} \otimes N$ is a fibrant (C, D) -bicomodule. Since the tower stabilizes in each degree, we have

$$(\lim_n M(n)) \otimes N \cong \lim_n (M(n) \otimes N),$$

and thus it is enough to show that $\{M(n) \otimes N\}$ is a fibrant tower of (C, D) -bicomodules. For $n = 0$, we have $M(0) \otimes N = 0$, which is fibrant. For $n = 1$, we have $M(1) \otimes N = (C \otimes M) \otimes N$, which is a fibrant (C, D) -bicomodule by Lemma A.8. For $n \geq 1$, we obtain the pullback in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$\begin{array}{ccc} M(n+1) \otimes N & \longrightarrow & (C \otimes P) \otimes N \\ \downarrow & \lrcorner & \downarrow \\ M(n) \otimes N & \longrightarrow & (C \otimes Q) \otimes N. \end{array}$$

The right vertical map is a fibration in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ by Lemma A.6 since its kernel $(C \otimes K) \otimes N$ is a fibrant (C, D) -bicomodule by Lemma A.8, where K is the kernel of $P \rightarrow Q$. Thus the left vertical map is a fibration. \square

Proposition A.10. *Let C , D and E be simply connected dg-coalgebras over \mathbb{k} . If M is a fibrant (D, C) -bicomodule and N is a fibrant (C, E) -bicomodule, then $M \square_C N$ is a fibrant (D, E) -bicomodule.*

Proof. Let $\{M(n)\}$ be the Postnikov tower of M in ${}_D\text{CoMod}_C(\text{Ch}_{\mathbb{k}}^{\geq 0})$. Then M is a retract of $\widetilde{M} \simeq \lim_n M(n)$. As $\widetilde{M} \square_C N = (\lim_n M(n) \square_C N) \cong \lim_n (M(n) \square_C N)$, it is enough to show that $\{M(n) \square_C N\}$ is a fibrant tower of (D, E) -bicomodules. For $n = 0$, then $M(0) \square_C N = 0$ and is thus fibrant. For $n = 1$, we get $M(1) \square_C N \cong (D \otimes M \otimes C) \square_C N \cong (D \otimes M) \otimes N$. By Lemmas A.7 and A.9, it is a fibrant (D, E) -bicomodule. As the functor $-\square_C Y$ preserves pullbacks, we obtain the pullback in ${}_D\text{CoMod}_E(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$\begin{array}{ccc} M(n+1) \square_C N & \longrightarrow & (D \otimes P) \otimes N \\ \downarrow & \lrcorner & \downarrow \\ M(n) \square_C N & \longrightarrow & (D \otimes Q) \otimes N. \end{array}$$

The right vertical map is a fibration by Lemma A.6 as its kernel is $(D \otimes K) \otimes N$, which is a fibrant (D, E) -bicomodule by Lemmas A.7 and A.9, where K is the kernel of $P \rightarrow Q$. Thus the left vertical map is a fibration. \square

One particularly nice characterizing algebraic property of fibrant comodules is that they are *coflat*, i.e., they interplay well with the cotensor product and exact sequences. The cotensor product is left-exact, as it preserves finite products and equalizers. We are interested in knowing the cases in which it preserves exactness, without any cocommutativity requirement.

Definition A.11. Let C and D be flat coalgebras in a symmetric monoidal abelian category, $(\mathbb{C}, \otimes, \mathbb{I})$. Let M be a (C, D) -bicomodule. We say M is *left coflat over C* if given any short exact sequence in CoMod_C

$$0 \longrightarrow N \longrightarrow N' \longrightarrow N'' \longrightarrow 0,$$

we obtain a short exact sequence in \mathbb{C}

$$0 \longrightarrow N \square_C M \longrightarrow N' \square_C M \longrightarrow N'' \square_C M \longrightarrow 0.$$

Similarly, we say M is *right coflat over D* if given any short exact sequence in ${}_D\text{CoMod}$

$$0 \longrightarrow N \longrightarrow N' \longrightarrow N'' \longrightarrow 0,$$

we obtain a short exact sequence in \mathbb{C}

$$0 \longrightarrow M \square_D N \longrightarrow M \square_D N' \longrightarrow M \square_D N'' \longrightarrow 0.$$

We say the bicomodule M is *two-sided coflat over (C, D)* if it is left coflat over C and right coflat over D . Using Definition 1.19, if M is right coflat over C , or if N is left coflat over C , then $\text{CoTor}_C^i(M, N) = 0$ for all $i \geq 1$.

Proposition A.12. *Let C and D be simply connected dg-coalgebras over \mathbb{k} . If a (C, D) -bicomodule is fibrant, then it is a two-sided coflat bicomodule over (C, D) .*

Proof. Notice that any cofree (C, D) -bicomodule $C \otimes V \otimes D$ is two-sided coflat. Let us show that coflatness is preserved under extensions. Consider a short exact sequence in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$,

$$0 \longrightarrow M \longrightarrow N \longrightarrow P \longrightarrow 0.$$

Suppose M and P are two-sided coflat. Let Q be a left D -comodule. We then obtain an exact sequence

$$\text{CoTor}_1^D(M, Q) \longrightarrow \text{CoTor}_1^D(N, Q) \longrightarrow \text{CoTor}_1^D(P, Q).$$

By exactness, we get that $\text{CoTor}_1^D(N, Q) = 0$ for any left D -comodule Q . Thus N is coflat as a right D -comodule. We argue similarly to show that N is coflat as a left C -comodule.

Now let F be a fibrant (C, D) -bicomodule. Let $\{F(n)\}$ be its Postnikov tower in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$ and write $\widetilde{F} = \lim_n F(n)$. Since F is a retract of \widetilde{F} , then for any left D -comodule Q , we get $\text{CoTor}_1^D(F, Q)$ is a retract of $\text{CoTor}_1^D(\widetilde{F}, Q)$. Thus if \widetilde{F} is coflat as a right D -comodule, so is F . We prove \widetilde{F} is right coflat by

induction. For $n = 0$, we see that $F(0) = 0$ is coflat. For $n = 1$, we get that $F(1) = C \otimes F \otimes D$, a cofree bicomodule, and is thus right coflat. Suppose we have shown that $F(n)$ is right coflat over D . Then from the short exact sequence in ${}_C\text{CoMod}_D(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$0 \longrightarrow C \otimes K \otimes D \longrightarrow F(n+1) \longrightarrow F(n) \longrightarrow 0,$$

we get that $F(n+1)$ is also right coflat over D . Given any short exact sequence in ${}_D\text{CoMod}(\text{Ch}_{\mathbb{k}}^{\geq 0})$:

$$0 \longrightarrow Q \longrightarrow Q' \longrightarrow Q'' \longrightarrow 0,$$

we obtain a short exact sequence of towers in $\text{Ch}_{\mathbb{k}}^{\geq 0}$:

$$0 \longrightarrow \{F(n) \square_D W\} \longrightarrow \{F(n) \square_D W'\} \longrightarrow \{F(n) \square_D W''\} \longrightarrow 0.$$

Each of these towers satisfies the Mittag-Leffler condition, and since limit of towers that stabilizes in each degree commute with cotensor product, we obtain a short exact sequence

$$0 \longrightarrow \tilde{F} \square_D Q \longrightarrow \tilde{F} \square_D Q' \longrightarrow \tilde{F} \square_D Q'' \longrightarrow 0.$$

Therefore \tilde{F} is right coflat over D . We can show \tilde{F} is left coflat over C in a similar fashion. \square

Remark A.13. In fact, the result of [Pér24b, 4.7] remains true in the non-commutative case. Thus one can show that a (C, D) -bicomodule is fibrant if and only if it is coflat as a right $(C^{\text{op}} \otimes D)$ -comodule. We shall not need this result here so we do not provide details, however we do mention below the relationship between two-sided coflat (C, D) -comodules and right coflat $(C^{\text{op}} \otimes D)$ -comodules. In particular, by Lemma A.15, a fibrant (C, D) -bicomodule is always two-sided coflat.

Lemma A.14. *Let $(\mathbb{C}, \otimes, \mathbb{I})$ be a symmetric monoidal category. Let $(C, \Delta_C, \varepsilon_C)$ and $(D, \Delta_D, \varepsilon_D)$ be flat coalgebras in \mathbb{C} . Let M be a right $(C \otimes D)$ -comodule. Let N be a left D -comodule. Then $C \otimes N$ is a left $(C \otimes D)$ -comodule, and we obtain an isomorphism in \mathbb{C} :*

$$M \square_{C \otimes D} (C \otimes N) \cong M \square_D N.$$

Proof. Let $\lambda : N \rightarrow D \otimes N$ be the left D -coaction on N . Then we obtain a left $(C \otimes D)$ -coaction on N via

$$C \otimes N \xrightarrow{\text{id}_C \otimes \lambda} C \otimes D \otimes N \xrightarrow{\Delta_C \otimes \text{id}_{D \otimes N}} C \otimes C \otimes D \otimes N \cong C \otimes D \otimes C \otimes N.$$

The above coaction, denoted $\tilde{\lambda} : C \otimes N \rightarrow (C \otimes D) \otimes C \otimes N$, is counital and coassociative because the left D -coaction on N and the coalgebra structure on C are both coassociative and counital.

To prove the desired isomorphism, we verify that $M \square_D N$ satisfies the universal property of the equalizer. From the right $(C \otimes D)$ -coaction $\rho : M \rightarrow M \otimes C \otimes D$, we obtain the underlying right C -coaction ρ_C on M as the composite

$$M \xrightarrow{\rho} M \otimes C \otimes D \xrightarrow{\text{id}_M \otimes \varepsilon_D} M \otimes C.$$

Now we obtain a morphism $\alpha : M \square_D N \rightarrow M \square_{C \otimes D} (C \otimes N)$ by functoriality of the equalizers:

$$\begin{array}{ccccc} M \square_D N & \longrightarrow & M \otimes N & \xrightarrow[\text{id}_M \otimes \lambda]{\rho_D \otimes \text{id}_N} & M \otimes D \otimes N \\ \alpha \downarrow \exists! & & \downarrow \rho_C \otimes \text{id}_N & & \downarrow \\ M \square_{C \otimes D} (C \otimes N) & \longrightarrow & M \otimes (C \otimes N) & \xrightarrow[\text{id}_M \otimes \tilde{\lambda}]{\rho \otimes \text{id}_{C \otimes N}} & M \otimes (C \otimes D) \otimes (C \otimes N). \end{array}$$

The right unlabeled vertical arrow is induced by applying the functor $- \otimes N$ on the map

$$M \otimes D \xrightarrow{\rho_C \otimes \text{id}_D} (M \otimes C) \otimes D \xrightarrow{\text{id}_M \otimes \Delta_C \otimes \text{id}_D} \underbrace{(M \otimes (C \otimes C)) \otimes D}_{\cong M \otimes (C \otimes D) \otimes C}.$$

Similarly, by applying the counit map $\varepsilon_C: C \rightarrow \mathbb{I}$ vertically on each C , we obtain the dashed map on the equalizers below:

$$\begin{array}{ccccc}
M \square_{C \otimes D}(C \otimes N) & \longrightarrow & M \otimes C \otimes N & \xrightarrow[\text{id}_M \otimes \tilde{\lambda}]{\rho \otimes \text{id}_{C \otimes N}} & M \otimes (C \otimes D) \otimes C \otimes N \\
\downarrow \beta \dashv \exists! & & \downarrow & & \downarrow \\
M \square_D N & \longrightarrow & M \otimes N & \xrightarrow[\text{id}_M \otimes \lambda]{\rho_D \otimes \text{id}_N} & M \otimes D \otimes N.
\end{array}$$

The induced morphism β is the inverse of the morphism α defined above. Therefore we obtain the desired isomorphism in \mathcal{C} . \square

Lemma A.15. *Let $(\mathcal{C}, \otimes, \mathbb{I})$ be a symmetric monoidal abelian category. Let C and D be flat coalgebras in \mathcal{C} . Let M be a (C, D) -bicomodule in \mathcal{C} . If M is right coflat as a right $(C^{\text{op}} \otimes D)$ -comodule, then it is two-sided coflat as a (C, D) -bicomodule.*

Proof. Suppose M is right coflat a $(C^{\text{op}} \otimes D)$ -comodule. Let us first show that M is right coflat over D . Consider a short exact sequence in ${}_D \text{CoMod}$:

$$0 \longrightarrow N \longrightarrow N' \longrightarrow N'' \longrightarrow 0.$$

By the previous lemma, we obtain that $C^{\text{op}} \otimes N$, $C^{\text{op}} \otimes N'$, and $C^{\text{op}} \otimes N''$ are left $(C^{\text{op}} \otimes D)$ -comodules. Since $C^{\text{op}} \otimes -$ preserves exactness because it is flat, we obtain a short exact sequence in $(C^{\text{op}} \otimes D) \text{CoMod}$:

$$0 \longrightarrow C^{\text{op}} \otimes N \longrightarrow C^{\text{op}} \otimes N' \longrightarrow C^{\text{op}} \otimes N'' \longrightarrow 0.$$

Since M is a right coflat over $(C^{\text{op}} \otimes D)$, we obtain that the sequence is exact in \mathcal{C} :

$$0 \longrightarrow M \square_{C^{\text{op}} \otimes D} C^{\text{op}} \otimes N \longrightarrow M \square_{C^{\text{op}} \otimes D} C^{\text{op}} \otimes N' \longrightarrow M \square_{C^{\text{op}} \otimes D} C^{\text{op}} \otimes N'' \longrightarrow 0.$$

By the previous lemma, this induces the short exact sequence in \mathcal{C} :

$$0 \longrightarrow M \square_D N \longrightarrow M \square_D N' \longrightarrow M \square_D N'' \longrightarrow 0.$$

Thus M is right coflat over D as desired. We prove that M is left coflat over C in a similar fashion. \square

Example A.16. The converse is not true: a two-sided coflat (C, D) -bicomodule need not to be a right coflat $(C^{\text{op}} \otimes D)$ -comodule, and thus need not be a fibrant (C, D) -bicomodule. This can already be seen for modules over an algebra (even commutative). Indeed, consider the polynomial ring, $\mathbb{k}[x]$, as a \mathbb{k} -algebra. It is a two-sided flat $\mathbb{k}[x]$ -module, but it is not flat as a right $(\mathbb{k}[x] \otimes \mathbb{k}[x])$ -module. Indeed, note first that $\mathbb{k}[x, y] \cong \mathbb{k}[x] \otimes \mathbb{k}[x]$. Consider the following short exact sequence of left $\mathbb{k}[x, y]$ -modules

$$0 \longrightarrow \mathbb{k}[x, y] \xrightarrow{\cdot y} \mathbb{k}[x, y] \longrightarrow \mathbb{k}[x] \longrightarrow 0.$$

If we apply $\mathbb{k}[x] \otimes_{\mathbb{k}[x, y]} -$, we obtain the exact sequence

$$\mathbb{k}[x] \xrightarrow{0} \mathbb{k}[x] \longrightarrow \mathbb{k}[x] \longrightarrow 0.$$

The above sequence is not a short exact sequence however, which shows that $\mathbb{k}[x]$ is not flat as a right $\mathbb{k}[x, y]$ -module.

We are now equipped to prove a crucial result in this paper: the cotensor product with a fibrant comodule preserves quasi-isomorphisms.

Proposition A.17. *Let C and D be simply connected dg-coalgebras over \mathbb{k} . Let M be a fibrant (C, D) -bicomodule. Then $M \square_D -: {}_D \text{CoMod}(\text{Ch}_{\mathbb{k}}^{\geq 0}) \rightarrow \text{Ch}_{\mathbb{k}}^{\geq 0}$ and $-\square_C M : \text{CoMod}_C(\text{Ch}_{\mathbb{k}}^{\geq 0}) \rightarrow \text{Ch}_{\mathbb{k}}^{\geq 0}$ preserve weak equivalences.*

Proof. We use an Eilenberg–Moore spectral sequence argument. Let N be any left D -comodule in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. As in Remark 3.11, the conormalized cobar complex $\underline{\Omega}^{\bullet}(M, D, N)$ is a second quadrant double chain complex that is bounded. We grade the row cohomologically, but the columns homologically. Hence its two associated spectral sequences converge to the same page, see [McC01, 2.15].

The first spectral sequence has its E^1 -page induced by the cohomology of the rows and therefore

$$E_{\bullet,q}^1 = H^q(\underline{\Omega}^\bullet(M, D, N)) \cong \text{CoTor}_D^q(M, N).$$

Since M is a fibrant (C, D) -bicomodule, then it is coflat as a right D -comodule. Thus $E_{\bullet,q}^1 = 0$ for all $q \geq 1$, and $E_{\bullet,0}^1 = M \square_D N$. Hence the spectral sequence collapses onto its second page, $E_{\bullet,0}^2 = H_*(M \square_D N)$.

The second spectral sequence has its E^1 -page induced by the homology of its columns and therefore

$$E_{\bullet,q}^1 = H_*(\underline{\Omega}^q(X, D, Y)) = \underline{\Omega}^q(H_*(M), H_*(D), H_*(N)).$$

Thus, as its E^2 -page is given by the cohomology of the induced cochain complex, we obtain

$$E_{\bullet,q}^2 = \text{CoTor}_{H_*(D)}^q(H_*(M), H_*(N)).$$

It converges to the page with trivial columns except its 0^{th} column, which is given by the cohomology $H_*(M \square_D N)$.

Combining the arguments above, we obtain a converging Eilenberg–Moore spectral sequence

$$E_{\bullet,q}^2 = \text{CoTor}_{H_*(D)}^q(H_*(M), H_*(N)) \Rightarrow E_{\bullet,0}^\infty = H_*(M \square_D N),$$

for any left D -comodule N . In particular, given a map of left D -comodules $N \rightarrow N'$ that is a quasi-isomorphism, we get

$$\text{CoTor}_{H_*(D)}^q(H_*(M), H_*(N)) \cong \text{CoTor}_{H_*(D)}^q(H_*(M), H_*(N')).$$

Thus the map $N \rightarrow N'$ induces an isomorphism, $H_*(M \square_D N) \cong H_*(M \square_D N')$. \square

Corollary A.18. *Let C , D and E be simply connected coalgebras in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. Let M be a fibrant (D, C) -bicomodule and N a fibrant (C, E) -bicomodule. Then we obtain a quasi-isomorphism of (D, E) -bicomodules*

$$M \square_C N \simeq \Omega(M, C, N).$$

Corollary A.19. *Let C , D , and E be simply connected coalgebras in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. Let M be a fibrant (C, D) -bicomodule, let N be a left fibrant D -comodule and P be a fibrant right E -comodule. Then there is a natural quasi-isomorphism of (C, E) -bicomodules*

$$\Omega(M, D, N) \otimes P \simeq \Omega(M, C, N \otimes P).$$

Proof. Since M and N are fibrant, we obtain

$$\begin{aligned} \Omega(M, D, N) \otimes P &\simeq (M \square_D N) \otimes P \\ &\cong M \square_D (N \otimes P) \\ &\simeq \Omega(M, D, N \otimes P). \end{aligned}$$

The isomorphism follows from the fact that P is flat in $\text{Ch}_{\mathbb{k}}^{\geq 0}$, and thus preserves equalizers. The last quasi-isomorphism follows from the fact that $N \otimes P$ remains a fibrant (D, E) -bicomodule by Lemma A.9. \square

Remark A.20. Corollary A.19 is not true if the comodules are not fibrant, since in general the tensor product does not preserve infinite homotopy limits.

Corollary A.21. *Let C , D , E , and F be simply connected coalgebras in $\text{Ch}_{\mathbb{k}}^{\geq 0}$. Let M be a fibrant (C, D) -bicomodule, let N be a fibrant (D, E) -bicomodule, and let P be a fibrant (E, F) -bicomodule. Then we obtain a quasi-isomorphism of (C, F) -bicomodules*

$$\Omega(\Omega(M, D, N), E, P) \simeq \text{holim}_{\Delta \times \Delta} \Omega^\bullet(\Omega^\bullet(M, D, N), E, P).$$

Proof. By the previous Corollary, for any $i \geq 0$

$$\begin{aligned} \text{holim}_{\Delta}(\Omega^\bullet(M, D, N) \otimes E^{\otimes i} \otimes P) &\simeq \text{holim}_{\Delta}(\Omega^\bullet(M, D, N \otimes E^{\otimes i} \otimes P)) \\ &\simeq \Omega(M, D, N \otimes E^{\otimes i} \otimes P) \\ &\simeq \Omega(M, D, N) \otimes E^{\otimes i} \otimes P \\ &\simeq \text{holim}_{\Delta}(\Omega^\bullet(M, D, N)) \otimes E^{\otimes i} \otimes P. \end{aligned}$$

In other words, we have proved that we obtain a quasi-isomorphism between fibrant (C, F) -bicomodules (recall that we fixed a model for the homotopy limit):

$$\mathrm{holim}_{\Delta}(\Omega^i(\Omega^{\bullet}(M, D, N), E, P)) \simeq \Omega^i(\mathrm{holim}_{\Delta}\Omega^{\bullet}(M, D, N), E, P).$$

This above equivalence is compatible with cofaces and codegeneracies of the cobar construction, and therefore we obtain by [Hir03, 18.5.2, 18.5.3]:

$$\begin{aligned} \Omega(\Omega(M, D, N), E, P) &\simeq \mathrm{holim}_{\Delta}\Omega^{\bullet}(\Omega(M, D, N), E, P) \\ &\simeq \mathrm{holim}_{\Delta}\Omega^{\bullet}(\mathrm{holim}_{\Delta}\Omega^{\bullet}(M, D, N), E, P) \\ &\simeq \mathrm{holim}_{\Delta}\mathrm{holim}_{\Delta}\Omega^{\bullet}(\Omega^{\bullet}(M, D, N), E, P) \\ &\simeq \mathrm{holim}_{\Delta \times \Delta}\Omega^{\bullet}(\Omega^{\bullet}(M, D, N), E, P). \end{aligned} \quad \square$$

Corollary A.22. *Let C , D , and E be simply connected coalgebras in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$. Let M be a fibrant (C, D) -bicomodule and N a fibrant (D, E) -bicomodule. Then we obtain a quasi-isomorphism*

$$\mathrm{coHH}(\Omega(M, D, N), C) \simeq \mathrm{holim}_{\Delta \times \Delta} \mathrm{coHH}^{\bullet}(\Omega^{\bullet}(M, D, N), E, P).$$

Proof. By Corollary A.19, for any $i \geq 0$

$$\mathrm{holim}_{\Delta}(\Omega^{\bullet}(M, D, N) \otimes C^{\otimes i}) \simeq \mathrm{holim}_{\Delta}(\Omega^{\bullet}(M, D, N)) \otimes C^{\otimes i}.$$

The above equivalence is compatible with cofaces and codegeneracies of the cobar construction, and therefore we obtain [Hir03, 18.5.2, 18.5.3]:

$$\begin{aligned} \mathrm{coHH}(\Omega(M, D, N), C) &\simeq \mathrm{holim}_{\Delta} \mathrm{coHH}^{\bullet}(\mathrm{holim}_{\Delta}\Omega^{\bullet}(M, D, N), C) \\ &\simeq \mathrm{holim}_{\Delta \times \Delta} \mathrm{coHH}^{\bullet}(\Omega^{\bullet}(M, D, N), C). \end{aligned} \quad \square$$

Proposition A.23. *Let C be a simply connected in $\mathrm{Ch}_{\mathbb{k}}^{\geq 0}$. Let M be a fibrant (C, C) -bicomodule. There is a quasi-isomorphism*

$$\mathrm{coHH}(M, C) \simeq \Omega(M, C \otimes C^{\mathrm{op}}, C).$$

Proof. We denote the enveloping coalgebra by $C^e = C \otimes C^{\mathrm{op}}$. Recall that we have the quasi-isomorphism $C \simeq \Omega(C, C, C)$ as (C, C) -bicomodules. Thus we obtain

$$M \hat{\square}_{C^e} C \simeq M \hat{\square}_{C^e} \Omega(C, C, C).$$

Notice that each object in the cosimplicial diagram $\Omega^{\bullet}(C, C, C)$ is a fibrant (C, C) -bicomodule by Lemma A.8. Thus $\Omega(C, C, C)$ is a fibrant (C, C) -bicomodule by [Hir03, 18.5.2]. Therefore

$$M \hat{\square}_{C^e} \Omega(C, C, C) \simeq M \square_{C^e} \Omega(C, C, C).$$

Since the functor $M \square_{C^e} -$ preserves towers that stabilize in each degree, we get

$$M \square_{C^e} \Omega(C, C, C) \simeq \mathrm{holim} \left(M \square_{C^e} \Omega^{\bullet}(C, C, C) \right).$$

We have an isomorphism, $\mathrm{coHH}^n(M, C) \cong M \square_{C^e} \Omega^n(C, C, C)$, for each $n \geq 0$, induced by

$$M \otimes C^n \cong M \square_{C^e} (C^e \otimes C^n) \cong M \square_{C^e} (C \otimes C^n \otimes C),$$

given by the permutation of $C^{\mathrm{op}} \cong C$ past C^n . These isomorphisms are compatible with the cosimplicial structures and thus provide an isomorphism, $\mathrm{coHH}^{\bullet}(M, C) \cong M \square_{C^e} \Omega^{\bullet}(C, C, C)$. \square

REFERENCES

- [Al-02] Khaled Al-Takhman, *Equivalences of comodule categories for coalgebras over rings*, J. Pure Appl. Algebra **173** (2002), no. 3, 245–271. MR 1916479
- [AR05] Vigleik Angeltveit and John Rognes, *Hopf algebra structure on topological Hochschild homology*, Algebr. Geom. Topol. **5** (2005), 1223–1290. MR 2171809
- [Bar23] Justin Barhite, *Bicategorical traces and cotraces*, 2023, arXiv2307.13070.v1.
- [BGH⁺18] Anna Marie Bohmann, Teena Gerhardt, Amalie Høgenhaven, Brooke Shipley, and Stephanie Ziegenhagen, *Computational tools for topological coHochschild homology*, Topology Appl. **235** (2018), 185–213. MR 3760202
- [BGS22] Anna Marie Bohmann, Teena Gerhardt, and Brooke Shipley, *Topological coHochschild homology and the homology of free loop spaces*, Math. Z. **301** (2022), no. 1, 411–454. MR 4405656

- [BHK⁺15] Marzieh Bayeh, Kathryn Hess, Varvara Karpova, Magdalena Kędziołek, Emily Riehl, and Brooke Shipley, *Left-induced model structures and diagram categories*, Women in topology: collaborations in homotopy theory, Contemp. Math., vol. 641, Amer. Math. Soc., Providence, RI, 2015, pp. 49–81. MR 3380069
- [BHM93] M. Bökstedt, W. C. Hsiang, and I. Madsen, *The cyclotomic trace and algebraic K-theory of spaces*, Invent. Math. **111** (1993), no. 3, 465–539. MR 1202133
- [BM11] Andrew J. Blumberg and Michael A. Mandell, *Derived Koszul duality and involutions in the algebraic K-theory of spaces*, J. Topol. **4** (2011), no. 2, 327–342. MR 2805994
- [BM12] ———, *Localization theorems in topological Hochschild homology and topological cyclic homology*, Geom. Topol. **16** (2012), no. 2, 1053–1120. MR 2928988
- [BP23] Özgür Haldun Bayındır and Maximilien Péroux, *Spanier-Whitehead duality for topological coHochschild homology*, J. Lond. Math. Soc. (2) **107** (2023), no. 5, 1780–1822. MR 4585302
- [Bun12] Ulrich Bunke, *Differential cohomology*, arXiv e-prints (2012), arXiv:1208.3961.
- [BW03] Tomasz Brzezinski and Robert Wisbauer, *Corings and comodules*, London Mathematical Society Lecture Note Series, vol. 309, Cambridge University Press, Cambridge, 2003. MR 2012570
- [CP19] Jonathan A. Campbell and Kate Ponto, *Topological Hochschild homology and higher characteristics*, Algebr. Geom. Topol. **19** (2019), no. 2, 965–1017. MR 3924181
- [Doi81] Yukio Doi, *Homological coalgebra*, J. Math. Soc. Japan **33** (1981), no. 1, 31–50. MR 597479
- [DP80] Albrecht Dold and Dieter Puppe, *Duality, trace, and transfer*, Proceedings of the International Conference on Geometric Topology (Warsaw, 1978), PWN, Warsaw, 1980, pp. 81–102. MR 656721
- [Dun97] Bjørn Ian Dundas, *Relative K-theory and topological cyclic homology*, Acta Math. **179** (1997), no. 2, 223–242. MR 1607556
- [EKMM97] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May, *Rings, modules, and algebras in stable homotopy theory*, Mathematical Surveys and Monographs, vol. 47, American Mathematical Society, Providence, RI, 1997, With an appendix by M. Cole. MR 1417719
- [FS98] Marco A. Farinati and Andrea Solotar, *Morita-Takeuchi equivalence, cohomology of coalgebras and Azumaya coalgebras*, Rings, Hopf algebras, and Brauer groups (Antwerp/Brussels, 1996), Lecture Notes in Pure and Appl. Math., vol. 197, Dekker, New York, 1998, pp. 119–146. MR 1615805
- [GKR20] Richard Garner, Magdalena Kędziołek, and Emily Riehl, *Lifting accessible model structures*, J. Topol. **13** (2020), no. 1, 59–76. MR 3999672
- [GP87] Luzius Grünenfelder and Robert Paré, *Families parametrized by coalgebras*, J. Algebra **107** (1987), no. 2, 316–375. MR 885801
- [Hat65] Akira Hattori, *Rank element of a projective module*, Nagoya Math. J. **25** (1965), 113–120. MR 175950
- [Heu20] Gijs Heuts, *Koszul duality and a conjecture of Francis–Gaitsgor*, Topologie (hybrid meeting), Oberwolfach Reports, vol. 17, 2020, pp. 1083–1137. MR 4281706
- [Hir03] Philip S. Hirschhorn, *Model categories and their localizations*, Mathematical Surveys and Monographs, vol. 99, American Mathematical Society, Providence, RI, 2003. MR 1944041
- [HKRS17] Kathryn Hess, Magdalena Kędziołek, Emily Riehl, and Brooke Shipley, *A necessary and sufficient condition for induced model structures*, J. Topol. **10** (2017), no. 2, 324–369. MR 3653314
- [Hov99] Mark Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999. MR 1650134
- [HPS09] Kathryn Hess, Paul-Eugène Parent, and Jonathan Scott, *CoHochschild homology of chain coalgebras*, J. Pure Appl. Algebra **213** (2009), no. 4, 536–556. MR 2483836
- [HR21a] Kathryn Hess and Nima Rasekh, *Shadows are Bicategorical Traces*, arXiv e-prints (2021), arXiv:2109.02144.
- [HR21b] Geoffroy Horel and Maxime Ramzi, *A multiplicative comparison of Mac Lane homology and topological Hochschild homology*, Ann. K-Theory **6** (2021), no. 3, 571–605. MR 4310330
- [HS16] Kathryn Hess and Brooke Shipley, *Waldhausen K-theory of spaces via comodules*, Adv. Math. **290** (2016), 1079–1137. MR 3451948
- [HS21] ———, *Invariance properties of coHochschild homology*, J. Pure Appl. Algebra **225** (2021), no. 2, Paper No. 106505, 27. MR 4128640
- [Kla22] Sarah Klanderma, *Computations of relative topological coHochschild homology*, J. Homotopy Relat. Struct. **17** (2022), no. 3, 393–417. MR 4470385
- [Lan02] Serge Lang, *Algebra*, third ed., Graduate Texts in Mathematics, vol. 211, Springer-Verlag, New York, 2002. MR 1878556
- [Lod98] Jean-Louis Loday, *Cyclic homology*, second ed., Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 301, Springer-Verlag, Berlin, 1998, Appendix E by María O. Ronco, Chapter 13 by the author in collaboration with Teimuraz Pirashvili. MR 1600246
- [Lur17] Jacob Lurie, *Higher algebra*, <https://www.math.ias.edu/~lurie/papers/HA.pdf>, 2017, electronic book.
- [Mal17] Cary Malkiewich, *Cyclotomic structure in the topological Hochschild homology of DX*, Algebr. Geom. Topol. **17** (2017), no. 4, 2307–2356. MR 3686399
- [McC01] John McCleary, *A user’s guide to spectral sequences*, second ed., Cambridge Studies in Advanced Mathematics, vol. 58, Cambridge University Press, Cambridge, 2001. MR 1793722
- [NS18] Thomas Nikolaus and Peter Scholze, *On topological cyclic homology*, Acta Math. **221** (2018), no. 2, 203–409. MR 3904731

- [Pér22a] Maximilien Péroux, *The coalgebraic enrichment of algebras in higher categories*, J. Pure Appl. Algebra **226** (2022), no. 3, Paper No. 106849, 11. MR 4291529
- [Pér22b] ———, *Coalgebras in the Dwyer-Kan localization of a model category*, Proc. Amer. Math. Soc. **150** (2022), no. 10, 4173–4190. MR 4470166
- [Pér24a] ———, *A monoidal Dold-Kan correspondence for comodules*, J. Pure Appl. Algebra **228** (2024), no. 8, Paper No. 107660. MR 4720531
- [Pér24b] ———, *Rigidification of connective comodules*, 2024, to appear in Proc. Amer. Soc., arXiv:2006.09398.v5.
- [Pon10] Kate Ponto, *Fixed point theory and trace for bicategories*, Astérisque (2010), xii+102. MR 2741967
- [Pos23] Leonid Positselski, *Homological full-and-faithfulness of comodule inclusion and contra-module forgetful functors*, 2023, arXiv:2301.09561.v2.
- [PS13] Kate Ponto and Michael Shulman, *Shadows and traces in bicategories*, J. Homotopy Relat. Struct. **8** (2013), no. 2, 151–200. MR 3095324
- [PS14] ———, *Traces in symmetric monoidal categories*, Expo. Math. **32** (2014), no. 3, 248–273. MR 3253568
- [PS19] Maximilien Péroux and Brooke Shipley, *Coalgebras in symmetric monoidal categories of spectra*, Homology Homotopy Appl. **21** (2019), no. 1, 1–18. MR 3852287
- [Qui88] Daniel Quillen, *Algebra cochains and cyclic cohomology*, Inst. Hautes Études Sci. Publ. Math. (1988), no. 68, 139–174 (1989). MR 1001452
- [Qui89] ———, *Cyclic cohomology and algebra extensions*, K-Theory **3** (1989), no. 3, 205–246. MR 1040400
- [Rav86] Douglas C. Ravenel, *Complex cobordism and stable homotopy groups of spheres*, Pure and Applied Mathematics, vol. 121, Academic Press, Inc., Orlando, FL, 1986. MR 860042
- [Sta65] John Stallings, *Centerless groups—an algebraic formulation of Gottlieb’s theorem*, Topology **4** (1965), 129–134. MR 202807
- [Swa70] Richard G. Swan, *K-theory of finite groups and orders*, Lecture Notes in Mathematics, vol. Vol. 149, Springer-Verlag, Berlin-New York, 1970. MR 308195
- [Tak77] Mitsuhiro Takeuchi, *Morita theorems for categories of comodules*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **24** (1977), no. 3, 629–644. MR 472967
- [Wal79] Friedhelm Waldhausen, *Algebraic K-theory of topological spaces. II*, Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978), Lecture Notes in Math., vol. 763, Springer, Berlin, 1979, pp. 356–394. MR 561230
- [Wal85] ———, *Algebraic K-theory of spaces*, Algebraic and geometric topology (New Brunswick, N.J., 1983), Lecture Notes in Math., vol. 1126, Springer, Berlin, 1985, pp. 318–419. MR 802796
- [Wei94] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR 1269324
- [Wei13] ———, *The K-book*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013, An introduction to algebraic K-theory. MR 3076731

DEPARTMENT OF MATHEMATICAL AND COMPUTATIONAL SCIENCES, MARIAN UNIVERSITY, 3200 COLD SPRING ROAD, INDIANAPOLIS, IN, 46222, USA

Email address: sklanderman@marian.edu

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, 619 RED CEDAR RD, EAST LANSING, MI 48824, USA

Email address: peroux@msu.edu