Understanding parton evolution in matter from renormalization group analysis

Weiyao Ke^{*} and Ivan Vitev^{\dagger}

Theoretical Division, Los Alamos National Laboratory, Los Alamos NM 87545, United States

(Dated: January 31, 2023)

We perform a renormalization group (RG) analysis of collinear hadron production in deep inelastic scattering on nuclei. We consider the limit where one of the dimensionless in-medium scale ratios $E/(\mu_D^2 L) \gg 1$, with L, μ_D, E being the medium size, inverse scattering range and the parton energy in the nuclear rest frame, while the opacity L/λ_g remains small. We identify the fixed order and leading $\ln[E/(\mu_D^2 L)]$ enhanced medium contributions to the semi-inclusive cross sections and derive RG equations which resum multiple emissions near the $x \to 0, 1$ endpoints of the splitting functions at first order in opacity. We find that the evolution equations obtained in this work treat the same type of radiation enhancement in matter as the modified DGLAP approach, but differ in the way one chooses to regulate the endpoint divergences and provide unique *analytic* insight into the problem of resummation. The new RG evolution framework is applied to study fragmentation in eA reactions.

Introduction. A common characteristic of many problems in science is that microscopic fluctuations in the system manifest themselves in macroscopic effects. Such problems arise in fields ranging from social networks [1] and turbulence [2] to particle [3] and nuclear physics [4, 5]. They are most prevalent in inherently divergent theories and efficiently addressed using renormalization group (RG) analysis [6, 7]. Effective theories of quantum chromodynamics (QCD) geared toward jet physics [8, 9] have provided new insights into renormalization and resummation, and given a modern perspective to the problem of parton production and propagation in nuclear matter [10–14]. These advances are key to the interpretation of the data from reactions with nuclei at current and future colliders [15].

Over the past two decades, medium-induced parton showers have been successfully implemented in jet quenching phenomenology to describe the modification of hadron and jet cross sections, and jet substructure in nuclear collisions [16–28]. Still, resummation of QCD radiation in nuclear matter remains challenging, especially lacking in analytic insight. We address this longstanding problem using RG techniques. If we consider semi-inclusive hadron production in deep inelastic scattering (DIS) on a nuclear target $(eA \rightarrow h + X)$, we encounter a number of energy and length scales (defined in the target rest frame) including: 1) the hard scale Q, 2) energy of the virtual photon/jet ν , 3) the path length L, 4) the mean free path $\lambda_{\rm g}$ and 5) the inverse interaction range μ_D . Therefore, observables in eA are functions of many dimensionless control parameters

$$Obs \equiv Obs\left(\frac{Q}{Q_0}, \frac{E}{\mu_D^2 L}, \frac{L}{\lambda_g}, \lambda_g \mu_D, \frac{\mu_D}{Q_0}, \cdots\right).$$
(1)

A simplified description with controlled accuracy is often possible when one (or more) of these dimensionless ratios become asymptotically large [29]. For example, the limit $\lambda_g \mu_D \gg 1$, implying independent multiple parton-medium scatterings, allows the use of timeordered perturbation theory to derive quark and gluon splitting functions in matter [13, 30]. A partonic transport picture emerges when $L/\lambda_g \gg 1$ in a thick and dense medium [31–33].

In this letter, we compute hadron production in a particular regime when Q/Q_0 , $E/(\mu_D^2 L)$, $\lambda_g \mu_D$ become asymptotically large while L/λ_g , μ_D/Q_0 stay at order unity/few. This limit is particularly relevant for high-energy hadron production in thin, dilute or fast-expanding media. Still, renormalization is needed to resum large $\ln[Q/Q_0]$ and $\ln[E/(\mu_D^2 L)]$ enhancements from the vacuum and medium-induced radiative corrections. We thus introduce two final-state renormalization scales μ_1 and μ_2 in the single-inclusive hadron cross section [34]

$$\frac{d\sigma_{eA\to h}}{dx_B dQ^2 dz_h} = \frac{2\pi\alpha_e^2}{Q^4} \sum_{i,j} e_q^2 \Big\{ \Big\{ f_{i/A} \otimes \Big[(1+(1-y)^2)C_{ij}^1 + 2(1-y)C_{ij}^L \Big] \Big\}_{x_B} \otimes d_{h/j} \Big\}_{z_h},$$
(2)

$$\{h \otimes g\}_x \equiv \int_x^1 h\left(\frac{x}{x'}\right) g(x') \frac{dx'}{x'} \,. \tag{3}$$

Here, $y = \nu/E_e$ with E_e and $\nu = Q^2/(2x_BM_p)$ being the energies of the incident electron and the virtual photon, respectively. $f_{i/A}(x, Q^2), d_{h/j}(z, \mu_1^2, \mu_2^2)$ and $C_{ii}^{1,L}(x,z,Q^2,\mu_1^2,\mu_2^2)$ are the parton distribution functions, fragmentation functions, and the hard coefficient functions with fractional electric charge e_q . The PDF is evaluated at scale Q^2 , and the dependence on Q^2 will not be written explicitly hereafter. The cross-section $f_{i/A} \otimes C_{ij}^{1,L} \otimes d_{j/h}$ must not depend on μ_1 and μ_2 , thus it is sufficient to study scale dependence of $F_{ij}^{1,L}(z) \equiv$ $f_{i/A} \otimes z C_{ij}^{1,L}$ [35]. This quantity can be interpreted as the invariant distribution of parton j, resolved at scales μ_1, μ_2 , after the hard process. This "parton shower" choice enables us to track the evolving parton energy $E = z\nu$, which is important for the most consistent implementation of medium-induced splitting functions. $F_{ij}^{1} = z\delta_{ij}f_{i/A}(x_B)\delta(1-z) + \alpha_s(\cdots), F_{ij}^{L} = \alpha_s(\cdots), \text{ and}$ we use the NLO expression for $C_{ij}^{1,L}$ [34, 36] without writing them explicitly.

As will become clear in a moment, the Q/Q_0 and $E/(\mu_D^2 L)$ enhancements have distinct physics origins. Therefore, in addition to the vacuum renormalization that leads to the DGLAP evolution, the "medium bare" F_{ij} needs to be further renormalized by a medium coefficient M_{kj} that only depends on μ_2 ,

$$F_{ij}(z,\mu_1^2,\mu_2^2) \to F_{ik}(y,\mu_1^2,\mu_2^2) \otimes M_{kj}\left(\frac{z}{y},\mu_2^2\right) + \mathcal{F}(z).$$
(4)

Here, the $M_{kj} = M_{kj}^{(0)} + M_{kj}^{(1)} + \cdots$ with $M_{kj}^{(0)} = y\delta_{kj}\delta(1-y)$ and the first non-trivial contribution $M_{kj}^{(1)}$. $\mathcal{F}(z)$ stands for medium contributions subleading in $\ln[E/(\mu_D^2 L)]$, i.e. other fixed order contributions.

Renormalization group analysis of endpoint divergences in collinear emission spectra. We consider the correction to $F_{ij}(z)$ from both vacuum and medium-induced collinear splittings and find that it has the form

$$\frac{\alpha_s(\mu_1^2)\mu_1^{2\epsilon}}{2\pi^2} \int^{Q^2} \frac{d^{2-2\epsilon}\mathbf{k}}{\mathbf{k}^2} F_{ik} \otimes x[P_{kj}]_+ + \Delta F_{ij}^{\mathrm{m}}(z,\mu_2^{2\epsilon}).$$
(5)

We work in $d = 4 - 2\epsilon$ dimensions and P_{kj} are the vacuum Altarelli-Parisi splitting functions [37, 38], with the

"+" prescription in Eq. (5) only applied to diagonal contributions. In the first term, μ_1 acts as an infrared cutoff of vacuum emissions. In this case RG analysis leads to DGLAP evolution of $F_{ij}(z)$ in μ_1 that resums $\ln[Q/Q_0]$, and we always evolve μ_1 from the hard scale Q down to Q_0 . Next, we will extract the leading logarithmic and fixed order contribution from the medium-induced correction in the second term.

For thin and uniform nuclear matter of length L we use the medium-induced splitting functions $P_{ij}^{(1)}(x)$ [13, 14, 30, 39] obtained in the opacity expansion approach using Soft-Collinear-Effective-Theory with Glauber Gluons (SCET_G) [10–12]. The full expressions involve integration over both the transverse momentum of the radiated parton \mathbf{k} and the transverse momentum of the Glauber gluon \mathbf{q} that mediates jet-medium interactions, and are included in the supplementary material for completeness. They contain transverse momentum propagator terms of the form $\frac{\mathbf{Va}}{\mathbf{Va}^2} \cdot \frac{\mathbf{Vb}}{\mathbf{Vb}^2}$, with $\mathbf{Va,b}$ being any vectors among $\mathbf{k}, \mathbf{k} - x\mathbf{q}, \mathbf{k} - \mathbf{q}, \mathbf{k} - (1 - x)\mathbf{q}$. Nevertheless, in the large parton phase space limit by shifting the integration variable \mathbf{k} we can cast $P_{ij}^{(1)}(x)$ into a generic form,

$$P_{ij}^{(1)}(x, E, \mu_2^2) = \frac{\alpha_s^{(0)} P_{ij}(x)}{2\pi^2} L \int \frac{\mu_2^{2\epsilon} d^{2-2\epsilon} \mathbf{k}}{(2\pi)^{-2\epsilon}} \frac{\Phi\left[\frac{\mathbf{k}^2 L}{2x(1-x)E}\right]}{\mathbf{k}^2} \sum_n \int \frac{\mu_2^{2\epsilon} d^{2-2\epsilon} \mathbf{q}}{(2\pi)^{-2\epsilon}} \frac{\rho_G \times 4\pi \alpha_s^{(0)} C_n^{ij} \Delta_n^{ij}(x)}{(2\pi)^2 (\mathbf{q}^2 + \mu_D^2)^2} \times \frac{\mathbf{q} \cdot [\mathbf{k} + \Delta_n^{ij}(x)\mathbf{q}]}{[\mathbf{k} + \Delta_n^{ij}(x)\mathbf{q}]^2} \\ = \frac{\alpha_s^2(\mu_2^2) L^2 \rho_G P_{ij}(x)}{8E[x(1-x)]^{1+2\epsilon}} \sum_n C_n^{ij} (\Delta_n^{ij})^{2-2\epsilon} \left[\frac{e^{\gamma_E} \mu_2^2 L}{2E}\right]^{2\epsilon} \frac{\epsilon \Gamma(\epsilon)}{\Gamma(1-\epsilon)} \int_0^{w_{\max}} dw \frac{4}{\pi} \frac{\Phi(w)}{w^{1+\epsilon}} \int_0^1 \frac{du}{(1-w)^{\epsilon}} \frac{-\epsilon uw + v \frac{1-\epsilon}{2} (\Delta_n^{ij})^2}{[uw + v(\Delta_n^{ij})^2]^{2+\epsilon}} \\ \approx \frac{A(\mu_2^2, E, w_{\max}) P_{ij}(x)}{[x(1-x)]^{1+2\epsilon}} \sum_n C_n^{ij} (\Delta_n^{ij})^{2-2\epsilon} \left[\frac{\mu_2^2 L}{\chi(w_{\max})E}\right]^{2\epsilon} (1 + \mathcal{O}(\epsilon^2)).$$

$$\tag{6}$$

Here, $E = z\nu$, $\alpha_s^{(0)}$ is the bare coupling constant, and $\Phi(u) = 1 - \sin(u)/u$ is the Landau-Pomeranchuk-Migdal interference phase. C_n^{ij} and $\Delta_n^{ij}(x)$ are color and kinematic factors of jet partons interacting with the Glauber gluon for different channels $i \to j$, listed in Table I. Because we focus on the radiative correction to collinear observables for the jet sector, it is sufficient to represent target properties by an effective Glauber gluon density ρ_G (see supplementary material for detailed definition).

In performing **q** and **k** integrals, we introduce $v = \mu_D^2 L/[2x(1-x)E]$ and an integration variable $w = \mathbf{k}^2 L/[2x(1-x)E]$ with $w_{\max} = Q^2 L/(2\nu)$ as bounded by the maximum virtuality of the parton. Even through we require $Q^2, \nu/L \gg \mu_D^2$, we do not have to assume any ordering between Q^2 and ν/L , so w_{\max} can be an order one quantity. Because we focus on modifications in the collinear sector using splitting function obtained in SCET_G, xE, $(1-x)E \gg \mu_D$ from power counting, which allows one to take v = 0 [40]. Doing so results in the final

TABLE I. Color (C_n^{ij}) and kinematic factors (Δ_n^{ij}) in Eq. (6)

| $i \rightarrow j$ | $C_1^{ij}, \ (\Delta_1^{ij})^2$ | $C_2^{ij}, \ (\Delta_2^{ij})^2$ | $C_3^{ij}, \ (\Delta_3^{ij})^2$ |
|-------------------|---------------------------------|---------------------------------|---------------------------------|
| $q \rightarrow q$ | C_A, x^2 | $C_A, 1$ | $2C_F - C_A, (1-x)^2$ |
| $q \rightarrow g$ | $C_A, 1$ | $C_A, \ (1-x)^2$ | $2C_F - C_A, x^2$ |
| $g \rightarrow q$ | $C_A, (1-x)^2$ | C_A, x^2 | $2C_F - C_A, 1$ |
| $g \rightarrow g$ | $C_A, 1$ | C_A, x^2 | $C_A, \ (1-x)^2$ |

expression for the medium-induced branching, where we denote for brevity

$$A(\mu_2^2, E, w_{\max}) = \alpha_s^2(\mu_2^2) L^2 B(w_{\max}) \rho_G / (8E) \,.$$
(7)

 $B(w_{\text{max}})$ and $\chi(w_{\text{max}})$ depend only weakly on Q^2 and ν/L (see supplementary material).

With the results from Eq. (6), taking the medium-

induced $q \to q$ contribution in Eq. (5) as an example,

$$\Delta F_{iq}^{\rm m}(z,\mu_2^2) = \left(F_{iq} + F_{iq} \otimes P_{qq}^{(1)}\right) \otimes \left(M_{qq}^{(0)} + M_{qq}^{(1)}\right) \quad (8)$$

$$\approx \int_0^1 dx \left[P_{qq}^{(1)}(x,\frac{z}{x}\nu,\mu_2^2)F_{iq}\left(\frac{z}{x}\right) - P_{qq}^{(1)}(x,z\nu,\mu_2^2)F_{iq}\left(z\right)\right]$$

$$+ \int_0^1 \frac{dx}{x}F_{iq}(x)M_{qq}^{(1)}\left(\frac{z}{x},\mu_2\right), \quad (9)$$

where we have used the expression for $M_{qq}^{(0)}$, while the NLO renormalization factor $M_{qq}^{(1)}$ will be determined after we identify the relevant poles. To do that, we note that $P_{ij}^{(1)}(x)$ in Eq. (6) contain additional $[x(1-x)]^{-1-2\epsilon}$ divergences as compared to the vacuum splitting functions, which do not cancel among real and virtual corrections in Eq. (9). These extra divergences at x = 0, 1 are consequences of dropping all screening effect in the collinear sector from power counting. If we take the $q \rightarrow q$ channel as an example (details for other channels can be found in the supplementary material), we can isolate these divergences, including the multiplicative $(1-x)^{-1}$ contribution from the vacuum $P_{qq}(x)$, using

the following decomposition for any well-behaved function G(x),

$$\int_{0}^{1} \frac{G(x)}{x^{1+2\epsilon}(1-x)^{2+2\epsilon}} dx = \int_{0}^{1} \frac{\{G(x)\}_{qq}}{x(1-x)^{2}} dx$$
$$-\frac{G(0)}{2\epsilon} + \frac{G'(1)}{2\epsilon} - G(1)\left(\frac{1}{2\epsilon} + 2\right) + \mathcal{O}(\epsilon).$$
(10)

The result has been expanded near $\epsilon = 0$, and the subtracted function is $\{G(x)\}_{qq} = G(x) - (1-x)^2 G(0) - x(2-x)G(1) - x(x-1)G'(1)$. It is straightforward to check that $\{G(x)\}_{qq} / [x^{1+2\epsilon}(1-x)^{2+2\epsilon}]$ is free from divergences at x = 0, 1, while the second line contains all singularities. Note that due to the double pole at x = 1one needs to subtract both G(x) and the derivative G'(x)at x = 1, and such derivative subtractions (a higher-order "plus" prescription) are also used in the study of subleading power corrections in SCET [41].

Following Eq. (10), the $q \rightarrow q$ contribution to the flavor non-singlet sector $\Delta F_{\rm NS} = \Delta F_{iq}^m - \Delta F_{i\bar{q}}^m$ can be decomposed into $1/\epsilon$ poles and $\ln[L\mu_2^2/(\chi z\nu)]$ enhanced terms plus fixed-order contributions

$$\Delta F_{\rm NS}(z,\mu_2^2) = A(\mu_2^2,\nu,w_{\rm max}) \left(\frac{1}{2\epsilon} + \ln\frac{\mu_2^2 L}{\chi z \nu}\right) 2C_F \left(\frac{2C_A + C_F}{z} - 2C_A \frac{d}{dz}\right) F_{\rm NS}(z) + \int_0^1 \frac{dy}{y} F_{\rm NS}(y) M_{qq}^{(1)} \left(\frac{z}{y},\mu_2,z\nu\right) + A(\mu_2^2,\nu,w_{\rm max}) \left\{\int_0^1 \frac{\left\{\sum_n C_n^{qq} [\Delta_n^{qq}(x)]^2 C_F(1+x^2) \left[\frac{x}{z} F_{\rm NS}\left(\frac{z}{x}\right) - \frac{F_{\rm NS}(z)}{z}\right]\right\}_{qq} dx + (4C_A - C_F) C_F \frac{F_{\rm NS}(z)}{z}\right\}, \quad (11)$$

shown in the first and second lines of Eq. (11), respectively. The medium contribution has a natural scale of $\mu_2^2 = \chi z \nu / L$. Divergences due to the $P_{qq}(x)$ factor in $P_{qq}^{(1)}(x)$ have canceled among the real and virtual terms. The remaining poles come from extra $x \to 0, 1$ divergences of the medium-induced emission spectra. We can now define the NLO medium renormalization factor $M_{qq}^{(1)}$ such that it cancels the $1/\epsilon$ pole in the first term. Note that $M_{qq}^{(1)}$ will contain generalized functions and only depend on μ_2 through the coupling constant.

The in-medium RG evolution. With the $1/\epsilon$ pole absorbed in the renormaliation factor, we take a derivative with respect to $\ln \mu_2^2$ on both sides of Eq. (11) and keeping leading terms in α_s and obtain an evolution equation for the μ_2 dependence of $F_{\rm NS}$

$$\frac{\partial F_{\rm NS}(\tau, z)}{\partial \tau} = 2C_F \left(2C_A \frac{\partial}{\partial z} - \frac{2C_A + C_F}{z} \right) F_{\rm NS} \,. \, (12)$$

We have defined a new evolution variable $\tau(z, \mu_2^2) = \frac{\pi B(w_{\max})\rho_G L^2}{2\beta_0 \nu} [\alpha_s(\mu_2^2) - \alpha_s(\frac{\chi z \nu}{L})]$ to take into account the running coupling effect with $\beta_0 = (11 - 2N_f/3)$. One can perform a similar RG analysis on the flavor-singlet sector

and obtain (see supplementary material for details),

$$\frac{\partial F_f}{\partial \tau} = 2C_F \left(2C_A \frac{\partial}{\partial z} - \frac{2C_A + C_F}{z} \right) F_f + C_F \frac{F_g}{z} , (13)$$
$$\frac{\partial F_g}{\partial \tau} = \left(4C_A^2 \frac{\partial}{\partial z} - \frac{2N_f C_F}{z} \right) F_g + 2C_F^2 \sum_f \frac{F_f}{z} , \quad (14)$$

where $F_g \equiv F_{ig}$ is the gluon spectrum and $F_f = F_{iq} + F_{i\bar{q}}$ for f = u, d, s are flavor-singlet quark spectra. Eqs. (12), (13) and (14) are the main results of this letter. Starting with initial condition at $\mu_2^2 = \chi z \nu / L$ and evolving down to $\mu_2^2 = \mu_D^2$ where screening effects become important, the non-singlet Eq. (12) has a very elegant traveling wave solution

$$F_{\rm NS}(\tau, z) = \frac{F_{\rm NS}(0, z + 4C_F C_A \tau)}{\left(1 + 4C_F C_A \tau/z\right)^{1 + \frac{C_F}{2C_A}}}.$$
 (15)

The main effect of the μ_2^2 evolution is to shift the distribution of partons by $\Delta z = -4C_F C_A \tau$. This way, an effective in-medium energy loss can be directly obtained from RG analysis. Neglecting the off-diagonal quarkgluon coupling terms in the flavor-singlet Eqs. (13), (14), similar traveling wave solutions can also be derived. For applications in this letter, we will solve them numerically including the off-diagonal coupling terms.

A widely used phenomenological approach for parton evolution in matter is based upon modified DGLAP (mDGLAP) framework [20, 42–44]. Consider the flavor non-singlet equation

$$\frac{\partial F_{\rm NS}(z)}{\partial \ln \mu^2} = \int_0^1 \mathbf{k}^2 \frac{d[P_{qq}(x, \mathbf{k}^2) + P_{qq}^{(1)}(x, \mathbf{k}^2)]}{dx d\mathbf{k}^2} \\ \times \left[F_{\rm NS}\left(\frac{z}{x}\right) - F_{\rm NS}(z) \right] dx \,, \qquad (16)$$

with $\mu^2 = \mathbf{k}^2/[x(1-x)]$ being the virtuality of the parton, and $d[P_{qq} + P_{qq}^{(1)}]/dxd\mathbf{k}^2$ the double differential splitting function including both vacuum and medium-induced contributions. Despite its apparent different form, we now show that the mDGLAP approach resums the same medium enhancement as the in-medium RG equation to leading logarithmic accuracy. Unlike the RG analysis that uses dimensional regularization, the mDGLAP equation evaluates the full splitting functions numerically and regulates the endpoint divergences of $P_{ij}^{(1)}$ such that $x, 1-x \ge \mu_D^2/\mu^2$ [45]. If we focus on the medium-induced contributions from the $x \approx 1$ region and use a fixed coupling $A(\mu_{\text{fix}}, \nu) = A_0$ for simplicity, the mDGLAP equation becomes

$$\frac{\partial F_{\rm NS}}{\partial \ln \mu^2} = 4C_F C_A A_0 \int_0^{1-\frac{\mu_D^2}{\mu^2}} \frac{4}{\pi} \frac{\Phi(u)}{u} \frac{\frac{x}{z} F_{\rm NS}(\frac{z}{x}) - \frac{F_{\rm NS}(z)}{z}}{(1-x)^2} dx$$
$$\approx \frac{4}{\pi} \frac{\Phi(u)}{u} 4C_F C_A A_0 \left[\frac{\partial F_{\rm NS}}{\partial z} - \frac{F_{\rm NS}}{z}\right] \ln \frac{\mu^2}{\mu_D^2}$$
$$\approx \delta \left(\mu^2 - \frac{2\pi E}{L}\right) 4C_F C_A A_0 \left[\frac{\partial F_{\rm NS}}{\partial z} - \frac{F_{\rm NS}}{z}\right] \ln \frac{\mu^2}{\mu_D^2}, \quad (17)$$

with $u = \mu^2 L/(2E)$. In the second line, we have performed a Taylor expansion of (x/z)F(z/x) near x = 1and omitted subleading terms in $\ln[E/(L\mu_D^2)]$.

The connection to the RG analysis is most easily illustrated by considering a specific case of scale separation $Q_0^2 \ll E/L \ll Q^2$. Because $\frac{4}{\pi} \frac{\Phi(u)}{u}$ peaks at $u = \pi$ and normalizes to unity $\int_0^\infty \frac{4}{\pi} \frac{\Phi(u)}{u} d\ln u = 1$, one can make an impulse approximation, as shown in the third line of Eq. (17). Then, to leading-log accuracy, one can perform vacuum DGLAP evolution above and below $\mu^2 = 2\pi E/L$. However, due to medium effects that sharply peak at $\mu^2 = 2\pi E/L$, the solution below $2\pi E/L$ (F_{NS}^-) and the solution above (F_{NS}^+) are related by

$$F_{NS}^{+}(z) = \frac{F_{NS}^{-}(z + 4C_F C_A \tau_{\text{fix}})}{1 + 4C_F C_A \tau_{\text{fix}}/z},$$
(18)

with $\tau_{\text{fix}} = A_0 \ln \frac{2\pi E}{\mu_D^2 L}$. This is the same traveling wave solution as in Eq. (15), but with fixed coupling and neglecting contributions from the x = 0 endpoint [46]. We, therefore, conclude that the mDGLAP approach with the



FIG. 1. Top panel: medium modifications to the π^+ fragmentation function compare to HERMES data, performed for the average $\nu = 12$ GeV, $Q^2 = 2.25$ GeV². Bottom panel: predictions for the modified pion fragmentation function at EIC with Pb nucleus for three (x_B, Q^2) combinations.

chioce $\mu^2 = \mathbf{k}^2/[x(1-x)]$ resums the same medium enhanced branchings as the RG equation derived in this letter. Of course, with a large separation of scales, it becomes computationally intensive to evaluate the RHS of the mDGLAP equation with an explicit cut-off and the analytic approach that we formulated here is not only more illuminating, but also easier to implement.

We demonstrate the new method by studying nuclear effects on pion fragmentation in SIDIS, with the cross section given in Eq. (2). We implement the fully-coupled RG evolution Eqs. (12), (13), (14), and the fixed order terms in Eq. (11). The nuclear modifications is defined as the ratio of inclusive-normalized cross sections between electron-nucleus (eA) and electron-deuterium (edfor HERMES) or electron-proton (ep for EIC) collisions

$$R_A(x_B, Q^2, z_h) = \frac{\frac{d\sigma_{eA \to h}}{dx_B dQ^2 dz_h} / \frac{d\sigma_{eA}}{dx_B dQ^2}}{\frac{d\sigma_{ed,ep \to h}}{dx_B dQ^2 dz_h} / \frac{d\sigma_{ed,ep}}{dx_B dQ^2}}.$$
 (19)

For parton distribution and fragmentation functions we use the nNNPDF30nlo [47] and NNFF10lo parametrizations [48] and perform the calculation for the averaged HERMES kinematics $Q^2 = 2.25 \text{ GeV}^2$ and $\nu =$ 12 GeV [49]. To numerically solve Eqs. (12) and (14), we smear the singular hard parton energy spectrum by a Gaussian with width parameter $\sigma_z = 0.05$ and the evolution from $\mu_1 = Q$ to $Q_0 = 1$ GeV is performed using standard vacuum DGLAP. In turn, the in-medium RG evolves μ_2^2 from $\chi(w_{\text{max}})z\nu/L$ to μ_D^2 . We take $\Lambda_{\text{QCD}} = 0.16$ GeV, the inverse range of the interaction $\mu_D^2 = 0.12$ GeV² and the central value of the effective medium density parameter $\rho_G = 0.4$ fm⁻³. These values yield a quark transport parameter $\hat{q}_F \approx 0.053$ GeV^2/fm at $\nu = 12$ GeV (see supplemental material), consistent with existing mDGLAP applications [50, 51]. An average over the geometry of the nucleus of radius $r_A = 1.2A^{1/3}$ fm is also performed.

The resulting nuclear modification factor $R_A(z_h)$ is compared to the HERMES data for ²⁰Ne and ¹³¹Xe targets [49] in the top row of Fig. 1. Qualitatively, inmedium evolution shifts hadron spectra towards lower z_h . Results including only RG contributions (red dashed lines) give a good description of R_A from small to intermediate z, but lead to a suppression that is too strong at large z_h . We remark that the region very close to $z_h = 1$ is dominated by soft emissions, where one should consider soft power counting and threshold type resummation in both vacuum [52, 53] and in-medium calculations. Therefore, we have exclude this region from our comparison. Blue solid lines include the fixed order (FO) contribution from Eq. (10) in the initial condition of the RG evolution, and the bands correspond to the density variation in the range $(\rho_G/1.5, 1.5\rho_G)$. The FO correction improves the description of HERMES data at large z, but remains subleading to the RG evolution effect. The nuclear size dependence of R_A for Ne, Kr, and Xe nuclei is naturally explained with the same set of transport parameters.

Using the same in-medium transport parameters, we present projections (lower panel of Fig. 1) for modified pion fragmentation functions at the future electron-ion collider (EIC) for ePb reactions at fixed $Q^2 = 20 \text{ GeV}^2$ and various Bjorken x_B values. We find that for $x_B >$ 0.3, where partons are less energetic in the nuclear rest frame, modifications become very large, consistent with existing predictions for heavy flavor and jets [50, 51]. Summary. In the limit $Q/Q_0, E/(\mu_D^2 L), \lambda_q \mu_D \gg 1$ and to first order in the opacity of QCD matter we performed a renormalization group (RG) analysis of medium effects for the SIDIS process on a nuclear target. We derived a set of in-medium RG equations that resum the leading $\ln[E/(\mu_D^2 L)]$ terms from multiple medium-induced emissions and identified the corresponding fixed order corrections. We further showed that such resummation is also contained in the modified DGLAP equations, which differ in the way of regulating the endpoint divergences of medium-induced emission spectra. Importantly, the new RG evolution in matter approach provides analytic insight into the salient features of parton showers responsible for the modification of hadron production in eAthat are not possible with numerical methods alone. It is a more efficient and systematically improvable way of treating the logarithmic enhancements in matter as compared to solving mDGLAP.

We applied the new method to study the cold nuclear matter (CNM) effects on pion fragmentation and found that it gives a good description of the HERMES SIDIS data. Predictions for the future EIC were also presented, where improved theoretical precision is especially important [15]. The semi-analytic framework derived here can be generalized to initial-state CNM effects, such as the ones observed in Drell-Yan production in proton-nucleus collisions, and to heavy ion collisions. This work further benefits future QCD studies by providing guidance on incorporating medium effects in Monte-Carlo event generators for the EIC, the Relativistic Heavy Ion Collider and the Large Hadron Collider.

Acknowledgments.— The authors would like to thank Duff Neill for helpful discussion. This work is supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics through Contract No. 89233218CNA000001 and by the Laboratory Directed Research and Development Program at LANL.

* weiyaoke@lanl.gov

- [†] ivitev@lanl.gov
- M. Newman and D. Watts, Physics Letters A 263, 341 (1999).
- [2] V. Yakhot and S. A. Orszag, Journal of Scientific Computing 1, 3 (1986).
- [3] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, JHEP 04, 159 (2014), arXiv:1312.2014 [hep-ph].
- [4] J. Jalilian-Marian, A. Kovner, and H. Weigert, Phys. Rev. D 59, 014015 (1998), arXiv:hep-ph/9709432.
- [5] S. K. Bogner, R. J. Furnstahl, and R. J. Perry, Phys. Rev. C 75, 061001 (2007), arXiv:nucl-th/0611045.
- [6] G. 't Hooft and M. J. G. Veltman, Nucl. Phys. B 44, 189 (1972).
- [7] K. G. Wilson and J. B. Kogut, Phys. Rept. 12, 75 (1974).
- [8] C. W. Bauer, S. Fleming, D. Pirjol, and I. W. Stewart, Phys. Rev. D 63, 114020 (2001), arXiv:hep-ph/0011336.
- [9] M. Beneke, A. P. Chapovsky, M. Diehl, and T. Feldmann, Nucl. Phys. B 643, 431 (2002), arXiv:hepph/0206152.
- [10] A. Idilbi and A. Majumder, Phys. Rev. D 80, 054022 (2009), arXiv:0808.1087 [hep-ph].
- [11] F. D'Eramo, H. Liu, and K. Rajagopal, Phys. Rev. D 84, 065015 (2011), arXiv:1006.1367 [hep-ph].
- [12] G. Ovanesyan and I. Vitev, Phys. Lett. B 706, 371 (2012), arXiv:1109.5619 [hep-ph].
- [13] G. Ovanesyan and I. Vitev, JHEP 06, 080 (2011), arXiv:1103.1074 [hep-ph].
- [14] Z.-B. Kang, F. Ringer, and I. Vitev, JHEP 03, 146 (2017), arXiv:1610.02043 [hep-ph].
- [15] R. Abdul Khalek *et al.*, Nucl. Phys. A **1026**, 122447 (2022), arXiv:2103.05419 [physics.ins-det].
- [16] G.-Y. Qin and X.-N. Wang, International Journal of Modern Physics E 24, 1530014 (2015), https://doi.org/10.1142/S0218301315300143.
- [17] I. Vitev and M. Gyulassy, Phys. Rev. Lett. 89, 252301 (2002), arXiv:hep-ph/0209161.
- [18] M. Gyulassy, I. Vitev, X.-N. Wang, and B.-W. Zhang, , 123 (2004), arXiv:nucl-th/0302077.
- [19] G.-Y. Qin, J. Ruppert, C. Gale, S. Jeon, G. D. Moore, and M. G. Mustafa, Phys. Rev. Lett. **100**, 072301 (2008).
- [20] W.-T. Deng and X.-N. Wang, Phys. Rev. C 81, 024902 (2010), arXiv:0910.3403 [hep-ph].
- [21] J.-P. Blaizot, E. Iancu, and Y. Mehtar-Tani, Phys. Rev. Lett. **111**, 052001 (2013), arXiv:1301.6102 [hep-ph].

- [22] Z.-B. Kang, R. Lashof-Regas, G. Ovanesyan, P. Saad, and I. Vitev, Phys. Rev. Lett. **114**, 092002 (2015), arXiv:1405.2612 [hep-ph].
- [23] Y.-T. Chien and I. Vitev, JHEP 05, 023 (2016), arXiv:1509.07257 [hep-ph].
- [24] J. Noronha-Hostler, B. Betz, J. Noronha, and M. Gyulassy, Phys. Rev. Lett. **116**, 252301 (2016), arXiv:1602.03788 [nucl-th].
- [25] N.-B. Chang and G.-Y. Qin, Phys. Rev. C 94, 024902 (2016), arXiv:1603.01920 [hep-ph].
- [26] D. Zigic, I. Salom, J. Auvinen, M. Djordjevic, and M. Djordjevic, Phys. Lett. B **791**, 236 (2019), arXiv:1805.04786 [nucl-th].
- [27] W. Chen, S. Cao, T. Luo, L.-G. Pang, and X.-N. Wang, Phys. Lett. B 810, 135783 (2020), arXiv:2005.09678 [hepph].
- [28] S. Schlichting and I. Soudi, (2020), arXiv:2008.04928 [hep-ph].
- [29] G. I. Barenblatt, Scaling, Self-similarity, and Intermediate Asymptotics: Dimensional Analysis and Intermediate Asymptotics, Cambridge Texts in Applied Mathematics (Cambridge University Press, 1996).
- [30] M. D. Sievert, I. Vitev, and B. Yoon, Phys. Lett. B 795, 502 (2019), arXiv:1903.06170 [hep-ph].
- [31] B. Schenke, C. Gale, and S. Jeon, Phys. Rev. C 80, 054913 (2009), arXiv:0909.2037 [hep-ph].
- [32] Y. He, T. Luo, X.-N. Wang, and Y. Zhu, Phys. Rev. C 91, 054908 (2015), [Erratum: Phys.Rev.C 97, 019902 (2018)], arXiv:1503.03313 [nucl-th].
- [33] J.-P. Blaizot, F. Dominguez, E. Iancu, and Y. Mehtar-Tani, JHEP 06, 075 (2014), arXiv:1311.5823 [hep-ph].
- [34] D. de Florian, M. Stratmann, and W. Vogelsang, Phys. Rev. D 57, 5811 (1998), arXiv:hep-ph/9711387.
- [35] $d_{j/h}$ will obey the same evolution, but with the opposite sign. We choose to keep fragmentation at the relevant non-perturbative scale and evolve the final-state distribution of partons down to them.
- [36] W. Furmanski and R. Petronzio, Z. Phys. C 11, 293 (1982).

- [37] G. Altarelli and G. Parisi, Nucl. Phys. B 126, 298 (1977).
- [38] L. N. Lipatov, Yad. Fiz. **20**, 181 (1974).
- [39] G. Ovanesyan, F. Ringer, and I. Vitev, Phys. Lett. B 760, 706 (2016), arXiv:1512.00006 [hep-ph].
- [40] Under DR, the integration of the dimensionless variables u and w receives vanishing contributions from the soft region, and therefore are treated as order-unity quantities.
- [41] M. A. Ebert, I. Moult, I. W. Stewart, F. J. Tackmann, G. Vita, and H. X. Zhu, JHEP 04, 123 (2019), arXiv:1812.08189 [hep-ph].
- [42] A. Majumder, Phys. Rev. C 88, 014909 (2013), arXiv:1301.5323 [nucl-th].
- [43] N.-B. Chang, W.-T. Deng, and X.-N. Wang, Phys. Rev. C 89, 034911 (2014), arXiv:1401.5109 [nucl-th].
- [44] Y.-T. Chien, A. Emerman, Z.-B. Kang, G. Ovanesyan, and I. Vitev, Phys. Rev. D 93, 074030 (2016), arXiv:1509.02936 [hep-ph].
- [45] W. Ke and I. Vitev, (2022), arXiv:2204.00634 [hep-ph].
- [46] One can drop contributions from the x = 0 endpoint in Eq. 12 by neglecting $C_F/(2C_A)$.
- [47] R. A. Khalek, R. Gauld, T. Giani, E. R. Nocera, T. R. Rabemananjara, and J. Rojo, "nNNPDF3.0: Evidence for a modified partonic structure in heavy nuclei," (2022), arXiv:2201.12363 [hep-ph].
- [48] V. Bertone, S. Carrazza, N. P. Hartland, E. R. Nocera, and J. Rojo (NNPDF), Eur. Phys. J. C 77, 516 (2017), arXiv:1706.07049 [hep-ph].
- [49] A. Airapetian *et al.* (HERMES), Nucl. Phys. B **780**, 1 (2007), arXiv:0704.3270 [hep-ex].
- [50] H. T. Li and I. Vitev, Phys. Rev. Lett. **126**, 252001 (2021), arXiv:2010.05912 [hep-ph].
- [51] H. T. Li, Z. L. Liu, and I. Vitev, Phys. Lett. B 816, 136261 (2021), arXiv:2007.10994 [hep-ph].
- [52] S. Catani and L. Trentadue, Nucl. Phys. B 327, 323 (1989).
- [53] D. P. Anderle, F. Ringer, and W. Vogelsang, Phys. Rev. D 87, 034014 (2013), arXiv:1212.2099 [hep-ph].
- [54] I. Vitev, Phys. Rev. C 75, 064906 (2007), arXiv:hepph/0703002.

SUPPLEMENTARY MATERIAL

In-medium scattering cross section and jet transport parameter estimate

The elastic cross section between jet and target partons in color representations R and T, respectively, is [13]

$$\frac{d\sigma_{TR}}{d^2\mathbf{q}} = \frac{1}{(2\pi)^2} \frac{1}{d_A} \frac{4\pi\alpha_s C_T \times 4\pi\alpha_s C_R}{(\mathbf{q}^2 + \mu_D^2)^2},$$
(20)

with $d_A = N_c^2 - 1$. The collision rate after summing over the medium color sources of representation T with density ρ_T then reads

$$\sum_{T} \rho_T \frac{d\sigma_{TR}}{d^2 \mathbf{q}} = \sum_{T} \frac{1}{(2\pi)^2} \frac{\rho_T}{d_A} \frac{4\pi \alpha_s C_T \times 4\pi \alpha_s C_R}{(\mathbf{q}^2 + \mu_D^2)^2} = \frac{\alpha_s C_R}{\pi} \frac{1}{(\mathbf{q}^2 + \mu_D^2)^2} \rho_G \,, \quad \rho_G \equiv \sum_{T} \rho_T \frac{4\pi \alpha_s^{\text{med}} C_T}{d_A} \,. \tag{21}$$

In other words, we have chosen to put kinematic factors, the square color charges and coupling to the medium in the effective medium gluon density ρ_G . With $\mu_D^2 = 0.12 \text{ GeV}^2$, $\rho_G = 0.4 \text{ fm}^{-3}$, the quark jet transport parameter is

$$\hat{q}_F = \int_0^{\mathbf{q}_{\max}^2 = \nu \mu_D/2} d^2 \mathbf{q} \, \mathbf{q}^2 \frac{\alpha_s C_F}{\pi} \frac{1}{(\mathbf{q}^2 + \mu_D^2)^2} \rho_G \approx 0.053 \, \text{GeV}^2/\text{fm}$$
(22)

for $\nu = Q^2/(2x_B M_p) = 12$ GeV in the nuclear rest frame. Here, the ultraviolet cut off of the \mathbf{q}^2 integration is chosen to be $\nu\mu_D/2$, as in Ref. [54]. The running coupling is cut-off when α_s reaches $2\pi/\beta_0$. The value of \hat{q} is further consistent with the analysis of [50, 51].

Full splitting functions in matter

The splitting functions in nuclear matter induced by final-state interactions are taken from Refs. [13, 14] (in d = 4 dimension). After performing the path length integration in a medium of uniform density and size L, the splitting functions become

$$P_{ij}^{(1)}(x) = \frac{\alpha_s}{2\pi^2} P_{ij}(x) L \int d^2 \mathbf{k} \sum_T \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{\rho_T}{d_A} \frac{4\pi \alpha_s C_T \times 4\pi \alpha_s}{(\mathbf{q}^2 + \mu_D^2)^2} W_{ij}(x, \mathbf{k}, \mathbf{q}, E/L)$$

$$\equiv \frac{\alpha_s}{2\pi^2} P_{ij}(x) L \int d^2 \mathbf{k} \int \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{\rho_G \times 4\pi \alpha_s}{(\mathbf{q}^2 + \mu_D^2)^2} W_{ij}(x, \mathbf{k}, \mathbf{q}, E/L) .$$
(23)

The continuous parts of the vacuum splitting functions in $d = 4 - 2\epsilon$ dimension, arising from real emissions, are

$$P_{qq}(x) = C_F \left[\frac{1+x^2}{1-x} - \epsilon(1-x) \right], \ P_{gq}(x) = T_R[x^2 + (1-x)^2 - 2\epsilon x(1-x)],$$
(24)

$$P_{qg}(x) = P_{qq}(1-x), \ P_{gg}(x) = C_A \frac{1+x^4+(1-x)^4}{x(1-x)}.$$
(25)

The diagonal terms also receive virtual corrections, which we determined from flavor and momentum sum rules [44].

For
$$P_{ij}^{(1)}(x)$$
, we define $\mathbf{A}_{\perp} = \mathbf{k}$, $\mathbf{B}_{\perp} = \mathbf{k} + (1-x)\mathbf{q}$, $\mathbf{C}_{\perp} = \mathbf{k} - x\mathbf{q}$, $\mathbf{D}_{\perp} = \mathbf{k} + \mathbf{q}$, and interference phase factors

$$\Phi_{A} = \Phi\left[\frac{\mathbf{A}_{\perp}^{2}L}{2x(1-x)E}\right], \ \Phi_{B} = \Phi\left[\frac{\mathbf{B}_{\perp}^{2}L}{2x(1-x)E}\right], \ \Phi_{C} = \Phi\left[\frac{\mathbf{C}_{\perp}^{2}L}{2x(1-x)E}\right],$$
$$\Phi_{AD} = \Phi\left[\frac{(\mathbf{A}_{\perp}^{2} - \mathbf{D}_{\perp}^{2})L}{2x(1-x)E}\right], \ \Phi_{CB} = \Phi\left[\frac{(\mathbf{C}_{\perp}^{2} - \mathbf{B}_{\perp}^{2})L}{2x(1-x)E}\right].$$
(26)

Then, W_{ij} , including the relevant quadratic Casimirs from the Glauber gluon-hard parton system interactions, are

$$W_{qq} = C_A \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_B + C_A \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \right) \Phi_B + (2C_F - C_A) \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \left(\frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_C + C_A \Delta W,$$
(27)

$$W_{gg} = C_A \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_B + C_A \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \right) \Phi_B + C_A \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \left(\frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_C + C_A \Delta W, \tag{28}$$

$$W_{gq} = C_A \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_B + (2C_F - C_A) \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} \left(\frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^2} - \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \right) \Phi_B + C_A \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} \left(\frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^2} - \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^2} \right) \Phi_C + (2C_F - C_A) \Delta W,$$
(29)

and $W_{qg} = W_{qq}(x \to 1 - x)$. For each channel, the first three terms can be cast into the form of the first line of Eq. (6) by shifting integration variables such that arguments of $\Phi(\cdot)$ become $\mathbf{k}^2 L/[2x(1-x)E]$. The remaining piece ΔW ,

$$\Delta W = \left[\frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^{2}} \left(\frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^{2}} - \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^{2}}\right) \Phi_{C} + \frac{\mathbf{B}_{\perp}}{\mathbf{B}_{\perp}^{2}} \cdot \frac{\mathbf{C}_{\perp}}{\mathbf{C}_{\perp}^{2}} \Phi_{CB}\right] - \left[\frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^{2}} \left(\frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^{2}} - \frac{\mathbf{D}_{\perp}}{\mathbf{D}_{\perp}^{2}}\right) \Phi_{A} + \frac{\mathbf{A}_{\perp}}{\mathbf{A}_{\perp}^{2}} \cdot \frac{\mathbf{D}_{\perp}}{\mathbf{D}_{\perp}^{2}} \Phi_{AD}\right], \quad (30)$$

is written as the difference of two terms. Note that the second term can be obtained from the first one by shifting $\mathbf{k} \to \mathbf{k} + x\mathbf{q}$, causing $\mathbf{C}_{\perp} \to \mathbf{A}_{\perp}$ and $\mathbf{B}_{\perp} \to \mathbf{D}_{\perp}$. Therefore, $\Delta W = 0$ under dimensional regularized integration of \mathbf{k} and \mathbf{q} . If one uses an explicit ultraviolet cut-off $\Lambda_{\rm UV}$, the integration of ΔW over \mathbf{q}, \mathbf{k} (or in general, any differences caused by a shift of $\mathbf{k} + \Delta \cdot \mathbf{q}$ of Eqs. (27),(28), and (29)) are further suppressed by $\Lambda_{\rm UV}^{-2}$ and do not contribute to the medium-induced logarithmic enhancement.

We can account for the virtuality of the collision Eq. (6) by introducing the variable $w_{\text{max}} = Q^2 L/(2\nu)$, which appears in the functions B and χ defined as

$$B(w_{\max}) = \frac{4}{\pi} \int_0^{w_{\max}} \Phi(x) \frac{dx}{x^2}, \quad \chi(w_{\max}) = 2 \exp\left\{\frac{1}{B(w_{\max})} \frac{4}{\pi} \int_0^{w_{\max}} \Phi(x) \ln(x) \frac{dx}{x^2} + \gamma_E\left(\frac{1}{B(w_{\max})} - 1\right)\right\}.$$
 (31)

For the SIDIS process at moderate x_B , $w_{\text{max}} \equiv x_B M_p L \approx 6.0 x_B A^{1/3}$ is of order few. For fragmentation at mid rapidity in hadronic collisions, $Q^2 L/(2E) \sim EL \rightarrow \infty$, and $B(\infty) = 1$, $\chi(\infty) = 2e^{3/2 - \gamma_E} \approx 5.0$.

The flavor singlet sector

Analogously to the treatment of the flavor non-singlet sector, we provide the details for the subtraction of divergences and renormalization of the flavor singlet sector. To isolate the extra $[x(1-x)]^{-1-2\epsilon}$ poles, we define the following decomposition for any well-behaved function G(x). For singularities associate to $P_{qg}^{(1)}(x)$,

$$\int_{0}^{1} \frac{G(x)}{x^{2+2\epsilon}(1-x)^{1+2\epsilon}} dx = \int_{0}^{1} \frac{\{G(x)\}_{qg}}{x^{2}(1-x)} dx - \frac{G(0)}{2\epsilon} - \frac{G'(0)}{2\epsilon} - G(1)\left(\frac{1}{2\epsilon} + 2\right) + \mathcal{O}(\epsilon),$$
(32)

with $\{G(x)\}_{qg} = G(x) - x^2 G(1) - (1 - x^2) G(0) - x(1 - x) G'(0)$. For singularities associate to $P_{gg}^{(1)}(x)$,

$$\int_{0}^{1} \frac{G(x)}{x^{2+2\epsilon}(1-x)^{2+2\epsilon}} dx = \int_{0}^{1} \frac{\{G(x)\}_{gg}}{x^{2}(1-x)^{2}} dx - G(0) \left[\frac{1}{\epsilon} + 2\right] - \frac{G'(0)}{2\epsilon} - G(1) \left[\frac{1}{\epsilon} + 2\right] + \frac{G'(1)}{2\epsilon} + \mathcal{O}(\epsilon), \quad (33)$$

with $\{G(x)\}_{gg} = G(x) - (1-x)^2 \left[(1+2x)G(0) + xG'(0)\right] - x^2 \left[(3-2x)G(1) + (x-1)G'(1)\right]$. Finally, for $P_{gq}^{(1)}(x)$

$$\int_{0}^{1} \frac{G(x)}{x^{1+2\epsilon}(1-x)^{1+2\epsilon}} dx = \int_{0}^{1} \frac{\{G(x)\}_{gq}}{x(1-x)} dx - \frac{G(0)}{2\epsilon} - \frac{G(1)}{2\epsilon} + \mathcal{O}(\epsilon) , \qquad (34)$$

with $\{G(x)\}_{gq} = G(x) - xG(1) - (1 - x)G(0)$. One can directly and explicitly check that the endpoint divergences are removed from the integration. With this procedure, the medium-induced NLO contributions from these channels can be decomposed into log-enhanced and fixed order contributions,

$$\frac{\Delta F_g}{A(\mu_2^2, E, w_{\max})} = \left(\frac{1}{2\epsilon} + \mathcal{L}\right) \left[-4C_A^2 \frac{dF_g(z)}{dz} + 2C_F N_f \frac{F_g(z)}{z} - 2C_F^2 \sum_f \frac{F_f(z)}{z} \right] \\ + \sum_f \int_0^1 dx \frac{\left\{\sum_n C_n^{qg} \left(\Delta_n^{qg}\right)^2 C_F [1 + (1 - x)^2] \frac{x}{z} F_f(\frac{z}{x})\right\}_{qg}}{x^2 (1 - x)} - N_f \frac{F_g(z)}{z} \int_0^1 dx \frac{\left\{\sum_i C_n^{gg} \left(\Delta_n^{gg}\right)^2 T_R [x^2 + (1 - x)^2]\right\}_{gq}}{x (1 - x)} \\ + \int_0^1 dx \frac{\left\{\sum_i C_n^{gg} \left(\Delta_n^{gg}\right)^2 C_A [1 + x^4 + (1 - x)^4] \left[\frac{x}{z} F_g(\frac{z}{x}) - x \frac{F_g(z)}{z}\right]\right\}_{gg}}{x^2 (1 - x)^2} + 14C_A^2 \frac{F_g(z)}{z} - 3C_F^2 \sum_f \frac{F_f(z)}{z}, \quad (35)$$

$$\frac{\Delta F_f}{A(\mu_2^2, E, w_{\max})} = \left(\frac{1}{2\epsilon} + \mathcal{L}\right) \left[-4C_F C_A \frac{dF_f(z)}{dz} + 2C_F (2C_A + C_F) \frac{F_f(z)}{z} - C_F \frac{F_g(z)}{z} \right] + \int_0^1 \frac{\left\{ \sum_n C_n \Delta_n^2(x) C_F (1+x^2) \left(\frac{x}{z} F_f\left(\frac{z}{x}\right) - \frac{F_f(z)}{z}\right) \right\}_{qq}}{x(1-x)^2} dx + (4C_A - C_F) C_F \frac{F_f(z)}{z} + \int_0^1 dx \frac{\left\{ \sum_i C_n^{gq} \left(\Delta_n^{gq}\right)^2 T_R [x^2 + (1-x)^2] \frac{x}{z} F_g\left(\frac{z}{x}\right) \right\}_{gq}}{x(1-x)} , \qquad (36)$$

where $\mathcal{L} \equiv \ln \frac{\mu_2^2}{\chi E/L}$ and $\{\cdots\}_{ij}$ stands for the subtraction for the corresponding channel $i \to j$. The $1/\epsilon$ poles are subtracted by the corresponding contribution from $M_{ij}^{(1)}$. Then, taking derivatives with respect to τ gives the in-medium RG Eqs. (13) and (14) for the flavor singlet sector.