Generalized Uncertainty Principle for Entangled States of Two Identical Particles

K. C. Lemos Filho^{*1}, B. B. Dilem^{†2}, J. C. Fabris^{‡1,3}, and J. A. Nogueira^{§1}

 1 Universidade Federal do Espírito Santo – Ufes, Vitória, Espírito Santo, 29.075-910, Brasil

 2 Instituto Federal do Espírito Santo – Ifes, Alegre, Espírito Santo, 29.520-000, Brasil

³National Research Nuclear University MEPhI, Kashirskoe sh. 31, Moscow 115409, Russia

Abstract

In this work we determine the consequences of the quantum entanglement of a system of two identical particles when the generalized uncertainty principle (GUP) is considered. GUP is usually associated with the existence of a minimal length. We focus on the main formulations of the GUP and then we determine the minimal uncertainties in position induced by those modified GUP's. Our results point out that the minimal uncertainty is reduced by half of its usual value independently of the GUP employed. This implies that the minimal length is also reduced by half. On the other hand, it is generally expected that the minimal length must not depend on physical system. We overcome this apparent paradox by realizing that the entangled system is composed by two particles so that an effective parameter related to the minimal length must be employed.

PACS numbers: 04.60.-m, 03.65.Ud, 03.65.Ta

Keywords: Minimal length; generalized uncertainty principle; quantum entanglement.

^{*}kim.vasco@gmail.com

[†]bernardob@ifes.edu.br

[‡]julio.fabris@cosmo-ufes.org

[§]jose.nogueira@ufes.br

1 Introduction

Amongst the (many) new concepts introduced by the quantum mechanics, the quantum entanglement [1, 2] is one that, probably, more contradicts our common sense. Although, in the beginning, the quantum entanglement were only associated to theoretical aspects of quantum mechanics, specially those related to the non-locality or the complementarity (hidden variables) [3], nowadays it is a key component of the applications and experiments on quantum information, quantum computation and quantum teleportation [4, 5].

The uncertainty principle is one of the fundamental cornerstones of the quantum mechanics. Nevertheless, it is a principle: its correct form can not be proven. That opens the way to the possibility that its canonical form, described by the Heisenberg's uncertainty principle (HUP), can be generalized¹. An example of the possible generalizations of HUP, whose origin can be traced back to quantum gravity, is given by introducing a non-zero minimal uncertainty in the measurement of position. That non-zero minimal uncertainty in position is, then, understood as a minimal-length scale, below which the necessary amount of energy to probe the position of a particle is so hight that it disturbs the space-time so that the concept of a length measurement loses its meaning. Hence, theories searching to describe a quantum approach for gravity lead generally to the existence of a minimal length. In fact, a minimal length actually appears in almost all proposed theories of the quantum gravity. For this reason theories formulated in a minimal-length scenario are considered to be effective theories of quantum gravity [17, 18, 19, 20].

In 1999, Yoon-Ho Kim and Yanhua Shih conducted an experiment whose results apparently suggested a violation of the HUP [21]. However, G. Rigolin pointed out that in fact there is no violation of the HUP, because the HUP is derived for particles non-correlated (non-entangled). In the Kim and Shih's experiment, the photons of the pair are correlated (entangled) when one of the physical slits is replaced by a virtual slit²: the canonical HUP is no longer applicable since the quantum entanglement modifies the canonical HUP [22, 23].

An immediate question arising from the previous considerations is how the quantum entanglement modifies a generalized uncertainty principle (GUP), in others words, which is the effect of the quantum entanglement in the minimal-length scale. The answer for that question is important in order to know the role of the quantum entanglement at the minimal-length scale (maybe appearing in the Planck scale) or in the early Universe. Unfortunately, this issue has been little considered in the literature. In [24], G. Blado, F. Herrera and J. Erwin have studied the inseparability conditions with the most usual GUP correction, whereas D. Park has used a coupled harmonic oscillator in order to find the effects of the quantum entanglement with a linear GUP in [25].

The purpose of this work is to answer that question considering the main proposals of generalization for the HUP which take into account the existence of a minimal length in

¹For more information on generalization of the Heisenberg's uncertainty principle see [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16].

²The interaction of the photon with a physical slit destroys the correlation between the photons of the pair.

nature. With this goal in mind, we will analyze the modifications in the HUP arising from the quantum entanglement of two identical particles determining the minimal uncertainty associated to them.

The outline of this paper is as follows. In Section 2 we obtain an expression of the uncertainty principle for entangled states which is independent of the chosen GUP. In Section 3 we find the modified uncertainty principle for a pair of identical particles regarding the main proposals of GUP's: Kempf, Mangano and Mann GUP (KMM-GUP), Ali, Das and Vagenas GUP (ADV-GUP), Pedram GUP and exponential all orders GUP. In Section 4 we estimate an upper bound for the minimal-length value. We present our conclusions in the Section 5.

2 Uncertainty principle for entangled states

The Hilbert space of the state vectors \mathcal{E} of a system of N particles is given by the tensor product of the Hilbert spaces of the state vectors \mathcal{E}_i of each particle [26, 27],

$$\mathcal{E} = \mathcal{E}_1 \otimes \cdots \otimes \mathcal{E}_N. \tag{1}$$

The position and momentum linear operators of the i-th particle which act on the state vectors $|\psi\rangle \in \mathcal{E}$ are the extensions \tilde{Q}_i and \tilde{P}_i defined as

$$\tilde{Q}_i = \mathbb{I}_1 \otimes \cdots \otimes \hat{x}_i \otimes \cdots \otimes \mathbb{I}_N, \tag{2}$$

$$\tilde{P}_i = \mathbb{I}_1 \otimes \cdots \otimes \hat{p}_i \otimes \cdots \otimes \mathbb{I}_N, \tag{3}$$

where \mathbb{I}_i is the identity operator in \mathcal{E}_i and \hat{x}_i and \hat{p}_i are the position and the momentum operators of the i-th particle acting on the state vectors $|\psi_i\rangle \in \mathcal{E}_i$.

The extensions \tilde{Q}_i and \tilde{P}_i do not satisfy the canonical uncertainty principle (HUP), because \tilde{Q}_i and \tilde{P}_i are not physical observables [22, 23, 26, 27]. Physical observables are operators which commute with every permutation operators of the particles system. Hence, the operators \tilde{Q} and \tilde{P} , defined as

$$\tilde{Q} := \sum_{i=1}^{N} \tilde{Q}_i \tag{4}$$

and

$$\tilde{P} := \sum_{i=1}^{N} \tilde{P}_i,\tag{5}$$

are physical observables and they satisfy the relation

$$\left(\Delta Q\right)^{2} \left(\Delta P\right)^{2} \geq \frac{1}{4} \left| \left\langle \left[\tilde{Q}, \tilde{P}\right] \right\rangle \right|^{2}.$$
(6)

The relation (6) is general. It does not depend whether the system of particle is entangled or not.

As it was showed by G. Rigolin [22, 23], if the state of the particles system is entangled then the operators \tilde{Q}_i and \tilde{P}_i do not satisfy the canonical Heisenberg uncertainty principle (HUP) - as previously stated.

We briefly review the Rigolin's result for a two particles system. From the definitions of ΔQ and ΔP we have³

$$(\Delta_{\psi}Q)^{2} = \left\langle \psi | \tilde{Q}^{2} | \psi \right\rangle - \left\langle \psi | \tilde{Q} | \psi \right\rangle^{2}.$$
(7)

From now on we omit the subscript ψ for the sake of simplicity, whenever this does not cause any confusion. Thus,

$$\left(\Delta Q\right)^{2} = \left(\Delta Q_{1}\right)^{2} + \left(\Delta Q_{2}\right)^{2} + 2\left(\left\langle \tilde{Q}_{1}\tilde{Q}_{2}\right\rangle - \left\langle \tilde{Q}_{1}\right\rangle \left\langle \tilde{Q}_{2}\right\rangle\right).$$
(8)

In the same way,

$$(\Delta P)^{2} = (\Delta P_{1})^{2} + (\Delta P_{2})^{2} + 2\left(\left\langle \tilde{P}_{1}\tilde{P}_{2}\right\rangle - \left\langle \tilde{P}_{1}\right\rangle \left\langle \tilde{P}_{2}\right\rangle\right).$$

$$(9)$$

Using the results (8) and (9) into Eq. (6) we obtain

$$\left[\left(\Delta Q_1 \right)^2 + \left(\Delta Q_2 \right)^2 + 2 \left(\left\langle \tilde{Q}_1 \tilde{Q}_2 \right\rangle - \left\langle \tilde{Q}_1 \right\rangle \left\langle \tilde{Q}_2 \right\rangle \right) \right] \times \left[\left(\Delta P_1 \right)^2 + \left(\Delta P_2 \right)^2 + 2 \left(\left\langle \tilde{P}_1 \tilde{P}_2 \right\rangle - \left\langle \tilde{P}_1 \right\rangle \left\langle \tilde{P}_2 \right\rangle \right) \right] \ge \frac{1}{4} \left| \langle [\tilde{Q}, \tilde{P}] \rangle \right|^2.$$
(10)

We now use the functions

$$C_Q(1,2) := \left\langle \tilde{Q}_1 \tilde{Q}_2 \right\rangle - \left\langle \tilde{Q}_1 \right\rangle \left\langle \tilde{Q}_2 \right\rangle, \tag{11}$$

$$C_P(1,2) := \left\langle \tilde{P}_1 \tilde{P}_2 \right\rangle - \left\langle \tilde{P}_1 \right\rangle \left\langle \tilde{P}_2 \right\rangle, \qquad (12)$$

which are called *quantum covariance functions* (QCF). By definition, QCF's vanish if and only if the system is separable [28]. Therefore (11) and (12) are zero for any not entangled quantum system.

Using the QCF's (11) and (12) we have

$$\left[(\Delta Q_1)^2 + (\Delta Q_2)^2 + 2C_Q(1,2) \right] \left[(\Delta P_1)^2 + (\Delta P_2)^2 + 2C_P(1,2) \right] \ge \frac{1}{4} \left| \langle [\tilde{Q}, \tilde{P}] \rangle \right|^2, \quad (13)$$

or

$$\sum_{i,j=1}^{2} C_Q(i,j) \sum_{k,l=1}^{2} C_P(k,l) \ge \frac{1}{4} \left| \langle [\tilde{Q}, \tilde{P}] \rangle \right|^2,$$
(14)

since $C_Q(i,i) = (\Delta Q_i)^2$, $C_P(i,i) = (\Delta P_i)^2$, $C_Q(i,j) = C_Q(j,i)$ and $C_P(i,j) = C_P(j,i)$.

³Note that $[\tilde{Q}_1, \tilde{Q}_2] = 0$ and $[\tilde{P}_1, \tilde{P}_2] = 0$.

In this work, we concern with the case of an entangled system of two identical particles, so we are going to handle Eq. (13) in order to express it in a more appropriate way. For this end, we define

$$|\psi\prime\rangle := \left(\tilde{Q}_1 - \tilde{Q}_2\right)|\psi\rangle,\tag{15}$$

with $|\psi \rangle, |\psi \rangle \in \mathcal{E}$ and $\langle \psi | \psi \rangle = 1$. Therefore,

$$\langle \psi \prime | \psi \prime \rangle = (\Delta_{\psi} Q_1)^2 + (\Delta_{\psi} Q_2)^2 - 2 \left\langle \tilde{Q}_1 \tilde{Q}_2 \right\rangle_{\psi} + \left\langle \tilde{Q}_1 \right\rangle_{\psi}^2 + \left\langle \tilde{Q}_2 \right\rangle_{\psi}^2.$$
(16)

Now, using the Schwarz inequality, $\langle \psi \mid \psi \rangle \langle \psi \prime \mid \psi \prime \rangle \geq \langle \psi \mid \psi \prime \rangle \langle \psi \prime \mid \psi \rangle$, we have

$$\left(\Delta_{\psi}Q_{1}\right)^{2} + \left(\Delta_{\psi}Q_{2}\right)^{2} \geq 2\left(\left\langle\tilde{Q}_{1}\tilde{Q}_{2}\right\rangle_{\psi} - \left\langle\tilde{Q}_{1}\right\rangle_{\psi}\left\langle\tilde{Q}_{2}\right\rangle_{\psi}\right).$$
(17)

In the same way

$$(\Delta_{\psi}P_1)^2 + (\Delta_{\psi}P_2)^2 \ge 2\left(\left\langle \tilde{P}_1\tilde{P}_2 \right\rangle_{\psi} - \left\langle \tilde{P}_1 \right\rangle_{\psi} \left\langle \tilde{P}_2 \right\rangle_{\psi}\right).$$
(18)

Finally, from inequalities (17), (18) and (13) we obtain

$$\left[(\Delta Q_1)^2 + (\Delta Q_2)^2 \right] \left[(\Delta P_1)^2 + (\Delta P_2)^2 \right] \ge \frac{1}{16} \left| \langle [\tilde{Q}, \tilde{P}] \rangle \right|^2.$$
(19)

In the case where $(\Delta Q_1)^2 = (\Delta Q_2)^2$ and $(\Delta P_1)^2 = (\Delta P_2)^2$ the inequality (19) becomes

$$\Delta Q_i \Delta P_i \ge \frac{1}{8} \left| \langle [\tilde{Q}, \tilde{P}] \rangle \right|.$$
⁽²⁰⁾

It is worth noting that the expression of the inequality (20) is independent of the chosen uncertainty principle that does not take into account the quantum correlation. This uncertainty principle is related to the commutation relation $[\tilde{Q}, \tilde{P}]$.

3 Uncertainty principle for entangled states in different minimallength scenarios

In this section we consider a system of two entangled identical particles whose momenta have the same value but opposite directions, that is, $\vec{p_1} = -\vec{p_2}$, just as in the Kim and Shih's experiment [21]. Therefore, in this case $\langle \hat{p}_1 \rangle + \langle \hat{p}_2 \rangle = 0$. Moreover, such a consideration also allows us to estimate, in the next section, an upper bound for the value of the minimal length based on the experimental results obtained by Kim and Shih.

3.1 Heisenberg uncertainty principle

Before we consider a minimal-length scenario it is appropriate to determine the change in the canonical HUP, that is, in a scenario in which effects of quantum gravity are not present. The canonical HUP for states of one simple-particle is

$$\Delta x \Delta p \ge \frac{\hbar}{2}.\tag{21}$$

The commutation related to the HUP is

$$[\hat{x}, \hat{p}] = i\hbar. \tag{22}$$

Hence

$$[\tilde{Q}, \tilde{P}] = [\tilde{Q}_1 + \tilde{Q}_2, \tilde{P}_1 + \tilde{P}_2] = 2i\hbar.$$
 (23)

Substituting Eq. (23) into Eq. (20) we get

$$\Delta Q_i \Delta P_i \ge \frac{\hbar}{4}.\tag{24}$$

The result (24) shows that for a system of two entangled identical particles the HUP is modified. Such an outcome is not new, it was already obtained by G. Rigolin in 2002 [22] and then in 2016 [23].

From a quick glance at the result (24) and recalling that dimensionally $(\Delta Q)_{min} \propto \hbar$, we expect the minimal uncertainty in the position will be reduced by half for all GUP's.

3.2 KMM GUP

The GUP

$$\Delta x_i \Delta p_i \ge \frac{\hbar}{2} \left[1 + \beta \left(\Delta p_i \right)^2 + \beta \left\langle \hat{p}_i \right\rangle^2 \right], \tag{25}$$

where β is a parameter related to the minimal length, has been proposed by A. Kempf, G. Mangano an R. B. Mann (KMM-GUP) [29] and it is the most used in the literature. The commutation relation related to it is given by

$$[\hat{x}_i, \hat{p}_i] = i\hbar \left(1 + \beta \hat{p}_i^2\right).$$
(26)

Hence

$$[\tilde{Q}, \tilde{P}] = i\hbar \left[1 + \beta \left(\tilde{P}_1^2 + \tilde{P}_2^2 \right) \right] = i\hbar \left(1 + 2\beta \tilde{P}_i^2 \right).$$
⁽²⁷⁾

Substituting Eq. (27) into Eq. (20) we get

$$\Delta Q_i \Delta P_i \ge \frac{\hbar}{4} \left[1 + \beta \left(\Delta P_i \right)^2 + \gamma \right], \qquad (28)$$

where $\gamma := \beta \left\langle \tilde{P}_i \right\rangle^2$.

The modified KMM-GUP (28) induces the existence of a minimal uncertainty given by

$$(\Delta Q_i)_{min} = \frac{\hbar}{2}\sqrt{\beta}.$$
(29)

The result above shows that the non-zero minimal uncertainty in position induced by the KMM-GUP for two entangled identical particles is twice smaller than for a separable system of two identical particles (non-entangled).

3.3 ADV GUP

A. Farag Ali, S. Das and E. C. Vagenas have proposed a GUP related to a commutation relation which has a linear and a quadratic term in the momentum operator [30],

$$[\hat{x}_i, \hat{p}_i] = i\hbar \left(1 - 2\alpha \hat{p}_i + 4\alpha^2 \hat{p}_i^2\right), \qquad (30)$$

where α is a parameter related to the minimal length. Besides the existence of a minimal length this linear approach induces a maximal uncertainty in the momentum, too. Then, from Eq. (30) we get

$$[\tilde{Q}, \tilde{P}] = 2i\hbar \left[1 - \alpha \left(\tilde{P}_1 + \tilde{P}_2 \right) + 2\alpha^2 \left(\tilde{P}_1^2 + \tilde{P}_2^2 \right) \right].$$
(31)

Therefore,

$$\langle [\tilde{Q}, \tilde{P}] \rangle = 2i\hbar \left[1 - \alpha \left(\left\langle \tilde{P}_1 \right\rangle + \left\langle \tilde{P}_2 \right\rangle \right) + 2\alpha^2 \left(\left\langle \tilde{P}_1^2 \right\rangle + \left\langle \tilde{P}_2^2 \right\rangle \right) \right], \tag{32}$$

$$\left\langle \left[\tilde{Q},\tilde{P}\right]\right\rangle = 2i\hbar \left[1 + 4\alpha^2 \left\langle \tilde{P}_i^2 \right\rangle\right],\tag{33}$$

Substituting Eq. (33) into Eq. (20) we obtain

$$\Delta Q_i \Delta P_i \ge \frac{\hbar}{4} \left[1 + 4\alpha^2 \left(\Delta P_i \right)^2 + \gamma \prime \right], \tag{34}$$

where $\gamma \prime := 4\alpha^2 \left\langle \tilde{P}_i \right\rangle^2$.

The modified ADV-GUP (34) induces the existence of a non-zero minimal uncertainty given by

$$(\Delta Q_i)_{min} = \hbar \alpha, \tag{35}$$

which once again is twice smaller than for a non-entangled system of two particles.

It is important to note that the linear GUP (ADV-GUP) becomes non-linear in this case and consequently a maximal uncertainty in the momentum is no longer induced.

3.4 Pedram GUP

In order to overcome some problems arising from KMM-GUP and ADV-GUP - such as incorporation of a maximal momentum required in doubly special relativity (DSR) theories, commutative geometry and an approach valid for all order in the parameter related to the minimal length - P. Pedram has proposed a GUP [31, 32] based on the commutation relation given by

$$[\hat{x}_i, \hat{p}_i] = \frac{i\hbar}{1 - \beta \hat{p}_i^2}.$$
(36)

Hence

$$[\tilde{Q}, \tilde{P}] = \frac{i\hbar}{1 - \beta \tilde{P}_1^2} + \frac{i\hbar}{1 - \beta \tilde{P}_2^2}.$$
(37)

Using the so-called Jensen's Inequality [33] we have

$$\langle [\tilde{Q}, \tilde{P}] \rangle \ge \frac{i\hbar}{1 - \beta \left\langle \tilde{P}_1^2 \right\rangle} + \frac{i\hbar}{1 - \beta \left\langle \tilde{P}_2^2 \right\rangle}$$
(38)

Substituting Eq. (38) into Eq. (20) we obtain

$$\Delta Q_i \Delta P_i \ge \frac{\hbar}{4} \frac{1}{\left[1 - \beta \left(\Delta P_i\right)^2 - \gamma\right]}.$$
(39)

Therefore, the modified Pedram-GUP (39) introduces a non-zero minimal uncertainty given by

$$\left(\Delta Q_i\right)_{min} = \frac{3\hbar}{8}\sqrt{3\beta},\tag{40}$$

which is also twice smaller than for a non-entangled system of a pair of identical particles.

3.5 Exponential all orders GUP

In the canonical field theory in the context of non-commutative coherent states representation and field theory on non-anticommutative superspace the Feynman propagator displays an ultra-violet (UV) cut-off of the form $e^{-\beta p^2}$ [34, 35, 36, 37]. In consequence, K. Nouicer has proposed an exponential all orders GUP [38, 39] based on the commutation relation given by

$$[\hat{x}_i, \hat{p}_i] = i\hbar e^{\beta \hat{p}_i^2}.$$
(41)

Hence

$$[\tilde{Q}, \tilde{P}] = i\hbar \left(e^{\beta \tilde{P}_1^2} + e^{\beta \tilde{P}_2^2} \right).$$
(42)

Again, using the so-called Jensen's Inequality [33] we have

$$\langle [\tilde{Q}, \tilde{P}] \rangle \ge i\hbar \left(e^{\beta \left\langle \tilde{P}_1^2 \right\rangle} + e^{\beta \left\langle \tilde{P}_2^2 \right\rangle} \right).$$
(43)

Substituting Eq. (42) into Eq. (20) we get

$$\Delta Q_i \Delta P_i \ge \frac{\hbar}{4} e^{\beta (\Delta P_i)^2 + \gamma}.$$
(44)

Therefore, the modified exponential all order GUP (44) introduces a minimal uncertainty given by

$$(\Delta Q_i)_{min} = \frac{\hbar}{2} \sqrt{\frac{e\beta}{2}}.$$
(45)

Finally, we note that $(\Delta Q_i)_{min}$ is also twice smaller than for a non-entangled system of a pair of identical particles, as we have already said.

3.6 Minimal length

If a non-zero minimal uncertainty in position can be interpreted as a minimal length then the previous results show that the minimal length for two entangled identical particles can be twice smaller than for a separable system. It is clear this statement must be taken with care because a minimal length should be a constant, that is, it must not depend on the physical system, it is a quantum gravitation effect. In fact, the minimal length should be an invariant as well as the light speed is. The answer for that apparent contradiction is possibly because the system is made up of two particles. In references [40], C. Quesne and V. M. Tkachuck have claimed that for a system composed by N particles the effective parameter related to the minimal length, β , is reduced by a factor $\frac{1}{N^2}$. Hence, $\frac{\hbar}{2}\sqrt{\beta}$ is not the correct minimal uncertainty in position $(\Delta Q_i)_{min}$ for the modified KMM-GUP, but $\frac{\hbar}{2}\sqrt{\beta_i}$, where [40, 41]

$$\beta = \frac{\beta_i}{2^2}.\tag{46}$$

Consequently, we find that $(\Delta Q_i)_{min} = \hbar \sqrt{\beta}$, therefore $l_{min} = \hbar \sqrt{\beta}$, as we expected⁴. In the same way for others GUP's.

The reader might want to claim that the minimal length is actually described by the minimal uncertainty of the entangled system. However, according to reference [23] we can presume that the minimal uncertainty for a system of N entangled identical particles is reduced by $\frac{1}{N}$. Since, in principle, there is no limit for the number of particles for a entangled system, there would also be no limit for the minimal length.

4 Upper bound for the minimal-length value

In the Kim and Shih's experiment entangled identical pairs of photons were produced by spontaneous parametric down conversion (SPDC) with momentum conservation. A narrow physical slit was placed along the trajectory of one of the photons, whereas the

 $^{^{4}}$ It is worth noting that if the minimal uncertainty was greater for the entangled system, we could suppose that the quantum entanglement decreases the accuracy of a position measurement thereby increasing the minimal uncertainty.

other photon of the pair (called 2) passed through a virtual slit. The ghost image experimental technique [42] ensured that the quantum correlation between the pair of photons was not destroyed. Then simultaneous detection of photons of the pairs were performed and data just for coincidence events were obtained in case when the photon 2 passed through a virtual slit (non-slit case) and in case when the photon 2 passed through a physical slit (slit case).

From the experimental data obtained by Kim and Shih one gets that [23]

$$\frac{\Delta P_2^{ns}}{\Delta P_2^s} = \frac{1.25}{2.15},\tag{47}$$

where ΔP_2^{ns} is uncertainty in momentum of the photon 2 in the non-slit case and ΔP_2^s is uncertainty in momentum of the photon 2 in slit case. Since the width of the slit was 0.16 mm ($\Delta Q_2 = 0.16$ mm), the uncertainty in momentum in the slit case for KMM-GUP can be find from⁵

$$0.16\Delta P_2^s = \frac{\hbar}{2} \left[1 + 4\beta \left(\Delta P_i^s \right)^2 + \gamma \right].$$
 (48)

Eq. (48) has real roots only if $\beta \leq \frac{0.16^2}{\hbar^2 \eta}$, where $\eta := 1 + \gamma$. Thus,

$$\Delta P_{2+}^s = \frac{0.32}{\hbar\beta} - \frac{\hbar\eta}{0.32} - \beta \frac{\hbar^3 \eta^2}{0.32^3} \tag{49}$$

and

$$\Delta P_{2-}^s = \frac{\hbar\eta}{0.32} + \beta \frac{\hbar^3 \eta^2}{0.32^3},\tag{50}$$

Now, using the above results we can estimate an upper bound for the minimal-length value induced by KMM-GUP. Hence, substituting the root (49) into Eq. (28) we have that

$$\beta \le \frac{3.58 \times 10^{-2}}{\hbar^2 \eta}.\tag{51}$$

Therefore,

$$l_{min} \le \hbar \sqrt{\frac{3.58 \times 10^{-2}}{\hbar^2 \eta}} \le \hbar \sqrt{\frac{3.58 \times 10^{-2}}{\hbar^2}} = 1.9 \times 10^{-4} m.$$
(52)

The substitution of the root (50) hold the inequality (28) for all $\beta > 0$.

Consequently, the upper bound for the minimal-length value is order 10^{-4} m. Therefore, using the experiment described above a result with less restrictions than those reported in the literature is obtained [41, 43, 44, 45, 46, 47, 48, 49].

⁵Note that in with-slit case the correlation between the photons of the pair was destroyed, therefore the photons were not entangled and the usual KMM-GUP, Eq. (25), is held.

5 Conclusion

In this work we find the non-zero minimal uncertainties induced by the main proposals of GUP's (KMM, ADV, Pedram and exponential) which are modified due to the quantum entanglement of a system of two identical particles. In principle, our results have pointed out that the minimal uncertainties are reduced at half for a system of two entangled identical particles independently of the GUP. Hence, if a non-zero minimal uncertainty in position can be interpreted as a minimal length then the quantum entanglement reduces by half the minimal length. However, the minimal length must not depend on the physical system. We overcome this apparent paradox by using the Quesne and Tkachuck's proposal for a system composed. Consequently, despite the quantum entanglement to change the GUP, the minimal length does not change. Based on our results and using the reference [23] we can expect that the apparent minimal uncertainty for a system of N entangled identical particles is reduced by $\frac{1}{N}$, nonetheless the minimal length does not change because the effective parameter β is also reduced by a factor $\frac{1}{N^2}$.

Finally, we have estimated from the data obtained from the Kim and Shih's experiment an upper bound value for the minimal length of the order of 10^{-4} m. Consequently, it is rather an inexpressive value (in the sense of leading to poor predictive power) as compared to ones have been found in the literature. This is due to the high imprecision of the experiment on the entangled system described above. However, we may expect that more refined version of the experiment may lead to more stringent bounds on the minimum length.

Acknowledgements

We would like to thank FAPES, CAPES and CNPq (Brazil) for financial support.

References

- R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki "Quantum entanglement", Rev. Mod. Phys. 81(2), 865 (2009). doi.org/10.1103/RevModPhys.81.865
- [2] J. Brody QUANTUM ENTANGLEMENT, (MIT Press, Cambridge, Massachusetts, 2020.)
- [3] A. Eisntein, B. Podolsky and N. Rose, "Can Quantum-Mechanical Description of Physics Reality Be Considered Complete?", Phys. Rev. 47(10), 777 (1935). doi.org/10.1103/PhysRev.47.777
- M. A. Nielsen and I. Chung, "Quantum Computation and Quantum Information", Am. J. Phys. 70(5), 558 (2002). doi.org/10.1119/1.1463744

- [5] M. A. Nielsen and I. Chung, *Quantum Computation and Quantum Information*, 10th Anniversary edition, (Cambridge University Press, Cambridge, 2010.)
- [6] K. Konishi, G. Paffuti and P. Provero, "Minimum physical length and the generalized uncertainty principle in string theory", Phys. Lett. B 234(3), 276 (1990). doi.org/10.1016/0370-2693(90)91927-4
- [7] M. Maggiore, "A Generalized uncertainty principle in quantum gravity", Phys. Lett. B 304(1-2), 65 (1993). doi.org/10.1016/0370-2693(93)91401-8
- [8] M. Maggiore, "The algebraic structure of the generalized uncertainty principle", Phys. Lett. B 319(1-3), 83 (1993). doi.org/10.1016/0370-2693(93)90785-G
- M. Maggiore, "Quantum groups, gravity, and the generalized uncertainty principle", Phys. Rev. D 49(10), 5182 (1994). doi.org/ 10.1103/PhysRevD.49.5182
- [10] R. J. Adler and D. I. Santiago, "On gravity and the uncertainty principle", Mod. Phys. Lett. A 14(20), 1371 (1999). doi.org/10.1142/S0217732399001462
- [11] F. Scardigli, "Generalized uncertainty principle in quantum gravity from micro-black hole gedanken experiment", Phys. Lett. B 452(1-2), 39 (1999). doi.org/10.1016/S0370-2693(99)00167-7
- [12] F. Scardigli, "Generalized uncertainty principle, extra dimensions and holography", Class. Quantum Grav. 20(18), 3915 (2003). doi.org/10.1088/0264-9381/20/18/305
- [13] S. Das and E. C.Vagenas, "Phenomelogical implications of the generalized uncertainty principle", Can. J. Phys. 87(3), 233 (2009). doi.org/10.1139/P08-105
- [14] A. Tawfik and A. Diab, "Generalized uncertainty principle: Approaches and applications", Int. J. Mod. Phys. D 23(12), 1430025 (2014). doi.org/10.1142/S0218271814300250
- [15] F. Scardigli and R. Casadio, "Gravitational tests of the generalized uncertainty principle", Eur. Phys. J. C 75, 425 (2015). doi.org/10.1140/epjc/s10052-015-3635-y
- [16] A. Tawfik and A. Diab, "Review on Generalized Uncertainty Principle", Rept. Prog. Phys. 78(12), 126001 (2015). doi.org/10.1088/0034-4885/78/12/126001
- [17] S. Hossenfelder, "A note on theories with a minimal length", Class. Quantum Grav.
 23, 1815 (2006). doi.org/10.1088/0264-9381/23/5/N01
- [18] M. Kober, "Generalized Uncertainty Principle in Canonical Quantum Gravity and Application to Quantum Cosmology", Int. J. Mod. Phys. A 27(20), 1250106 (2012). doi.org/10.1142/S0217751X12501060

- [19] Y. Sabri and Kh. Nouicer, "Phase transitions of a GUP-corrected Schwarzschild black hole within isothermal cavities", Class. Quantum Grav. 29(21), 215015 (2012). doi:10.1088/0264-9381/29/21/215015.
- Y. C. Ong and Y. Yao, "Generalized uncertainty principle and white dwarfs redux: How the cosmological constant protects the Chandrasekhar limit", Phys. Rev. D 98(12), 126018 (2018). doi.org/10.1103/PhysRevD.98.126018
- [21] Yoon-Ho Kim and Y. Shih, "Experimental Realization of Popper's Experiment: Violation of the Uncertainty Principle?", Found. Phys. 29(12), 1849 (1999). doi.org/10.1023/A:1018890316979
- [22] G. Rigolin, "Uncertainty Relations for Entangled States", Found. Phys. Lett. 15(3), 293 (2002). doi.org/10.1023/A:1021039822206
- [23] G. Rigolin, "Entanglement, Identical Particles and the Uncertainty Principle", Commun. Theor. Phys. 66, 201 (2016). doi.org/10.1088/0253-6102/66/2/201
- [24] G. Blado, F. Herrera and J. Erwin, "Entanglement and the generalized uncertainty principle", Phys. Essays 31(4), 397 (2018). doi.org/10.4006/0836-1398-31.4.397
- [25] D. Park, "Quantum entanglement with generalized uncertainty principle", Nucl. Phys. B 977(7), 115736 (2022). doi.org/10.1016/j.nuclphysb.2022.115736
- [26] C. Cohen-Tannoudji, B. Dui and F. laloe, *Quantum Mechanics*, (Jhon Wiley and Sons, Paris, 1977).
- [27] A. Messiah, *Quantum Mechanics*, (North-Holland Publishing, Amsterdan, 1965).
- [28] A. C. de la Torre, P. Catuogno and S. Ferrando, "Uncertainty and nonseparability", Found. Phys. Lett. 2, 235 (1989). doi.org/10.1007/BF00692669
- [29] A. Kempf, G. Mangano and R. B. Mann, "Hilbert Space Representation Of The Minimal Length Uncertainty Relation", Phys. Rev. D 52(2), 1108 (1995). doi.org/10.1103/PhysRevD.52.1108
- [30] A. Farag Ali, S. Das and E. C. Vagenas, "Discreteness of space from the generalized uncertainty principle", Phys. Lett. B 678(5), 497 (2009). doi.org/10.1016/j.physletb.2009.06.061
- [31] P. Pedram, "A higher order GUP with minimal length uncertainty and maximal momentum", Phys. Lett. B 714(2-5), 317 (2012). doi.org/10.1016/j.physletb.2012.07.005
- [32] P. Pedram, "A higher order GUP with minimal length uncertainty and maximal momentum II: Applications", Phys. Lett. B 718(2), 638 (2012). doi.org/10.1016/j.physletb.2012.10.059

- [33] J. L. W. V. Jensen "Sur les fonctions convexes et les inégalités entre les valeurs moyennes", Acta Math. 30, 175 (1906). doi.org/10.1007/BF02418571
- [34] J. W. Moffat, "Noncommutative and non-anticommutative quantum field theory", Phys. Lett. B 506(1-2), 193 (2001). doi.org/10.1016/S0370-2693(01)00409-9
- [35] A. Smailagic and E. Spallucci, "Feynman path integral on the non-commutative plane", J. Phys. A: Math. Gen. 36(33), L467 (2003). doi.org/10.1088/0305-4470/36/33/101
- [36] A. Smailagic and E. Spallucci, "UV divergence-free QFT on noncommutative plane", J. Phys. A: Math. Gen. 36(39), L517 (2003). doi.org/10.1088/0305-4470/36/39/103
- [37] A. Smailagic and E. Spallucci, "Lorentz invariance, unitarity and UV-finiteness of QFT on noncommutative spacetime", J. Phys. A: Math. Gen. 37(28), 7169 (2004). doi.org/10.1088/0305-4470/37/28/008
- [38] Kh. Nouicer, "Quantum-corrected black hole thermodynamics toallPlanck length", Phys. Lett. В 626(2-3),(2007).orders inthe 63 doi.org/10.1016/j.physletb.2006.12.072
- [39] Kh. Nouicer, "Black hole thermodynamics to all orders in the Planck length in extra dimensions", Class. Quantum Grav. 24(23), 5917 (2007). doi.org/10.1088/0264-9381/24/23/014
- [40] C. Quesne and V. M. Tkachuk, "Composite system in deformed space with minimal", Phys. Rev. A 81, 012106 (2010). doi.org/ 10.1103/PhysRevA.81.012106
- [41] T. S. Quintela Jr., J. C. Fabris and J. A. Nogueira "The Harmonic Oscillator in the Classical Limit of a Minimal-Length Scenario", Braz. J. Phys. 46, 777 (2016). doi.org/10.1007/s13538-016-0457-9
- [42] T. B. Pittman, Y. H. Shih, D. V. Strekalov, and A. V. Sergienko, "Optical imaging by means of two-photon quantum entanglement", Phys. Rev. A 52(5), R3429 (1995). doi.org/10.1103/PhysRevA.52.R3429
- [43] M. Hemeda, H. Alshal, A. F. Ali, E. C. Vagenas "Gravitational Observations and LQGUP", [arXiv:gr-qc/arXiv:2208.04686v1]. doi.org/10.48550/arXiv.2208.04686
- [44] S. Sen, S. Bhattacharyya and S. Gangopadhyay "Probing the generalized uncertainty principle through quantum noises in optomechanical systems", Class. Quantum Grav. 39(7), 075020 (2022). doi.org/10.1088/1361-6382/abe758
- [45] S. Bhattacharyya, S. Gangopadhyay and A. Saha "Generalized uncertainty principle in resonant detectors of gravitational waves", Class. Quantum Grav. 37(19), 195006 (2020). doi.org/10.1088/1361-6382/abac45

- [46] M. F. Gusson, A. Oakes O. Gonçalves, R. O. Francisco, R. G. Furtado, J. C. Fabris, J. A. Nogueira "Dirac δ-function potential in quasiposition representation of a minimal-length scenario", Eur. Phys. J. C 78, 179 (2018). doi.org/10.1140/epjc/s10052-018-5659-6
- [47] K. Zeynali, F. Darabi and H. Motavalli, "Probing Planck-scale physics with quantum optics", Nat. Phys. 12, 033 (2012). doi.org/10.1088/1475-7516/2012/12/033
- [48] I. Pikovski, M. R. Vanner, M Aspelmeyer, M. S. Kim and C. Brukner "Probing Planck-scale physics with quantum optics", Nat. Phys. 8, 393 (2012). doi.org/10.1038/nphys2262
- [49] F. L. Antonacci Oakes, R. O. Francisco, J. C. Fabris and J. A. Nogueira, "Ground state of the hydrogen atom via Dirac equation in a minimal-length scenario", Eur. Phys. J. C 73, 2495 (2013). doi.org/10.1140/epjc/s10052-013-2495-6