The gauge-invariant formulation of the local expansion rate driven by the local average density in an inhomogeneous universe

Masanori Tomonaga $^{*1},$ Masumi Kasai $^{\dagger 2},$ and Toshifumi Futamase $^{\ddagger 1}$

¹Faculty of Science, Kyoto Sangyo University, Motoyama, Kamigamo, Kita-ku, Kyoto, 603-8555 Japan ²Graduate school of Science and Technology, Hirosaki University, Hirosaki 036-8561, Japan

Abstract

The Hubble tension casts a blight on the standard cosmology. As a possible solution to the problem, the local variation of the expansion rate has been proposed where the spatial averaging over a finite domain was introduced in order to restore the local Friedmannian behavior in an inhomogeneous cosmology. So far, however, the approaches are limited to the particular choices of the gauges, and it has been unclear whether the results are gauge-invariant. In this paper, we present the gauge-invariant formulation of the local expansion rate which is driven by the spatial average of the gauge-invariant inhomogeneous density. We show that the local cosmological parameters in the finite domain may change from the global parameters, and the relations between them are expressed by the gauge-invariant averaged density.

1 Introduction

The Hubble constant H_0 is one of the most important cosmological parameter since it characterizes the global properties of our universe. The standard cosmology is based on the assumption of the homogeneity and isotropy. Thus, the Hubble parameter H_0 is regarded as a constant over at least the horizon scale which is also the prediction of the inflationary scenario. However recent observations suggest a non negligible difference between local and global (or recent

^{*}email:i2185088@cc.kyoto-su.ac.jp

[†]email:kasai@hirosaki-u.ac.jp

 $^{^\}ddagger$ email:tof@cc.kyoto-su.ac.jp

and old) Hubble parameter [1, 2].

There has been a large number of studies which try to resolve the discrepancy [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18]. We regard that the difference of the local and global Hubble parameter is real and be explained by the inhomogeneous distribution of the matter. In fact, the observation of the K-band luminosity density seems to suggest that region with several hundred Mpc around us has low density with density contrast $\delta_K \sim -0.5$ compared with the globally averaged density [19]. Furthermore, there is some indication that the voids are actually low density by weak lensing observation. Thus, it will be meaningful to pursue the indication of the cosmological inhomogeneity.

The homogeneous and isotropic universe (here we call Friedmann Universe) appears as the result of some kind of averaging procedure since the universe is actually very inhomogeneous. There are various ways to averaging inhomogeneous universe (such as the light-corn averaging that is directly related with observational quantities). In this paper, we only consider the scalar perturbations in the linear order and the spatial averaging [20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35]. Our purpose is not studying the averaging itself in general inhomogeneous spacetimes, but rather the gauge dependence of the relationship between locally averaged and globally averaged spacetime in the linearly perturbed universe using the spatial averaging. By adopting the spatial averaging defined below, we were able to derive a locally averaged Friedmann universe and have obtained the following relation between the locally average Hubble parameter and the globally averaged Hubble parameter within the framework of the general relativistic perturbation theory [36, 37]

$$H_{D0} = H_0 \left(1 - \frac{1}{3} f(t_0) \langle \Delta \rangle_{Dt_0} \right), \tag{1}$$

where H_{D0} is the averaged Hubble parameter at the present time t_0 over a finite domain D, and H_0 is the global, or the horizon scale Hubble parameter, and $\langle \Delta \rangle_{Dt_0}$ is the present density contrast average over the domain D. $f(t) = d \log \Delta/d \log a$ is the growth function of the density contrast. However, the treatment is carried out in the comoving synchronous and Newtonian gauge, and there is some question if the averaging and the result are gauge invariant or not.

In order to answer the question, we study the spatial averaging in the framework of the gauge invariant cosmological perturbation theory, and find that local Hubble parameter (and local cosmological parameter) can be describe used by gauge-invariant physical quantities averaged in the local region D.

2 Gauge-invariant linear perturbation theory

In this section, we briefly summarize the gauge-invariant perturbation theory [38, 39]. We assume the flat background with dust fluid. Then the background metric is

$$ds^2 = -dt^2 + a^2(t)\delta_{ij}dx^idx^j$$
(2)

and the energy momentum tensor is

$$T^{\mu} = \rho_b(t) u^{\mu} u^{\nu}, \quad u^{\mu} = (1, 0, 0, 0).$$
 (3)

From the Einstein equations, we obtain the following Friedmann equation

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho_b + \frac{\Lambda}{3} \tag{4}$$

and from $T^{\mu\nu}_{;\nu} = 0$, we obtain the energy conservation equation

$$\dot{\rho}_b + 3\frac{\dot{a}}{a}\rho_b = 0. \tag{5}$$

Next, we write the metric and the energy-momentum tensor in the perturbed universe as follows:

$$ds^2 = g_{\mu\nu}dx^{\mu}dx^{\nu} \tag{6}$$

$$g_{00} = -(1+2A) \tag{7}$$

$$g_{0i} = -B_{,i} \tag{8}$$

$$g_{ij} = a^2(t) \left(\delta_{ij} + 2E_{,ij} + 2F\delta_{ij} \right) \tag{9}$$

$$T^{\mu\nu} = \rho \, u^{\mu} u^{\nu} \tag{10}$$

$$\rho = \rho_b \left(1 + \delta \right) \tag{11}$$

$$u^{\mu} = (u^{0}, u^{i}) = (1 - A, a^{-2} \delta^{ij} v_{,i}), \qquad (12)$$

where we consider only scalar perturbations and the scalar perturbation variables A, B, E, F, δ and v are arbitrary functions of t and x^i , and assumed to be small quantities.

Now consider the scalar type infinitesimal gauge transformation

$$\bar{t} = t + \alpha \,, \tag{13}$$

$$\bar{x}^i = x^i + \delta^{ij}\beta_i \,, \tag{14}$$

where α and β are arbitrary functions of t and x^i , which are regarded as small as the perturbation variables. The gauge dependence of the perturbed quantities are

$$\bar{A} = A - \dot{\alpha} \,, \tag{15}$$

$$\bar{B} = B - \alpha + a^2 \dot{\beta} \,, \tag{16}$$

$$\bar{E} = E - \beta \,, \tag{17}$$

$$\bar{F} = F - \frac{\dot{a}}{a}\alpha\,,\tag{18}$$

$$\bar{\delta} = \delta + 3\frac{\dot{a}}{a}\alpha\,,\tag{19}$$

$$\bar{v} = v + a^2 \dot{\beta} \,. \tag{20}$$

Then, the following gauge invariant quantities are defined in the usual manner.

$$\Phi \equiv A - \left(B + a^2 \dot{E}\right)^{\cdot}, \tag{21}$$

$$\Psi \equiv F - \frac{\dot{a}}{a} \left(B + a^2 \dot{E} \right) \,, \tag{22}$$

$$\Delta \equiv \delta - 3\frac{\dot{a}}{a}(v - B) , \qquad (23)$$

$$V \equiv v + a^2 \dot{E} \,. \tag{24}$$

Using these quantities, we can obtain the first-order equations in terms of the gauge invariant quantities of linearized Einstein equation as follows:

$$-\frac{1}{a^2}\nabla^2\Psi = 4\pi G\rho_b\Delta\tag{25}$$

$$\frac{\dot{a}}{a}\Phi - \dot{\Psi} = -4\pi G \rho_b V \tag{26}$$

$$\Psi + \Phi = 0. \tag{27}$$

$$\Psi + \Phi = 0. \tag{27}$$

Using (27), the equations (25) and (26) are re-written as

$$\frac{1}{a^2}\nabla^2\Phi = 4\pi G\rho_b\Delta,\tag{28}$$

$$\dot{\Phi} + \frac{\dot{a}}{a}\Phi = -4\pi G \rho_b V. \tag{29}$$

From $T^{\mu\nu}_{;\nu} = 0$, we obtain

$$\dot{\Delta} + \frac{1}{a^2} \nabla^2 V = 0, \tag{30}$$

$$\dot{V} + \Phi = 0. \tag{31}$$

Differentiating (30) with respect to t and using (28) and (31), we obtain

$$\ddot{\Delta} + 2\frac{\dot{a}}{a}\dot{\Delta} - 4\pi G\rho_b \Delta = 0. \tag{32}$$

The solution of the second-order differential equation (32) generally has two independent modes as follows:

$$\Delta(t, x^{i}) = \mathcal{D}_{+}(t)Q_{+}(x^{i}) + \mathcal{D}_{-}(t)Q_{-}(x^{i}), \qquad (33)$$

where

$$\mathcal{D}_{+}(t) = H \int^{t} \frac{dt'}{\left(aH\right)^{2}},\tag{34}$$

$$\mathcal{D}_{-}(t) = H = \frac{\dot{a}}{a},\tag{35}$$

 $Q_{+}(x)$ and $Q_{-}(x)$ represent the spatially dependent part of the growing and decaying mode of the density contrast, respectively.

In summary, from (4) and (28) multiplied by 2/3, we obtain the following equation

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{2}{3}\frac{1}{a^2}\nabla^2\Phi = \frac{8\pi G}{3}\left(\rho_b + \rho_b\Delta\right) + \frac{\Lambda}{3}$$
 (36)

as the perturbed version of the Friedmann equation, and from (5) including $\rho_b \Delta$,

$$\frac{\partial}{\partial t} (\rho_b + \rho_b \Delta) + 3 \frac{\dot{a}}{a} (\rho_b + \rho_b \Delta) - \rho_b \dot{\Delta} = 0$$
 (37)

as the perturbed version of the energy conservation equation.

3 Spatial averaging over a local domain in the perturbed universe

In the previous section, we have employed the standard assumption that the inhomogeneous matter density ρ can be decomposed into the homogeneous background part $\rho_b(t)$ and the small perturbed part δ . In the actual inhomogeneous universe, however, we need to extract the homogeneous part through the averaging procedure.

We define the spatial volume V_D of a finite small domain D in the t= const. hypersurface Σ_t as

$$V_D \equiv \int_D \sqrt{\det(g_{ij})} \, d^3x \,. \tag{38}$$

D is sufficiently smaller than the horizon scale but more than the scale at which the picture of the Hubble expansion is valid, e.g. more than several 10 Mpc. Using the metric described in the previous section, Σ_t is specified by the normal vector

$$n^{\mu} = \left(1 - A, \frac{1}{a^2} \delta^{ij} B_{,j}\right). \tag{39}$$

Contrary to those in [36] and [37], no gauge-fixing is made in (38) in order to specify the t= const. hypersurface Σ_t . Fixing the gauge A=B=v=0 reproduces the results in [36], and another gauge B=E=0 leads to those in [37]. The spatial average of a scalar quantity $Q(t,x^i)$ over the domain D is in general

$$\langle Q \rangle \equiv \frac{1}{V_D} \int_D Q \sqrt{\det(g_{ij})} \, d^3 x \,.$$
 (40)

Therefore, the average density in this domain is

$$\langle \rho \rangle \equiv \frac{1}{V_D} \int_D \rho \sqrt{\det(g_{ij})} \, d^3 x \,.$$
 (41)

Since we can observe only a finite nearby region of the entire space, it is likely that the average density $\langle \rho \rangle$ in the nearby region does not always coincides with the background density ρ_b .

Spatially averaging (37) over a local domain D, we obtain

$$\left\langle \frac{\partial \varrho}{\partial t} \right\rangle + 3 \frac{\dot{a}}{a} \langle \varrho \rangle - \rho_b \langle \dot{\Delta} \rangle = 0,$$
 (42)

where we have defined the gauge-invariant inhomogeneous density

$$\varrho \equiv \rho_b + \rho_b \Delta \tag{43}$$

in order to distinguish it from $\rho = \rho_b + \rho_b \delta$.

Note that the time derivative does not commute with the spatial averaging in general. In fact, for a physical quantity Q we have

$$\left\langle \frac{\partial Q}{\partial t} \right\rangle - \frac{d}{dt} \left\langle Q \right\rangle = \left\langle \frac{1}{2} g^{ij} \dot{g}_{ij} \right\rangle \left\langle Q \right\rangle - \left\langle \frac{1}{2} g^{ij} \dot{g}_{ij} Q \right\rangle. \tag{44}$$

However, if we consider the case $Q \to \rho$ up to the linear order, we obtain

$$\left\langle \frac{\partial \varrho}{\partial t} \right\rangle - \frac{d}{dt} \left\langle \varrho \right\rangle = \left\langle \frac{1}{2} g^{ij} \dot{g}_{ij} \right\rangle \left\langle \rho_b + \rho_b \Delta \right\rangle - \left\langle \frac{1}{2} g^{ij} \dot{g}_{ij} (\rho_b + \rho_b \Delta) \right\rangle$$

$$= \left\langle 3 \frac{\dot{a}}{a} \right\rangle \left\langle \rho_b \Delta \right\rangle - \left\langle 3 \frac{\dot{a}}{a} \rho_b \Delta \right\rangle$$

$$= 0. \tag{45}$$

Therefore, using the relation (45), it is straightforward to show from (42) the following equation holds up to the linear order of the perturbations:

$$\frac{d}{dt} \langle \varrho \rangle + 3 \frac{\dot{a}_D}{a_D} \langle \varrho \rangle = 0, \qquad (46)$$

where

$$\frac{\dot{a}_D}{a_D} \equiv \frac{\dot{a}}{a} - \frac{1}{3} \langle \dot{\Delta} \rangle \tag{47}$$

can be regarded as the local expansion rate driven by the local average density $\langle \varrho \rangle$.

In order to express (36) in terms of a_D , we rewrite as

$$\left(\frac{\dot{a}}{a}\right)^2 - \frac{2}{3}\frac{\dot{a}}{a}\dot{\Delta} + \frac{2}{3}\frac{1}{a^2}\left(\nabla^2\Phi + a^2\frac{\dot{a}}{a}\dot{\Delta}\right) = \frac{8\pi G}{3}\left(\rho_b + \rho_b\Delta\right) + \frac{\Lambda}{3}.\tag{48}$$

Spatially averaging (48) and substituting (47), we obtain

$$\left(\frac{\dot{a}_D}{a_D}\right)^2 + \frac{K_{\text{eff}}}{a_D^2} = \frac{8\pi G}{3} \langle \varrho \rangle + \frac{\Lambda}{3} \,, \tag{49}$$

where

$$K_{\text{eff}} \equiv \frac{2}{3} \left\langle \nabla^2 \Phi + a^2 \frac{\dot{a}}{a} \dot{\Delta} \right\rangle \tag{50}$$

$$= \frac{2}{3} \left\langle -a^2 \dot{H} \Delta + a^2 H \dot{\Delta} \right\rangle \tag{51}$$

$$= \frac{2}{3}a^2H^2\left(\frac{\mathcal{D}_+}{H}\right) \cdot \langle Q_+(x_i)\rangle \tag{52}$$

$$=\frac{2}{3}\langle Q_{+}(x_{i})\rangle\tag{53}$$

is a constant which can be regarded as the effective curvature constant on the local domain in the averaged sense.

Although (48) looks similar to that obtained in [36], we emphasize the following advantages of our analysis in this paper over that in [36].

- (1) [36]'s result is heavily dependent on the solution of $\delta \propto a$ in the Einstein-de Sitter background. In particular all of the averaged quantities are defined and calculated directly using the growing mode solution in the Einstein-de Sitter background described in eqs.(14-17) in [36]. So it is unclear whether it holds in any other background. In this paper, we explicitly showed that this averaged picture holds backgrounds other than the Einstein-de Sitter background, especially even if $\Lambda \neq 0$ background.
- (2) If we don't ignore the decaying mode of δ , [36] does not work. However, our discussion has no problem even if we consider the decaying mode.
- (3) It was unclear that [36]'s result is valid gauges other than comoving synchronous gauge. Therefore, we explicitly showed that we can describe using the spatial average of gauge-invariant variables all the averaged density, expansion rate, and (effective) curvature constant in an inhomogeneous universe.

4 The cosmological parameters in the nearby regions expressed by the gauge-invariant variables

We define the global Hubble parameter as

$$H_0 \equiv \frac{\dot{a}}{a}\Big|_{t_0} \tag{54}$$

and the global density parameters as

$$\Omega_m \equiv \frac{8\pi G \rho_b(t_0)}{3H_0^2} \tag{55}$$

and

$$\Omega_{\Lambda} \equiv \frac{\Lambda}{3H_0^2} \,, \tag{56}$$

where $\Omega_m + \Omega_{\Lambda} = 1$ since we have assumed the flat background. These global parameters are supposed to be determined by the very large-scale and distant observations such as the cosmic microwave background.

On the other hand, the cosmological parameters which are obtained from the observations in the local nearby regions are certainly determined by the local average density $\langle \varrho \rangle$, rather than by the background density ρ_b . We define the local Hubble parameter as

$$\tilde{H}_0 \equiv \frac{\dot{a}_D}{a_D}\Big|_{t_0} = H_0 \left(1 - \frac{1}{3}f(t_0)\langle \Delta \rangle_{t_0}\right), \tag{57}$$

where

$$f(t) \equiv \frac{d \ln \mathcal{D}_{+}}{d \ln a} \tag{58}$$

is the growth function of the gauge-invariant density perturbation Δ , and the local density parameters as

$$\tilde{\Omega}_m \equiv \frac{8\pi G \langle \varrho \rangle}{3\tilde{H}_0^2} = \Omega_m \left\{ 1 + \left(1 + \frac{2}{3} f(t_0) \right) \langle \Delta \rangle_{t_0} \right\}$$
(59)

and

$$\tilde{\Omega}_{\Lambda} \equiv \frac{\Lambda}{3\tilde{H}_0^2} = \Omega_{\Lambda} \left(1 + \frac{2}{3} f(t_0) \langle \Delta \rangle_{t_0} \right) , \tag{60}$$

which are valid up to the linear order in the gauge-invariant variable Δ .

The local cosmological parameters coincide with the global ones if and only if $\langle \Delta \rangle = 0$. Otherwise, the local parameters may change. Let us show a simple estimation in the case $\Lambda = 0$, where f(t) = 1. If the local nearby region is, say, 30% under dense, namely $\langle \Delta \rangle_{t_0} = -0.3$, the local Hubble parameter \tilde{H}_0 can be 10% larger than the global H_0 .

5 Conclusion and Discussion

Motivated by the Hubble tension, there have been many studies on the possible resolutions. One of them is the local variation of the cosmological parameters due to inhomogeneous matter distribution. We have also studied the inhomogeneous universe by spatial averaging and obtained an interesting result on the relation between the local and global Hubble parameters which might explain the Hubble tension.

However, the question of the gauge invariance of the result is not fully understood. In this paper we address this question. We employ the gauge-invariant linear cosmological perturbation theory to show that the relationship between local and global cosmological parameter can be describe used by the gauge-invariant physical quantities that averaged in the local region.

It is of some interest to develop this treatment to the second order since the density contrast report by the observation of the K-band luminosity density is of the order -0.5. Although we gave an argument based on the order of magnitude discussion of the cosmological Poisson equation, it is clearly not sufficient. Another direction of this study is to consider the possible interpretation by the inhomogeneity of the observation of m-z relation of Type Ia supernovae and CMB Power spectrum. We hope to study this possibility in future.

Acknowledgment

This work is supported by a Grant-in-Aid for Scientific Research from JSPS (No. 20K03937) for TF.

References

- Nabila Aghanim, Yashar Akrami, Mark Ashdown, J Aumont, C Bacci-galupi, M Ballardini, AJ Banday, RB Barreiro, N Bartolo, S Basak, et al. Planck 2018 results-vi. cosmological parameters. Astronomy & Astro-physics, 641:A6, 2020.
- [2] Adam G Riess, Lucas M Macri, Samantha L Hoffmann, Dan Scolnic, Stefano Casertano, Alexei V Filippenko, Brad E Tucker, Mark J Reid, David O Jones, Jeffrey M Silverman, et al. A 2.4% determination of the local value of the hubble constant. The Astrophysical Journal, 826(1):56, 2016.
- [3] Kiyotomo Ichiki, Chul-Moon Yoo, and Masamune Oguri. Relationship between the cmb, sunyaev-zel'dovich cluster counts, and local hubble parameter measurements in a simple void model. *Physical Review D*, 93(2):023529, 2016.
- [4] Krzysztof Bolejko. Emerging spatial curvature can resolve the tension between high-redshift cmb and low-redshift distance ladder measurements of the hubble constant. *Physical Review D*, 97(10):103529, 2018.
- [5] Kenji Tomita. Cosmological models with the energy density of random fluctuations and the hubble-constant problem. *Progress of Theoretical and Experimental Physics*, 2017(8), 2017.
- [6] Kenji Tomita. Super-horizon second-order perturbations for cosmological random fluctuations and the hubble-constant problem. *Progress of Theoretical and Experimental Physics*, 2018(2):021E01, 2018.
- [7] Micol Benetti, Welber Miranda, Humberto A Borges, Cassio Pigozzo, Saulo Carneiro, and Jailson S Alcaniz. Looking for interactions in the cosmological dark sector. *Journal of Cosmology and Astroparticle Physics*, 2019(12):023, 2019.

- [8] Özgür Akarsu, Suresh Kumar, Shivani Sharma, and Luigi Tedesco. Constraints on a bianchi type i spacetime extension of the standard λ cdm model. *Physical Review D*, 100(2):023532, 2019.
- [9] Paolo Cea. The ellipsoidal universe and the hubble tension. $arXiv\ preprint\ arXiv:2201.04548,\ 2022.$
- [10] Nikita Blinov, Celeste Keith, and Dan Hooper. Warm decaying dark matter and the hubble tension. *Journal of Cosmology and Astroparticle Physics*, 2020(06):005, 2020.
- [11] MC Bento, Orfeu Bertolami, and Anjan A Sen. Generalized chaplygin gas, accelerated expansion, and dark-energy-matter unification. *Physical Review D*, 66(4):043507, 2002.
- [12] Eleonora Di Valentino, Olga Mena, Supriya Pan, Luca Visinelli, Weiqiang Yang, Alessandro Melchiorri, David F Mota, Adam G Riess, and Joseph Silk. In the realm of the hubble tension—a review of solutions. *Classical* and Quantum Gravity, 38(15):153001, 2021.
- [13] Micol Benetti, HA Borges, Cassio Pigozzo, Saulo Carneiro, and JS Alcaniz. Dark sector interactions and the curvature of the universe in light of planck's 2018 data. *Journal of Cosmology and Astroparticle Physics*, 2021(08):014, 2021.
- [14] Kim V Berghaus and Tanvi Karwal. Thermal friction as a solution to the hubble tension. *Physical Review D*, 101(8):083537, 2020.
- [15] Karsten Jedamzik and Levon Pogosian. Relieving the hubble tension with primordial magnetic fields. *Physical Review Letters*, 125(18):181302, 2020.
- [16] Hayley J Macpherson, Paul D Lasky, and Daniel J Price. The trouble with hubble: Local versus global expansion rates in inhomogeneous cosmological simulations with numerical relativity. The Astrophysical Journal Letters, 865(1):L4, 2018.
- [17] Rong-Gen Cai, Zong-Kuan Guo, Li Li, Shao-Jiang Wang, and Wang-Wei Yu. Chameleon dark energy can resolve the hubble tension. *Physical Review D*, 103(12):L121302, 2021.
- [18] Marco San Martín and Carlos Rubio. Hubble tension and matter inhomogeneities: a theoretical perspective. arXiv preprint arXiv:2107.14377, 2021.
- [19] Ryan C Keenan, Amy J Barger, and Lennox L Cowie. Evidence for a 300 megaparsec scale under-density in the local galaxy distribution. The Astrophysical Journal, 775(1):62, 2013.
- [20] Thomas Buchert and Mauro Carfora. Regional averaging and scaling in relativistic cosmology. Classical and Quantum Gravity, 19(23):6109, 2002.

- [21] David L Wiltshire. Cosmic clocks, cosmic variance and cosmic averages. New Journal of Physics, 9(10):377, 2007.
- [22] Thomas Buchert and Jürgen Ehlers. Averaging inhomogeneous newtonian cosmologies. arXiv preprint astro-ph/9510056, 1995.
- [23] Ido Ben-Dayan, Maurizio Gasperini, Giovanni Marozzi, Fabien Nugier, and Gabriele Veneziano. Backreaction on the luminosity-redshift relation from gauge invariant light-cone averaging. *Journal of Cosmology and Astropar*ticle Physics, 2012(04):036, 2012.
- [24] Maurizio Gasperini, G Marozzi, F Nugier, and Gabriele Veneziano. Light-cone averaging in cosmology: Formalism and applications. *Journal of Cosmology and Astroparticle Physics*, 2011(07):008, 2011.
- [25] Giuseppe Fanizza, Maurizio Gasperini, Giovanni Marozzi, and Gabriele Veneziano. Generalized covariant prescriptions for averaging cosmological observables. *Journal of Cosmology and Astroparticle Physics*, 2020(02):017, 2020.
- [26] Jaiyul Yoo and Ruth Durrer. Gauge-transformation properties of cosmological observables and its application to the light-cone average. *Journal of Cosmology and Astroparticle Physics*, 2017(09):016, 2017.
- [27] Mikołaj Korzyński. Covariant coarse graining of inhomogeneous dust flow in general relativity. Classical and quantum gravity, 27(10):105015, 2010.
- [28] R Zalaletdinov. Space-time averages of classical physical fields, annals. eur. acad. sci.(2003) 344 [arxiv: gr-qc/0411004] aa coley, n. pelavas and rm zalaletdinov, cosmological solutions in macroscopic gravity. *Phys. Rev. Lett*, 95:151102, 2005.
- [29] Chris Clarkson, Timothy Clifton, Alan Coley, and Rockhee Sung. Observational constraints on the averaged universe. *Physical Review D*, 85(4):043506, 2012.
- [30] Alan A Coley and N Pelavas. Averaging spherically symmetric spacetimes in general relativity. *Physical Review D*, 74(8):087301, 2006.
- [31] Asta Heinesen, Pierre Mourier, and Thomas Buchert. On the covariance of scalar averaging and backreaction in relativistic inhomogeneous cosmology. *Classical and Quantum Gravity*, 36(7):075001, 2019.
- [32] Thomas Buchert. On average properties of inhomogeneous fluids in general relativity: perfect fluid cosmologies. *General Relativity and Gravitation*, 33(8):1381–1405, 2001.
- [33] Thomas Buchert. On average properties of inhomogeneous fluids in general relativity: dust cosmologies. *General Relativity and Gravitation*, 32(1):105–125, 2000.

- [34] Jelle P Boersma. Averaging in cosmology. *Physical Review D*, 57(2):798, 1998.
- [35] Toyokazu Sekiguchi and Tomo Takahashi. Early recombination as a solution to the H_0 tension. *Phys. Rev. D*, 103:083507, Apr 2021.
- [36] Masumi Kasai and Toshifumi Futamase. A possible solution to the hubble constant discrepancy: Cosmology where the local volume expansion is driven by the domain average density. *Progress of Theoretical and Experimental Physics*, 2019(7):073E01, 2019.
- [37] Masanori Tomonaga and Toshifumi Futamase. A comment on the averaging in an inhomogeneous cosmology and the hubble constant problem. *Progress of Theoretical and Experimental Physics*, 2021(10):103E02, 2021.
- [38] James M Bardeen. Gauge-invariant cosmological perturbations. *Physical Review D*, 22(8):1882, 1980.
- [39] H Kodama and M Sasaki. Multiple fluid density perturbations. Prog. Theor. Phys. Suppl, 78, 1984.