Localized-Diffusive and Ballistic-Diffusive Transitions in Kicked Incommensurate lattice

Hiroaki S. Yamada
Yamada Physics Research Laboratory, Aoyama 5-7-14-205, Niigata 950-2002, Japan

Kensuke S. Ikeda
College of Science and Engineering, Ritsumeikan University, Noji-higashi 1-1-1, Kusatsu 525-8577, Japan
(Dated: January 31, 2023)

By using the kicked Harper model, the effect of dynamical perturbations to the localized and ballistic phases in aperiodic lattice systems is investigated. The transition from the localized phase to diffusive phase via a critical sub-diffusion t^α(t:time) with $0 < \alpha < 1$ is observed. In addition, we first confirmed the existence of the transition from the ballistic phase to the diffusive phase via a critical super-diffusion with $1 < \alpha < 2$.

PACS numbers: 05.45.Mt,71.23.An,72.20.Ee

arXiv:2301.12415v1 [cond-mat.dis-nn] 29 Jan 2023

Introduction.- In 1D tight-binding model with any disorder the dynamical quantum states are always localized [1] But if it is coupled with noisy source such as irregular thermal lattice vibrations, the localization is immediately destroyed and is taken the place by normal diffusion [2, 3]. Even very simple time-dependent harmonic perturbations consisting only a few number of periods may convert the localized wave packet into irreversible diffusion in Anderson model and kicked Anderson model (Anderson map) [4–6]. The characteristic feature of such coherent perturbations is that it realizes irreversible diffusion from the localized state via a critical phase transition we call localization-delocalization transition (LDT).

1D aperiodic lattice systems have the both features of disordered lattice and periodoic lattice. Indeed it has ballistic states and critical states in addition to the localized states. Our question arising here is what would be the effect of the coherent perturbations on the dynamical states of 1D aperiodic lattice systems, i.e.., the localized states, the critical states and the ballistic states

To answer the above question, we investigate the effects of the coherent perturbation by using kicked Harper model (KHM) that facilitate an efficient study of the dynamics in aperiodic lattice. In fact, the KHM shows the above three states intrinsic localized, critical, and ballistic states. The first question is whether the LDT observed for the coherently perturbed random system also exists in the aperiodic lattice system. If this is the case, considering the remarkable duality property of the KHM, a transition from the ballistic to the diffusive state is also expected. The purpose of the present paper is to confirm above conjectures.

As stated above, there are three types of main dynamical states of the quantum wave packet, localized, normal diffusion, and ballistic spread, in the KHM depending upon the potential strength. In that respect, it is the same as the Harper model without the kicks. The phase diagram of the localized/delocalized state in KHM has a nested structure and is quite complicated [7–10]. As for the LDT, its presence was reported by some authors for the KHM driven by the doublly periodic kick or the intensity modulated periodic kicks. It is also feasible to experimentally observe the dynamical LDT through the diffusion of wave packets in the pulsed 1D incommensurate optical lattice such as the KHM [11].

The purpose of this paper is to show realization of the normal diffusion by simple harmonic perturbations via the above two transitions paths, where the harmonic perturbation is polychromatic and is composed of M frequencies mode. We first show the localized state of KHM responds to such perturbation in quite similar ways to that of kicked disordered lattice systems such as the Anderson map [4, 5] and quantum standard map [5, 12–14]: for monochromatic perturbation $M = 1$ the localizaion still remains and it is $M \geq 2$ that the KHM localized state undergoes the LDT with increase in the perturbation strength ϵ. We next show the response of the extended (ballistic) state of KHM to the harmonic perturbation. We report the wave packet dynamics exhibiting the ballistic spreading changes into the normal diffusion when the perturbation strength is increased. The transition occurs through a novel anomalous diffusion as the critical state, which is characterized by a "super" diffusion exponent $1 < \alpha_c < 2$. Hereinafter, this transition is referred to as a ballistic-diffusive transition (BDT) of the wave packet spreading.

Model.- We deal with the dynamically perturbed kicked Harper model (KHM),

$$\begin{aligned} H(t) &= J\sum_{n}^{N}(\hat{b}_n^\dagger \hat{b}_{n+1} + H.C.) \\ &+ 2V\left[1 + f_\epsilon(t)\right]\delta_1(t)\sum_{n}^{N}\cos(2\pi Q n)\hat{b}_n^\dagger \hat{b}_n, \quad (1) \end{aligned}$$

where $\hat{b}_n(\hat{b}_n^\dagger)$ is the creation(annihilation) operator of the n-th site, and N is the lattice size. $\delta_1(t) = \sum_{m\in\mathbb{Z}}\delta(t - m)$. The $Q(= \frac{\sqrt{5}-1}{2})$ is an irrational number. V and $J(= -1)$ denote the potential strength and the hopping energy between adjacent sites, respectively.

The harmonic perturbation $f_\epsilon(t)$ is the sum of harmonic functions

$$f_\epsilon(t) = \epsilon f(t) = \frac{\epsilon}{\sqrt{M}} \sum_i^M \cos(\omega_i t), \quad (2)$$

where M and ϵ are the number of frequency components and the relative strength of the perturbation, respectively. The frequencies $\{\omega_i\}(i = 1, ..., M)$ are taken as mutually incommensurate numbers of order $O(1)$. Note that the long-time average of the total power of the perturbation is normalized to $\overline{f_\epsilon(t)^2} = \epsilon^2/2$.

In the spatial continuous limit, $H(t)$ is given by

$$H_c(t) = 2J\cos(\hat{p}/\hbar) + 2V\cos(2\pi Q\hat{q})[1 + f_\epsilon(t)]\delta_1(t), \quad (3)$$

where $\hat{p}$ and $\hat{q}$ denote the momentum and position operators, respectively. The unperturbed KHM ($\epsilon = 0$) is known to take a localized state ($|V| >> |J|$), critical state ($V = J$), and extended ($|V| << |J|$) state depending on V.

The initial wave packet $< n|\Psi(t = 0) >= \delta_{n,N/2}$ is localized at the site $N/2$, and we calculate the time evolution of the wavefunction $|\Psi(t) >$ using Schr$\acute{}$odinger equation. We monitor the spread of the wave function in the site space by the mean square displacement (MSD): $m_2(t) = \sum_n (n - N/2)^2 \langle|\phi(n,t)|^2\rangle$, where $\phi(n,t) =< n|\Psi(t) >$ is the site representation of the wave function. The number of steps is $10^5 \sim 10^6$. We mainly use the system size $N = 2^{16} - 2^{17}$, and $\hbar = 1/8$.

At the critical state of the transition, an anomalous diffusion $m_2(t) \sim t^\alpha$ characterized by diffusion index α is expected. To observe such a behavior directly we introduce time-dependent the instantaneous diffusion index $\alpha_{ins}(t)$ defined by $\alpha_{ins}(t) := \frac{d\log\overline{m_2(t)}}{d\log t}$, where the locally time-averaged MSD $\overline{m_2(t)}$ is used in order to reduce fluctuation in a shorter time-scale.

Localization-delocalization transition (LDT).- We investigate the delocalization in the KHM with $V = 5$. In the case of $M = 1$ the localization maintains for the harmonic perturbaion, and for $M \geq 2$ the delocalization is observed on a finite time scale, which agrees with the case of kicked Anderson model [5]. Here we focus on the case $M = 3$. In Fig.1(a), the MSD are shown for $M = 3$. They indicate localization when ϵ is small, but the LDT occurs at a certain critical value ϵ_c, and ϵ exceeds ϵ_c, the plot of MSD warping upward (upward deviation) can be seen in the double-logarithmic plots. For $\epsilon > \epsilon_c$ the normal diffusive behavior $m_2 \sim t^1$ appears as $t \to \infty$. Around $\epsilon = \epsilon_c$, the time-dependence of MSD can be approximately described by the sub-diffusive spreading. As a result, the time-dependent MSD displays opposite behaviors for $\epsilon < \epsilon_c$ and for $\epsilon > \epsilon_c$, which can directly be confirmed by $\alpha_{ins}(t)$ vs t plot. As shown in Fig.1(b), in the case of $V >> 1$, we can see that there exist an ϵ_c above which the plot of $\alpha_{ins}(t)$ increases and below which it decreases. The limiting tendency of $\alpha_{ins}(t)$ for $t \to \infty$ seems to be $\alpha_{ins} \to 1$ or $\alpha_{ins}(t) \to 0$ for $\epsilon > \epsilon_c$ or for $\epsilon > \epsilon_c$, respectively. At $\epsilon = \epsilon_c$, $\alpha_{ins}(t)$ fluctuates around $\alpha_{ins}(t) \sim 1/2$, which indicate the anomalous diffusion of $m_2 \propto t^{\alpha_c}$ with $\alpha_c \sim 0.5$.

As shown in Fig.1(c), the scaled MSD $\Lambda(t) = m_2(t)/t^{1/2}$ for various ϵ has trumpet-shaped pattern that suggests the existence of the LDT [5].




FIG. 1: (Color online) (a)The double-logarithmic plots of $m_2(t)$ as a function of time for different values of the perturbation strength ϵ increasing from $\epsilon = 0.0001$(bottom) to $\epsilon = 0.002$(top) in the polychromatically perturbed KHM of $V = 5$ with $M = 3$.. $\hbar = 1/8$. (b)The instantaneous diffusion index $\alpha_{ins}(t)$ for some ϵ. The broken line indicates the critical sub-diffusion line $\alpha_{ins}(t) = \alpha_c = 0.5$ predicted by the scaling theory. (c) The scaled MSD $\Lambda(\epsilon, t) = m_2(t)/t^{1/2}$ as functions of time for increasing perturbation strengths. The results for $\epsilon_c \simeq 0.0009$ are shown in thick blue lines.

As shown in Fig.2, for $M \geq 2$ the $M-$dependence of the critical strength ϵ_c indicates the inverse power-law

$$\epsilon_c \propto \frac{1}{V}\frac{1}{(M-1)}. \quad (4)$$

The same $M-$dependence even for $V = 10$ can be obtained, as seen in Fig.2. This difference in ϵ_c due to V can be interpreted by the Maryland transform in appendix. According to the Maryland transform, the effect of V in the diagonal term is saturated in the region $V > V^*(\equiv 0.38)$ for $\hbar = 1/8$, and the nature of the off-diagonal term depends on ϵV, so the critical value ϵ_c for $V = 10$ can be interpreted as a half of the transition point for $V = 5$.

We plotted in Fig.2 dependence of the substantial critical perturbation strength $\epsilon_c V$ upon the color number, which leads to

$$\epsilon_c V \propto \frac{1}{(M-1)}. \quad (5)$$

An interesting fact is that the result means that $\epsilon_c V$(more precisely $\epsilon_c V/\hbar$) does no longer dependent upon V for $|V| \gg 1$. It implies the effect of the quasi-periodic perturbation on the localization effect saturates as $|V|$ increases. This can immediately be understood by the Maryland transformed scheme of the quasi-energy eigenvalue equation for the time-periodic Hamiltonian (1), which is shown in Appendix. It is interesting that the LDT occurs at a very small relative amplitude of perturbation, ie, $\epsilon_c \sim 10^{-3}$, in contrast to the BDT to be given later.




FIG. 2: (Color online) The critical perturbation strength $V\epsilon_c$ as a function of $(M-1)$ for $V=5$ and $V=10$ in the KHM. The black solid line with slope -1 is shown as a reference.

Ballistic-Diffusive transition (BDT).- Next, we study the effect of the harmonic perturbation on the extended states in the KHM which occurs in the case $V \ll 1$. In such a regime the initial wave packet spreads ballistically as $m_2(t) \sim t^2$ without the harmonic perturbation. In view of such dynamical behavior, we henceforth refer to the extended state as a ballistic state. In the case of $M=1$ the ballistic spreading does not change, and we investigate the case of $M \geq 2$ in accordance with the case of LDT. We take the case of $V=0.2$ and $M=3$ as the typical case in accordance with the case where LDT was observed in Fig.1. The time-dependence of MSD for various values of ϵ is displayed in Fig.3(a). When ϵ is small, the growth of $m_2(t)$ slightly deviates from ballistic spreading in the initial time regime, but it reaches to ballistic increase as $t \to \infty$. However as ϵ is taken large enough, it gradually approaches to normal diffusion obeying $m_2(t) \sim t^1$ and finally it increases as $m_2(t) \sim t$ even from the initial stage.

The above observation suggests a transition from the ballistic spreading to a normal diffusion. We confirm this directly by the $\alpha_{ins}(t)$ vs t-plot shown in Fig.3(b). In all cases $\alpha_{ins}(t)$ increases in the initial stage. For ϵ large enough the increase continues to $t \to \infty$, whereas $\alpha_{ins}(t)$ decreases as $t \to \infty$ for ϵ small enough . There exist a certain $\epsilon = \epsilon_b$ at which $\alpha_{ins}(t)$ approaches to a constant value $\alpha_b \sim 1.73$, indicating the presence of the




FIG. 3: (Color online) (a)The double-logarithmic plots of $m_2(t)$ as a function of time for different values of the perturbation strength ϵ increasing from $\epsilon = 0.02$(top) to $\epsilon = 0.8$(bottom) in the polychromatically perturbed KHM of $V=0.2$ with $M=3$. $\hbar = 1/8$. (b)The instantaneous diffusion index $\alpha_{ins}(t)$. The broken line indicates the critical super-diffusion line $\alpha_{ins}(t) = \alpha_c = 1.73$. (c) The scaled MSD $\Lambda(t) = m_2(t)/t^{1.73}$ as functions of time for increasing strengths ϵ. The results for $\epsilon_b \simeq 0.10$ are shown in thick blue lines.

asymptotic super-diffusion $m_2 \propto t^{\alpha_c}$ as the critical state.

Figure 4 is an estimate of the diffusion coefficient D in the normal diffusion region ($\epsilon > \epsilon_b$). The time-dependence of MSD shows the normal diffusion for $\epsilon > \epsilon_b$ and the diffusion coefficient decreases by increasing ϵ. On the other hand, as seen in the previous part, for $\epsilon > \epsilon_c$ $m_2(t)$ gradually approaches to the normal diffusion for $t \to \infty$. Here, in the KHM the $\epsilon-$dependence of the diffusion coefficient in the normal diffusion region in a wide region of $\epsilon(> \max\{\epsilon_c, \epsilon_b\})$ is observed in Fig.4.

In the localized case ($V=5$), the diffusion coefficient increases with an increase of ϵ. As ϵ is increased far beyond ϵ_c and the decrease saturates at a certain level of the order of $O(1)$. On the other hand, in the extended case ($V=0.2$), the diffusion coefficient decreases monotonically as ϵ increases and for ϵ far beyond ϵ_b, D falls down to the same level as the localized cases for $\epsilon >> 1$. This fact implies that for any V there always exist a critical point ϵ_c beyond which an irreversible diffusion switches on, whose diffusion coefficient converges to the same level.

Summary and discussion.- We investigated the dynamical property of the initially localized wave packet in coherently perturbed KHM. In the localized cases ($V >> 1$), the LDT appeared with increasing the per-




FIG. 4: (Color online) Diffusion coefficient D as a function of ϵ of the localized ($V = 5$) and extended ($V = 0.2$) states in the perturbed KHM. $\hbar = 1/8$.

turbation strength ϵ. The critical value M_c which appear the LDT is $M_c = 2$. It also describes the localized and delocalized transitions in the multidimensional Anderson model that corresponds to the 1D Anderson map and kicked rotors with the quasi-periodic perturbation. If $M + 1$ can be identified with the spatial dimension d, the existence of the LDT is a qualitatively consistent result with those of the LDT in $d-$dimensional Anderson model [15–18]. It was also shown that the ballistic dynamics without the perturbation makes transition to the normal diffusion with increase in the perturbation strength. It will be interesting to see if similar results presented in this paper can be obtained for the original Harper model with localized, critical and delocalized states as well as the KHM,

As a result, it was shown that whether the unperturbed state is localized or ballistic, the increase in the strength of the coherent harmonic perturbation causes a transition to the normal diffusion. The asymptotic diffusive behavior is expected as a generic feature of decoherence, when noise is not introduced into a system. The observed facts presented above implies that in order to induce quantum diffusion, which is apparently an irreversible quantum dynamics, we need neither spatial nor temporal stochasticity and only spatial and temporal aperiodicity is sufficient. This work may give some hints leading to a deeper understanding of dynamical localization and quantum diffusion in quasi-periodic systems. It also provides insight into the control of localized and delocalized states by coherent perturbations in the Floquet engineering.

Acknowledgments: This work is partly supported by Japanese people's tax via JPSJ KAKENHI 15H03701, and the authors would like to acknowledge them. They are also very grateful to Dr. T.Tsuji and Koike memorial house for using the facilities during this study. The author (H.Y.) would like to acknowledge the hospitality of the Physics Division of the Nippon Dental University at Niigata, where part of this work was completed.

Appendix: Maryland transform.- We can regard the time-dependent harmonic perturbation $f_\epsilon(t)$ as the dynamical degrees of freedom. To show this we introduce the classically canonical action-angle operators ($\hat{J}_j = -i\hbar\frac{\partial_j}{\partial_j\phi_j}, \phi_j$) representing the harmonic perturbation as the linear modes, and we call them the color modes. Each quantum oscillator has the action eigenstates $|n_j >$ with the action eigenvalue $J_j = n_j\hbar$ (n_j :integer) and the energy $n_j\hbar\omega_j$, where $\hat{J}_j|m_j\rangle = m_j\hbar|m_j\rangle (m_j \in \mathbb{Z})$. Thus the system (1) is regarded as a quantum system of $(M + 1)$-degrees of freedom spanned by the quantum states $|n > \prod_{j=1}^{M} |n_j >$. Then the Hamiltonian $\tilde{H}_{kick}$ that include the color modes becomes

$$\tilde{H}_{kick}(\hat{p}, \hat{q}, \{\hat{J}_j\}, \{\hat{\phi}_j\}) = 2\cos(\hat{p}/\hbar) + 2V\cos(2\pi Q\hat{q})\left[1 + \frac{\epsilon}{\sqrt{M}}\sum_j^M \cos\phi_j\right]\delta_1(t) + \sum_{j=1}^{M}\omega_j\hat{J}_j, \quad (6)$$

where γ and $|u\rangle$ are the quasi-eigenvalue and quasi-eigenstate. Here, if the eigenstate representation of $\hat{J}_j$ is used, we can obtain the following $(M + 1)-$dimensional tight-binding expression by the Maryland transform [5, 19]:

$$D(n, \{m_j\})u(n, \{m_j\}) + \sum_{n', \{m_j^{'}\}} \langle n, \{m_j\}|\hat{t}_{KHM}|n', \{m_j^{'}\}\rangle u(n', \{m_j^{'}\}) = 0, \quad (7)$$

where $\{m_j\} = (m_1, ..., m_M)$. Here the diagonal term is

$$D(n, \{m_j\}) = \tan\left[\frac{2V\cos(2\pi Qn) + \hbar\sum_j^M m_j\omega_j}{2\hbar} - \frac{\gamma}{2}\right], \quad (8)$$

and the $\hat{t}_{KHM}$ of the off-diagonal term is

$$\hat{t}_{KHM} = i\frac{e^{-i\frac{\epsilon 2V}{\sqrt{M}}\cos(2\pi Q\hat{q})(\sum_j^M\cos\phi_j)/\hbar} - e^{i2\cos(\hat{p}/\hbar)/\hbar}}{e^{-i\frac{\epsilon 2V}{\sqrt{M}}\cos(2\pi Q\hat{q})(\sum_j^M\cos\phi_j)/\hbar} + e^{i2\cos(\hat{p}/\hbar)/\hbar}}. \quad (9)$$

The effect of V in the on-site-potential $D(n, \{m_j\})$ is saturated for $|V/\hbar| >> \pi$, since the aperiodicity is expressed through the tangent function. The off-diagonal component controlling inter-site-hopping is governed by a single parameter $\epsilon V/\hbar$. This is consistent with the fact that when $|V|$ is sufficiently large, it is the combined parameter ϵV (more precisely $\epsilon V/\hbar$) that governs the feature of the transition.

[1] K. Ishii, Localization of Eigenstates and Transport Phenomena in the One-Dimensional Disordered System, Prog. Theor. Phys. Suppl. **53**, 77 (1973).

[2] H. Haken and P. Reineker, The coupled coherent and incoherent motion of excitons and its influence on the line shape of optical absorption, Z. Phys. **249**, 253 (1972);

[3] H. Haken and G. Strobl, An exactly solvable model for coherent and incoherent exciton motion, Z. Phys. **262**, 135 (1973).

[4] H.S.Yamada, F. Matsui and K.S.Ikeda, Scaling Properties of Dynamical Localization in Monochromatically Perturbed Quantum Maps: standard map and Anderson map, Phys.Rev.E **97**, 012210(2018).

[5] H.S.Yamada, and K.S. Ikeda, Critical phenomena of dynamical delocalization in quantum maps: Standard map and Anderson map, Phys.Rev.E **101**, 032210(2020).

[6] H.S.Yamada, and K.S. Ikeda, Localization and delocalization properties in quasi-periodically-driven one-dimensional disordered systems, Phys.Rev.E **105**, 054201(2022).

[7] R. Artuso, G. Casati, F. Borgonovi, L. Rebuzzini and I. Guarneri, Fractal and dynamical properties of the kicked Harper model, International Journal of Modern Physics B **8**, 207-235 (1994).

[8] T. Prosen, I. I Satija, N. R. Shah, Dimer Decimation and Intricately Nested Localized-Ballistic Phases of Kicked Harper, Phys.Rev.Lett.**87**, 066601(2001).

[9] A. R. Kolovsky and H. J. Korsch, Quantum diffusion in a biased kicked Harper system, Phys. Rev. E **68**, 046202(2003).

[10] B. Lévi and B. Georgeot, Quantum computation of a complex system: The kicked Harper model, Phys. Rev. E **70**, 056218(2004).

[11] S. Sarkar, S. Paul, C. Vishwakarma, S. Kumar, G. Verma, M. Sainath, U. D. Rapol, and M. S. Santhanam, Nonexponential Decoherence and Subdiffusion in Atom-Optics Kicked Rotor, Phys. Rev. Lett.**118**, 174101 (2017).

[12] G. Casati, I. Guarneri, and D. L. Shepelyansky, Phys. Rev. Lett., Anderson Transition in a One-Dimensional System with Three Incommensurate Frequencies, **62**, 345 (1989).

[13] J. Chabe, G. Lemarie, B. Gremaud, D. Delande, P. Szriftgiser, and J. C. Garreau, Experimental Observation of the Anderson Metal-Insulator Transition with Atomic Matter Waves, Phys. Rev. Lett. **101**, 255702(2008).

[14] M. Lopez, J.-F. Clement, G. Lemarie, D. Delande, P. Szriftgiser, and J. C. Garreau, Phase diagram of the Anderson transition with atomic matter waves, New J. Phys. **15**, 065013 (2013).

[15] P. W. Anderson, Absence of diffusion in certain random lattices, Phys. Rev. **109**, 1492-1505 (1958).

[16] E.Abrahams, P.W.Anderson, D.C.Licciardello, and T.V.Ramakrishnan, Scaling Theory of Localization: Absence of Quantum Diffusion in Two Dimensions, Phys. Rev.Lett. **42**, 673 (1979).

[17] L.M.Lifshiz, S.A.Gredeskul and L.A.Pastur, *Introduction to the theory of Disordered Systems*, (Wiley, New York,1988).

[18] E. Abrahams (Editor), *50 Years of Anderson Localization*, (World Scientific 2010).

[19] S. Fishman, D. R. Grempel, and R. E. Prange, Chaos, Quantum Recurrences, and Anderson Localization, Phys. Rev. Lett. **49**, 509 (1982).