Measuring Lorentz Violation in Weak Gravity Fields

1

Zonghao Li

Department of Physics, Indiana University, Bloomington, IN 47405, USA

Many new linearized coefficients for Lorentz violation are discovered in our recent work on the construction of a generic Lorentz-violating effective field theory in curved spacetime. The new coefficients can be constrained by experiments in weak gravity fields. In this work, we compare experiments in different gravitational potentials and study three types of gravity-related experiments: free-fall, gravitational interferometer, and gravitational bound-state experiments. First constraints on the new coefficients for Lorentz violation are extracted from those experiments.

1. Lorentz violation in gravity

In recent years, Lorentz violation has been a popular topic in the search for physics beyond the Standard Model (SM) and General Relativity (GR). The Standard-Model Extension (SME)^{1,2} has been widely used as a comprehensive framework to study Lorentz violation in the context of effective field theory. The minimal terms in the Lagrange density of the SME in curved spacetime were constructed by Kostelecký in 2004,² and the nonminimal terms were systematically constructed in our recent work.^{3,4} The linearizations of those terms in weak gravity fields were also obtained.⁵ The present contribution to the proceedings of CPT'22 studies the experimental implications of the linearized terms with a focus on matter–gravity couplings in weak gravity fields. This work is based on the results in Ref. [5].

2. Potential-dependent experiments

An interesting implication of the linearized Lagrange density constructed in our recent work⁵ is that the measured SME coefficients for Lorentz violation can depend on the gravitational potential of the laboratory. Coefficients for Lorentz violation have been measured in many experiments under the assumption that spacetime is flat.⁶ However, these experiments are typically performed at different elevations and hence at different gravitational potentials, so the SME coefficients can depend on the potentials. Taking the *b*-type coefficients as an example, we know that the term in the Lagrange density containing the minimal b^{κ} coefficient is $\mathcal{L} \supset b^{\kappa} \overline{\psi} \gamma_{\kappa} \psi$ in flat spacetime.¹ Adding couplings with the gravitational field, we can write the generalization of the term in a weak gravity field as⁵

$$\mathcal{L} \supset (b_{\text{asy}}^{\kappa} + (b^{\text{L}})^{\kappa\mu\nu} h_{\mu\nu} + \cdots) \overline{\psi} \gamma_{\kappa} \psi \equiv b_{\text{expt}}^{\kappa} \overline{\psi} \gamma_{\kappa} \psi, \qquad (1)$$

where $h_{\mu\nu} \equiv g_{\mu\nu} - \eta_{\mu\nu}$ is the linearized gravitational field. In a nonrelativistic weak gravity field, $h_{\mu\nu}$ can be approximated by $h_{00} \approx -2\phi$, $h_{0j} \approx 0$, and $h_{jk} \approx -2\phi \delta_{jk}$, where ϕ is the gravitational potential. The actual b^{κ} coefficients measured in experiments should be the effective value

$$b_{\text{expt}}^{\kappa} = b_{\text{asy}}^{\kappa} + (b^{\text{L}})^{\kappa\mu\nu} h_{\mu\nu} + \dots \approx b_{\text{asy}}^{\kappa} - 2(b^{\text{L}})^{\kappa\Sigma\Sigma}\phi, \qquad (2)$$

where $\Sigma\Sigma$ in the index means a summation over space and time indices in the Sun-centered frame,⁶ i.e., $(b^{L})^{\kappa\Sigma\Sigma} = (b^{L})^{\kappa TT} + (b^{L})^{\kappa XX} + (b^{L})^{\kappa YY} + (b^{L})^{\kappa ZZ}$. We see that this effective coefficient depends on the gravitational potential. Moreover, the combination $(b^{L})^{\kappa\Sigma\Sigma}$ can be constrained by comparing experiments measuring b^{κ} at different elevations.

As an example, an experiment in Seattle constrained the b_e^X coefficient, a combination of b^{κ} and other SME coefficients in the electron sector,⁶ as $|\tilde{b}_e^X| < 3.7 \times 10^{-31} \text{ GeV.}^7$ Another experiment in Taiwan measured the same combination and got $|\tilde{b}_e^X| < 3.1 \times 10^{-29}$ GeV.⁸ The results are obtained at different elevations with different gravitational potentials, so we can compare them to get a constraint on the linearized coefficient $\tilde{b}_e^{X\Sigma\Sigma}$ as $|\tilde{b}_e^{X\Sigma\Sigma}| < 3.2 \times 10^{-15} \text{ GeV}$. Similar analyses can be done for other coefficients using experimental data summarized in the data tables⁶ for the SME.⁵ More experiments can be done to measure SME coefficients in different gravitational potentials, and those can be used to constrain the linearized coefficients.

3. Free-fall experiments

2

Aside from comparing results in different experiments, we can also constrain the linearized coefficients by single gravity-related experiments. To better analyze those experiments, we derived the nonrelativistic Hamiltonian from the linearized Lagrange density. The Hamiltonian can be written as

$$H = H_0 + H_{\phi} + H_{\sigma\phi} + H_g + H_{\sigma g} + \dots,$$
(3)

where H_0 is the Hamiltonian without background fields, and other components are the corrections from background fields. Components with subscript σ are spin dependent, and those without are spin independent. The exact terms in the components can be found in Ref. [5]. The Hamiltonian can modify the gravitational acceleration experienced by a dynamical system on the Earth's surface. The spin-dependent components permits us to study spin–gravity couplings in the SME framework for the first time. In this section, we use free-fall experiments to test the spin–gravity couplings.

One experiment⁹ compares the effective gravitational accelerations of two isotopes of strontium atoms, the spin-zero bosonic ⁸⁸Sr and the spin-9/2 fermionic ⁸⁷Sr. Unpolarized ⁸⁷Sr atoms were used there, so if effective gravitational accelerations depend on spin orientations, the measured gravitational accelerations of ⁸⁷Sr atoms should span a broader range than those of ⁸⁸Sr atoms. They found no such effect to a sensitivity of 10^{-7} . Analyzing the result in our framework, we get bounds on nonrelativistic SME coefficients as

$$\left| (k_{\sigma\phi}^{\rm NR})_n^Z \right| < 1 \times 10^{-4} \text{ GeV},$$
$$\left| (k_{\sigma\phi pp}^{\rm NR})_n^{ZJJ} - 0.4 (k_{\sigma\phi pp}^{\rm NR})_n^{ZZZ} \right| < 5 \times 10^{-2} \text{ GeV}^{-1}, \tag{4}$$

where the subscript n means the coefficients are for neutrons, and repeated J indices mean a summation over special coordinates J = X, Y, Z in the Sun-centered frame.

Another experiment¹⁰ compares the gravitational acceleration experienced by ⁸⁷Rb atoms with different spin orientations. They found no difference to a sensitivity of 10^{-7} . This can be translated to constraints on nonrelativistic coefficients as

$$\left| (k_{\sigma\phi}^{\mathrm{NR}})_p^Z - 0.6 (k_{\sigma\phi}^{\mathrm{NR}})_e^Z \right| < 2 \times 10^{-5} \text{ GeV},$$
$$\left| (k_{\sigma\phi pp}^{\mathrm{NR}})_p^{ZJJ} + 0.3 (k_{\sigma\phi pp}^{\mathrm{NR}})_p^{JJZ} \right| < 7 \times 10^{-3} \text{ GeV}^{-1}, \tag{5}$$

where the subscripts p and e mean proton and electron flavor, respectively.

Another type of interesting free-fall experiment is to compare the falls of hydrogen H and antihydrogen \overline{H} . This can provide insights on CPT symmetry. Several groups have been designing experiments for that.¹¹ A detailed theoretical analysis of the falls in our framework can be found in Ref. [5]. We expect new results from those experiments in the near future.

4. Gravitational interferometer experiments

Our nonrelativistic Hamiltonian can also modify the gravity-induced phase shift in gravitational interferometer experiments. In this section, we analyze several interferometer experiments with neutrons and use them to extract bounds on the nonrelativistic coefficients.

4

The first gravitational interferometer experiment was performed by Colella, Overhauser, and Werner (COW).¹² They used Bragg diffraction to split a coherent neutron beam into two paths at different heights and measured the relative gravity-induced phase shift between the two paths. Unpolarized neutron beams were used in the experiment, so it is mainly sensitive to the spin-independent terms in our Hamiltonian.

The effective gravitational acceleration measured in the original COW experiment attains an accuracy of 10%. From this, we deduce a constraint on a nonrelativistic coefficient as

$$(k_{\phi}^{\rm NR})_n < 1 \times 10^{-1} \,\,{\rm GeV},$$
 (6)

where $(k_{\phi}^{\text{NR}})_n$ is a spin-independent coefficient in the neutron sector. More recent versions of the COW experiment can improve this result.⁵

The next type of interferometer experiments we consider is the OffSpec experiment, which uses polarized nonrelativistic neutron beams and splits the beams by magnetic fields.¹³ This is sensitive to spin–gravity couplings. The experiment measured the effective gravitational acceleration to an accuracy of 2.5%. After some analysis,⁵ we get the constraint

$$\left| (k_{\phi}^{\text{NR}})_n + (k_{\sigma\phi}^{\text{NR}})_n^j \hat{s}^j \right| < 2.5 \times 10^{-2} \text{ GeV},$$
 (7)

where $(k_{\phi}^{\text{NR}})_n$ and $(k_{\sigma\phi}^{\text{NR}})_n^j$ are coefficients in the neutron sector, and \hat{s}^j is the initial polarization direction of the neutron beams. We expect a more precise result to be obtained from a more detailed analysis of the experiment. Also, our understanding of spin–gravity couplings and Lorentz violation can be further improved by future experiments using similar setups with the OffSpec experiment. For example, the coefficients $(k_{\sigma g}^{\text{NR}})_n^{jk}$ in our Hamiltonian can be constrained by comparing the phase shifts between horizontally split neutron beams with different spin orientations.

5. Gravitational bound-state experiments

Another application of the nonrelativistic Hamiltonian concerns gravitational bound-state experiments,^{15,16} where the bounds states of neutrons in the Earth's gravitational field are measured. In those experiments, our nonrelativistic Hamiltonian can modify the energy states by changing the potential experienced by the neutrons. Specifically, the spin-independent terms in the Hamiltonian shift the energy levels, and the spin-dependent terms split the energy levels. The first gravitation bound-state experiment¹⁵ measured the critical heights of the bound states, which are related to the energy levels. The precision of the measurement is around 10%. A later experiment¹⁶ improved the precision to around 0.3% by measuring the transition frequencies between different energy levels. From those results, constraints on nonrelativistic SME coefficients are found to be^{5,17}

$$\left| (k_{\phi}^{\mathrm{NR}})_n \right| < 1 \times 10^{-3} \text{ GeV},$$
$$\left/ \overline{\left[(k_{\sigma\phi}^{\mathrm{NR}})_n^J \right]^2} < 8 \times 10^{-3} \text{ GeV},$$
(8)

5

where $(k_{\phi}^{\text{NR}})_n$ and $(k_{\sigma\phi}^{\text{NR}})_n^J$ are nonrelativistic coefficients in the neutron sector, and the square implies a summation over J = X, Y, Z in the Suncentered frame. We expect these results to be improved by more precise future measurements.

Acknowledgments

This work was supported in part by the U.S. Department of Energy and by the Indiana University Center for Spacetime Symmetries (IUCSS).

References

- D. Colladay and V.A. Kostelecký, Phys. Rev. D 55, 6760 (1997); Phys. Rev. D 58, 116002 (1998).
- 2. V.A. Kostelecký, Phys. Rev. D 69, 105009 (2004).
- 3. V.A. Kostelecký and Z. Li, Phys. Rev. D 99, 056016 (2019).
- 4. V.A. Kostelecký and Z. Li, Phys. Rev. D 103, 024059 (2021).
- 5. V.A. Kostelecký and Z. Li, Phys. Rev. D 104, 044054 (2021).
- Data Tables for Lorentz and CPT Violation, V.A. Kostelecký and N. Russell, 2023 edition, arXiv:0801.0287v16.
- 7. B.R. Heckel et al., Phys. Rev. D 78, 092006 (2008).
- 8. L.-S. Hou et al., Phys. Rev. Lett. 90, 201101 (2003).
- 9. M.G. Tarallo et al., Phys. Rev. Lett. 113, 023005 (2014).
- 10. X.C. Duan et al., Phys. Rev. Lett. 117, 023001 (2016).
- S. Aghion et al., Nat. Commun. 5, 4538 (2014); C. Amole et al., Phys. Rev. Lett. 112, 121102 (2014); P. Indelicato et al., Nat. Commun. 4, 1787 (2013).
- R. Colella, A.W. Overhauser, and S.A. Werner, Phys. Rev. Lett. 34, 23 (1975).
- 13. V.-O. de Haan et al., Phys. Rev. A 89, 063611 (2014).
- 14. V. de Haan et al., Phys. Rev. A 89, 063661 (2014).
- 15. V.V. Nesvizhevsky et al., Nature 415, 297 (2002).
- 16. G. Cronenberg et al., Nat. Phys. 14, 1022 (2018).
- 17. A.N. Ivanov, M. Wellenzohn, and H. Abele, Phys. Lett. B 822 136640 (2021).