

# Determining the normalization of the quantum field theory vacuum, with implications for quantum gravity

Philip D. Mannheim  
*Department of Physics, University of Connecticut,  
 Storrs, CT 06269, USA  
 philip.mannheim@uconn.edu*

(Dated: February 27 2023)

In a standard quantum field theory the norm  $\langle \Omega | \Omega \rangle$  of the vacuum state is taken to be finite. In this paper we provide a procedure, based on constructing an equivalent wave mechanics, for determining whether or not  $\langle \Omega | \Omega \rangle$  actually is finite. We provide an example based on a second-order plus fourth-order scalar field theory, a prototype for quantum gravity, in which it is not. In this example the Minkowski path integral with a real measure diverges though the Euclidean path integral does not. Thus in this example contributions from the Wick rotation contour cannot be ignored. Since  $\langle \Omega | \Omega \rangle$  is not finite, use of the standard Feynman rules is not valid. And while these rules not only lead to states with negative norm, they in fact lead to states with infinite negative norm. However, if the fields in that theory are continued into the complex plane, we show that then there is a domain in the complex plane known as a Stokes wedge in which one can define an appropriate time-independent, positive and finite inner product, viz. the  $\langle L | R \rangle$  overlap of left-eigenstates and right-eigenstates of the Hamiltonian; with the vacuum state then being normalizable, and with there being no states with negative or infinite  $\langle L | R \rangle$  norm. In this Stokes wedge it is the Euclidean path integral that diverges while the Minkowski path integral does not. The concerns that we raise in this paper only apply to bosons since the matrices associated with their creation and annihilation operators are infinite dimensional. Since the ones associated with fermions are finite dimensional, the fermion theory vacuum is automatically normalizable. We discuss some general implications of our results for quantum gravity studies, and show that they are relevant to the construction of a consistent, unitary and renormalizable quantum theory of gravity.

## I. THE HIDDEN ASSUMPTION OF QUANTUM FIELD THEORY

Consider a free relativistic neutral scalar field with action

$$I_S = \int d^4x \frac{1}{2} [\partial_\mu \phi \partial^\mu \phi - m^2 \phi^2], \quad (1.1)$$

and wave equation, Hamiltonian, and equal time commutation relation of the form

$$\begin{aligned} [\partial_\mu \partial^\mu + m^2] \phi &= 0, \\ H &= \int d^3x \frac{1}{2} [\dot{\phi}^2 + \bar{\nabla} \phi \cdot \bar{\nabla} \phi + m^2 \phi^2], \\ [\phi(\bar{x}, t), \dot{\phi}(\bar{x}', t)] &= i \delta^3(\bar{x} - \bar{x}'). \end{aligned} \quad (1.2)$$

With  $\omega_k = +(\bar{k}^2 + m^2)^{1/2}$  solutions to the wave equation obey

$$\phi(\bar{x}, t) = \int \frac{d^3k}{\sqrt{(2\pi)^3 2\omega_k}} [a(\bar{k}) e^{-i\omega_k t + i\bar{k} \cdot \bar{x}} + a^\dagger(\bar{k}) e^{i\omega_k t - i\bar{k} \cdot \bar{x}}], \quad (1.3)$$

and with  $[a(\bar{k}), a^\dagger(\bar{k}')] = \delta^3(\bar{k} - \bar{k}')$  the Hamiltonian is given by

$$H = \frac{1}{2} \int d^3k [\bar{k}^2 + m^2]^{1/2} [a^\dagger(\bar{k}) a(\bar{k}) + a(\bar{k}) a^\dagger(\bar{k})]. \quad (1.4)$$

Given (1.4) we can introduce a no-particle state  $|\Omega\rangle$  that obeys  $a(\bar{k})|\Omega\rangle = 0$  for each  $\bar{k}$ , and can identify it as the ground state of  $H$ . This procedure does not specify the value of  $\langle \Omega | \Omega \rangle$ .

For the theory the associated c-number propagator obeys

$$(\partial_t^2 - \bar{\nabla}^2 + m^2)D(x) = -\delta^4(x), \quad (1.5)$$

so that

$$D(x) = \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot x}}{(k^2 - m^2 + i\epsilon)}. \quad (1.6)$$

If we identify the propagator as a vacuum matrix element of q-number fields, viz.

$$D(x) = -i\langle\Omega|T[\phi(x)\phi(0)]|\Omega\rangle, \quad (1.7)$$

then use of the equal time commutation relation gives

$$(\partial_t^2 - \bar{\nabla}^2 + m^2)(-i)\langle\Omega|T[\phi(x)\phi(0)]|\Omega\rangle = -\langle\Omega|\Omega\rangle\delta^4(x). \quad (1.8)$$

Comparing with (1.5) we see that we can only identify  $D(x)$  as the matrix element  $-i\langle\Omega|T[\phi(x)\phi(0)]|\Omega\rangle$  if the vacuum is normalized to one, viz.  $\langle\Omega|\Omega\rangle = 1$ . Now if the normalization of the vacuum is finite we of course can always rescale it to one. However, that presupposes that the normalization of the vacuum is not infinite. We are not aware of any proof in the literature that the normalization of the vacuum is not infinite (either in this particular case or in general), and taking it to be finite is a hidden assumption. So in this paper we shall present a procedure for determining whether the normalization of the vacuum state is finite or infinite. The procedure is based on generalizing to quantum field theory what we know from quantum mechanics.

This paper is organized as follows. In Sec. II we discuss the quantum-mechanical harmonic oscillator. In Sec. III we present an example, the above second-order derivative scalar field theory, in which  $\langle\Omega|\Omega\rangle$  is finite. And in Secs. IV to VI we present an example, a second-order plus fourth-order derivative scalar field theory, in which  $\langle\Omega|\Omega\rangle$  is not finite. In Sec. VII we show that by continuing the dynamical variables into the complex plane we can construct a second-order plus fourth-order derivative scalar field theory inner product that is finite. In Secs. VIII and XII we provide an analogous path integral analysis of second-order plus fourth-order derivative theories and reach the same conclusions. In Sec. IX we discuss the effect of interactions. In Sec. X we discuss fermion field theories. In Secs. XI and XII we discuss the relevance of our study to the construction of a consistent, unitary and renormalizable quantum gravity theory. In Sec. XIII we make some final comments.

## II. THE QUANTUM-MECHANICAL SIMPLE HARMONIC OSCILLATOR

For a simple harmonic oscillator with Hamiltonian  $H = \frac{1}{2}[p^2 + q^2]$  and commutator  $[q, p] = i$ , there are two sets of bases, the wave function basis and the occupation number space basis. The wave function basis is obtained by setting  $p = -i\partial/\partial q$  in  $H$  and then solving the Schrödinger wave equation  $H\psi(q) = E\psi(q)$ . In this way we obtain a ground state with energy  $E_0 = \frac{1}{2}$  and wave function  $\psi_0(q) = e^{-q^2/2}$ . For occupation number space we set  $q = (a + a^\dagger)/\sqrt{2}$  and  $p = i(a^\dagger - a)/\sqrt{2}$ . This yields  $[a, a^\dagger] = 1$  and  $H = a^\dagger a + 1/2$ . We introduce a no-particle state  $|\Omega\rangle$  that obeys  $a|\Omega\rangle = 0$ , with  $|\Omega\rangle$  being the occupation number space ground state with energy  $E_0 = \frac{1}{2}$ . However, in and of itself this does not fix the norm  $\langle\Omega|\Omega\rangle$  of the no-particle state or oblige it to be finite.

To fix the  $\langle\Omega|\Omega\rangle$  norm we need to relate the ground states of the two bases. With  $a = (q + ip)/\sqrt{2}$  we set

$$\langle q|a|\Omega\rangle = \frac{1}{\sqrt{2}} \left( q + \frac{\partial}{\partial q} \right) \langle q|\Omega\rangle = 0, \quad (2.1)$$

and find that  $\langle q|\Omega\rangle = e^{-q^2/2}$ . We thus identify  $\psi_0(q) = \langle q|\Omega\rangle$ . We now calculate the standard Dirac norm for vacuum, and obtain

$$\langle\Omega|\Omega\rangle = \int_{-\infty}^{\infty} dq \langle\Omega|q\rangle \langle q|\Omega\rangle = \int_{-\infty}^{\infty} dq \psi_0^*(q) \psi_0(q) = \int_{-\infty}^{\infty} dq e^{-q^2} = \sqrt{\pi}. \quad (2.2)$$

We thus establish that the Dirac norm of the no-particle state is finite. And on setting  $\psi_0(q) = e^{-q^2/2}/\pi^{1/4}$  we normalize it to one. That we are able to do this is because we know the form of the wave function  $\psi_0(q)$ .

While this procedure is both straightforward and familiar, it works because both the wave function basis approach and occupation number basis approach have something in common, namely that they are both based on an infinite number of degrees of freedom. For the occupation number basis we can represent the creation and annihilation operators as infinite-dimensional matrices labeled by  $|\Omega\rangle$ ,  $a^\dagger|\Omega\rangle$ ,  $a^2|\Omega\rangle$  and so on. For the wave function basis the coordinate  $q$  is a continuous variable that varies between  $-\infty$  and  $\infty$ . The two sets of bases are both infinite dimensional, one discrete and the other continuous. The advantage of the continuous basis is that it enables us to express the normalization of the vacuum state as an integral with an infinite range, an integral that is then either finite or infinite.

### III. THE QUANTUM FIELD THEORY OSCILLATOR

In the quantum field theory case we do not know the form of the wave function solutions to  $H|\psi\rangle = E|\psi\rangle$ , since we cannot realize the canonical commutator given in (1.2) as a differential relation. Specifically, we cannot satisfy (1.2) by setting  $\dot{\phi}(\bar{x}, t)$  equal to  $-i\partial/\partial\phi(\bar{x}, t)$  (though we could introduce a functional derivative  $\dot{\phi}(\bar{x}, t) = -i\delta/\delta\phi(\bar{x}, t)$ ).

However, we can express the Hamiltonian in terms of creation and annihilation operators. So what we can then do is reverse engineer what we did in the quantum-mechanical case. We thus introduce

$$a(\bar{k}) = \frac{1}{\sqrt{2}}[q(\bar{k}) + ip(\bar{k})], \quad a^\dagger(\bar{k}) = \frac{1}{\sqrt{2}}[q(\bar{k}) - ip(\bar{k})], \quad (3.1)$$

so that

$$\begin{aligned} [q(\bar{k}), p(\bar{k}')] &= i\delta^3(\bar{k} - \bar{k}'), \quad H = \frac{1}{2} \int d^3k [\bar{k}^2 + m^2]^{1/2} [p^2(\bar{k}) + q^2(\bar{k})], \\ \phi(\bar{x}, t) &= \frac{1}{\sqrt{2}} \int \frac{d^3k}{\sqrt{(2\pi)^3 2\omega_k}} \left[ [q(\bar{k}) + ip(\bar{k})] e^{-i\omega_k t + i\bar{k}\cdot\bar{x}} + [q(\bar{k}) - ip(\bar{k})] e^{i\omega_k t - i\bar{k}\cdot\bar{x}} \right]. \end{aligned} \quad (3.2)$$

These  $q(\bar{k})$  and  $p(\bar{k})$  operators bear no relation to any physical position or momentum operators. Their only role here is to enable us to convert the discrete infinite-dimensional basis associated with each  $a(\bar{k})$  and  $a^\dagger(\bar{k})$  into a continuous one. Specifically, we can realize the  $[q(\bar{k}), p(\bar{k}')]$  commutator by  $p(\bar{k}') = -i\partial/\partial q(\bar{k}')$ , with  $H$  then becoming a wave operator. In this way for each  $\bar{k}$  we obtain a solution to the Schrödinger equation of the form  $\psi(\bar{k}) = e^{-q^2(\bar{k})/2}/\pi^{1/4}$ . We can define a no-particle vacuum that obeys  $a(\bar{k})|\Omega\rangle$  for each  $\bar{k}$ . For each  $\bar{k}$  we have

$$\langle q(\bar{k})|a(\bar{k})|\Omega\rangle = \frac{1}{\sqrt{2}} \left[ q(\bar{k}) + \frac{\partial}{\partial q(\bar{k})} \right] \langle q(\bar{k})|\Omega\rangle = 0, \quad (3.3)$$

so that  $\langle q(\bar{k})|\Omega\rangle = e^{-q^2(\bar{k})/2}/\pi^{1/4}$ , and thus

$$\langle \Omega|\Omega\rangle = \prod_{\bar{k}} \int dq(\bar{k}) \langle \Omega|q(\bar{k})\rangle \langle q(\bar{k})|\Omega\rangle = \prod_{\bar{k}} \int dq(\bar{k}) \frac{e^{-q^2(\bar{k})}}{\pi^{1/2}} = \prod_{\bar{k}} 1 = 1. \quad (3.4)$$

Thus the vacuum for the full  $H$  obeys  $\langle \Omega|\Omega\rangle = 1$ , to thus have a finite normalization. In this way we establish that the vacuum state of the free relativistic scalar field is normalizable.

The general prescription then is to convert the occupation number space Hamiltonian into a product of individual occupation number spaces each with its own  $\bar{k}$ , and then determine whether the equivalent wave mechanics ground state wave functions constructed this way have a finite normalization in the conventional Schrödinger wave mechanics theory sense. If they do, then so does the full vacuum  $|\Omega\rangle$  of the full  $H$ . If on the other hand the equivalent wave mechanics wave functions are not normalizable, then neither is the full  $|\Omega\rangle$ . Our results for the second-order scalar field theory are also presented in [1], and in this paper we apply them to the higher-derivative theories that are of interest to quantum gravity, showing that for them the standard Dirac norm  $\langle \Omega|\Omega\rangle$  is not finite.

Once we are able to show that the vacuum state of the free theory is normalizable, this will remain true in the presence of interactions if the interacting theory is renormalizable. Specifically, if the free theory  $D(x) = -i\langle \Omega|T[\phi(x)\phi(0)]|\Omega\rangle$  is finite, which it will be if the free theory  $\langle \Omega|\Omega\rangle$  is, then the interacting  $D(x) = -i\langle \Omega|T[\phi(x)\phi(0)]|\Omega\rangle$  propagator will equally be finite after renormalization. Consequently, the renormalized  $\langle \Omega|\Omega\rangle$  will be finite too. Thus to establish the finiteness of the vacuum normalization of a renormalizable interacting theory, we only need to be able to make a creation and annihilation representation of the free theory. We discuss the role of interactions further in Sec. IX.

As well as providing a procedure for determining whether or not  $\langle \Omega|\Omega\rangle$  is finite, since the procedure enables us to express the free second-order-derivative Hamiltonian  $H$  as an ordinary derivative operator, it does so for interactions as well. Specifically, from (3.2) we can write  $\phi(\bar{x}, t)$  as a derivative operator, viz.

$$\phi(\bar{x}, t) = \frac{1}{\sqrt{2}} \int \frac{d^3k}{\sqrt{(2\pi)^3 2\omega_k}} \left[ \left[ q(\bar{k}) + \frac{\partial}{\partial q(\bar{k})} \right] e^{-i\omega_k t + i\bar{k}\cdot\bar{x}} + \left[ q(\bar{k}) - \frac{\partial}{\partial q(\bar{k})} \right] e^{i\omega_k t - i\bar{k}\cdot\bar{x}} \right]. \quad (3.5)$$

Thus the insertion of (3.5) into an interaction Hamiltonian of the form  $H_I = \lambda \int d^3x \phi^4(\bar{x}, t)$  enables us to write  $H_I$ , and thus  $H + H_I$ , as a derivative operator. While this procedure enables us to in principle set up the Schrödinger problem for  $H + H_I$  as a wave mechanics problem, it is still quite a formidable one, just as interacting field theories

always have been. However, since it would write the theory in terms of fourth-order derivatives, and since it has an interacting vacuum that is normalized to one and excited states that all have positive norm, it does provide an example of a theory with higher derivatives that is free of negative-norm ghost states. Thus we can anticipate (and in fact find) that the second-order plus fourth-order quantum field theory that we discuss below in Secs. [IV](#) - [VII](#) will be ghost free too.

Since the above analysis is driven by the fact that the dimension of the occupation number space is infinite, the analysis can be carried out for any bosonic field. However, because of the Pauli principle, the occupation number space basis for fermions of any given  $\bar{k}$  is finite dimensional. Thus we have to treat fermions separately, and do so in Sec. [X](#).

#### IV. HIGHER-DERIVATIVE QUANTUM FIELD THEORIES

Having presented an example of a theory whose vacuum state is normalizable, we now present an example for which  $\langle\Omega|\Omega\rangle$  is not finite. The example is based on a second-order plus fourth-order derivative neutral scalar field theory with action and equation of motion

$$I_S = \frac{1}{2} \int d^4x \left[ \partial_\mu \partial_\nu \phi \partial^\mu \partial^\nu \phi - (M_1^2 + M_2^2) \partial_\mu \phi \partial^\mu \phi + M_1^2 M_2^2 \phi^2 \right],$$

$$(\partial_t^2 - \bar{\nabla}^2 + M_1^2)(\partial_t^2 - \bar{\nabla}^2 + M_2^2)\phi(x) = 0, \quad (4.1)$$

with  $\text{diag}[\eta_{\mu\nu}] = (1, -1, -1, -1)$ . While we now study this particular model just for illustrative purposes, we note that it actually arises in quantum gravity studies, and in Secs. [XI](#) and [XII](#) we shall explore the implications of this study for quantum gravity.

For [\(4.1\)](#) the associated propagator obeys

$$(\partial_t^2 - \bar{\nabla}^2 + M_1^2)(\partial_t^2 - \bar{\nabla}^2 + M_2^2)D(x) = -\delta^4(x),$$

$$D(x) = - \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot x}}{(k^2 - M_1^2)(k^2 - M_2^2)} = - \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot x}}{(M_1^2 - M_2^2)} \left[ \frac{1}{(k^2 - M_1^2)} - \frac{1}{(k^2 - M_2^2)} \right]. \quad (4.2)$$

The energy-momentum tensor  $T_{\mu\nu}$ , the canonical momenta  $\pi^\mu$  and  $\pi^{\mu\lambda}$ , and the equal-time commutators appropriate to the higher-derivative theory are given by [\[2\]](#)

$$T_{\mu\nu} = \pi_\mu \phi_{,\nu} + \pi_\mu^{\lambda} \phi_{,\nu,\lambda} - \eta_{\mu\nu} \mathcal{L},$$

$$\pi^\mu = \frac{\partial \mathcal{L}}{\partial \phi_{,\mu}} - \partial_\lambda \left( \frac{\partial \mathcal{L}}{\partial \phi_{,\mu,\lambda}} \right) = -\partial_\lambda \partial^\mu \partial^\lambda \phi - (M_1^2 + M_2^2) \partial^\mu \phi,$$

$$\pi^{\mu\lambda} = \frac{\partial \mathcal{L}}{\partial \phi_{,\mu,\lambda}} = \partial^\mu \partial^\lambda \phi,$$

$$T_{00} = \frac{1}{2} \pi_{00}^2 + \pi_0 \dot{\phi} + \frac{1}{2} (M_1^2 + M_2^2) \dot{\phi}^2 - \frac{1}{2} M_1^2 M_2^2 \phi^2 - \frac{1}{2} \pi_{ij} \pi^{ij} + \frac{1}{2} (M_1^2 + M_2^2) \phi_{,i} \phi^{,i}$$

$$= \frac{1}{2} \ddot{\phi}^2 - \frac{1}{2} (M_1^2 + M_2^2) \dot{\phi}^2 - \ddot{\phi} \dot{\phi} - [\partial_i \partial^i \dot{\phi}] \dot{\phi} - \frac{1}{2} M_1^2 M_2^2 \phi^2 - \frac{1}{2} \partial_i \partial_j \phi \partial^i \partial^j \phi + \frac{1}{2} (M_1^2 + M_2^2) \partial_i \phi \partial^i \phi,$$

$$[\phi(\bar{0}, t), \dot{\phi}(\bar{x}, t)] = 0, \quad [\phi(\bar{0}, t), \ddot{\phi}(\bar{x}, t)] = 0, \quad [\phi(\bar{0}, t), \ddot{\phi}(\bar{x}, t)] = -i\delta^3(x). \quad (4.3)$$

With the use of these commutation relations we find that

$$D(x) = i \langle \Omega | T[\phi(x)\phi(0)] | \Omega \rangle \quad (4.4)$$

indeed satisfies the first equation given in [\(4.2\)](#), provided that is that  $\langle \Omega | \Omega \rangle = 1$  [\[3\]](#).

To check whether  $\langle \Omega | \Omega \rangle$  actually is finite, we need to express the scalar field Hamiltonian  $H_S = \int d^3x T_{00}$  in terms of creation and annihilation operators and then construct an equivalent wave mechanics. Given that the solutions to [\(4.1\)](#) are plane waves, we set

$$\phi(\bar{x}, t) = \int \frac{d^3k}{(2\pi)^{3/2}} \left[ a_1(\bar{k}) e^{-i\omega_1 t + i\bar{k} \cdot \bar{x}} + a_1^\dagger(\bar{k}) e^{i\omega_1 t - i\bar{k} \cdot \bar{x}} + a_2(\bar{k}) e^{-i\omega_2 t + i\bar{k} \cdot \bar{x}} + a_2^\dagger(\bar{k}) e^{i\omega_2 t - i\bar{k} \cdot \bar{x}} \right]. \quad (4.5)$$

where  $\omega_1 = +(\bar{k}^2 + M_1^2)^{1/2}$ ,  $\omega_2 = +(\bar{k}^2 + M_2^2)^{1/2}$ . Given the commutators in [\(4.3\)](#) we obtain

$$[a_1(\bar{k}), a_1^\dagger(\bar{k}')] = [2(M_1^2 - M_2^2)(\bar{k}^2 + M_1^2)^{1/2}]^{-1} \delta^3(\bar{k} - \bar{k}'),$$

$$\begin{aligned}
[a_2(\bar{k}), a_2^\dagger(\bar{k}')] &= -[2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2)^{1/2}]^{-1} \delta^3(\bar{k} - \bar{k}'), \\
[a_1(\bar{k}), a_2(\bar{k}')] &= 0, \quad [a_1(\bar{k}), a_2^\dagger(\bar{k}')] = 0, \quad [a_1^\dagger(\bar{k}), a_2(\bar{k}')] = 0, \quad [a_1^\dagger(\bar{k}), a_2^\dagger(\bar{k}')] = 0,
\end{aligned} \tag{4.6}$$

with the Hamiltonian then taking the form

$$\begin{aligned}
H_S &= \frac{1}{2} \int d^3k \left[ 2(M_1^2 - M_2^2)(\bar{k}^2 + M_1^2) \left[ a_1^\dagger(\bar{k}) a_1(\bar{k}) + a_1(\bar{k}) a_1^\dagger(\bar{k}) \right] \right. \\
&\quad \left. - 2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2) \left[ a_2^\dagger(\bar{k}) a_2(\bar{k}) + a_2(\bar{k}) a_2^\dagger(\bar{k}) \right] \right] \\
&= \int d^3k \left[ 2(M_1^2 - M_2^2)(\bar{k}^2 + M_1^2) a_1^\dagger(\bar{k}) a_1(\bar{k}) - 2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2) a_2^\dagger(\bar{k}) a_2(\bar{k}) \right. \\
&\quad \left. + \frac{1}{2}(\bar{k}^2 + M_1^2)^{1/2} \delta^3(0) + \frac{1}{2}(\bar{k}^2 + M_2^2)^{1/2} \delta^3(0) \right],
\end{aligned} \tag{4.7}$$

where  $(2\pi)^3 \delta^3(0)$  is a quantization box volume  $V$ . We note that with  $M_1^2 - M_2^2 > 0$  for definitiveness, we see negative signs in both  $H_S$  and the  $[a_2(\bar{k}), a_2^\dagger(\bar{k}')]$  commutator, while noting that despite this the zero-point energy is positive. We shall see below that the negative sign concerns will be resolved once we settle the issue of the normalization of the vacuum. To do that we now descend to the quantum-mechanical limit of the theory, the Pais-Uhlenbeck oscillator model.

## V. HIGHER-DERIVATIVE QUANTUM MECHANICS

In order to study the Pauli-Villars regulator, in [4] Pais and Uhlenbeck (PU) introduced a fourth-order quantum-mechanical oscillator model with action and equation of motion

$$I_{\text{PU}} = \frac{1}{2} \int dt [\ddot{z}^2 - (\omega_1^2 + \omega_2^2) \dot{z}^2 + \omega_1^2 \omega_2^2 z^2], \quad \ddot{\ddot{z}} + (\omega_1^2 + \omega_2^2) \ddot{z} + \omega_1 \omega_2 z^2 = 0, \tag{5.1}$$

where for definitiveness in the following we take  $\omega_1 > \omega_2$ . As constructed this action possesses three variables  $z$ ,  $\dot{z}$  and  $\ddot{z}$ . This is too many for one oscillator but not enough for two. The system is thus a constrained system. And so we introduce a new variable  $x = \dot{z}$  and its conjugate  $p_x$ . And using the method of Dirac constraints obtain the time-independent Hamiltonian [5, 6]

$$H_{\text{PU}} = \frac{p_x^2(t)}{2} + p_z(t)x(t) + \frac{1}{2}(\omega_1^2 + \omega_2^2)x^2(t) - \frac{1}{2}\omega_1^2\omega_2^2 z^2(t), \tag{5.2}$$

with two sets of canonical equal-time commutators of the form

$$[z(t), p_z(t)] = i, \quad [x(t), p_x(t)] = i. \tag{5.3}$$

The terms in  $H_{\text{PU}}$  are in complete parallel to the first four terms in the field theory  $T_{00}$  given in (4.3), with the PU oscillator model being the nonrelativistic limit of the relativistic scalar field theory, with the spatial dependence having been frozen out. Since canonical commutators only involve time derivatives, freezing out the spatial dependence will still give the full dynamical content of the relativistic theory. In fact we can set  $i = [z, p_z] \equiv [\phi, \pi_0] = [\phi, -\dot{\phi} - (M_1^2 + M_2^2)\phi] = i\delta^3(x)$ , to thus parallel the commutators given in (4.3).

On setting  $p_z = -i\partial_z$ ,  $p_x = -i\partial_x$  the Schrödinger problem for  $H_{\text{PU}}$  can be solved analytically, with the state with energy  $(\omega_1 + \omega_2)/2$  having a wave function that is of the form [7]

$$\psi_0(z, x) = \exp\left[\frac{1}{2}(\omega_1 + \omega_2)\omega_1\omega_2 z^2 + i\omega_1\omega_2 z x - \frac{1}{2}(\omega_1 + \omega_2)x^2\right]. \tag{5.4}$$

While this wave function is well behaved at large  $x$ , it diverges at large  $z$ , and consequently as a wave function it is not normalizable.

To relate this wave function to the no-particle vacuum  $|\Omega\rangle$  we second quantize the theory. And with the wave equation given in (5.1), and with  $\dot{z} = i[H_{\text{PU}}, z] = x$ ,  $\dot{x} = p_x$ ,  $\dot{p}_x = -p_z - (\omega_1^2 + \omega_2^2)x$ ,  $\dot{p}_z = \omega_1^2\omega_2^2 z$ , we obtain

$$\begin{aligned}
z(t) &= a_1 e^{-i\omega_1 t} + a_1^\dagger e^{i\omega_1 t} + a_2 e^{-i\omega_2 t} + a_2^\dagger e^{i\omega_2 t}, \\
p_x(t) &= i\omega_1\omega_2^2 [a_1 e^{-i\omega_1 t} - a_1^\dagger e^{i\omega_1 t}] + i\omega_1^2\omega_2 [a_2 e^{-i\omega_2 t} - a_2^\dagger e^{i\omega_2 t}],
\end{aligned}$$

$$\begin{aligned}
x(t) &= -i\omega_1[a_1e^{-i\omega_1t} - a_1^\dagger e^{i\omega_1t}] - i\omega_2[a_2e^{-i\omega_2t} - a_2^\dagger e^{i\omega_2t}], \\
p_x(t) &= -\omega_1^2[a_1e^{-i\omega_1t} + a_1^\dagger e^{i\omega_1t}] - \omega_2^2[a_2e^{-i\omega_2t} + a_2^\dagger e^{i\omega_2t}],
\end{aligned} \tag{5.5}$$

and a Hamiltonian and commutator algebra of the form [5]

$$H_{\text{PU}} = 2(\omega_1^2 - \omega_2^2)(\omega_1^2 a_1^\dagger a_1 - \omega_2^2 a_2^\dagger a_2) + \frac{1}{2}(\omega_1 + \omega_2), \tag{5.6}$$

$$[a_1, a_1^\dagger] = \frac{1}{2\omega_1(\omega_1^2 - \omega_2^2)}, \quad [a_2, a_2^\dagger] = -\frac{1}{2\omega_2(\omega_1^2 - \omega_2^2)}. \tag{5.7}$$

We note the similarity to (4.7) and (4.6).

As constructed, (5.6) and (5.7) admit of two inequivalent realizations as one can take  $a_1$  and either  $a_2$  or  $a_2^\dagger$  to annihilate the vacuum. Thus one can define a Hilbert space in which  $a_1|\Omega\rangle = 0$ ,  $a_2|\Omega\rangle = 0$ , or one can define a separate and distinct Hilbert space in which  $a_1|\Omega\rangle = 0$ ,  $a_2^\dagger|\Omega\rangle = 0$ . In the Hilbert space in which  $a_1|\Omega\rangle = 0$ ,  $a_2|\Omega\rangle = 0$ , we note that even though the  $a_2^\dagger a_2$  term appears in  $H_{\text{PU}}$  with a minus sign, there is a compensating minus sign in the  $[a_2, a_2^\dagger]$  commutator. In consequence, in this realization all energy eigenvalues of  $H_{\text{PU}}$  are positive, with the no-particle state  $|\Omega\rangle$  that both  $a_1$  and  $a_2$  annihilate being the state of lowest energy. However, in this Hilbert space the matrix element  $\langle\Omega|a_2 a_2^\dagger|\Omega\rangle$  is negative, the ghost problem of higher-derivative theories. This problem has been solved in the literature [2, 8], and we shall return to it in detail below.

Alternatively, if one takes  $a_2^\dagger \equiv b_2$  to annihilate the vacuum, we obtain a commutator  $[b_2, b_2^\dagger] = 1/[2\omega_1(\omega_1^2 - \omega_2^2)]$  that is positive. However then the energy spectrum of the Hamiltonian becomes unbounded from below, the familiar Ostrogradski instability of higher-derivative theories.

While we thus have to deal with a negative-norm problem or a negative energy problem, we note that the two problems do not occur in one and the same Hilbert space. Thus in any given Hilbert space we at most only have to deal with one. And since an energy spectrum that is unbounded from below is not physical, we shall work solely in the Hilbert space in which  $a_1|\Omega\rangle = 0$ ,  $a_2|\Omega\rangle = 0$ . In Sec. XII, which provides a quick explanation of our results, we shall show that the  $a_1|\Omega\rangle = 0$ ,  $a_2|\Omega\rangle = 0$  and  $a_1|\Omega\rangle = 0$ ,  $a_2^\dagger|\Omega\rangle = 0$  realizations correspond to different and thus inequivalent Feynman  $i\epsilon$  prescriptions.

In the Hilbert space in which both  $a_1$  and  $a_2$  annihilate the vacuum the energy spectrum is bounded from below, and the energy of the ground state is  $(\omega_1 + \omega_2)/2$ . On solving the time-dependent Schrödinger equation the wave function of the ground state is  $\psi_0(z, x)e^{-i(\omega_1 + \omega_2)t/2}$ , where  $\psi_0(z, x)$  is given in (5.4). For this wave function the normalization of  $|\Omega\rangle$  is then given by

$$\langle\Omega|\Omega\rangle = \int_{-\infty}^{\infty} dz \int_{-\infty}^{\infty} dx \langle\Omega|z, x\rangle \langle z, x|\Omega\rangle = \int_{-\infty}^{\infty} dz \int_{-\infty}^{\infty} dx \psi_0^*(z, x) \psi_0(z, x). \tag{5.8}$$

With  $\psi_0(z, x)$  diverging at large  $z$ , this normalization integral is infinite. Thus we see that through our knowledge of the form of the ground state wave function as given in (5.4) we are able to determine the normalization of the PU theory vacuum and establish that it is infinite. We can thus anticipate and will immediately show in Sec. VI that this is also the case for the second-order plus fourth-order scalar quantum field theory as well. Then in Sec. VII we will discuss what to do about it, with there actually being a mechanism for obtaining a finite normalization [2, 8], one that also takes care of the fact that according to (5.7)  $\langle\Omega|a_2 a_2^\dagger|\Omega\rangle$  is negative.

Already in [7] it was known that for the ground state of the second-order plus fourth-order quantum-mechanical PU oscillator model the vacuum  $\langle\Omega|\Omega\rangle$  norm was infinite. However, it was not known whether the vacuum  $\langle\Omega|\Omega\rangle$  norm of the second-order plus fourth-order quantum field theory was finite or infinite. Using the procedure developed in this paper we can now determine whether it is finite or infinite, and in Sec. VI we show that  $\langle\Omega|\Omega\rangle$  is in fact infinite. Then in Sec. VII we show that there is another inner product, viz. the  $\langle L|R\rangle$  overlap of left-eigenstates and right-eigenstates of the Hamiltonian, that is finite.

## VI. THE NONNORMALIZABLE VACUUM OF HIGHER-DERIVATIVE FIELD THEORIES

To determine the second-order plus fourth-order scalar field theory vacuum normalization we first need to invert (5.5). This yields

$$a_1 e^{-i\omega_1 t} = \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ -\omega_2^2 z(t) - p_x(t) + i\omega_1 x(t) + i\frac{p_z(t)}{\omega_1} \right],$$

$$\begin{aligned}
a_1^\dagger e^{i\omega_1 t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ -\omega_2^2 z(t) - p_x(t) - i\omega_1 x(t) - i\frac{p_z(t)}{\omega_1} \right], \\
a_2 e^{-i\omega_2 t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ \omega_1^2 z(t) + p_x(t) - i\omega_2 x(t) - i\frac{p_z(t)}{\omega_2} \right], \\
a_2^\dagger e^{i\omega_2 t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ \omega_1^2 z(t) + p_x(t) + i\omega_2 x(t) + i\frac{p_z(t)}{\omega_2} \right].
\end{aligned} \tag{6.1}$$

On generalizing to each  $\bar{k}$  and setting  $\omega_1(\bar{k}) = +(\bar{k}^2 + M_1^2)^{1/2}$ ,  $\omega_2(\bar{k}) = +(\bar{k}^2 + M_2^2)^{1/2}$ , we obtain

$$\begin{aligned}
a_1(\bar{k}) e^{-i\omega_1(\bar{k})t} &= \frac{1}{2(M_1^2 - M_2^2)} \left[ -\omega_2^2(\bar{k})z(\bar{k}, t) - p_x(\bar{k}, t) + i\omega_1(\bar{k})x(\bar{k}, t) + i\frac{p_z(\bar{k}, t)}{\omega_1(\bar{k})} \right], \\
a_1^\dagger(\bar{k}) e^{i\omega_1(\bar{k})t} &= \frac{1}{2(M_1^2 - M_2^2)} \left[ -\omega_2^2(\bar{k})z(\bar{k}, t) - p_x(\bar{k}, t) - i\omega_1(\bar{k})x(\bar{k}, t) - i\frac{p_z(\bar{k}, t)}{\omega_1(\bar{k})} \right], \\
a_2(\bar{k}) e^{-i\omega_2(\bar{k})t} &= \frac{1}{2(M_1^2 - M_2^2)} \left[ \omega_1^2(\bar{k})z(\bar{k}, t) + p_x(\bar{k}, t) - i\omega_2(\bar{k})x(\bar{k}, t) - i\frac{p_z(\bar{k}, t)}{\omega_2(\bar{k})} \right], \\
a_2^\dagger(\bar{k}) e^{i\omega_2(\bar{k})t} &= \frac{1}{2(M_1^2 - M_2^2)} \left[ \omega_1^2(\bar{k})z(\bar{k}, t) + p_x(\bar{k}, t) + i\omega_2(\bar{k})x(\bar{k}, t) + i\frac{p_z(\bar{k}, t)}{\omega_2(\bar{k})} \right].
\end{aligned} \tag{6.2}$$

Inverting (6.2) gives

$$\begin{aligned}
z(\bar{k}, t) &= a_1(\bar{k}) e^{-i\omega_1(\bar{k})t} + a_1^\dagger(\bar{k}) e^{i\omega_1(\bar{k})t} + a_2(\bar{k}) e^{-i\omega_2(\bar{k})t} + a_2^\dagger(\bar{k}) e^{i\omega_2(\bar{k})t}, \\
p_z(\bar{k}, t) &= i\omega_1(\bar{k})\omega_2^2(\bar{k})[a_1(\bar{k}) e^{-i\omega_1(\bar{k})t} - a_1^\dagger(\bar{k}) e^{i\omega_1(\bar{k})t}] + i\omega_1^2(\bar{k})\omega_2(\bar{k})[a_2(\bar{k}) e^{-i\omega_2(\bar{k})t} - a_2^\dagger(\bar{k}) e^{i\omega_2(\bar{k})t}], \\
x(\bar{k}, t) &= -i\omega_1(\bar{k})[a_1(\bar{k}) e^{-i\omega_1(\bar{k})t} - a_1^\dagger(\bar{k}) e^{i\omega_1(\bar{k})t}] - i\omega_2(\bar{k})[a_2(\bar{k}) e^{-i\omega_2(\bar{k})t} - a_2^\dagger(\bar{k}) e^{i\omega_2(\bar{k})t}], \\
p_x(\bar{k}, t) &= -\omega_1^2(\bar{k})[a_1(\bar{k}) e^{-i\omega_1(\bar{k})t} + a_1^\dagger(\bar{k}) e^{i\omega_1(\bar{k})t}] - \omega_2^2(\bar{k})[a_2(\bar{k}) e^{-i\omega_2(\bar{k})t} + a_2^\dagger(\bar{k}) e^{i\omega_2(\bar{k})t}].
\end{aligned} \tag{6.3}$$

From (6.3) and the commutation relations given in (4.6) it follows that

$$\begin{aligned}
[z(\bar{k}, t), p_z(\bar{k}', t)] &= \delta^3(\bar{k} - \bar{k}'), & [x(\bar{k}, t), p_x(\bar{k}', t)] &= \delta^3(\bar{k} - \bar{k}'), \\
[z(\bar{k}, t), x(\bar{k}', t)] &= 0, & [z(\bar{k}, t), p_x(\bar{k}', t)] &= 0, & [p_z(\bar{k}, t), x(\bar{k}', t)] &= 0, & [p_z(\bar{k}, t), p_x(\bar{k}', t)] &= 0.
\end{aligned} \tag{6.4}$$

Insertion of (6.2) into the Hamiltonian given in (4.7) then yields an equivalent, time-independent Hamiltonian

$$H_S = \int d^3k \left[ \frac{p_x^2(\bar{k}, t)}{2} + p_z(\bar{k}, t)x(\bar{k}, t) + \frac{1}{2} [\omega_1^2(\bar{k}) + \omega_2^2(\bar{k})] x^2(\bar{k}, t) - \frac{1}{2} \omega_1^2(\bar{k})\omega_2^2(\bar{k}) z^2(\bar{k}, t) \right]. \tag{6.5}$$

For each momentum state we recognize the quantum field theory Hamiltonian  $H_S$  given in (6.5) as being of precisely the form of the quantum-mechanical  $H_{\text{PU}}$  Hamiltonian that is given in (5.2).

We can now proceed as in the second-order scalar quantum field theory discussed above and represent the commutators by

$$\left[ z(\bar{k}, t), -i\frac{\partial}{\partial z(\bar{k}', t)} \right] = \delta^3(\bar{k} - \bar{k}'), \quad \left[ x(\bar{k}, t), -i\frac{\partial}{\partial x(\bar{k}', t)} \right] = \delta^3(\bar{k} - \bar{k}'). \tag{6.6}$$

With the vacuum obeying  $a_1(\bar{k})|\Omega\rangle = 0$ ,  $a_2(\bar{k})|\Omega\rangle = 0$  for each  $\bar{k}$ , from (6.2) we obtain

$$\begin{aligned}
\langle z(\bar{k}), x(\bar{k}) | a_1(\bar{k}) | \Omega \rangle &= \frac{1}{2(M_1^2 - M_2^2)} \left[ -\omega_2^2(\bar{k})z(\bar{k}) + i\frac{\partial}{\partial x(\bar{k})} + i\omega_1(\bar{k})x(\bar{k}) + \frac{1}{\omega_1(\bar{k})} \frac{\partial}{\partial z(\bar{k})} \right] \langle z(\bar{k}), x(\bar{k}) | \Omega \rangle = 0, \\
\langle z(\bar{k}), x(\bar{k}) | a_2(\bar{k}) | \Omega \rangle &= \frac{1}{2(M_1^2 - M_2^2)} \left[ \omega_1^2(\bar{k})z(\bar{k}) - i\frac{\partial}{\partial x(\bar{k})} - i\omega_2(\bar{k})x(\bar{k}) - \frac{1}{\omega_2(\bar{k})} \frac{\partial}{\partial z(\bar{k})} \right] \langle z(\bar{k}), x(\bar{k}) | \Omega \rangle = 0,
\end{aligned} \tag{6.7}$$

for each  $\bar{k}$ . From (6.7) it follows that for each  $\bar{k}$  we can identify each  $\langle z(\bar{k}), x(\bar{k}) | \Omega \rangle$  with the PU oscillator ground state wave function  $\psi_0(z(\bar{k}), x(\bar{k}))$ , which, analogously to (5.4), is given by

$$\psi_0(z(\bar{k}), x(\bar{k})) = \exp\left[\frac{1}{2}[\omega_1(\bar{k}) + \omega_2(\bar{k})]\omega_1(\bar{k})\omega_2(\bar{k})z^2(\bar{k}) + i\omega_1(\bar{k})\omega_2(\bar{k})z(\bar{k})x(\bar{k}) - \frac{1}{2}[\omega_1(\bar{k}) + \omega_2(\bar{k})]x^2(\bar{k})\right]. \tag{6.8}$$

Consequently, the normalization of the vacuum is given by

$$\begin{aligned} \langle \Omega | \Omega \rangle &= \Pi_{\bar{k}} \int_{-\infty}^{\infty} dz(\bar{k}) \int_{-\infty}^{\infty} dx(\bar{k}) \langle \Omega | z(\bar{k}), x(\bar{k}) \rangle \langle z(\bar{k}), x(\bar{k}) | \Omega \rangle \\ &= \Pi_{\bar{k}} \int_{-\infty}^{\infty} dz(\bar{k}) \int_{-\infty}^{\infty} dx(\bar{k}) \psi_0^*(z(\bar{k}), x(\bar{k})) \psi_0(z(\bar{k}), x(\bar{k})). \end{aligned} \quad (6.9)$$

With each  $\psi_0(z(\bar{k}), x(\bar{k}))$  diverging at large  $z(\bar{k})$ , we thus establish that the normalization of the field theory vacuum is infinite. Thus whatever is the normalization of the vacuum in the associated wave-mechanical limit translates into the same normalization in the quantum field theory.

## VII. HOW TO OBTAIN A NORMALIZABLE VACUUM

In analyzing the second-order plus fourth-order scalar field theory we note that with a conventional Hermitian field  $\phi(x)$ , and thus with  $a_1^\dagger(\bar{k})$  and  $a_2^\dagger(\bar{k})$  being the Hermitian conjugates of  $a_1(\bar{k})$  and  $a_2(\bar{k})$ , the  $a_2^\dagger(\bar{k})a_2(\bar{k})$  product would be positive definite and the energy spectrum of  $H_S$  as given in (4.7) would initially be unbounded from below, this being the familiar Ostrogradski instability of higher-derivative theories with Hermitian fields. However, from (4.6) we see that  $\langle \Omega | a_2(\bar{k}) a_2^\dagger(\bar{k}) | \Omega \rangle$  would be negative. This would imply the potential presence of ghost states of negative norm, with it then not being the case that a product such as  $a_2(\bar{k}) a_2^\dagger(\bar{k})$  could be positive definite. If one accepts this then matrix elements of the  $-2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2) a_2^\dagger(\bar{k}) a_2(\bar{k})$  term in  $H_S$  would be compensated for by the ghost signature, and the energy spectrum of  $H_S$  would then be bounded from below. While this takes care of the unboundedness from below of the energy spectrum, it appears to do so at a high price, namely the potential presence of unitarity-violating ghost states. But if  $a_2^\dagger(\bar{k})$  is the Hermitian conjugate of  $a_2(\bar{k})$  then  $\langle \Omega | a_2(\bar{k}) a_2^\dagger(\bar{k}) | \Omega \rangle$  would have to be positive. Thus despite the dagger notation  $a_2^\dagger(\bar{k})$  could not be the Hermitian conjugate of  $a_2(\bar{k})$ . Hence our starting assumption that  $\phi(x)$  is Hermitian could not be valid. Consequently, the Hamiltonian that is built out of the  $\phi(x)$  field could not be Hermitian either. And in fact we have actually established that it is not, since the diverging of  $\psi_0(z(\bar{k}), x(\bar{k}))$  at large  $z(\bar{k})$  means that in an integration by parts we could not drop surface terms, with the presence of such surface terms preventing Hermiticity or self-adjointness. With the eigenstates of the Hamiltonian not being normalizable, there not only are potential negative-norm states present, they are infinitely negative.

Surprisingly, it is this very inability to drop surface terms in an integration by parts that actually saves the theory [2, 8]. Specifically, we have seen that we are working with a Hamiltonian  $H_S$  (and likewise  $H_{\text{PU}}$ ) that is not Hermitian. However, all the energy eigenvalues associated with  $H_S$  and  $H_{\text{PU}}$  are real. Now Hermiticity is only sufficient for real eigenvalues, with the necessary condition [9, 10] being that the Hamiltonian have an antilinear symmetry [11]. The theory thus falls into the class of  $PT$  theories ( $P$  is the linear parity operator and  $T$  is the antilinear time reversal operator) developed by Bender and collaborators [12, 13]. Critical to the  $PT$  program is that the wave functions be normalizable in some domain in the complex plane, a domain known technically as a Stokes wedge. Since the wave functions are not normalizable with real  $z$  or real  $z(\bar{k})$ , we have to continue  $z$  and  $z(\bar{k})$  into the complex plane in order to make them normalizable. Then the theory is well-defined, with, as we discuss in Sec. VIII, the domain of the measure needed for the Minkowski path integral accordingly also having to be continued into the complex plane in order to make it be well-defined too [2, 10]. For the particular case of  $\psi_0(z, x)$  and  $\psi_0(z(\bar{k}), x(\bar{k}))$ , replacing  $z$  by  $-iz$  and  $z(\bar{k})$  by  $-iz(\bar{k})$  would then make both  $\psi_0(z, x)$  and  $\psi_0(z(\bar{k}), x(\bar{k}))$  normalizable. (We have no need to modify  $x$  or  $x(\bar{k})$  since the wave functions already are well behaved when these quantities become large.)

To achieve the continuation of  $z$  or  $z(\bar{k})$  at the level of operators we effect similarity transformations as they preserve both energy eigenvalues and canonical commutators. We introduce

$$S(\text{PU}) = e^{\pi p_z z/2}, \quad S(S) = e^{\pi \int d^3 x \pi_0(\bar{x}, t) \phi(\bar{x}, t)/2}, \quad (7.1)$$

and obtain

$$\begin{aligned} S(\text{PU})zS(\text{PU})^{-1} &= -iz \equiv y, & S(\text{PU})p_zS(\text{PU})^{-1} &= ip_z \equiv q, \\ S(S)z(\bar{k})S(S)^{-1} &= -iz(\bar{k}) \equiv y(\bar{k}), & S(S)p_z(\bar{k})S(S)^{-1} &= ip_z(\bar{k}) \equiv q(\bar{k}). \end{aligned} \quad (7.2)$$

### A. The PU case

For the PU oscillator this leads to

$$S(\text{PU})H_{\text{PU}}S(\text{PU})^{-1} = \bar{H}_{\text{PU}} = \frac{1}{2}p_x^2(t) - iq(t)x(t) + \frac{1}{2}(\omega_1^2 + \omega_2^2)x^2(t) + \frac{1}{2}\omega_1^2\omega_2^2y^2(t),$$



$$[y(t), q(t)] = i, \quad [x(t), p_x(t)] = i. \quad (7.3)$$

Since the  $\psi(y, x)$  wave functions are now well behaved at infinity, the  $\bar{H}_{\text{PU}}$  Hamiltonian is self-adjoint. However, because of the  $-iq(t)x(t)$  term it is not Hermitian. Instead it is  $PT$  symmetric. Specifically, with  $p$  and  $q$  being taken to be  $PT$  even and  $y$  and  $x$  being taken to be  $PT$  odd [2], the  $PT$  invariance of  $\bar{H}_{\text{PU}}$  and of the  $[y, q] = i$  and  $[x, p_x] = i$  commutators follows. Now when a Hamiltonian is not Hermitian the action of it to the right and the action of it to the left are not related by Hermitian conjugation. Thus in general one must distinguish between right and left eigenstates, both for the vacuum and the states that can be excited out of it. Thus we represent the  $[y, q] = i$  and  $[x, p_x] = i$  commutators by  $q = -i\overrightarrow{\partial}_y$ ,  $p_x = -i\overrightarrow{\partial}_x$  when acting to the right, and by  $q = i\overleftarrow{\partial}_y$ ,  $p_x = i\overleftarrow{\partial}_x$  when acting to the left. This then leads to right and left ground state wave functions of the form [2]

$$\begin{aligned} \psi_0^R(y, x) &= \exp[-\frac{1}{2}(\omega_1 + \omega_2)\omega_1\omega_2y^2 - \omega_1\omega_2yx - \frac{1}{2}(\omega_1 + \omega_2)x^2], \\ \psi_0^L(y, x) &= \exp[-\frac{1}{2}(\omega_1 + \omega_2)\omega_1\omega_2y^2 + \omega_1\omega_2yx - \frac{1}{2}(\omega_1 + \omega_2)x^2]. \end{aligned} \quad (7.4)$$

Given these wave functions the vacuum normalization is given by [2]

$$\begin{aligned} \langle \Omega^L | \Omega^R \rangle &= \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dx \langle \Omega^L | y, x \rangle \langle y, x | \Omega^R \rangle = \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dx \psi_0^L(y, x) \psi_0^R(y, x) \\ &= \int_{-\infty}^{\infty} dy \int_{-\infty}^{\infty} dx \exp[-(\omega_1 + \omega_2)\omega_1\omega_2y^2 - (\omega_1 + \omega_2)x^2] = \frac{\pi}{(\omega_1\omega_2)^{1/2}(\omega_1 + \omega_2)}, \end{aligned} \quad (7.5)$$

with the vacuum state thus being normalizable. In the following we shall understand the wave functions to have been normalized to one, so that  $\int dy dx \psi_0^L(y, x) \psi_0^R(y, x) = 1$  and  $\langle \Omega^L | \Omega^R \rangle = 1$  [14].

With the above  $PT$  assignments and with  $\dot{y} = i[\bar{H}_{\text{PU}}, y] = -ix$ ,  $\dot{x} = p_x$ ,  $\dot{p}_x = iq - (\omega_1^2 + \omega_2^2)x$ ,  $\dot{q} = -\omega_1^2\omega_2^2y$ , we set

$$\begin{aligned} y(t) &= -ia_1e^{-i\omega_1t} + a_2e^{-i\omega_2t} - i\hat{a}_1e^{i\omega_1t} + \hat{a}_2e^{i\omega_2t}, \\ x(t) &= -i\omega_1a_1e^{-i\omega_1t} + \omega_2a_2e^{-i\omega_2t} + i\omega_1\hat{a}_1e^{i\omega_1t} - \omega_2\hat{a}_2e^{i\omega_2t}, \\ p_x(t) &= -\omega_1^2a_1e^{-i\omega_1t} - i\omega_2^2a_2e^{-i\omega_2t} - \omega_1^2\hat{a}_1e^{i\omega_1t} - i\omega_2^2\hat{a}_2e^{i\omega_2t}, \\ q(t) &= \omega_1\omega_2[-\omega_2a_1e^{-i\omega_1t} - i\omega_1a_2e^{-i\omega_2t} + \omega_2\hat{a}_1e^{i\omega_1t} + i\omega_1\hat{a}_2e^{i\omega_2t}], \\ a_1e^{-i\omega_1t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ -i\omega_2^2y(t) - p_x(t) + i\omega_1x(t) + \frac{q(t)}{\omega_1} \right], \\ \hat{a}_1e^{+i\omega_1t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ -i\omega_2^2y(t) - p_x(t) - i\omega_1x(t) - \frac{q(t)}{\omega_1} \right], \\ ia_2e^{-i\omega_2t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ i\omega_1^2y(t) + p_x(t) - i\omega_2x(t) - \frac{q(t)}{\omega_2} \right], \\ i\hat{a}_2e^{+i\omega_2t} &= \frac{1}{2(\omega_1^2 - \omega_2^2)} \left[ i\omega_1^2y(t) + p_x(t) + i\omega_2x(t) + \frac{q(t)}{\omega_2} \right]. \end{aligned} \quad (7.6)$$

In (7.6) we have introduced  $a_1, a_2, \hat{a}_1$  and  $\hat{a}_2$ , with the four creation and annihilation operators obeying  $PTa_1TP = a_1$ ,  $PTa_2TP = -a_2$ ,  $PT\hat{a}_1TP = \hat{a}_1$ ,  $PT\hat{a}_2TP = -\hat{a}_2$ , so as to enforce the  $PT$  assignments of  $y, x, p_x$  and  $q$ . Comparing with (5.5) we have  $(a_1, a_2, a_1^\dagger, a_2^\dagger) \rightarrow (a_1, ia_2, \hat{a}_1, i\hat{a}_2)$ .

With (7.3) and (7.6) the Hamiltonian is given by

$$\bar{H}_{\text{PU}} = 2(\omega_1^2 - \omega_2^2) (\omega_1^2\hat{a}_1a_1 + \omega_2^2\hat{a}_2a_2) + \frac{1}{2}(\omega_1 + \omega_2), \quad (7.7)$$

and the operator commutation algebra is given by

$$\begin{aligned} [a_1, \hat{a}_1] &= \frac{1}{2\omega_1(\omega_1^2 - \omega_2^2)}, \quad [a_2, \hat{a}_2] = \frac{1}{2\omega_2(\omega_1^2 - \omega_2^2)}, \\ [a_1, a_2] &= 0, \quad [a_1, \hat{a}_2] = 0, \quad [\hat{a}_1, a_2] = 0, \quad [\hat{a}_1, \hat{a}_2] = 0. \end{aligned} \quad (7.8)$$

With the  $PT$  assignments of  $a_1, a_2, \hat{a}_1$  and  $\hat{a}_2$ , we confirm the  $PT$  invariance of (7.7) and (7.8). In (7.7) and (7.8) the relative signs are all positive (we take  $\omega_1 > \omega_2 > 0$  for definitiveness), so these equations define a standard positive energy, positive norm, two-dimensional harmonic oscillator system. Given the creation and annihilation operators the left and right vacua are defined by

$$\langle \Omega^L | \hat{a}_1 = 0, \quad \langle \Omega^L | \hat{a}_2 = 0, \quad a_1 | \Omega^R \rangle = 0, \quad a_2 | \Omega^R \rangle = 0. \quad (7.9)$$

By exciting modes out of the left and right vacua we can build excited states that have positive norm [8], viz.  $\langle n^L | m^R \rangle = \delta_{nm}$ , and obey a completeness relation

$$\sum |n_1^R\rangle \langle n_1^L| + \sum |n_2^R\rangle \langle n_2^L| = I. \quad (7.10)$$

Even though these norms are all positive, the insertion of (7.10) into  $-i\langle \Omega^L | T[y(t)y(0)] | \Omega^R \rangle$  (corresponding to  $+i\langle \Omega^L | T[z(t)z(0)] | \Omega^R \rangle$ ) generates the relative minus sign in the nonrelativistic limit of the  $-[1/(k^2 - M_1^2) - 1/(k^2 - M_2^2)]/(M_1^2 - M_2^2)$  propagator given in (4.2), viz.  $-[1/(\omega^2 - \omega_1^2) - 1/(\omega^2 - \omega_2^2)]/(\omega_1^2 - \omega_2^2)$ . We thus establish the consistency and physical viability of the similarity transformed *PU* oscillator theory.

## B. The relativistic case

For  $S(S)H_S S(S)^{-1} = \bar{H}_S$  we introduce creation and annihilation operators for  $\bar{\phi} = S(S)\phi S(S)^{-1} = -i\phi(x)$  of the form

$$\bar{\phi}(x) = \int \frac{d^3k}{(2\pi)^{3/2}} \left[ -ia_1(\bar{k})e^{-i\omega_1(\bar{k})t + i\bar{k}\cdot\bar{x}} + a_2(\bar{k})e^{-i\omega_2(\bar{k})t + i\bar{k}\cdot\bar{x}} - i\hat{a}_1(\bar{k})e^{i\omega_1(\bar{k})t - i\bar{k}\cdot\bar{x}} + \hat{a}_2(\bar{k})e^{i\omega_2(\bar{k})t - i\bar{k}\cdot\bar{x}} \right]. \quad (7.11)$$

Comparing with (4.5) we have  $(a_1(\bar{k}), a_2(\bar{k}), a_1^\dagger(\bar{k}), a_2^\dagger(\bar{k})) \rightarrow (a_1(\bar{k}), ia_2(\bar{k}), \hat{a}_1(\bar{k}), i\hat{a}_2(\bar{k}))$ . Like  $y(t)$ ,  $\bar{\phi}(x)$  is *PT* odd. As constructed the Hamiltonian is *PT* even. (In general, even without Hermiticity one still has *CPT* symmetry [10], but since the fields are neutral *C* is separately conserved, so in this case *CPT* defaults to *PT*.) The field-theoretic *PT*-even Hamiltonian and *PT*-preserving commutation relations are given by [2]

$$\begin{aligned} S(S)H_S S(S)^{-1} = \bar{H}_S = & \frac{1}{2} \int d^3k \left[ 2(M_1^2 - M_2^2)(\bar{k}^2 + M_1^2) [\hat{a}_1(\bar{k})a_1(\bar{k}) + a_1(\bar{k})\hat{a}_1(\bar{k})] \right. \\ & \left. + 2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2) [\hat{a}_2(\bar{k})a_2(\bar{k}) + a_2(\bar{k})\hat{a}_2(\bar{k})] \right], \end{aligned} \quad (7.12)$$

and

$$\begin{aligned} [\dot{\bar{\phi}}(\bar{x}, t), \bar{\phi}(0)] &= 0, & [\ddot{\bar{\phi}}(\bar{x}, t), \bar{\phi}(0)] &= 0, & [\ddot{\bar{\phi}}(\bar{x}, t), \bar{\phi}(0)] &= i\delta^3(x), \\ [a_1(\bar{k}), \hat{a}_1(\bar{k}')] &= [2(M_1^2 - M_2^2)(\bar{k}^2 + M_1^2)^{1/2}]^{-1} \delta^3(\bar{k} - \bar{k}'), \\ [a_2(\bar{k}), \hat{a}_2(\bar{k}')] &= [2(M_1^2 - M_2^2)(\bar{k}^2 + M_2^2)^{1/2}]^{-1} \delta^3(\bar{k} - \bar{k}'), \\ [a_1(\bar{k}), a_2(\bar{k}')] &= 0, & [a_1(\bar{k}), \hat{a}_2(\bar{k}')] &= 0, & [\hat{a}_1(\bar{k}), a_2(\bar{k}')] &= 0, & [\hat{a}_1(\bar{k}), \hat{a}_2(\bar{k}')] &= 0. \end{aligned} \quad (7.13)$$

With all relative signs being positive (we take  $M_1^2 > M_2^2$  for definitiveness), there are no states of negative norm or of negative energy. The discussion completely parallels that of the *PU* oscillator model given above. We introduce

$$\begin{aligned} y(\bar{k}, t) &= -ia_1(\bar{k})e^{-i\omega_1(\bar{k})t} + a_2(\bar{k})e^{-i\omega_2(\bar{k})t} - i\hat{a}_1(\bar{k})e^{i\omega_1(\bar{k})t} + \hat{a}_2(\bar{k})e^{i\omega_2(\bar{k})t}, \\ x(\bar{k}, t) &= -i\omega_1(\bar{k})a_1(\bar{k})e^{-i\omega_1(\bar{k})t} + \omega_2(\bar{k})a_2(\bar{k})e^{-i\omega_2(\bar{k})t} + i\omega_1(\bar{k})\hat{a}_1(\bar{k})e^{i\omega_1(\bar{k})t} - \omega_2(\bar{k})\hat{a}_2(\bar{k})e^{i\omega_2(\bar{k})t}, \\ p_x(\bar{k}, t) &= -\omega_1^2(\bar{k})a_1(\bar{k})e^{-i\omega_1(\bar{k})t} - i\omega_2^2(\bar{k})a_2(\bar{k})e^{-i\omega_2(\bar{k})t} - \omega_1^2(\bar{k})\hat{a}_1(\bar{k})e^{i\omega_1(\bar{k})t} - i\omega_2^2(\bar{k})\hat{a}_2(\bar{k})e^{i\omega_2(\bar{k})t}, \\ q(\bar{k}, t) &= \omega_1(\bar{k})\omega_2(\bar{k})[-\omega_2(\bar{k})a_1(\bar{k})e^{-i\omega_1(\bar{k})t} - i\omega_1(\bar{k})a_2(\bar{k})e^{-i\omega_2(\bar{k})t} + \omega_2(\bar{k})\hat{a}_1(\bar{k})e^{i\omega_1(\bar{k})t} + i\omega_1(\bar{k})\hat{a}_2(\bar{k})e^{i\omega_2(\bar{k})t}], \end{aligned} \quad (7.14)$$

with  $\bar{H}_S$  then taking the form

$$\bar{H}_S = \int d^3k \left[ \frac{p_x^2(\bar{k}, t)}{2} - iq(\bar{k}, t)x(\bar{k}, t) + \frac{1}{2} [\omega_1^2(\bar{k}) + \omega_2^2(\bar{k})] x^2(\bar{k}, t) + \frac{1}{2} \omega_1^2(\bar{k})\omega_2^2(\bar{k})y^2(\bar{k}, t) \right]. \quad (7.15)$$

Introducing left and right vacua that obey

$$\langle \Omega^L | \hat{a}_1(\bar{k}) = 0, \quad \langle \Omega^L | \hat{a}_2(\bar{k}) = 0, \quad a_1(\bar{k}) | \Omega^R \rangle = 0, \quad a_2(\bar{k}) | \Omega^R \rangle = 0 \quad (7.16)$$

for all  $\bar{k}$ , we find that

$$\langle \Omega^L | \bar{H}_S | \Omega^R \rangle = \int d^3k \left[ \frac{1}{2} (\bar{k}^2 + M_1^2)^{1/2} + \frac{1}{2} (\bar{k}^2 + M_2^2)^{1/2} \right] \delta^3(0),$$

$$\begin{aligned}
\langle \Omega^L | \Omega^R \rangle &= \Pi_{\bar{k}} \int_{-\infty}^{\infty} dy(\bar{k}) \int_{-\infty}^{\infty} dx(\bar{k}) \langle \Omega^L | y(\bar{k}), x(\bar{k}) \rangle \langle y(\bar{k}), x(\bar{k}) | \Omega^R \rangle \\
&= \Pi_{\bar{k}} \int_{-\infty}^{\infty} dy(\bar{k}) \int_{-\infty}^{\infty} dx(\bar{k}) \psi_0^L(y(\bar{k}), x(\bar{k})) \psi_0^R(y(\bar{k}), x(\bar{k})) = \Pi_{\bar{k}} 1 = 1.
\end{aligned} \tag{7.17}$$

We thus confirm that the vacuum normalization is both finite and positive, while the vacuum energy has the conventional zero-point infinity associated with an infinite number of modes. (This infinity occurs because  $\bar{H}_S$  contains an infinite number of modes and not because  $\langle \Omega^L | \Omega^R \rangle$  itself is infinite.) We thus establish the consistency and physical viability of the similarity-transformed higher-derivative scalar field theory. And we note that even though all the norms are positive, the insertion of (7.10) into  $-i\langle \Omega^L | T[\bar{\phi}(x)\bar{\phi}(0)] | \Omega^R \rangle$  (corresponding to  $+i\langle \Omega^L | T[\phi(x)\phi(0)] | \Omega^R \rangle$ ) generates the relative minus sign in  $-[1/(k^2 - M_1^2) - 1/(k^2 - M_2^2)]/(M_1^2 - M_2^2)$  [2]. Thus with one similarity transform into an appropriate Stokes wedge we solve both the vacuum normalization problem and the negative-norm problem.

At this point we can see the key aspect of our study. Ordinarily in quantum field theory it is taken as a given that one should use the Dirac inner product  $\langle \Omega | \Omega \rangle$ , viz.  $\langle \Omega^R | \Omega^R \rangle$ , for the vacuum. And also it is taken as a given that this inner product is finite. In this paper we have provided a procedure for checking whether this is in fact the case, and presented a second-order plus fourth-order derivative model in which it explicitly is not finite. For this particular model we have found a different inner product, viz.  $\langle \Omega^L | \Omega^R \rangle$ , that is finite. (For a Hamiltonian that is Hermitian  $|\Omega^R\rangle = |\Omega\rangle$ ,  $\langle \Omega^L| = \langle \Omega|$ , and  $\langle \Omega^L | \Omega^R \rangle = \langle \Omega | \Omega \rangle$ .) And thus in general one has to determine whether or not  $\langle \Omega^R | \Omega^R \rangle$  is finite on case by case basis.

We should note that in general there is no requirement that a quantum field theory must use the Dirac inner product  $\langle \Omega | \Omega \rangle$ . What one does need of an acceptable inner product is that it be finite, positive and time independent. And as long as we can find one we have a well-defined Hilbert space. However, if the Hamiltonian is not Hermitian then the Dirac inner product obeys  $\langle \Omega(t) | \Omega(t) \rangle = \langle \Omega(t=0) | e^{iH^\dagger t} e^{-iHt} | \Omega(t=0) \rangle \neq \langle \Omega(t=0) | \Omega(t=0) \rangle$ , to thus not be time independent, and thus not acceptable.

In regard to time independence, we also note that in the nonrelativistic case the time derivative of the  $\int d^3x \psi^*(x)\psi(x)$  probability is related to an asymptotic spatial surface term. The time independence of  $\int d^3x \psi^*(x)\psi(x)$  is then secured if the asymptotic spatial surface term vanishes, i.e., the time independence of  $\int d^3x \psi^*(x)\psi(x)$  is determined by asymptotic spatial boundary conditions. If it turns out that the theory is well behaved asymptotically for Hermitian fields then one can use the Dirac norm. If it is not well behaved asymptotically for Hermitian fields, then the insight of the  $PT$  program is to look to see if instead there then might be some domain in the complex plane where the theory is well behaved asymptotically. If there is, then one can have a good Hilbert space, but the inner product will not then be the Dirac norm. If the theory has a  $PT$  symmetry, then the norm will be the  $PT$  theory norm, and when the Hamiltonian is not Hermitian the left-right inner product obeys  $\langle \Omega^L(t) | \Omega^R(t) \rangle = \langle \Omega^L(t=0) | e^{iH^\dagger t} e^{-iHt} | \Omega^R(t=0) \rangle = \langle \Omega^L(t=0) | \Omega^R(t=0) \rangle$  to thus be time independent. And this is the case for the second-order plus fourth-order theory.

With the finiteness or otherwise of the path integral also being determined by spatial boundary conditions, in Sec. VIII we now discuss our findings from the perspective of path integrals. In Sec. XII we augment our path integral results with a quick and straightforward analysis of Feynman's  $i\epsilon$  prescription for propagators.

## VIII. PATH INTEGRALS AND THE NORMALIZATION OF THE VACUUM

We are interested in the path integral approach for both the PU oscillator and the second-order plus fourth-order field theory. We treat the field theory path integral in Sec. XII. For the PU oscillator the Minkowski path integral ( $PI$ ) associated with  $I_{PU}$  is of the form

$$PI(MINK) = \int D[z] D[dz/dt] \exp \left[ \frac{i}{2} \int_{-\infty}^{\infty} dt \left( \left( \frac{d^2 z}{dt^2} \right)^2 - (\omega_1^2 + \omega_2^2) \left( \frac{dz}{dt} \right)^2 + \omega_1^2 \omega_2^2 z^2 \right) \right], \tag{8.1}$$

as initially integrated over paths with real  $z$  and  $x$ . Since the theory is fourth order we need four pieces of information to solve the equations of motion. The pieces that are the most convenient for path integral purposes are two initial and two final conditions, hence the path integral measure is over both  $z$  and  $dz/dt$ . However, we had noted in Sec. V that the PU theory is a constrained theory. Thus we must treat the  $z$  and  $dz/dt$  path integrations as independent. We can do this directly as shown in the measure in (8.1), or replace (8.1) by

$$PI(MINK, x, z) = \int D[z] D[x] \exp \left[ \frac{i}{2} \int_{-\infty}^{\infty} dt \left( \left( \frac{dx}{dt} \right)^2 - (\omega_1^2 + \omega_2^2) x^2 + \omega_1^2 \omega_2^2 z^2 \right) \right]. \tag{8.2}$$

To make the path integral converge rather than just oscillate we first use the Feynman  $i\epsilon$  prescription and replace  $\omega_1^2$  and  $\omega_2^2$  by  $\omega_1^2 - i\epsilon$  and  $\omega_2^2 - i\epsilon$ . This yields

$$PI(MINK, x, z) = \int D[z]D[x] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} dt \left( i \left( \frac{dx}{dt} \right)^2 - i (\omega_1^2 + \omega_2^2) x^2 + i\omega_1^2\omega_2^2 z^2 - 2\epsilon x^2 + \epsilon (\omega_1^2 + \omega_2^2) z^2 \right) \right], \quad (8.3)$$

as integrated over paths with real  $x$  and real  $z$ . However, while the  $x^2$  term is now damped the  $z^2$  term is not. Consequently, as integrated with a real measure the path integral does not exist. Now the path integral is used to generate time-ordered Green's functions such as  $D(x) = i\langle\Omega|T[\phi(x)\phi(0)]|\Omega\rangle$  (hence the  $i\epsilon$  prescription). And thus these Green's functions will not be finite, with the vacuum in which the Green's function matrix elements are evaluated thus not being normalizable. Study of the Minkowski path integral thus gives us an alternate way to determine whether or not  $\langle\Omega|\Omega\rangle$  is finite: the path integral with a real measure either exists or does not exist.

To make (8.2) exist we need to damp the  $z^2$  term, but not modify the  $x^2$  term. Thus we continue  $z$  into the complex plane and replace it by  $y = -iz$ , while leaving  $x$  real. The path integral for Minkowski time then takes the form

$$PI(MINK, x, y) = \int D[y]D[x] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} dt \left( i \left( \frac{dx}{dt} \right)^2 - i (\omega_1^2 + \omega_2^2) x^2 - i\omega_1^2\omega_2^2 y^2 - 2\epsilon x^2 - \epsilon (\omega_1^2 + \omega_2^2) y^2 \right) \right]. \quad (8.4)$$

This puts us into a domain in the complex plane (known as a Stokes wedge) in which the path integral is now fully defined, and now the vacuum state is normalizable. This completely parallels the discussion of  $\psi_0(z, x)$  that we gave in Sec. V.




FIG. 1: Wick contour

However, our concern here could be missed in a Euclidean time path integral approach. Specifically, if we disperse in  $t$  (assuming of course that we can, i.e., that the Cauchy-Riemann equations for complex  $t$  are obeyed), we can, as shown in Fig. 1, write

$$\int_{-\infty}^{\infty} + \int_{\infty}^{i\infty} + \int_{i\infty}^{-i\infty} + \int_{-i\infty}^{-\infty} = \text{pole terms plus cut contributions}, \quad (8.5)$$

i.e., along the real axis, then upper-half-plane quarter circle, then down the imaginary axis, and then lower-half-plane quarter circle. Assuming no pole, cut or circle contributions, and on setting  $\tau = it$  and letting  $I$  denote the action, from (8.1) and (8.2) we obtain

$$\begin{aligned} I(MINK, z, x) &\equiv \int_{-\infty}^{\infty} idt \equiv - \int_{i\infty}^{-i\infty} idt = - \int_{-\infty}^{\infty} d\tau \equiv I(EUCL, z, x), \\ PI(EUCL, z, x) &= \int D[z]D[dz/d\tau] \exp \left[ -\frac{1}{2} \int_{-\infty}^{\infty} d\tau \left( \left( \frac{d^2z}{d\tau^2} \right)^2 + (\omega_1^2 + \omega_2^2) \left( \frac{dz}{d\tau} \right)^2 + \omega_1^2\omega_2^2 z^2 \right) \right] \\ &= \int D[z]D[x] \exp \left[ -\frac{1}{2} \int_{-\infty}^{\infty} d\tau \left( \left( \frac{dx}{d\tau} \right)^2 + (\omega_1^2 + \omega_2^2) x^2 + \omega_1^2\omega_2^2 z^2 \right) \right]. \end{aligned} \quad (8.6)$$

Given the overall minus sign that multiplies the Euclidean action on every path, we see that with real  $z$  and real  $x = dz/d\tau$  the Euclidean path integral is well behaved. (The same is true of the analog relativistic second-order plus

fourth-order scalar field theory path integral [15].) However, the Minkowski time path integral with a real measure is not. Thus we conclude that the pole and/or cut and/or circle contributions are not only not ignorable, they generate an infinite contribution. Hence their contribution in a Wick rotation cannot be ignored and the Euclidean time path integral does not correctly describe the situation.

In parallel, if we set  $y = -iz$ , then (8.6) is replaced by

$$I(MINK, y, x) \equiv I(EUCL, y, x),$$

$$PI(EUCL, y, x) = \int D[y]D[x] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} d\tau \left( - \left( \frac{dx}{d\tau} \right)^2 - (\omega_1^2 + \omega_2^2) x^2 + \omega_1^2 \omega_2^2 y^2 \right) \right]. \quad (8.7)$$

And since the  $y$  and  $x$  path integrations are independent, now it is the Euclidean time path integral that is not well defined. Thus with either  $z$  or  $y = -iz$ , in neither case are the Minkowski time and Euclidean time path integrals simultaneously finite.

As we discuss in Sec. XII, these same results carry over directly to the field theory case, and thus we see that even if finite, a Euclidean time path integral approach is only valid if the vacuum state of the theory (as determined in a Minkowski time analysis) is normalizable.

## IX. INTERACTIONS

In developing Wick's contraction theorem in quantum field theory one needs to put the time-ordered product of Heisenberg fields  $\phi(x)$ , viz.

$$\tau(x_1, \dots, x_n) = \langle \Omega | T[\phi(x_1) \dots \phi(x_n)] | \Omega \rangle, \quad (9.1)$$

into a form that can be developed perturbatively. To this end one introduces a set of in-fields  $\phi_{in}(x)$  that satisfy free field equations with Hamiltonian  $H_{in}$ . And one also introduces an evolution operator  $U(t)$  that evolves with the interaction Hamiltonian  $H_I(t)$  according to

$$i \frac{\partial U(t)}{\partial t} = H_I(t) U(t). \quad (9.2)$$

With this  $U(t)$  we can relate  $\phi(x)$  and  $\phi_{in}(x)$  according to

$$\phi(\bar{x}, t) = U^{-1}(t) \phi_{in}(\bar{x}, t) U(t). \quad (9.3)$$

If one introduces  $U(t, t') = U(t) U^{-1}(t')$ , then  $U(t, t')$  is given by

$$U(t, t') = 1 - i \int_{t'}^t dt_1 H_I(t_1) U(t_1, t') = T \left[ \exp \left( -i \int_{t'}^t dt_1 H_I(t_1) \right) \right]. \quad (9.4)$$

Using these relations we obtain (see e.g. [16]) viz.

$$\tau(x_1, \dots, x_n) = \langle \Omega | U^{-1}(t) T \left[ \phi_{in}(x_1) \dots \phi_{in}(x_n) \exp \left( -i \int_{-t}^t dt_1 H_I(t_1) \right) \right] U(-t) | \Omega \rangle. \quad (9.5)$$

The contributions due to the  $U(t)|\Omega\rangle$  and  $\langle\Omega|U^{-1}(t)$  terms lead to

$$\begin{aligned} \tau(x_1, \dots, x_n) &= \langle \Omega | T \left[ \phi_{in}(x_1) \dots \phi_{in}(x_n) \exp \left( -i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle \\ &\times \langle \Omega | T \left[ \exp \left( i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle. \end{aligned} \quad (9.6)$$

After inverting the last term we obtain the standard form [16]

$$\tau(x_1, \dots, x_n) = \frac{\langle \Omega | T \left[ \phi_{in}(x_1) \dots \phi_{in}(x_n) \exp \left( -i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle}{\langle \Omega | T \left[ \exp \left( -i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle}. \quad (9.7)$$

If one starts with (9.7) it would appear that the normalization of the vacuum state is actually irrelevant since it would drop out of the ratio. And so it would not appear to matter if it did happen to be infinite. However, this is not the case since we could only go from (9.6) to (9.7) if  $\langle \Omega | T \left[ \exp \left( i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle$  is finite. And it would not be if the vacuum state is not normalizable. If we expand  $\langle \Omega | T \left[ \exp \left( i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle$  out as a power series in  $H_I$  the first term is  $\langle \Omega | \Omega \rangle$  as calculated in a free theory. Thus, as we had noted in Sec. III, for finiteness we need this term to be finite and need the power series expansion in  $H_I$  to be renormalizable in order for  $\langle \Omega | T \left[ \exp \left( i \int_{-t}^t dt_1 H_I(t_1) \right) \right] | \Omega \rangle$  to be finite. However, for a nonnormalizable vacuum the standard Wick expansion and Feynman rules are not valid. Since this concern is of relevance to radiative corrections to Einstein gravity we return to this point in Secs. XI and XII below.

## X. FERMIONS

For fermions we have to deal with anticommutators such as

$$bb^\dagger + b^\dagger b = 1. \quad (10.1)$$

Also, because of the Pauli principle we have

$$b^2 = 0, \quad b^{\dagger 2} = 0. \quad (10.2)$$

We can represent (10.1) and (10.2) by matrices of the form

$$b = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad b^\dagger = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}. \quad (10.3)$$

Thus, unlike the infinite-dimensional matrix representation of the bosonic  $a$  and  $a^\dagger$  that obey  $aa^\dagger - a^\dagger a = 1$ , the fermionic  $b$  and  $b^\dagger$  matrices are finite dimensional. Thus with a finite number of degrees of freedom, the fermion vacuum that obeys  $b|\Omega\rangle = 0$  has a finite  $\langle \Omega | \Omega \rangle$  norm.

## XI. IMPLICATIONS FOR RADIATIVE CORRECTIONS IN QUANTUM EINSTEIN GRAVITY

As a quantum theory the standard second-order derivative Einstein gravitational theory with its  $1/k^2$  propagator is not renormalizable since graviton loops cannot be made finite by counterterms that are themselves of the same generic form as the Einstein-Hilbert Ricci scalar action. Since these graviton loops generate higher-derivative gravity terms, one can construct a candidate theory of quantum gravity by augmenting the Einstein Ricci scalar action with a term that is quadratic in the Ricci scalar. This gives a much studied quantum gravity action of the generic form

$$I_{\text{GRAV}} = \int d^4x (-g)^{1/2} [6M^2 R^\alpha{}_\alpha + (R^\alpha{}_\alpha)^2]. \quad (11.1)$$

This same action also appears in Starobinsky's inflationary universe model [17].

On adding on a matter source with energy-momentum tensor  $T_{\mu\nu}$ , variation of this action with respect to the metric generates a gravitational equation of motion of the form

$$-6M^2 G^{\mu\nu} + V^{\mu\nu} = -\frac{1}{2} T^{\mu\nu}. \quad (11.2)$$

Here  $G_{\mu\nu}$  is the Einstein tensor and  $V_{\mu\nu}$  may for instance be found in [18], with these various terms being of the form

$$\begin{aligned} G^{\mu\nu} &= R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} g^{\alpha\beta} R_{\alpha\beta}, \\ V^{\mu\nu} &= 2g^{\mu\nu} \nabla_\beta \nabla^\beta R^\alpha{}_\alpha - 2\nabla^\nu \nabla^\mu R^\alpha{}_\alpha - 2R^\alpha{}_\alpha R^{\mu\nu} + \frac{1}{2} g^{\mu\nu} (R^\alpha{}_\alpha)^2. \end{aligned} \quad (11.3)$$

If we now linearize about flat spacetime with background metric  $\eta_{\mu\nu}$  and fluctuation metric  $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ , to first perturbative order we obtain

$$\delta G_{\mu\nu} = \frac{1}{2} (\partial_\alpha \partial^\alpha h_{\mu\nu} - \partial_\mu \partial^\alpha h_{\alpha\nu} - \partial_\nu \partial^\alpha h_{\alpha\mu} + \partial_\mu \partial_\nu h) - \frac{1}{2} \eta_{\mu\nu} (\partial_\alpha \partial^\alpha h - \partial^\alpha \partial^\beta h_{\alpha\beta}),$$

$$\delta V_{\mu\nu} = [2\eta_{\mu\nu}\partial_\alpha\partial^\alpha - 2\partial_\mu\partial_\nu][\partial_\beta\partial^\beta h - \partial_\lambda\partial_\kappa h^{\lambda\kappa}], \quad (11.4)$$

where  $h = \eta^{\mu\nu}h_{\mu\nu}$ . On taking the trace of the fluctuation around a background (11.2) we obtain

$$[M^2 + \partial_\beta\partial^\beta](\partial_\lambda\partial^\lambda h - \partial_\kappa\partial_\lambda h^{\kappa\lambda}) = -\frac{1}{12}\eta^{\mu\nu}\delta T_{\mu\nu}. \quad (11.5)$$

In the convenient transverse gauge where  $\partial_\mu h^{\mu\nu} = 0$ , the propagator for  $h$  is given by

$$D(h, k^2) = -\frac{1}{k^2(k^2 - M^2)} = \frac{1}{M^2} \left( \frac{1}{k^2} - \frac{1}{k^2 - M^2} \right). \quad (11.6)$$

As we see, in this case the  $1/k^2$  graviton propagator for  $h$  that would be associated with the Einstein tensor  $\delta G_{\mu\nu}$  alone is replaced by a  $D(h, k^2) = [1/k^2 - 1/(k^2 - M^2)]/M^2$  propagator. And now the leading behavior at large momenta is  $-1/k^4$ . In consequence, the theory is thought to be renormalizable [19, 20]. But since  $\langle\Omega|\Omega\rangle$  is not finite the proof of renormalizability has a flaw in it. Fortunately, the flaw is not fatal, and we rectify it below.

We recognize  $D(h, k^2)$  as being of the same form as the second-order plus fourth-order scalar field theory propagator that was given in (4.2), with  $\phi$  being replaced by  $h$  and with  $M_1^2 = M^2$ ,  $M_2^2 = 0$ . We can thus give  $h$  an equivalent effective action of the form

$$I_h = \frac{1}{2} \int d^4x \left[ \partial_\mu\partial_\nu h \partial^\mu\partial^\nu h - M^2 \partial_\mu h \partial^\mu h \right]. \quad (11.7)$$

The action given in (11.7) thus shares the same vacuum state normalization and negative norm challenges as the scalar field action given in (4.1).

Thus if, as is conventional, we take  $h$  to be Hermitian we would immediately encounter the negative-norm problem associated with the relative minus sign in (11.6). However, since  $M^2$  is Planck scale in magnitude, this difficulty can be postponed until observations can reach that energy scale. However, the lack of normalizability of the vacuum state has consequences at all energies and cannot be postponed at all. Specifically, with  $\langle\Omega|\Omega\rangle$  being infinite we cannot even identify the propagator as  $i\langle\Omega|T[h(x)h(0)]|\Omega\rangle$  since in analog to (1.8) it will obey

$$(\partial_t^2 - \bar{\nabla}^2)(\partial_t^2 - \bar{\nabla}^2 + M^2)D(h, x) = -\langle\Omega|\Omega\rangle\delta^4(x). \quad (11.8)$$

Consequently, we cannot make the standard Wick contraction expansion. And thus both the Feynman rules that are used presupposing that  $\langle\Omega|\Omega\rangle$  is finite, and the renormalizability that is thought to then follow from them are therefore not valid. Additionally, with  $\langle\Omega|\Omega\rangle$  being infinite, we cannot treat the Einstein theory with its  $1/k^2$  propagator as an effective field theory that holds for momenta that obey  $k^2 \ll M^2$ .

However, as noted above, we can resolve all of these concerns by dropping the requirement that  $h$  be Hermitian, and set it equal to  $i\bar{h}$ . Then, with the theory being recognized as a  $PT$  theory, vacuum state normalization and negative-norm problems are resolved and the theory is consistent. Moreover, the propagator is given by  $-i\langle\Omega^L|T[\bar{h}(x)h(0)]|\Omega^R\rangle$  (corresponding to  $+i\langle\Omega^L|T[h(x)h(0)]|\Omega^R\rangle$ ). And with the propagator still being given by (11.6) as it satisfies  $(\partial_t^2 - \bar{\nabla}^2 + M^2)(\partial_t^2 - \bar{\nabla}^2)[-i\langle\Omega^L|T[\bar{h}(x)h(0)]|\Omega^R\rangle] = -\delta^4(x)$ , all the steps needed to prove renormalizability are now valid. At this point the only concern is that even though the  $M^2$  field now has positive norm, it still remains in the spectrum and would eventually have to be observed.

As can be seen from (11.7), the only reason that there is an  $M^2$  term at all is because we are considering an action that has both second-order and fourth-order terms. With a pure fourth-order theory there would be no dimensionful parameter in the action and the theory would be scale invariant. If like the gauge theories of  $SU(3) \times SU(2) \times U(1)$  this scale symmetry is also local, we would be led to conformal gravity, a metric theory of gravity in which the action is left invariant under local changes of the metric of the form  $g_{\mu\nu}(x) \rightarrow e^{2\alpha(x)}g_{\mu\nu}(x)$ , where  $\alpha(x)$  is a local function of the coordinates. The conformal gravity theory has been advocated and explored in [18, 21] and references therein. And in [22] 't Hooft has also argued that there should be an underlying local conformal symmetry in nature.

In the conformal gravity theory an action that is to be a polynomial function of the metric has the unique form

$$I_W = -\alpha_g \int d^4x (-g)^{1/2} C_{\lambda\mu\nu\kappa} C^{\lambda\mu\nu\kappa} \equiv -2\alpha_g \int d^4x (-g)^{1/2} \left[ R_{\mu\kappa} R^{\mu\kappa} - \frac{1}{3} (R^\alpha{}_\alpha)^2 \right], \quad (11.9)$$

where  $\alpha_g$  is a dimensionless gravitational coupling constant, and  $C_{\lambda\mu\nu\kappa}$  is the conformal Weyl tensor. The perturbative propagator has a  $-1/k^4$  behavior at all  $k^2$ , and with its large  $k^2$  behavior the theory is renormalizable [23]. With a  $-1/k^4$  propagator it would initially appear that there would be two massless particles at  $k^2 = 0$ . However, we cannot use the partial fraction decomposition given in (11.6) as a guide since its  $1/M^2$  prefactor is singular in the  $M^2 \rightarrow 0$

limit. Because of this singular behavior the  $M^2 = 0$  Hamiltonian becomes of nondiagonalizable Jordan-block form and only has one massless eigenstate, with the other would-be massless eigenstate becoming nonstationary [2].

To understand why we have lost an eigenstate let us consider two positive frequency mode solutions to

$$[M^2 + \partial_\beta \partial^\beta] \partial_\lambda \partial^\lambda h = 0, \quad (11.10)$$

viz. modes with time dependence  $e^{-i\omega_1 t}$  and  $e^{-i\omega_2 t}$ , where  $\omega_1 = |\bar{k}|$ ,  $\omega_2 = +(\bar{k}^2 + M^2)^{1/2}$ . These modes are solutions to the  $M^2 \neq 0$  wave equation associated with (11.5) (as are the two negative frequency solutions with time dependence  $e^{i\omega_1 t}$  and  $e^{i\omega_2 t}$ .) If we now let  $M^2$  go to zero the two positive frequency mode solutions become equal and we only have one positive frequency mode solution. However, we cannot lose any solutions to the differential equation given in (11.10) by setting  $M^2 = 0$  in it since the wave equation is still fourth order. To find the other solution in the limit we consider

$$\lim_{M^2 \rightarrow 0} \frac{(e^{-i\omega_1 t} - e^{-i\omega_2 t})}{M^2} = \frac{ite^{-i\omega_1 t}}{2|\bar{k}|}. \quad (11.11)$$

As we see, this second solution is not stationary and is thus not an eigenstate of the Hamiltonian.

Since we have lost a massless eigenstate the propagator should be constructed not as the  $M^2 \rightarrow 0$  limit of (11.6) but as the manifestly ghost-free limit

$$-\frac{1}{(k^2 + i\epsilon)^2} = -\lim_{M^2 \rightarrow 0} \frac{d}{dM^2} \left( \frac{1}{k^2 - M^2 + i\epsilon} \right), \quad (11.12)$$

a limit that shows that there is only one  $k^2 = 0$  pole not two. With the Hamiltonian not being diagonalizable, it could not be Hermitian. It does however have a  $PT$  symmetry, with its ground state being normalizable. Conformal gravity is thus a fully consistent theory of quantum gravity, one which despite its fourth-order character only possesses one massless particle, not two [24].

To understand why both second-order plus fourth-order theories and pure fourth-order theories must be ghost free consider the Dirac action for a fermion coupled to a background geometry of the form

$$I_D = \int d^4x (-g)^{1/2} [i\bar{\psi} \gamma^c V_c^\mu(x) (\partial_\mu + \Gamma_\mu(x)) \psi - M \bar{\psi}(x) \psi(x)]. \quad (11.13)$$

Here the  $V_a^\mu(x)$  are vierbeins,  $\Gamma_\mu(x) = -(1/8)[\gamma_a, \gamma_b](V_\nu^b(x) \partial_\mu V^{a\nu}(x) + V_\lambda^b(x) \Gamma_{\nu\mu}^\lambda(x) V^{a\nu}(x))$  is the spin connection, and  $\Gamma_{\nu\mu}^\lambda(x)$  is the geometric Levi-Civita connection. We introduce the path integral  $\int D[\psi] D[\bar{\psi}] \exp(iI_D) = \exp(iI_{\text{EFF}})$ . With  $I_D$  being linear in both  $\psi$  and  $\bar{\psi}$  the fermion path integration can be performed analytically, to thereby yield an effective action  $I_{\text{EFF}}$  whose leading term is of the form [25], [21]

$$I_{\text{EFF}} = \int d^4x (-g)^{1/2} C \left[ \frac{1}{20} \left[ R_{\mu\nu} R^{\mu\nu} - \frac{1}{3} (R^\alpha{}_\alpha)^2 \right] - M^4 + \frac{1}{6} M^2 R^\alpha{}_\alpha \right], \quad (11.14)$$

where  $C$  is a log divergent constant. We recognize  $I_{\text{EFF}}$  as containing none other than second-order and fourth-order gravitational terms. Now the  $I_D$  action is a completely standard action of a fermion coupled to a background gravitational field, and as such it is not only ghost free, it would remain so if the fermion is given some internal quantum numbers and coupled to some gauge fields. Since the fermion path integral is equivalent to a one loop Feynman diagram and since one cannot change the signature of a Hilbert space in perturbation theory, higher-derivative gravity must be ghost free too. And if it were not, then the standard  $SU(3) \times SU(2) \times U(1)$  model would not remain unitary when coupled to gravity.

Thus despite the negative-norm issue, higher-derivative gravity cannot contain any negative-norm states. And for the standard model not to be destabilized when coupled to gravity, the subsequent  $g_{\mu\nu}$  path integration must be conducted with a measure  $D[g_{\mu\nu}]$  that is continued into the complex plane, just as found with  $PT$  theory. Since, as noted in Sec. VIII, this continuation is required by the  $i\epsilon$  prescription, we now comment on this prescription in more detail.

## XII. $PT$ SYMMETRY AND THE $i\epsilon$ PRESCRIPTION

The  $i\epsilon$  prescription is central to both propagators and path integrals, and discussion of its role provides a quick explanation of our results. For the second-order plus fourth-order theory discussed in Sec. IV the identification of the  $D(k)$  propagator in (4.2) as

$$D(k) = -\frac{1}{(k^2 - M_1^2)(k^2 - M_2^2)} = -\frac{1}{(M_1^2 - M_2^2)} \left[ \frac{1}{(k^2 - M_1^2)} - \frac{1}{(k^2 - M_2^2)} \right] \quad (12.1)$$



is only formal since  $D(k)$  is singular. To give it a meaning we need to define it via a contour integral and specify the appropriate complex  $k_0$  plane contour. If, as is conventional, we take all of the operators in  $H = \int d^3x T_{00}$  to be Hermitian, where as given in (4.3)

$$T_{00} = \frac{1}{2}\pi_{00}^2 + \pi_0\dot{\phi} + \frac{1}{2}(M_1^2 + M_2^2)\dot{\phi}^2 - \frac{1}{2}M_1^2M_2^2\phi^2 - \frac{1}{2}\pi_{ij}\pi^{ij} + \frac{1}{2}(M_1^2 + M_2^2)\phi_{,i}\phi^{,i}, \quad (12.2)$$

we immediately find that because of the  $-(1/2)M_1^2M_2^2\phi^2$  term the Hamiltonian  $H$  is unbounded from below, the Ostrogradski instability that is characteristic of higher-derivative theories. Now the standard Feynman contour  $k^2 + i\epsilon$  prescription with

$$D(k) = -\frac{1}{(k^2 - M_1^2 + i\epsilon)(k^2 - M_2^2 + i\epsilon)} = -\frac{1}{(M_1^2 - M_2^2)} \left[ \frac{1}{(k^2 - M_1^2 + i\epsilon)} - \frac{1}{(k^2 - M_2^2 + i\epsilon)} \right] \quad (12.3)$$

is chosen so that positive energy states propagate forward in time (viz.  $\omega_1(\bar{k}) = +(\bar{k}^2 + M_1^2)^{1/2}$ ,  $\omega_2(\bar{k}) = +(\bar{k}^2 + M_2^2)^{1/2}$  located below the real  $k_0$  axis), while negative energy states propagate backwards ( $-\omega_1(k)$ ,  $-\omega_2(k)$  above the real  $k_0$  axis); with this corresponding to an energy spectrum that is bounded from below while leading to negative residues that appear to correspond to states of negative norm. In contrast, having an energy spectrum that is unbounded from below would instead entail that negative energies in one of sectors ( $\omega_2(k)$  say) are propagating forward in time ( $-\omega_2(k)$  below the real  $k_0$  axis), and positive energies are propagating backward in time ( $\omega_2(k)$  above the real  $k_0$  axis), and would require using an unconventional  $i\epsilon$  prescription [2] of the form

$$D(k) = -\frac{1}{(k^2 - M_1^2 + i\epsilon)(k^2 - M_2^2 - i\epsilon)} = -\frac{1}{(M_1^2 - M_2^2)} \left[ \frac{1}{(k^2 - M_1^2 + i\epsilon)} - \frac{1}{(k^2 - M_2^2 - i\epsilon)} \right]. \quad (12.4)$$

While this possibility is unacceptable physically, it does have the feature that because of the way the singularities are traversed all pole residues are positive. Thus the two options are: bounded energies and negative residues, or unbounded energies and positive residues. These two realizations are inequivalent and correspond to different Feynman contours and different Hilbert spaces. As discussed in Sec. V and in [2] the first option corresponds to working in a Hilbert space in which  $a_2$  effects  $a_2|\Omega\rangle = 0$ , while the latter corresponds to working in a Hilbert space in which  $a_2^\dagger|\Omega\rangle = 0$ . Thus in no Hilbert space do we have both negative energies and negative residues. Thus if we work in the Hilbert space in which  $a_2|\Omega\rangle = 0$  (which we do in this paper) and use the standard Feynman  $i\epsilon$  prescription given in (12.3), our results will not be affected by the Ostrogradski instability at all. Nor will it affect the continuation into the complex plane that resolves the negative-norm issue.

In analog to the quantum-mechanical (8.2) and the field theory study given in [15] the Minkowski path integral associated with the field theory action given in (4.1) is of the form

$$PI(MINK) = \int D[\phi]D[\sigma_\mu] \exp \left[ \frac{i}{2} \int_{-\infty}^{\infty} d^4x \left[ \partial_\nu \sigma_\mu \partial^\nu \sigma^\mu - (M_1^2 + M_2^2) \sigma_\mu \sigma^\mu + M_1^2 M_2^2 \phi^2 \right] \right], \quad (12.5)$$

where  $\sigma_\mu = \partial_\mu \phi$ . In order to damp out oscillations we choose the Feynman  $i\epsilon$  prescription in which we replace  $M_1^2$  and  $M_2^2$  by  $M_1^2 - i\epsilon$ ,  $M_2^2 - i\epsilon$ . For the path integral this yields

$$PI(MINK) = \int D[\phi]D[\sigma_\mu] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} d^4x \left[ i\partial_\nu \sigma_\mu \partial^\nu \sigma^\mu - i(M_1^2 + M_2^2) \sigma_\mu \sigma^\mu + iM_1^2 M_2^2 \phi^2 - 2\epsilon \sigma_\mu \sigma^\mu + (M_1^2 + M_2^2)\epsilon \phi^2 \right] \right]. \quad (12.6)$$

With  $\phi$  and  $\sigma_\mu$  being taken to be real and with  $\sigma_\mu \sigma^\mu$  being taken to be timelike on every path, the  $\sigma_\mu$  path integration is damped but the  $\phi$  path integration is not.

For the unconventional  $i\epsilon$  prescription in which we replace  $M_1^2$  and  $M_2^2$  by  $M_1^2 - i\epsilon$ ,  $M_2^2 + i\epsilon$  the path integral takes the form

$$PI(MINK) = \int D[\phi]D[\sigma_\mu] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} d^4x \left[ i\partial_\nu \sigma_\mu \partial^\nu \sigma^\mu - i(M_1^2 + M_2^2) \sigma_\mu \sigma^\mu + iM_1^2 M_2^2 \phi^2 - (M_1^2 - M_2^2)\epsilon \phi^2 \right] \right], \quad (12.7)$$

and has no damping on the  $\sigma_\mu$  path integration at all. The unconventional  $i\epsilon$  prescription for the Feynman contour that leads to an unbounded from below energy spectrum thus cannot be associated with a well-defined path integral, and we cannot consider it at all.

Thus the only Feynman  $i\epsilon$  prescription that can be relevant is the standard one with  $M_1^2 - i\epsilon$ ,  $M_2^2 - i\epsilon$ . However even with this choice the  $\phi$  path integration is not damped if  $\phi$  is real. It becomes damped if we do not require  $\phi$  to be real, but instead take it to be pure imaginary (though  $(\text{Im}[\phi])^2 > (\text{Re}[\phi])^2$  would suffice). With  $\bar{\phi} = -i\phi$  we replace (12.6) by

$$PI(MINK) = \int D[\bar{\phi}]D[\sigma_\mu] \exp \left[ \frac{1}{2} \int_{-\infty}^{\infty} d^4x [i\partial_\nu \sigma_\mu \partial^\nu \sigma^\mu - i(M_1^2 + M_2^2) \sigma_\mu \sigma^\mu - iM_1^2 M_2^2 \bar{\phi}^2 - 2\epsilon \sigma_\mu \sigma^\mu - (M_1^2 + M_2^2) \epsilon \bar{\phi}^2] \right], \quad (12.8)$$

With  $\bar{\phi}$  and  $\sigma_\mu$  being taken to be real and with  $\sigma_\mu \sigma^\mu$  being taken to be timelike on every path, the path integral is now well defined and the theory is consistent.

In classical mechanics Poisson bracket relations are unaffected by symplectic transformations, while in quantum mechanics commutation relations are unaffected by similarity transformations. With path integrals being based on classical fields, the utility of making a symplectic transformation on the classical fields is that it can take a badly-behaved path integral with a real measure into a complex domain for the measure known as a Stokes wedge in which the path integral then is well behaved. Analogously, as described in Sec. VII, a similarity transformation can take us into a complex domain in which nonnormalizable quantum wave functions become normalizable. In both the classical and quantum cases we thus seek appropriate complex domains that have good boundary behavior.

While one is always free to make symplectic transformations or similarity transformations, for standard positive Dirac norm Hermitian theories with energies that are bounded from below, these transformations contain no new information. For such Hermitian theories the quantum fields are Hermitian, the classical fields that appear in the path integral are their real eigenvalues, the path integral with a real measure and Feynman  $i\epsilon$  prescription exists, the Dirac norm of the vacuum is finite, and the path integral can be associated with matrix elements of the form  $i\langle \Omega | T[\phi(x)\phi(0)] | \Omega \rangle$ .

However, it can happen that the path integral with real measure does not exist, but that it does exist for some appropriately chosen complex domain for the measure. In that case the Dirac norm of the vacuum is not finite. Also the Hamiltonian will not be Hermitian, but if all the poles in the propagator are real, the Hamiltonian will instead be  $PT$  symmetric. In that case the propagator is given by  $i\langle \Omega^L | T[(i\bar{\phi}(x))(i\bar{\phi}(0))] | \Omega^R \rangle = -i\langle \Omega^L | T[\bar{\phi}(x)\bar{\phi}(0)] | \Omega^R \rangle$ , where  $\phi$  is transformed into  $-i\phi = \bar{\phi}$ . And the vacuum norm is given by  $\langle \Omega^L | \Omega^R \rangle$ , a norm that is finite, positive definite and time independent. Thus when we find for the second-order plus fourth-order derivative quantum field theory that propagator pole residues are negative or that the path integral with real measure does not exist, it does not mean that the theory does not exist, but that it has to be formulated in a Hilbert space with an inner product other than the standard Dirac one.

Since for the second-order plus fourth-order derivative theory the vacuum Dirac norm is not finite, the standard derivation of the Feynman rules from the Wick contraction procedure is invalid, as is then the renormalizability that would follow from these Feynman rules. Interestingly, since the pole structure of the propagator is not affected by a similarity transformation, even after the transformation we can still use the propagator given in (12.3) with its  $-1/k^4$  short-distance behavior. The only thing that changes is the quantum field matrix element that we identify it with. This leads to the same Feynman rules, only now validly derived from a Wick contraction procedure in which everything is finite, to thus provide for an a posteriori derivation of renormalizability, while deriving it now in a Hilbert space that possesses no states with negative norm. The renormalizability and unitarity of the second-order plus fourth-order field theory is thus established, just as needed for a consistent theory of quantum gravity.

### XIII. FINAL COMMENTS

For a quantum field theory to be physically relevant it must be formulatable in a Hilbert space with an inner product that is time independent, finite and nonnegative. However, in and of itself, specifying an action and a set of canonical commutators is not enough to either fix the Hilbert space or specify the appropriate inner product. Ordinarily, one supplements these requirements with the additional (generally regarded as self-evident) requirements that the fields and the Hamiltonian of the theory be Hermitian, and that the inner product be the standard, presumed finite, Dirac  $\langle n|n \rangle$  one. (In the axiomatic approach to quantum field theory [26] the finiteness of  $\langle \Omega | \Omega \rangle$  is one of the starting assumptions.) However, it is not automatic for an arbitrary theory that  $\langle \Omega | \Omega \rangle$  be finite, and so one needs to check on a case by case basis. And in this paper we have presented a procedure for doing so. The procedure is based on using the occupation number space representation to construct an equivalent wave mechanics representation, from which we can then check for the normalizability of the vacuum state, and accordingly of the states that can be excited out

of it. An alternative but equivalent approach is to check whether or not the Minkowski time path integral with a real measure exists. If it does not, then the standard Dirac inner product is not finite.

Using the occupation number space representation procedure we have found a case, a second-order plus fourth-order scalar field theory, in which the standard Dirac inner product  $\langle n|n\rangle$  actually is not finite. In this example the Minkowski time path integral with a real measure diverges even though the Euclidean time path integral does not. In this case contributions from the Wick rotation contour cannot be ignored, and the use of a Euclidean time path integral can be misleading. Thus even if a Euclidean time path integral is well behaved, it only gives a good description of the theory if the Minkowski time path integral is well behaved too. Since  $\langle\Omega|\Omega\rangle$  is not finite for the second-order plus fourth-order scalar field theory, use of the standard Feynman rules is not valid, with these rules not only leading to states with negative norm, they lead to states with infinite negative norm. This lack of finiteness means that the Hamiltonian is not self-adjoint when acting on these particular states.

However, the Hamiltonian of the second-order plus fourth-order scalar field theory is  $PT$  symmetric, so we can use the techniques of the  $PT$  symmetry program and continue the fields and the Hamiltonian in this theory into the complex plane. There is then a domain in the complex plane in which one can define an appropriate time-independent, positive and finite inner product, viz. the  $\langle L|R\rangle$  overlap of left-eigenstates and right-eigenstates of the resulting Hamiltonian, with the resulting vacuum state then being normalizable, and with there being no states with negative or infinite  $\langle L|R\rangle$  norm [27]. In this domain it is the Euclidean time path integral that diverges while the Minkowski time path integral does not. So again there are contributions from the Wick rotation contour. In this domain the second-order plus fourth-order scalar field theory is fully consistent, unitary and renormalizable, with this analysis being relevant to the construction of a consistent, unitary and renormalizable quantum theory of gravity.

### Acknowledgments

The author wishes to thank Dr. C. M. Bender, Dr. J. Feinberg and Dr. O. Lechtenfeld for helpful comments.

- 
- [1] P. D. Mannheim, [arXiv:2209.15047](https://arxiv.org/abs/2209.15047). Eur. Phys. J. Plus (in press).
- [2] C. M. Bender and P. D. Mannheim, [Phys. Rev. D \*\*78\*\*, 025022 \(2008\)](https://arxiv.org/abs/1802.02502).
- [3] The difference in sign between the fourth-order (4.4) and the second-order (1.7) is due to the fact that  $[\phi, \ddot{\phi}]$  and  $[\phi, \dot{\phi}]$  as respectively given in (4.3) and (1.2) have opposite signs.
- [4] A. Pais and G. E. Uhlenbeck, [Phys. Rev. \*\*79\*\*, 145 \(1950\)](https://arxiv.org/abs/1950.0145).
- [5] P. D. Mannheim and A. Davidson, [arXiv:hep-th/0001115](https://arxiv.org/abs/hep-th/0001115).
- [6] P. D. Mannheim and A. Davidson, [Phys. Rev. A \*\*71\*\*, 042110 \(2005\)](https://arxiv.org/abs/2005.04211).
- [7] P. D. Mannheim, [Found. Phys. \*\*37\*\*, 532 \(2007\)](https://arxiv.org/abs/2007.0532).
- [8] C. M. Bender and P. D. Mannheim, [Phys. Rev. Lett. \*\*100\*\*, 110402 \(2008\)](https://arxiv.org/abs/2008.11040).
- [9] C. M. Bender and P. D. Mannheim, [Phys. Lett. A \*\*374\*\*, 1616 \(2010\)](https://arxiv.org/abs/2010.1616).
- [10] P. D. Mannheim, [J. Phys. A: Math. Theor. \*\*51\*\*, 315302 \(2018\)](https://arxiv.org/abs/2018.31530).
- [11] If  $AHA^{-1} = H$ , where  $A$  is a general antilinear operator and  $H|\psi\rangle = E|\psi\rangle$ , then  $AH|\psi\rangle = AE|\psi\rangle = E^*A|\psi\rangle = AHA^{-1}A|\psi\rangle = HA|\psi\rangle$ . Thus for every eigenvalue  $E$  there is an eigenvalue  $E^*$ , with  $H_{PU}$  and  $H_S$  being in the  $E = E^*$  realization in which all energies are real. While this analysis would apply to any antilinear symmetry, since it was first discovered in the  $PT$  context, even in the general case theories with antilinear symmetry are referred to as  $PT$  theories. As discussed in [10], under the requirements solely of complex Lorentz invariance and probability conservation one can show that a theory must be  $CPT$  invariant, where  $C$  denotes charge conjugation. For the theories discussed in this paper  $C$  is separately conserved, with  $CPT$  then defaulting to  $PT$ .
- [12] C. M. Bender, [Rep. Prog. Phys. \*\*70\*\*, 947 \(2007\)](https://arxiv.org/abs/2007.0947).
- [13] C. M. Bender, *PT Symmetry in Quantum And Classical Physics*. World Scientific, Singapore (2019).
- [14] The left-right inner product for general left-eigenstates and right-eigenstates of a Hamiltonian  $H$  obeys  $\langle L(t)|R(t)\rangle = \langle L(0)|e^{iHt}e^{-iHt}|R(0)\rangle = \langle L(0)|R(0)\rangle$ , to thus be time independent. When  $H$  is Hermitian then  $\langle L(t)| = \langle R(t)|$  and the left-right inner product reduces to the standard  $\langle R(t)|R(t)\rangle = \langle R(0)|R(0)\rangle$ . The left-right inner product is thus more general than the Dirac inner product as it also applies when the Hamiltonian is not Hermitian. As noted in [10], the left-right inner product is actually the most general inner product that can be time independent. Within the context of  $PT$  theory the left-right inner product is also the overlap of a ket with its  $PT$  theory conjugate bra  $\langle R(t)^{PT}|R(t)\rangle$  (see e.g. [10]).
- [15] S. W. Hawking and T. Hertog, [Phys. Rev. D \*\*65\*\*, 103515 \(2002\)](https://arxiv.org/abs/2002.10351).
- [16] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields*. McGraw-Hill, New York (1965).
- [17] A. A. Starobinsky, [JETP Lett. \*\*30\*\*, 682 \(1979\)](https://arxiv.org/abs/1979.030).
- [18] P. D. Mannheim, [Prog. Part. Nucl. Phys. \*\*56\*\*, 340 \(2006\)](https://arxiv.org/abs/2006.056).
- [19] K. S. Stelle, [Phys. Rev. D \*\*16\*\*, 953 \(1977\)](https://arxiv.org/abs/1977.0953).
- [20] K. S. Stelle, [Gen. Rel. Gravit. \*\*9\*\*, 353 \(1978\)](https://arxiv.org/abs/1978.0353).

- [21] P. D. Mannheim, *Prog. Part. Nucl. Phys.* **94**, 125 (2017).
- [22] G. 't Hooft, *Int. J. Mod. Phys. D* **24**, 1543001 (2015).
- [23] E. S. Fradkin and A. A. Tseytlin, *Phys. Rep.* **119**, 233 (1985).
- [24] As well as the quantum gravity problem, conformal gravity also addresses the dark matter and dark energy problems. It has been shown (see [21] and references therein) able to account for the systematics of galactic rotation curves without the need for any dark matter or its two free parameters per galactic dark matter halo, and able to account for the accelerating universe data without fine tuning. It is a theory in which local and global physics are connected, with it recently having been shown that there is an imprint of galactic rotation curves on the recombination era cosmic microwave background (P. D. Mannheim, [arXiv:2212.13942](https://arxiv.org/abs/2212.13942)).
- [25] G. 't Hooft, [arXiv:1009.0669](https://arxiv.org/abs/1009.0669) [gr-qc].
- [26] R. F. Streater and A. S. Wightman, *PCT, Spin and Statistics, and All That*, W. A. Benjamin, New York (1964).
- [27] While the axiomatic approach to quantum field theory was developed in [26] within the context of the Dirac inner product, it would be of interest to determine how many of the results presented in [26] might continue to hold if one uses the *PT*-theory inner product instead. A step in this direction has already been taken in [10], where it was shown that the *CPT* theorem could be derived without using the Dirac inner product or taking the Hamiltonian to be Hermitian.