

Massive Gravity and Lorentz Symmetry

R. Potting^{1,2}

¹*Departamento de Física, Faculdade de Ciências e Tecnologia,
Universidade do Algarve, 8005-139 Faro, Portugal*

²*Centro de Astrofísica e Gravitação, Instituto Superior Técnico,
Universidade de Lisboa, Avenida Rovisco Pais, 1049-001 Lisbon, Portugal*

We consider Lorentz-symmetry properties of the ghost-free massive gravity theory proposed by de Rham, Gabadadze, and Tolley. In particular, we present potentially observable effects in gravitational-wave propagation and in Newton's law, including Lorentz-violating signals.

1. The de Rham–Tolley–Gabadadze action

One of the motivations for considering massive gravity is the possibility that modifications of General Relativity over large distances may yield a solution to the cosmological constant problem. A ghost-free theory of non-interacting gravitons was constructed in 1939 by Fierz and Pauli.¹ However, attempts to generalize it to the nonlinear level failed during decades, with the work of Boulware and Deser showing that generically such theories will suffer from ghost instabilities.² Recently, however, de Rham, Gabadadze, and Tolley (dRGT) showed that there exists a non-linear extension of Fierz–Pauli massive gravity that does not suffer from ghosts.³ It can be shown that their model can be maximally extended to the action:⁴

$$S = \frac{1}{2\kappa} \int d^4x \sqrt{-g} \left(R - 2m^2 \sum_{n=0}^4 \beta_n e_n(\mathbb{X}) \right), \quad (1)$$

where R is the Ricci scalar, β_i are free parameters and $\kappa = 8\pi G$. The 4×4 matrix $\mathbb{X}^\mu{}_\nu$ equals $(\sqrt{g^{-1}}f)^\mu{}_\nu$, where, in addition to the usual physical metric $g_{\mu\nu}$, one assumes a given, nondynamical “fiducial” background metric $f_{\mu\nu}$. The invariant polynomials $e_n(\mathbb{X})$ are defined through the relation $\det(\mathbb{1} + \lambda\mathbb{X}) = \sum_{n=0}^4 \lambda^n e_n(\mathbb{X})$, yielding $e_0(\mathbb{X}) = 1$, $e_1(\mathbb{X}) = [\mathbb{X}]$, $e_2(\mathbb{X}) = \frac{1}{2}([\mathbb{X}]^2 - [\mathbb{X}^2])$, \dots

By expressing the metric in ADM form,⁵ using as the dynamical variables the spatial metric together with lapse and shift variables it can be shown that the equations of motion arising from the action (1) yield five local degrees of freedom, corresponding to the helicity states of a massive spin-2 particle. Crucial in the counting is the presence of the so-called Hamiltonian constraint.

2. Spacetime symmetries

In this talk I will report on recent work with Alan Kostelecký⁶ in which we studied the role of Lorentz symmetry in dRGT massive gravity. Defining the vierbein e_μ^a as usual through $g_{\mu\nu} = \eta_{ab}e_\mu^a e_\nu^b$, one can identify local Lorentz transformations as well as diffeomorphisms. As was explained in the talk by Alan Kostelecký at this conference, suitable local Lorentz transformations and diffeomorphisms can be combined to yield the so-called manifold Lorentz transformations, defined by

$$\begin{aligned} x^\mu &\rightarrow (\Lambda^{-1})^\mu{}_\nu x^\nu, & g_{\mu\nu} &\rightarrow (\Lambda^{-1})^\rho{}_\mu (\Lambda^{-1})^\sigma{}_\nu g_{\rho\sigma}, \\ e_\mu^a &\rightarrow (\Lambda^{-1})^\rho{}_\mu \Lambda^a{}_\rho e_\rho^b, & f_{\mu\nu} &\rightarrow f_{\mu\nu}, \end{aligned} \quad (2)$$

for spacetime-independent Lorentz transformations Λ . These are the analogues in approximately Minkowski spacetime of global Lorentz transformations in Minkowski spacetime. The dRGT action is invariant under manifold Lorentz transformations if $f_{\mu\nu} \propto \eta_{\mu\nu}$, but is otherwise Lorentz violating⁷ due to the presence of the background $f_{\mu\nu}$.

In Ref. [6] a careful study was done of the static solutions of the dRGT potential, for flat fiducial metrics. It was shown that the four-parameter potential has a highly nontrivial structure of extrema and saddle points, depending on the values of the parameters. Stability of these solutions was investigated by using the technique of bordered Hessians. The surface generated by the Hamiltonian constraint and the positions of the solutions on its connected sheets were used to establish global and absolute stability properties. We concluded that extrema of the potential are invariant under manifold Lorentz transformations, while the saddle point solutions are Lorentz violating, with maximally four broken Lorentz generators.

3. Linearized massive gravity

The action (1) yields the equations of motion

$$G_{\mu\nu} + \frac{m^2}{2} \sum_{n=0}^3 (-1)^n \beta_n \left(g_{\mu\alpha} Y_{(n)\nu}^\alpha + g_{\nu\alpha} Y_{(n)\mu}^\alpha \right) = \kappa T_{\mu\nu}, \quad (3)$$

where $Y_{(n)}(\mathbb{X}) = \sum_{k=0}^n (-1)^k \mathbb{X}^{n-k} e_k(\mathbb{X})$. Writing $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ and $f_{\mu\nu} = \eta_{\mu\nu} + \delta f_{\mu\nu}$ it follows that, to first order in $h_{\mu\nu}$ and $\delta f_{\mu\nu}$,

$$\mathbb{X} \approx \mathbb{1} + \frac{1}{2}\eta^{-1}\delta f - \frac{1}{2}\eta^{-1}h + \frac{1}{8}\eta^{-1}\delta f \eta^{-1}h - \frac{3}{8}\eta^{-1}h \eta^{-1}\delta f. \quad (4)$$

The equations of motion (3) become

$$\begin{aligned} G_{\mu\nu}^L + \frac{m^2}{2} \sum_{n=0}^3 \beta_n \left\{ 2 \binom{3}{n} (\eta_{\mu\nu} + h_{\mu\nu}) + \binom{2}{n-1} (h_{\mu\nu} - \delta f_{\mu\nu} - \eta_{\mu\nu} [h - \delta f]) \right. \\ + \left(\frac{1}{2} \binom{1}{n-2} + \binom{2}{n-1} \right) [\delta f] h_{\mu\nu} + \frac{1}{2} \binom{1}{n-2} [h] \delta f_{\mu\nu} \\ - \frac{1}{2} \binom{1}{n-1} [\delta f \eta^{-1} h] \eta_{\mu\nu} - \frac{1}{2} \binom{1}{n-2} [\delta f] [h] \eta_{\mu\nu} \\ \left. - \left(\frac{3}{4} \binom{2}{n-1} - \frac{1}{2} \binom{1}{n-1} \right) (h \eta^{-1} \delta f + \delta f \eta^{-1} h)_{\mu\nu} \right\} = \kappa T_{\mu\nu}, \quad (5) \end{aligned}$$

where $G_{\mu\nu}^L$ is the linearized Einstein tensor and $[X] = \eta^{\mu\nu} X_{\mu\nu}$. It is usual to require that $h_{\mu\nu} = \delta f_{\mu\nu} = T_{\mu\nu} = 0$ satisfies the equations of motion (5), yielding the constraint $\beta_0 + 3\beta_1 + 3\beta_2 + \beta_3 = 0$. Moreover, we normalize the mass m such that $\beta_1 + 2\beta_2 + \beta_3 = 1$.

Consider now a nontrivial fiducial background metric $\delta f_{\mu\nu} \neq 0$, it follows that spacetime now has nonzero curvature in the absence of matter! In order to simplify the further analysis we will assume a special constant background energy–momentum tensor

$$\kappa T_{\mu\nu} = -\frac{m^2}{2} (\delta f_{\mu\nu} - \eta_{\mu\nu} [\delta f]). \quad (6)$$

For this special choice, $h_{\mu\nu} = 0$ solves the equations of motion.

4. Gravitational waves

In order to investigate the propagation of gravitational waves, we define the Fourier transform of $h_{\mu\nu}$ through $h_{\mu\nu}(x) = (2\pi)^{-4} \int d^4p e^{-ip \cdot x} \tilde{h}_{\mu\nu}(p)$. For $\delta f_{\mu\nu} = 0$, $\tilde{h}_{\mu\nu}$ satisfies the conditions $\tilde{h}_{\mu\alpha} p^\alpha = 0$ and $\tilde{h}_\mu{}^\mu = 0$, thus yielding $10 - 5 = 5$ propagating modes. These can be identified with the helicity eigenstates of a massive spin-two field, $\tilde{h}_{\mu\nu}^{(n)}$, with $n = 0, \pm 1, \pm 2$. All modes satisfy the massive dispersion relation $(p^2 + m^2) \tilde{h}_{\mu\nu}^{(n)} = 0$.

For nonzero $\delta f_{\mu\nu}$ the situation becomes more complicated. The equations of motion (5) can be cast in the form

$$(p^2 + m^2) \tilde{h}_{\mu\nu} = \frac{c_2 m^2}{2} S_{\mu\nu}{}^{\alpha\beta} \tilde{h}_{\alpha\beta} \quad (7)$$

where $c_2 = \sum_n \beta_n \left(\binom{1}{n-1} - \frac{3}{2} \binom{2}{n-1} \right)$. The quantities $S_{\mu\nu}^{\alpha\beta}$ are tensor coefficients depending on the momentum p^μ and on $\delta f_{\mu\nu}$. It is convenient to expand them in a momentum-dependent orthonormal basis spanned by p^μ and three other, spacelike vectors. It is then straightforward to work out the expressions $S_{\mu\nu}^{\alpha\beta} \tilde{h}_{\alpha\beta}^{(n)}$ for any helicity mode n , which are well-defined linear combinations of the five helicity modes.

Equation (7) can be solved by constructing the eigenstates of $S_{\mu\nu}^{\alpha\beta}$. For general δf , we find a ‘‘pentarefringence’’ effect: each of these eigenstates solves Eq. (7) with a (slightly) different dispersion relation. These pentarefringence effects are momentum (and direction) dependent. The modes can be sub- or superluminal, a result typical of Lorentz-violating theories. For details, see Ref. [6].

5. Corrections to Newton’s law

Next we study the effects of the extra terms in the equation of motion (5) on Newton’s law. Writing the momentum-space linearized modified Einstein equation as $\tilde{O}_{\mu\nu}^{\alpha\beta} \tilde{h}^{\alpha\beta} = 0$, the corresponding propagator is defined to satisfy $\tilde{D}_{\mu\nu}^{\sigma\tau} \tilde{O}_{\sigma\tau}^{\alpha\beta} = \delta_{(\mu}^{\alpha} \delta_{\nu)}^{\beta}$. At first order in δf it has the form

$$\begin{aligned} \tilde{D}_{\mu\nu}^{\alpha\beta} = & \frac{1}{p^2 + m^2} \left[\delta_{(\mu}^{\alpha} \delta_{\nu)}^{\beta} - \frac{1}{3} \eta_{\mu\nu} \eta^{\alpha\beta} + \frac{2}{m^2} p_{(\mu} p^{\alpha} \delta_{\nu)}^{\beta} \right. \\ & \left. - \frac{1}{3m^2} \left(p_\mu p_\nu \eta^{\alpha\beta} + \eta_{\mu\nu} p^\alpha p^\beta - \frac{2}{m^2} p_\mu p_\nu p^\alpha p^\beta \right) \right] \\ & - \frac{m^2}{(p^2 + m^2)^2} \left[\rho_1 \delta f_{(\mu}^{\alpha} \delta_{\nu)}^{\beta} + \rho_2 \delta f_{\mu\nu} \eta^{\alpha\beta} + \rho_4 p_\mu p_\nu \delta f^{\alpha\beta} + \dots \right], \end{aligned} \quad (8)$$

where $\rho_i = \rho_i(p^2)$ are momentum-dependent scalars (parentheses in the indices indicate symmetrization). For given energy–momentum tensor the linearized solution for the metric can then be expressed as

$$h_{\mu\nu}(x) = 2\kappa \int \frac{d^4 p}{(2\pi)^4} e^{-ip \cdot x} D_{\mu\nu}^{\alpha\beta} \tilde{T}_{\alpha\beta}(p), \quad (9)$$

where $T^{\mu\nu}(x) = (2\pi)^{-4} \int d^4 p e^{-ip \cdot x} \tilde{T}^{\mu\nu}(p)$.

Consider now the gravitational potential energy between two stationary point masses with energy–momentum tensors

$$T_1^{\mu\nu}(x) = M_1 \delta_0^\mu \delta_0^\nu \delta^3(\vec{x}), \quad T_2^{\mu\nu}(x) = M_2 \delta_0^\mu \delta_0^\nu \delta^3(\vec{x} - \vec{r}). \quad (10)$$

At linear order in $h_{\mu\nu}$ the matter Lagrangian corresponding to a given energy–momentum tensor is $\mathcal{L}_m \approx -\frac{1}{2} h^{\mu\nu} T_{\mu\nu}$. From this it follows that

the potential energy corresponding to the energy–momentum tensors (10) is given by $U(\vec{r}) = \int d^3x h_1^{\mu\nu}(\vec{x})T_{2,\mu\nu}(\vec{x}) = M_2 h_{1,00}(\vec{r})$. Using Eqs. (8) and (9) we obtain

$$\begin{aligned}
 U(\vec{r}) = & -\frac{GM_1M_2e^{-mr}}{9r} \left[24 - \delta f_{00}((4c_2 + 9)mr + 8c_2 + 8) \right. \\
 & - \delta f_i^i \left((4c_2 - 9)mr + 2c_2 + 4 + \frac{2}{mr} + \frac{4}{m^2r^2} \right) \\
 & \left. - \frac{\delta f_{ij}x^i x^j}{r^2} \left(2c_2mr + 2c_2 - 4 - \frac{6}{mr} - \frac{12}{m^2r^2} \right) \right] \\
 & + \frac{2\kappa M_1M_2}{9m^2} \delta^3(\vec{r}) \left[\delta f_{00} - \frac{2}{3}\delta f_i^i \right]. \tag{11}
 \end{aligned}$$

The exponential suppression factor e^{-mr} is as expected due to the massive graviton. The term independent of $\delta f_{\mu\nu}$ is scaled by 4/3 relative to the gravitational potential in General Relativity, in concordance with the van Dam–Veltman–Zakharov discontinuity. Moreover, note that $U(\vec{r})$ acquires terms that generally violate rotational invariance.

Acknowledgments

This work was supported in part by the Portuguese Fundação para a Ciência e a Tecnologia under grants SFRH/BSAB/150324/2019 and UID/FIS/00099/2019, and by the Indiana University Center for Spacetime Symmetries.

References

1. M. Fierz, *Helv. Phys. Acta* **12**, 3 (1939); M. Fierz and W. Pauli, *Proc. Roy. Soc. (London)* **A 173**, 211 (1939).
2. D.G. Boulware and S. Deser, *Phys. Rev. D* **6**, 3368 (1972).
3. C. de Rham, G. Gabadadze, and A.J. Tolley, *Phys. Rev. Lett.* **106**, 231101 (2011); *Phys. Lett. B* **711**, 190 (2012).
4. S.F. Hassan and R.A. Rosen, *JHEP* **04**, 123 (2012); *Phys. Rev. Lett.* **108**, 041101 (2012); S.F. Hassan, R.A. Rosen and A. Schmidt-May, *JHEP* **02**, 026 (2012).
5. R. Arnowitt, S. Deser, and C. Misner, *Phys. Rev.* **116**, 1322 (1959).
6. V.A. Kostelecký and R. Potting, *Phys. Rev. D* **104**, 104046 (2021).
7. V.A. Kostelecký, *Phys. Rev. D* **69**, 105009 (2004).
8. H. van Dam and M. Veltman, *Nucl. Phys.* **22**, 397 (1970).
9. V.I. Zakharov, *JETP Lett. (Sov. Phys.)*, **12**, 312 (1970).