

A peak in the power spectrum of primordial gravitational waves induced by primordial magnetic fields

Sugumi Kanno^{*}, Ann Nakato[†], Jiro Soda^{†,‡}, and Kazushige Ueda^{*}

^{*} *Department of Physics, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka 819-0395, Japan*

[†] *Department of Physics, Kobe University, Kobe 657-8501, Japan*

[‡] *International Center for Quantum-field Measurement Systems for Studies of the Universe and Particles (QUP), KEK, Tsukuba 305-0801, Japan*

Abstract

It has been argued that there exist extragalactic magnetic fields in the range from 10^{-17} to 10^{-9} Gauss with a cosmological coherence length. One plausible explanation for the origin of the extragalactic magnetic fields would be quantum fluctuations of the electromagnetic fields during inflation. It is also believed that primordial gravitational waves (PGWs) arise out of quantum fluctuations during inflation. We study the graviton-photon conversion process in the presence of background magnetic fields and find that the process induces the tachyonic instability of the PGWs. As a consequence, a peak appears in the power spectrum of PGWs. It turns out that the peak height depends on the direction of observation. The peak frequency could be in the range from 10^{-5} to 10^3 Hertz for GUT scale inflation. Hence, the observation of PGWs could provide a new window for probing primordial magnetic fields.

Contents

1	Introduction	1
2	Graviton-photon interaction in the early universe	3
2.1	Primordial GWs	4
2.2	Primordial magnetic fields	4
2.3	Graviton-photon conversion	5
2.4	Diagonal equations	7
3	Generation and evolution of primordial GWs	8
3.1	Cosmological setup	8
3.2	Bogoliubov Transformations	10
4	Tachyonic instability and the PGW power spectrum	12
5	Conclusion	15

1 Introduction

Since the first direct detection of gravitational waves [1], gravitational waves have been the most important tool for investigating fundamental physics. Gravitational waves interact with matter very weakly and travel through the universe virtually unimpeded, so they can be a probe of the early inflationary universe. The inflationary scenario predicts that primordial gravitational waves (PGWs) are generated from quantum fluctuations of spacetime. In terms of the density parameter, the spectrum $\Omega_{\text{GW}}(f)$ for the GUT scale inflation is independent of frequency f , that is, of wave number k for the gravitons created at the de Sitter to radiation dominant transition:

$$\Omega_{\text{GW}}(f) = 10^{-14} \left(\frac{H}{10^{-4} M_{\text{pl}}} \right)^2, \quad (1.1)$$

where H is the Hubble parameter, M_{pl} is the reduced Planck mass. This scale-invariant power spectrum comes from the time translation invariance of de Sitter space. There are several experimental projects for detecting PGWs [2, 3, 4, 5]. If the nonclassicality of the PGWs is found, it implies the existence of gravitons. Notably, due to the particle creation, the quantum state of gravitons becomes squeezed during inflation [6, 7, 8, 9, 10]. Since the squeezing tends to enhance the observability of gravitons, it is expected that we may be able to probe quantum gravity through observations of PGWs. There are new ideas for detecting the quantum nature of PGWs. One method is to utilize the Hanbury Brown-Twiss interferometry developed in quantum optics for the PGWs, which can distinguish nonclassical particles from classical ones by measuring intensity-intensity correlations [11, 12, 13]. Moreover, the squeezed state of gravitons can be measured indirectly through their noise in the interferometers [14, 15, 16, 17] or by measuring the decoherence time of a quantum object caused by the surrounding primordial gravitons [18].

On top of the PGWs, there might exist primordial magnetic fields (PMFs). Indeed, it is argued that observations of GeV gamma-ray from blazars give rise to the lower bound of the extragalactic magnetic field 10^{-17} Gauss [19, 20, 21, 22, 23, 24]. We also have the upper bound 10^{-9} Gauss from CMB data [25, 26]. It is important to seek the origin of the extragalactic magnetic fields. If a coherence length of magnetic fields is around mega-parsec scales, it would be natural to consider magnetic fields originated from quantum fluctuations during inflation [27, 28]. In the presence of PMFs during inflation, the effect of the PMFs on PGWs is studied in [29]. In the case of scale-invariant electromagnetic fields, there arises an entanglement between gravitons and photons [30]. Therefore, the presence of scale-invariant electromagnetic fields may affect the squeezed state of gravitons. Thus, it is important to clarify the effect of PMFs on PGWs.

In this paper, we consider the background PMFs during inflation and the scale-invariant perturbed electromagnetic fields. It is known that the presence of background magnetic fields causes the conversion of gravitons into photons and vice versa [31, 32]. We focus on the effect of the graviton-photon conversion process on the PGW power spectrum. We show that there occurs the tachyonic instability of PGWs due to graviton-photon conversion in the presence of PMFs. Because of this instability, there arises a peak in the PGW spectrum

(1.1). Moreover, since the magnetic fields specify a direction in the space, there appears statistical anisotropy in the power spectrum of PGWs. Remarkably, it turns out that the peak frequency could be in the range from 10^{-5} to 10^3 Hertz. The presence of the peak turns out to enhance the squeezing of gravitons and makes it easy to observe the PGWs. More interestingly, we may be able to probe the PMFs by observing the spectrum of PGWs.

The organization of the paper is as follows. In section 2, we present the setup. In section 3, we calculate Bogoliubov coefficients. In section 4, we calculate the PGW power spectrum. We see the tachyonic instability leads to a peak in the spectrum. We also discuss the observability of the peak. Section 5 is devoted to the conclusion.

2 Graviton-photon interaction in the early universe

We consider the Einstein-Hilbert action and the action for a $U(1)$ gauge field coupled with a scalar field:

$$S = S_g + S_\phi + S_A = \int d^4x \sqrt{-g} \left[\frac{M_{\text{pl}}^2}{2} R - \frac{1}{2} (\partial_\mu \phi) (\partial^\mu \phi) - V(\phi) - \frac{1}{4} f^2(\phi) F^{\mu\nu} F_{\mu\nu} \right], \quad (2.1)$$

where $M_{\text{pl}} = 1/\sqrt{8\pi G}$ is the Planck mass. The gauge field A_μ represents photons and the field strength is defined by $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$.

The background inflationary dynamics is determined by the metric

$$ds^2 = a^2(\eta) [-d\eta^2 + \delta_{ij} dx^i dx^j], \quad (2.2)$$

and the inflaton $\phi(\eta)$. Once the background is given, the coupling function can be regarded as a function of the conformal time η ; $f = f(\eta)$. We also assume the presence of constant magnetic fields $B_i = \text{constant}$. It should be emphasized that the physical magnetic fields are not B_i but fB_i . In the following, we consider the quantum evolution of gravitons and photons in the above background.

2.1 Primordial GWs

We consider gravitons in a spatially flat expanding background represented by tensor mode perturbations in the three-dimensional metric h_{ij} ,

$$ds^2 = a^2(\eta) [-d\eta^2 + (\delta_{ij} + h_{ij}) dx^i dx^j] , \quad (2.3)$$

where h_{ij} satisfies the transverse traceless conditions $h_{ij}{}^{;j} = h^i{}_i = 0$. The spatial indices i, j, k, \dots are raised and lowered by δ^{ij} and δ_{kl} . In the case of de Sitter space, the scale factor is given by $a(\eta) = -1/(H\eta)$ where $-\infty < \eta < 0$.

Expanding the Einstein-Hilbert action up to the second order in perturbations h_{ij} , we have

$$\delta S_g = \frac{M_{\text{pl}}^2}{8} \int d^4x a^2 [h^{ij'} h'_{ij} - h^{ij,k} h_{ij,k}] , \quad (2.4)$$

where a prime denotes the derivative with respect to the conformal time. At this quadratic order of the action, it is convenient to expand $h_{ij}(\eta, x^i)$ in Fourier modes,

$$h_{ij}(\eta, x^i) = \frac{2}{M_{\text{pl}}} \sum_P \frac{1}{(2\pi)^{3/2}} \int d^3k h_{\mathbf{k}}^P(\eta) e_{ij}^P(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} , \quad (2.5)$$

where three-vectors are denoted by bold math type and $e_{ij}^P(\mathbf{k})$ are the polarization tensors for the \mathbf{k} mode normalized as $e^{ijP}(\mathbf{k}) e_{ij}^Q(\mathbf{k}) = \delta^{PQ}$ with $P, Q = +, \times$. Then the action (2.4) in the Fourier space becomes

$$\delta S_g = \frac{1}{2} \sum_P \int d^3k d\eta a^2 [|h_{\mathbf{k}}^{P'}|^2 - k^2 |h_{\mathbf{k}}^P|^2] . \quad (2.6)$$

2.2 Primordial magnetic fields

Next, we consider the action for the photon up to the second order in perturbations A_i , which is given by

$$\delta S_A = \frac{1}{2} \int d^4x f^2 [A_i'^2 - A_{k,i}^2] , \quad (2.7)$$

where the photon field satisfies the Coulomb gauge $A_0 = 0$ and $A^i{}_{,i} = 0$.

Expanding the field $A_i(\eta, x^i)$ by the Fourier modes, we find

$$A_i(\eta, x^i) = \sum_P \frac{\pm i}{(2\pi)^{3/2}} \int d^3k A_{\mathbf{k}}^P(\eta) e_i^P(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{x}} , \quad (2.8)$$

where $e_i^P(\mathbf{k})$ are the polarization vectors for the \mathbf{k} mode normalized as $e^{iP}(\mathbf{k})e_i^Q(\mathbf{k}) = \delta^{PQ}$ with $P, Q = +, \times$. The sign of $\pm i$ corresponds to the $P, Q = +, \times$. The action (2.7) in terms of the Fourier modes reads

$$\delta S_A = \frac{1}{2} \sum_P \int d^3k d\eta f^2 [|A_{\mathbf{k}}^{P'}|^2 - k^2 |A_{\mathbf{k}}^P|^2] . \quad (2.9)$$

2.3 Graviton-photon conversion

The action for the interaction between the graviton and the photon up to second order in perturbations h_{ij}, A^i is found to be

$$\delta S_I = \int d^4x [\varepsilon_{ilm} f^2 B_m h^{ij} (\partial_j A_\ell - \partial_\ell A_j)] , \quad (2.10)$$

where $B_m = \varepsilon_{mj\ell} \partial_j A_\ell$ is a constant background magnetic field.

In terms of the Fourier modes defined in Eqs. (2.5) and (2.8), the interaction term reads

$$\delta S_I = \frac{2}{M_{\text{pl}}} \sum_{P,Q} \int d^3k d\eta f^2 [\varepsilon_{ilm} B_m h_{\mathbf{k}}^P A_{-\mathbf{k}}^Q e_{ij}^P(\mathbf{k}) \{ ik_\ell e_j^Q(-\mathbf{k}) - ik_j e_\ell^Q(-\mathbf{k}) \}] , \quad (2.11)$$

where $k = |\mathbf{k}|$. Polarization vectors $e^{i+}, e^{i\times}$ and a vector k^i/k constitute an orthonormal basis. Without loss of generality, we assume the constant background magnetic field is in the $(k^i, e^{i\times})$ -plane as depicted in FIG. 1.




FIG 1. Configuration of the polarization vector $e^P(\mathbf{k})$, wave number \mathbf{k} , and background magnetic field \mathbf{B} . The angle between the magnetic field and the wavenumber vector is represented by θ .

The polarization tensors can be written in terms of polarization vectors e^{i+} and $e^{i\times}$ as

$$e_{ij}^+(\mathbf{k}) = \frac{1}{\sqrt{2}} \left\{ e_i^+(\mathbf{k}) e_j^+(\mathbf{k}) - e_i^\times(\mathbf{k}) e_j^\times(\mathbf{k}) \right\}, \quad (2.12)$$

$$e_{ij}^\times(\mathbf{k}) = \frac{1}{\sqrt{2}} \left\{ e_i^+(\mathbf{k}) e_j^\times(\mathbf{k}) + e_i^\times(\mathbf{k}) e_j^+(\mathbf{k}) \right\}. \quad (2.13)$$

Below, we assume $e_i^\times(-\mathbf{k}) = -e_i^\times(\mathbf{k})$. The action (2.11) is then written as

$$\delta S_I = \int d^3k d\eta f^2 \lambda k \left[h_{\mathbf{k}}^+(\eta) A_{-\mathbf{k}}^+(\eta) + h_{\mathbf{k}}^\times(\eta) A_{-\mathbf{k}}^\times(\eta) \right], \quad (2.14)$$

where we defined the coupling between graviton and photon as

$$\lambda \equiv \frac{\sqrt{2}}{M_{\text{pl}}} \varepsilon^{ilm} e_i^+ \frac{k_\ell}{k} B_m. \quad (2.15)$$

Here, the conditions for the graviton and photon to be real are read, $h_{-\mathbf{k}}^{+,\times}(\eta) = h_{\mathbf{k}}^{*+,\times}(\eta)$ and $A_{-\mathbf{k}}^{+,\times}(\eta) = -A_{\mathbf{k}}^{*+,\times}(\eta)$. Below, we focus on the plus polarization and omit the index P unless there may be any confusion.

2.4 Diagonal equations

If we use the canonical variable $y_{\mathbf{k}}(\eta) = a h_{\mathbf{k}}(\eta)$ and $x_{\mathbf{k}}(\eta) = f A_{\mathbf{k}}(\eta)$, the total action of Eqs. (2.6), (2.9) and (2.14) are written as

$$\begin{aligned}
\delta S &= \delta S_y + \delta S_x + \delta S_I \\
&= \frac{1}{2} \int d^3k d\eta \left[|y'_{\mathbf{k}}|^2 - \left(k^2 - \left(\frac{a'}{a} \right)^2 \right) |y_{\mathbf{k}}|^2 - \frac{a'}{a} (y_{\mathbf{k}} y'_{-\mathbf{k}} + y_{-\mathbf{k}} y'_{\mathbf{k}}) \right] \\
&\quad + \frac{1}{2} \int d^3k d\eta \left[|x'_{\mathbf{k}}|^2 - \left(k^2 - \left(\frac{f'}{f} \right)^2 \right) |x_{\mathbf{k}}|^2 - \frac{f'}{f} (x_{\mathbf{k}} x'_{-\mathbf{k}} + x_{-\mathbf{k}} x'_{\mathbf{k}}) \right] \\
&\quad + \int d^3k d\eta \left[\frac{f}{a} \lambda k y_{\mathbf{k}} x_{-\mathbf{k}} \right]. \tag{2.16}
\end{aligned}$$

The variation of the action (2.16) with respect to the graviton and the photon fields gives

$$y''_{\mathbf{k}} + \left(k^2 - \frac{a''}{a} \right) y_{\mathbf{k}} = -\lambda k f \frac{x_{\mathbf{k}}}{a}, \quad x''_{\mathbf{k}} + \left(k^2 - \frac{f''}{f} \right) x_{\mathbf{k}} = -\lambda k f \frac{y_{\mathbf{k}}}{a}. \tag{2.17}$$

We assume the gauge kinetic function in the form

$$f(\eta) = a(\eta)^{-2c}, \tag{2.18}$$

where c is a constant parameter. We take $c = -1/2$ to make the analysis easier. For this parameter, the power spectrum of the electromagnetic fields A_{μ} is scale-invariant [33]. While the spectrum of magnetic fields becomes $P(B) \propto k^2$. Then Eqs. (2.17) become

$$y''_{\mathbf{k}} + \left(k^2 - \frac{a''}{a} \right) y_{\mathbf{k}} = -\lambda k x_{\mathbf{k}}, \quad x''_{\mathbf{k}} + \left(k^2 - \frac{a''}{a} \right) x_{\mathbf{k}} = -\lambda k y_{\mathbf{k}}. \tag{2.19}$$

If we define new variables $X_{\mathbf{k}}$ and $Y_{\mathbf{k}}$ such as

$$Y_{\mathbf{k}} = \frac{1}{\sqrt{2}} (y_{\mathbf{k}} + x_{\mathbf{k}}), \quad X_{\mathbf{k}} = \frac{1}{\sqrt{2}} (y_{\mathbf{k}} - x_{\mathbf{k}}). \tag{2.20}$$

Eq. (2.19) are diagonalized in the form

$$Y_{\mathbf{k}}'' + \left(k^2 + \lambda k - \frac{a''}{a} \right) Y_{\mathbf{k}} = 0 \quad , \quad X_{\mathbf{k}}'' + \left(k^2 - \lambda k - \frac{a''}{a} \right) X_{\mathbf{k}} = 0 \quad . \quad (2.21)$$

3 Generation and evolution of primordial GWs

In this section, we present a cosmological setup in order to calculate Bogoliubov coefficients.

3.1 Cosmological setup

We consider instantaneous reheating after inflation approximated by the de Sitter phase leading to a radiation-dominated phase. The scale factor evolves as follows

$$a(\eta) = \begin{cases} \frac{1}{-H(\eta-2\eta_1)} & \text{for } \eta < \eta_1 \text{ ,} \\ \frac{\eta}{H\eta_1^2} & \text{for } 0 < \eta_1 < \eta \text{ .} \end{cases} \quad (3.1)$$

where H is the Hubble parameter during inflation and η_1 is the time of reheating. The scale factor is smoothly connected up to the first order of derivative at the reheating. We assume that magnetic fields arise instantaneously from the Bunch-Davies vacuum at the time η_* .




FIG 2. During inflation, magnetic fields B_i arise instantaneously at η_* . Instantaneous reheating at η_1 leads to a radiation-dominated phase.

Before the time η_* (phase I), the equations of motion read

$$Y_{\mathbf{k}}'' + \left(k^2 - \frac{2}{(\eta - 2\eta_1)^2} \right) Y_{\mathbf{k}} = 0, \quad X_{\mathbf{k}}'' + \left(k^2 - \frac{2}{(\eta - 2\eta_1)^2} \right) X_{\mathbf{k}} = 0. \quad (3.2)$$

After the η_* , the coupling λ is switched on until η_1 (phase II). There, gravitons and photons are coupled as

$$Y_{\mathbf{k}}'' + \left(k^2 + \lambda k - \frac{2}{(\eta - 2\eta_1)^2} \right) Y_{\mathbf{k}} = 0, \quad X_{\mathbf{k}}'' + \left(k^2 - \lambda k - \frac{2}{(\eta - 2\eta_1)^2} \right) X_{\mathbf{k}} = 0. \quad (3.3)$$

After the reheating time η_1 (phase III), we assume $f = \text{const}$. Then, the equations of motion become

$$Y_{\mathbf{k}}'' + k^2 Y_{\mathbf{k}} = 0, \quad X_{\mathbf{k}}'' + k^2 X_{\mathbf{k}} = 0. \quad (3.4)$$

We assume the vacuum state can be specified by the Bunch-Davies vacuum which behaves in the same way as the Minkowski vacuum in the remote past in the de Sitter phase (Phase I). In the Phase II, the basis is properly normalized but deformed due to the presence of the magnetic field. In the Phase III, we choose the positive frequency mode for the adiabatic vacuum as a basis in radiation dominated phase in order to discuss particle creation. Thus we choose the properly normalized basis in each phase as the solution of Eqs. (3.2), (3.3) and (3.4).

$$u_{\mathbf{k}} = \frac{1}{\sqrt{2k}} \left(1 - \frac{i}{k(\eta - 2\eta_1)} \right) e^{-ik(\eta - 2\eta_1)} \quad \text{for Phase I} \quad (3.5)$$

$$w_{\mathbf{k}}^+ = \frac{1}{\sqrt{2\sqrt{k^2 + \lambda k}}} \left(1 - \frac{i}{\sqrt{k^2 + \lambda k}(\eta - 2\eta_1)} \right) e^{-i\sqrt{k^2 + \lambda k}(\eta - 2\eta_1)} \quad \text{for Phase II} \quad (3.6)$$

$$w_{\mathbf{k}}^- = \frac{1}{\sqrt{2\sqrt{k^2 - \lambda k}}} \left(1 - \frac{i}{\sqrt{k^2 - \lambda k}(\eta - 2\eta_1)} \right) e^{-i\sqrt{k^2 - \lambda k}(\eta - 2\eta_1)} \quad \text{for Phase II} \quad (3.7)$$

$$v_{\mathbf{k}} = \frac{1}{\sqrt{2k}} e^{-ik\eta} \quad \text{for Phase III} \quad (3.8)$$

The above mode functions are normalized as

$$u_{\mathbf{k}} u_{\mathbf{k}}'^* - u_{\mathbf{k}}^* u_{\mathbf{k}}' = i, \quad w_{\mathbf{k}}^{\pm} w_{\mathbf{k}}^{\pm*'} - w_{\mathbf{k}}^{\pm*} w_{\mathbf{k}}^{\pm'} = i, \quad v_{\mathbf{k}} v_{\mathbf{k}}'^* - v_{\mathbf{k}}^* v_{\mathbf{k}}' = i. \quad (3.9)$$

3.2 Bogoliubov Transformations

Through the history from Phase I to III, the fields $X_{\mathbf{k}}$ and $Y_{\mathbf{k}}$ in Eqs. (2.21) can be expanded by the mode functions $U_{\mathbf{k}}$ and $V_{\mathbf{k}}$, respectively such as

$$Y_{\mathbf{k}} = \hat{a}_{\mathbf{k}} U_{\mathbf{k}} + \hat{a}_{-\mathbf{k}}^{\dagger} U_{\mathbf{k}}^* , \quad X_{\mathbf{k}} = \hat{b}_{\mathbf{k}} V_{\mathbf{k}} + \hat{b}_{-\mathbf{k}}^{\dagger} V_{\mathbf{k}}^* . \quad (3.10)$$

The commutation relations $[\hat{a}_{\mathbf{k}}, \hat{a}_{\mathbf{k}'}^{\dagger}] = \delta(\mathbf{k} - \mathbf{k}')$ and $[\hat{b}_{\mathbf{k}}, \hat{b}_{\mathbf{k}'}^{\dagger}] = \delta(\mathbf{k} - \mathbf{k}')$ guarantee the canonical commutation relations. We write the mode function $U_{\mathbf{k}}/V_{\mathbf{k}}$ that satisfies Eqs. (3.2), (3.3) and (3.4) as $U_{\mathbf{k}}^{\text{I}}/V_{\mathbf{k}}^{\text{I}}$, $U_{\mathbf{k}}^{\text{II}}/V_{\mathbf{k}}^{\text{II}}$ and $U_{\mathbf{k}}^{\text{III}}/V_{\mathbf{k}}^{\text{III}}$.

As the initial state in Phase I, we consider the Bunch-Davies vacuum $\hat{a}_{\mathbf{k}}|0\rangle_{\text{BD}} = 0$, $\hat{b}_{\mathbf{k}}|0\rangle_{\text{BD}} = 0$. Then the mode functions $U_{\mathbf{k}}$ and $V_{\mathbf{k}}$ are expressed by the positive frequency mode Eq. (3.5):

$$U_{\mathbf{k}}^{\text{I}} = V_{\mathbf{k}}^{\text{I}} = u_{\mathbf{k}}(\eta). \quad (3.11)$$

In phase II, the mode function $U_{\mathbf{k}}$ and $V_{\mathbf{k}}$ are written by the solutions Eqs. (3.6) and (3.7) of Eqs. (3.3) and their complex conjugate,

$$U_{\mathbf{k}}^{\text{II}} = \alpha_{\text{IIk}}^+ w_{\mathbf{k}}^+(\eta) + \beta_{\text{IIk}}^+ w_{\mathbf{k}}^{+*}(\eta) , \quad (3.12)$$

$$V_{\mathbf{k}}^{\text{II}} = \alpha_{\text{IIk}}^- w_{\mathbf{k}}^-(\eta) + \beta_{\text{IIk}}^- w_{\mathbf{k}}^{-*}(\eta) , \quad (3.13)$$

where $\alpha_{\text{IIk}}^{\pm}$ and β_{IIk}^{\pm} are integration constants which can be interpreted as the Bogoliubov coefficients. Similarly, in phase III, the mode functions are written by the solution Eq. (3.8) of Eq. (3.4):

$$U_{\mathbf{k}}^{\text{III}} = \alpha_{\text{IIIk}}^+ v_{\mathbf{k}}(\eta) + \beta_{\text{IIIk}}^+ v_{\mathbf{k}}^*(\eta) , \quad (3.14)$$

$$V_{\mathbf{k}}^{\text{III}} = \alpha_{\text{IIIk}}^- v_{\mathbf{k}}(\eta) + \beta_{\text{IIIk}}^- v_{\mathbf{k}}^*(\eta) , \quad (3.15)$$

where $\alpha_{\text{IIIk}}^{\pm}$ and $\beta_{\text{IIIk}}^{\pm}$ are integration constants interpreted as the Bogoliubov coefficients.

Since the solutions of $Y_{\mathbf{k}}$ and $X_{\mathbf{k}}$ must be continuous and continuously differentiable, the boundary conditions for the above mode functions are required at η_* and η_1 respectively.

Let us focus on $Y_{\mathbf{k}}$. The boundary conditions at η_* are given by,

$$U_{\mathbf{k}}^{\text{I}}(\eta_*) = U_{\mathbf{k}}^{\text{II}}(\eta_*) \quad (3.16)$$

$$U_{\mathbf{k}}^{\text{I}' }(\eta) \Big|_{\eta=\eta_*} = U_{\mathbf{k}}^{\text{II}' }(\eta) \Big|_{\eta=\eta_*} \quad (3.17)$$

The same is true for the $V_{\mathbf{k}}^{\text{I}}(\eta_0)$. Plugging Eqs. (3.11) and (3.12) into the above, we have

$$\begin{pmatrix} u(\eta_*) \\ u'(\eta_*) \end{pmatrix} = \begin{pmatrix} w_{\mathbf{k}}^+(\eta_*) & w_{\mathbf{k}}^{+*}(\eta_*) \\ w_{\mathbf{k}}^{+'}(\eta_*) & w_{\mathbf{k}}^{+'*}(\eta_*) \end{pmatrix} \begin{pmatrix} \alpha_{\text{III}\mathbf{k}}^+ \\ \beta_{\text{III}\mathbf{k}}^+ \end{pmatrix}. \quad (3.18)$$

Similarly, the boundary conditions at η_1 are

$$U_{\mathbf{k}}^{\text{II}}(\eta_1) = U_{\mathbf{k}}^{\text{III}}(\eta_1) \quad (3.19)$$

$$U_{\mathbf{k}}^{\text{II}' }(\eta) \Big|_{\eta=\eta_1} = U_{\mathbf{k}}^{\text{III}' }(\eta) \Big|_{\eta=\eta_1} \quad (3.20)$$

Plugging Eqs. (3.12) and (3.14) into the above, we have

$$\begin{pmatrix} w_{\mathbf{k}}^+(\eta_1) & w_{\mathbf{k}}^{+*}(\eta_1) \\ w_{\mathbf{k}}^{+'}(\eta_1) & w_{\mathbf{k}}^{+'*}(\eta_1) \end{pmatrix} \begin{pmatrix} \alpha_{\text{III}\mathbf{k}}^+ \\ \beta_{\text{III}\mathbf{k}}^+ \end{pmatrix} = \begin{pmatrix} v_{\mathbf{k}}(\eta_1) & v_{\mathbf{k}}^*(\eta_1) \\ v_{\mathbf{k}}'(\eta_1) & v_{\mathbf{k}}'^*(\eta_1) \end{pmatrix} \begin{pmatrix} \alpha_{\text{III}\mathbf{k}}^+ \\ \beta_{\text{III}\mathbf{k}}^+ \end{pmatrix}. \quad (3.21)$$

Combining Eqs. (3.18) and (3.21), we find

$$\begin{pmatrix} \alpha_{\text{III}\mathbf{k}}^+ \\ \beta_{\text{III}\mathbf{k}}^+ \end{pmatrix} = \begin{pmatrix} v_{\mathbf{k}}(\eta_1) & v_{\mathbf{k}}^*(\eta_1) \\ v_{\mathbf{k}}'(\eta_1) & v_{\mathbf{k}}'^*(\eta_1) \end{pmatrix}^{-1} \begin{pmatrix} w_{\mathbf{k}}^+(\eta_1) & w_{\mathbf{k}}^{+*}(\eta_1) \\ w_{\mathbf{k}}^{+'}(\eta_1) & w_{\mathbf{k}}^{+'*}(\eta_1) \end{pmatrix} \begin{pmatrix} w_{\mathbf{k}}^+(\eta_*) & w_{\mathbf{k}}^{+*}(\eta_*) \\ w_{\mathbf{k}}^{+'}(\eta_*) & w_{\mathbf{k}}^{+'*}(\eta_*) \end{pmatrix}^{-1} \begin{pmatrix} u_{\mathbf{k}}(\eta_*) \\ u_{\mathbf{k}}'(\eta_*) \end{pmatrix}. \quad (3.22)$$

By using the normalization in Eq. (3.9), the above can be written as

$$\begin{pmatrix} \alpha_{\text{III}\mathbf{k}}^+ \\ \beta_{\text{III}\mathbf{k}}^+ \end{pmatrix} = \begin{pmatrix} -v_{\mathbf{k}}'^*(\eta_1) & v_{\mathbf{k}}^*(\eta_1) \\ v_{\mathbf{k}}'(\eta_1) & -v_{\mathbf{k}}(\eta_1) \end{pmatrix} \begin{pmatrix} w_{\mathbf{k}}^+(\eta_1) & w_{\mathbf{k}}^{+*}(\eta_1) \\ w_{\mathbf{k}}^{+'}(\eta_1) & w_{\mathbf{k}}^{+'*}(\eta_1) \end{pmatrix} \begin{pmatrix} w_{\mathbf{k}}^{+'*}(\eta_*) & -w_{\mathbf{k}}^{+*}(\eta_*) \\ -w_{\mathbf{k}}^{+'}(\eta_*) & w_{\mathbf{k}}^+(\eta_*) \end{pmatrix} \begin{pmatrix} u_{\mathbf{k}}(\eta_*) \\ u_{\mathbf{k}}'(\eta_*) \end{pmatrix}. \quad (3.23)$$

For the $V_{\mathbf{k}}$, the result is obtained by replacing the superscript $+$ by $-$. Using the above results, we calculate the power spectrum of energy density in the next section.

4 Tachyonic instability and the PGW power spectrum

Let us discuss the power spectrum of the energy density of gravitons. From Eq. (2.19), the graviton is given by

$$y_{\mathbf{k}} = \frac{1}{\sqrt{2}} (Y_{\mathbf{k}} + X_{\mathbf{k}}) . \quad (4.1)$$

By using Eq. (3.10), the graviton operator in radiation dominated phase is written as

$$\begin{aligned} y_{\mathbf{k}} &= \frac{1}{\sqrt{2}} \left(\hat{a}_{\mathbf{k}} U_{\mathbf{k}}^{\text{III}} + \hat{a}_{-\mathbf{k}}^\dagger U_{\mathbf{k}}^{\text{III}*} + \hat{b}_{\mathbf{k}} V_{\mathbf{k}}^{\text{III}} + \hat{b}_{-\mathbf{k}}^\dagger V_{\mathbf{k}}^{\text{III}*} \right) \\ &= \frac{1}{\sqrt{2}} \left[\left(\alpha_{\text{III}\mathbf{k}}^+ a_{\mathbf{k}} + \beta_{\text{III}\mathbf{k}}^{+*} a_{\mathbf{k}}^\dagger + \alpha_{\text{III}\mathbf{k}}^- b_{\mathbf{k}} + \beta_{\text{III}\mathbf{k}}^{-*} b_{\mathbf{k}}^\dagger \right) v_{\mathbf{k}} + \left(\beta_{\text{III}\mathbf{k}}^+ a_{\mathbf{k}} + \alpha_{\text{III}\mathbf{k}}^{+*} a_{\mathbf{k}}^\dagger + \beta_{\text{III}\mathbf{k}}^- b_{\mathbf{k}} + \alpha_{\text{III}\mathbf{k}}^{-*} b_{\mathbf{k}}^\dagger \right) v_{\mathbf{k}}^* \right] \\ &\equiv A_{\mathbf{k}} v_{\mathbf{k}} + A_{\mathbf{k}}^\dagger v_{\mathbf{k}}^* , \end{aligned} \quad (4.2)$$

where we defined natural annihilation and creation operators $A_{\mathbf{k}}, A_{\mathbf{k}}^\dagger$ in the radiation-dominated phase. Thus, we can identify the number operator of gravitons as

$${}_{\text{BD}} \langle 0 | A_{\mathbf{k}}^\dagger A_{\mathbf{k}} | 0 \rangle_{\text{BD}} = \frac{1}{2} \left(|\beta_{\text{III}\mathbf{k}}^+|^2 + |\beta_{\text{III}\mathbf{k}}^-|^2 \right) . \quad (4.3)$$

Then, the GW spectral density parameter can be calculated as

$$\Omega_{\text{GW}}(f) = \frac{k^4}{2\pi^2 \rho_c} \left(|\beta_{\text{III}\mathbf{k}}^+|^2 + |\beta_{\text{III}\mathbf{k}}^-|^2 \right) , \quad (4.4)$$

where ρ_c is the critical energy density of the universe and $f = k/2\pi$ is a frequency of GWs. The GW spectral density parameter is plotted in FIG. 3 where we considered the GUT scale inflation $H = 10^{14}$ GeV and $B_0 = 2\sqrt{3} \times 10^{-16}$ G, $\eta_* = -1$ GeV $^{-1}$. We see a peak appears at a particular frequency and no more scale-invariant spectrum as in Eq. (1.1).




FIG 3. GW power spectrum is plotted for $H = 10^{14}$ GeV and $B_0 = 2\sqrt{3} \times 10^{-16}$ G. Parameters are set as $\eta_* = -1 \text{ GeV}^{-1}$, $\eta_1 = 10^{-14} \text{ GeV}^{-1}$, and $\lambda = 2\sqrt{3} \text{ GeV}$. The GUT scale inflation predicts a flat spectrum of the amplitude $\Omega_{\text{GW}} = 10^{-14}$.

The origin of the peak comes from the equation for the $X_{\mathbf{k}}$ in Eq. (3.3). If we rewrite it in the form

$$X_{\mathbf{k}}'' + \left(\left(k - \frac{\lambda}{2} \right)^2 - \frac{\lambda^2}{4} - \frac{2}{(\eta - 2\eta_1)^2} \right) X_{\mathbf{k}} = 0, \quad (4.5)$$

we see that the contribution from the coupling λ causes a tachyonic instability around $k \simeq \lambda/2$. From Eq. (2.15), the strength of the coupling constant can be estimated as

$$\lambda = \frac{\sqrt{2} |\mathbf{B}|}{M_{\text{pl}}} \sin \theta, \quad (4.6)$$

where θ is the angle between the direction of the constant background magnetic field and the wave number vector of GWs. We assume $\sin \theta > 0$ below for simplicity. In the case of $\sin \theta < 0$, we just switch the role of $X_{\mathbf{k}}$ to $Y_{\mathbf{k}}$. From the point of view of observation, we consider the magnitude of current extragalactic magnetic fields of the range $10^{-17} \text{ G} < |\mathbf{B}_0| < 10^{-9} \text{ G}$. After inflation, the energy density of radiation decays as $\rho_\gamma \propto a^{-4}$. By combining the Stefan-Boltzman law $\rho_\gamma = \sigma T^4$, we find that the temperature of radiation decays as the Universe expands such as $T \propto 1/a$. Then the ratio of the present scale factor a_0 to that of reheating time a_1 is found to be $a_0/a_1 = T_{\text{reh}}/T_0$. Here, T_{reh} is the reheating temperature. In our model, the physical magnetic fields decay as $B_{\text{phy}} \propto 1/a$ during inflation and then decay as

$B_{\text{phy}} \propto 1/a^2$ after the reheating. Hence, the ratio of the magnetic field at the reheating time B_{reh} to that at present B_0 becomes $B_{\text{reh}}/B_0 = (a_0/a_1)^2 = (T_{\text{reh}}/T_0)^2 \sim 10^{54} (T_{\text{reh}}/10^{-4}M_{\text{pl}})^2$ where we used $T_0 \sim 2.7\text{K} \sim 10^{-4}\text{eV}$. Accordingly, the magnitude of magnetic fields at the end of the inflation is found to be

$$10^{37} \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right)^2 \text{G} < B_{\text{reh}} < 10^{45} \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right)^2 \text{G}, \quad (4.7)$$

where $B_{\text{reh}} = |\mathbf{B}|/a_1 = |\mathbf{B}|$ because we used $a_1 = 1$ as a convention. By using Eq. (4.6), $1\text{G} \sim 10^{-2}\text{eV}^2$ and $M_{\text{pl}} \sim 10^{18}\text{GeV}$, the above range gives the range of maximum coupling $\lambda_{\text{max}} = \sqrt{2}B_{\text{reh}}/M_{\text{pl}}$ at $\theta = \pi/2$ such as

$$10^{-1} \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right)^2 \text{GeV} < \lambda_{\text{max}} < 10^7 \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right)^2 \text{GeV}. \quad (4.8)$$

The wavenumber at the peak of tachyonic instability $k \simeq \lambda/2$ is translated into the peak frequency observed today and the above lowest value is translated into

$$f_0 = \frac{\lambda}{4\pi} \frac{a_1}{a_0} = 10^{-5} \sin\theta \left(\frac{B_0}{10^{-17}\text{G}} \right) \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right) \text{Hz}, \quad (4.9)$$

where we used $1\text{eV} = 10^{15}\text{Hz}$. Note that the peak frequency depends on the direction of observation. Thus, in terms of maximum frequency at $\theta = \pi/2$, the current range $10^{-17}\text{G} < B_0 < 10^{-9}\text{G}$ corresponds to the frequency range

$$10^{-5} \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right) \text{Hz} < f_{0,\text{max}} < 10^3 \left(\frac{T_{\text{reh}}}{10^{-4}M_{\text{pl}}} \right) \text{Hz}. \quad (4.10)$$

Without the magnetic field, the density parameter of PGWs is known to be $\Omega_{\text{GW}} = 10^{-14}$ for the GUT scale inflation as in Eq. (1.1). In the presence of the magnetic field, Eq. (4.4) tells us that the peak height of the Ω_{GW} is given by

$$\Omega_{\text{GW}}(f_{\text{peak}}) = 10^{-14} \left(\frac{H}{10^{-4}M_{\text{pl}}} \right)^2 \frac{1}{2} \exp[-\lambda\eta_*]. \quad (4.11)$$

We find that the PGWs power spectrum depends on the direction of observation due to the

dependence of θ in λ . The peak height can be determined by

$$-\lambda_{\max}\eta_* = -\frac{\sqrt{2}B}{M_{\text{pl}}}\eta_* = -2\sqrt{3}\frac{B/\sqrt{2}}{\sqrt{3}M_{\text{pl}}H}\frac{\eta_*}{\eta_1} = 2\sqrt{3}\sqrt{\frac{\rho_{B^*}}{\rho_{\text{inf}}}}, \quad (4.12)$$

where we used $a(\eta_1) = 1/(H\eta_1) = 1$ in the second equality. In the third equality, the energy density of the magnetic field at η_* and that of the inflaton field at η_1 are written by $\rho_{B^*} = (a_*/a_1)^2 |\mathbf{B}|^2/2$ and $\rho_{\text{inf}} = 3M_{\text{pl}}^2 H^2$ respectively. Thus, the extra factor $\exp[-\lambda\eta_*]/2$ could be 16 for $\rho_{B^*} \simeq \rho_{\text{inf}}$. We see that the peak height also depends on the direction of observation. The dependence on the direction of observation appears in the presence of the PMFs. Hence, our results would offer new vistas to probe the PMFs through the observations of primordial GWs.

5 Conclusion

In this paper, we studied the effect of PMFs generated during inflation on the PGW power spectrum. Assuming that there exist extragalactic magnetic fields in the range from 10^{-17} to 10^{-9} Gauss with a cosmological coherence length, we showed that the graviton photon conversion process affects the PGW power spectrum. The point is that the graviton photon conversion process induces tachyonic instability. As a consequence, a peak appeared in the spectrum and the peak height depends on the direction of observation. Remarkably, it turned out that the peak frequency lies in the range from 10^{-5} to 10^3 Hertz in the case of GUT scale inflation. Thus, we have a possibility to probe the PMFs by observing the PGW spectrum. Although we considered the graviton photon conversion process, the same argument applies to the graviton dark photon conversion [34]. In this case, the PGW power spectrum could be a probe of vector-type ultralight dark matter.

In [35, 36, 37, 38], the statistical anisotropy in the PGW power spectrum is calculated in the presence of background electric fields. There, no peak appeared in the spectrum. It would be interesting to study what happens when both electric and magnetic fields coexist in the background during inflation.

In this work, we have focused on the coupling parameter $c = -1/2$. In the case of

$c = -1/2$, the physical magnetic fields decay slowly as $fB \propto 1/a$ during inflation compared to the case of constant f where the physical magnetic fields decay as $fB \propto 1/a^2$. On the other hand, in the case of $c = -1$, the physical magnetic fields fB do not decay during inflation. Hence, we expect a different shape of the PGW power spectrum. We leave the analysis of this issue for future work.

Acknowledgments

S. K. was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Number JP22K03621. J. S. was in part supported by JSPS KAKENHI Grant Numbers JP17H02894, JP17K18778, JP20H01902, JP22H01220. K. U. was supported by the JSPS KAKENHI Grant Number 20J22946.

References

- [1] B. P. Abbott et al. “Observation of Gravitational Waves from a Binary Black Hole Merger”. In: *Phys. Rev. Lett.* 116.6 (2016), p. 061102. arXiv: 1602.03837 [gr-qc].
- [2] M. Hazumi et al. “LiteBIRD: A Satellite for the Studies of B-Mode Polarization and Inflation from Cosmic Background Radiation Detection”. In: *J. Low Temp. Phys.* 194.5-6 (2019), pp. 443–452.
- [3] E. Allys et al. “Probing Cosmic Inflation with the LiteBIRD Cosmic Microwave Background Polarization Survey”. In: (Feb. 2022). arXiv: 2202.02773 [astro-ph.IM].
- [4] Seiji Kawamura et al. “The Japanese space gravitational wave antenna: DECIGO”. In: *Class. Quant. Grav.* 28 (2011). Ed. by Sasha Buchman and Ke-Xun Sun, p. 094011.
- [5] Pau Amaro-Seoane et al. “eLISA/NGO: Astrophysics and cosmology in the gravitational-wave millihertz regime”. In: *GW Notes* 6 (2013), pp. 4–110. arXiv: 1201.3621 [astro-ph.CO].
- [6] L. P. Grishchuk and Yu. V. Sidorov. “On the Quantum State of Relic Gravitons”. In: *Class. Quant. Grav.* 6 (1989), pp. L161–L165.

- [7] L. P. Grishchuk and Yu. V. Sidorov. “Squeezed quantum states of relic gravitons and primordial density fluctuations”. In: *Phys. Rev. D* 42 (1990), pp. 3413–3421.
- [8] Andreas Albrecht et al. “Inflation and squeezed quantum states”. In: *Phys. Rev. D* 50 (1994), pp. 4807–4820. arXiv: astro-ph/9303001.
- [9] David Polarski and Alexei A. Starobinsky. “Semiclassicality and decoherence of cosmological perturbations”. In: *Class. Quant. Grav.* 13 (1996), pp. 377–392. arXiv: gr-qc/9504030.
- [10] Sugumi Kanno and Jiro Soda. “Squeezed quantum states of graviton and axion in the universe”. In: (Dec. 2021). arXiv: 2112.14496 [gr-qc].
- [11] Massimo Giovannini. “Hanbury Brown-Twiss interferometry and second-order correlations of inflaton quanta”. In: *Phys. Rev. D* 83 (2011), p. 023515. arXiv: 1011.1673 [astro-ph.CO].
- [12] Sugumi Kanno and Jiro Soda. “Detecting nonclassical primordial gravitational waves with Hanbury-Brown–Twiss interferometry”. In: *Phys. Rev. D* 99.8 (2019), p. 084010. arXiv: 1810.07604 [hep-th].
- [13] Sugumi Kanno. “Nonclassical primordial gravitational waves from the initial entangled state”. In: *Phys. Rev. D* 100.12 (2019), p. 123536. arXiv: 1905.06800 [hep-th].
- [14] Maulik Parikh, Frank Wilczek, and George Zahariade. “The Noise of Gravitons”. In: *Int. J. Mod. Phys. D* 29.14 (2020), p. 2042001. arXiv: 2005.07211 [hep-th].
- [15] Sugumi Kanno, Jiro Soda, and Junsei Tokuda. “Noise and decoherence induced by gravitons”. In: *Phys. Rev. D* 103.4 (2021), p. 044017. arXiv: 2007.09838 [hep-th].
- [16] Maulik Parikh, Frank Wilczek, and George Zahariade. “Quantum Mechanics of Gravitational Waves”. In: *Phys. Rev. Lett.* 127.8 (2021), p. 081602. arXiv: 2010.08205 [hep-th].
- [17] Maulik Parikh, Frank Wilczek, and George Zahariade. “Signatures of the quantization of gravity at gravitational wave detectors”. In: *Phys. Rev. D* 104.4 (2021), p. 046021. arXiv: 2010.08208 [hep-th].

- [18] Sugumi Kanno, Jiro Soda, and Junsei Tokuda. “Indirect detection of gravitons through quantum entanglement”. In: *Phys. Rev. D* 104.8 (2021), p. 083516. arXiv: 2103.17053 [gr-qc].
- [19] Shin’ichiro Ando and Alexander Kusenko. “Evidence for Gamma-Ray Halos Around Active Galactic Nuclei and the First Measurement of Intergalactic Magnetic Fields”. In: *Astrophys. J. Lett.* 722 (2010), p. L39. arXiv: 1005.1924 [astro-ph.HE].
- [20] A. Neronov and I. Vovk. “Evidence for strong extragalactic magnetic fields from Fermi observations of TeV blazars”. In: *Science* 328 (2010), pp. 73–75. arXiv: 1006.3504 [astro-ph.HE].
- [21] Warren Essey, Shin’ichiro Ando, and Alexander Kusenko. “Determination of intergalactic magnetic fields from gamma ray data”. In: *Astropart. Phys.* 35 (2011), pp. 135–139. arXiv: 1012.5313 [astro-ph.HE].
- [22] F. Tavecchio et al. “Extreme TeV blazars and the intergalactic magnetic field”. In: *Mon. Not. Roy. Astron. Soc.* 414 (2011), p. 3566. arXiv: 1009.1048 [astro-ph.HE].
- [23] Justin D. Finke et al. “Constraints on the Intergalactic Magnetic Field with Gamma-Ray Observations of Blazars”. In: *Astrophys. J.* 814.1 (2015), p. 20. arXiv: 1510.02485 [astro-ph.HE].
- [24] Tanmay Vachaspati. “Progress on cosmological magnetic fields”. In: *Reports on Progress in Physics* 84.7, 074901 (July 2021), p. 074901. arXiv: 2010.10525 [astro-ph.CO].
- [25] Dario Grasso and Hector R. Rubinstein. “Magnetic fields in the early universe”. In: *Phys. Rept.* 348 (2001), pp. 163–266. arXiv: astro-ph/0009061.
- [26] Daniela Paoletti and Fabio Finelli. “Constraints on primordial magnetic fields from magnetically-induced perturbations: current status and future perspectives with LiteBIRD and future ground based experiments”. In: *JCAP* 11 (2019), p. 028. arXiv: 1910.07456 [astro-ph.CO].
- [27] Michael S. Turner and Lawrence M. Widrow. “Inflation Produced, Large Scale Magnetic Fields”. In: *Phys. Rev. D* 37 (1988), p. 2743.

- [28] Bharat Ratra. “Cosmological ‘seed’ magnetic field from inflation”. In: *Astrophys. J. Lett.* 391 (1992), pp. L1–L4.
- [29] Sugumi Kanno, Jiro Soda, and Kazushige Ueda. “Conversion of squeezed gravitons into photons during inflation”. In: (July 2022). arXiv: 2207.05734 [hep-th].
- [30] Sugumi Kanno, Jiro Soda, and Kazushige Ueda. “Impact of quantum entanglement induced by magnetic fields on primordial gravitational waves”. In: (Nov. 2022). arXiv: 2211.05576 [hep-th].
- [31] M.E. Gertsenshtein. “Wave Resonance of Light and Gravitational Waves”. In: *J.Exp.Theor.Phys.* 14 (1962), p. 84.
- [32] Georg Raffelt and Leo Stodolsky. “Mixing of the Photon with Low Mass Particles”. In: *Phys. Rev. D* 37 (1988), p. 1237.
- [33] Sugumi Kanno, Jiro Soda, and Masa-aki Watanabe. “Cosmological Magnetic Fields from Inflation and Backreaction”. In: *JCAP* 12 (2009), p. 009. arXiv: 0908.3509 [astro-ph.CO].
- [34] Emi Masaki and Jiro Soda. “Conversion of Gravitons into Dark Photons in Cosmological Dark Magnetic Fields”. In: *Phys. Rev. D* 98.2 (2018), p. 023540. arXiv: 1804.00458 [astro-ph.CO].
- [35] Masa-aki Watanabe, Sugumi Kanno, and Jiro Soda. “Inflationary Universe with Anisotropic Hair”. In: *Phys. Rev. Lett.* 102 (2009), p. 191302. arXiv: 0902.2833 [hep-th].
- [36] Masa-aki Watanabe, Sugumi Kanno, and Jiro Soda. “The Nature of Primordial Fluctuations from Anisotropic Inflation”. In: *Prog. Theor. Phys.* 123 (2010), pp. 1041–1068. arXiv: 1003.0056 [astro-ph.CO].
- [37] Timothy R. Dulaney and Moira I. Gresham. “Primordial Power Spectra from Anisotropic Inflation”. In: *Phys. Rev. D* 81 (2010), p. 103532. arXiv: 1001.2301 [astro-ph.CO].
- [38] A. E. Gumrukcuoglu, Burak Himmetoglu, and Marco Peloso. “Scalar-Scalar, Scalar-Tensor, and Tensor-Tensor Correlators from Anisotropic Inflation”. In: *Phys. Rev. D* 81 (2010), p. 063528. arXiv: 1001.4088 [astro-ph.CO].