

PHILIP SALTENBERGER
Institute for Numerical Analysis, TU Braunschweig
Braunschweig, Germany
 (E-Mail: philip.saltenberger@tu-bs.de)

STRUCTURE-PRESERVING EIGENVALUE MODIFICATION OF SYMPLECTIC MATRICES AND MATRIX PENCILS

Abstract

A famous theorem by R. Brauer shows how to modify a single eigenvalue of a matrix A by a rank-one update without changing the remaining eigenvalues. A generalization of this theorem (due to R. Rado) is used to change a pair of eigenvalues $\lambda, 1/\lambda$ of a symplectic matrix S in a structure-preserving way to desired target values $\mu, 1/\mu$. Universal bounds on the relative distance between S and the newly constructed symplectic matrix \hat{S} with modified spectrum are given. The eigenvalues Segre characteristics of \hat{S} are related to those of S and a statement on the eigenvalue condition numbers of \hat{S} is derived. The main results are extended to matrix pencils.

1. INTRODUCTION

In numerical linear algebra and matrix analysis one occasionally encounters the necessity of modifying special eigenvalues of a matrix without altering its remaining eigenvalues. Techniques for changing certain eigenvalues of a matrix have, for instance, been applied to solve nonnegative inverse eigenvalue problems [13, 16] or, in form of deflation methods, to remove dominant eigenvalues in eigenvalue computations [14, Sec. 4.2]. Furthermore, the task of modifying eigenvalues of matrices is of interest in stability and feedback of linear systems [4, § 25], [3, Sec. 2.3] or for passivity and eigenvalue assignment in control design [1]. One basic result on how a single eigenvalue of a matrix may be changed without modifying any other eigenvalues is due to R. Brauer and can be found in [3, Sec. 1], [16].

Theorem 1 (Brauer). *Let $A \in \mathbf{M}_n(\mathbb{C})$ have eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and let $x_1 \in \mathbb{C}^n$ be an eigenvector for λ_1 . Then, for any $c \in \mathbb{C}^n$, the matrix $\hat{A} = A + x_1 c^T \in \mathbf{M}_n(\mathbb{C})$ has the eigenvalues $\lambda_1 + c^T x_1, \lambda_2, \dots, \lambda_n$.*

This work is concerned with the purposive change of certain eigenvalues of matrices with symplectic structure. A complex $2n \times 2n$ matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$

is called *symplectic*, if¹

$$S^T J_{2n} S = J_{2n} =: J, \quad \text{where } J_{2n} = \begin{bmatrix} 0_{n \times n} & I_n \\ -I_n & 0_{n \times n} \end{bmatrix}. \quad (1)$$

Defining $S^* := J^T S^T J$, we see that (1) is equivalent to $S^* S = I_{2n}$. Therefore, a symplectic matrix S is always nonsingular and $S^* = S^{-1}$. In consequence, as S^* is similar² to S , the eigenvalues of a symplectic matrices arise in pairs $\lambda_j, \lambda_j^{-1}$, $j = 1, \dots, n$, where λ_j and λ_j^{-1} have the same Segre characteristic. Recall that for an eigenvalue λ of S , its Segre characteristic is the sequence of sizes of the Jordan blocks of S with eigenvalue λ in non-increasing order [15]. We denote the Segre characteristic of an eigenvalue by $((\cdot, \dots, \cdot))$. It is now immediate that Theorem 1 can in general not be used for a structure-preserving, symplectic change of eigenvalues. In fact, for a structure-preserving eigenvalue modification, the change of λ_j and λ_j^{-1} must take place simultaneously.

Without any structure-preservation in mind, changing two (or more) eigenvalues simultaneously is possible with the following generalization of Theorem 1 attributed to R. Rado. It can be found in [13], see also [3, Sec. 3].

Theorem 2 (Rado). *Let $A \in \mathbf{M}_n(\mathbb{C})$ have eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ and let $x_1, \dots, x_k \in \mathbb{C}^n$ be linearly independent eigenvectors for $\lambda_1, \dots, \lambda_k$. Set $X = [x_1 \ \dots \ x_k] \in \mathbf{M}_{n \times k}(\mathbb{C})$. Then, for any matrix $C \in \mathbf{M}_{n \times k}(\mathbb{C})$, the matrix $\hat{A} = A + XC^T$ has the eigenvalues $\mu_1, \dots, \mu_k, \lambda_{k+1}, \dots, \lambda_n$, where μ_j , $j = 1, \dots, k$, are the eigenvalues of $\Omega = \text{diag}(\lambda_1, \dots, \lambda_n) + C^T X$.*

In this work, we investigate how Theorem 2 can be utilized to change a pair of eigenvalues $\lambda_j, \lambda_j^{-1}$ of a symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ (or a symplectic matrix pencil) to desired target values μ, μ^{-1} in a structure-preserving way without modifying any other eigenvalues of S . Considering Theorem 2, the starting point of our discussion is thus the following question:

Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with eigenvalues $\lambda_1, \lambda_1^{-1}, \dots, \lambda_n, \lambda_n^{-1}$, linearly independent eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ for λ_1 and λ_1^{-1} , respectively, and $X = [x_1 \ x_2]$. How has $C \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ to be chosen, such that $\hat{S} := S + XC^T$ is symplectic with eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ for some given value $\mu \in \mathbb{C} \setminus \{0\}$?

The above-mentioned problem will be discussed in Section 2. In Section 3 we investigate whether we can find an upper bound $b > 0$ that only depends on λ_1, μ, x_1 and x_2 that assures the existence of a symplectic matrix $\hat{S} \in \mathbf{M}_{2n}(\mathbb{C})$ with eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ and relative distance $\|\hat{S} - S\|/\|S\| \leq b$. We derive distinguished matrices \hat{S}_1, \hat{S}_2 for which such a bound b can be neatly expressed and related to the relative change

¹Here and in the following, T denotes the transpose of a (maybe complex) matrix or vector, not its conjugate transpose.

²By definition, S^* is similar to S^T and by the Taussky-Zassenhaus Theorem [17], S^T is similar to S .

in the eigenvalue, i.e. $|\lambda_1 - \mu|/|\lambda_1|$. We discuss commutativity relations between S and \hat{S} in Section 4.1 and characterize the Segre characteristics of the eigenvalues of \hat{S} in Section 4.2. The results of Section 4.1 will come in handy here to find a condition on the simultaneous diagonalizability of S and \hat{S} . In Section 6 we partially extend our results from Section 2 to symplectic matrix pencils.

1.1. NOTATION

The set of all $m \times n$ matrices over \mathbb{K} (where we use either $\mathbb{K} = \mathbb{C}$ or $\mathbb{K} = \mathbb{R}$) is denoted by $\mathbf{M}_{m \times n}(\mathbb{K})$. Whenever $n = m$ we write $\mathbf{M}_n(\mathbb{K})$ instead of $\mathbf{M}_{n \times n}(\mathbb{C})$. For $J_{2n} \in \mathbf{M}_{2n}(\mathbb{R})$, see (1), we simply write J and add the index whenever it is necessary to specify the size of J . The range of a matrix $A \in \mathbf{M}_{m \times n}(\mathbb{C})$ is the vector space spanned by its columns and is denoted $\text{range}(A)$. For $A \in \mathbf{M}_{m \times n}(\mathbb{C})$, we denote the Moore-Penrose pseudoinverse of A by A^+ . In case $m > n$ and $\text{rank}(A) = n$, we have $A^+ = (A^H A)^{-1} A^H$ so that $A^+ A = I_n$, while for $n > m$ and $\text{rank}(A) = m$, $A^+ = A^H (A A^H)^{-1}$ yields $A A^+ = I_m$. The superscript H always denotes the conjugate transpose of a matrix or vector while T is used for the pure transposition. Whenever $\lambda \in \mathbb{C}$ is some complex number, we denote by $\Re(\lambda)$ and $\Im(\lambda)$ its real and imaginary part, respectively. Complex conjugation of a number $x = a + ib \in \mathbb{C}$ is denoted by a bar, i.e. $\bar{x} = a - ib$.

2. SYMPLECTIC EIGENVALUE MODIFICATION

Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be a symplectic matrix (see (1)) with eigenvalues $\lambda_1, \lambda_1^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Furthermore, assume $x_1, x_2 \in \mathbb{C}^{2n}$ are linearly independent³ eigenvectors of S for λ_1 and λ_1^{-1} , respectively, and define $X = [x_1 \ x_2] \in \mathbf{M}_{2n \times 2}(\mathbb{C})$. In this section, our goal is to determine all possible matrices $C \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ such that $\hat{S} := S + X C^T$ is again symplectic and has the eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$. To this end, we will make use of Rado's theorem and derive a structure-preserving version of Theorem 2 (see Theorem 3).

As it will become clear later, it seems appropriate to consider the situations $x_1^T J x_2 \neq 0$ and $x_1^T J x_2 = 0$ separately. First, we assume that for the eigenvalues $\lambda_1, \lambda_1^{-1}$ there exist eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ such that $x_1^T J x_2 \neq 0$ (this immediately implies x_1 and x_2 to be linearly independent). In this case, we can assume w. l. o. g. $x_1^T J x_2 = 1$, which can be achieved by a scaling of x_1 and/or x_2 . That is, we have $X^T J X = J_2$. For the matrix $\hat{S} := S + X C^T$ to be symplectic, it has to hold that

$$\hat{S}^T J \hat{S} = (S + X C^T)^T J (S + X C^T) = J. \quad (2)$$

³If $\lambda_1 \neq \lambda_1^{-1}$, then x_1 and x_2 are necessarily linear independent. Therefore, the linear independence is only a restrictive requirement if $\lambda_1 = \lambda_1^{-1}$, i.e. $\lambda_1 = \pm 1$.

Using $S^T JS = J$, (2) is equivalent to the matrix equation

$$CX^T JS + S^T JXC^T + CX^T JXC^T = 0 \quad (3)$$

for the unknown matrix $C \in \mathbf{M}_{2n \times 2}(\mathbb{C})$. Notice that (3) can be rewritten as

$$C(X^T JS + J_2 C^T) = -S^T JXC^T \quad (4)$$

using $X^T JX = J_2$. Since $S^T JX \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ is a matrix of full rank, (4) immediately implies $\text{range}(C) \subseteq \text{range}(S^T JX)$ for any solution C . Thus, for every C satisfying (3), there is a matrix $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ such that $C = S^T JXR^T$. Plugging this ansatz into (3), we obtain a $2n \times 2n$ equation for R , namely

$$S^T JXR^T X^T JS + S^T JXR^T X^T J^T S + S^T JXR^T J_2 R X^T J^T S = 0.$$

Replacing $X^T JS$ by $-X^T J^T S$ this can be rewritten as

$$S^T JX(R - R^T + R^T J_2 R)X^T J^T S = 0. \quad (5)$$

Finally, we may multiply (5) with the pseudo inverses $(S^T JX)^+$ from the left and with $(X^T J^T S)^+$ from the right to obtain

$$R - R^T + R^T J_2 R = 0, \quad (6)$$

which is a matrix equation for R of size 2×2 that is equivalent to (5). As $R - R^T$ and $R^T J_2 R$ are both skew-symmetric, their diagonals are identically zero. Comparing the entries of $R - R^T$ and $R^T J_2 R$ in the (1,2) position, we obtain the condition

$$r_{12} - r_{21} + r_{11}r_{22} - r_{12}r_{21} = 0 \quad (7)$$

for (6) to hold (comparing the elements in the (2,1) position certainly gives the same condition with a minus sign). In summary, *a matrix of the form $\hat{S} = S + XC^T$ is symplectic if and only if $C = S^T JXR^T$ for some matrix $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ whose entries satisfy (7).*

Next, to achieve the desired eigenvalue modification, according to Theorem 2 we need to assure that the eigenvalues of

$$\Omega := \Lambda + C^T X = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} + C^T X = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} + RX^T J^T SX \quad (8)$$

become equal to μ and μ^{-1} . To this end, recall that $SX = X \text{diag}(\lambda_1, \lambda_1^{-1})$ (by construction of X). Thus $\text{diag}(\lambda_1, \lambda_1^{-1}) + RX^T J^T SX = \text{diag}(\lambda_1, \lambda_1^{-1}) - RX^T JX \text{diag}(\lambda_1, \lambda_1^{-1})$ which, since $X^T JX = J_2$, yields

$$\Omega = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} - RJ_2 \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} = \begin{bmatrix} \lambda_1 + \lambda_1 r_{12} & -\lambda_1^{-1} r_{11} \\ \lambda_1 r_{22} & \lambda_1^{-1} - \lambda_1^{-1} r_{21} \end{bmatrix}. \quad (9)$$

The characteristic polynomial of Ω is

$$p(z) = z^2 - (\lambda_1(1 + r_{12}) + \lambda_1^{-1}(1 - r_{21}))z + (1 + r_{12})(1 - r_{21}) + r_{11}r_{22},$$

which should, by Theorem 2, be equal to $q(z) = (z - \mu)(z - \mu^{-1}) = z^2 - (\mu + \mu^{-1})z + 1$ to achieve that \hat{S} will have the eigenvalues μ and μ^{-1} . This gives two more conditions: one the one hand $\lambda_1(1 + r_{12}) + \lambda_1^{-1}(1 - r_{21}) = \mu + \mu^{-1}$, i.e.

$$\lambda_1 r_{12} - \lambda_1^{-1} r_{21} = (\mu + \mu^{-1}) - (\lambda_1 + \lambda_1^{-1}). \quad (10)$$

One the other hand, $(1 + r_{12})(1 - r_{21}) + r_{11}r_{22} = 1$. The latter condition, however, is equal to condition (7) obtained for the symplectic structure above. Thus, *additionally to (7), which is required for $S + XC^T$ to be symplectic, the equation (10) has to hold to achieve that μ, μ^{-1} become eigenvalues of $S + XC^T$* . In conclusion, we obtain the following version of Theorem 2 that answers the question stated in Section 1 on the eigenvalue modification for symplectic matrices.

Theorem 3. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with eigenvalues $\lambda_1, \lambda_1^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Let $x_1, x_2 \in \mathbb{C}^{2n}$ be eigenvectors for λ_1 and λ_1^{-1} , respectively, normalized such that $X^T J X = J_2$ for $X = [x_1 \ x_2] \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ and set $d := (\mu + \mu^{-1}) - (\lambda_1 + \lambda_1^{-1})$. Then the matrix*

$$\hat{S} := S + XC^T \in \mathbf{M}_{2n}(\mathbb{C}) \quad (11)$$

is symplectic and has the eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ if and only if $C^T = RX^T J^T S$ for some matrix $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ whose entries satisfy the conditions

$$d = \lambda_1 r_{12} - \lambda_1^{-1} r_{21}, \text{ and} \quad (12)$$

$$0 = r_{12} - r_{21} + r_{11}r_{22} - r_{12}r_{21}. \quad (13)$$

Notice that the matrix $\hat{S} = S + XRX^T J^T S$ in (11) can also be expressed as $\hat{S} = (I_{2n} + XRX^T J^T)S$ or as

$$\hat{S} = S + XRA^{-1}X^T J^T = S + XR \begin{bmatrix} \lambda_1^{-1} & 0 \\ 0 & \lambda_1 \end{bmatrix} X^T J^T \quad (14)$$

according to the relation $\Lambda^{-1}X^T J^T = X^T J^T S$ (where $\Lambda = \text{diag}(\lambda_1, \lambda_1^{-1})$). Furthermore, we see from (12) and (13) that there exist infinitely many possible choices for R that realize the desired eigenvalue modification.

Next, we discuss the case that the eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ of the symplectic matrix S for λ_1 and λ_1^{-1} , respectively, satisfy $x_1^T J x_2 = 0$ and how this condition effects the result from Theorem 3. To this end, first notice that a symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ need in fact not have eigenvectors x_1, x_2 for

λ_1 and λ_1^{-1} that satisfy $x_1^T J x_2 \neq 0$. A situation of this kind arises for the symplectic matrix

$$S = \begin{bmatrix} \lambda_1 & 1 & 0 & 0 \\ 0 & \lambda_1 & 0 & 0 \\ 0 & 0 & \lambda_1^{-1} & 0 \\ 0 & 0 & -\lambda_1^{-2} & \lambda_1^{-1} \end{bmatrix}$$

and its eigenvalue λ_1 . The only eigenvectors for λ_1 and λ_1^{-1} are e_1 and e_4 , respectively, and we have $e_1^T J_4 e_4 = 0$. Thus, Theorem 3 cannot be applied. A simple sufficient (but not necessary) criterion to assure that eigenvectors with $x_1^T J x_2 \neq 0$ must exist, is that S is a diagonalizable matrix, cf. [10, Lem. 3, Cor. 3.1] and Corollary 1 below.

Whenever $x_1, x_2 \in \mathbb{C}^{2n}$ are eigenvectors of S for λ_1 and λ_1^{-1} with $x_1^T J x_2 = 0$, then $X^T J X = 0$ follows for $X = [x_1 \ x_2]$. In this case, it follows from (3) that (4) takes the form

$$C X^T J S = -S^T J X C^T.$$

Again we obtain $\text{range}(C) \subseteq \text{range}(S^T J X)$, so there has to exist some matrix $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ such that $C = S^T J X R^T$. However, despite the concrete form of R , analogously to (8) we obtain

$$\Omega = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} + C^T X = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix} - R X^T J S X = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{bmatrix}$$

since $SX = X \text{diag}(\lambda, \lambda^{-1})$ and $X^T J X = 0$. Thus, *even if R is chosen according to (13) such that $\tilde{S} = S + X R X^T J^T S$ is symplectic, no change in the eigenvalues can be achieved.* In consequence, a change of an eigenvalue pair $\lambda_1, \lambda_1^{-1}$ of a symplectic matrix by Rado's theorem in a structure-preserving way is only possible if there exist eigenvectors x_1 and x_2 for λ_1 and λ_1^{-1} , respectively, such that $x_1^T J x_2 \neq 0$. In the next section, we derive a universal criterion on the existence of such eigenvectors.

2.1. APPLYING THEOREM 3: A CRITERION

We will now characterize those symplectic matrices $S \in \mathbf{M}_{2n}(\mathbb{C})$, for which an eigenvalue adjustment according to Theorem 3 is possible. The condition derived below involves the Segre characteristic of the eigenvalue $\lambda_1 \in \sigma(S)$ to be modified.

First, let $\lambda_1 \in \sigma(S)$ and $Sx_1 = \lambda_1 x_1$ and $Sx_2 = \lambda_1^{-1} x_2$. Now suppose at least one of both vectors, e.g. x_1 , belongs to a nontrivial⁴ Jordan chain, that is, there is some $z \in \mathbb{C}^{2n}$ such that $(S - \lambda_1 I_{2n})z = x_1$ (and possibly more

⁴By nontrivial, we mean a Jordan chain of length ≥ 2 while a trivial Jordan chain refers to a chain of length one.

generalized eigenvectors beside z). Then we have

$$\begin{aligned}
x_1^T J x_2 &= ((S - \lambda_1 I_{2n})z)^T J x_2 = z^T S^T J x_2 - \lambda_1 z^T J x_2 \\
&= z^T J J^T S^T J x_2 - \lambda_1 z^T J x_2 \\
&= z^T J S^{-1} x_2 - \lambda_1 z^T J x_2 \\
&= \lambda_1 z^T J x_2 - \lambda_1 z^T J x_2 = 0
\end{aligned} \tag{15}$$

as $J^T S^T J = S^* = S^{-1}$ and $S^{-1} x_2 = \lambda_1 x_2$. In consequence, $x_1^T J x_2 = 0$ whenever x_1, x_2 are eigenvectors of S for λ_1 and λ_1^{-1} , respectively, and at least one of them belongs to a nontrivial Jordan chain. *In other words, we may have $x_1^T J x_2 \neq 0$ only in case both x_1 and x_2 belong to trivial Jordan chains.* Next, we show that in case x_1 belongs to a trivial Jordan chain there must exist x_2 (also from a trivial Jordan chain) such that $x_1^T J x_2 \neq 0$.

To this end, assume that $\lambda_1 \in \mathbb{C}$ is an eigenvalue of the symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ with $p \geq 1$ ones in its Segre characteristic (that is, there are p Jordan blocks of size 1×1 , i.e. p trivial Jordan chains, and possibly other Jordan blocks of size ≥ 2). Then there exists a matrix $F \in \mathbf{M}_{2n}(\mathbb{C})$ transforming S to the following Jordan form

$$F^{-1} S F =: G = \left[\begin{array}{ccc|c} \lambda_1 & & & 0 \\ & \ddots & & \\ & & \lambda_1 & \\ \hline & & 0 & \hat{G} \end{array} \right] \tag{16}$$

where the upper-left block is $\lambda_1 I_p$ and \hat{G} contains all other Jordan blocks (note that there might also be other Jordan blocks for λ_1 of size ≥ 2 contained in \hat{G}). Now define

$$x_1 := F e_1 \quad \text{and} \quad \tilde{f}^H := e_1^T F^{-1}. \tag{17}$$

Then x_1 is a right eigenvector of S for λ_1 ($S x_1 = \lambda_1 x_1$) and \tilde{f} is a left eigenvector of S for λ_1 ($\tilde{f}^H S = \lambda_1 \tilde{f}^H$). Certainly, $\tilde{f}^H x_1 = 1$. Now we define $x_2^T := \tilde{f}^H J$. Then we have

$$\begin{aligned}
\tilde{f}^H S = \lambda_1 \tilde{f}^H &\Leftrightarrow x_2^T J^T S = \lambda_1 x_2^T J^T \\
&\Leftrightarrow x_2^T J^T S J = \lambda_1 x_2^T \\
&\Leftrightarrow x_2^T S^{-T} = \lambda_1 x_2^T \Leftrightarrow S x_2 = \lambda_1^{-1} x_2.
\end{aligned} \tag{18}$$

It follows that x_2 is an eigenvector of S for λ_1^{-1} . Now we obtain

$$x_1^T J x_2 = x_2^T J^T x_1 = \tilde{f}^H x_1 = 1 \neq 0.$$

In conclusion, for any eigenvector x_1 of S for λ_1 belonging to a trivial Jordan chain, there always exists an eigenvector x_2 of S for λ_1^{-1} such that $x_1^T J x_2 \neq 0$. Recall from our observation (15) above, that x_2 must also be a vector from a trivial Jordan chain. We conclude our findings in the following theorem.

Theorem 4. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Then Theorem 3 is applicable to S for λ_1 and μ , i.e. there exist eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ for λ_1 and λ_1^{-1} , respectively, with $x_1^T J x_2 \neq 0$, if and only if the Segre characteristic of S for λ_1 contains a one, that is, it has the form $((\star, \star, \dots, \star, 1))$. In particular, eigenvectors x_1 for λ_1 and x_2 for λ_1^{-1} with $x_1^T J x_2 \neq 0$ always belong to trivial Jordan chains of S .*

Do not overlook that Theorem 4 applies also for $\lambda_1 = \lambda_1^{-1}$, i.e. $\lambda_1 = \pm 1$. In this case $\lambda_1 \in \sigma(S)$ necessarily has an even multiplicity and an even number of Jordan blocks of the same size, so a Segre characteristic of the form $((\star, \star, \dots, \star, 1))$ implies that there appears at least another one, i.e. $((\star, \star, \dots, \star, 1, 1))$. Then the reasoning in (16), (17) and (18) applies in the same way. If S is diagonalizable, the Segre characteristic of S for any eigenvalue $\lambda_j \in \sigma(S)$ consists only of ones. So we immediately obtain the following corollary.

Corollary 1. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Then Theorem 3 is applicable to S for λ_1 and μ if S is diagonalizable.*

3. BOUNDING THE RELATIVE CHANGE

Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic. For the matrix $\hat{S} = S + X R X^T J^T S$ in (11) we immediately obtain a bound on its (absolute or relative) change in norm with respect to S . That is,

$$\|S - \hat{S}\| \leq \|R\| \|X\| \|X^T\| \|S\| \quad \text{and} \quad \frac{\|S - \hat{S}\|}{\|S\|} \leq \|R\| \|X\| \|X^T\| \quad (19)$$

hold for any submultiplicative and unitarily invariant matrix norm $\|\cdot\|$. In this section, we intend to derive explicit bounds of the relative distance between S and \hat{S} for certain choices of $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$. To this end, we assume $\|\cdot\| = \|\cdot\|_F$ so that $\|X\|_F = \|X^T\|_F$ holds and the upper bound in (19) reduces to $\|R\|_F \|X\|_F^2$.

To bound the relative change $\|\hat{S} - S\|_F / \|S\|_F$ with respect to S consider again (12) and (13). The solution set to (12) is an affine subspace of \mathbb{C}^2 and all solutions may be parameterized as

$$r_{12} = \eta \lambda_1^{-1}, \quad r_{21} = -\lambda_1 d + \eta \lambda_1, \quad \eta \in \mathbb{C}. \quad (20)$$

Plugging these expressions for r_{12} and r_{21} into (13) yields a polynomial in η , i.e.

$$p(\eta) = -\eta^2 + \eta (\lambda_1^{-1} + d - \lambda_1) + d \lambda_1 + r_{11} r_{22}. \quad (21)$$

Thus, depending on r_{11} and r_{22} (which can both be arbitrary), in (21) there are always two solutions $\eta_1, \eta_2 \in \mathbb{C}$ of $p(\eta) = 0$ and, in consequence, two

matrices

$$R(\eta_j, r_{11}, r_{22}) := \begin{bmatrix} r_{11} & \eta_j \lambda_1^{-1} \\ \lambda_1(\eta_j - d) & r_{22} \end{bmatrix}, \quad j = 1, 2, \quad (22)$$

so that their entries satisfy (12) and (13).

To find some $R \in \mathbf{M}_2(\mathbb{C})$ that yields a small norm $\|R\|_F$ and thus a small bound in (19), it seems natural to consider the case $r_{11}r_{22} = 0$, in particular $r_{11} = r_{22} = 0^5$. The two possible roots of $p(\eta)$ for $r_{11}r_{22} = 0$ are $\eta_1 = \mu - \lambda_1$ and $\eta_2 = \mu^{-1} - \lambda_1$. The matrices $R_1 := R(\eta_1, 0, 0)$ and $R_2 := R(\eta_2, 0, 0)$ that arise according to (22) are thus given by

$$R_1 = \begin{bmatrix} 0 & \lambda_1^{-1}(\mu - \lambda_1) \\ \mu^{-1}(\mu - \lambda_1) & 0 \end{bmatrix}, \quad R_2 = \begin{bmatrix} 0 & \mu^{-1}(\lambda_1^{-1} - \mu) \\ \lambda_1(\lambda_1^{-1} - \mu) & 0 \end{bmatrix}. \quad (23)$$

Using R_1 and R_2 in (23), explicit bounds can be found on $\|\hat{S} - S\|_F / \|S\|_F$. According to (19) and (23) such a bound $b \geq 0$ only depends on λ_1 , μ and the eigenvectors of S for λ_1 and λ_1^{-1} and guarantees the existence of a symplectic matrix $\hat{S} \in \mathbf{M}_{2n}(\mathbb{C})$ that solves the problem from Section 1 with $\|\hat{S} - S\|_F / \|S\|_F \leq b$. To formulate these bounds, we impose a condition on X to estimate $\|X\|_F$ without computing the norm. In particular, we assume the eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ of S for λ_1 and λ_1^{-1} , respectively, to be normalized, i.e. $\|x_1\|_2 = \|x_2\|_2 = 1$ and $X \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ to be of the form

$$X = \frac{1}{\sqrt{x_1^T J x_2}} \begin{bmatrix} x_1 & x_2 \end{bmatrix}. \quad (24)$$

Then $X^T J X = J_2$ holds and it follows that

$$\|X\|_F^2 = \left| \frac{1}{\sqrt{x_1^T J x_2}} \right|^2 \left\| \begin{bmatrix} x_1 & x_2 \end{bmatrix} \right\|_F^2 = \frac{1}{|x_1^T J x_2|} \left\| \begin{bmatrix} x_1 & x_2 \end{bmatrix} \right\|_F^2 = \frac{2}{|x_1^T J x_2|} \quad (25)$$

The value $1/|x_1^T J x_2|$ has a nice interpretation whenever λ_1 is a *simple* eigenvalue of S and $\|x_1\|_2 = \|x_2\|_2 = 1$ holds. To see this, recall that, whenever $A \in \mathbf{M}_n(\mathbb{C})$ has a simple eigenvalue $\lambda \in \mathbb{C}$ (i.e. its algebraic multiplicity equals one), then

$$\kappa(A, \lambda) := \frac{\|u\|_2 \|v\|_2}{|v^H u|}$$

is called its condition number, where $u \in \mathbb{C}^{2n}$ and $v \in \mathbb{C}^{2n}$ are right and left eigenvectors of A for λ (i.e. $Au = \lambda u$ and $v^H A = \lambda v^H$). It is a measure on how sensitive λ reacts to small changes in the matrix A , see [14, Sec. 3.3]. As we have seen in (18) above, $(x_2^T J^T)S = \lambda_1(x_2^T J^T)$ whenever x_2 satisfies

⁵Certainly, choosing $r_{11} = 0$ and $r_{22} \neq 0$ gives the same roots of $p(\eta) = 0$ in (21), and thus the same values for r_{12} and r_{21} , but a larger Frobenius norm of R than choosing $r_{11} = r_{22} = 0$.

$Sx_2 = \lambda_1^{-1}x_2$. Thus, for simple λ_1 (which implies that λ_1^{-1} is simple as well) we can choose $u = x_1$ and $v^H = x_2^T J^T$ so that

$$\kappa(S, \lambda_1) = \frac{\|u\|_2 \|v\|_2}{|v^H u|} = \frac{\|x_1\|_2 \|J\bar{x}_2\|_2}{|x_2^T J^T x_1|} = \frac{1}{|x_1^T J x_2|}$$

since $\|x_1\|_2 = 1$ and $\|J\bar{x}_2\|_2 = \|\bar{x}_2\|_2 = \|x_2\|_2 = 1$. We can now formulate the following theorem which follows directly from the bound in (19), the observation in (25) and the Frobenius norms of the matrices R_1, R_2 in (23).

Theorem 5. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1, \lambda_1^{-1} \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Let $x_1, x_2 \in \mathbb{C}^{2n}$ be normalized eigenvectors for λ_1 and λ_1^{-1} , respectively, and $X \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ as in (24). Define*

$$\Phi := \frac{2}{|x_1^T J x_2|} \quad (= 2\kappa(S, \lambda_1) \text{ if } \lambda_1 \text{ is simple}).$$

(i) *Let $\hat{S}_1 = S + X R_1 X^T J^T S$ be constructed according to Theorem 3 with R_1 from (23). Then*

$$\frac{\|\hat{S}_1 - S\|_F}{\|S\|_F} \leq \frac{|\lambda_1 - \mu|}{|\lambda_1|} \left[\Phi \sqrt{1 + \frac{|\lambda_1|^2}{|\mu|^2}} \right]. \quad (26)$$

(ii) *Let $\hat{S}_2 = S + X R_2 X^T J^T S$ be constructed according to Theorem 3 with R_2 from (23). Then*

$$\frac{\|\hat{S}_2 - S\|_F}{\|S\|_F} \leq \frac{|\lambda_1^{-1} - \mu|}{|\lambda_1^{-1}|} \left[\Phi \sqrt{1 + \frac{|\lambda_1^{-1}|^2}{|\mu|^2}} \right]. \quad (27)$$

As the following example shows, similar easy bounds can be found with the use of R_1 and R_2 when $\|\cdot\|_2$ is considered. In fact, in the 2-norm, such a bound can be sharp.

Example 1. For a symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$, the bound (19) for $\hat{S}_1 = S + X R_1 X^T J^T S$ with respect to $\|\cdot\|_2$ can easily be determined as

$$\frac{\|\hat{S}_1 - S\|_2}{\|S\|_2} \leq \frac{|\lambda_1 - \mu|}{|\lambda_1|} \cdot \max \left\{ 1, \frac{|\lambda_1|}{|\mu|} \right\} \|X\|_2^2, \quad (28)$$

It can be seen for $S = \text{diag}(\Lambda, \Lambda^{-1})$ with $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ that the bound in (28) can be sharp. In particular, with eigenvectors $e_1, e_{n+1} \in \mathbb{R}^{2n}$ for $\lambda_1, \lambda_1^{-1}$, respectively, and $X = [e_1 \ e_{n+1}]$ we have

$$X R_1 X^T J^T S = \text{diag}(r_{12}\lambda_1, 0, \dots, 0, -r_{21}\lambda_1^{-1}, 0, \dots, 0)$$

with nonzero entries in the first and $(n+1)$ st position. As $r_{12}\lambda_1 = \mu - \lambda_1$ and $-r_{21}\lambda_1^{-1} = \mu^{-1} - \lambda_1^{-1}$ we obtain under the assumption $|\lambda_1 - \mu| \geq |\lambda_1^{-1} - \mu^{-1}|$

$$\|\hat{S} - S\|_2 = \|XR_1X^TJ^TS\|_2 = |\lambda_1 - \mu|$$

and so $\|\hat{S} - S\|_2/\|S\|_2 = |\lambda_1 - \mu|/|\lambda_1|$ if λ_1 is the largest eigenvalue of S in absolute value (i.e. $\|S\|_2 = |\lambda_1|$). On the other hand, for X we certainly have $\|X\|_2^2 = 1$ and thus, whenever $|\mu| \geq |\lambda_1|$, the bound on the right hand side in (28) also reduces to $|\lambda_1 - \mu|/|\lambda_1|$.

3.1. IMPROVED DISTANCE BOUNDS

Although the bound in (26) nicely relates $\|\hat{S}_1 - S\|_F/\|S\|_F$ to the relative value change $|\lambda_1 - \mu|/|\lambda_1|$ and the condition number $\kappa(S, \lambda_1)$, it can be quite bad⁶, see e.g. Fig. 1 in Section 5. In this section we derive sharper bounds under the additional assumption that $\|S\|_F$ is known.

As before, let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with eigenvectors $x_1, x_2 \in \mathbb{C}^{2n}$ for $\lambda_1, \lambda_1^{-1} \in \sigma(S)$, respectively, such that $x_1^T J x_2 \neq 0$ (and set $X = [x_1 \ x_2]$). As seen in (14), we have for $\Lambda := \text{diag}(\lambda_1, \lambda_1^{-1})$ and $R \in \mathbf{M}_2(\mathbb{C})$ that satisfies (12) and (13)

$$\hat{S} = S + XR_1X^TJ^TS = S + XRA^{-1}X^TJ^T$$

and therefore $\|S - \hat{S}\|_F = \|XRA^{-1}X^TJ^T\|_F = \|XRA^{-1}X^T\|_F$. Whenever $R = R_j$ ($j = 1, 2$) from (23), then

$$XR_j\Lambda^{-1}X^T = X \begin{bmatrix} 0 & \eta_j \\ \eta_j - d & 0 \end{bmatrix} X^T =: X\tilde{R}_jX^T, \quad \tilde{R}_j = R_j\Lambda^{-1},$$

according to (20). Recall the solutions of $p(\eta) = 0$ in (21), i.e. $\eta_1 = \mu - \lambda_1$ (corresponding to R_1) and $\eta_2 = \mu^{-1} - \lambda_1$ (corresponding to R_2). Instead of estimating $\|X\tilde{R}_jX^T\|_F$ by $\|\tilde{R}_j\|_F\|X\|_F^2$ we now intend to estimate $\|X\tilde{R}_jX^T\|_F$ directly. To this end, assume that $X \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ has the form (24) and $\hat{S}_j = S + XR_jX^TJ^TS$, $j = 1, 2$. Then we have for $Y = \sqrt{x_1^T J x_2} X = [x_1 \ x_2]$

$$\begin{aligned} \|S - \hat{S}_j\|_F^2 &= \|X\tilde{R}_jX^T\|_F^2 = \frac{\text{tr}(Y\tilde{R}_jY^T(Y\tilde{R}_jY^T)^H)}{|x_1^T J x_2|^2} \\ &= \frac{\text{tr}(Y\tilde{R}_jY^T\bar{Y}\tilde{R}_j^HY^H)}{|x_1^T J x_2|^2} = \frac{\text{tr}(Y^HY\tilde{R}_j\overline{(Y^HY)}\tilde{R}_j^H)}{|x_1^T J x_2|^2}. \end{aligned}$$

⁶The same is true for the bound in (27).

Now we further obtain

$$\begin{aligned}
|x_1^T J x_2|^2 \|\hat{S}_j - S\|_F^2 &= \text{tr}(Y^H Y \tilde{R}_j \overline{(Y^H Y)} \tilde{R}_j^H) \\
&= \text{tr} \left(\begin{bmatrix} 1 & x_1^H x_2 \\ x_2^H x_1 & 1 \end{bmatrix} \begin{bmatrix} 0 & \eta_j \\ \eta_j - d & 0 \end{bmatrix} \begin{bmatrix} 1 & x_2^H x_1 \\ x_1^H x_2 & 1 \end{bmatrix} \begin{bmatrix} 0 & \bar{\eta}_j - \bar{d} \\ \bar{\eta}_j & 0 \end{bmatrix} \right) \\
&= |x_1^H x_2|^2 \bar{\eta}_j (\eta_j - d) + |\eta_j|^2 + |\eta_j - d|^2 + |x_1^H x_2|^2 \eta_j (\bar{\eta}_j - \bar{d}) \\
&= |\eta_j|^2 + |\eta_j - d|^2 + 2|x_1^H x_2|^2 \cdot \Re(\eta_j(\bar{\eta}_j - \bar{d})). \tag{29}
\end{aligned}$$

Note that $|x_1^H x_2| \leq \|x_1\|_2 \|x_2\|_2 = 1$ as x_1 and x_2 are normalized. Furthermore, $\Re(\eta_j(\bar{\eta}_j - \bar{d})) = \Re(|\eta_j|^2 - \eta_j \bar{d}) = |\eta_j|^2 - \Re(\eta_j \bar{d})$, and we may now derive upper (and lower) bounds for (29) depending on whether this term is positive or negative.

- (i) Suppose $\Re(\eta_j \bar{d}) < |\eta_j|^2$. Then $\Re(\eta_j(\bar{\eta}_j - \bar{d})) > 0$ follows and, since $|x_1^H x_2|^2 \leq 1$, we can estimate from (29), setting $|x_1^H x_2|^2 = 1$,

$$\begin{aligned}
|x_1^T J x_2|^2 \|S - \hat{S}_j\|_F^2 &\leq |\eta_j|^2 + |\eta_j - d|^2 + 2 \cdot \Re(\eta_j(\bar{\eta}_j - \bar{d})) \\
&= (\eta_j + (\eta_j - d))(\bar{\eta}_j + (\bar{\eta}_j - \bar{d})) = |2\eta_j - d|^2.
\end{aligned}$$

On the other hand, changing the sign of $\Re(\eta_j(\bar{\eta}_j - \bar{d}))$ we certainly have

$$\begin{aligned}
|x_1^T J x_2|^2 \|S - \hat{S}_j\|_F^2 &\geq |\eta_j|^2 + |\eta_j - d|^2 - 2 \cdot \Re(\eta_j(\bar{\eta}_j - \bar{d})) \\
&= (\eta_j - (\eta_j - d))(\bar{\eta}_j - (\bar{\eta}_j - \bar{d})) = |d|^2.
\end{aligned}$$

- (ii) Suppose $\Re(\eta_j \bar{d}) \geq |\eta_j|^2$. Then $\Re(\eta_j(\bar{\eta}_j - \bar{d})) \leq 0$ follows and we can estimate from (29), setting again $|x_1^H x_2|^2 = 1$,

$$|x_1^T J x_2|^2 \|S - \hat{S}_j\|_F^2 \geq |\eta_j|^2 + |\eta_j - d|^2 + 2 \cdot \Re(\eta_j(\bar{\eta}_j - \bar{d})) = |2\eta_j - d|^2$$

while on the other hand, with a change of sign, we obtain

$$|x_1^T J x_2|^2 \|S - \hat{S}_j\|_F^2 \leq |\eta_j|^2 + |\eta_j - d|^2 - 2 \cdot \Re(\eta_j(\bar{\eta}_j - \bar{d})) = |d|^2.$$

Before we state our findings in the next theorem, notice that there are neat expressions for the terms $2\eta_j - d$, $j = 1, 2$, arising above, i.e.

$$2\eta_1 - d = \frac{\mu - \lambda_1}{\lambda_1} (\lambda_1 + \mu^{-1}), \quad 2\eta_2 - d = \frac{\lambda_1^{-1} - \mu}{\lambda_1^{-1}} (\mu^{-1} + \lambda_1^{-1}).$$

As it turns out, also d can be rewritten in a similar fashion as

$$d = \frac{\mu - \lambda_1}{\lambda_1} (\lambda_1 - \mu^{-1}) = \frac{\lambda_1^{-1} - \mu}{\lambda_1^{-1}} (\mu^{-1} - \lambda_1^{-1}). \tag{30}$$

Finally, the two conditions to be checked in (i) and (ii) above can be simplified. For $\eta_1 = \mu - \lambda_1$ one finds, after some reformulations,

$$\begin{aligned}
\Re(\eta_1 \bar{d}) - |\eta_1|^2 &= \frac{1}{2}((\mu - \lambda_1)\bar{d} + (\bar{\mu} - \bar{\lambda}_1)d) - |\mu - \lambda_1|^2 \\
&= \frac{1}{2} \left(\frac{|\mu - \lambda_1|^2 \bar{d}}{\bar{\mu} - \bar{\lambda}_1} + \frac{|\mu - \lambda_1|^2 d}{\mu - \lambda_1} \right) - |\mu - \lambda_1|^2 \\
&= -|\mu - \lambda_1|^2 \left(1 - \frac{1}{2} \left(\frac{\bar{d}}{\bar{\mu} - \bar{\lambda}_1} + \frac{d}{\mu - \lambda_1} \right) \right) \\
&= -|\mu - \lambda_1|^2 \left(1 - \Re \left(\frac{d}{\mu - \lambda_1} \right) \right) \\
&= -|\mu - \lambda_1|^2 \left(1 - \Re \left(\frac{\lambda_1 - \mu^{-1}}{\lambda_1} \right) \right) \\
&= -|\mu - \lambda_1|^2 \Re((\mu \lambda_1)^{-1})
\end{aligned}$$

where we used the first expression for d in (30) in the second-last equation.

Thus $\Re(\eta_1 \bar{d}) \geq |\eta_1|^2$ holds if and only if $-|\mu - \lambda_1|^2 \Re((\mu \lambda_1)^{-1}) \geq 0$, which is the case if and only if $\Re(\lambda_1 \mu) \leq 0$ as $(\lambda_1 \mu)^{-1}$ and $\lambda_1 \mu$ are located in the same half plane. For $\eta_2 = \mu^{-1} - \lambda_1$ we obtain analogously

$$\Re(\eta_2 \bar{d}) - |\eta_2|^2 = -|\lambda_1 \mu - 1|^2 \Re((\bar{\lambda}_1 \mu)^{-1})$$

and so $\Re(\eta_2 \bar{d}) \geq |\eta_2|^2$ holds if and only if $\Re((\bar{\lambda}_1 \mu)^{-1}) \leq 0$. This, in turn, holds if and only if $\Re(\bar{\lambda}_1 \mu) \leq 0$. In conclusion, we have proven the following theorem.

Theorem 6. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1, \lambda_1^{-1} \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Let $x_1, x_2 \in \mathbb{C}^{2n}$ be normalized eigenvectors for λ_1 and λ_1^{-1} , respectively, with $x_1^T J x_2 \neq 0$ and $X \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ as in (24). Define*

$$\Phi := \frac{1}{|x_1^T J x_2| \cdot \|S\|_F} \left(= \frac{\kappa(S, \lambda_1)}{\|S\|_F} \text{ if } \lambda_1 \text{ is simple} \right)$$

(i) *Let $\hat{S}_1 = S + X R_1 X^T J^T S$ be constructed according to Theorem 3 with R_1 from (23). Whenever $\Re(\lambda_1 \mu) \leq 0$, then*

$$\frac{|\lambda_1 - \mu|}{|\lambda_1|} (|\lambda_1 + \mu^{-1}| \Phi) \leq \frac{\|\hat{S}_1 - S\|_F}{\|S\|_F} \leq \frac{|\lambda_1 - \mu|}{|\lambda_1|} (|\lambda_1 - \mu^{-1}| \Phi).$$

If $\Re(\lambda_1 \mu) > 0$ the upper and lower bounds interchange.

(ii) *Let $\hat{S}_2 = S + X R_2 X^T J^T S$ be constructed according to Theorem 3 with R_2 from (23). Whenever $\Re(\bar{\lambda}_1 \mu) \leq 0$, then*

$$\frac{|\lambda_1^{-1} - \mu|}{|\lambda_1^{-1}|} (|\lambda_1^{-1} + \mu^{-1}| \Phi) \leq \frac{\|\hat{S}_2 - S\|_F}{\|S\|_F} \leq \frac{|\lambda_1^{-1} - \mu|}{|\lambda_1^{-1}|} (|\lambda_1^{-1} - \mu^{-1}| \Phi).$$

If $\Re(\bar{\lambda}_1 \mu) > 0$ the upper and lower bounds interchange.

It is shown in Section 5 (see Fig. 1) that the bounds in Theorem 6 are significantly sharper compared to the bounds in (26) and (27).

Remark 1. For any matrix $R = [r_{ij}] \in \mathbf{M}_2(\mathbb{C})$ that satisfies the conditions (12) and (13) the bounds in (19) can easily be calculated. However, there are several reasons for not considering other choices of R (beside R_1 and R_2 from (23)) in this section in detail:

- (a) If $r_{11} \neq 0, r_{22} \neq 0$, there are two possibilities for R whose entries r_{12} and r_{21} of R depend on $c = r_{11}r_{22}$ through (one of) the zeros of $p(\eta) = 0$, see (21). Thus, r_{12} and r_{21} involve the expression of a complex square root and there is no neat and compact expression for $\|R\|_F$ compared to (26) and (27) or to the formulas in Theorem 6. Furthermore, minimizing $\|S - \hat{S}\|_F$ with respect to the entries of R under the side conditions (12) and (13) results in a difficult complex optimization problem for which the author is not aware of a closed form solution.
- (b) Among all matrices R that satisfy the conditions (12) and (13) the matrices R_1 and R_2 from (23) are the only possible choice when $\hat{S} = S + XRX^T J^T S$ should inherit desirable properties (e.g. related to diagonalizability) from S . These distinguishing features of R_1 and R_2 are discussed in the upcoming sections.
- (c) All numerical experiments that have been performed indicate that rarely a matrix $R' \in \mathbf{M}_2(\mathbb{C})$ different from R_1 and R_2 that satisfies (12) and (13) was detected such that $\|\hat{S} - S\|_F / \|S\|_F$ for $\hat{S} = S + X R' X^T J^T S$ was smaller than the minimum of $\|\hat{S}_1 - S\|_F / \|S\|_F$ and $\|\hat{S}_2 - S\|_F / \|S\|_F$. This is visualized in Section 5, see Figure 3.

4. SEGRE CHARACTERISTICS AND COMMUTATIVITY RELATIONS

In this section we discuss how the Segre characteristics of eigenvalues are effected by a change of a symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ to $\hat{S} \in \mathbf{M}_{2n}(\mathbb{C})$ according to Theorem 3. In particular, we will show that the Segre characteristics of the eigenvalues of S and \hat{S} are either the same or connected in a direct way. Furthermore, we make a statement on eigenvectors of S and \hat{S} that remain unchanged. Notice that, in the context of Theorem 2, the eigenvectors of A and $\hat{A} = A + XC^T$ are in general all different and *not* related in an immediate fashion [3] if no further restrictions are imposed on the form of C . In this section we show that, in the structure-preserving context of Theorem 3, the particular form of C allows for some explicit statements. Furthermore, we derive statements on the diagonalizability of \hat{S} and the simultaneous diagonalizability of S and \hat{S} . To this end, we begin in Section 4.1 with a result on the commutativity of S and \hat{S} .

4.1. THE COMMUTATIVITY OF S AND \hat{S}

Recall that the matrix \hat{S} in (11) can also be expressed as

$$\hat{S} = (I_{2n} + XRX^T J^T)S. \quad (31)$$

Since \hat{S} and S are both symplectic, the matrix $\hat{S}S^{-1} = \hat{S}S^* = I_{2n} + XRX^T J^T$ is symplectic, too. As for any $A, B \in \mathbf{M}_n(\mathbb{C})$ the matrices AB and BA always have the same eigenvalues [9], beside \hat{S} , we may also define the symplectic matrix $\tilde{S} := S(I_{2n} + XRX^T J^T) \in \mathbf{M}_{2n}(\mathbb{C})$ that solves the eigenvalue modification problem stated in Section 1. A question naturally arising is whether there is a connection between \hat{S} from (31) and \tilde{S} . Such a connection is revealed in Theorem 7 which shows a distinguishing feature of the matrices from (23) among all matrices R that satisfy (12) and (13), see Remark 1 (b). The result from Theorem 7 will be used when the diagonalizability of \hat{S} is investigated in Section 4.2.

Theorem 7. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and assume $\hat{S} = S + XRX^T J^T S$ has been constructed according to Theorem 3. Then the following is true:*

1. In case $\lambda_1 \neq \pm 1$,

$$\hat{S} = (I_{2n} + XRX^T J^T)S = S(I_{2n} + XRX^T J^T) = \tilde{S} \quad (32)$$

holds if and only if $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ is one of the matrices in (23).

2. In case $\lambda_1 = \pm 1$, (32) holds for any $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ satisfying (12) and (13).

Proof. First, notice that $\hat{S} = \tilde{S}$ is equivalent to

$$XRX^T J^T S = SXRX^T J^T. \quad (33)$$

Multiplying both equations with J (from the right) and using the relations $SX = X\Lambda$ (where $\Lambda = \text{diag}(\lambda_1, \lambda_1^{-1})$) and $J^T S J = S^{-T}$ yields $XRX^T S^{-T} = X\Lambda R X^T$. Moreover, $X^T S^{-T} = (S^{-1} X)^T = (X\Lambda^{-1})^T = \Lambda^{-1} X^T$ and, equivalently to (33), it suffices to investigate the equation

$$X R \Lambda^{-1} X^T = X \Lambda R X^T. \quad (34)$$

Now, as $X \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ has full rank, (34) is (by the multiplication with X^+ from the left and $(X^T)^+$ from the right) equivalent to $R\Lambda^{-1} = \Lambda R$, that is, $\Lambda R \Lambda = R$. For $R = [r_{ij}]_{ij}$ we obtain

$$\Lambda R \Lambda = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_1^{-1} \end{bmatrix} = \begin{bmatrix} \lambda_1^2 r_{11} & r_{12} \\ r_{21} & \lambda_1^{-2} r_{22} \end{bmatrix}.$$

This shows that $\Lambda R \Lambda = R$ holds, in case $\lambda_1 \neq \pm 1$, if and only if $r_{11} = r_{22} = 0$. The two possibilities for R that satisfy the conditions (12) and (13) when $r_{11} = r_{22} = 0$ are the matrices in (23). Furthermore, if $\lambda_1 = \pm 1$, the equation always holds. This completes the proof. \square

Theorem 7 shows that, in general (i.e. for $\lambda_1 \neq \pm 1$), only the two possible choices for R in (23) produce commutativity of S and $I_{2n} + XRX^T J^T$ (i.e. to have $\hat{S} = \tilde{S}$). In the special case $\lambda_1 = \pm 1$, any matrix R determined from (12) and (13) will cause this commutativity relation.

4.2. SEGRE CHARACTERISTICS

Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with eigenvalues $\lambda_1, \lambda_1^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. To analyse the consequences of the change $S \mapsto \hat{S} = S + XRX^T J^T S$ on the Segre characteristics of the eigenvalues of S and \hat{S} , we discuss the cases of $\lambda_1, \lambda_1^{-1}$ (the eigenvalues that are changed), μ, μ^{-1} (the values λ_1 and λ_1^{-1} are changed to) and all other eigenvalues (which are the same for S and \hat{S}) separately. As before, let $x_1, x_2 \in \mathbb{C}^{2n}$ be eigenvectors for λ_1 and λ_1^{-1} , respectively, normalized such that $X^T J_{2n} X = J_2$ for $X = [x_1 \ x_2] \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ and assume $\hat{S} = S + XRX^T J^T S$ has been constructed as in Theorem 3.

We first consider eigenvalues different from $\lambda_1, \lambda_1^{-1}, \mu$ and μ^{-1} . These eigenvalues and their algebraic multiplicities are the same for S and \hat{S} and we show that their eigenspaces and Jordan chains (thus, in consequence, their Segre characteristics) remain completely unchanged. To prove this, we need the following fact about the matrix S and its (generalized) eigenvectors (see also [10, Sec. 2]): assume that λ is some eigenvalue of S different from λ_1 and λ_1^{-1} and let $y_1, y_2, \dots, y_p \in \mathbb{C}^{2n}$ ($p \geq 1$) be a Jordan chain for S and λ , i.e. it holds that $(S - \lambda I_{2n})y_1 = 0$ and $(S - \lambda I_{2n})y_{k+1} = y_k$ for $k = 1, \dots, p-1$. Then $X^T J y_k = 0$ follows for any $k = 1, \dots, p$. To see this, first consider the eigenvector y_1 of S for λ . We have

$$\lambda_1 x_1^T J y_1 = x_1^T S^T J y_1 = x_1^T J J^T S^T J y_1 = x_1^T J S^{-1} y_1 = \lambda^{-1} x_1^T J y_1$$

This shows that $x_1^T J y_1 = 0$ if $\lambda \neq \lambda_1^{-1}$. Similarly, $x_2^T J y_1 = 0$ follows for $\lambda \neq \lambda_1$. Now assume that $x_i^T J y_\ell = 0$ holds for $i = 1, 2$ and $\ell = 1, \dots, k$. For $(S - \lambda I_{2n})y_{k+1} = y_k$ we thus obtain

$$\begin{aligned} 0 &= x_1^T J y_k = x_1^T J (S - \lambda I_{2n}) y_{k+1} = x_1^T J S y_{k+1} - \lambda x_1^T J y_{k+1} \\ &= x_1^T S^{-T} J y_{k+1} - \lambda x_1^T J y_{k+1} \\ &= \lambda_1^{-1} x_1^T J y_{k+1} - \lambda x_1^T J y_{k+1} \\ &= (\lambda_1^{-1} - \lambda) x_1^T J y_{k+1}. \end{aligned}$$

Therefore, again $x_1^T J y_{k+1} = 0$ follows whenever $\lambda \neq \lambda_1^{-1}$. With the same reasoning we obtain $x_2^T J y_{k+1} = 0$ for $\lambda \neq \lambda_1$. In conclusion we have $X^T J y_k = 0$ for any $k = 1, \dots, p$ whenever $\lambda_1 \neq \lambda \neq \lambda_1^{-1}$.

Lemma 1. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Suppose that $\hat{S} \in \mathbf{M}_{2n}(\mathbb{C})$ has been constructed according to Theorem*

3. Then for any $\lambda \in \sigma(\hat{S})$ which is neither equal to λ_1 or λ_1^{-1} nor equal to μ or μ^{-1} the Segre characteristics of λ as an eigenvalue of S and \hat{S} and their corresponding Jordan chains, respectively, are identical.

Proof. Assume $\lambda \in \sigma(\hat{S})$ is neither equal to λ_1 or λ_1^{-1} nor equal to μ or μ^{-1} . By construction of $\hat{S} = S + XRX^T J^T S$, λ is an eigenvalue of both S and \hat{S} with the same algebraic multiplicities. Whenever $y_1 \in \mathbb{C}^{2n}$ is an eigenvector of S for λ it is also an eigenvector of \hat{S} for λ since $X^T J y_1 = 0$, which implies

$$\hat{S}y_1 = Sy_1 + XRX^T J^T S y_1 = Sy_1 - \lambda XRX^T J y_1 = Sy_1 = \lambda y_1. \quad (35)$$

Next, let $y_1, \dots, y_p \in \mathbb{C}^{2n}$ be a Jordan chain for S and λ . Then

$$\begin{aligned} (\hat{S} - \lambda I_{2n})y_{i+1} &= (S + XRX^T J^T S)y_{i+1} - \lambda y_{i+1} \\ &= (y_i + \lambda y_{i+1}) + XRX^T J^T S y_{i+1} - \lambda y_{i+1} \\ &= y_i - XRX^T J(y_i + \lambda y_{i+1}) = y_i \end{aligned}$$

since $X^T J y_k = 0$ for any y_k , $k = 1, \dots, p$, from the Jordan chain. Inductively, this shows that y_1, \dots, y_p remains to be a Jordan chain of \hat{S} for λ . Therefore, the Segre characteristic for λ of S and \hat{S} and the corresponding Jordan chains are the same. \square

Next we consider λ_1 and λ_1^{-1} . When $S \in \mathbf{M}_{2n}(\mathbb{C})$ is transformed to $\hat{S} = S + XRX^T J^T S$ and (one instance of) $\lambda_1, \lambda_1^{-1}$ is replaced by μ and μ^{-1} , the Segre characteristic of λ_1 for \hat{S} is necessarily different from its Segre characteristic for S due to the eigenvalue modification that has taken place (if λ_1 is a simple eigenvalue of S , then it is not even an eigenvalue of \hat{S} anymore). However, if the algebraic multiplicity of λ_1 as an eigenvalue of S is ≥ 2 , then the Segre characteristics of λ_1 as an eigenvalue of S and \hat{S} are connected in an easy fashion (see Theorem 8 below). This is obviously false in the general context of Rado's Theorem, where nontrivial Jordan blocks may arise, as the following counterexample for $A = I_4$ and $\hat{A} = A + XC^T$ shows:

$$\hat{A} = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & & & \\ & 1 & 1 & \\ & & 1 & \\ & & & 1 \end{bmatrix}.$$

In this example, the Segre characteristic of $1 \in \sigma(A)$ is $((1, 1, 1, 1))$ while it is $((2, 1))$ for \hat{A} .

Theorem 8. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0, \lambda_1, \lambda_1^{-1}\}$ be given. Suppose that $\hat{S} = S + XRX^T J^T S \in \mathbf{M}_{2n}(\mathbb{C})$ has been constructed according to Theorem 3. Then the following hold:*

(i) If the Segre characteristic of $\lambda_1 \neq \pm 1$ as an eigenvalue of S is

$$((s_k, s_{k-1}, \dots, s_2, s_1)) \quad (36)$$

with⁷ $s_k \geq s_{k-1} \geq \dots \geq s_2 \geq s_1 = 1$, then the Segre characteristic of λ_1 as an eigenvalue of \hat{S} is $((s_k, s_{k-1}, \dots, s_2))$. Moreover, if $\lambda_1 = \pm 1$ and (36) is its Segre characteristic of S with⁷ $s_2 = s_1 = 1$, then the Segre characteristic of λ_1 as an eigenvalue of \hat{S} is $((s_k, s_{k-1}, \dots, s_3))$.

(ii) Let $\mu \notin \sigma(S)$. Then the Segre characteristic of μ as an eigenvalue of \hat{S} is always ((1)) if $\mu \neq \mu^{-1}$. If $\mu = \mu^{-1}$ its Segre characteristic is ((1, 1)) if and only if $R = R_1$ or $R = R_2$ from (23), otherwise it is ((2)).

Proof. (i) We first assume $\lambda_1 \neq \lambda_1^{-1}$. According to the Segre characteristic $((s_k, \dots, s_1))$ of $\lambda_1 \in \sigma(S)$ there are $k \geq 1$ Jordan blocks L_k, \dots, L_1 of sizes s_k, \dots, s_1 . As $s_1 = 1$ let x_1 be the corresponding eigenvector. We denote the generalized eigenvectors corresponding to the ℓ -th Jordan block L_ℓ by $x_1^\ell, \dots, x_{s_\ell}^\ell$ and set $X_\ell = [x_1^\ell \ \dots \ x_{s_\ell}^\ell]$ and $\tilde{X} := [X_2 \ \dots \ X_k]$. Since $\lambda_1^{-1} \in \sigma(S)$ has the same Segre characteristic as λ_1 , there are also k Jordan blocks G_k, \dots, G_1 of S for λ_1^{-1} of sizes s_k, \dots, s_1 . Let $y_1, Y_\ell = [y_1^\ell \ \dots \ y_{s_\ell}^\ell]$ and $\tilde{Y} := [Y_2 \ \dots \ Y_k]$ be defined analogously from the (generalized) eigenvectors for λ_1^{-1} . We now define the matrix $U := [x_1 \ y_1 \ \tilde{X} \ \tilde{Y}] \in \mathbf{M}_{2n \times 2p}(\mathbb{C})$. Then the matrix $\hat{S} = S + XRX^T J^T S$ can be written as

$$\hat{S} = S + \begin{bmatrix} x_1 & y_1 & \tilde{X} & \tilde{Y} \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \\ 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \end{bmatrix} X^T J^T S =: S + U \tilde{R} X^T J^T S.$$

As $SU = UP'$ for $P' := \text{diag}(\lambda_1, \lambda_1^{-1}, L_2, \dots, L_k, G_2, \dots, G_k)$ we therefore obtain $\hat{S}U = SU + U \tilde{R} X^T J^T S U$ and thus $\hat{S}U = U(P' + \tilde{R} X^T J^T U P')$. Now set $P = [p_{ij}]_{ij} := P' + \tilde{R} X^T J^T U P'$ and notice that the third to last row of $\tilde{R} X^T J^T U P'$ are identically zero (due to the form of \tilde{R}). Therefore, the form of P can be explicitly determined (with \star indicating zero or nonzero entries

⁷Notice that for Theorem 3 to be applicable to $\lambda_1 \neq \pm 1$, $s_1 = 1$ is a necessary condition according to Theorem 4. If $\lambda_1 = \pm 1$, then $s_1 = 1$ implies $s_2 = 1$ since then Jordan blocks of a particular size must appear an even number of times in the Jordan structure of S .

that are not of further interest):

$$P = \left[\begin{array}{cc|ccc} \lambda_1(1+r_{12}) & -\lambda_1^{-1}r_{11} & \star & \cdots & \star & \star & \cdots & \star \\ \lambda_1 r_{22} & \lambda_1^{-1}(1-r_{21}) & \star & \cdots & \star & \star & \cdots & \star \\ \hline 0 & 0 & L_2 & & & & & 0 \\ \vdots & \vdots & & \ddots & & & & \\ \vdots & \vdots & & & L_k & & & \\ \hline \vdots & \vdots & & & & G_2 & & \\ \vdots & \vdots & 0 & & & & \ddots & \\ 0 & 0 & & & & & & G_k \end{array} \right]. \quad (37)$$

Now, it is easily seen that $L_2, \dots, L_k, G_2, \dots, G_k$ are part of the Jordan structure of P , hence they also arise in the Jordan structure of \hat{S} . This shows that the Segre characteristic of $\lambda_1 \neq \pm 1 \in \sigma(\hat{S})$ and $\lambda_1^{-1} \in \sigma(\hat{S})$ is both $((s_k, s_{k-1}, \dots, s_2))$. If $\lambda = \pm 1$ the proof follows the same lines without the use of \tilde{Y} . To prove (ii) we note that the upper-left 2×2 block of P is exactly Ω from (9), hence its eigenvalues are μ, μ^{-1} . If $\mu \neq \mu^{-1}$, then Ω is semisimple. Therefore, we obtain the Segre characteristic of μ and μ^{-1} as eigenvalues of \hat{S} both as ((1)). On the other hand, if $\mu = \mu^{-1}$, then Ω is semisimple if and only if its minimal polynomial is $p(z) = z - \mu$. Now

$$p(\Omega') = \begin{bmatrix} \lambda_1(1+r_{12}) - \mu & -\lambda_1^{-1}r_{11} \\ \lambda_1 r_{22} & \lambda_1^{-1}(1-r_{21}) - \mu \end{bmatrix}.$$

Thus, for $p(\Omega) = 0$ we must have $r_{11} = r_{22} = 0$ and the only possible choices for Ω to be semisimple are R_1 and R_2 from (23). It is easy to check that in fact both choices result in $p(\Omega) = 0$. This proves that the Segre characteristic of μ as an eigenvalue of \hat{S} is $((1, 1))$ if $R = R_1, R_2$ and that it has to be ((2)) otherwise. \square

Finally, assume that μ was already an eigenvalue of S and Theorem 3 is applied for $\lambda_1 \in \sigma(S)$. Then the arguments of the proof of Lemma 1 apply to μ (as an 'old' eigenvalue of S) as well as the result from Theorem 8 (ii) (for μ as the 'new' eigenvalue appearing in the spectrum of \hat{S}). Thus, the Segre characteristic of $\mu \in \sigma(\hat{S})$ is its old Segre characteristic from S extended by one of the cases described in Theorem 8 (ii).

As another consequence of Theorem 8 it follows that, if λ_1 is semisimple for S , then λ_1 remains semisimple for \hat{S} is case its multiplicity was ≥ 2 . Moreover, assume the symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ is diagonalizable (i.e. all its eigenvalues are semisimple) and let \hat{S} be constructed according to Theorem 3. As a consequence of Lemma 1 and Theorem 8, the diagonalizability of \hat{S} can then only be circumvented in case a 2×2 Jordan block arises for $\mu = \mu^{-1}$. As seen above, a 2×2 Jordan block for μ will arise if R is different from R_1

and R_2 . In other words, if R_1 or R_2 from (23) are chosen in Theorem 3, the matrix \hat{S} will be semisimple in case S was. In fact, this is the only situation in which S and \hat{S} are simultaneously diagonalizable since the simultaneous diagonalizability of S and \hat{S} is only possible if S and \hat{S} commute. According to Theorem 7 this is the case if and only if $R = R_1, R_2$ are chosen. We summarize this result in the following corollary.

Corollary 2. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with $\lambda_1 \in \sigma(S)$ and let $\mu \in \mathbb{C} \setminus \{0, \lambda_1, \lambda_1^{-1}\}$ be given. Suppose that $\hat{S} = S + XRX^T J^T S \in \mathbf{M}_{2n}(\mathbb{C})$ has been constructed according to Theorem 3 and that S is semisimple. Then \hat{S} is semisimple if and only if $R = R_1$ or $R = R_2$ for one of the matrices in (23). Moreover, in this case, S and \hat{S} are simultaneously diagonalizable.*

Example 2 below shows how the simultaneous diagonalization looks like if R_1 or R_2 are used to construct \hat{S} .

Example 2. Whenever $T^{-1}ST = \text{diag}(\Lambda, \Lambda^{-1})$ with $T = [x_1 \ Y \ x_2 \ Z] \in \mathbf{M}_{2n}(\mathbb{C})$ and $Y, Z \in \mathbf{M}_{2n \times (n-1)}(\mathbb{C})$ and $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$, then for $S_j = S + XR_j X^T J^T S$ with R_1, R_2 from (23) we have

$$T^{-1}\hat{S}_1 T = \begin{bmatrix} \Lambda_1 & \\ & \Lambda_1^{-1} \end{bmatrix}, \quad \text{and} \quad T^{-1}\hat{S}_2 T = \begin{bmatrix} \Lambda_2 & \\ & \Lambda_2^{-1} \end{bmatrix}$$

where $\Lambda_1 = \text{diag}(\mu, \lambda_2, \dots, \lambda_n)$ and $\Lambda_2 = \text{diag}(\mu^{-1}, \lambda_2, \dots, \lambda_n)$. In fact, for S being diagonal, R_1 and R_2 are the only possible choice such that \hat{S} is diagonal, too.

4.3. A NOTE ON CONDITION NUMBERS

We conclude this section with a result on eigenvalue condition numbers. It is a surprising fact, that we can apply Theorem 3 to a symplectic matrix $S \in \mathbf{M}_{2n}(\mathbb{C})$ without changing any eigenvalue condition number (for simple eigenvalues) at all. For Rado's theorem this is not true: an unstructured application of Theorem 2 typically changes all eigenvalue condition numbers, even those of eigenvalues that remain unchanged. The main result on the behavior of condition numbers is stated in Theorem 9. For its proof, we need the following well-known fact that we state without proof in Lemma 2.

Lemma 2. *Let $A \in \mathbf{M}_n(\mathbb{C})$ and suppose x is a right eigenvector of A for λ (i.e. $Ax = \lambda x$) and y is a left eigenvector of A for μ (i.e. $y^H A = \mu y^H$). Then $y^H x = 0$ if $\lambda \neq \mu$.*

Theorem 9. *Let $S \in \mathbf{M}_{2n}(\mathbb{C})$ be symplectic with eigenvalues $\lambda_1, \lambda_1^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ and let $\mu \in \mathbb{C} \setminus \{0\}$ with $\mu \notin \sigma(S)$ be given. Assume that λ_1 is simple and let $\hat{S} = S + XRX^T J^T S$ be constructed according to Theorem 3 so that $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ are the eigenvalues of \hat{S} . Then the following holds:*

(i) If ν is a simple eigenvalue of \hat{S} , $\mu \neq \nu \neq \mu^{-1}$, then $\kappa(\hat{S}, \nu) = \kappa(S, \nu)$.

(ii) For $R = R_1$, where R_1 is the matrix given in (23), it holds that

$$\kappa(\hat{S}, \mu) = \kappa(S, \lambda_1) \quad \text{and} \quad \kappa(\hat{S}, \mu^{-1}) = \kappa(S, \lambda_1^{-1}).$$

Proof. (i) Under the assumption $\mu^{-1} \neq \nu \neq \mu$ and the simplicity of λ_1 , it follows that $\nu = \lambda_j$ or $\nu = \lambda_j^{-1}$ for some $j = 2, \dots, n$. Assume w.l.o.g. that $\nu = \lambda_2$ and let $y_1 \in \mathbb{C}^{2n}$ be some corresponding eigenvector of S . According to Lemma 1, $\hat{S}y_1 = \lambda_2 y_1$ holds. Next suppose $z_1 \in \mathbb{C}^{2n}$ is some left eigenvector of S for λ_2 , i.e. $z_1^H S = \lambda_2 z_1^H$. Then

$$z_1^H \hat{S} = z_1^H (S + X R X^T J^T S) = z_1^H S = \lambda_2 z_1^H$$

as $z_1^H X = 0$ according to Lemma 2. Thus the left and right eigenvectors for λ_2 of S and \hat{S} coincide, which directly implies $\kappa(\hat{S}, \lambda_2) = \kappa(S, \lambda_2)$.

(ii) Now assume $R = R_1$ for R_1 in (23) and suppose $\nu = \mu$. The condition $\mu \notin \sigma(S)$ implies μ to be simple for \hat{S} . If $x_1, x_2 \in \mathbb{C}^{2n}$ are eigenvectors of S for λ_1 and λ_1^{-1} , respectively, then one directly obtains

$$\begin{aligned} \hat{S}x_1 &= Sx_1 + X R X^T J^T S x_1 = \lambda_1 x_1 - \lambda_1 X R X^T J x_1 = \lambda_1 x_1 - \lambda_1 X R \begin{bmatrix} 0 \\ -1 \end{bmatrix} \\ &= \lambda_1 (1 + r_{12}) x_1 = \lambda_1 \cdot (1 + \lambda_1^{-1} (\mu - \lambda_1)) x_1 = (\lambda_1 + \mu - \lambda_1) x_1 = \mu x_1 \end{aligned}$$

(similarly, $\hat{S}x_2 = \mu^{-1} x_2$ follows). Analogously we have $x_2^T J^T \hat{S} = \mu x_2^T J^T$, which follows from $x_2^T J^T S = \lambda_1 x_2^T J^T$ (see (18)). Thus, the left and right eigenvectors of S for λ_1 and those of \hat{S} for μ coincide and the statement follows. The proof is analogous for $\nu = \mu^{-1}$. \square

Remark 2. One can proceed as in the above proof to see that, if $R = R_2$ is used,

$$\kappa(\hat{S}, \mu) = \kappa(S, \lambda_1^{-1}) \quad \text{and} \quad \kappa(\hat{S}, \mu^{-1}) = \kappa(S, \lambda_1).$$

In fact, it is easy to show that now $\hat{S}x_1 = \mu^{-1} x_1$ and $\hat{S}x_2 = \mu x_2$ hold.

5. EXPERIMENTS

To perform numerical experiments in MATLAB R2021b, we used the code available in [8]. To obtain symplectic matrices $S \in \mathbf{M}_{200}(\mathbb{C})$, a symplectic matrix $S' \in \mathbf{M}_{200}(\mathbb{C})$ constructed from [8], was modified as

$$S = \begin{bmatrix} D_1^{-1} & \\ & D_1 \end{bmatrix} S' \begin{bmatrix} D_2 & \\ & D_2^{-1} \end{bmatrix}, \quad \text{where} \quad \begin{aligned} D_1 &= \text{diag}(\alpha_1, \dots, \alpha_{100}), \\ D_2 &= \text{diag}(\beta_1, \dots, \beta_{100}) \end{aligned} \quad (38)$$

and α_i, β_j are random numbers: `rand(1) + 1i*rand(1)`. Compared to S' , S has a more widespread spectrum ranging in magnitude from about 10^{-3} to 10^2 .




Figure 1: The plots show the effect of an eigenvalues modification when $\lambda_1 \in \sigma(S)$ undergoes a relative change of $|\lambda_1|$. Fifty experiments have been performed with $S \in \mathbf{M}_{200}(\mathbb{C})$. The plots show the use of $\hat{S}_1 = (I_{2n} + X R_1 X^T J^T)S$ for R_1 in (23), the coarse bound in (26) (left plot), the upper and lower bounds from Theorem 5 (right plot) and the relative change in the eigenvalue $|\lambda_1 - \mu|/|\lambda_1|$.

In Figure 1, 50 examples are shown where a randomly selected eigenvalue λ_1 of S has been changed to $\mu = \lambda_1(1 + \gamma z)$, where $z \in \mathbb{C}$ is a random complex number with $|z| = 1$ and $\gamma = |\lambda_1|$. Therefore, $|\mu - \lambda_1|/|\lambda_1| = |\lambda_1|$. The plot shows the relative change $\|S - \hat{S}_1\|_F/\|S\|_F$, the coarse bound in (26), the upper and lower bounds from Theorem 5 and the relative change in the eigenvalue $|\lambda_1 - \mu|/|\lambda_1|$. The plots show clearly that the bounds from Theorem 5 are significantly sharper than the bound from (26). If R_2 is used instead of R_1 the plots do not alter essentially.

The most significant difference between using \hat{S}_1 and \hat{S}_2 can be seen when an eigenvalue is subject to a small or large relative change compared to its absolute value. This is shown in Figure 2 where the experimental set-up is the same as before but now $\gamma = 10^{-3}|\lambda_1|$ (left plot) and $\gamma = 10^3|\lambda_1|$ (right plot) are chosen. Both plots show the relative change $\|S - \hat{S}_j\|_F/\|S\|_F$ for $j = 1, 2$ and the relative change in the eigenvalue $|\lambda_1 - \mu|/|\lambda_1|$. It is seen that for a small change in the eigenvalue λ_1 , the relative change $\|S - \hat{S}_1\|_F/\|S\|_F$ is significantly smaller than $\|S - \hat{S}_2\|_F/\|S\|_F$. However, when λ_1 undergoes a large change, then there is no big difference in using R_1 or R_2 .

In Figure 3 we show the use of matrices R different from R_1 and R_2 in (23). To this end, we fix a symplectic matrix $S \in \mathbf{M}_{200}(\mathbb{C})$ and an eigenvalue λ_1 of S that is to be modified by a relative change of $|\lambda_1|$. We consider a mesh grid on $[-1, 1] \times [-1, 1]$ with 150 discretization points in each direction. Each point (x_j, y_k) is associated with the complex number $c_{jk} := x_j + iy_k$ from which we made up $r_{11} = r_{22} = \sqrt{c_{jk}}$. The values for r_{12} and r_{21} are found according to (20) and (21) so that we obtain two different matrices $\hat{R}_1 = R(\eta_1, r_{11}, r_{22})$




Figure 2: The plots show the effect of an eigenvalues modification when $\lambda_1 \in \sigma(S)$ undergoes a relative change of $0.001 \cdot |\lambda_1|$ (left plot) and $1000 \cdot |\lambda_1|$ (right plot). Fifty experiments have been performed with $S \in \mathbf{M}_{200}(\mathbb{C})$. The plots show the use of \hat{S}_1 and \hat{S}_2 and the relative change in the eigenvalue $|\lambda_1 - \mu|/|\lambda_1|$.

and $\tilde{R}_2 = R(\eta_2, r_{11}, r_{22})$. The matrices $\tilde{S}_j = (I_{2n} + X\tilde{R}_jX^TJ^T)S$, $j = 1, 2$, where constructed and in Figure 3 the minimum of $\|S - \tilde{S}_j\|_F / \|S\|_F$ is shown for each point $c_{jk} \in [-1, 1] \times [-1, 1]$. The plot indicates that the minimum numerically found among all values $\|S - \tilde{S}_j\|_F / \|S\|_F$ is attained for $c_{jk} = 0$, i.e. $r_{11} = r_{22} = 0$. Thus, the minimum is obtained for one of the matrices R in (23). In most examples that have been considered a plot similar to the one in Figure 3 arose. However, seldom the numerical minimum was detected somewhere near $c_{jk} = 0$.

6. SYMPLECTIC MATRIX PENCILS

In this section we analyse the eigenvalue modification problem of Section 1 for symplectic matrix pencils. We call a matrix pencil $P(\lambda) = A - \lambda B$, $A, B \in \mathbf{M}_{2n}(\mathbb{C})$ symplectic (see [6, Sec. 2.1.1]) if it holds that

$$AJA^T = BJB^T. \quad (39)$$

From (39) it follows that for a symplectic pencil $P(\lambda) = A - \lambda B$ either A and B are both regular or singular, cf. [6, Sec. 2]. A scalar $\nu \in \mathbb{C}$ is called an eigenvalue of $P(\lambda)$ if there exists some nonzero $x \in \mathbb{C}^{2n}$ with $P(\nu)x = 0$, i.e. $Ax = \nu Bx$ [2, Sec. 2]. Eigenvalues of symplectic pencils also arise in pairs (ν, ν^{-1}) [6]. In particular, if A and B are singular, it is also possible to have $\nu = 0$ as an eigenvalue of $P(\lambda)$ which implies that $\lambda^{-1} = \infty$ is also an eigenvalue of $P(\lambda)$ (see, e.g., [5] for more information on the finite and infinite eigenstructure of matrix pencils). In the following, we focus on symplectic




Figure 3: The plot shows the minimum of $\|S - \tilde{S}_j\|_F / \|S\|_F$ for $j = 1, 2$ on the square $[-1, 1] \times [-1, 1]$ with 150 discretization points in each direction. The minimum on the grid is attained for $c_{jk} = 0$, i.e. $r_{11} = r_{22} = 0$. For random experiments the plots always look similar to the plot shown above. Notice that the surface has a sharp edge which arises due to the complex square root that has to be computed.

pencils $P(\lambda) = A - \lambda B$ where both A and B are regular. Thus, $P(\lambda)$ has neither the eigenvalue zero nor the eigenvalue infinity.

In general, for an arbitrary matrix pencil $P(\lambda) = A - \lambda B$, $A, B \in \mathbf{M}_n(\mathbb{C})$, where B is nonsingular, we can easily derive an adapted version of Rado's theorem.

Theorem 10. *Let $P(\lambda) = A - \lambda B$, $A, B \in \mathbf{M}_n(\mathbb{C})$ and B nonsingular, be a matrix pencil with eigenvalues $\lambda_1, \dots, \lambda_n \in \mathbb{C}$. Let $x_1, \dots, x_k \in \mathbb{C}^n$ be eigenvectors for $\lambda_1, \dots, \lambda_k$ such that $\text{rank}(X) = k$ for $X = [x_1 \ x_2 \ \dots \ x_k] \in \mathbf{M}_{n \times k}(\mathbb{C})$. Furthermore, let $C \in \mathbf{M}_{n \times k}(\mathbb{C})$ be arbitrary. Then the matrix pencil*

$$\tilde{P}(\lambda) = (A + BXC^T) - \lambda B$$

has the eigenvalues $\mu_1, \dots, \mu_k, \lambda_{k+1}, \dots, \lambda_n$, where μ_1, \dots, μ_k are the eigenvalues of the matrix $\Lambda + C^T X$ with $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_k)$.

Proof. As $B^{-1}P(\lambda) = B^{-1}A - \lambda I_n$, the eigenvalues of $P(\lambda)$ coincide with those of the matrix $M := B^{-1}A \in \mathbf{M}_{2n}(\mathbb{C})$. In particular, $P(\lambda)$ does not have $\lambda = \infty$ as an eigenvalue. Assume that $AX = BXC\Lambda$ with $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_k)$ we obtain $B^{-1}AX = X\Lambda$ and Theorem 2 implies that $\hat{M} := B^{-1}A + XC^T$ has the eigenvalues $\mu_1, \dots, \mu_k, \lambda_{k+1}, \dots, \lambda_n$, where μ_1, \dots, μ_k are the eigenvalues of the matrix $\Lambda + C^T X$. As \hat{M} and the matrix pencil $\hat{P}(\lambda) = (A + BXC^T) - \lambda B$ have the same eigenvalues, the statement follows. \square

Now let $P(\lambda) = A - \lambda B$ be a symplectic matrix pencil according to (39) with nonsingular A, B and eigenvalues $\lambda_1, \lambda_1^{-1}, \dots, \lambda_n, \lambda_n^{-1}$. We set

$$\hat{P}(\lambda) := (A + BXC^T) - \lambda B$$

and intend to determine $C \in \mathbf{M}_{2n \times 2}(\mathbb{C})$ such that $\hat{P}(\lambda)$ is again symplectic and has the eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ for a given value $\mu \in \mathbb{C} \setminus \{0\}$. A direct calculation reveals that (39) is equivalent to $B^{-1}A$ being a symplectic matrix, i.e. $J^T(B^{-1}A)^T J = (B^{-1}A)^{-1}$. Thus, Theorem 3 can be applied to the matrix $S := B^{-1}A$ that has the same eigenvalues as $P(\lambda)$. Now suppose

$$AX = BX \begin{bmatrix} \lambda_1 & \\ & \lambda_1^{-1} \end{bmatrix}, \quad X = [x_1 \quad x_2] \in \mathbf{M}_{2n \times 2}(\mathbb{C}),$$

i.e. $x_1, x_2 \in \mathbb{C}^{2n}$ are generalized eigenvectors for λ_1 and λ_1^{-1} , respectively. Furthermore, assume $x_1^T J x_2 \neq 0$. Then

$$\hat{S} := B^{-1}A + XRX^T J^T B^{-1}A$$

is symplectic with eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ provided that R is chosen according to the conditions in Theorem 3 for μ . Then, the matrix pencil $\hat{P}(\lambda) := B(\hat{S} - \lambda I_{2n})$, i.e.

$$\hat{P}(\lambda) = (A + BXR X^T J^T B^{-1}A) - \lambda B, \quad (40)$$

has the same eigenvalues as \hat{S} [5, Sec. 3.1]. In fact, $\hat{P}(\lambda)$ is again a symplectic pencil. To show this, we have to check that

$$(A + BXR X^T J^T B^{-1}A)J(A + BXR X^T J^T B^{-1}A)^T = BJB^T$$

holds. To this end, it only remains to prove that

$$\begin{aligned} AJ(BXR X^T J^T B^{-1}A)^T + (BXR X^T J^T B^{-1}A)JA^T \\ + (BXR X^T J^T B^{-1}A)J(BXR X^T J^T B^{-1}A)^T = 0 \end{aligned} \quad (41)$$

since $AJA^T = BJB^T$ holds by assumption. Using this relation, (41) simplifies to

$$-BXR^T X^T B^T + BXR X^T B^T + BXR J_2 R^T X^T B^T = 0$$

which can be rewritten as $BX(R - R^T + R J_2 R^T)X^T B^T = 0$. As in Section 2 this relation holds if and only if $R - R^T + R J_2 R^T = 0$. As for any $R \in \mathbf{M}_2(\mathbb{C})$ we have $R J_2 R^T = R^T J_2 R$, it follows that the condition $R - R^T + R J_2 R^T = 0$ is equivalent to (6). Therefore, since R was constructed according to (13) so that $R - R^T + R^T J_2 R = 0$ holds, this finally shows that (41) is true, so $\hat{P}(\lambda)$ is symplectic. As already mentioned above, $\hat{P}(\lambda)$ and \hat{S} have the same eigenvalues, so the eigenvalues of $\hat{P}(\lambda)$ are $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$.

Notice that $\hat{P}(\lambda)$ in (40) can be rewritten in various ways, e.g.

$$\hat{P}(\lambda) = (A + BXR X^T B^T A^{-T} J^T) - \lambda B \quad (42)$$

$$= (A + BXR X^T B^T J^T A) - \lambda B \quad (43)$$

using the relation $J^T B^{-1} A = B^T A^{-T} J^T$ that follows from $B^{-1} A$ being symplectic in (42) and $A^{-T} = J^T A J$ in (43). Using $X^T B^T = \Lambda^{-1} X^T A^T$ and exchanging $X^T B^T$ with $\Lambda^{-1} X^T A^T$ in (42) yields

$$\begin{aligned} \hat{P}(\lambda) &= (A + BXR \Lambda^{-1} X^T A^T A^{-T} J^T) - \lambda B \\ &= \left(A + BXR \begin{bmatrix} \lambda_1^{-1} & \\ & \lambda_1 \end{bmatrix} X^T J^T \right) - \lambda B. \end{aligned} \quad (44)$$

which is an expression for $\hat{P}(\lambda)$ similar to (14). We conclude our finding in the following theorem.

Theorem 11. *Let $P(\lambda) = A - \lambda B$, $A, B \in \mathbf{M}_{2n}(\mathbb{C})$ nonsingular, be a symplectic pencil with eigenvalues $\lambda_1, \lambda_1^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1} \in \mathbb{C}$ and let $\mu \in \mathbb{C} \setminus \{0\}$ be given. Let $x_1, x_2 \in \mathbb{C}^{2n}$ be eigenvectors for λ_1 and λ_1^{-1} , respectively, normalized such that $X^T J_{2n} X = J_2$ for $X = [x_1 \ x_2] \in \mathbf{M}_{2n}(\mathbb{C})$ and set $d := (\mu + \mu^{-1}) - (\lambda_1 + \lambda_1^{-1})$. Then the matrix pencil*

$$\hat{P}(\lambda) := \left(A + BXR \begin{bmatrix} \lambda_1^{-1} & \\ & \lambda_1 \end{bmatrix} X^T J \right) - \lambda B \quad (45)$$

is again symplectic and has the eigenvalues $\mu, \mu^{-1}, \lambda_2, \lambda_2^{-1}, \dots, \lambda_n, \lambda_n^{-1}$ provided that $R = [r_{ij}]_{ij} \in \mathbf{M}_2(\mathbb{C})$ is chosen such that (12) and (13) hold.

Regarding $\hat{P}(\lambda)$ in (40), let $\hat{P}(\lambda) = \hat{A} - \lambda \hat{B}$ with $\hat{A} = A + BXR X^T J^T B^{-1} A$ and $\hat{B} = B$. Then certainly $\|\hat{B} - B\|/\|B\| = 0$ while

$$\frac{\|\hat{A} - A\|}{\|A\|} \leq \kappa(B) \|R\| \|X\|^2,$$

where $\kappa(B) = \|B\| \|B^{-1}\|$ is the condition number of the matrix B . Thus, using Theorem 5 we may immediately bound $\|\hat{A} - A\|/\|A\|$. Furthermore, from the form of $\hat{P}(\lambda)$ in (45), statements similar to those in Section 4.2 can directly be derived.

7. SUMMARY

In this work we showed how to modify a pair of eigenvalues $\lambda, 1/\lambda$ of a symplectic matrix S to desired target values $\mu, 1/\mu$ for a symplectic matrix \hat{S} in a structure-preserving way. Universal bounds on the relative distance between S and \hat{S} with modified spectrum were given. The eigenvalues Segre characteristics of S were related to those of \hat{S} and some statements on eigenvalue condition numbers have been derived. The main results have been extended to matrix pencils.

8. ACKNOWLEDGEMENT

The author is grateful to Thomas Richter as this work was in parts developed during the authors employment in Thomas Richter's group at the Otto-von-Guericke-Universität Magdeburg.

REFERENCES

- [1] A. T. Alexandridis, H. E. Psillakis. The inverse optimal LQR problem and its relation to passivity and eigenvalue assignment. *International Journal of Tomography & Statistics*, Vol. 5, 2007.
- [2] T. Betcke, N. J. Higham, V. Mehrmann, C. Schröder, F. Tisseur. NLEVP: A Collection of Nonlinear Eigenvalue Problems. *ACM Transactions on Mathematical Software*, Vol. 39(2), 2013.
- [3] R. Bru, R. Canto, R. L. Soto and A. M. Urbano *A Brauer's theorem and related Results*. Cent. Eur. J. Math. 10(1), 2012.
- [4] D. F. Delchamps. *State Space and Input-Output Linear Systems*. Springer Verlag, New York, USA, 1988.
- [5] F. De Terán, F. M. Dopico, D. S. Mackey. Spectral equivalence of matrix polynomials and the Index Sum Theorem. *Linear Algebra Appl.*, Vol. 459, 2014.
- [6] H. Fassbender. *Symplectic Methods for the Symplectic Eigenproblem*. Kluwer Academic Publishers, New York, USA, 2002.
- [7] R. A. Horn, C. R. Johnson. *Matrix Analysis (Second Edition)*. Cambridge University Press, New York, USA, 2013.
- [8] D. P. Jagger. MATLAB Toolbox for Classical Matrix Groups. Masters thesis, University of Manchester, 2003. MIMS EPrint 2007.99.
- [9] C. R. Johnson, E. A. Schreiner. The Relationship between AB and BA . *The American Mathematical Monthly*, Vol. 103(7), 1996.
- [10] A. J. Laub, K. Meyer. Canonical Forms for Symplectic and Hamiltonian Matrices. *Celestial Mechanics*, Vol. 9, 1974.
- [11] D. G. Luenberger. *Optimization by Vector Space Methods*. John Wiley & Sons, Inc., New York, USA, 1969.
- [12] T. Lyche. *Numerical Linear Algebra and Matrix Factorizations*. Springer Nature, Cham, Switzerland, 2020.
- [13] H. Perfect. *Methods of constructing certain stochastic matrices. II* Duke Math. J. 22(2), 1955.
- [14] Y. Saad. *Numerical Methods for Large Eigenvalue Problems (Revised Edition)*. Society for Industrial and Applied Mathematics, Philadelphia, USA, 2011.

- [15] H. Shapiro. *Linear Algebra and Matrices. Topics for a Second Course.* American Mathematical Society, Providence, USA, 2015.
- [16] R. L. Soto and O. Rojo. Applications of a Brauer theorem in the nonnegative inverse eigenvalue problem. *Linear Algebra Appl.* 416(2-3), 2006.
- [17] O. Taussky, H. Zassenhaus. On the similarity transformation between a matrix and its transpose. *Pacific J. Math.* 9(3), 1959.