

FINITE SETS CONTAINING ZERO ARE MAPPING DEGREE SETS

CRISTINA COSTOYA, VICENTE MUÑOZ, AND ANTONIO VIRUEL

ABSTRACT. In this paper we solve in the positive the question of whether any finite set of integers A , containing 0, is the mapping degree set between two oriented closed connected manifolds of the same dimension. We extend this question to the rational setting, where an affirmative answer is also given.

1. INTRODUCTION

In this paper, we settle in the positive various questions which have been raised about $D(M, N)$, the set of mapping degrees between two oriented closed connected manifolds M and N of the same dimension:

$$D(M, N) = \{d \in \mathbb{Z} \mid \exists f : M \rightarrow N, \deg(f) = d\}.$$

C. Neofytidis, S. Wang, and Z. Wang [22, Problem 1.1] discuss the problem of finding, for any set $A \subset \mathbb{Z}$ containing 0, two oriented closed connected manifolds M and N of the same dimension such that $A = D(M, N)$. Note that $0 \in A$ is a necessary condition as the constant map $M \rightarrow N$ is of degree zero.

A cardinality argument shows that when A is an infinite set, the problem above is solved in the negative [22, Theorem 1.3]. Indeed, there are uncountably many infinite subsets of \mathbb{Z} containing 0, compared to the countably many mapping degree sets that exist for pairs of oriented closed connected manifolds with the same dimension. Hence, *most* of the infinite sets are not realizable as mapping degree sets. Now, using a computability argument, C. Löh and M. Uschold [18, Proposition A.1] prove that $D(M, N)$ is a recursively enumerable set. This provides with a sufficient condition for infinite sets of integers (containing 0) not being realizable as mapping degree sets (see [18, Example A.2]).

Thus, one might ask:

Question 1.1 ([22, Problem 1.4]). *Let A be a finite set of integers containing 0. Is $A = D(M, N)$ for some oriented closed manifolds M, N of the same dimension?*

Remark 1.2. It is important to notice that if $\{0\} \subsetneq A = D(M, N)$, A finite, for some manifolds M and N , then $D(M, M)$ and $D(N, N)$ must both be contained in $\{0, 1, -1\}$. Otherwise, if there exists $g : M \rightarrow M$ with $|\deg(g)| > 1$, then for any non-zero degree $f : M \rightarrow N$ (which exists by assumption), the subset $\{\deg(f \circ g^m) \mid m \in \mathbb{N}\}$ of $D(M, N)$ is unbounded. This leads to a contradiction as $A = D(M, N)$ is finite. The same follows for $D(N, N)$.

2020 *Mathematics Subject Classification.* 55M25, 57N65, 55P62, 55R10.

Key words and phrases. Mapping degree sets, inflexible manifold, fiber bundle, unstable Adams operation.

An oriented closed manifold M satisfying $D(M, M) \subset \{0, 1, -1\}$ is called an *inflexible* manifold [9, Definition 1.4]. This condition is equivalent to asking that $D(M, M)$ is bounded: since it is a multiplicative semi-group, if there exists any $\ell \in D(M, M)$ with $|\ell| > 1$, then $D(M, M)$ is unbounded. Simply connected inflexible manifolds are rare objects that have appeared quite recently in literature using rational homotopy theory and surgery theory (see [7] for an account on the simply connected inflexible manifolds that are known at present). Not surprisingly, and in lights of Remark 1.2, part of our key constructions will use rational homotopy methods.

The main result in this work answers Question 1.1 positively:

Theorem A. *Let A be a finite set of integers containing 0. Then, $A = D(M, N)$ for some oriented closed connected 3-manifolds M, N .*

The proof of this theorem will be carried out at the end of Section 3. Appealing to [22, Example 1.5], we point out that the 3-dimension of the manifolds is the lowest possible.

A second problem related to Question 1.1 is also treated in this paper. More precisely, let the *rational mapping degree set* between oriented closed connected n -manifolds M, N be the following set

$$D_{\mathbb{Q}}(M, N) = \{d \in \mathbb{Q} \mid \exists f: (M_{(0)}, [M]_{\mathbb{Q}}) \rightarrow (N_{(0)}, [N]_{\mathbb{Q}}), \deg(f) = d\},$$

where $[M] \in H^n(M; \mathbb{Z})$ denotes the cohomological fundamental class of M , $[M]_{\mathbb{Q}} \in H^n(M; \mathbb{Q})$ denotes the rational cohomological fundamental class of M , and $M_{(0)}$ the rationalization of M . Then, we raise the following question, which can be thought of as a rational version of [22, Problem 1.4]:

Question 1.3. *Let A be a finite set of rational numbers containing 0. Is $A = D_{\mathbb{Q}}(M, N)$ for some oriented closed connected manifolds M, N of the same dimension?*

In Section 4 we solve this problem in the positive by proving:

Theorem B. *Let A be a finite set of rational numbers containing 0. Then $A = D_{\mathbb{Q}}(M, N)$ for some oriented closed manifolds M, N . Moreover, given any integer $k \geq 1$, the manifolds M, N above can be chosen $(30k + 17)$ -connected.*

The proofs of Theorem A and Theorem B consist of mainly two steps:

- *Arithmetical decomposition of finite sets:* In Section 2 we show how to decompose the candidate A to be realized as the mapping degree set of manifolds, as an intersection of sums over specifically designed sequences of integers S_{B_i} , $i = 0, \dots, n$ (see Definition 2.1). Each of those sums gradually approaches A (Proposition 2.2, Corollary 2.3).
- *Spherical fibrations:* In Sections 3 and 4, we use certain inflexible manifolds (resp. inflexible Sullivan algebras) as the basis of spherical fibrations where the total spaces are also inflexible manifolds (resp. inflexible Sullivan algebras). Relations between connected sums and mapping degree sets (see Propositions 3.5, 4.3, and 5.8) allow one to consider iterated connected sums of the total spaces, in a first stage to realize the sums S_{B_i} above mentioned, and in a second stage to realize the candidate A .

Looking at the connectivity, while manifolds from Theorem B are simply connected (indeed, they are as highly connected as desired) the ones from Theorem A have non-trivial fundamental group. In Section 5 we will use unstable Adams operations to prove the following results that guarantee that manifolds realizing finite sets of integers can be chosen simply connected:

Theorem C. *Suppose that there exists an oriented closed k -connected $2m$ -manifold Σ , $m > 1$, verifying that $\Sigma_{(0)}$ is inflexible and $\pi_j(\Sigma_{(0)}) = 0$ for $j \geq 2m - 1$. Then any finite set of integers A containing 0 can be realized as $A = D(M, N)$ for some oriented closed k -connected $(4m - 1)$ -manifolds M, N .*

Examples of simply connected manifolds fulfilling the hypotheses of Theorem C can be found in [1, Example 3.8] and [9, Theorem 6.8, Theorem II.5] (see also [21, Theorem 1.4]). Hence, the following holds:

Corollary D. *Any finite set of integers A containing 0 can be realized as $A = D(M, N)$ for some oriented closed simply connected manifolds M, N of the same dimension.*

Also, we would like to point out that in every dimension $n > 6$, there exist infinitely many nilmanifolds Σ verifying that $\Sigma_{(0)}$ is inflexible, as it follows from [2], [4] and [14]. Hence, we have the following:

Corollary E. *For every $m > 3$, any finite set of integers A containing 0 can be realized as $A = D(M, N)$ for some oriented closed simply connected $(4m - 1)$ -manifolds M, N .*

Recall that an oriented closed connected manifold is said *strongly chiral* if it does not admit self-maps of degree -1 (see [20]). Also recall that a subset $A \subset \mathbb{Z}$ is said *symmetric* if $A = -A$ where $-A = \{-a \mid a \in A\}$. We finish the introduction by mentioning that our results provide us with a method for obtaining strongly chiral manifolds:

Remark 1.4. Let A be a finite set of integers strictly containing 0. Theorem A and Theorem C illustrate how to construct oriented closed connected manifolds M, N verifying that $A = D(M, N)$. Those manifolds need to be inflexible, as mentioned in Remark 1.2. Moreover, if we choose a non-symmetric set A , then M and N need also to be strongly-chiral manifolds. Otherwise, if $-1 \in D(M, M)$ (resp. $-1 \in D(N, N)$), since $D(M, M)$ (resp. $D(N, N)$) acts on $A = D(M, N)$ by multiplication on the left (resp. right), the set A would be symmetric, leading thus to a contradiction.

Funding. This work was partially supported by MCIN/AEI/10.13039/501100011033 [PID2020-115155GB-I00 to C.C., PID2020-118452GB-I00 to V.M., and PID2020-118753GB-I00 to A.V.]. The third author was also partially supported by PAIDI 2020 (Andalusia) grant PROYEXCEL-00827.

Acknowledgments. We would like to express our sincere gratitude to C. Neofytidis for pointing out an error in a previous version of this manuscript.

2. SOME ARITHMETIC COMBINATORICS

In this section we show that every finite set $A \subset \mathbb{Z}$ (resp. $\subset \mathbb{Q}$) containing 0 can be expressed as the intersection of sums over certain sequences of integers, that gradually

approach A. The sequences have an additional property (see Proposition 2.2) that will be crucial to prove Theorem C in Section 5 below.

Following the notation in [9, Section II.1], [22, Section 3], given $A, B \subset \mathbb{Z}$ (resp. $\subset \mathbb{Q}$) we write $A + B := \{a + b \mid a \in A, b \in B\} \subset \mathbb{Z}$ (resp. $\subset \mathbb{Q}$).

Definition 2.1. *Let B be a finite sequence of not necessarily pairwise distinct non-zero integers (resp. rational numbers). We write*

$$S_B := \sum_{b \in B} \{0, b\} \subset \mathbb{Z} \text{ (resp. } \subset \mathbb{Q}\text{),}$$

and we refer to it as the sum over the sequence B .

Proposition 2.2. *Let d_1, \dots, d_n be pairwise distinct non-zero integers. For any positive integer $m \geq 1$, there exist finite sequences $B(i)$, $i = 0, \dots, n$, of not necessarily pairwise distinct non-zero integers, such that*

$$\{0, d_1, \dots, d_n\} = \bigcap_{i=0}^n S_{B(i)}.$$

Moreover, every element in $B(i)$ can be written as a power $\pm k^m$ for some positive integer k coprime to $m!$.

Proof. Fix $m \geq 1$. Since the construction of $B(i)$, $i = 0, \dots, n$, depends on the sign of the pairwise distinct $d_i \in \mathbb{Z}$, $i = 1, \dots, n$, we write them as an ordered sequence

$$\{-a_r < \dots < -a_1 < 0 < e_1 < \dots < e_s\}$$

where $n = r + s$. We assume $a_0 = 0 = e_0$.

In the first place, let $B(0)$ be the sequence consisting of a_r copies of $-1 = -1^m$ and e_s copies of $1 = 1^m$. Thus

$$\{-a_r < \dots < e_s\} \subset S_{B(0)} = [-a_r, e_s] \cap \mathbb{Z}.$$

In the second place, for $j = 1, \dots, s$, choose a positive $k_j \in \mathbb{Z}$ coprime with $m!$ such that $k_j^m > \max\{e_s, e_j + a_r\}$. Then, let $B(j)$ be the sequence consisting of $k_j^m - e_j$ copies of $-1 = -1^m$, e_{j-1} copies of $1 = 1^m$, and one copy of k_j^m . Hence,

$$\{-a_r < \dots < e_s\} \subset S_{B(j)} = ([-(k_j^m - e_j), e_{j-1}] \cup [e_j, k_j^m + e_{j-1}]) \cap \mathbb{Z}.$$

Finally, for $j = s + 1, \dots, n$, choose a positive $k_j \in \mathbb{Z}$ coprime with $m!$ such that $k_j^m > \max\{a_r, a_{j-s} + e_s\}$. Then, let $B(j)$ be the sequence consisting of $k_j^m - a_{j-s}$ copies of $1 = 1^m$, a_{j-s-1} copies of $-1 = -1^m$, and one copy of $-k_j^m$. Hence,

$$\{-a_r < \dots < e_s\} \subset S_{B(j)} = ([-k_j^m - a_{j-s-1}, -a_{j-s}] \cup [-a_{j-s-1}, k_j^m - a_{j-s-1}]) \cap \mathbb{Z}.$$

All of the above implies that

$$\{0, d_1, \dots, d_n\} = \{-a_r < \dots < e_s\} = \bigcap_{i=0}^n S_{B(i)}.$$

□

For $A \subset \mathbb{Q}$ and $\lambda \in \mathbb{Q}$ we write

$$\lambda A := \{\lambda a \mid a \in A\}.$$

Notice that if $B(i)$ is a finite sequence of not necessarily pairwise distinct non-zero rational numbers, $i = 0, \dots, n$, for any $\lambda \in \mathbb{Q}$, we have that

$$\lambda \left(\bigcap_{i=0}^n S_{B(i)} \right) = \bigcap_{i=0}^n S_{\lambda B(i)}.$$

Therefore, the following is a direct consequence of Proposition 2.2:

Corollary 2.3. *Let d_1, d_2, \dots, d_n be pairwise distinct non-zero rational numbers. Then, there exist finite sequences $B(i)$, $i = 0, \dots, n$, of not necessarily pairwise distinct non-zero rational numbers, such that*

$$\{0, d_1, \dots, d_n\} = \bigcap_{i=0}^n S_{B(i)}.$$

3. CIRCLE BUNDLES OVER INFLEXIBLE 2-MANIFOLDS: MAPPING DEGREE SET

This section is devoted to prove Theorem A. As explained in the introduction (see Remark 1.2) if we want to realize a finite set of integers, strictly containing 0, as a mapping degree set $D(M, N)$, then both M and N need to be inflexible manifolds. We are going to consider circle bundles over certain inflexible 2-manifolds, with prescribed Euler class, whose total space is again an inflexible 3-manifold. These 3-manifolds will be used as building blocks to construct, by means of iterated connected sums, manifolds M and N .

We first collect from literature a couple of results that are needed:

Lemma 3.1 ([7, Lemma 7.8], [22, Lemma 3.5]). *Let M_1, M_2 and N be oriented closed connected n -manifolds. Then*

$$D(M_1, N) + D(M_2, N) \subset D(M_1 \# M_2, N)$$

Moreover, if $\pi_{n-1}(N) = 0$, then

$$D(M_1, N) + D(M_2, N) = D(M_1 \# M_2, N).$$

We reformulate [22, Lemma 4.3] as follows:

Lemma 3.2. *Let M and N_1, N_2 be oriented closed manifolds of the same dimension. Then*

$$D(M, N_1 \# N_2) \subset D(M, N_1) \cap D(M, N_2).$$

Whenever we need to enlarge the fundamental group of our manifolds to construct mapping degree sets we shall use handle-bodies:

Definition 3.3. *Given integers $n, k > 0$, we denote by $H(n, k)$ the oriented closed n -manifold arising from the k -fold connected sum of oriented handle bodies $S^{n-1} \times S^1$, that is*

$$H(n, k) := \#_{i=1}^k (S^{n-1} \times S^1).$$

Lemma 3.4. *Let $n > 2$ and N be an oriented closed n -manifold such that $\pi_{n-1}(N) = 0$. Then $D(H(n, k), N) = \{0\}$, for every integer $k > 0$.*

Proof. Given a space X , and a positive integer m , let $X^{(m)}$ and $X\langle m \rangle$ respectively denote the m th Postnikov stage and the m -connected cover of X .

Since Postnikov stages and connected covers can be constructed in a functorial way [10, Example 1.A.1.1], given $f: H(n, 1) \rightarrow N$ we can take the $(n-2)$ th Postnikov stage and the $(n-2)$ -connected cover to obtain the following commutative diagram

$$\begin{array}{ccccc} S^{n-1} & \longrightarrow & H(n, 1) & \longrightarrow & S^1 \\ \downarrow f^{(n-2)} & & \downarrow f & & \downarrow f^{(n-2)} \\ N\langle n-2 \rangle & \longrightarrow & N & \longrightarrow & N^{(n-2)}. \end{array} \quad (1)$$

We now compare the homological Serre spectral sequences (Sss) associated to the two fiber sequences above. On the one hand, the Sss associated to the top-row fiber sequence in Equation (1) collapses at the $E_{*,*}^2$ -page and the fundamental class $[H(n, 1)] \in H_n(H(n, 1))$ is represented by $[S^{n-1}] \otimes [S^1]$. Therefore, using the edge morphisms, we get that the class $H_n(f)([H(n, 1)]) \in H_n(N)$ is represented by $H_{n-1}(f\langle n-2 \rangle)([S^{n-1}]) \otimes H_1(f^{(n-2)})([S^1])$ in the Sss associated to the bottom-row fiber sequence in Equation (1).

On the other hand, since $\pi_{n-1}(N) = 0$, then $N\langle n-2 \rangle$ is indeed $(n-1)$ -connected and therefore $H_{n-1}(N\langle n-2 \rangle) = 0$. Thus $H_{n-1}(f\langle n-2 \rangle)$ is the trivial morphism, and the class $H_{n-1}(f\langle n-2 \rangle)([S^{n-1}]) \otimes H_1(f^{(n-2)})([S^1])$ is trivial. In other words $D(H(n, 1), N) = \{0\}$.

Finally, for the general case, we apply Lemma 3.1, to $H(n, k) = \#_{i=1}^k H(n, 1)$ and N to get that

$$D(H(n, k), N) = \sum_{i=1}^k D(H(n, 1), N) = \{0\}.$$

□

Using the lemmas above, we prove the following result:

Proposition 3.5. *Let M_1, M_2 and N_1, N_2 be oriented closed n -manifolds, $n > 2$, verifying:*

- (1) $\pi_{n-1}(N_i) = 0$ for $i = 1, 2$;
- (2) $D(M_i, N_j) = \{0\}$ for $i \neq j$;
- (3) $D(M_1, N_1) \cap D(M_2, N_2)$ is finite.

Then there exists an integer $\ell \geq 0$ such that

$$D(M_1 \# M_2 \# H(n, \ell), N_1 \# N_2) = D(M_1, N_1) \cap D(M_2, N_2).$$

Proof. Let us define the integers

$$\{0, d_1, \dots, d_r\} := D(M_1, N_1) \cap D(M_2, N_2).$$

Since every continuous $f: M_i \rightarrow N_i$ is homotopic to a smooth map, for each

$$d_j \in D(M_1, N_1) \cap D(M_2, N_2) \subset D(M_i, N_i),$$

we can choose $f_{i,j}: M_i \rightarrow N_i$ such that $f_{i,j}$ is smooth, and $\deg(f_{i,j}) = d_j$. Then by Sard's Theorem [24], the set of singular values of a smooth map has measure zero, so we can choose a regular value $y_{i,j} \in N_i$ for $f_{i,j}$ such that $y_{i,j} \neq y_{i,j'}$, and $f_{i,j}^{-1}(y_{i,j}) \cap f_{i,j'}^{-1}(y_{i,j'}) = \emptyset$, for $j \neq j'$. Let us set

$$f_{i,j}^{-1}(y_{i,j}) := \{x_{i,j,1}, \dots, x_{i,j,s_{i,j}}\},$$

where $s_{i,j} \geq |d_j|$.

Now, according to the Stack of Records Theorem [23, Theorem 9.1], [12, p. 26], there exists an n -dimensional open disc $U_{i,j} \subset N_i$ such that $y_{i,j} \in U_{i,j}$ and

$$f_{i,j}^{-1}(U_{i,j}) = \bigcup_{k=1}^{s_{i,j}} V_{i,j,k}$$

where $x_{i,j,k} \in V_{i,j,k} \subset M_i$, $V_{i,j,k} \cap V_{i,j,k'} = \emptyset$ for $k \neq k'$, and $f_{i,j}|_{V_{i,j,k}}: V_{i,j,k} \rightarrow U_{i,j}$ is a diffeomorphism. Therefore $V_{i,j,k} \subset M_i$ is an open disc, $\deg(f_{i,j}|_{V_{i,j,k}}) = \pm 1$, and

$$\deg(f_{i,j}) = \sum_{k=1}^{s_{i,j}} \deg(f_{i,j}|_{V_{i,j,k}}).$$

We may assume that $\deg(f_{i,j}|_{V_{i,j,k}}) = \text{sign}(\deg(f_{i,j}))$, for $k = 1, \dots, |d_j|$ while the following $s_{i,j} - |d_j|$ diffeomorphisms, which are on even number, have alternating signs.

Since we are dealing with compact Hausdorff spaces, we can shrink all the discs and assume that the closures $\overline{V_{i,j,k}} \subset M_i$ are pairwise disjoint, and $f_{i,j}|_{\overline{V_{i,j,k}}}: \overline{V_{i,j,k}} \rightarrow \overline{U_{i,j}}$ are diffeomorphisms.

Let M be the n -manifold obtained by removing discs and adding handles $S^{n-1} \times [0, 1]$ to the disjoint union of M_1 and M_2 as follows:

- (1) First, we remove all discs $V_{i,j,k} \subset M_i$, $i = 1, 2$, from the disjoint union of M_1 and M_2 so we obtain K_0 , a disconnected n -manifold with boundary;
- (2) Then, for each $k = 1, \dots, |d_j|$, we add to K_0 a handle, that we label $(d_j, k, 0)$, by identifying $S^{n-1} \times \{0\}$ with $\partial(V_{1,j,k})$ and $S^{n-1} \times \{1\}$ with $\partial(V_{2,j,k})$, thus we obtain K_1 , a connected n -manifold with boundary when $s_{i,j} > |d_j|$;
- (3) Finally, for each $i = 1, 2$, $j = 1, \dots, r$, if $s_{i,j} > |d_j|$, for $m = 1, \dots, \frac{s_{i,j} - |d_j|}{2}$, we add to K_1 a handle, that we label $(d_j, m, 1)$, identifying $S^{n-1} \times \{0\}$ with $\partial(V_{i,j,|d_j|+2m-1})$ and $S^{n-1} \times \{1\}$ with $\partial(V_{i,j,|d_j|+2m})$, so obtaining M .

Observe that $M = M_1 \# M_2 \# H(n, \ell)$ for some $\ell \geq 0$.

We now show that

$$D(M, N_1 \# N_2) = \{0, d_1, \dots, d_r\} = D(M_1, N_1) \cap D(M_2, N_2).$$

Given any d_j , for $j = 1, \dots, r$, we construct a map $F_j: M \rightarrow N_1 \# N_2$ of degree d_j as follows:

- (1) $M \rightarrow \widetilde{M}$ is obtained by applying pinching-stretching-collapsing (see Figure 1) to all the handles in M with labels $(d_{j'}, k, l)$ for $j' \neq j$, so \widetilde{M} can be identified with the manifold constructed from the disjoint union of M_1 and M_2 where the discs $V_{i,j,k} \subset M_i$, $i = 1, 2$, have been removed and handles with labels $(d_j, k, 0)$ and $(d_j, m, 1)$ have been added as above.




FIGURE 1. A graphical description of the pinching-stretching-collapsing procedure

- (2) Think of $N_1 \# N_2$ as the manifold obtained by removing the discs $U_{i,j} \subset N_i$, $i = 1, 2$ and identifying $\partial(U_{1,j})$ with $\partial(U_{2,j})$.
- (3) Then, $\widetilde{M} \rightarrow N_1 \# N_2$ is obtained by contracting all existing handles to $\partial(U_{1,j}) \equiv \partial(U_{2,j})$ while points in M_i are mapped by f_i for $i = 1, 2$.

Let F_j be the composition of the maps $M \rightarrow \widetilde{M}$ and $\widetilde{M} \rightarrow N_1 \# N_2$ constructed above. Then, $\deg(F_j) = d_j$ and therefore $D(M_1, N_1) \cap D(M_2, N_2) \subset D(M, N_1 \# N_2)$.

Conversely, to show that $D(M, N_1 \# N_2) \subset D(M_1, N_1) \cap D(M_2, N_2)$, we first observe that according to Lemma 3.1:

$$D(M, N_i) = D(M_1 \# M_2 \# H(n, \ell), N_i) = D(M_1, N_i) + D(M_2, N_i) + D(H(n, \ell), N_i).$$

Thus by hypothesis and Lemma 3.4, we obtain $D(M, N_i) = D(M_i, N_i)$. Therefore, applying Lemma 3.2 we obtain:

$$D(M, N_1 \# N_2) \subset D(M, N_1) \cap D(M, N_2) = D(M_1, N_1) \cap D(M_2, N_2).$$

□

Although we will not give the details, Lemma 3.1 and Lemma 3.2 can be easily generalized to finite iterated connected sums. Hence, following along the lines of the proof in Proposition 3.5 we obtain:

Corollary 3.6. *Let M_i, N_i , $i = 1, \dots, r$, be oriented closed connected n -manifolds, $n > 2$, verifying:*

- (1) $\pi_{n-1}(N_i) = 0$ for $i = 1, \dots, r$;
- (2) $D(M_i, N_j) = \{0\}$ for $i \neq j$;
- (3) $D(M_i, N_i) \cap D(M_j, N_j)$ is finite for $i \neq j$.

Then there exists an integer $\ell \geq 0$ such that

$$D(M_1 \# \dots \# M_r \# H(n, \ell), N_1 \# \dots \# N_r) = \bigcap_{i=1}^r D(M_i, N_i).$$

We now have all the ingredients to prove our main theorem.

Proof of Theorem A. Let $A = \{0, d_1, \dots, d_n\}$ be a finite set of pairwise distinct integers. We need to show that A is realized by two oriented closed 3-manifolds M, N in the sense that $A = D(M, N)$.

For this purpose, we consider an oriented closed hyperbolic surface of genus $g > 1$, Σ_g . Then, for any $i \in \mathbb{Z}$, let K_i be the total space in the circle bundle

$$S^1 \rightarrow K_i \rightarrow \Sigma_g$$

with Euler number $e(K_i) = i$. Observe that $K_i, i \in \mathbb{Z}$, is an aspherical 3-manifold. The mapping degree set between these 3-manifolds is fully described in [22, Lemma 3.4]:

$$D(K_i, K_j) = \begin{cases} \{0, j/i\}, & \text{if } i|j, \\ \{0\}, & \text{if } i \nmid j. \end{cases} \quad (2)$$

According to Proposition 2.2, for any positive integer $m > 0$ that we fix, there exist finite sequences, $B(i), i = 0, \dots, n$, of not necessarily pairwise distinct non-zero integers, satisfying that

$$A = \bigcap_{i=0}^n S_{B(i)}.$$

Now, we choose particular pairwise distinct primes q_0, q_1, \dots, q_n fulfilling the condition

$$q_i > \max\{|b| \mid b \in B(i)\}, \quad i = 0, \dots, n,$$

and we denote

$$\alpha_i = q_i \prod_{b \in B(i)} b, \quad i = 0, \dots, n.$$

Then, we construct the following ‘‘intermediate’’ manifolds (that will serve us to realize each of the sums $S_{B(i)}$), for $i = 0, \dots, n$:

$$\begin{aligned} M_i &= \#_{b \in B(i)} K_{\alpha_i/b} \\ N_i &= K_{\alpha_i}. \end{aligned}$$

Because K_{α_i} are aspherical 3-manifolds, for $i = 0, \dots, n$, we have that $\pi_2(K_{\alpha_i}) = 0$, and conditions to apply Lemma 3.1 hold. Therefore:

$$D(M_i, N_j) = D(\#_{b \in B(i)} K_{\alpha_i/b}, K_{\alpha_j}) = \sum_{b \in B(i)} D(K_{\alpha_i/b}, K_{\alpha_j}).$$

Using (2), we then get that, for $i = 0, \dots, n$,

$$\begin{aligned} D(M_i, N_i) &= S_{B(i)}, \text{ and} \\ D(M_i, N_j) &= \{0\}, \text{ for } i \neq j. \end{aligned}$$

Since all the conditions to apply Corollary 3.6 plainly hold for the manifolds M_i and N_i , there exists an integer $\ell \geq 0$ such that, for

$$\begin{aligned} M &= M_0 \# M_1 \# \dots \# M_n \# H(3, \ell), \\ N &= N_0 \# N_1 \# \dots \# N_n, \end{aligned}$$

we have that

$$D(M, N) = \bigcap_{i=0}^n S_{B(i)} = A,$$

and the proof of Theorem A is complete. \square

Remark 3.7. We claim that the 3-manifolds K_i , M_i , N_i , $\#_{i=0}^n M_i$, and N involved in the previous proof are inflexible (see also Remark 1.2). It is clear, by (2), that K_i , $i \in \mathbb{Z}$, are inflexible. Now, proceeding along the lines of Theorem A, we apply repeatedly Lemma 3.2 and Lemma 3.1 to get the inflexibility property. On the one hand, we obtain that $D(M_i, M_j) = \{0\}$ for $i \neq j$, and on the other hand

$$D(\#_{i=0}^n M_i, \#_{i=0}^n M_i) \subset \bigcap_{i=0}^n D(M_i, M_i).$$

Also, by Lemma 3.2,

$$D(M_i, M_i) = D(M_i, \#_{b \in B(i)} K_{\alpha_i/b}) \subset \bigcap_{b \in B(i)} D(M_i, K_{\alpha_i/b})$$

and using Lemma 3.1,

$$D(M_i, K_{\alpha_i/b}) = D(\#_{b' \in B(i)} K_{\alpha_i/b'}, K_{\alpha_i/b}) = \sum_{b' \in B(i)} D(K_{\alpha_i/b'}, K_{\alpha_i/b}).$$

Now, by Equation (2), $D(K_{\alpha_i/b'}, K_{\alpha_i/b})$ is either $\{0\}$ or $\{0, b'/b\}$ whenever $b|b'$. Hence, $D(M_i, K_{\alpha_i/b})$ is bounded, and so $D(M_i, M_i)$ and $D(\#_{i=0}^n M_i, \#_{i=0}^n M_i)$ are bounded. Hence M_i , $i = 1, \dots, n$ and $\#_{i=0}^n M_i$ are inflexible manifolds. The same arguments work for N_i and N , and we conclude.

Remark 3.8. The 3-manifolds involved in Theorem A are not unique. Let P be any 3-manifold such that $\pi_1(P)$ is finite (for example the Poincaré sphere $S^3/\mathrm{SL}(2, 5)$, or any lens space $L(p, q)$). Let K_i, K_j be the building blocks in the proof of Theorem A. Then, according to Lemma 3.1,

$$D(P \# K_i, K_j) = D(P, K_j) + D(K_i, K_j) = \{0\} + D(K_i, K_j) = D(K_i, K_j)$$

since K_j is aspherical, and $\pi_1(K_j)$ torsion free while $\pi_1(P)$ is finite. Therefore, in the construction of M , we can replace any K_i by $P \# K_i$ to obtain a new 3-manifold M' that verifies $D(M, N) = D(M', N)$.

4. SPHERICAL FIBRATIONS OVER INFLEXIBLE SULLIVAN MODELS: RATIONAL MAPPING DEGREE SET

In this section we prove Theorem B, which can be thought of as the rational version of Theorem A. Rational homotopy theory provides an equivalence of categories between the category of simply connected rational spaces and the category of certain differential graded algebras, the so-called Sullivan minimal models. We refer to [11] for basics facts in rational homotopy theory.

More concretely, if V is a graded rational vector space, we write ΛV for the free commutative graded algebra on V . A Sullivan model $(\Lambda V, \partial)$ is a commutative differential

graded algebra (cdga for short) which is free as commutative graded algebra on a simply connected graded vector space V of finite dimension in each degree, and such that V admits a basis x_α indexed by a well-ordered set with $\partial(x_\alpha) \in \Lambda(x_\beta)_{\beta < \alpha}$. It is minimal if in addition $\partial(W) \subset \Lambda^{\geq 2}W$.

Now, if M is an oriented closed simply connected manifold, then the cohomology of the associated minimal model A_M coincides with the rational cohomology of M . In particular A_M has a cohomological fundamental class $[A_M] \in H^*(A_M) \cong H^*(M; \mathbb{Q})$ which is isomorphic to the rational cohomological fundamental class $[M]_{\mathbb{Q}}$ of M .

Ellipticity for a Sullivan minimal model $(\Lambda V, \partial)$ means that both V and $H^*(\Lambda V)$ are finite-dimensional. Hence, the cohomology is a Poincaré duality algebra [13] and one can easily compute the degree of its fundamental cohomological class [11, Theorem 32.6]. In particular one can introduce the notion of mapping degree between elliptic Sullivan minimal models and also translate the notion of inflexibility:

Let $(\Lambda V, \partial)$ be an elliptic Sullivan minimal model. Let $\mu \in (\Lambda V)^n$ be a representative of its cohomological fundamental class. Then $(\Lambda V, \partial)$ is *inflexible* if for every cdga-morphism

$$\varphi: (\Lambda V, \partial) \rightarrow (\Lambda V, \partial)$$

we have $\deg(\varphi) = 0, \pm 1$, where $H([\mu]) = \deg(\varphi)[\mu]$.

4.1. Rational mapping degree set and connected sums. The following results establish, under certain restrictions, the relationship between rational mapping degree sets and connected sums of manifolds:

Lemma 4.1 ([9, Lemma II.2]). *Let M_1, M_2 and N be oriented closed n -manifolds with $\pi_{n-1}(N_{(0)}) = 0$. Then*

$$D_{\mathbb{Q}}(M_1 \# M_2, N) = D_{\mathbb{Q}}(M_1, N) + D_{\mathbb{Q}}(M_2, N).$$

Proof. Under the same assumptions, in [9, Lemma II.2] is asserted that the following holds:

$$D(M_1 \# M_2, N) \subset D_{\mathbb{Q}}(M_1, N) + D_{\mathbb{Q}}(M_2, N).$$

However, a stronger result is demonstrated in the proof. Namely,

$$D_{\mathbb{Q}}(M_1 \# M_2, N) \subset D_{\mathbb{Q}}(M_1, N) + D_{\mathbb{Q}}(M_2, N).$$

Hence, it suffices to prove the other inclusion. To that end, one can apply the same arguments as in [7, Lemma 7.8]: let $q_{(0)}: (M_1)_{(0)} \# (M_2)_{(0)} \rightarrow (M_1)_{(0)} \vee (M_2)_{(0)}$ denote the rationalization of the pinching map. Then for any given maps $f_i: (M_i)_{(0)} \rightarrow N_{(0)}$, the composition

$$(f_1 \vee f_2) \circ q: (M_1)_{(0)} \# (M_2)_{(0)} \rightarrow N_{(0)}$$

has degree $\deg(f_1) + \deg(f_2)$ and the result follows. □

We give a precise definition of connected sum in the world of cdgas:

Definition 4.2. *Let $A_i, i = 1, 2$, be connected cdgas and let $a_i \in A_i, i = 1, 2$, be elements of the same degree. The connected sum of the pairs $(A_i, [a_i]), i = 1, 2$, is the cdga*

$$(A_1, [a_1]) \# (A_2, [a_2]) \stackrel{\text{def}}{=} (A_1 \oplus_{\mathbb{Q}} A_2) / I,$$

where $A_1 \oplus_{\mathbb{Q}} A_2 \stackrel{\text{def}}{=} (A_1 \oplus A_2)/\mathbb{Q}\{(1, -1)\}$, and $I \subset A_1 \oplus_{\mathbb{Q}} A_2$ is the differential ideal generated by $a_1 - a_2$.

Connected sums of cdgas provide rational models for connected sums of oriented manifolds. Indeed, for M_i , $i = 1, 2$ oriented closed simply connected n -manifold, with Sullivan minimal model A_{M_i} , let m_i be a representative of the cohomological fundamental class of A_{M_i} , for $i = 1, 2$. By [7, Theorem 7.12]

$$(A_{M_1}, [m_1]) \# (A_{M_2}, [m_2]) \quad (3)$$

is a rational model of $M_1 \# M_2$.

We use (3) above to prove a rational version of Proposition 3.5 that does not involve handle-bodies:

Proposition 4.3. *Let M_1, M_2 and N_1, N_2 be oriented closed simply connected n -manifolds such that $\pi_{n-1}(N_j) \otimes \mathbb{Q} = 0$, $j = 1, 2$, and $D_{\mathbb{Q}}(M_i, N_j) = \{0\}$, $i \neq j$. Then*

$$D_{\mathbb{Q}}(M_1 \# M_2, N_1 \# N_2) = D_{\mathbb{Q}}(M_1, N_1) \cap D_{\mathbb{Q}}(M_2, N_2).$$

Proof. According to Lemma 4.1 and the rational version of Lemma 3.2 (which can be proved following the same arguments as in [22, Lemma 4.3]), we get that

$$D_{\mathbb{Q}}(M_1 \# M_2, N_1 \# N_2) \subset D_{\mathbb{Q}}(M_1, N_1) \cap D_{\mathbb{Q}}(M_2, N_2).$$

Conversely, let $(A_{M_i}, [m_i])$ and $(A_{N_i}, [n_i])$ be Sullivan minimal models of $(M_i, [M_i])$ and $(N_i, [N_i])$ respectively, $i = 1, 2$. For

$$0 \neq d \in D_{\mathbb{Q}}(M_1, N_1) \cap D_{\mathbb{Q}}(M_2, N_2)$$

there exists $f_i: A_{N_i} \rightarrow A_{M_i}$ with $f_i(n_i) = d \cdot m_i + \alpha_i$ and α_i a coboundary, $i = 1, 2$. Because $(A_{M_1}, [m_1 + \alpha_1/d]) \# (A_{M_2}, [m_2 + \alpha_2/d])$ and $(A_{N_1}, [n_1]) \# (A_{N_2}, [n_2])$ are rational models for $M_1 \# M_2$ and $N_1 \# N_2$ respectively, then f_1 and f_2 give rise to a well defined cdga-morphism

$$f_1 \# f_2: (A_{N_1}, [n_1]) \# (A_{N_2}, [n_2]) \rightarrow (A_{M_1}, [m_1 + \alpha_1/d]) \# (A_{M_2}, [m_2 + \alpha_2/d])$$

defined by

$$(f_1 \# f_2)(x) = \begin{cases} f_1(x), & \text{if } x \in A_{N_1}, \\ f_2(x), & \text{if } x \in A_{N_2} \end{cases}$$

and whose degree is $\deg(f_1 \# f_2) = d$. □

Remark 4.4. The previous result can be generalized to arbitrary finite iterated connected sums, as in Corollary 3.6. Namely, if M_i, N_i , $i = 1, \dots, r$, are oriented closed simply connected n -manifolds such that $\pi_{n-1}(N_j) = 0$, $j = 1, \dots, r$, and $D_{\mathbb{Q}}(M_i, N_j) = \{0\}$, for $i \neq j$, then

$$D_{\mathbb{Q}}(M_1 \# \dots \# M_r, N_1 \# \dots \# N_r) = \bigcap_{i=1}^r D_{\mathbb{Q}}(M_i, N_i).$$

4.2. Inflexible Sullivan minimal models of inflexible manifolds. Following the same strategy as in Section 3, we consider spherical fibrations over certain elliptic and inflexible Sullivan minimal models (Definition 4.5), whose total spaces are Sullivan minimal models of inflexible manifolds (see Lemma 4.7). These manifolds will be the building blocks to construct, by means of iterated connected sums, manifolds that realize finite sets of rational numbers.

Definition 4.5. *Let (A, ∂) be an elliptic, inflexible Sullivan minimal model of formal dimension $2m$, $m \geq 1$, such that $\pi_j(A) = 0$ for $j \geq 2m - 1$. Fix $\mu \in A$ a representative of its cohomological fundamental class. Then, for any non-zero $q \in \mathbb{Q}$, we define the following Sullivan minimal model*

$$(K_q(A), \partial) := (A \otimes \Lambda(y_{2m-1}), \partial)$$

that extends the differential of A by $\partial(y_{2m-1}) = q\mu$.

Remark 4.6. Notice that $(K_q(A), \partial)$ is the total space in the rational S^{2m-1} -fiber sequence:

$$(\Lambda(y_{2m-1}), 0) \longleftarrow (K_q(A), \partial) \longleftarrow (A, \partial),$$

whose Euler class is $q[\mu]$.

Lemma 4.7. *Let (A, ∂) be an elliptic, inflexible Sullivan minimal model of formal dimension $2m$, $m \geq 1$, such that $\pi_j(A) = 0$ for $j \geq 2m - 1$. Fix $\mu \in A$ a representative of the fundamental class of A , and let $x \in A$ such that $\partial(x) = \mu^2$. Then, for any non-zero $q \in \mathbb{Q}$, $(K_q(A), [y_{2m-1}\mu - qx])$ is the Sullivan minimal model of an oriented closed inflexible $(4m - 1)$ -manifold M_{K_q} , with the same connectivity as (A, ∂) .*

Proof. According to [8, Proposition 3.1], $(K_q(A), \partial)$ is an elliptic Sullivan model of formal dimension $4m - 1$ where $y_{2m-1}\mu - qx$ is a representative of its cohomological fundamental class. By [8, Lemma 3.2], $(K_q(A), \partial)$ is an inflexible algebra because (A, ∂) is so. Now, since its formal dimension is $4m - 1 \equiv 3 \pmod{4}$, the obstruction theory of Sullivan [25, Theorem (13.2)] and Barge [3, Théorème 1] guarantees that $(K_q(A), [y_{2m-1}\mu - qx])$ is the Sullivan minimal model of an oriented closed simply connected manifold M_{K_q} . Finally, by [6, Proposition A.1], M_{K_q} and (A, ∂) have the same connectivity. \square

We compute the rational mapping degree set between the manifolds appearing in the previous lemma:

Lemma 4.8. *For any non-zero $q \in \mathbb{Q}$, let M_{K_q} be the oriented closed manifold from Lemma 4.7 whose Sullivan minimal model is $(K_q(A), \partial)$ from Definition 4.5. Then*

$$D_{\mathbb{Q}}(M_{K_p}, M_{K_q}) = \{0, q/p\}.$$

Proof. We follow the ideas in [8, Lemma 3.2]. Let $f: (K_q(A), \partial) \rightarrow (K_p(A), \partial)$ be a morphism of non-trivial degree $d \in \mathbb{Q}$, that is,

$$f(y_{2m-1}\mu - qx) = d(y_{2m-1}\mu - px) + \alpha \tag{4}$$

where α is a coboundary. By a degree argument, f induces a non-trivial degree morphism $f|_A: (A, \partial) \rightarrow (A, \partial)$. On the one hand $f(\mu) = \tilde{d}\mu + \beta_1$ and $f(x) = \tilde{d}^2x + \beta_2$ where β_1, β_2 are coboundaries, and $\tilde{d} \in \{-1, 1\}$. On the other hand, $f(y_{2m-1}) = ay_{2m-1} + \gamma$ where $a \in \mathbb{Q}$ and γ is a coboundary.

Because $f(\partial y_{2m-1}) = \partial f(y_{2m-1})$, we get that $ap = q\tilde{d}$ and $\beta_1 = 0$. Hence $a = \tilde{d}(q/p)$ and

$$\begin{aligned} f(y_{2m-1}\mu - qx) &= (\tilde{d}q/p y_{2m-1} + \gamma)(\tilde{d}\mu) - q(\tilde{d}^2x + \beta_2) \\ &= (\tilde{d}^2q/p)(y_{2m-1}\mu - px) - q\beta_2 \\ &= (q/p)(y_{2m-1}\mu - px) - q\beta_2 \quad (\text{recall } \tilde{d} \in \{-1, 1\}). \end{aligned}$$

By comparing this equation to (4), we obtain that $d = q/p$ and the proof is complete. \square

We illustrate the existence of elliptic, inflexible Sullivan minimal models satisfying the conditions from Definition 4.5 and Lemma 4.7:

Definition 4.9. *Let Γ be a connected finite simple graph with more than one vertex, i.e., $|V(\Gamma)| > 1$. Given an integer $k \geq 1$, let $(A_k(\Gamma), \partial)$ be the $(30k + 17)$ -connected elliptic and inflexible Sullivan algebra constructed in [6, Definition 2.1], whose formal dimension is $2m = 540k^2 + 984k + 396 + |V(\Gamma)|(360k^2 + 436k + 132)$ and $\pi_j(A_k(\Gamma)) = 0$ for $j \geq 2m - 1$. Fix $\mu \in A_k(\Gamma)$ a representative of the cohomological fundamental class. Then for any non-zero $q \in \mathbb{Q}$, define the following Sullivan minimal model*

$$(K_q(\Gamma, k), \partial) := (A_k(\Gamma) \otimes \Lambda(y_{2m-1}), \partial)$$

that extends the differential of $A_k(\Gamma)$ by $\partial(y_{2m-1}) = q\mu$.

Remark 4.10. Because conditions from Lemma 4.7 hold, $(K_q(\Gamma, k), \partial)$ is a Sullivan model of an oriented closed $(30k + 17)$ -connected inflexible $(4m - 1)$ -manifold $M_{K_q(\Gamma, k)}$, where $2m = 540k^2 + 984k + 396 + |V(\Gamma)|(360k^2 + 436k + 132)$.

Lemma 4.11. *Let Γ_1 and Γ_2 be connected finite simple graphs with $|V(\Gamma_1)| = |V(\Gamma_2)| > 1$. Given a positive integer $k \geq 1$, and a non-zero $p_i \in \mathbb{Q}$, $i = 1, 2$, consider the manifold $M_{K_{p_i}(\Gamma_i, k)}$, $i = 1, 2$ as in Remark 4.10. Then*

$$D_{\mathbb{Q}}(M_{K_{p_1}(\Gamma_1, k)}, M_{K_{p_2}(\Gamma_2, k)}) = \begin{cases} \{0, p_2/p_1\}, & \text{if } \Gamma_1 \cong \Gamma_2, \\ \{0\}, & \text{otherwise.} \end{cases}$$

Proof. Let $(K_{p_i}(\Gamma_i, k), \partial) = (A_k(\Gamma_i) \otimes \Lambda(y_i), \partial)$, introduced in Definition 4.9, be the Sullivan model of the manifold $M_{K_{p_i}(\Gamma_i, k)}$, where $\partial(y_i) = p_i\mu_i$ for μ_i a representative of the cohomological fundamental class of $A_k(\Gamma_i)$, $i = 1, 2$. Recall from Lemma 4.7 that for $x_i \in A_k(\Gamma_i)$ satisfying $\partial(x_i) = \mu_i^2$, the element $y_i\mu_i - p_ix_i$ is a representative of the cohomological fundamental class of $(K_{p_i}(\Gamma_i, k), \partial)$, $i = 1, 2$.

With these constructions in mind, we follow the ideas from [8, Lemma 3.2]. Consider a morphism of non-trivial degree $d \in \mathbb{Q}$:

$$f: (K_{p_2}(\Gamma_2, k), \partial) \rightarrow (K_{p_1}(\Gamma_1, k), \partial).$$

Then $f(y_2\mu_2 - p_2x_2) = d(y_1\mu_1 - p_1x_1) + \alpha$ with α a coboundary. By a degree argument, the morphism f induces a non-trivial degree morphism

$$f|_{A_k(\Gamma_2)}: (A_k(\Gamma_2), \partial) \rightarrow (A_k(\Gamma_1), \partial).$$

Focusing specifically on this former morphism, the arguments in [6, Lemma 2.12] (see also [8, Remark 2.8]), show that it is induced by a graph full monomorphism $\sigma: \Gamma_1 \rightarrow \Gamma_2$.

Now, since $|V(\Gamma_1)| = |V(\Gamma_2)|$, σ is indeed an isomorphism of graphs, and $f(\mu_2) = \mu_1 + \beta_1$ and $f(x_2) = x_1 + \beta_2$ with β_1, β_2 coboundaries, by [6, Lemma 2.12].

Finally, by another degree reasoning argument, one obtains that $f(y_2) = ay_1 + \gamma$ where a is a non-zero rational number, and γ is a coboundary. We conclude as in the proof of Lemma 4.8. \square

4.3. Proof of Theorem B. Let $A = \{0, d_1, \dots, d_n\}$ where d_1, d_2, \dots, d_n are pairwise different non-zero rational numbers. Fix an integer $k \geq 1$. According to Corollary 2.3, there exist finite sequences of not necessarily pairwise distinct non-zero rational numbers $B(i)$, $i = 0, \dots, n$, such that

$$A = \bigcap_{i=0}^n S_{B(i)}.$$

Choose $\Gamma_0, \Gamma_1, \dots, \Gamma_n$, pairwise non-isomorphic connected finite simple graphs, such that $|V(\Gamma_i)| = |V(\Gamma_j)| > 1$ for every $i, j = 0, \dots, n$. According to Remark 4.10, we define the $(30k + 17)$ -connected manifolds

$$M_i = \#_{b \in B(i)} M_{K_{b-1}(\Gamma_i, k)}$$

$$N_i = M_{K_1(\Gamma_i, k)},$$

for $i = 0, \dots, n$. By Lemmas 4.11 and the rational version of Lemma 3.2 (which can be proved following the same arguments as in [22, Lemma 4.3]), we have that

$$D_{\mathbb{Q}}(M_i, N_i) = S_{B(i)}, \text{ and}$$

$$D_{\mathbb{Q}}(M_i, N_j) = \{0\}, \text{ for } i \neq j.$$

Finally, define

$$M = M_0 \# M_1 \# \dots \# M_n,$$

$$N = N_0 \# N_1 \# \dots \# N_n.$$

and use Proposition 4.3 (see also Remark 4.4) to get

$$D_{\mathbb{Q}}(M, N) = \bigcap_{i=0}^n S_{B(i)} = A.$$

5. FROM UNSTABLE ADAMS OPERATIONS TO MAPPING DEGREE SETS

We recall the basics on unstable Adams operations following Jackowski-McClure-Oliver's work [15, 16]. Given a compact connected Lie group G , a self-map $f: BG \rightarrow BG$ is called an unstable Adams operation of degree $r \geq 0$, if $H^{2i}(f; \mathbb{Q})$ is the multiplication by r^i for each $i > 0$ [15, p. 183]. For a given simple Lie group G with Weyl group W_G , an unstable Adams operation of degree $r > 0$ exists if and only if $\gcd(r, |W_G|) = 1$, and moreover, this operation is unique up to homotopy [15, Theorem 2]. In particular, when $G = \text{SO}(2m - 1)$ or $G = \text{SO}(2m)$, $m > 1$, unstable Adams operations of degree $r > 0$ exist if $\gcd(r, m!) = 1$. In what follows, we denote by φ^r the unstable Adams operation of degree $r > 0$ on $B\text{SO}(2m - 1)$ and $B\text{SO}(2m)$. Notice that since they are unique, then $\varphi^s \circ \varphi^r = \varphi^{sr}$.

Henceforward, $(\Sigma, [\Sigma])$ is a fixed oriented closed connected $2m$ -manifold whose rationalization $(\Sigma_{(0)}, [\Sigma]_{\mathbb{Q}})$ is inflexible and $\pi_j(\Sigma_{(0)}) = 0$ for $j \geq 2m - 1$. Let (A_{Σ}, ∂) be a Sullivan minimal model of Σ . Denote by $\pi: \Sigma \rightarrow S^{2m}$ the map obtained by collapsing the $(2m - 1)$ -skeleton of Σ .

Lemma 5.1. *Let $X_{2m} \in H^{2m}(BSO(2m); \mathbb{Z})$ be the Euler class of the spherical fiber sequence*

$$S^{2m-1} \rightarrow BSO(2m-1) \rightarrow BSO(2m),$$

thus X_{2m} is a torsion free integral cohomology class [5, Theorem 1.5, Equation (2.1)], and S^{2m} is thought of as an oriented closed manifold. Then, there exists $\iota: S^{2m} \rightarrow BSO(2m)$, a torsion free element in $\pi_{2m}(BSO(2m))$, and a non-zero integer $\kappa \in \mathbb{Z}$ such that $H^(\iota; \mathbb{Z})(X_{2m}) = \kappa[S^{2m}]$.*

Proof. Recall that $\pi_{2m}(BSO(2m)) \cong \pi_{2m-1}(SO(2m))$. By [19, Corollary IV.6.14] (see also [17, p. 161]), $\pi_{2m-1}(SO(2m))$ contains a copy of \mathbb{Z} inducing the p -local (thus rational) splitting $SO(2m) \simeq_{(p)} SO(2m-1) \times S^{2m-1}$ [19, Corollary IV.6.21]. Let ι be a generator of such a copy of \mathbb{Z} in $\pi_{2m}(BSO(2m))$.

By construction, $H^*(\iota; \mathbb{Q})$ is non-trivial on the Euler class of the rational fiber sequence

$$S^{2m-1}_{(0)} \rightarrow BSO(2m-1)_{(0)} \rightarrow BSO(2m)_{(0)},$$

which is just $X_{2m} \otimes_{\mathbb{Q}} 1$. Therefore, $H^*(\iota; \mathbb{Z})(X_{2m}) = \kappa[S^{2m}]$ for some non-zero $\kappa \in \mathbb{Z}$. \square

Definition 5.2. *Given any integers $r > 0, m > 1$, with r coprime to $m!$, we define:*

- (1) *The oriented $(4m - 1)$ -manifold E_{r^m} as the total space in the principal spherical $SO(2m)$ -fiber bundle*

$$\begin{array}{ccc} S^{2m-1} & \xlongequal{\quad} & S^{2m-1} \\ \downarrow & & \downarrow \\ E_{r^m} & \longrightarrow & BSO(2m-1) \\ \downarrow & \lrcorner & \downarrow \\ \Sigma & \xrightarrow{\phi_r} & BSO(2m), \end{array} \quad (5)$$

where $\phi_r = \varphi^r \circ \iota \circ \pi$.

- (2) *The oriented $(4m - 1)$ -manifold E_{-r^m} obtained by reversing the original orientation on the manifold E_{r^m} introduced above.*

Remark 5.3. By construction, the Euler class of the spherical fiber bundle over Σ given in diagram (5) is $\kappa r^m[\Sigma]$.

Recall from the beginning of this section that $(\Sigma, [\Sigma])$ is a fixed oriented closed connected $2m$ -manifold where (A_{Σ}, ∂) is its Sullivan minimal model.

Lemma 5.4. *Let E_{r^m} be the manifold introduced in Definition 5.2. A Sullivan minimal model of E_{r^m} is $K_{\kappa r^m}(A_{\Sigma})$ as given in Definition 4.5. Therefore E_{r^m} is rationally equivalent to $M_{K_{\kappa r^m}}$, the manifold given in Lemma 4.7.*

Proof. As it was pointed out in Remark 4.6, $K_{\kappa r^m}(A_\Sigma)$ is a Sullivan minimal model for the total space in a rational S^{2m-1} -fiber sequence whose Euler class is $\kappa r^m[\Sigma]_{\mathbb{Q}}$. It coincides with the Euler class of the rationalization of the spherical $\mathrm{SO}(2m)$ -fiber bundle in diagram (5). Therefore $K_{\kappa r^m}(A_\Sigma)$ is a Sullivan minimal model for E_{r^m} . \square

Lemma 5.5. *Let i, j, m be positive integers, $m > 1$, such that $\gcd(i, m!) = \gcd(j, m!) = 1$, and let E_{r^m} , $r = i, j$, be the $(4m - 1)$ -manifold introduced in Definition 5.2. Then*

$$D(E_{i^m}, E_{j^m}) = \begin{cases} \{0, (j/i)^m\}, & \text{if } i|j, \\ \{0\}, & \text{if } i \nmid j. \end{cases}$$

Proof. By Lemma 5.4, the manifolds E_{r^m} and $M_{K_{\kappa r^m}}$ are rationally equivalent, for every $0 < r \in \mathbb{Z}$. Therefore:

$$\begin{aligned} D(E_{i^m}, E_{j^m}) &\subset D_{\mathbb{Q}}(E_{i^m}, E_{j^m}) \cap \mathbb{Z} = D_{\mathbb{Q}}(M_{K_{\kappa i^m}}, M_{K_{\kappa j^m}}) \cap \mathbb{Z} \\ &= \{0, (j/i)^m\} \cap \mathbb{Z} \quad (\text{by Lemma 4.8}) \\ &= \begin{cases} \{0, (j/i)^m\}, & \text{if } i|j, \\ \{0\}, & \text{if } i \nmid j. \end{cases} \end{aligned}$$

The proof will be completed if we construct a map $f: E_{i^m} \rightarrow E_{j^m}$ of degree $(j/i)^m$ when $i|j$. To this end, let us suppose that $j = di$, $d \in \mathbb{Z}$, and recall that unstable Adams operations satisfy that $\varphi^j = \varphi^d \circ \varphi^i$. Therefore, by construction (see Definition 5.2) $\phi_j = \varphi^d \circ \phi_i$. Let $f: E_{i^m} \rightarrow E_{j^m}$ be the map obtained by the universal property of pullbacks in the following commutative diagram:

$$\begin{array}{ccccc} E_{i^m} & \longrightarrow & B\mathrm{SO}(2m-1) & \xrightarrow{\varphi^d} & B\mathrm{SO}(2m-1) \\ & \searrow f & \downarrow & & \downarrow \\ & & E_{j^m} & \longrightarrow & B\mathrm{SO}(2m-1) \\ & & \downarrow \lrcorner & & \downarrow \\ & & \Sigma & \xrightarrow{\phi_i} & B\mathrm{SO}(2m) \xrightarrow{\varphi^d} B\mathrm{SO}(2m) \end{array} \quad (6)$$

Diagram (6) gives rise to a commutative diagram of spherical fiber sequences

$$\begin{array}{ccc} S^{2m-1} & \xrightarrow{\tilde{f}} & S^{2m-1} \\ \downarrow & & \downarrow \\ E_{i^m} & \xrightarrow{f} & E_{j^m} \\ \downarrow & & \downarrow \\ \Sigma & \xlongequal{\quad} & \Sigma, \end{array} \quad (7)$$

whose associated Serre spectral sequences (Sss) can be compared via the edge morphisms given by naturality: the Sss associated to the left (resp. right) side of diagram (7) is fully determined by the differential

$$d_{2m}([S^{2m-1}]) = \kappa i^m[\Sigma] \quad (\text{resp. } d_{2m}([S^{2m-1}]) = \kappa j^m[\Sigma]),$$

and since by naturality

$$d_{2m}(H^*(\tilde{f})([S^{2m-1}])) = H^*(Id_\Sigma)(d_{2m}([S^{2m-1}]))$$

we obtain that $\deg(\tilde{f}) = (j/i)^m$.

Now, the cohomological fundamental class $[E_i^m]$ (resp. $[E_j^m]$) is represented by the class $[S^{2m-1}] \otimes [\Sigma]$ in the $E_\infty^{2m-1, 2m}$ -term of the Sss associated to the left (resp. right) fiber sequence in diagram (7). Hence by naturality

$$\begin{aligned} H^*(f)([E_j^m]) &= H^*(\tilde{f})([S^{2m-1}]) \otimes H^*(Id_\Sigma)([\Sigma]) \\ &= ((j/i)^m [S^{2m-1}]) \otimes [\Sigma] \\ &= (j/i)^m [E_i^m] \end{aligned}$$

and therefore $\deg(f) = (j/i)^m$. \square

Remark 5.6. Notice that manifolds E_{r^m} and E_{-r^m} differ in just the orientation. Hence, for any other oriented closed connected $(4m-1)$ -manifold N , the mapping set degree is $D(E_{-r^m}, N) = -D(E_{r^m}, N)$ and $D(N, E_{-r^m}) = -D(N, E_{r^m})$.

Prior to proving Theorem C, we need an integral version of Proposition 4.3 (see Proposition 5.8). First, we prove the following result:

Lemma 5.7. *Let M, N be oriented closed n -manifolds, $n > 2$, and $d \in D(M, N)$, $d \neq 0$. If N is simply connected, then there exists a map $h: M \rightarrow N$ that induces a map between discs $h|_{D_M}: D_M \rightarrow D_N$ such that $D_M = h^{-1}(D_N)$, $h|_{\partial D_M}: \partial D_M \rightarrow \partial D_N$, and $\deg(h) = \deg(h|_{\partial D_M}) = d$.*

Proof. We take a map $f: M \rightarrow N$ of degree d , that we can assume to be smooth. Take a regular value $y_0 \in N$ of f , and let x_1, \dots, x_s be the preimages. We take a collection of smooth regular curves L_j from x_1 to x_j , $j = 2, \dots, s$, which do not intersect each other except at the original point x_1 , and write $L = \bigcup_{j=2}^s L_j$.

Around each L_j we are going to build a spindle as follows: Take a parametrization $\gamma_j: [0, 1] \rightarrow L_j$, with $\gamma_j(0) = x_1$, $\gamma_j(1) = x_j$, and a tubular closed neighbourhood $\Gamma_j: [0, 1] \times D_\epsilon^{n-1} \rightarrow U_j \subset M$ around L_j . The spindle is the subset

$$S_j^\epsilon := \{\Gamma_j(t, x) \mid |x| \leq \epsilon t(1-t), t \in [0, 1]\}.$$

Taking $\epsilon > 0$ small enough, we can assume $S_j^\epsilon \cap S_{j'}^\epsilon = \{x_1\}$ for $j \neq j'$. Then, there exists a retraction $r: M \rightarrow M$ such that it is the identity on $M \setminus \bigcup S_j^\epsilon$, and sends $S_j^{\epsilon/2}$ to L_j so that $r(\Gamma_j(t, t(1-t)\bar{x})) = \gamma_j(t)$ for $\bar{x} \in D_{\epsilon/2}^{n-1}$. Indeed, if $\Gamma_j(t, x) \in S_j^\epsilon$ write $(t, x) = (t, t(1-t)\bar{x})$ where $\bar{x} \in D_\epsilon^{n-1}$, and define

$$r(\Gamma_j(t, t(1-t)\bar{x})) = \begin{cases} \gamma_j(t), & |\bar{x}| \leq \epsilon/2. \\ \Gamma_j(t, (\frac{2}{\epsilon}|\bar{x}| - 1)t(1-t)\bar{x}), & \epsilon/2 \leq |\bar{x}| \leq \epsilon. \end{cases}$$

Now let $\hat{f} = f \circ r: M \rightarrow N$, which has degree d and maps $S_j^{\epsilon/2}$ to the curve $R_j = f(L_j)$ in N . Note that $\hat{f}^{-1}(y_0) = \{x_1, \dots, x_s\}$.

Since N is simply connected we can construct a contraction from R_j to y_0 being careful not to cross y_0 . The curve R_j is parametrized by $\beta_j = f \circ \gamma_j$. Take a small ball $B_\delta(y_0)$

and two points $0 < a < b < 1$ such that $\beta_j([0, a]), \beta_j([b, 1]) \subset B_\delta(y_0)$ and $\beta_j(a), \beta_j(b) \in \partial B_\delta(y_0)$. Using that $N \setminus B_\delta(y_0)$ is simply-connected, we construct a homotopy of $\beta_j|_{[a, b]}$ to a path in $\partial B_\delta(y_0)$ from $\beta_j(a)$ to $\beta_j(b)$. This gives a homotopy from β_j to a loop in $B_\delta(y_0)$ not crossing y_0 . Next we take a radial retraction to y_0 , and by juxtaposition, we get a homotopy $H_j(t, s)$ from $\beta_j(t)$ to the constant path $H_j(t, 1) = y_0$, such that $H_j(t, s) \neq y_0$, if $s < 1$ and $t \in (0, 1)$.

Now we define $\tilde{f}: M \rightarrow N$ as follows. First $\tilde{f} = \hat{f}$ on $M \setminus S_j^{\epsilon/2}$. Now on $S_j^{\epsilon/2}$, we set

$$\tilde{f}(\Gamma_j(t, t(1-t)\bar{x})) = H_j(t, 1 - 2|\bar{x}|/\epsilon),$$

for $\bar{x} \in D_{\epsilon/2}^{n-1}$. This is continuous, because at the boundary $|\bar{x}| = \epsilon/2$, we have $\tilde{f} = \beta_j(t)$. We also have $\tilde{f}|_{L_j} = y_0$, when $|\bar{x}| = 0$, and these are the only possible points such that they take value y_0 . Therefore

$$\tilde{f}^{-1}(y_0) = \bigcup L_j = L.$$

Finally, we contract L in M and use the degree 1 homeomorphism $M \cong M/L$, to define $h: M \cong M/L \xrightarrow{\tilde{f}} N$ verifying $h^{-1}(y_0)$ is one point, namely $p \in M$ and $\deg(h) = d$. Then, taking small enough $\delta, \xi > 0$ we may homotope h so that $D_\delta^n(p) = h^{-1}(D_\xi^n(y_0))$ and $h|_{\partial D_\delta(p)}: S^{n-1} \rightarrow \partial D_\xi^n(y_0) \cong S^{n-1}$, which has degree d . \square

Proposition 5.8. *Let M_1, M_2 and N_1, N_2 be oriented closed n -manifolds, $n > 2$, verifying:*

- (1) N_i is simply connected, $i = 1, 2$;
- (2) $\pi_{n-1}(N_i) = 0$ for $i = 1, 2$;
- (3) $D(M_i, N_j) = \{0\}$ for $i \neq j$.

Then

$$D(M_1 \# M_2, N_1 \# N_2) = D(M_1, N_1) \cap D(M_2, N_2).$$

Proof. By combining Lemmas 3.1 and 3.2, it follows directly that:

$$D(M_1 \# M_2, N_1 \# N_2) \subset D(M_1 \# M_2, N_1) \cap D(M_1 \# M_2, N_2) = D(M_1, N_1) \cap D(M_2, N_2).$$

Conversely, let $d \in D(M_1, N_1) \cap D(M_2, N_2)$. Then, by Lemma 5.7, there exists $g_i: M_i \rightarrow N_i$ that induces a map between discs $g_i|_{D_{M_i}}: D_{M_i} \rightarrow D_{N_i}$ such that $D_{M_1} = g_1^{-1}(D_{N_1})$ and $g_i|_{\partial D_{M_i}}: \partial D_{M_i} \rightarrow \partial D_{N_i}$, and $\deg(g_i) = \deg(g_i|_{\partial D_{M_i}}) = d$, $i = 1, 2$. Hence, gluing $M_1 \# M_2$ along ∂D_{M_i} , $i = 1, 2$, and $N_1 \# N_2$ along ∂D_{N_i} , $i = 1, 2$, gives us a well defined a map

$$g_1 \# g_2: M_1 \# M_2 \rightarrow N_1 \# N_2$$

whose degree, by construction, is precisely d . Therefore

$$D(M_1, N_1) \cap D(M_2, N_2) \subset D(M_1 \# M_2, N_1 \# N_2)$$

and we conclude the proof. \square

As it was mentioned in Section 3, Lemma 3.1 and Lemma 3.2 can be easily generalized to finite iterated connected sums. Hence, following along the lines of the proof in Proposition 5.8 we obtain:

Corollary 5.9. *Let $M_i, N_i, i = 1, \dots, r$, be oriented closed connected n -manifolds, $n > 2$, verifying:*

- (1) N_i is simply connected, $i = 1, \dots, r$;
- (2) $\pi_{n-1}(N_i) = 0$ for $i = 1, \dots, r$;
- (3) $D(M_i, N_j) = \{0\}$ for $i \neq j$.

Then

$$D(M_1 \# \dots \# M_r, N_1 \# \dots \# N_r) = \bigcap_{i=1}^r D(M_i, N_i).$$

Proof of Theorem C. Let Σ be an oriented closed k -connected $2m$ -manifold verifying that $\Sigma_{(0)}$ is inflexible and $\pi_j(\Sigma_{(0)}) = 0$ for $j \geq 2m - 1$. Let $A = \{0, d_1, \dots, d_n\}$ where d_1, d_2, \dots, d_n are pairwise different non-zero integers.

According to Proposition 2.2, there exist finite sequences $B(i), i = 0, \dots, n$, of not necessarily pairwise distinct non-zero integers, such that every element in $B(i)$ can be written as $\pm r^m$ for $0 < r \in \mathbb{Z}$ with $\gcd(r, m!) = 1$, and

$$A = \bigcap_{i=0}^n S_{B(i)}.$$

Choose pairwise distinct prime numbers q_0, q_1, \dots, q_n , in such a way that

$$q_j > \max\{|b| \mid b \in B(i), i = 0, \dots, n\}$$

and $\gcd(q_j, m!) = 1$, for $j = 0, \dots, n$. Let $\alpha_i = q_i^m \prod_{b \in B(i)} b$, for every $i = 0, \dots, n$. Notice that α_i and $\alpha_i/b, b \in B(i)$, are integers that can be written up to a sign as r^m for some positive integer r such that $\gcd(r, m!) = 1$.

Following the notation in Definition 5.2, we define the following $(4m - 1)$ -manifolds

$$\begin{aligned} M_i &= \#_{b \in B(i)} E_{\alpha_i/b} \\ N_i &= E_{\alpha_i} \end{aligned}$$

for $i = 0, \dots, n$. According to Lemma 5.5 and Lemma 3.1, we deduce that

$$\begin{aligned} D(M_i, N_i) &= S_{B(i)}, \text{ and} \\ D(M_i, N_j) &= \{0\}, \text{ for } i \neq j. \end{aligned}$$

Finally, we construct

$$\begin{aligned} M &= M_0 \# M_1 \# \dots \# M_n, \\ N &= N_0 \# N_1 \# \dots \# N_n. \end{aligned}$$

If $k > 0$, thus the N_i are simply connected, according to Corollary 5.9, we obtain that

$$D(M, N) = \bigcap_{i=0}^n S_{B(i)} = A.$$

If $k = 0$, we apply Corollary 3.6 to get that there exists $\ell \geq 0$ such that

$$D(M\#H(n, \ell), N) = \bigcap_{i=0}^N S_{B(i)} = A.$$

REFERENCES

- [1] M. Amann. Degrees of self-maps of simply connected manifolds. *Int. Math. Res. Not.*, 2015(18):8545–8589, 2015.
- [2] J. M. Ancochea and R. Campoamor. Characteristically nilpotent Lie algebras: A survey. *Extr. Math.*, 16(2):153–210, 2001.
- [3] J. Barge. Structures différentiables sur les types d’homotopie rationnelle simplement connexes. *Ann. Sci. Éc. Norm. Supér. (4)*, 9:469–501, 1976.
- [4] I. Belegradek. On co-Hopfian nilpotent groups. *Bull. Lond. Math. Soc.*, 35(6):805–811, 2003.
- [5] E. H. j. Brown. The cohomology of BSO_n and BO_n with integer coefficients. *Proc. Am. Math. Soc.*, 85:283–288, 1982.
- [6] C. Costoya, D. Méndez, and A. Viruel. Homotopically rigid Sullivan algebras and their applications. In *An alpine bouquet of algebraic topology*, volume 708 of *Contemp. Math.*, pages 103–121. Amer. Math. Soc., Providence, RI, 2018.
- [7] C. Costoya, V. Muñoz, and A. Viruel. On strongly inflexible manifolds. *Int. Math. Res. Not.*, 2023(9):7355–7390, 2023.
- [8] C. Costoya and A. Viruel. Every finite group is the group of self-homotopy equivalences of an elliptic space. *Acta Math.*, 213(1):49–62, 2014.
- [9] D. Crowley and C. Löh. Functorial seminorms on singular homology and (in)flexible manifolds. *Algebr. Geom. Topol.*, 15(3):1453–1499, 2015.
- [10] E. Dror Farjoun. *Cellular spaces, null spaces and homotopy localization*, volume 1622 of *Lect. Notes Math.* Berlin: Springer-Verlag, 1995.
- [11] Y. Félix, S. Halperin, and J.-C. Thomas. *Rational homotopy theory*, volume 205 of *Springer-Verlag*. Springer-Verlag, New York, 2001.
- [12] V. Guillemin and A. Pollack. *Differential topology*. Prentice-Hall, Inc., Englewood Cliffs, NJ, 1974.
- [13] S. Halperin. Finiteness in the minimal models of sullivan. *Trans. Amer. Math. Soc.*, 230:173–199, 1977.
- [14] K. Hasegawa. Minimal models of nilmanifolds. *Proc. Am. Math. Soc.*, 106(1):65–71, 1989.
- [15] S. Jackowski, J. E. McClure, and B. Oliver. Homotopy classification of self-maps of BG via G -actions. I. *Ann. Math. (2)*, 135(1):183–226, 1992.
- [16] S. Jackowski, J. E. McClure, and B. Oliver. Homotopy classification of self-maps of BG via G -actions. II. *Ann. Math. (2)*, 135(2):227–270, 1992.
- [17] M. A. Kervaire. Some non-stable homotopy groups of Lie groups. *Ill. J. Math.*, 4:161–169, 1960.
- [18] C. Loeh and M. Uschold. L^2 -betti numbers and computability of reals, 2022. arXiv:2202.03159.
- [19] M. Mimura and H. Toda. *Topology of Lie groups, I and II. Transl. from the Jap. by Mamoru Mimura and Hiroshi Toda*, volume 91 of *Transl. Math. Monogr.* Providence, RI: American Mathematical Society, 1991.
- [20] D. Müllner. Orientation reversal of manifolds. *Algebr. Geom. Topol.*, 9(4):2361–2390, 2009.
- [21] C. Neofytidis. Degrees of self-maps of products. *Int. Math. Res. Not. IMRN*, (22):6977–6989, 2017.
- [22] C. Neofytidis, S. Wang, and Z. Wang. Realising sets of integers as mapping degree sets. *Bull. Lond. Math. Soc. (to appear)*, 2022.
- [23] P. Renteln. *Manifolds, tensors, and forms. An introduction for mathematicians and physicists*. Cambridge: Cambridge University Press, 2014.
- [24] A. Sard. The measure of the critical values of differentiable maps. *Bull. Am. Math. Soc.*, 48:883–890, 1942.
- [25] D. Sullivan. Infinitesimal computations in topology. *Publ. Math., Inst. Hautes Étud. Sci.*, 47:269–331, 1977.

CITMAGA, DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDADE DE SANTIAGO DE COMPOSTELA,
15705-A CORUÑA, SPAIN.

Email address: `crisrina.costoya@usc.es`

INSTITUTO DE MATEMÁTICA INTERDISCIPLINAR AND DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y
TOPOLOGÍA, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA DE LAS CIENCIAS, 3, 28040-MADRID,
SPAIN

Email address: `vicente.munoz@ucm.es`

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, UNIVERSIDAD DE MÁLAGA, CAMPUS
DE TEATINOS, S/N, 29071-MÁLAGA, SPAIN

Email address: `viruel@uma.es`