

Quantum Gravity Signatures in the Late-Universe

Michael W. Toomey,^{1,2} Savvas Koushiappas,^{1,2} Bruno Alexandre,³ and João Magueijo³

¹ *Department of Physics, Brown University, Providence, RI 02912-1843, USA*

² *Brown Theoretical Physics Center, Brown University, Providence, RI 02912-1843, USA*

³ *Theoretical Physics Group, The Blackett Laboratory, Imperial College,
Prince Consort Rd., London, SW7 2BZ, UK*

We calculate deviations in cosmological observables as a function of parameters in a class of connection-based models of quantum gravity. In this theory non-trivial modifications to the background cosmology can occur due to a distortion of the wave function of the Universe at the transition from matter to dark energy domination (which acts as a “reflection” in connection space). We are able to exclude some regions of parameter space and show with projected constraints that future experiments like DESI will be able to further constrain these models. An interesting feature of this theory is that there exists a region of parameter space that could naturally alleviate the S_8 tension.

I. INTRODUCTION

One of the most striking features of Λ CDM cosmology is the recent transition from decelerated to accelerated expansion. While the dynamics of this transition through the lens of the concordance model is well understood and has been studied *ad nauseam*, it has recently been pointed out [1] that from the perspective of some theories of quantum gravity, such a transition can result in a novel evolution of the background cosmology. The reason is because the expansion rate \dot{a} directly maps to the (connection or gauge field) degree of freedom of the gravitational theory and the semi-classical limit at the transition breaks down. This manifests as a brief enhancement in the Hubble parameter around dark energy — matter equality.

This feature is restricted to first order gravity, a connection based formulation of general relativity where the metric is no longer a fundamental degree of freedom. This “connection-based” representation of general relativity is classically entirely equivalent to the metric formulation but can have several advantages over second order gravity, e.g., coupling gravity to fermions, a feature that played a pivotal role in the development of loop quantum gravity. Here, we are interested in the effects of the recent dark energy—matter transition in the context of the connection representation because of the predictive effects on the dynamical evolution of the connection variable, $b = \dot{a}$. Classically, b decreases before the transition (decelerated expansion, $\dot{b} = \ddot{a} < 0$), reaches a minimum, and then increases (accelerated expansion, $\dot{b} = \ddot{a} > 0$). During the dark energy—matter transition, interference of incident and reflected waves manifests a brief enhancement in the Hubble parameter, but more importantly, still, the wave function is squashed against the minimum- b -wall, creating a skewed wave function, and so a bias towards larger Hubble parameters [1].

While its possible to directly constrain deviations to the background cosmology, if modifications of this nature are large enough they can also appreciably alter the growth of large-scale structure (LSS) and low- ℓ features in the cosmic microwave background (CMB) via the in-

tegrated Sachs-Wolfe (ISW) effect. In this work, we explicitly calculate the novel signatures present in this class of quantum gravity models.

In Section II we review the model of quantum cosmology considered in this work, in Section III the relevant cosmological signatures and in Section IV we discuss the implication of our results.

II. THEORY OF QUANTUM COSMOLOGY

The model studied here is a generalization of unimodular gravity [2–7] based on [8]. In this formulation the cosmological constant is no longer a coupling constant, but rather an integration constant, or a constant as a result of the equations of motion: a constant on-shell only. Since Λ appears in the phase space (rather than being a parameter) it has a dual variable, which acts as a physical time variable (this turns out to be unimodular or 4-volume time on-shell [9, 10]) As a result, the wave function of the Universe in these theories does not obey the standard Wheeler-DeWitt equation, but rather a Schrödinger-like equation. One benefit of this approach is that one is not faced with the same normalization issues that typically plague solutions to the Wheeler-DeWitt equation [11]¹ since we can now superpose waves with different Λ to form wave packets.

Critically, this approach also avoids the issue of time in quantum gravity [13, 14], because as already mentioned, the canonical conjugates to the integration constants in these theories are great candidates for a cosmic time. For example, the cosmological constant is conjugate to the 4D volume of past observers. This approach can also be extended more generally to constants derived from a perfect fluid which will also have a conjugate corresponding to a cosmic time. When one fluid dominates over the other, we can think of that fluid’s conjugate serving the

¹ Recent progress suggest this problem may actually be surmountable [12].

role of a clock for the Universe. We will now outline some of the details of the model considered in this work. For more details see [1].

A. Classical Theory

The Einstein-Cartan action in the connection based formulation and a homogeneous and isotropic minisuperspace model is given by,

$$S_{GR} = \frac{3V_c}{8\pi G} \int dt \left(\dot{b}a^2 + Na(b^2 + k) \right), \quad (1)$$

where the connection variable is $b = \dot{a}/N$, N is the lapse function, $V_c = \int d^3x$ is the coordinate 3-volume from the slicing and $k = 0, \pm 1$ is the standard curvature term. In this paper we are interested in the effects at the transition from matter to dark energy domination. Thus, we consider a cosmological model for two perfect fluids with equations of state $w = 0$ and $w = -1$. A perfect fluid in the minisuperspace representation is defined at the level of the action as [15, 16],

$$S_{\text{fl}} = \int dt (U\dot{\tau} - Na^3V_c\rho n) \quad (2)$$

with U the total (conserved) particle number, $n = U/a^3V_c$ is the number density, and τ a Lagrange multiplier. Assuming the standard evolution of a fluid with equation of state w one can define a new conserved variable

$$m = \frac{8\pi G\rho_0}{3V_c} \left(\frac{U}{V_c} \right)^{1+w} \quad (3)$$

from which we can write down an alternative form of Eq. 2 (see also [17–19])

$$S_{\text{fl}}^{(w)} = \frac{3V_c}{8\pi G} \int dt \left(\dot{m}\chi - N \frac{m}{a^{3w}} \right). \quad (4)$$

With this in hand we can now write down our full action for gravity with dark matter and dark energy,

$$S_{GR} = \frac{3V_c}{8\pi G} \int dt \left[\dot{b}a^2 + \dot{m}\chi_1 + \dot{\Lambda}\chi_2 - Na \left(-(b^2 + k) + \frac{m}{a} + \frac{\Lambda}{3}a^2 \right) \right]. \quad (5)$$

This action implies the corresponding set of non-vanishing Poisson brackets,

$$\{b, a^2\} = \{m, \chi_1\} = \{\Lambda, \chi_2\} = \frac{8\pi G}{3V_c}, \quad (6)$$

where we can view b as a coordinate and a^2 as its conjugate momentum. Additionally, we have our “constants” m and Λ along with their conjugate time variable. Now we can write down the Hamiltonian density for the theory,

$$\mathcal{H} = \frac{3V_c}{8\pi G} Na \left(-(b^2 + k) + \frac{m}{a} + \frac{\Lambda}{3}a^2 \right). \quad (7)$$

Hamilton’s equations for N give the Hamiltonian constraint,

$$-(b^2 + k) + \frac{m}{a} + \frac{\Lambda}{3}a^2 = 0, \quad (8)$$

which is just the Friedmann equation. Going forward we will now make two simplifications. First we will abandon the variables for dark energy in favor of a canonically transformed pair, $\phi = 3/\Lambda$ and $T_\phi = -3\chi_2/\phi^2$, and second we will define a potential $V(b) = (b^2 + k)$. As demonstrated in [1, 20–22] the Hamiltonian constraint can then be recast in the form of two constraints,

$$h_\pm(b)a^2 - m^2 = 0, \quad (9)$$

where one will hold before the bounce and the other afterwards.

B. Quantum Theory

We can now quantize the theory by promoting the classical Poisson bracket (Eq. 6) to a commutation relation,

$$[\hat{b}, \hat{a}^2] = i \frac{l_P^2}{3V_c} \quad (10)$$

with the reduced Planck length $l_p = \sqrt{8\pi G_N \hbar}$. Making the choice of representation that diagonalizes \hat{b} we can write its conjugate operator as,

$$\hat{a}^2 = -i\hbar \frac{\partial}{\partial b} \quad (11)$$

with $\hbar = l_P^2/3V_c$ the “effective Planck parameter” [23]. We now have ingredients to define a two-branch first order formulation of a quantum theory of gravity after combining Eqs. 9 and 11,

$$\left[-i\hbar h_\pm(b) \frac{\partial}{\partial b} - m^2 \right] \psi = 0, \quad (12)$$

which is a Wheeler-DeWitt equation for the connection representation. To generalize the Chern-Simons functional the following linearizing variable is defined,

$$X_\pm(b) = \int \frac{db}{h_\pm(b)}. \quad (13)$$

The general solution is a superposition of different valued constants $\alpha = (m^2, \phi)$ for spatial monochromatic functions $\psi_s(b; \alpha)$ which are normalized such that $|\psi_s|^2 = 1/(2\pi\hbar)^D$ and the total solution is of the form,

$$\psi(b, \mathbf{T}) = \int d\alpha \mathcal{A}(\alpha) \exp \left[-\frac{i}{\hbar} \alpha \cdot \mathbf{T} \right] \psi_s(b; \alpha), \quad (14)$$

where \mathbf{T} is the conjugate time to α and D is the number of conserved parameters α . Here ψ_s is made up of an incident and a reflected wave, associated with X_\pm .




FIG. 1: *Left:* The fractional change in the Hubble parameter from the base ΛCDM model, with best-fit Planck 2018 parameters from Eq. 20, for different parameter values of the quantum cosmology model. *Right:* Evolution of the expansion rate for cosmologies considered in this work along with various experimental constraints.

Since we would want to reproduce a semiclassical limit at late times, the choice for $\mathcal{A} = \sqrt{\mathcal{N}}(\alpha_{i0}, \sigma_{\alpha_i})$ corresponding to wave packets with Gaussian amplitudes

$$\mathcal{A}(\alpha_i) = \frac{1}{(2\pi\sigma_i^2)^{1/4}} \exp\left[-\frac{(\alpha_i - \alpha_{i0})^2}{4\sigma_i^2}\right] \quad (15)$$

leads to wave packets of the form,

$$\psi_{\pm i}(b, T_i) = \frac{1}{(2\pi\sigma_{T_i}^2)^{1/4}} \exp\left[-\frac{(X_{\pm i}^{\text{eff}}(b) - T_i)^2}{4\sigma_{T_i}^2}\right] \quad (16)$$

where $\sigma_{T_i} = \hbar/2\sigma_i$ and $X_{\pm i}^{\text{eff}}(b)$ is an effective functional adapted to wave packets in the saddle point approximation (see [1, 21, 22] for details). The probability for a given b at T_i in the semi-classical limit can be found to be

$$\mathcal{P}(b, T_i) = \left|\frac{dX_{+i}^{\text{eff}}}{db}\right| |\psi_+|^2 + \left|\frac{dX_{-i}^{\text{eff}}}{db}\right| |\psi_-|^2 \quad (17)$$

(see [1, 22] for more details; no interference terms appear in the semiclassical approximation). From this, one can then calculate modifications induced in the Hubble parameter as,

$$\bar{b}(T_\phi) = \int_{b_\Lambda}^{\infty} db \mathcal{P}(b, T_\phi) b, \quad (18)$$

where b_Λ is the value at dark energy-matter equality and the contribution to the Hubble parameter originating

from quantum effects is explicitly given as,

$$H_Q = \frac{\bar{b} - b}{a} \quad (19)$$

such that we can think of the Hubble parameter as the sum of the classical background solution and this new quantum addition: $H = H_C + H_Q$.

III. COSMOLOGICAL IMPRINTS

Deviations from the standard ΛCDM expansion history in the late-Universe will impact the growth of structure and can be constrained from observational probes ranging from measurements of the growth factor, $f\sigma_8$, and the late-ISW effect. However, before we can calculate quantum effects our first step is to calculate the standard, classical ΛCDM cosmology. For this we use the best-fit Planck 2018 parameters given by [24],

$$\begin{aligned} H_0 &= 67.32 \text{ km/s/Mpc} & 100\omega_b &= 2.283, \\ \omega_{\text{cdm}} &= 0.1201 & 10^9 A_s &= 2.210, \\ n_s &= 0.9661 & \tau_{\text{reio}} &= 0.0543, \end{aligned} \quad (20)$$

to calculate the cosmology with the Boltzmann code CLASS [25]. Separately, we then calculate the expected deviations in the expansion rate due to quantum effects in the very late-Universe for a range of values for σ_{T_ϕ} . Since the Λ clock becomes unreliable as we enter deeper into matter domination, we smoothly interpolate between the




FIG. 2: *Left:* Deviation in the growth factor from the base Λ CDM cosmology with projected constraints from DESI. *Right:* Evolution of $f\sigma_8$ for Λ CDM (dashed), with parameters from Eq. 20, and for the quantum cosmology model with differing values of σ_{T_ϕ} (solid). Over plotted are constraints on $f\sigma_8(z)$ from RSD measurements from galaxy surveys.

quantum effects at the bounce and the classical solution at higher redshifts, i.e. $H_Q \rightarrow 0$. This is to be expected since the relevant semi-classical wave function will correspond to the matter clock as we move to higher redshifts and away from the bounce.

In the left panel of Fig. 1 we show the evolution of the fractional change in the Hubble parameter for a set of different values of $\sigma_{T_\phi}/b_\Lambda^3$, while in the right panel of Fig. 1 we show the evolution of $H/(1+z) = \dot{a}$, i.e. the expansion rate, against direct BAO measurements from BOSS DR12 [26], BOSS DR14 quasars [27], and BOSS Ly- α [28] as well as the best fit H_0 values from SH0ES [29] and for Planck 2018 [24]. Notice that the deviation from the classical evolution in the Hubble parameter becomes more pronounced for larger values of $\sigma_{T_\phi}/b_\Lambda^3$, which also results in a characteristic flattening of the evolution of the expansion rate. This effect can be understood as directly stemming from the increase in variance for the wave packets which results in a more pronounced skewness of the wave function. In turn, this lends itself to a bias for larger values of the Hubble parameter and an impact on the cosmology for a larger range of redshifts around the bounce.

It is worth noting that the region of parameter space suggested in [1] that could alleviate the Hubble tension ($\sigma_{T_\phi}/b_\Lambda^3 \approx 0.30$) is not favored (due to the BOSS DR12 measurements).

A. Signatures in the matter power spectrum

Beyond modifications at the background level, there will also be an impact on the evolution of perturbations. Changes in the abundance and evolution of the various constituents of the Universe is readily encoded in the matter power spectrum. Current and future probes of the matter power spectrum can then be leveraged to distinguish between different models of dark matter, extensions of general relativity, and dark energy.

The growth of linear perturbations can be derived from the time evolution of the growth factor,

$$D''(\tau) = -aHD'(\tau) + \frac{3}{2}a^2\rho_M D(\tau), \quad (21)$$

where primes denote derivatives with respect to conformal time. To establish the sensitivity of the cosmology from varying σ_{T_ϕ} we have plotted the deviation in the growth factor from Λ CDM in Fig. 2. We also include the projected constraints for the Dark Energy Spectroscopic Instrument (DESI) [30] for changes in the growth factor relative to Λ CDM from the Bright Galaxy Survey (BGS) at very low redshifts, with the Main Survey (MS) around $z \approx 1$, and at higher redshifts with the Lyman- α survey (Ly- α). As we anticipate, before dark energy-matter equality the evolution is identical to the classical expectation until the bounce where the semi-classical limit breaks down and we see values of $\sigma_{T_\phi}/b_\Lambda^3 \gtrsim 0.10$ are likely to be constrained by DESI.

Another important probe of new physics is the growth

rate of perturbations, $f \equiv d \log D / d \log a$, which can be measured directly with supernova (but only for very low redshifts) and with galaxies surveys like BOSS through redshift-space distortions (RSD) [31]. For the latter, this stems from distortions along the line-of-sight in the distribution of galaxies (in redshift space) stemming from an induced peculiar velocity as they are drawn toward the center of potential wells. This can be measured from the galaxy power spectrum which can be decomposed as,

$$P_g(k, \mu_k, z) = P(k, z) (b_1 + f \mu_k^2)^2 + P_N, \quad (22)$$

where $P(k, z)$ is the standard linear matter power spectrum, b_1 is the galaxy bias, $\mu_k = \hat{\mathbf{e}}_z \cdot \hat{\mathbf{k}}$ the angle between the wavevector and line-of-sight, and P_N is shot noise. From Eq. 22 it is clear that its not possible to disentangle the growth rate from the matter power spectrum, thus the actual observable from galaxy surveys is the redshift dependant quantity $f \sigma_8$ where σ_8 can be thought of as the amplitude of the power spectrum but is defined as the RMS linear-theory *mass* fluctuation in a sphere of radius $8 h^{-1} \text{Mpc}$. In k-space this is calculated as an integral over the matter power spectrum,

$$\sigma_8(z)^2 = \int d \log k W(kR)^2 k^3 P(k, z), \quad (23)$$

where $W(kR)$ is a window function of scale $8 h^{-1} \text{Mpc}$.

In Fig. 2 we plot the evolution of $f \sigma_8(z)$ relative to constraints from RSD measurements from BOSS DR11 [32], WiggleZ [33], SDSS LRG [34], VVDS [35], VIPERS [36], FastSound [37], 2dFGRS [38], and 6dFGRS [39]. As we would expect from a model that results in an enhancement in the Hubble parameter, there is a corresponding suppression in the growth of structure relative to the baseline ΛCDM model. This is rather unique among models that make modifications in the gravitational sector as typically this leads to an enhancement in the growth rate, e.g. scalar-tensor theories. For the curves in Fig. 2 we calculate the χ^2 values for the plotted data points. For ΛCDM we find that $\chi^2 \sim 14$ and we find for $\sigma_{T_\phi}/b_\Lambda^3 = 0.05, 0.10, 0.20$ that $\chi^2 \sim 9, 7, 19$ respectively. This implies that there could be a region in parameter space that improves the fit between data and model for $0 \lesssim \sigma_{T_\phi}/b_\Lambda^3 \lesssim 0.20$ but more careful data analysis would be required to make a more definitive statement. Quantifying in terms of the S_8 parameter ($S_8 \equiv \sigma_8 \sqrt{\Omega_m/0.3}$) for $\sigma_{T_\phi}/b_\Lambda^3 = 0.05, 0.10, 0.20$, this results in a suppression $\sim 3\%, 5\%,$ and 12% respectively. Interestingly, the suppression in this region of parameter space is enough to help alleviate the S_8 tension [40].

B. Signatures in the CMB

While CMB power spectra encode rich information about the physics around recombination, it is also a probe of other processes at intervening epochs leading up to today. Relevant to the model considered here, changes




FIG. 3: CMB temperature power spectrum including contribution from the late-ISW effect for ΛCDM (dashed), with parameters from Eq. 20, and for the quantum cosmology model with differing values of σ_{T_ϕ} (solid). The gray band corresponds to cosmic variance.

in the growth of structure in the late-Universe can be probed via the late-ISW effect [41] defined as,

$$\Theta^{\text{ISW}}(\hat{\mathbf{n}}) = 2 \int_{\tau_{\text{dec}}}^{\tau_0} d\tau \Phi', \quad (24)$$

which is an integral from recombination to today over the rate of change of the potential and can serve as a sensitive probe of new physics beyond the ΛCDM model. The angular power spectrum for the late-ISW effect is given by,

$$C_\ell^{\text{ISW}} = \frac{8}{\pi} \int d \log k k^3 \left\langle \left| \int d\tau \frac{\partial \Phi_{\mathbf{k}}}{\partial \tau} j_\ell(k\tau) \right|^2 \right\rangle, \quad (25)$$

where we have transitioned to Fourier space and $j_\ell(k\tau)$ is the spherical Bessel function. The Poisson equation in k-space gives the Newtonian potential which, ignoring the velocity perturbations, can be written as

$$\Phi_{\mathbf{k}} = \frac{3H_0^2 \Omega_m D}{2k^2 a} \delta(k). \quad (26)$$

Eq. 26 can then be differentiated with respect to conformal time to get,

$$\frac{\partial \Phi_{\mathbf{k}}}{\partial \tau} = \frac{3H_0^2 \Omega_m}{2k^2} \delta(k) \left(\frac{\partial D}{\partial a} \dot{a} - \frac{D}{a} H \right), \quad (27)$$

and then be combined with Eqs. 25 and 27 to yield the expression of the full angular power spectrum,

$$C_\ell^{\text{ISW}} = \frac{18}{\pi} H_0^4 \Omega_m^2 \int \frac{d \log k}{k} P(k) \left[\int d\tau \mathcal{H} D (f-1) j_\ell(k\tau) \right]^2 \quad (28)$$

where $P(k)$ is the matter power spectrum. Having solved for the background cosmology and evolution of perturbations it is then straightforward to numerically calculate the late-ISW contribution to the CMB using Eq. 28.

In Fig. 3 we show the full temperature power spectrum for different values of σ_{T_ϕ} together with the base Λ CDM model. We see the impact of the quantum effects as an excess of power at low ℓ relative to Λ CDM. This behavior is easy to understand given the explicit dependence on the Hubble parameter in Eq. 28. However, from these results it is evident that the late-ISW effect in this model is not as constraining relative to the other observables as it is limited by cosmic variance – only values of $\sigma_{T_\phi}/b_\Lambda^3 \gtrsim 0.30$ are likely to be distinguishable from the standard concordance model prediction.

IV. DISCUSSION AND CONCLUSION

In this work we explored the cosmological signatures of a recently proposed model of quantum cosmology [1], where quantum uncertainties in the physical clock are particularly evident about the time of the transition to accelerated expansion. One of the features of this model that was identified in the original work is that there is an enhancement in the Hubble parameter around the transition from matter to dark energy domination. Extending the previous work, we have shown that the growth of structure can strongly constrain the parameter space of this model. In particular, future measurements of the growth factor by DESI will be able constrain the parameter space for values of $\sigma_{T_\phi}/b_\Lambda^3 \gtrsim 0.10$. Furthermore, the redshift evolution of $f\sigma_8$ measured from RSD will be a sensitive probe of modification to the cosmology in this theory. While this model can also be constrained by the late-ISW effect, the parameter space that can be probed is limited due to cosmic variance.

In [1] it was suggested that the model of quantum cosmology considered here could potentially address the Hubble tension. While we haven't explored this suggestion rigorously through an MCMC analysis, it seems unlikely given current data that this can be accommodated

by the parameter space. Indeed, it is well known that late-time resolutions to the Hubble tension tend to be very well constrained relative to early-Universe modifications - namely those which work on the basis of injecting energy density before recombination to modify the size of the sound horizon [42]. That said, this motivates exploring the possible quantum effects at a transition between matter and radiation domination. Indeed, one of the “unsettling” features of models like early dark energy [43] (that are quite successful at addressing the Hubble tension) is their why now problem, i.e. why do they turn on at matter-radiation equality. It would be interesting to study if quantum effects in a model like the one studied here could work as an early Universe solution to the Hubble tension – though since there will be no reflection in the connection, it is not clear that it will be a nearly as strong of an effect as observed here. At most there will be a refraction/diffraction of the wave.

While the prospects for the model considered here to address the H_0 tension are bleak, it does show some promise for potentially alleviating another growing tension in cosmology, the S_8 tension. A particularly interesting feature of this tension is that low-redshift probes ($z \leq 1$) of the clustering of galaxies is lower than that inferred from the CMB and CMB lensing which probes from today to recombination [44]. Given that this period is around the same time that dark energy becomes dominate in the Universe, it would be interesting if the observed suppression was somehow connected. Indeed, in the context of the model considered here there is no *why now?* problem, i.e. the suppression in structure growth is a direct consequence of the dark energy-matter transition. Recently, some success was seen with a similar idea [45] where the transition triggers new dynamics in the dark sector to alter the growth of structure and alleviate the tension. We leave further study of the implications of this model for the growth of LSS and the possibility of alleviating the Hubble tension with similar dynamics to those studied here but at matter-radiation equality for future work.

V. ACKNOWLEDGEMENTS

We thank Stephon Alexander and Marc Kamionkowski for useful conversations. SMK is partially supported by NSF PHY-2014052.

-
- [1] Bruno Alexandre and Joao Magueijo. Possible quantum effects at the transition from cosmological deceleration to acceleration. *Phys. Rev. D*, 106(6):063520, 2022.
 - [2] W. G. Unruh. Unimodular theory of canonical quantum gravity. *Phys. Rev. D*, 40:1048–1052, Aug 1989.
 - [3] Lee Smolin. The Quantization of unimodular gravity

- and the cosmological constant problems. *Phys. Rev. D*, 80:084003, 2009.
- [4] Karel V. Kuchař. Does an unspecified cosmological constant solve the problem of time in quantum gravity? *Phys. Rev. D*, 43:3332–3344, May 1991.
- [5] Alan Daughton, Jorma Louko, and Rafael D. Sorkin. Ini-

- tial conditions and unitarity in unimodular quantum cosmology. *arXiv: General Relativity and Quantum Cosmology*, 1993.
- [6] R.D. Sorkin. Role of time in the sum-over-histories framework for gravity. *Int J Theor Phys*, 33:523–534, 1994.
- [7] R.D. Sorkin. Forks in the road, on the way to quantum gravity. *Int J Theor Phys*, 36:2759–2781, 1997.
- [8] Marc Henneaux and Claudio Teitelboim. The cosmological constant and general covariance. *Physics Letters B*, 222(2):195–199, 1989.
- [9] Luca Bombelli, W. E. Couch, and R. J. Torrence. Time as spacetime four-volume and the ashtekar variables. *Phys. Rev. D*, 44:2589–2592, Oct 1991.
- [10] L. Smolin. Unimodular loop quantum gravity and the problems of time. *Phys Rev D*, 84:044047, 2011.
- [11] Edward Witten. A Note on the Chern-Simons and Kodama wave functions. 6 2003.
- [12] Stephon Alexander, Laurent Freidel, and Gabriel Herzceg. An Inner Product for 4D Quantum Gravity and the Chern-Simons-Kodama State. 12 2022.
- [13] C. J. Isham. Canonical quantum gravity and the problem of time. *NATO Sci. Ser. C*, 409:157–287, 1993.
- [14] K. V. Kuchar. Time and interpretations of quantum gravity. *Int. J. Mod. Phys. D*, 20:3–86, 2011.
- [15] J. David Brown. Action functionals for relativistic perfect fluids. *Class. Quant. Grav.*, 10:1579–1606, 1993.
- [16] J. David Brown. Tunneling in perfect-fluid (minisuper-space) quantum cosmology. *Phys. Rev. D*, 41:1125–1141, Feb 1990.
- [17] Steffen Gielen and Neil Turok. Quantum propagation across cosmological singularities. *Phys. Rev. D*, 95(10):103510, 2017.
- [18] Steffen Gielen and Lucía Menéndez-Pidal. Singularity resolution depends on the clock. *Class. Quant. Grav.*, 37(20):205018, 2020.
- [19] Steffen Gielen and Lucía Menéndez-Pidal. Unitarity, clock dependence and quantum recollapse in quantum cosmology. *Class. Quant. Grav.*, 39(7):075011, 2022.
- [20] João Magueijo. Cosmological time and the constants of nature. *Physics Letters B*, 820:136487, 2021.
- [21] Joao Magueijo. Connection between cosmological time and the constants of nature. *Phys. Rev. D*, 106(8):084021, 2022.
- [22] Steffen Gielen and João Magueijo. Quantum analysis of the recent cosmological bounce in comoving Hubble length. 1 2022.
- [23] John D. Barrow and Joao Magueijo. A contextual Planck parameter and the classical limit in quantum cosmology. *Found. Phys.*, 51(1):22, 2021.
- [24] N. Aghanim et al. Planck 2018 results. I. Overview and the cosmological legacy of Planck. *Astron. Astrophys.*, 641:A1, 2020.
- [25] Diego Blas, Julien Lesgourgues, and Thomas Tram. The Cosmic Linear Anisotropy Solving System (CLASS). Part II: Approximation schemes. *JCAP*, 2011(7):034, July 2011.
- [26] Shadab Alam et al. The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample. *Mon. Not. Roy. Astron. Soc.*, 470(3):2617–2652, 2017.
- [27] Pauline Zarrouk et al. The clustering of the SDSS-IV extended Baryon Oscillation Spectroscopic Survey DR14 quasar sample: measurement of the growth rate of structure from the anisotropic correlation function between redshift 0.8 and 2.2. *Mon. Not. Roy. Astron. Soc.*, 477(2):1639–1663, 2018.
- [28] Julian E. Bautista et al. Measurement of baryon acoustic oscillation correlations at $z = 2.3$ with SDSS DR12 Ly α -Forests. *Astron. Astrophys.*, 603:A12, 2017.
- [29] Adam G. Riess et al. A Comprehensive Measurement of the Local Value of the Hubble Constant with 1 km s $^{-1}$ Mpc $^{-1}$ Uncertainty from the Hubble Space Telescope and the SH0ES Team. *Astrophys. J. Lett.*, 934(1):L7, 2022.
- [30] Amir Aghamousa et al. The DESI Experiment Part I: Science, Targeting, and Survey Design. 10 2016.
- [31] Nick Kaiser. Clustering in real space and in redshift space. *Monthly Notices of the Royal Astronomical Society*, 227(1):1–21, 07 1987.
- [32] Lado Samushia, Beth A. Reid, Martin White, Will J. Percival, Antonio J. Cuesta, Lucas Lombriser, Marc Manera, Robert C. Nichol, Donald P. Schneider, Dmitry Bizyaev, Howard Brewington, Elena Malanushenko, Viktor Malanushenko, Daniel Oravetz, Kaike Pan, Audrey Simmons, Alaina Shelden, Stephanie Snedden, Jeremy L. Tinker, Benjamin A. Weaver, Donald G. York, and Gong-Bo Zhao. The clustering of galaxies in the SDSS-III DR9 Baryon Oscillation Spectroscopic Survey: testing deviations from Λ and general relativity using anisotropic clustering of galaxies. *MNRAS*, 429(2):1514–1528, February 2013.
- [33] Chris Blake, Sarah Brough, Matthew Colless, Carlos Contreras, Warrick Couch, Scott Croom, Tamara Davis, Michael J. Drinkwater, Karl Forster, David Gilbank, Mike Gladders, Karl Glazebrook, Ben Jelliffe, Russell J. Jurek, I. Hui Li, Barry Madore, D. Christopher Martin, Kevin Pimblett, Gregory B. Poole, Michael Pracy, Rob Sharp, Emily Wisnioski, David Woods, Ted K. Wyder, and H. K. C. Yee. The WiggleZ Dark Energy Survey: the growth rate of cosmic structure since redshift $z=0.9$. *MNRAS*, 415(3):2876–2891, August 2011.
- [34] L. Samushia, W. J. Percival, and A. Raccanelli. Interpreting large-scale redshift-space distortion measurements. *MNRAS*, 420(3):2102–2119, March 2012.
- [35] L. Guzzo et al. A test of the nature of cosmic acceleration using galaxy redshift distortions. *Nature*, 451:541–545, 2008.
- [36] S. de la Torre et al. The VIMOS Public Extragalactic Redshift Survey (VIPERS). Galaxy clustering and redshift-space distortions at $z=0.8$ in the first data release. *Astron. Astrophys.*, 557:A54, 2013.
- [37] Teppei Okumura et al. The Subaru FMOS galaxy redshift survey (FastSound). IV. New constraint on gravity theory from redshift space distortions at $z \sim 1.4$. *Publ. Astron. Soc. Jap.*, 68(3):38, 2016.
- [38] Will J. Percival et al. The 2dF Galaxy Redshift Survey: Spherical harmonics analysis of fluctuations in the final catalogue. *Mon. Not. Roy. Astron. Soc.*, 353:1201, 2004.
- [39] Florian Beutler, Chris Blake, Matthew Colless, D. Heath Jones, Lister Staveley-Smith, Gregory B. Poole, Lachlan Campbell, Quentin Parker, Will Saunders, and Fred Watson. The 6dF Galaxy Survey: $z \approx 0$ measurements of the growth rate and σ_8 . *MNRAS*, 423(4):3430–3444, July 2012.
- [40] Elcio Abdalla et al. Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies. *JHEAp*, 34:49–211, 2022.
- [41] Wayne Hu and Scott Dodelson. Cosmic Microwave Back-

- ground Anisotropies. *Ann. Rev. Astron. Astrophys.*, 40:171–216, 2002.
- [42] Marc Kamionkowski and Adam G. Riess. The Hubble Tension and Early Dark Energy. 11 2022.
- [43] Tanvi Karwal and Marc Kamionkowski. Dark energy at early times, the Hubble parameter, and the string axiverse. *Phys. Rev. D*, 94(10):103523, 2016.
- [44] Shi-Fan Chen, Martin White, Joseph DeRose, and Nickolas Kokron. Cosmological analysis of three-dimensional BOSS galaxy clustering and Planck CMB lensing cross correlations via Lagrangian perturbation theory. *JCAP*, 07(07):041, 2022.
- [45] Vivian Poulin, José Luis Bernal, Ely Kovetz, and Marc Kamionkowski. The Sigma-8 Tension is a Drag. 9 2022.