




Phase space analysis of the bouncing universe with stringy effects

Alfredo D. Millano ^{1,*}, Kimet Jusufi ^{2,†} and Genly Leon ^{1,3‡}

¹*Departamento de Matemáticas, Universidad Católica del Norte,
Avda. Angamos 0610, Casilla 1280 Antofagasta, Chile*

²*Physics Department, State University of Tetovo,
Ilinden Street nn, 1200, Tetovo, North Macedonia and*

³*Institute of Systems Science, Durban University of Technology,
PO Box 1334, Durban 4000, South Africa*

Abstract

We use the recently modified Friedmann equations obtained from string T-duality effects that encode the zero-point length (Phys. Lett. B 836 (2023), 137621) to study the phase space analyses of a bouncing early universe. An important implication of such stringy effects is that they can alleviate the initial singularity since the Raychaudhuri equation is modified. We investigate if the Universe can undergo an accelerated expansion phase for a specific domain of the equation of state parameter. The stringy effects are encoded in the parameter Γ , which depends not only on the zero-point length but also on the state parameter ω . We construct two dynamical systems depending on whether $-1 < \omega \leq -1/3$ and $\omega \geq -1/3$, and we classify the equilibrium points of each system. Exact solutions and cosmological implications are discussed.

*Electronic address: alfredo.millano@alumnos.ucn.cl

†Electronic address: kimet.jusufi@unite.edu.mk

‡Electronic address: genly.leon@ucn.cl

I. INTRODUCTION

Although the tremendous success of the cosmological Λ CDM model in describing the current state and evolution of the Universe, the deep origin of the accelerating expansion of the Universe still needs to be well understood [1–3]. We often refer to the cosmological constant as dark energy - a hypothetical source of energy permeating the Universe and exerting a negative pressure, causing, in this way, the expansion of the Universe to accelerate [4]. Nevertheless, dark matter is another mysterious form of matter revealed by different observations. There are a lot of observational facts that support the presence of dark matter and dark energy. It is well known that dark matter is needed to explain the galaxy rotation curves. In particular, one can use the rotation of stars in galaxies to infer the presence of matter/dark matter in a given galaxy. In this direction, the astrophysical measurements of the rotation of stars have revealed the presence of dark matter [5]. Another essential fact is cosmic microwave background (CMB) radiation. The CMB is thought to be remnant radiation of the early Universe; it is characterised by highly uniform temperature, but, importantly, there are slight variations in temperature, which can be used to infer many properties of the early universe [6–9]. Let us mention here the observational fact related to the Type Ia supernovae. We can use this explosive death of stars to measure the distance to distant galaxies and discover that the Universe is expanding. That, again, implies the role of the cosmological constant in explaining the expansion of the Universe. Among other things, dark matter and dark energy are needed to explain the Universe’s large-scale structure. Other findings in modern cosmology, like the baryon acoustic oscillations, have revealed further arguments supporting the existence of dark matter and dark energy [10]. The variations in the density of matter in the early Universe can infer the properties of the Universe. Finally, we can use general relativistic effects like the gravitational lensing effect to infer the distribution of matter in large-scale structures. The Λ CDM model fits very well with the above observational facts.

Other attempts to explain the Universe’s accelerated expansion in terms of the quintessence field have been proposed [11–15]. Unlike the cosmological constant, which is assumed to be a constant amount of energy density everywhere, quintessence is a dynamic field and, therefore, can vary with time. Another possibility to explain the acceleration rate of the Universe might be the modified gravity theories (see [16–74] and references therein).

According to general relativity, our Universe started with the Big Bang, a cosmological singularity that can be understood in the Raychaudhuri equation. If we study the evolution of the Universe, one can show that the expansion scalar can become infinite at some point, indicating in this way the presence of the singularity. However, we now believe that such a singularity can be avoided in these extreme conditions if we consider quantum effects.

In the 1970s, Hawking and Bekenstein showed a profound connection between black hole physics and the law of thermodynamics. In particular, it was argued that thermodynamic quantities such as entropy and temperature are proportional to the horizon area and surface gravity, which are pure geometrical quantities. An exciting finding was achieved by Jacobson, who derived the Einstein field equation of gravity as the equation of state for spacetime [75]. This idea was further used in several studies; see in particular ([76–89]). Recently, Verlinde argued that gravity is not a fundamental force but can be regarded as an entropic force [90]. In Verlinde’s gravity, gravity is an emergent effect, and dark matter is an apparent effect; an entropy displacement caused by matter [91]. Verlinde’s argument is based on two important principles. The first principle is the equipartition law of energy for the system’s degrees of freedom, and the second is the holographic principle. In this line of thought, Padmanabhan made further steps to understand the nature of gravity. He argued that the spatial expansion of our Universe could be understood as a consequence of the emergence of space. By equating the difference between the number of degrees of freedom in bulk and on the boundary with the volume change, he was able to obtain the Friedmann equation describing the evolution of the Universe [92, 93] (see also [94–96]).

Padmanabhan introduced the concept of duality and zero point length as a tool to obtain quantum gravity effect [97] and was applied to black hole physics to cure the black hole singularity [98–101]. In the present work, we will study the phase space analyses of the recently modified Friedmann’s equation found in [102]. The modified Friedmann’s was obtained from Verlinde’s emergent gravity with zero-point length corrections. The quantum effect is essential in the early Universe; in particular, it was shown how the singularity could be avoided, and the Universe can perform a bouncing scenario [102].

This paper is structured as follows. In Sec. II, we review the modified Friedmann’s equation in string T-duality derived from Verlinde’s entropic force scenario. In Sec. III, we investigate the phase space analyses of the early Universe. Finally, we comment on our findings in Sec. IV.

II. MODIFIED FRIEDMANN EQUATIONS WITH STRINGY EFFECTS

First, we review the modified Friedmann's equation reported in [102]. The gravitational potential is modified, and, consequently, Newton's law reads [102]

$$F = -\frac{GMm}{R^2} \left[1 + \frac{l_0^2}{R^2} \right]^{-3/2}, \quad (1)$$

here l_0 is the zero-point length with a value close to the Planck length (see, [98]). We shall assume a spatially homogeneous and isotropic background described by the Friedmann-Robertson-Walker (FRW) metric

$$ds^2 = h_{\mu\nu} dx^\mu dx^\nu + R^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (2)$$

in which $R = a(t)r$, $x^0 = t$, $x^1 = r$. For the metric functions, we have

$$h_{\mu\nu} = \text{diag} \left(-1, \frac{a^2}{1 - kr^2} \right), \quad (3)$$

with k being the the curvature of space; in particular, for $k = 1, 0, -1$, we have a closed, flat, and open FLRW model, respectively. For the FRW metric, we can find the dynamical apparent horizon, by the following equation

$$h^{\mu\nu} \partial_\mu R \partial_\nu R = 0. \quad (4)$$

It is easy to show that the apparent horizon radius for the FRW Universe reads

$$R = ar = \frac{1}{\sqrt{H^2 + k/a^2}}. \quad (5)$$

In addition, for the matter source in the FRW Universe, we shall assume a perfect fluid described in terms of the stress-energy tensor as follows

$$T_{\mu\nu} = (\rho + p)u_\mu u_\nu + pg_{\mu\nu}, \quad (6)$$

along with the continuity equation

$$\dot{\rho} + 3H(\rho + p) = 0. \quad (7)$$

The last equation $H = \dot{a}/a$ defines the Hubble parameter. First, we can derive the dynamical equation and, for this goal, let us consider a compact spatial region V with a compact

boundary \mathcal{S} , namely we have a sphere with a radius $R = a(t)r$, where r is a dimensionless quantity. By combining the second law of Newton with the gravitational force (1), we get

$$F = m\ddot{R} = m\ddot{a}r = -\frac{GMm}{R^2} \left[1 + \frac{l_0^2}{R^2}\right]^{-3/2}. \quad (8)$$

Next, to derive the Friedmann equations of the FRW Universe, we need to use \mathcal{M} rather than the total mass M . By replacing M with \mathcal{M} , it follows [102]

$$\mathcal{M} = -\frac{\ddot{a}a^2}{G} r^3 \left[1 + \frac{l_0^2}{R^2}\right]^{3/2} \quad (9)$$

and for the gravitational mass, we have

$$\mathcal{M} = 2 \int_V dV \left(T_{\mu\nu} - \frac{1}{2} T g_{\mu\nu} \right) u^\mu u^\nu. \quad (10)$$

From these equations, it is easy to obtain the modified dynamical evolution of the FRW universe [102]

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} (\rho + 3p) \left[1 + \frac{l_0^2}{R^2}\right]^{-3/2}. \quad (11)$$

Furthermore, since l_0 is a very small number, we can consider a series expansion around l_0 [102]

$$\frac{\ddot{a}}{a} = -\left(\frac{4\pi G}{3}\right) (\rho + 3p) \left[1 - \frac{3}{2} \frac{l_0^2}{r^2 a^2} + \dots\right], \quad (12)$$

then by multiplying $2\dot{a}a$ on both sides of Eq. (11), and making use of the continuity equation

$$\rho = \rho_0 a^{-3(1+w)}, \quad (13)$$

it was found that the modified Friedmann's and Raychaudhuri's equations [102]

$$H^2 + \frac{k}{a^2} = \frac{8\pi G}{3} \rho [1 - \Gamma\rho], \quad (14)$$

and

$$\dot{H} - \frac{k}{a^2} = -4\pi G(\rho + p)(1 - 2\Gamma\rho), \quad (15)$$

respectively, with $p = \omega\rho$ and Γ defined by

$$\Gamma \equiv \frac{4\pi G l_0^2}{3} \left(\frac{1 + 3\omega}{1 + \omega}\right). \quad (16)$$

For $\Gamma > 0$, one can discover that the string T-duality modified Friedmann equations have a similar form with the modified Friedmann's equation found in Loop quantum gravity (LQG) [112]

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G\rho}{3} \left[1 - \frac{\rho}{\rho_c}\right], \quad (17)$$

where ρ_c is the critical energy density defined by

$$\rho_c \equiv \frac{3}{8\pi\gamma^2\lambda^2}, \quad (18)$$

with $\lambda \sim 5.2l_{Pl}^2$ [112] being the area gap that sets the discreteness scale of Loop quantum gravity, and γ is the Immirzi parameter. The correspondence is achieved by identifying $\Gamma = \rho_c^{-1}$ [101]. One difference, however, is that the Γ depends not only on the zero-point length but also on the state parameter ω . An interesting result can be found if we set $\dot{a} = 0$ ($H = 0$); in that case, we can solve the equation for the critical density we get [102]

$$\rho_c = \frac{1}{2\Gamma} \left(1 \pm \frac{\sqrt{2G\pi(2G\pi a^2 - 2\Gamma k)}}{2G\pi a} \right). \quad (19)$$

The last equation imposes the constraint $2G\pi a^2 - 2\Gamma k \geq 0$. Solving for the scale factor $a = a_{\min}$, we get

$$a_{\min} = \sqrt{2} l_0 \sqrt{\frac{(1 + 3\omega) k}{1 + \omega}}. \quad (20)$$

This equation shows that for a closed universe, there exists a minimal length for the scale factor and, in particular, for $\omega = 0$, we get $a_{\min} = \sqrt{2} l_0$. For a flat universe with $k = 0$, the condition $\dot{a} = 0$ ($H = 0$) gives $\rho_c = 1/\Gamma$.

On the other hand, when $\Gamma < 0$, the modified Friedmann equation corresponds to the one of Brane-cosmology [113–119], particularly, on the RS2 scenario with a single Z_2 symmetric brane [113]. Eqs. (656) and (657) in [119] are

$$H^2 + \frac{k}{a^2} = \Lambda_4 + \frac{8\pi G_4}{3} \rho \left(1 + \frac{\rho}{2\sigma} \right) + \frac{\mu}{a^4}, \quad (21)$$

$$\dot{H} - \frac{k}{a^2} = -4\pi G(\rho + p) \left(1 + \frac{\rho}{\sigma} \right) - 2\frac{\mu}{a^4}, \quad (22)$$

where H the Hubble parameter along the brane, a is the scale factor, and $k = 0, \pm 1$ describes the spatial curvature. A tension, σ source the brane, and a cosmological fluid with energy density, ρ , and pressure, $p = \omega\rho$. The parameters Λ_4 and G_4 denote the effective cosmological constant and Newton's constant on the brane, respectively.

To compare with equations (14) and (15), we set the dark radiation term to zero and the 4-dimensional cosmological constant to zero $\Lambda_4 = 0$. The brane tension is reinterpreted as $\sigma = 1/(2|\Gamma|)$.

An interesting result can be found if we set $\dot{a} = 0$ ($H = 0$); in that case, we can solve the equation for the brane-tension to get

$$\sigma_c = -\frac{1}{4\Gamma} \left(1 \pm \frac{\sqrt{2G\pi(2G\pi a^2 - 2\Gamma k)}}{2G\pi a} \right). \quad (23)$$

The last equation imposes the constraint $2G\pi a^2 - 2\Gamma k \geq 0$. Solving for the scale factor $a = a_{\min}$, we get

$$a_{\min} = \sqrt{2} l_0 \sqrt{\frac{(1+3\omega)k}{1+\omega}}, \quad \sigma|_{a_{\min}} = -\frac{1}{4\Gamma}. \quad (24)$$

This equation shows that for a closed universe, there exists a minimal length for the scale factor and, in particular, for $\omega = 0$, we get $a_{\min} = \sqrt{2} l_0$. For a flat universe with $k = 0$, the condition $\dot{a} = 0$ ($H = 0$) gives $\sigma = 1/(2|\Gamma|)$.

Following (23) for an open Universe with $k = -1$, we can get the critical brane-tension provided $\Gamma \leq 0$. Since l_0 is of Planck length order, i.e., $l_0 \sim l_{Pl}$, its effect becomes important only in short distances. The critical density that corresponds to the minimal scale goes like $\sigma_c \sim l_{Pl}^2$, and it may prove a bound for the maximum brane-tension in nature.

In the present work, we consider the interval $\omega \in [-1, 1]$ as the physical one. According to (16), we have two regimes

$$\begin{cases} \Gamma \leq 0, & -1 < \omega \leq -\frac{1}{3}, & \text{Brane-like scenario} \\ \Gamma \geq 0, & \omega \geq -\frac{1}{3}, & \text{LQG-like scenario} \end{cases}. \quad (25)$$

We see that in the limit $l_0 \rightarrow 0$, the standard Friedmann equation is obtained. Furthermore, we can derive the modified Raychaudhuri equation using

$$\dot{H} = -H^2 + \frac{\ddot{a}}{a}. \quad (26)$$

and the deceleration parameter is defined as

$$q = -1 - \frac{\dot{H}}{H^2}, \quad (27)$$

where depending on the sign of q we can have deceleration or acceleration scenario, respectively. As shown in [101], the modified Friedmann's equations have interesting implications. Take for simplicity a flat universe with $k = 0$ along with the definition $H_0^2 = 8\pi G\rho/3$ and can write [101]

$$H \simeq H_0 (1 - \Delta_{GUP} H_0^2), \quad (28)$$

which has the form of Generalized Uncertainty Principle (GUP) modified Hubble parameter with $\Delta_{GUP} = \alpha l_0^2/2$ [105–109]. Since we believe that Cosmic Microwave Background (CMB) carries quantum gravity fingerprints, this equation can have implications for the Hubble tension problem. For H , we can take the value reported by the Planck collaboration, $H = 67.40 \pm 0.50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [103]. On the other hand, H_0 can be viewed as the unmodified Hubble parameter, and we can take the value reported by the Hubble Space Telescope (HST), $H_0 = 74.03 \pm 1.42 \text{ km s}^{-1} \text{ Mpc}^{-1}$ [104], see also [110, 111].

III. PHASE SPACE ANALYSIS

Now we present the evolution equations

$$\dot{H} = \frac{k}{a^2} + 4\pi G(\omega + 1)\rho(2\Gamma\rho - 1), \quad (29)$$

$$\dot{\rho} = -3H(1 + \omega)\rho. \quad (30)$$

as an autonomous dynamical system. We construct two dynamical systems depending on whether $-1 < \omega \leq -1/3$ and $\omega \geq -1/3$, and we classify the equilibrium points of each system, and cosmological implications are discussed.

Firstly, we find exact solutions and find qualitative results. Exact solutions can play an important role in the evolution of whole classes of models acting as asymptotic or intermediate stages in the evolution of the Universe. Additionally, one can use qualitative techniques of dynamical systems to obtain relevant information about the flow properties associated with an autonomous system of ordinary differential equations. For example, the qualitative theory of differential equations [122–124] has some applications in cosmology [125–127].

Notice that, by substituting (13) and $H = \dot{a}/a$ in (29) we obtain

$$-a\ddot{a} + \dot{a}^2 - 4\pi\rho_0 G(\omega + 1)a^{-6\omega-4} (a^{3\omega+3} - 2\Gamma\rho_0) + k = 0, \quad (31)$$

which admits the first integral

$$t - t_U = \int_1^a \left(\frac{8}{3} G\pi\rho_0 (s^{3\omega+3} - \Gamma\rho_0) s^{-6\omega-4} + c_1 s^2 - k \right)^{-\frac{1}{2}} ds \quad (32)$$

where we have denoted t_U the Universe's age, and we set $a(t_U) = 1$ for the current value of the scale factor and ρ_0 for the current value of the energy density. The integration constant

c_1 satisfies

$$c_1 = \dot{a}^2(t_U) + \frac{8}{3}\pi G\rho_0(\Gamma\rho_0 - 1) + k. \quad (33)$$

Therefore, the system is integrable.

Now we are interested in the asymptotic behaviour of the solutions. We consider two separate cases according to the sign of Γ .

A. $\Gamma \leq 0$

This regime corresponds to $-1 < \omega \leq -\frac{1}{3}$. Then, the Friedmann equation (14) becomes

$$H^2 + \frac{k}{a^2} = \frac{8\pi G}{3}\rho + \frac{8\pi G}{3}|\Gamma|\rho^2, \quad (34)$$

where for the closed Universe, we set $k = +1$.

The quantities in the left-hand and right-hand sides of (34) are non-negative. Then, we can define the dimensionless variables

$$\Omega = \frac{8\pi G}{3D^2}\rho, \quad \Omega_k = \frac{k}{a^2D^2}, \quad Q = \frac{H}{D}, \quad D = \sqrt{H^2 + \frac{k}{a^2}}, \quad (35)$$

where $(\Omega, \Omega_k, Q) \in [0, 1] \times [0, 1] \times [-1, 1]$, and satisfy

$$\frac{3|\Gamma|\Omega^2H^2}{8\pi G} = (1 - \Omega)(1 - \Omega_k), \quad (36)$$

$$Q^2 + \Omega_k = 1. \quad (37)$$

Introducing the time derivative

$$\frac{df}{d\tau} = \frac{1}{\sqrt{H^2 + ka^{-2}}} \dot{f}, \quad (38)$$

we obtain the dynamical system

$$\frac{d\Omega_k}{d\tau} = Q\Omega_k(-3\omega(\Omega - 2) - 3\Omega + 4), \quad (39)$$

$$\frac{d\Omega}{d\tau} = -3Q(\omega + 1)(\Omega - 1)\Omega, \quad (40)$$

$$\frac{dQ}{d\tau} = \frac{1}{2}\Omega_k(3\omega(\Omega - 2) + 3\Omega - 4). \quad (41)$$

Using the equation (37) as a definition of Ω_k we obtain the reduced dynamical system

$$\frac{d\Omega}{d\tau} = -3Q(\omega + 1)(\Omega - 1)\Omega, \quad (42)$$

$$\frac{dQ}{d\tau} = \frac{1}{2}(1 - Q^2)(3\omega(\Omega - 2) + 3\Omega - 4). \quad (43)$$

defined on the phase plane

$$\{(\Omega, Q) \in [0, 1] \times [-1, 1]\}. \quad (44)$$

The system system (42) and (43) admits the first integral

$$Q(\tau)^2 - 1 = e^{-\frac{c_1}{3(\omega+1)}} (1 - \Omega(\tau))^{-\frac{3\omega+1}{3(\omega+1)}} \Omega(\tau)^{\frac{2(3\omega+2)}{3(\omega+1)}}, \quad (45)$$

where c_1 is an integration constant.

Then, we obtain from (42) the equation

$$\Omega'(\tau) = -3(\omega + 1)\epsilon(\Omega(\tau) - 1)\Omega(\tau)\sqrt{1 + e^{-\frac{c_1}{3\omega+3}}(1 - \Omega(\tau))^{\frac{2}{3(\omega+1)}-1}\Omega(\tau)^{2-\frac{2}{3(\omega+1)}}}, \quad (46)$$

where $\epsilon = \pm 1$ is the sign of the initial value $H(0)$. Finally, we have the quadrature:

$$\tau = \pm \frac{c_2 - \int_1^\Omega \frac{1}{(\xi-1)\xi\sqrt{e^{-\frac{c_1}{3\omega+3}}(1-\xi)^{\frac{2}{3(\omega+1)}-1}\xi^{2-\frac{2}{3(\omega+1)}}+1}} d\xi}{3(\omega + 1)}, \quad (47)$$

where c_2 is an integration constant.

Equation (45) suggests to define the function

$$Z(\Omega, Q) := (1 - \Omega)^{\frac{3\omega+1}{3(\omega+1)}} \Omega^{-\frac{2(3\omega+2)}{3(\omega+1)}} [1 - Q^2], \quad (48)$$

which satisfies $Z' = 0$, implies $Z(\Omega, Q) = c$ where c is a constant. From this equation, it follows that asymptotic solutions satisfy $\Omega \rightarrow 1$, or $\Omega \rightarrow 0$ or $Q^2 \rightarrow 1$.

The equilibrium points of system (42) and (43) for the parameter region $-1 < \omega \leq -\frac{1}{3}$ are presented in Tab. I

The equilibrium points are the following.

P_1 : $(\Omega, Q) = (0, -1)$, always exists. It corresponds to a flat contracting solution that is a saddle for $-1 < \omega < -\frac{2}{3}$ or a sink for $-\frac{2}{3} < \omega < -\frac{1}{3}$.

P_2 : $(\Omega, Q) = (0, 1)$, always exists. It corresponds to an expanding flat solution that is a saddle for $-1 < \omega < -\frac{2}{3}$ or a source for $-\frac{2}{3} < \omega < -\frac{1}{3}$.

P_3 : $(\Omega, Q) = (1, -1)$, always exists. It corresponds to a flat contracting matter-dominated solution that is a source for $-1 < \omega < -\frac{1}{3}$, and it mimics a quintessence fluid.

P_4 : $(\Omega, Q) = (1, 1)$, always exists. It corresponds to a flat expanding matter-dominated solution that is a sink for $-1 < \omega < -\frac{1}{3}$, and it mimics a quintessence fluid.

P_5 : $(\Omega, Q) = \left(2 - \frac{2}{3(\omega+1)}, 0\right)$ exists for $-\frac{2}{3} < \omega \leq -\frac{1}{3}$. It corresponds to the Einstein static universe, and it is a saddle.

Label	Ω	Q	Existence	k_1	k_2	Stability
P_1	0	-1	always	$-3(\omega + 1)$	$-2(3\omega + 2)$	sink for $-\frac{2}{3} < \omega < -\frac{1}{3}$ saddle for $-1 < \omega < -\frac{2}{3}$
P_2	0	1	always	$3(\omega + 1)$	$2(3\omega + 2)$	source for $-\frac{2}{3} < \omega < -\frac{1}{3}$ saddle for $-1 < \omega < -\frac{2}{3}$
P_3	1	-1	always	$-(3\omega + 1)$	$3(\omega + 1)$	source for $-1 < \omega < -\frac{1}{3}$
P_4	1	1	always	$3\omega + 1$	$-3(\omega + 1)$	sink for $-1 < \omega < -\frac{1}{3}$
P_5	$2 - \frac{2}{3(\omega+1)}$	0	$-\frac{2}{3} < \omega \leq -\frac{1}{3}$	$-\sqrt{-(3\omega + 1)(3\omega + 2)}$	$\sqrt{-(3\omega + 1)(3\omega + 2)}$	saddle

TABLE I: Equilibrium points of system (42) and (43) for the parameter region $-1 < \omega \leq -\frac{1}{3}$.

Finally, for $\omega = -\frac{2}{3}$ we have the line of equilibrium points $\Omega = 0$, $-1 \leq Q \leq 1$ with eigenvalues $\{0, Q\}$ this line is stable for $Q < 0$ and unstable for $Q > 0$. For $\omega = -\frac{1}{3}$ we have the line of equilibrium points $\Omega = 1$, $-1 \leq Q \leq 1$ with eigenvalues $\{0, -2Q\}$ this line is stable for $Q > 0$ and unstable for $Q < 0$.

B. $\Gamma \geq 0$

This regime corresponds to $\omega \geq -\frac{1}{3}$. Then, the Friedmann equation (14) becomes

$$H^2 + \frac{k}{a^2} + \frac{8\pi G}{3}\Gamma\rho^2 = \frac{8\pi G}{3}\rho, \quad (49)$$

where for the closed Universe, we set $k = +1$. The evolution equations are (29) and (30).

The quantities in the left-hand and right-hand sides of (49) are non-negative. Then, we can define the dimensionless variables

$$\tilde{Q} = \frac{H}{\tilde{D}}, \quad \tilde{\Omega} = \Gamma\rho, \quad \tilde{\Omega}_k = \frac{k}{a^2\tilde{D}^2}, \quad \tilde{D} = \sqrt{\frac{8\pi G}{3}}\rho, \quad (50)$$




FIG. 1: Phase plot for system (42), (43) for different choices of the parameter ω .

Constraint (49) becomes

$$\tilde{Q}^2 + \tilde{\Omega}_k + \tilde{\Omega} = 1. \quad (51)$$

Introducing the time derivative

$$\frac{df}{d\eta} = \tilde{D}^{-1}f, \quad (52)$$

Label	\tilde{Q}	$\tilde{\Omega}_k$	Existence	k_1	k_2	Stability
Q_1	-1	0	always	$3(\omega + 1)$	$-(3\omega + 1)$	saddle for $\omega > -\frac{1}{3}$
Q_2	1	0	always	$-3(\omega + 1)$	$3\omega + 1$	saddle for $\omega > -\frac{1}{3}$
Q_3	0	$\frac{3(\omega+1)}{6\omega+4}$	$\omega \geq -\frac{1}{3}$	$-\sqrt{\frac{3}{2}}\sqrt{-((\omega+1)(3\omega+1))}$	$\sqrt{\frac{3}{2}}\sqrt{-((\omega+1)(3\omega+1))}$	centre for $\omega > -\frac{1}{3}$

TABLE II: Equilibrium points of system (56) and (57) for the parameter region $\omega \geq -\frac{1}{3}$. N.H. stands for nonhyperbolic

we obtain the dynamical system

$$\frac{d\tilde{Q}}{d\eta} = \frac{3}{2}(\omega + 1) \left(\tilde{Q}^2 + 2\tilde{\Omega} - 1 \right) + \tilde{\Omega}_k, \quad (53)$$

$$\frac{d\tilde{\Omega}_k}{d\eta} = \tilde{Q}(3\omega + 1)\tilde{\Omega}_k, \quad (54)$$

$$\frac{d\tilde{\Omega}}{d\eta} = -3\tilde{Q}(\omega + 1)\tilde{\Omega}. \quad (55)$$

Using (51) as a definition of $\tilde{\Omega}$, we obtain the reduced system

$$\frac{d\tilde{Q}}{d\eta} = \tilde{\Omega}_k - \frac{3}{2}(\omega + 1) \left(\tilde{Q}^2 + 2\tilde{\Omega}_k - 1 \right), \quad (56)$$

$$\frac{d\tilde{\Omega}_k}{d\eta} = (3\omega + 1)\tilde{Q}\tilde{\Omega}_k, \quad (57)$$

defined in the phase space

$$\left\{ (\tilde{Q}, \tilde{\Omega}_k) \in [-1, 1] \times [0, 1] : \tilde{Q}^2 + \tilde{\Omega}_k \leq 1 \right\}. \quad (58)$$

The equilibrium points of system (56) and (57) for the parameter region $\omega \geq -\frac{1}{3}$ are presented in Tab. II.

The equilibrium points for this case are the following.

Q_1 : $(\tilde{Q}, \tilde{\Omega}_k) = (-1, 0)$, always exists. It corresponds to a flat contracting solution. The eigenvalues are $3(\omega + 1)$ and $-(3\omega + 1)$, therefore, it is a saddle for $\omega > -\frac{1}{3}$.

Q_2 : $(\tilde{Q}, \tilde{\Omega}_k) = (1, 0)$, always exists. It corresponds to a flat expanding solution. The eigenvalues are $-3(\omega + 1)$ and $3\omega + 1$, therefore, it is a saddle for $\omega > -\frac{1}{3}$.

Q_3 : $(\tilde{Q}, \tilde{\Omega}_k) = \left(0, \frac{3(\omega+1)}{6\omega+4} \right)$, exists for $\omega \geq -\frac{1}{3}$. The eigenvalues are $-\sqrt{\frac{3}{2}}\sqrt{-((\omega+1)(3\omega+1))}$ and $\sqrt{\frac{3}{2}}\sqrt{-((\omega+1)(3\omega+1))}$. It corresponds to the Einstein static universe, a centre (nonhyperbolic) for $\omega > -\frac{1}{3}$.

Finally, for $\omega = -1/3$ we obtain a line of equilibrium points $\tilde{Q}^2 + \tilde{\Omega}_k = 1$ with eigenvalues $\{0, -2Q\}$ that is unstable for $Q < 0$ and stable for $Q > 0$.




FIG. 2: Phase plot for system (56), (57) for different choices of the parameter w .

IV. CONCLUDING REMARKS

In the present work, we provided detailed phase space analyses of the modified Friedmann's equations that describe the evolution of the early Universe. The modified Friedmann's equations are obtained from Verlinde's entropic and emergent force scenario and encode the string T-duality effect using the minimal length effect. One of the main implications of such modifications is the initial singularity of the early Universe. In particular, as was shown in [102], the initial singularity is avoided due to the quantum gravity effects.

That can be explained by the fact that there exists a minimal length scale l_0 . An interesting result can be found if we set $\dot{a} = 0$ ($H = 0$); for $\Gamma > 0$, one can find the contraction, or the collapsing matter stops at some critical distance $\sqrt{2}l_0$ and some critical density ρ_c (in the loop-quantum gravity scenario). Such a state is expected to be unstable and provides a natural way of generating an early expanding universe. However, in that case, for $\Gamma < 0$, we can solve the equation for the brane-tension to get (23), that imposes the constraint $2G\pi a^2 - 2\Gamma k \geq 0$. Solving for the scale factor $a = a_{\min}$, we get that for a closed universe, there exists a minimal length for the scale factor and, in particular, for $\omega = 0$, we get $a_{\min} = \sqrt{2}l_0$. For a flat universe with $k = 0$, the condition $\dot{a} = 0$ ($H = 0$) gives $\sigma = 1/(2|\Gamma|)$.

Following (23) for an open Universe with $k = -1$, we can get the critical brane-tension provided $\Gamma < 0$. Since l_0 is of Planck length order, i.e., $l_0 \sim l_{Pl}$, its effect becomes important only in short distances. The critical density that corresponds to the minimal scale goes like $\sigma_c \sim l_{Pl}^2$, and it may prove a bound for the maximum density in nature.

Moreover, we construct two dynamical systems depending on whether $-1 < \omega \leq -1/3$ and $\omega \geq -1/3$, and we classify the equilibrium points of each system. Finally, exact solutions and cosmological implications are discussed.

Here we have first studied the phase space analyses for the case $\Gamma \leq 0$, or equivalently with the regime corresponding to $-1 < \omega \leq -\frac{1}{3}$. In this regime, when $\Gamma < 0$, we can interpret the modified Friedmann equation as the one corresponding to the one of Brane-cosmology [113–119], remarkably, on the RS2 scenario with a single Z_2 symmetric brane. Moreover, the matter field behaves as a quintessence fluid for the region of the equation of state presented. Our model mimics the effect of a scalar field trapped on an FLRW-RS2 braneworld [120, 121].

Among other things, we found the following equilibrium points: P_1 that corresponds to a flat contracting solution that is a saddle for $-1 < \omega < -\frac{2}{3}$ or a sink for $-\frac{2}{3} < \omega < -\frac{1}{3}$; P_2 , that corresponds to an expanding flat solution that is a saddle for $-1 < \omega < -\frac{2}{3}$ or a source for $-\frac{2}{3} < \omega < -\frac{1}{3}$; P_3 , that corresponds to a flat contracting matter-dominated solution that is a source for $-1 < \omega < -\frac{1}{3}$, and it mimics a quintessence fluid; P_4 , that corresponds to a flat expanding matter-dominated solution that is a sink for $-1 < \omega < -\frac{1}{3}$, and it mimics a quintessence fluid and P_5 that exists for $-\frac{2}{3} < \omega \leq -\frac{1}{3}$ and corresponds to the Einstein static universe, and it is a saddle.

Then, we elaborated in more detail the domain $\Gamma \geq 0$, or equivalently the region $\omega \geq -\frac{1}{3}$ (Loop quantum gravity cosmology). In this case, we found the equilibrium points: Q_1 , which corresponds to a flat contracting solution, and it is a saddle for $\omega > -\frac{1}{3}$; Q_2 , that corresponds to a flat expanding solution and it is a saddle for $\omega > -\frac{1}{3}$ and Q_3 , which corresponds to the Einstein static universe and it is a centre (nonhyperbolic) for $\omega > -\frac{1}{3}$. The important finding is that the quantum gravity effects can provide a natural way of generating a dynamical early universe.

Acknowledgments

ADM was supported by Agencia Nacional de Investigación y Desarrollo (ANID) Subdirección de Capital Humano/Doctorado Nacional/año 2020 folio 21200837, Gastos operacionales Proyecto de tesis/2022 folio 242220121, and by Vicerrectoría de Investigación y Desarrollo Tecnológico (Vridt) at Universidad Católica del Norte. GL was funded through Concurso De Pasantías De Investigación Año 2022, Resolución Vridt No. 040/2022 and Resolución Vridt No. 054/2022. He also thanks the support of Núcleo de Investigación Geometría Diferencial y Aplicaciones, Resolución Vridt N°096/2022.

-
- [1] A. G. Riess *et al.* [Supernova Search Team], *Astron. J.* **116** (1998), 1009-1038.
 - [2] S. Perlmutter *et al.* [Supernova Cosmology Project], *Nature* **391** (1998), 51-54.
 - [3] S. Perlmutter *et al.* [Supernova Cosmology Project], *Astrophys. J.* **517** (1999), 565-586.
 - [4] S. M. Carroll, W. H. Press and E. L. Turner, *Ann. Rev. Astron. Astrophys.* **30** (1992), 499-542.
 - [5] P. Salucci, *Found. Phys.* **48** (2018) no.10, 1517-1537
 - [6] J. C. Mather, E. S. Cheng, R. A. Shafer, C. L. Bennett, N. W. Boggess, E. Dwek, M. G. Hauser, T. Kelsall, S. H. Moseley, Jr. and R. F. Silverberg, *et al.* *Astrophys. J. Lett.* **354** (1990), L37-L40.
 - [7] J. C. Mather, D. J. Fixsen, R. A. Shafer, C. Mosier and D. T. Wilkinson, *Astrophys. J.* **512** (1999), 511-520.
 - [8] C. L. Bennett *et al.* [WMAP], *Astrophys. J. Suppl.* **148** (2003), 1-27.

- [9] P. A. R. Ade *et al.* [Planck], *Astron. Astrophys.* **571** (2014), A23.
- [10] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, L. Campbell, Q. Parker, W. Saunders and F. Watson, *Mon. Not. Roy. Astron. Soc.* **416** (2011), 3017-3032.
- [11] B. Ratra and P. J. E. Peebles, *Phys. Rev. D* **37** (1988), 3406.
- [12] P. J. E. Peebles and B. Ratra, *Astrophys. J. Lett.* **325** (1988), L17.
- [13] K. Coble, S. Dodelson and J. A. Frieman, *Phys. Rev. D* **55** (1997), 1851-1859.
- [14] M. S. Turner and M. J. White, *Phys. Rev. D* **56** (1997) no.8, R4439.
- [15] R. R. Caldwell, R. Dave and P. J. Steinhardt, *Phys. Rev. Lett.* **80** (1998), 1582-1585.
- [16] K. Dutta, S. Panda and A. Patel, *Phys. Rev. D* **94** (2016) no.2, 024016.
- [17] E. N. Saridakis and M. Tsoukalas, *Phys. Rev. D* **93** (2016) no.12, 124032.
- [18] H. Sheikahmadi, E. N. Saridakis, A. Aghamohammadi and K. Saaidi, *JCAP* **10** (2016), 021.
- [19] R. Garcia-Salcedo and N. Breton, *Int. J. Mod. Phys. A* **15** (2000), 4341-4354.
- [20] C. S. Camara, M. R. de Garcia Maia, J. C. Carvalho and J. A. S. Lima, *Phys. Rev. D* **69** (2004), 123504
- [21] E. Elizalde, J. E. Lidsey, S. Nojiri and S. D. Odintsov, *Phys. Lett. B* **574** (2003), 1-7.
- [22] C. Quercellini, M. Bruni, A. Balbi and D. Pietrobon, *Phys. Rev. D* **78** (2008), 063527.
- [23] Y. F. Cai, E. N. Saridakis, M. R. Setare and J. Q. Xia, *Phys. Rept.* **493** (2010), 1-60
- [24] E. N. Saridakis and S. V. Sushkov, *Phys. Rev. D* **81** (2010), 083510.
- [25] E. N. Saridakis, *Eur. Phys. J. C* **67** (2010), 229-235
- [26] G. Leon and E. N. Saridakis, *JCAP* **11** (2009), 006
- [27] N. Breton, R. Lazkoz and A. Montiel, *JCAP* **10** (2012), 013
- [28] C. Xu, E. N. Saridakis and G. Leon, *JCAP* **07** (2012), 005.
- [29] G. Leon and E. N. Saridakis, *JCAP* **03** (2013), 025.
- [30] G. Leon, J. Saavedra and E. N. Saridakis, *Class. Quant. Grav.* **30** (2013), 135001.
- [31] G. Kofinas, G. Leon and E. N. Saridakis, *Class. Quant. Grav.* **31** (2014), 175011.
- [32] G. Leon, Y. Leyva and J. Socorro, *Phys. Lett. B* **732** (2014), 285-297.
- [33] C. R. Fadrakas, G. Leon and E. N. Saridakis, *Class. Quant. Grav.* **31** (2014), 075018.
- [34] G. Leon and E. N. Saridakis, *JCAP* **11** (2015), 009.
- [35] G. Pulgar, J. Saavedra, G. Leon and Y. Leyva, *JCAP* **05** (2015), 046.
- [36] G. Leon and E. N. Saridakis, *JCAP* **04** (2015), 031.

- [37] A. Giacomini, S. Jamal, G. Leon, A. Paliathanasis and J. Saavedra, *Phys. Rev. D* **95** (2017) no.12, 124060.
- [38] A. Joyce, L. Lombroiser and F. Schmidt, *Ann. Rev. Nucl. Part. Sci.* **66** (2016), 95-122.
- [39] S. Tsujikawa, *Class. Quant. Grav.* **30** (2013), 214003.
- [40] S. Nojiri, S. D. Odintsov and V. K. Oikonomou, *Phys. Rept.* **692** (2017), 1-104.
- [41] S. Lepe, G. Otalora and J. Saavedra, *Phys. Rev. D* **96** (2017) no.2, 023536.
- [42] T. Harko, F. S. N. Lobo, G. Otalora and E. N. Saridakis, *JCAP* **12** (2014), 021.
- [43] G. Otalora, *Phys. Rev. D* **88** (2013), 063505.
- [44] G. Otalora, *JCAP* **07** (2013), 044.
- [45] L. Karpathopoulos, S. Basilakos, G. Leon, A. Paliathanasis and M. Tsampanlis, *Gen. Rel. Grav.* **50** (2018) no.7, 79.
- [46] Ö. Akarsu and T. Dereli, *Gen. Rel. Grav.* **45** (2013), 1211-1226.
- [47] Ö. Akarsu, T. Dereli, N. Katırcı and M. B. Sheftel, *Gen. Rel. Grav.* **47** (2015) no.5, 61.
- [48] O. Akarsu and T. Dereli, *Int. J. Theor. Phys.* **51** (2012), 612-621.
- [49] D. Grasso and H. R. Rubinstein, *Phys. Rept.* **348** (2001), 163-266.
- [50] D. Garrison, [arXiv:1608.01005 [astro-ph.CO]].
- [51] J. D. Finke, L. C. Reyes, M. Georganopoulos, K. Reynolds, M. Ajello, S. J. Fegan and K. McCann, *Astrophys. J.* **814** (2015) no.1, 20.
- [52] Z. Haba, A. Stachowski and M. Szydlowski, *JCAP* **07** (2016), 024.
- [53] A. Stachowski and M. Szydlowski, *Eur. Phys. J. C* **76** (2016) no.11, 606.
- [54] O. Hrycyna, M. Szydlowski and M. Kamionka, *Phys. Rev. D* **90** (2014) no.12, 124040.
- [55] M. Szydlowski, *Astrophys. Space Sci.* **339** (2012), 389-399.
- [56] M. Szydlowski and O. Hrycyna, *JCAP* **01** (2009), 039.
- [57] M. P. Dabrowski, T. Stachowiak and M. Szydlowski, *Phys. Rev. D* **68** (2003), 103519.
- [58] M. Szydlowski, A. Krawiec, A. Kurek and M. Kamionka, *Eur. Phys. J. C* **75** (2015) no.99, 5.
- [59] M. Szydlowski, A. Kurek and A. Krawiec, *Phys. Lett. B* **642** (2006), 171-178.
- [60] W. Godlowski and M. Szydlowski, *Phys. Lett. B* **623** (2005), 10-16.
- [61] M. Szydlowski, *Phys. Lett. B* **632** (2006), 1-5.
- [62] R. Garcia-Salcedo, T. Gonzalez and I. Quiros, *Phys. Rev. D* **89** (2014) no.8, 084047.
- [63] J. D. Barrow, *Phys. Lett. B* **235** (1990), 40-43 doi:10.1016/0370-2693(90)90093-L

- [64] J. D. Barrow and P. Saich, *Phys. Lett. B* **249** (1990), 406-410.
- [65] J. D. Barrow and N. J. Nunes, *Phys. Rev. D* **76** (2007), 043501.
- [66] S. del Campo and R. Herrera, *Phys. Lett. B* **670** (2009), 266-270.
- [67] A. Cid and S. del Campo, *JCAP* **01** (2011), 013.
- [68] A. Cid and S. del Campo, *AIP Conf. Proc.* **1471** (2012), 114-117.
- [69] R. Herrera, M. Olivares and N. Videla, *Eur. Phys. J. C* **73** (2013) no.1, 2295.
- [70] A. Cid, G. Leon and Y. Leyva, *JCAP* **02** (2016), 027.
- [71] M. Kierdorf et al., *Astronomy & Astrophysics* **600**, id.A18 (2017).
- [72] R. Jimenez and A. Loeb, *Astrophys. J.* **573** (2002), 37-42.
- [73] M. Moresco, A. Cimatti, R. Jimenez, L. Pozzetti, G. Zamorani, M. Bolzonella, J. Dunlop, F. Lamareille, M. Mignoli and H. Pearce, *et al. JCAP* **08** (2012), 006.
- [74] J. Magana, M. H. Amante, M. A. Garcia-Aspeitia and V. Motta, *Mon. Not. Roy. Astron. Soc.* **476** (2018) no.1, 1036-1049.
- [75] T. Jacobson, *Phys. Rev. Lett.* **75** (1995), 1260-1263.
- [76] M. Akbar and R. G. Cai, *Phys. Rev. D* **75** (2007), 084003.
- [77] R. G. Cai and L. M. Cao, *Phys. Rev. D* **75** (2007), 064008.
- [78] R. G. Cai and S. P. Kim, *JHEP* **02** (2005), 050.
- [79] B. Wang, E. Abdalla and R. K. Su, *Mod. Phys. Lett. A* **17** (2002), 23-30.
- [80] B. Wang, E. Abdalla and R. K. Su, *Phys. Lett. B* **503** (2001), 394-398.
- [81] R. G. Cai and Y. S. Myung, *Phys. Rev. D* **67** (2003), 124021.
- [82] R. G. Cai, L. M. Cao and Y. P. Hu, *JHEP* **08** (2008), 090.
- [83] S. Nojiri and S. D. Odintsov, *Gen. Rel. Grav.* **38** (2006), 1285-1304.
- [84] A. Sheykhi, B. Wang and R. G. Cai, *Nucl. Phys. B* **779** (2007), 1-12.
- [85] A. Sheykhi, B. Wang and R. G. Cai, *Phys. Rev. D* **76** (2007), 023515.
- [86] A. Sheykhi, *JCAP* **05** (2009), 019.
- [87] A. Sheykhi and B. Wang, *Phys. Lett. B* **678** (2009), 434-437.
- [88] A. Sheykhi, *Eur. Phys. J. C* **69** (2010), 265-269.
- [89] A. Sheykhi, *Class. Quant. Grav.* **27** (2010), 025007.
- [90] E. P. Verlinde, *JHEP* **04** (2011), 029.
- [91] E. P. Verlinde, *SciPost Phys.* **2** (2017) no.3, 016.
- [92] T. Padmanabhan, *Mod. Phys. Lett. A* **25** (2010), 1129-1136.

- [93] T. Padmanabhan, [arXiv:1206.4916 [hep-th]].
- [94] R. G. Cai, JHEP **11** (2012), 016.
- [95] K. Yang, Y. X. Liu and Y. Q. Wang, Phys. Rev. D **86** (2012), 104013.
- [96] A. Sheykhi, Phys. Rev. D **87** (2013) no.6, 061501
- [97] T. Padmanabhan, Phys. Rev. Lett. **78** (1997), 1854-1857.
- [98] P. Nicolini, E. Spallucci and M. F. Wondrak, Phys. Lett. B **797** (2019), 134888.
- [99] P. Nicolini, Gen. Rel. Grav. **54** (2022) no.9, 106.
- [100] P. Gaete, K. Jusufi and P. Nicolini, Phys. Lett. B **835** (2022), 137546.
- [101] K. Jusufi, Universe **9** (2023) no.1, 41.
- [102] K. Jusufi and A. Sheykhi, Phys. Lett. B **836** (2023), 137621.
- [103] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641** (2020), A6 [erratum: Astron. Astrophys. **652** (2021), C4]
- [104] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri and D. Scolnic, Astrophys. J. **876** (2019) no.1, 85.
- [105] M. Maggiore, Phys. Lett. B **304** (1993), 65-69.
- [106] S. Hossenfelder, Living Rev. Rel. **16** (2013), 2
- [107] S. Aghababaei, H. Moradpour and E. C. Vagenas, Eur. Phys. J. Plus **136** (2021) no.10, 997.
- [108] A. Giacomini, G. Leon, A. Paliathanasis and S. Pan, Eur. Phys. J. C **80** (2020) no.10, 931
- [109] A. Paliathanasis and G. Leon, Gen. Rel. Grav. **55** (2023) no.1, 12
- [110] M. G. Dainotti, B. De Simone, T. Schiavone, G. Montani, E. Rinaldi and G. Lambiase, Astrophys. J. **912** (2021) no.2, 150.
- [111] M. G. Dainotti, B. De Simone, T. Schiavone, G. Montani, E. Rinaldi, G. Lambiase, M. Bogdan and S. Ugale, Galaxies **10** (2022) no.1, 24.
- [112] A. Ashtekar and D. Sloan, Gen. Rel. Grav. **43** (2011), 3619-3655.
- [113] L. Randall and R. Sundrum, Phys. Rev. Lett. **83** (1999), 3370-3373
- [114] P. Binetruy, C. Deffayet, U. Ellwanger and D. Langlois, Phys. Lett. B **477** (2000), 285-291.
- [115] P. Binetruy, C. Deffayet and D. Langlois, Nucl. Phys. B **565** (2000), 269-287
- [116] P. Bowcock, C. Charmousis and R. Gregory, Class. Quant. Grav. **17** (2000), 4745-4764
- [117] R. Maartens, Living Rev. Rel. **7** (2004), 7
- [118] P. Brax, C. van de Bruck and A. C. Davis, Rept. Prog. Phys. **67** (2004), 2183-2232
- [119] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, Phys. Rept. **513** (2012), 1-189

- [120] Y. Leyva, D. Gonzalez, T. Gonzalez, T. Matos and I. Quiros, Phys. Rev. D **80** (2009), 044026
- [121] D. Escobar, C. R. Fadrugas, G. Leon and Y. Leyva, Class. Quant. Grav. **29** (2012), 175005
- [122] S. Wiggins. “Introduction to Applied Nonlinear dynamical systems and Chaos”. Springer (2003).
- [123] L. Perko, “Differential equations and dynamical systems, third edition” (Springer-Verlag, New York, 2001). pp 272-273 & pp 281-282.
- [124] M. W. Hirsch and S. Smale. “Differential equations, dynamical systems, and linear algebra”. New York: Academic Press (1974).
- [125] R. Tavakol, “Introduction to dynamical systems”, ch 4. Part one, pp. 84–98, Cambridge University Press, Cambridge, England, (1997).
- [126] A.A. Coley, 2003, “Dynamical systems and cosmology” (Kluwer Academic, Dordrecht: ISBN 1-4020-1403-1).
- [127] A. A. Coley, [arXiv:gr-qc/9910074 [gr-qc]].