

Classical Spin Gravitational Compton Scattering

N. E. J. Bjerrum-Bohr, Gang Chen, Marcos Skowronek

*Niels Bohr International Academy, Niels Bohr Institute,
University of Copenhagen, Blegdamsvej 17,
DK-2100 Copenhagen Ø, Denmark*

E-mail:

bjbohr@nbi.dk; gang.chen@nbi.ku.dk; marcos.santos@nbi.ku.dk

ABSTRACT: We introduce a novel bootstrap method for classical Compton scattering amplitudes involving two massless gluon/graviton particles and two arbitrary-spin infinite-mass particles in a heavy-mass effective field theory context. Using a suitable ansatz, we deduce new and explicit classical spin results for gluon four and five-point infinite mass processes that exhibit exponentiated three-point factorizations to all orders in spin and feature no spurious poles. We discuss the generalization of our bootstrap to higher multiplicities and summarize future potential applications.

KEYWORDS: Scattering amplitudes, color-kinematics duality and the double copy, general relativity from amplitudes

Contents

1	Introduction	1
2	Classical Compton amplitudes	3
2.1	Three-point amplitude	3
2.2	Classical gravity amplitude decomposition in the heavy mass limit	4
3	Spinning Compton Amplitudes	5
3.1	Four-point Amplitude	5
3.2	Five-point Compton scattering	7
4	Building blocks for an all-multiplicity amplitude ansatz	10
5	Conclusion	11
6	Acknowledgements	12
A	Classical spin theory with finite mass	13
B	Removable singularities in the G functions	16

1 Introduction

Observations of gravitational waves from black hole mergers have fast evolved into an exciting avenue for testing Einstein’s classical theory. To facilitate this, a modern and efficient computational program inspired by various practical quantum field theory implementations of general relativity, see, for instance, [1–7], was developed in [8, 9] prompted by the state of current theoretical analysis and precision requirements of observations [10]. It has made quantum scattering amplitudes for superheavy black holes – taken as point particles – a powerful resource for delivering much-needed analytic precision in fully relativistic post-Minkowskian frameworks and promises new computational efficiency, particularly in analytical regions of the near-merger regimes.

Several post-Minkowskian spinless calculations have already made new landmarks by

extending the expansion in Newton’s constant G_N [11–29] with [30–32] pioneering fourth post-Minkowskian order. Another research attraction is the inclusion of spin effects which can lead to significant quantitative augmentations of observed signals for events. For spinning black holes, improvement till now has been gradual because of the more complicated and specialized framework, but one finds several vital advancements using spinor helicity formalism in [33–47], using covariant and world-line field theory in [48–51] and with various phenomenological applications in [52–65].

For addressing classical spins, it is customary to operate in the context of a Lagrangian formalism that involves higher spin particles systematized in expansions with increasing multi-pole interactions as pioneered in [66]. Unfortunately, the formalism behind this is relatively elaborate, and derived amplitudes usually contain intricate redundancies. In this presentation, we take an alternative route and bootstrap covariant arbitrary-spin amplitudes from factorization limits.


Our main focus here will be classical spin Compton scattering amplitudes, where an infinite mass spinning particle interacts with gluons. As we explain, we can extend such amplitudes into results for gravitational Compton scattering amplitudes utilizing the double-copy [67–69]. The rationale for mainly considering gravitational Compton scattering amplitudes in the infinite mass limit is that such amplitudes are convenient building blocks in generalized unitarity approaches [25–28, 70, 71] that identify the gravitational classical radial action computing from an integrand determined by generalized unitarity and constrained by delta functions [23, 24, 26].

In the bootstrap method, we use minimal physical input constraints such as the exponentiated Kerr black hole three-point amplitude [34], and we will introduce a convenient ansatz for the amplitude that avoids spurious poles and is fixed uniquely from imposing factorization identities. This suggests that we can derive classical Kerr black hole scattering from our amplitudes, as we will discuss later in the presentation. We find support for our classical amplitudes by comparing them to results derived at four-point linear and quadratic spin order [49, 72].

We arrange this paper as follows: in section 2, we will quickly review infinite mass Compton gluon and graviton tree amplitudes to establish a convenient and efficient framework for computation. Then we will introduce our new bootstrap method and demonstrate the calculation of four and five-point classical amplitudes in section 3. An essential inspiration in the bootstrap method is the formalism [73]. Finally, we will outline a generalization of the proposed computational technology in section 4 and discuss possible applications and questions for the future in section 5.

2 Classical Compton amplitudes

The tree amplitudes considered in this paper involve an infinity mass and spin particle with finite classical ring radius $a = s/m$ ($s \rightarrow \infty$, $m \rightarrow \infty$), which interacts with an arbitrary $n - 2$ number of gluons in the context of a heavy-mass effective field theory (HEFT) [26, 48, 73–78]. We depict the processes considered in the following way:



$$(2.1)$$

Here p_1, \dots, p_{n-2} denotes momenta of gluons while \bar{p}_{n-1} and \bar{p}_n are momenta for the heavy spinning particle. In the heavy mass limit the velocity v is fixed by the on-shell condition $v \cdot v = 1$ and $v \cdot q = 0$ which follows from $\bar{p}_n^2 = \bar{p}_{n-1}^2 = (mv - q)^2 = m^2$. We define $q \equiv (p_1 + \dots + p_{n-2})$.

2.1 Three-point amplitude

We start with the infinite mass three-point Compton amplitude, which we define as follows (using shorthand notation for momenta)

$$\lim_{m \rightarrow \infty} A_a(1, \bar{2}, \bar{3}) \equiv A_a(1, v). \quad (2.2)$$

It can conveniently be written in terms of the three-point color-kinematic numerator $\mathcal{N}_a(1, v)$

$$A_a(1, v) = \mathcal{N}_a(1, v), \quad (2.3)$$

that has a closed form for an arbitrary spin with minimal couplings as shown in [34–36, 79]

$$\mathcal{N}_a(1, v) = m (v \cdot \varepsilon_1) \exp \left(i \frac{p_1 \cdot S \cdot \varepsilon_1}{m v \cdot \varepsilon_1} \right). \quad (2.4)$$

We can rewrite the exponential by separating the even and odd powers in the spin variable and using the identity $\epsilon^{\mu\nu\rho\sigma} \epsilon_{\alpha\beta\gamma\delta} = -\delta_{\alpha\beta\gamma\delta}^{\mu\nu\rho\sigma}$

$$m v^\mu \exp \left(i \frac{p_1 \cdot S \cdot \varepsilon_1}{m v \cdot \varepsilon_1} \right) = m \cosh(p_1 \cdot a) v^\mu - i \left(\frac{\sinh(p_1 \cdot a)}{p_1 \cdot a} \right) (p_1 \cdot S)^\mu. \quad (2.5)$$

In the above expressions ε_i denotes gluon polarizations and $S^{\mu\nu} \equiv -m \epsilon^{\mu\nu\rho\sigma} v_\rho a_\sigma$. Since the singularity in the second term of Eq. (2.5) is removable, we can define an analytic

extension that is holomorphic everywhere in the complex plane by considering the function

$$G_1(x_1) = \frac{\sinh(x_1)}{x_1} \quad \forall \quad x_1 \neq 0 \quad \text{with} \quad G_1(0) = \lim_{x_1 \rightarrow 0} \frac{\sinh(x_1)}{x_1} = 1. \quad (2.6)$$

Thus we can define

$$w_1^\mu \equiv m \cosh(p_1 \cdot a) v^\mu - i \left(G_1(p_1 \cdot a) \right) (p_1 \cdot S)^\mu, \quad (2.7)$$

which defines a three-point numerator which has the advantage of manifestly featuring no pole terms in $a \cdot p_1$

$$\mathcal{N}_a(1, v) = w_1 \cdot \varepsilon_1. \quad (2.8)$$

We note that gauge invariance is manifest in Eq. (2.8) from the anti-symmetric property of the spin tensor and the three point on-shell condition $v \cdot p_1 = 0$, which imply that $w_1 \cdot p_1 = 0$. We conclude this section by noting that three-point gravity amplitudes can be computed from the double copy using a product of one scalar gluon numerator, $\mathcal{N}_0(1, v)$, and one arbitrary spin numerator factor, $\mathcal{N}_a(1, v)$ [80],

$$M_a(1, v) = \mathcal{N}_0(1, v) \mathcal{N}_a(1, v) = m (v \cdot \varepsilon_1) (w_1 \cdot \varepsilon_1). \quad (2.9)$$

2.2 Classical gravity amplitude decomposition in the heavy mass limit

Before we discuss the four-point gravity amplitude, we will review some material related to convenient amplitude decompositions in the heavy mass limit. We will here refer to the discussion in [26] (see also [27]), and start by noting that it is convenient to decompose gravity amplitudes in the following way when considering classical black hole scattering perturbatively:

$$\begin{array}{c} \begin{array}{c} 1 \quad 2 \\ \diagdown \quad / \\ \text{blob} \\ / \quad \diagdown \\ \bar{4} \quad \bar{3} \end{array} \xrightarrow{m \rightarrow \infty} \begin{array}{c} \begin{array}{c} 1 \quad 2 \\ | \quad | \\ \bar{4} \quad \bar{3} \end{array} + \begin{array}{c} \begin{array}{c} 1 \quad 2 \\ \diagdown \quad / \\ \text{H} \\ / \quad \diagdown \\ \bar{4} \quad \bar{3} \end{array} \end{array}, \quad (2.10) \end{array}$$

The first term in this diagrammatical expression corresponds to a velocity cut imposed by a δ -function constraint. This contribution is of order $\mathcal{O}(m^3)$ ¹. The second term, which we will refer to as the classical tree amplitude, is of order $\mathcal{O}(m^2)$ ($\mathcal{O}(\hbar^0)$). Since the first term is an iteration in the infinite mass limit, the following section focuses solely on computing the second term of (2.10). In Appendix A, we will discuss calculating the first contribution directly when considering a finite heavy mass.

¹Equivalently, if we count in powers of \hbar , of order $\mathcal{O}(\hbar^{-1})$

One can follow the same reasoning at five-point, and focus only on the last term of the following expansion:

$$(2.11)$$

3 Spinning Compton Amplitudes

3.1 Four-point Amplitude

We will now consider how to compute the classical four-point infinity-mass amplitude and start by writing the amplitude in terms of color-kinematic massless pole numerator factors:

$$A_a(12, v) = \frac{\mathcal{N}_a(1, 2, v) - \mathcal{N}_a(2, 1, v)}{p_{12}^2} \equiv \frac{\mathcal{N}_a([1, 2], v)}{p_{12}^2}. \quad (3.1)$$

Here $\mathcal{N}_a(1, 2, v)$ and $\mathcal{N}_a(2, 1, v)$ are numerators, and in the last step, we have used a commutator form $\mathcal{N}_a([1, 2], v)$ to write the amplitude in shorthand form. The color-kinematic numerator $\mathcal{N}_a([1, 2], v)$ has a massive pole in $v \cdot p_1$ [81], and thus it follows from its factorization behavior that we have

$$\mathcal{N}_a([1, 2], v) \rightarrow \frac{p_1 \cdot p_2}{m v \cdot p_1} \mathcal{N}_a(1, v) \mathcal{N}_a(2, v) = \frac{1}{m v \cdot p_1} (p_1 \cdot p_2) (w_1 \cdot \varepsilon_1) (w_2 \cdot \varepsilon_2), \quad (3.2)$$

which we will rewrite as

$$\mathcal{N}_a([1, 2], v) = -\frac{w_1 \cdot F_1 \cdot F_2 \cdot w_2}{m v \cdot p_1} + \mathcal{N}'_a([1, 2], v), \quad (3.3)$$

where $F_i^{\mu\nu} \equiv p_i^\mu \varepsilon_i^\nu - \varepsilon_i^\mu p_i^\nu$ denotes the abelian field strength.

This separates the color-kinematic numerator into a $-\frac{w_1 \cdot F_1 \cdot F_2 \cdot w_2}{m v \cdot p_1}$ with a pole, and an analytic and finite part $\mathcal{N}'_a([1, 2], v)$.

Now factorizing the numerator for $p_{12}^2 \rightarrow 0$ we see that

$$\mathcal{N}_a([1, 2], v)|_{p_{12}^2 \rightarrow 0} = \sum_{\lambda_i} \mathcal{N}_{\text{YM}}([1, 2], i) \times \mathcal{N}_a(i, v), \quad (3.4)$$

where we can write the three-point Yang-Mills numerator as, see *e.g.* [73],

$$\mathcal{N}_{\text{YM}}([1, 2], i) = -\frac{\varepsilon_i \cdot F_1 \cdot F_2 \cdot \varepsilon_i}{\varepsilon_i \cdot p_1}. \quad (3.5)$$

We consider an ansatz form to deduce $\mathcal{N}'_a([1, 2], v)$ at four points. We first introduce the function $G_2(x_1; x_2)$ (a generalization of the function $G_1(x_1)$) where one again can prove that all apparent singularities are removable²

$$G_2(x_1; x_2) \equiv \frac{1}{x_2} \left(\frac{\sinh(x_{12})}{x_{12}} - \cosh(x_2) \frac{\sinh(x_1)}{x_1} \right) \equiv \frac{1}{x_2} \left(G_1(x_{12}) - \cosh(x_2) G_1(x_1) \right). \quad (3.6)$$

We then work out which types of terms we can have in the ansatz considering the above functions and evaluate the factorization in Eq. (3.4). We find that any ansatz piece has to be of the following universal form (depending on the power in the spin variable),

$$\begin{aligned} \text{even} &: (v \cdot X \cdot a)(p \cdot Y \cdot a) G_1(x_1) G_1(x_2) \\ \text{odd} &: \text{tr}(S \cdot X) G_1(x_{12}), (p \cdot X \cdot S \cdot Y \cdot a) G_2(x_1; x_2), \end{aligned} \quad (3.7)$$

where we have defined $x_i \equiv p_i \cdot a$ and $x_{i\dots j} \equiv p_{i\dots j} \cdot a$. We follow the logic that we can only have a product of at most two G -functions since we only have two gluons at four points, and we insist on preserving the scaling in the spin variable of the three-point amplitudes by requiring that factors of a present in the denominators of the G functions cancel out with field strength monomial terms in the ansatz terms. Using the property of symmetry under parity ($a \rightarrow -a$) of the G -functions, it is furthermore easy to infer that only products $G_1(x_1)G_1(x_2)$ are possible for the even part. For the odd part, one can have either $G_1(x_{12})$ or $G_2(x_1; x_2)$ with a factor of S or $\dots \cdot S \cdot X \cdot a$, respectively, with X denoting a field strength tensor product.

Thus, we consider all the 14 possible independent monomials in the ansatz and bootstrap its form using the fact that all massless factorizations are consistent. Thereby we entirely fix the factorization behavior in Eq. (3.4), which leads to the expression:

$$\begin{aligned} \mathcal{N}'_a(12, v) = m & \left((a \cdot F_2 \cdot F_1 \cdot v)(a \cdot p_1) - (a \cdot F_2 \cdot v)(a \cdot F_1 \cdot p_2) \right) G_1(x_1) G_1(x_2) \\ & + i \left((a \cdot F_1 \cdot F_2 \cdot S \cdot p_2 + a \cdot F_2 \cdot F_1 \cdot S \cdot p_1) G_2(x_1; x_2) + \text{tr}(S \cdot F_2 \cdot F_1) G_1(x_{12}) \right). \end{aligned} \quad (3.8)$$

To fix the factorization, we sum over gluon states using


$$\sum_i \varepsilon_i^{*\mu} \varepsilon_i^\nu = \eta^{\mu\nu}.$$

²It is observed that the function $G_2(x_2; x_1)$ is independent since one can verify that $G_2(x_2; x_1) + G_2(x_1; x_2) = 0$.

Although the form Eq. (3.8) is not manifestly anti-symmetric, one can straightforwardly check its anti-symmetry using the relation:

$$\begin{aligned} & (a \cdot F_1 \cdot p_2) (a \cdot F_2 \cdot v) + (a \cdot F_2 \cdot p_1) (a \cdot F_1 \cdot v) \\ & - (a \cdot p_2) (a \cdot F_1 \cdot F_2 \cdot v) - (a \cdot p_1) (a \cdot F_2 \cdot F_1 \cdot v) = 0. \end{aligned} \quad (3.9)$$

Using the double copy relations, it follows that we have the classical spinning gravitational Compton amplitude



$$\equiv M_a(12, v) = \frac{\mathcal{N}_a([1, 2], v) \mathcal{N}_0([1, 2], v)}{p_{12}^2}, \quad (3.10)$$

where:

$$\mathcal{N}_0([1, 2], v) = -m \frac{v \cdot F_1 \cdot F_2 \cdot v}{v \cdot p_1}. \quad (3.11)$$

3.2 Five-point Compton scattering

We will now consider the five-point Compton amplitude. We can write this in the form

$$A_a(123, v) = \frac{\mathcal{N}_a([1, 2], 3], v)}{p_{12}^2 p_{123}^2} + \frac{\mathcal{N}_a([1, [2, 3]], v)}{p_{23}^2 p_{123}^2}, \quad (3.12)$$

$$(3.13)$$

$$(3.14)$$

again expressed in terms of color-kinematic numerators solely organized in terms of the massless poles. Compared to the four-point case, we now have three massive poles in the color-kinematic numerators and thus arrive at the following factorization behavior

$$\begin{aligned} \mathcal{N}_a([1, 2], 3], v) & \rightarrow \frac{p_{12} \cdot p_3}{m p_{12} \cdot v} \mathcal{N}_a([1, 2], v) \mathcal{N}_a(3, v) \\ \mathcal{N}_a([1, [2, 3]], v) & \rightarrow \frac{p_1 \cdot p_2}{m p_{13} \cdot v} \mathcal{N}_a([1, 3], v) \mathcal{N}_a(2, v) \\ \mathcal{N}_a([1, 2], 3], v) & \rightarrow \frac{p_1 \cdot p_2}{m p_1 \cdot v} \mathcal{N}_a(1, v) \mathcal{N}_a([2, 3], v). \end{aligned} \quad (3.15)$$

Again as we did for the four-point case, we use factorization to recast the expression into one where we can identify the finite piece $\mathcal{N}'_a([[1, 2], 3], v)$ with no pole³

$$\begin{aligned}
\mathcal{N}_a([[1, 2], 3], v) &= -\frac{(w_1 \cdot F_1 \cdot F_2 \cdot w_2)(p_{12} \cdot F_3 \cdot w_3)}{m^2 v \cdot p_1 v \cdot p_{12}} - \frac{(w_1 \cdot F_1 \cdot F_3 \cdot w_3)(p_1 \cdot F_2 \cdot w_2)}{m^2 v \cdot p_1 v \cdot p_{13}} \\
&+ \frac{(w_1 \cdot F_1 \cdot F_2 \cdot F_3 \cdot w_3) \cosh(p_2 \cdot a)}{m v \cdot p_1} + \frac{\mathcal{N}'_a([1, 2], v)(p_{12} \cdot F_3 \cdot w_3)}{m v \cdot p_{12}} \\
&+ \frac{\mathcal{N}'_a([1, 3], v)(p_1 \cdot F_2 \cdot w_2)}{m v \cdot p_{13}} + \frac{(w_1 \cdot F_1 \cdot p_2) \mathcal{N}'_a([2, 3], v)}{m v \cdot p_1} + \mathcal{N}'_a([[1, 2], 3], v). \tag{3.16}
\end{aligned}$$

In this case, we find that it can be composed of the following building blocks (extending the logic from the four-point case)

$$\begin{aligned}
\text{even} : & (v \cdot X \cdot a)(p \cdot Y \cdot a) G_1(x_i) G_1(x_{jk}) \\
& (v \cdot X \cdot a)(p \cdot Y \cdot a)(p \cdot Z \cdot a) G_1(x_i) G_2(x_j; x_k) \\
\text{odd} : & \text{tr}(S \cdot X) G_1(x_{123}), (p \cdot X \cdot S \cdot Y \cdot a) G_2(x_{ij}; x_k) \\
& (p \cdot X \cdot S \cdot Y \cdot a)(p \cdot Z \cdot a) G_3(x_i; x_j, x_k) \\
& (p \cdot X \cdot S \cdot Y \cdot a)(p \cdot Z \cdot a) G_1(x_1) G_1(x_2) G_1(x_3), \tag{3.17}
\end{aligned}$$

where $\{i, j, k\}$ are elements in the possible permutations of the set $\{1, 2, 3\}$ and

$$\begin{aligned}
G_3(x_1; x_2, x_3) := & \frac{1}{x_2 x_3} \left(\frac{\sinh(x_{123})}{x_{123}} - \cosh(x_2) \frac{\sinh(x_{13})}{x_{13}} - \cosh(x_3) \frac{\sinh(x_{12})}{x_{12}} \right. \\
& \left. + \cosh(x_2) \cosh(x_3) \frac{\sinh(x_1)}{x_1} \right). \tag{3.18}
\end{aligned}$$

As we discussed in the four-point case, we have some redundancy since the following type of identities hold

$$G_3(x_1; x_2, x_3) + G_3(x_2; x_1, x_3) + G_3(x_3; x_1, x_2) = -G_1(x_1) G_1(x_2) G_1(x_3). \tag{3.19}$$

The ansatz for the finite piece at five points has 981 independent (non-redundant) terms. Again we fix it completely by imposing the massless cut conditions,

$$\begin{aligned}
A_a(123, v) \Big|_{p_{123}^2 \rightarrow 0} &= \sum_{\lambda_i} A_{\text{YM}}(123, i) \times \mathcal{N}_a(i, v), \tag{3.20} \\
\mathcal{N}_a([[1, 2], 3], v) \Big|_{p_{12}^2 \rightarrow 0} &= \sum_{\lambda_i} \mathcal{N}_{\text{YM}}([1, 2], i) \times \mathcal{N}_a([i, 3], v),
\end{aligned}$$

³For the part of the numerator with poles, we see that the first three terms follow a similar structure as in the spinless case [73] but with the change $v \cdot F_i \rightarrow w_i \cdot F_i$, which accounts for the inclusion of spin.

$$+ \frac{\mathcal{N}_a([[1, 2], 3], v) \mathcal{N}_0([[1, 2], 3], v)}{p_{12}^2 p_{123}^2} + \frac{\mathcal{N}_a([[1, 3], 2], v) \mathcal{N}_0([[1, 3], 2], v)}{p_{13}^2 p_{123}^2}, \quad (3.25)$$

where the five-point numerator for the scalar case is [73]:

$$\mathcal{N}_0([[1, 2], 3], v) = m \left(\frac{v \cdot F_1 \cdot F_2 \cdot F_3 \cdot v}{v \cdot p_1} - \frac{v \cdot F_1 \cdot F_2 \cdot v \cdot p_{12} \cdot F_3 \cdot v}{v \cdot p_1 v \cdot p_{12}} - \frac{v \cdot F_1 \cdot F_3 \cdot v \cdot p_1 \cdot F_2 \cdot v}{v \cdot p_1 v \cdot p_{13}} \right). \quad (3.26)$$

4 Building blocks for an all-multiplicity amplitude ansatz

Considering the four and five-point numerators, we will now make some general remarks about extensions of the formalism to all-multiplicity Compton amplitudes for massive classical spinning particles. Starting with amplitude representations with numerators organized in massless poles similar to the lower point examples, we can again decompose the color-kinematic numerators into pieces with massive poles and a finite term. We will focus mainly on bootstrapping the finite contribution. Similar to the lower point demonstrations, one can set up a convenient ansatz based on a generalization of the G_1 , G_2 , and G_3 functions. One can prove in all generality that the function $G_r(x_1; x_2, \dots, x_r)$ only features removable singularities, see Appendix B.

$$G_r(x_1; x_2, \dots, x_r) \equiv \frac{1}{x_2 \cdots x_r} \times \left(\sum_{\tau_a \cup \tau_b = \tau} \frac{\sinh(x_{1\tau_a})}{x_{1\tau_a}} \prod_{i \in \tau_b} (-\cosh(x_i)) \right), \quad (4.1)$$

Here τ is the $\{2, \dots, r\}$ and we have a summation over all the partitions of τ into subsets τ_a and τ_b , including the empty set. It is easy to show that the following generic recursive relation holds

$$G_r(x_1; x_2, \dots, x_r) = \frac{1}{x_r} \left(G_{r-1}(x_1 + x_r; x_2, \dots, x_{r-1}) - G_{r-1}(x_1; x_2, \dots, x_{r-1}) \cosh(x_r) \right). \quad (4.2)$$

These functions also obey the following surprising relations:

$$\sum_{i=1}^n G_n(x_i; x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) = \begin{cases} 0 & n \text{ even} \\ (-1)^{\frac{n-1}{2}} G_1(x_1) \dots G_1(x_n) & n \text{ odd.} \end{cases} \quad (4.3)$$

The next step is to constrain the form of the finite term for the numerator. Again as in the lower point cases, to preserve the scaling in the spin variable of the three-point amplitudes, we require that the factors of a present in the denominators of the

G functions cancel out with the elements in the monomials. In particular, if we define the degree of an object, *e.g.*


$$\deg(G_r) = r, \quad \deg(\cosh(x_j)) = 0, \quad \deg(a) = -1, \quad (4.4)$$

by its inverse scale order, (*i.e.*, the number of factors of $1/\lambda$ when taking $a \rightarrow \lambda a$ and $\lambda \rightarrow i\infty$). Thus, for a general number of points, the following general ansatz for contributions is conjectured:

$$\begin{cases} (v \cdot X \cdot a) \mathcal{P}^{(r-1)} \mathcal{M}^{(r)}(G_1, \dots, G_r) & \text{even} \\ \text{tr}(S \cdot X)_{G_1}, (p \cdot X \cdot S \cdot Y \cdot a) \mathcal{P}^{(r-2)} \mathcal{M}^{(r)}(G_1, \dots, G_r) & \text{odd} \end{cases} \quad (4.5)$$

where $\mathcal{M}^{(r)}$ denotes a product of functions $G_i(x)$ such that the total degree is r , and $\mathcal{P}^{(r)}$ is a product of monomials of the arbitrary scalar products $p \cdot Z \cdot a$ with total degree $-r$ (X, Y, Z all denote products of field strength tensors). The building blocks Eq. (4.5) are all the possible combinations that respect the scaling in a and that include the correct factors of v . These terms can be further constrained by considering parity requirements, so that for the even(odd) part, the ansatz should be even(odd) under the transformation $a \rightarrow -a$.

After constructing the ansatz, the free parameters are fixed by the massless cut conditions:



$$(4.6)$$

The unique four and five points results indicate that our bootstrap procedure leads to a universal arbitrary-spin infinite-mass amplitude determined solely from exponentiated three-point amplitudes and factorization constraints. We expect this to be the case at any multiplicity.

5 Conclusion

This paper discusses a new systematic bootstrap procedure for the computation of the leading in mass classical Compton scattering of arbitrary spinning infinity mass particles and gluons/gravitons that agrees with linear and quadratic results published in [49, 72] but in principle extends to any order in the spin. We have demonstrated this procedure by deducing new explicit classical results for four and five-point processes

and discussed all multiplicity generalizations.

Our results for four- and five-point scattering are already suitable for concrete computations beyond this paper’s scope; in particular, it opens an avenue for efficient calculation of the gravitational attraction of spinning black holes. For instance, by considering the following type of diagrammatic contributions illustrated below

$$(5.1)$$

we can compute non-spinning observables at the tree and one-loop post-Minkowskian order, see *e.g.* [26, 73, 81] for examples using heavy-mass effective field theory or [27, 28] using an analogous diagrammatic formulation in terms of velocity cuts. This context is very suggestive of a generalization to arbitrary spin. A related question is working out a connection between the numerators explored with three legs fixed and the formalism with numerators with two legs fixed used discussed in [84, 85] using momentum kernel identities that link the two formalisms [86]. While we have only computed the leading in mass part of the Compton amplitude, it is interesting to extend the results of this paper to the total amplitude, considering that the spinning particle has finite mass. To handle such amplitudes, one has to go beyond the simple factorization procedure regarded in the infinite mass limit. The reason is for high spin fields, where the mass is finite, there are different non-trivial effects from the complete spin propagator, which is a consequence of the massless particles interacting with the spinning particle and thereby altering its angular momentum. We will discuss this question further in appendix A.

It is also an intriguing question to produce a Lagrangian formalism [46, 47, 87–90] and kinematic algebra [73, 78, 82, 83, 91–104] for numerators that potentially could supply a general and fundamental framework for Compton scattering of Kerr black holes valid to all orders in spin. This question is beyond this presentation, so we leave it for the future.

6 Acknowledgements

G.C. would like to thank Jung-Wook Kim for several profound discussions. We also thank Marco Chiodaroli, Poul H. Damgaard, Henrik Johansson, Kays Haddad, Yutin Huang and Alexander Ochirov for their comments. The work of N.E.J.B.-B. and G.C. was supported by DFF grant 1026-00077B and in part by the Carlsberg Foundation.

G.C. has received funding from the European Union Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No. 847523 INTERACTIONS. M.S. was supported in part by the Stefan and Hanna Kobylinski Rozental Foundation.

A Classical spin theory with finite mass

Extending the amplitudes for classically spinning particles beyond the infinite mass limit is highly non-trivial since all the possible spin states for the internal particles in the propagators have to be summed over. Such effects are sub-leading when doing an expansion in the mass m [45, 72], which is why we did not consider them previously. However, for the sake of clarity, we will compute the piece of the full expression with a non-zero contribution in the HEFT.

We start at three points, where the numerator has been calculated [34]

$$[\mathcal{N}_a(1, \bar{2}, \bar{3})] = [w_1^{(3)}] \cdot \varepsilon_1. \quad (\text{A.1})$$

Here $([w_i^{(j)}])^\mu = [\cosh(x_i)] p_j^\mu - i [\frac{\sinh(x_i)}{x_i}] (p_i \cdot S_j)^\mu$ and $S_j^{\mu\nu} = -\epsilon^{\mu\nu\rho\sigma} (p_j)_\rho a_\sigma$. The $[]$ denotes formally truncated pieces in spin order. In the following, we will omit all the $[]$ symbols and work under the assumption that we have formally expanded to some finite order. As in the main text, we denote all massive particles with a bar.

For the four-point amplitude, one needs to consider the propagator of the finite spin theory, as studied in [72, 105]. However, if we are only interested in the pieces of the amplitude which contribute to the heavy mass limit, then we can separate it as

$$A_a^{(\text{total})}(1, 2, \bar{3}, \bar{4}) = A_a(1, 2, \bar{3}, \bar{4}) + \text{sub-leading finite spin effect}, \quad (\text{A.2})$$

where the additional contribution is sub-dominant in the $1/m$ expansion. Assuming that $A_a(1, 2, \bar{3}, \bar{4})$ satisfies the color-kinematics duality, we can write it in numerator form

$$A_a(1, 2, \bar{3}, \bar{4}) = \frac{\mathcal{N}_a(1, 2, \bar{3}, \bar{4})}{p_{12}^2}, \quad (\text{A.3})$$

and we will here construct the color-kinematic numerator directly.

First, we note that from the factorization behavior in the massive pole, we have

$$\mathcal{N}_a(1, 2, \bar{3}, \bar{4}) = -\frac{w_1^{(4)} \cdot F_1 \cdot F_2 \cdot w_2^{(4)}}{p_4 \cdot p_1} + \frac{p_4 \cdot p_2}{p_4 \cdot p_1} \mathcal{N}'_1(1, 2, \bar{3}, \bar{4}) + \mathcal{N}'_2(1, 2, \bar{3}, \bar{4}). \quad (\text{A.4})$$

It is then easy to see that the first term factorizes into a product of two 3-point amplitudes in the cut

$$-\frac{w_1^{(4)} \cdot F_1 \cdot F_2 \cdot w_2^{(4)}}{p_4 \cdot p_1} \xrightarrow{p_4 \cdot p_1 \rightarrow 0} \frac{p_1 \cdot p_2}{p_4 \cdot p_1} w_1^{(4)} \cdot \varepsilon_1 w_2^{(4)} \cdot \varepsilon_2. \quad (\text{A.5})$$

The second term is a new non-trivial contribution from the finite spin effect. The factor $p_4 \cdot p_2 / p_4 \cdot p_1$ is the only possible object that transforms into a non-singular function in the heavy mass limit, *i.e.*, $p_4 \cdot p_2 \rightarrow v \cdot p_2 = -v \cdot p_1$. We see that the last term fixes the factorization behavior for the massless pole term.

Similarly to what happens in the heavy mass limit, the denominator of the singular term cancels when taking the cut $p_1 \cdot p_2 \rightarrow 0$

$$-\frac{w_1^{(4)} \cdot F_1 \cdot F_2 \cdot w_2^{(4)}}{p_4 \cdot p_1} \Big|_{p_1 \cdot p_2 \rightarrow 0} = -\cosh(x_1) \varepsilon_1 \cdot F_2 \cdot w_2^{(4)} + w_1^{(4)} \cdot \varepsilon_1 p_1 \cdot \varepsilon_2 \cosh(x_2). \quad (\text{A.6})$$

It follows that from the factorization behavior on the massless pole,

$$\mathcal{N}_a(1, 2, \bar{3}, \bar{4})|_{p_1 \cdot p_2 \rightarrow 0} = -\varepsilon_1 \cdot F_2 \cdot w_{12}^{(4)} + w_{12}^{(4)} \cdot \varepsilon_1 p_1 \cdot \varepsilon_2, \quad (\text{A.7})$$

we have

$$\begin{aligned} & -\mathcal{N}'_1(1, 2, \bar{3}, \bar{4}) + \mathcal{N}'_2(1, 2, \bar{3}, \bar{4})|_{p_1 \cdot p_2 \rightarrow 0} \quad (\text{A.8}) \\ & = -\varepsilon_1 \cdot F_2 \cdot w_{12}^{(4)} + w_{12}^{(4)} \cdot \varepsilon_1 p_1 \cdot \varepsilon_2 - \left(w_1^{(4)} \cdot \varepsilon_1 p_1 \cdot \varepsilon_2 \cosh(x_2) - \cosh(x_1) \varepsilon_1 \cdot F_2 \cdot w_2^{(4)} \right) \\ & = (a \cdot p_2 a \cdot p_1 \varepsilon_1 \cdot F_2 \cdot p_4 - a \cdot p_2 a \cdot p_1 p_1 \cdot \varepsilon_2 \varepsilon_1 \cdot p_4) G_1(x_1) G_1(x_2) \\ & + i(-\varepsilon_1 \cdot F_2 \cdot S_4 \cdot p_1 + p_1 \cdot \varepsilon_2 \varepsilon_1 \cdot S_4 \cdot p_2) G_1(x_{12}) \\ & + i(a \cdot p_1 \varepsilon_1 \cdot F_2 \cdot S_4 \cdot p_2 + p_1 \cdot \varepsilon_2 a \cdot p_2 \varepsilon \cdot S_4 \cdot p_1) G_2(x_1; x_2). \quad (\text{A.9}) \end{aligned}$$

Now, one can show that to satisfy the color-kinematics duality (*i.e.*, the Jacobi relations between the half-ladder numerators), the following must hold

$$\mathcal{N}'_1(1, 2, \bar{3}, \bar{4}) = -\mathcal{N}'_2(1, 2, \bar{3}, \bar{4}) \equiv \frac{1}{2} \mathcal{N}'_a(1, 2, \bar{3}, \bar{4}), \quad (\text{A.10})$$

so that

$$\begin{aligned} \mathcal{N}'_a(1, 2, \bar{3}, \bar{4}) & = \left(\frac{p_4 \cdot p_2 - p_4 \cdot p_1}{2} a \cdot F_1 \cdot F_2 \cdot a + a \cdot F_1 \cdot p_2 a \cdot F_2 \cdot p_4 \right. \\ & \quad \left. - a \cdot F_2 \cdot p_1 a \cdot F_1 \cdot p_4 \right) G_1(x_1) G_1(x_2) \quad (\text{A.11}) \\ & + i(a \cdot F_1 \cdot F_2 \cdot S_4 \cdot p_2 + a \cdot F_2 \cdot F_1 \cdot S_4 \cdot p_1) G_2(x_1; x_2) \\ & + i \operatorname{tr}(F_1 \cdot F_2 \cdot S_4) G_1(x_{12}) \quad (\text{A.12}) \end{aligned}$$

is determined by requirements of gauge invariance and the factorization behavior of massless pole terms. We conclude by expressing the full numerators as follows

$$\mathcal{N}_a(1, 2, \bar{3}, \bar{4}) = -\frac{w_1^{(4)} \cdot F_1 \cdot F_2 \cdot w_2^{(4)}}{p_4 \cdot p_1} + \frac{p_4 \cdot p_1 - p_4 \cdot p_2}{2p_4 \cdot p_1} \mathcal{N}'_a(1, 2, \bar{3}, \bar{4}). \quad (\text{A.13})$$

A direct check of the color-kinematic duality is from the BCJ relations with a massive particle [106]

$$p_4 \cdot p_1 A_a(1, 2, \bar{3}, \bar{4}) = p_4 \cdot p_2 A_a(2, 1, \bar{3}, \bar{4}). \quad (\text{A.14})$$

Finally, we obtain the gravity amplitude by performing the double copy with a scalar Compton amplitude [49, 80] yielding

$$M_a(1, 2, \bar{3}, \bar{4}) = -\frac{1}{p_{12}^2} \mathcal{N}_a(1, 2, \bar{3}, \bar{4}) \mathcal{N}_0(1, 2, \bar{4}, \bar{3}), \quad (\text{A.15})$$

where \mathcal{N}_0 is the color-kinematic numerator for the interaction of a massive scalar line with two gluons [26, 94]:

$$\mathcal{N}_0(1, 2, \bar{4}, \bar{3}) = -\frac{p_3 \cdot F_1 \cdot F_2 \cdot p_3}{p_3 \cdot p_1}. \quad (\text{A.16})$$

We can check the consistency of the derived numerators and their corresponding infinite mass effective theory expressions. We see that

$$\begin{aligned} \mathcal{N}_0(1, 2, \bar{4}, \bar{3})|_{m \rightarrow \infty} &= -\mathcal{N}_0([1, 2], v) \\ \mathcal{N}_a(1, 2, \bar{3}, \bar{4})|_{m \rightarrow \infty} &= \mathcal{N}_a([1, 2], v) \end{aligned} \quad (\text{A.17})$$

where $\mathcal{N}_a([1, 2], v)$ is given in Eq. (3.3) and Eq. (3.8), and $\mathcal{N}_0([1, 2], v)$ by Eq. (3.11).

In order to see that this amplitude has the correct factorization behaviour on the massive cut, we first take the limit $p_4 \cdot p_1 \rightarrow 0$ in Eq. (A.15), and only then perform an expansion in $1/m$, obtaining the following result:

$$\begin{aligned} M_a(1, 2, \bar{3}, \bar{4}) &\xrightarrow{p_4 \cdot p_1 \rightarrow 0} \frac{\left(p_4 \cdot \varepsilon_1 p_4 \cdot \varepsilon_2 + \mathcal{O}\left(\frac{1}{m}\right)\right) \left(w_1^{(4)} \cdot \varepsilon_1 w_2^{(4)} \cdot \varepsilon_2 + \mathcal{O}\left(\frac{1}{m}\right)\right)}{p_4 \cdot p_1} \\ &\xrightarrow{m \rightarrow \infty} m^3 \frac{M_a(1, v) M_a(2, v)}{v \cdot p_1}. \end{aligned} \quad (\text{A.18})$$

This is consistent with our reasoning in section 2.2 since, taking into account the regulators $i\varepsilon$, the decomposition of the amplitude contains a term of order $\mathcal{O}(m^3)$ where the massive cut is imposed by a δ -function. For example, the bottom graph in Eq. (5.1) is deduced from

$$\begin{array}{c} \begin{array}{ccc} 1 & 2 & \\ \text{wavy} & \text{wavy} & \\ \bullet & \bullet & \\ \hline \bar{4} & & \bar{3} \end{array} + \begin{array}{ccc} 1 & 2 & \\ \text{wavy} & \text{wavy} & \\ \bullet & \bullet & \\ \hline \bar{4} & & \bar{3} \end{array} = \begin{array}{ccc} 1 & 2 & \\ \text{wavy} & \text{wavy} & \\ \bullet & | & \bullet \\ \hline \bar{4} & & \bar{3} \end{array} . \end{array} \quad (\text{A.19})$$

References

- [1] Y. Iwasaki, *Quantum theory of gravitation vs. classical theory - fourth-order potential*, *Prog. Theor. Phys.* **46** (1971) 1587.
- [2] J.F. Donoghue, *General relativity as an effective field theory: The leading quantum corrections*, *Phys. Rev. D* **50** (1994) 3874 [[gr-qc/9405057](#)].
- [3] N.E.J. Bjerrum-Bohr, J.F. Donoghue and B.R. Holstein, *Quantum gravitational corrections to the nonrelativistic scattering potential of two masses*, *Phys. Rev. D* **67** (2003) 084033 [[hep-th/0211072](#)].
- [4] B.R. Holstein and J.F. Donoghue, *Classical physics and quantum loops*, *Phys. Rev. Lett.* **93** (2004) 201602 [[hep-th/0405239](#)].
- [5] B.R. Holstein and A. Ross, *Spin Effects in Long Range Gravitational Scattering*, [0802.0716](#).
- [6] D. Neill and I.Z. Rothstein, *Classical Space-Times from the S Matrix*, *Nucl. Phys.* **B877** (2013) 177 [[1304.7263](#)].
- [7] N.E.J. Bjerrum-Bohr, J.F. Donoghue and P. Vanhove, *On-shell Techniques and Universal Results in Quantum Gravity*, *JHEP* **02** (2014) 111 [[1309.0804](#)].
- [8] N.E.J. Bjerrum-Bohr, P.H. Damgaard, G. Festuccia, L. Plante and P. Vanhove, *General Relativity from Scattering Amplitudes*, *Phys. Rev. Lett.* **121** (2018) 171601 [[1806.04920](#)].
- [9] C. Cheung, I.Z. Rothstein and M.P. Solon, *From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion*, *Phys. Rev. Lett.* **121** (2018) 251101 [[1808.02489](#)].
- [10] T. Damour, *High-energy gravitational scattering and the general relativistic two-body problem*, *Phys. Rev.* **D97** (2018) 044038 [[1710.10599](#)].
- [11] D.A. Kosower, B. Maybee and D. O'Connell, *Amplitudes, Observables, and Classical Scattering*, *JHEP* **02** (2019) 137 [[1811.10950](#)].
- [12] A. Cristofoli, N.E.J. Bjerrum-Bohr, P.H. Damgaard and P. Vanhove, *Post-Minkowskian Hamiltonians in general relativity*, *Phys. Rev. D* **100** (2019) 084040 [[1906.01579](#)].
- [13] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order*, *Phys. Rev. Lett.* **122** (2019) 201603 [[1901.04424](#)].
- [14] A. Antonelli, A. Buonanno, J. Steinhoff, M. van de Meent and J. Vines, *Energetics of two-body Hamiltonians in post-Minkowskian gravity*, *Phys. Rev. D* **99** (2019) 104004 [[1901.07102](#)].

- [15] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M.P. Solon and M. Zeng, *Black Hole Binary Dynamics from the Double Copy and Effective Theory*, *JHEP* **10** (2019) 206 [[1908.01493](#)].
- [16] J. Parra-Martinez, M.S. Ruf and M. Zeng, *Extremal black hole scattering at $\mathcal{O}(G^3)$: graviton dominance, eikonal exponentiation, and differential equations*, *JHEP* **11** (2020) 023 [[2005.04236](#)].
- [17] P. Di Vecchia, C. Heissenberg, R. Russo and G. Veneziano, *Universality of ultra-relativistic gravitational scattering*, *Phys. Lett. B* **811** (2020) 135924 [[2008.12743](#)].
- [18] T. Damour, *Radiative contribution to classical gravitational scattering at the third order in G* , *Phys. Rev. D* **102** (2020) 124008 [[2010.01641](#)].
- [19] G. Kälin, Z. Liu and R.A. Porto, *Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach*, *Phys. Rev. Lett.* **125** (2020) 261103 [[2007.04977](#)].
- [20] P. Di Vecchia, C. Heissenberg, R. Russo and G. Veneziano, *Radiation Reaction from Soft Theorems*, *Phys. Lett. B* **818** (2021) 136379 [[2101.05772](#)].
- [21] P. Di Vecchia, C. Heissenberg, R. Russo and G. Veneziano, *The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$* , *JHEP* **07** (2021) 169 [[2104.03256](#)].
- [22] E. Herrmann, J. Parra-Martinez, M.S. Ruf and M. Zeng, *Radiative classical gravitational observables at $\mathcal{O}(G^3)$ from scattering amplitudes*, *JHEP* **10** (2021) 148 [[2104.03957](#)].
- [23] N.E.J. Bjerrum-Bohr, P.H. Damgaard, L. Planté and P. Vanhove, *Classical gravity from loop amplitudes*, *Phys. Rev. D* **104** (2021) 026009 [[2104.04510](#)].
- [24] N.E.J. Bjerrum-Bohr, P.H. Damgaard, L. Planté and P. Vanhove, *The Amplitude for Classical Gravitational Scattering at Third Post-Minkowskian Order*, [2105.05218](#).
- [25] P.H. Damgaard, L. Plante and P. Vanhove, *On an exponential representation of the gravitational S -matrix*, *JHEP* **11** (2021) 213 [[2107.12891](#)].
- [26] A. Brandhuber, G. Chen, G. Travaglini and C. Wen, *Classical gravitational scattering from a gauge-invariant double copy*, *JHEP* **10** (2021) 118 [[2108.04216](#)].
- [27] N.E.J. Bjerrum-Bohr, L. Planté and P. Vanhove, *Post-Minkowskian radial action from soft limits and velocity cuts*, *JHEP* **03** (2022) 071 [[2111.02976](#)].
- [28] N.E.J. Bjerrum-Bohr, L. Planté and P. Vanhove, *Effective Field Theory and Applications: Weak Field Observables from Scattering Amplitudes in Quantum Field Theory*, [2212.08957](#).
- [29] C. Dlapa, G. Kälin, Z. Liu and R.A. Porto, *Dynamics of binary systems to fourth*

- Post-Minkowskian order from the effective field theory approach*,
Phys. Lett. B **831** (2022) 137203 [2106.08276].
- [30] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al.,
Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$,
Phys. Rev. Lett. **126** (2021) 171601 [2101.07254].
- [31] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al.,
Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $\mathcal{O}(G^4)$,
Phys. Rev. Lett. **128** (2022) 161103 [2112.10750].
- [32] Z. Bern, J. Parra-Martinez, R. Roiban, M.S. Ruf, C.-H. Shen, M.P. Solon et al.,
Scattering amplitudes and conservative dynamics at the fourth post-Minkowskian order,
PoS **LL2022** (2022) 051.
- [33] A. Guevara, *Holomorphic Classical Limit for Spin Effects in Gravitational and Electromagnetic Scattering*,
JHEP **04** (2019) 033 [1706.02314].
- [34] N. Arkani-Hamed, T.-C. Huang and Y.-t. Huang, *Scattering amplitudes for all masses and spins*,
JHEP **11** (2021) 070 [1709.04891].
- [35] A. Guevara, A. Ochirov and J. Vines, *Scattering of Spinning Black Holes from Exponentiated Soft Factors*,
JHEP **09** (2019) 056 [1812.06895].
- [36] M.-Z. Chung, Y.-T. Huang, J.-W. Kim and S. Lee, *The simplest massive S -matrix: from minimal coupling to Black Holes*,
JHEP **04** (2019) 156 [1812.08752].
- [37] A. Guevara, A. Ochirov and J. Vines, *Black-hole scattering with general spin directions from minimal-coupling amplitudes*,
Phys. Rev. **D100** (2019) 104024 [1906.10071].
- [38] N. Arkani-Hamed, Y.-t. Huang and D. O’Connell, *Kerr black holes as elementary particles*,
JHEP **01** (2020) 046 [1906.10100].
- [39] R. Aoude, K. Haddad and A. Helset, *On-shell heavy particle effective theories*,
JHEP **05** (2020) 051 [2001.09164].
- [40] M.-Z. Chung, Y.-t. Huang, J.-W. Kim and S. Lee, *Complete Hamiltonian for spinning binary systems at first post-Minkowskian order*,
JHEP **05** (2020) 105 [2003.06600].
- [41] A. Guevara, B. Maybee, A. Ochirov, D. O’Connell and J. Vines, *A worldsheet for Kerr*,
JHEP **03** (2021) 201 [2012.11570].
- [42] W.-M. Chen, M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *The 2PM Hamiltonian for binary Kerr to quartic in spin*,
JHEP **08** (2022) 148 [2111.13639].
- [43] D. Kosmopoulos and A. Luna, *Quadratic-in-spin Hamiltonian at $\mathcal{O}(G^2)$ from scattering amplitudes*,
JHEP **07** (2021) 037 [2102.10137].
- [44] M. Chiodaroli, H. Johansson and P. Pichini, *Compton black-hole scattering for $s \leq 5/2$* ,
JHEP **02** (2022) 156 [2107.14779].

- [45] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *From Scattering in Black Hole Backgrounds to Higher-Spin Amplitudes: Part I*, [2107.10179](#).
- [46] L. Cangemi, M. Chiodaroli, H. Johansson, A. Ochirov, P. Pichini and E. Skvortsov, *Kerr Black Holes Enjoy Massive Higher-Spin Gauge Symmetry*, [2212.06120](#).
- [47] A. Ochirov and E. Skvortsov, *Chiral approach to massive higher spins*, *Physical Review Letters* **129** (2022) .
- [48] P.H. Damgaard, K. Haddad and A. Helset, *Heavy Black Hole Effective Theory*, *JHEP* **11** (2019) 070 [[1908.10308](#)].
- [49] Z. Bern, A. Luna, R. Roiban, C.-H. Shen and M. Zeng, *Spinning black hole binary dynamics, scattering amplitudes, and effective field theory*, *Phys. Rev. D* **104** (2021) 065014 [[2005.03071](#)].
- [50] G. Mogull, J. Plefka and J. Steinhoff, *Classical black hole scattering from a worldline quantum field theory*, *JHEP* **02** (2021) 048 [[2010.02865](#)].
- [51] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *SUSY in the sky with gravitons*, *JHEP* **01** (2022) 027 [[2109.04465](#)].
- [52] J. Vines, *Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings*, *Class. Quant. Grav.* **35** (2018) 084002 [[1709.06016](#)].
- [53] J. Vines, J. Steinhoff and A. Buonanno, *Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order*, *Phys. Rev. D* **99** (2019) 064054 [[1812.00956](#)].
- [54] B. Maybee, D. O’Connell and J. Vines, *Observables and amplitudes for spinning particles and black holes*, *JHEP* **12** (2019) 156 [[1906.09260](#)].
- [55] K. Haddad, *Exponentiation of the leading eikonal phase with spin*, *Phys. Rev. D* **105** (2022) 026004 [[2109.04427](#)].
- [56] G.U. Jakobsen, G. Mogull, J. Plefka and J. Steinhoff, *Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **126** (2021) 201103 [[2101.12688](#)].
- [57] Z. Liu, R.A. Porto and Z. Yang, *Spin Effects in the Effective Field Theory Approach to Post-Minkowskian Conservative Dynamics*, *JHEP* **06** (2021) 012 [[2102.10059](#)].
- [58] W.-M. Chen, M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *Gravitational Faraday effect from on-shell amplitudes*, *JHEP* **12** (2022) 058 [[2205.07305](#)].
- [59] G.U. Jakobsen and G. Mogull, *Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory*, *Phys. Rev. Lett.* **128** (2022) 141102 [[2201.07778](#)].

- [60] G. Menezes and M. Sergola, *NLO deflections for spinning particles and Kerr black holes*, *JHEP* **10** (2022) 105 [[2205.11701](#)].
- [61] F. Febres Cordero, M. Kraus, G. Lin, M.S. Ruf and M. Zeng, *Conservative Binary Dynamics with a Spinning Black Hole at $O(G^3)$ from Scattering Amplitudes*, *Phys. Rev. Lett.* **130** (2023) 021601 [[2205.07357](#)].
- [62] F. Alessio and P. Di Vecchia, *Radiation reaction for spinning black-hole scattering*, *Phys. Lett. B* **832** (2022) 137258 [[2203.13272](#)].
- [63] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban and F. Teng, *Binary Dynamics Through the Fifth Power of Spin at $\mathcal{O}(G^2)$* , [2203.06202](#).
- [64] R. Aoude, K. Haddad and A. Helset, *Classical Gravitational Spinning-Spinless Scattering at $O(G^2S^\infty)$* , *Phys. Rev. Lett.* **129** (2022) 141102 [[2205.02809](#)].
- [65] R. Aoude, K. Haddad and A. Helset, *Searching for Kerr in the 2PM amplitude*, *JHEP* **07** (2022) 072 [[2203.06197](#)].
- [66] L. Singh and C. Hagen, *Lagrangian formulation for arbitrary spin. 1. The boson case*, *Phys. Rev. D* **9** (1974) 898.
- [67] Z. Bern, J.J.M. Carrasco and H. Johansson, *New Relations for Gauge-Theory Amplitudes*, *Phys. Rev.* **D78** (2008) 085011 [[0805.3993](#)].
- [68] Z. Bern, J.J.M. Carrasco and H. Johansson, *Perturbative Quantum Gravity as a Double Copy of Gauge Theory*, *Phys. Rev. Lett.* **105** (2010) 061602 [[1004.0476](#)].
- [69] Z. Bern, J.J. Carrasco, M. Chiodaroli, H. Johansson and R. Roiban, *The Duality Between Color and Kinematics and its Applications*, [1909.01358](#).
- [70] G. Travaglini et al., *The SAGEX review on scattering amplitudes**, *J. Phys. A* **55** (2022) 443001 [[2203.13011](#)].
- [71] N.E.J. Bjerrum-Bohr, P.H. Damgaard, L. Plante and P. Vanhove, *Chapter 13: Post-Minkowskian expansion from scattering amplitudes*, *J. Phys. A* **55** (2022) 443014 [[2203.13024](#)].
- [72] Y.F. Bautista, A. Guevara, C. Kavanagh and J. Vines, *Scattering in Black Hole Backgrounds and Higher-Spin Amplitudes: Part II*, [2212.07965](#).
- [73] A. Brandhuber, G. Chen, H. Johansson, G. Travaglini and C. Wen, *Kinematic Hopf Algebra for Bern-Carrasco-Johansson Numerators in Heavy-Mass Effective Field Theory and Yang-Mills Theory*, *Phys. Rev. Lett.* **128** (2022) 121601 [[2111.15649](#)].
- [74] H. Georgi, *An Effective Field Theory for Heavy Quarks at Low-energies*, *Phys. Lett. B* **240** (1990) 447.
- [75] M.E. Luke and A.V. Manohar, *Reparametrization invariance constraints on heavy particle effective field theories*, *Phys. Lett. B* **286** (1992) 348 [[hep-ph/9205228](#)].

- [76] M. Neubert, *Heavy quark symmetry*, *Phys. Rept.* **245** (1994) 259 [[hep-ph/9306320](#)].
- [77] A.V. Manohar and M.B. Wise, *Heavy quark physics*, vol. 10, Cambridge University Press (2000).
- [78] A. Brandhuber, G. Chen, G. Travaglini and C. Wen, *A new gauge-invariant double copy for heavy-mass effective theory*, *JHEP* **07** (2021) 047 [[2104.11206](#)].
- [79] W.-M. Chen, M.-Z. Chung, Y.-t. Huang and J.-W. Kim, *The 2PM Hamiltonian for binary Kerr to quartic in spin*, *JHEP* **08** (2022) 148 [[2111.13639](#)].
- [80] H. Johansson and A. Ochirov, *Double copy for massive quantum particles with spin*, *JHEP* **09** (2019) 040 [[1906.12292](#)].
- [81] A. Brandhuber, G.R. Brown, G. Chen, J. Gowdy, G. Travaglini and C. Wen, *Amplitudes, Hopf algebras and the colour-kinematics duality*, *JHEP* **12** (2022) 101 [[2208.05886](#)].
- [82] G. Chen, H. Johansson, F. Teng and T. Wang, *On the kinematic algebra for BCJ numerators beyond the MHV sector*, *JHEP* **11** (2019) 055 [[1906.10683](#)].
- [83] G. Chen, H. Johansson, F. Teng and T. Wang, *Next-to-MHV Yang-Mills kinematic algebra*, *JHEP* **10** (2021) 042 [[2104.12726](#)].
- [84] N.E.J. Bjerrum-Bohr, A. Cristofoli, P.H. Damgaard and H. Gomez, *Scalar-Graviton Amplitudes*, *JHEP* **11** (2019) 148 [[1908.09755](#)].
- [85] N.E.J. Bjerrum-Bohr, T.V. Brown and H. Gomez, *Scattering of Gravitons and Spinning Massive States from Compact Numerators*, *JHEP* **04** (2021) 234 [[2011.10556](#)].
- [86] N.E.J. Bjerrum-Bohr, P.H. Damgaard, T. Sondergaard and P. Vanhove, *The Momentum Kernel of Gauge and Gravity Theories*, *JHEP* **01** (2011) 001 [[1010.3933](#)].
- [87] D. Sorokin, *Introduction to the classical theory of higher spins*, *AIP Conf. Proc.* **767** (2005) 172 [[hep-th/0405069](#)].
- [88] R.R. Metsaev, *Cubic interaction vertices of massive and massless higher spin fields*, *Nucl. Phys. B* **759** (2006) 147 [[hep-th/0512342](#)].
- [89] A. Fotopoulos and M. Tsulaia, *Gauge Invariant Lagrangians for Free and Interacting Higher Spin Fields. A Review of the BRST formulation*, *Int. J. Mod. Phys. A* **24** (2009) 1 [[0805.1346](#)].
- [90] I.L. Buchbinder and A. Reshetnyak, *General Lagrangian Formulation for Higher Spin Fields with Arbitrary Index Symmetry. I. Bosonic fields*, *Nucl. Phys. B* **862** (2012) 270 [[1110.5044](#)].
- [91] R. Monteiro and D. O’Connell, *The Kinematic Algebra From the Self-Dual Sector*, *JHEP* **07** (2011) 007 [[1105.2565](#)].

- [92] C. Cheung and C.-H. Shen, *Symmetry for Flavor-Kinematics Duality from an Action*, *Phys. Rev. Lett.* **118** (2017) 121601 [[1612.00868](#)].
- [93] M. Ben-Shahar and H. Johansson, *Off-shell color-kinematics duality for Chern-Simons*, *JHEP* **08** (2022) 035 [[2112.11452](#)].
- [94] G. Chen, G. Lin and C. Wen, *Kinematic Hopf algebra for amplitudes and form factors*, [2208.05519](#).
- [95] C.-H. Fu and K. Krasnov, *Colour-Kinematics duality and the Drinfeld double of the Lie algebra of diffeomorphisms*, *JHEP* **01** (2017) 075 [[1603.02033](#)].
- [96] M. Reiterer, *A homotopy BV algebra for Yang-Mills and color-kinematics*, [1912.03110](#).
- [97] M. Tolotti and S. Weinzierl, *Construction of an effective Yang-Mills Lagrangian with manifest BCJ duality*, *JHEP* **07** (2013) 111 [[1306.2975](#)].
- [98] S. Mizera, *Kinematic Jacobi Identity is a Residue Theorem: Geometry of Color-Kinematics Duality for Gauge and Gravity Amplitudes*, *Phys. Rev. Lett.* **124** (2020) 141601 [[1912.03397](#)].
- [99] L. Borsten, B. Jurčo, H. Kim, T. Macrelli, C. Saemann and M. Wolf, *Becchi-Rouet-Stora-Tyutin-Lagrangian Double Copy of Yang-Mills Theory*, *Phys. Rev. Lett.* **126** (2021) 191601 [[2007.13803](#)].
- [100] L. Borsten and S. Nagy, *The pure BRST Einstein-Hilbert Lagrangian from the double-copy to cubic order*, *JHEP* **07** (2020) 093 [[2004.14945](#)].
- [101] P. Ferrero and D. Francia, *On the Lagrangian formulation of the double copy to cubic order*, *JHEP* **02** (2021) 213 [[2012.00713](#)].
- [102] L. Borsten, H. Kim, B. Jurčo, T. Macrelli, C. Saemann and M. Wolf, *Double Copy from Homotopy Algebras*, *Fortsch. Phys.* **69** (2021) 2100075 [[2102.11390](#)].
- [103] C. Cheung and J. Mangan, *Covariant color-kinematics duality*, *JHEP* **11** (2021) 069 [[2108.02276](#)].
- [104] M. Ben-Shahar, L. Garozzo and H. Johansson, *Lagrangians Manifesting Color-Kinematics Duality in the NMHV Sector of Yang-Mills*, [2301.00233](#).
- [105] M.D. Scadron, *Covariant Propagators and Vertex Functions for Any Spin*, *Phys. Rev.* **165** (1968) 1640.
- [106] H. Johansson and A. Ochirov, *Color-Kinematics Duality for QCD Amplitudes*, *JHEP* **01** (2016) 170 [[1507.00332](#)].