

Anomaly-Induced Quenching of g_A in Nuclear Matter

Mannque Rho^{1,*}

¹ *Université Paris-Saclay, CNRS, CEA, Institut de Physique Théorique, 91191, Gif-sur-Yvette, France*
(Dated: February 2, 2023)

How to disentangle the possible quenching of g_A caused by scale anomaly of QCD parameterized by the “scale-symmetry-breaking” quenching factor q_{ssb} from nuclear correlation effects is described. This is done by “matching” the Fermi-liquid fixed point theory to the “extreme single particle (shell) model” (acronym ESPM) in superallowed Gamow-Teller transitions in heavy doubly-magic shell nuclei. The current observed indication for $(1 - q_{ssb}) \neq 0$ in certain experiments seems non-negligibly significant. How to confirm or refute this observation is discussed in terms of the superallowed Gamow-Teller transition in the doubly-magic nucleus ^{100}Sn and in the spectral shape in the multifold forbidden β decay of ^{115}In .

Introduction.— In a recent Letter article [1], it was argued that the long-standing “puzzle” of the quenching of the axial coupling constant g_A from the free-space value of $g_A = 1.276$ to $g_A^Q \simeq 1$ observed in Gamow-Teller transitions in light nuclei [2] is resolved almost entirely by nuclear correlations.¹ The quenching does not involve, if any, a significant “fundamental quenching” of the axial vector constant g_A , but it is the *single-particle* $\sigma\tau$ operator that is quenched, that is, purely nuclear correlation effect. This conclusion was arrived at by many authors as listed e.g. in the reviews [2]. To quote one example out of many, in 1977 Les Houches Lectures, Wilkinson concluded, based on his analysis, that with the shell-model wave-functions for light nuclei $A \leq 21$ that *include* “full mixing,” the “effective” g_{Ae} accounting for $g_A^Q = 1$ comes out to be [3]

$$g_{Ae}/g_A = 1 + (2 \pm 6) \quad (1)$$

implying a suspicion that there was nothing “fundamental” in g_A^Q going down to 1 from $g_A = 1.276$ listed in the Particle Physics Booklet. What was intriguing however was that the “preferred” g_A^Q was tantalizingly close to 1, resembling something fundamental like $g_V = 1$ in CVC. Indeed the modern high-power many-body calculations such as Quantum Monte Carlo calculations explain the β decay rates for light nuclei $A < 10$ with the free-space g_A , modulo some small corrections coming from two-body exchange currents [4].

The situation in heavier nuclei looks rather different, leaning toward (1) on the larger error bar side [2]. It will be argued that due to hidden symmetries in nuclear correlations, in the entire range of densities, g_{Ae}/g_A will be near 1 unless a so-far unsuspected quantum anomaly intervenes. This issue gets sharpened in the recent experimental result in the superallowed Gamow-Teller transition in the doubly magic nucleus ^{100}Sn [5] which appears to signal a significant renormalization of g_A in nuclear

medium *induced by the vacuum change by density*. If this is confirmed, it would seriously impact not only certain intrinsic properties of nuclear dynamics and astrophysical processes but also the searches for going beyond the Standard Model (BSM) such as in neutrinoless double β decay processes.

In this short note, we describe what the solution to the “ g_A puzzle,” $g_{Ae}/g_A = 1$ could be, what it implies how hidden (or spontaneously broken) scale symmetry manifests in nuclear medium and what fundamental information vis-à-vis with the quenched g_A it could encode.

What’s new in the solution to the g_A puzzle was that it involved a novel mechanism, so far totally unexplored, based on a symmetry hidden in QCD that “emerges” in strong nuclear correlations. It is argued that $g_A^Q = 1$ should apply not only to light nuclei as observed [2] but also to heavy nuclei as well as to dense matter: It is found to permeate from nuclear-matter density $n_0 \simeq 0.16 \text{ fm}^{-3}$ to high densities $n_{star} \sim 7n_0$ relevant to compact-star physics, and, more significantly, beyond to what is referred to as “dilaton-limit fixed point (DLFP)” $n_{dlfp} \gtrsim 25n_0$. The deviation of g_A^Q from 1 observed in the ^{100}Sn decay would, therefore, represent an anomaly in scale symmetry manifested specifically in nuclear medium.

Scale symmetry with the “genuine dilaton (GD)” in nuclear axial current.— We first describe how the scale (a.k.a. trace) anomaly of QCD can enter in the nuclear weak current. For this we adopt the scale symmetry in QCD proposed by Crewther and Tunstal [6, 7] that involves the notion of “genuine dilaton (GD)” which we believe is most appropriate to what’s being treated in nuclear medium.² The GD scheme is characterized by the existence of an infrared fixed point (IRFP) α_{IR} with the β function $\beta(\alpha_{IR}) = 0$ at which both scale symmetry and chiral symmetry (in the chiral limit) are realized in the Nambu-Goldstone (NG) mode populated by the massless NG bosons π and dilaton σ_d whose decay constants are

* mannque.rho@ipht.fr

¹ To avoid confusion, the “quenched g_A ” observed experimentally or given theoretically will be denoted by g_A^Q . The superscript with * will stand for density dependence.

² There is another approach involving “conformal dilaton (CD)” [8] that shares more or less the same properties with that of GD in the density regime we are dealing with, so we won’t distinguish the two schemes for this problem.

non-zero. What is characteristic of this notion and crucially relevant to us is that it accommodates the massive nucleons ψ and vector mesons V_μ at the IR fixed point. This scheme – and also CD [8] – is generally considered ruled out by those working on dilatonic Higgs theories for large N_f . The notion advocated here can be justified in nuclear medium where scale symmetry emerges from strong nuclear correlations.

Our approach is anchored on an effective field theory, the only reliable tool known to be available for the problem, given that what’s involved is nuclear-matter density inaccessible from both lattice and perturbative QCD. The relevant degrees of freedom that figure are, apart from the nucleons and NG bosons (π , σ_d), the vector mesons $V = (\rho, \omega)$ as flavor gauge fields. It therefore involves hidden local symmetry [9, 10] and hidden scale symmetry [6]. These are the minimal number of degrees of freedom that enable us to go with no serious conflict – up to now – with nature – from normal nuclear matter to, with the intervention of a topology change at $n_{1/2} \sim (2 - 3)n_0$, near the center of massive compact stars, $n \sim 6n_0$ [11].

It has been feasible to formulate an EFT with systematic power counting in both chiral and scale symmetries (“CS” for short) taken into account, both without [12] and with [13] the HLS and nucleon fields. In principle one should be able to address the g_A problem at high CS orders as one does with success in nuclear dynamics at $\lesssim n_0$ [14]. In practice this is not feasible at present with CS Lagrangians because there are too many unknown constants to control.

In fact we assert it is not needed for our purpose to go beyond the leading CS order. At our disposal is a powerful “effective” way to bypass the jungle of complications by treating nucleons on a Fermi surface.³ In renormalization-group (RG) approach to interacting fermions [15], one can transform the mean fields of the leading CS-order effective Lagrangian with hidden local symmetry and scale symmetry incorporated that we denote as $\mathcal{L}_{\psi\chi\text{HLS}}$ – with ψ standing for the nucleon fields, χ for the “conformal compensator field” for the dilaton $\chi = f_\chi e^{\sigma_d/f_\chi}$ and HLS for the vector fields – to the Landau-Fermi-liquid (LFL) fixed point theory of many-nucleon systems. At the fixed point, one obtains the nuclear matter at its equilibrium [16]. Now with the nucleons put on a Fermi sphere, the LFL fixed point approximation corresponds to taking $1/\bar{N}$ to zero where $\bar{N} = k_F/(\Lambda_{fs} - k_F)$ with Λ_{fs} the cut-off on top of the Fermi-surface measured with respect to the origin. It becomes more reliable as density increases. As a bonafide EFT, one can do higher-order corrections in $1/\bar{N}$ in what’s known as “ V_{lowK} RG-expansion” as done in finite nuclei. For the g_A problem explicit $1/\bar{N}$ corrections are found to be unnecessary.

It is easy to justify this approach in terms of DFT (density-functional theory) à la Hohenberg-Kohn theorem applied to nuclear matter. In fact the present LFL fixed point approach can be thought of as a refined version of Walecka’s relativistic mean field theory of nuclear matter [17] with the refinement brought in by the hidden symmetries and the intrinsic density dependence [18] via the dilaton condensate $\langle\chi\rangle$.⁴ The link to the Fermi-liquid structure of the Walecka’s model has been known since a long time [19].

Quenching of g_A in $GnEFT$.— The LFL fixed point approach briefly defined above, referred to as $GnEFT$ in our work, is the framework that was exploited to address compact-star physics at high density in [11]. Here we will focus on the nuclear axial current.

The power of the $GnEFT$, even when substantially simplified, has been that low-energy theorems, both iso-vector and iso-scalar, can be encoded in $\mathcal{L}_{\psi\chi\text{HLS}}$ -type Lagrangians and when applied to nuclear matter with suitable Ward identities, give very accurate results for nuclear EW response functions. For instance in a much simpler form, it has been shown [16] that for a quasiparticle on top of the Fermi surface, the iso-scalar orbital gyromagnetic ratio $g_l^0 = 1$ – with the nucleon mass undergoing the BR scaling – comes out to be consistent with the Kohn theorem [20] and the anomalous iso-vector gyromagnetic ratio δg_l^i , reproducing exactly the Migdal formula [21] in terms of the Landau-Migdal parameter F_1 , agrees exactly with the available Pb data. To the best of the author’s knowledge, there has not been as simple a derivation of, and with such a quantitative agreement with nature, for these and other EW functions in (heavy) nuclei obtained in standard nuclear chiral perturbative (NchiPT) approach [22].

We now apply this same formalism to the axial-vector transition in nuclei.

For this we consider the quasiparticle propagating with near zero momentum on top of the Fermi sea in interaction with charged pionic and vector-mesonic fluctuations on the surface. Via low-energy theorems, i.e., through the Goldberger-Treiman relation, the coupling constant involved can be written with the axial coupling g_A . In the same LFL fixed-point approximation applied in the EW case, the “effective” axial coupling of the quasiparticle can be calculated in the mean-field approximation as a Landau fixed-point quantity [16]

$$g_A^L = g_A q_{snc}^{\text{Landau}} \simeq 1 \quad (2)$$

with

$$q_{snc}^{\text{Landau}} = \left(1 - \frac{1}{3}\Phi^* \bar{F}_1^\pi\right)^{-2}. \quad (3)$$

³ The formation of which is considered to be a quantum critical phenomenon in condensed matter physics.

⁴ The BR-scaling masses of the vector mesons account for higher meson-field terms consistently with the symmetries involved – which are missed in nonlinear Walecka models.

Here

$$\Phi^* = f_\chi^*/f_\chi \quad (4)$$

is the BR scaling and \bar{F}_1^π is the pionic contribution to the Landau mass. As defined, the factor q_{snc}^{Landau} is to capture the full nuclear correlations (or full mixing in the sense of (1)). It can be unambiguously calculated and was found to be (with the currently accepted $g_A = 1.276$)

$$q_{snc}^{\text{Landau}} = g_A^L/g_A \simeq 1/g_A \simeq 0.78. \quad (5)$$

This factor turns out to be remarkably insensitive to density between $n = n_0/2$ and n_0 . We note that the underlying assumption for (2) is that N_c is large⁵, thus the double limiting process $N_c \rightarrow \infty$ and $\bar{N} \rightarrow \infty$ is invoked.

In the LFL fixed point approximation, the quenching of g_A to 1 by the factor (5) is accounted entirely by nuclear quasiparticle correlations on Fermi surface. Now the question that was left un-clarified in [16] was: How is this $g_A^L \simeq 1$ related to the quenching of g_A in nuclear processes, in particular, Gamow-Teller transitions in finite nuclei [2]? To answer this question, we look at the g_A problem in the nuclear axial response function to the external weak field \mathcal{W}_μ . For this we focus on the iso-vector nuclear axial current $J_{5\mu}^a$.

At the classical level, the nuclear axial current coupling to the external weak field in the CS Lagrangian is scale-invariant. However there is an anomalous dimension contribution from the trace anomaly of QCD that enters nonperturbatively at the leading-order (LO) chiral-scale perturbation expansion [7]. The full LO axial current is given by [6]

$$J_5^{a\mu} = g_A q_{\text{ssb}} \bar{\psi} \gamma^\mu \gamma_5 \frac{\tau^a}{2} \psi \quad (6)$$

where

$$q_{\text{ssb}}(\beta') = c_A + (1 - c_A) \left(\frac{\chi}{f_\chi} \right)^{\beta'}. \quad (7)$$

Here c_A is a constant that cannot be calculated perturbatively and β' is the derivative of the $\beta(\alpha_s)$ at the IR fixed point

$$\beta'|_{\alpha_s=\alpha_{\text{IR}}} > 0. \quad (8)$$

One sees that β' brings in an anomalous dimension, representing scale-symmetry explicit breaking, to the current.

We should mention that q_{ssb} cannot at present be calculated: β' is not known for $N_f \leq 3$ appropriate for nuclear physics and the coefficient c_A is not accessible perturbatively even in the matter-free vacuum. This forces us to rely on experiments to make any progress.

Now here is the principal point in our argument.

In the matter-free space, the vacuum expectation value (VeV) is $\langle \chi \rangle = f_\chi$, so if one ignores the fluctuating dilaton field that one is justified to do for the nucleon axial current where the dilaton does not figure at the leading order, one can set $q_{\text{ssb}} = 1$ and the current is scale symmetric: There is no β' effect. However in nuclear matter, $\langle \chi \rangle^* = f_\chi^* \neq f_\chi$ brings in scale symmetry breaking, both explicit and spontaneous, hence bringing in the β' dependence. Applied to nuclear matter, one then has the density-dependent factor that we will call ‘‘anomaly-induced quenching’’ (*AIQ* for short)

$$q_{\text{ssb}}^* = c_A + (1 - c_A) (\Phi^*)^{\beta'} \quad (9)$$

where

$$\Phi^* = f_\chi^*/f_\chi \simeq f_\pi^*/f_\pi \quad (10)$$

with $*$ standing for density dependence. Within the *GnEFT* scheme, $q_{\text{ssb}}^* < 1$ is the only ‘‘fundamental quenching’’ factor that can intervene.

We should mention that this *AIQ* was missing in the Goldberger-Treiman-type result in [16]. Setting $q_{\text{ssb}}^* = 1$ was referred to as *LOSS* (leading-order-scale-symmetric) approximation [11]. In fact its presence – and other β' effects – has never been observed or even suspected in the literature with an exception to be mentioned below.

ESPM in doubly-magic-shell nuclei.— In the absence of *reliable and full ab initio* numerical treatments of nuclear many-body problems for complex nuclei – apart possibly from very light nuclei $A < 10$, how to translate the LFL fixed-point result given above for nuclear matter to finite nuclei has not been obvious. However there is one case, we argue, where this translation is feasible and that is the superallowed Gamow-Teller transitions in doubly-magic-shell nuclei that are heavy enough to be treated as a nuclear matter.

The crucial point to note is that the LFL fixed-point limit in the Fermi-liquid system obtained above with $q_{\text{snc}}^{\text{Landau}}$ corresponds to a Gamow-Teller transition undergoing on the Fermi surface with the kinematics $\omega \rightarrow 0$, $q/\omega \rightarrow 0$ – where (q, ω) are the (momentum, energy) carried by the weak field. What is required for this is that the quasiparticle-quasihole bubble contributions entering at higher orders in $1/\bar{N}$ are as completely suppressed as possible. To see how the mapping of this description to that of the *ESPM* in doubly-magic-shell systems proceeds, the most illustrative case is the transition in the ^{100}Sn nucleus which has the proton and neutron shells completely filled at 50/50. There may be other nuclei of similar structure or perhaps even better but we will focus on this nucleus because it has the advantage of being the heaviest nucleus with the equal magic shells that has also been studied extensively both experimentally and theoretically [5, 23, 24] with the results directly relevant to the issue concerned.

The process involved is a pure superallowed GT transition of a proton (π) $\pi 0g_{9/2}$ in the completely filled orbital into a neutron (ν) in the empty spin-orbit partner, the $\nu 0g_{7/2}$ orbital of ^{100}In . This offers the most favorable

⁵ We are assuming that $N_c \rightarrow \infty$ is valid in nuclear medium as it is in the matter-free Goldberger-Treiman relation.

structure of the daughter state that is of a pure ($\nu g_{7/2}$) particle-($\pi g_{9/2}$) hole state to which the ESPM can be applied.

Evidence for AIQ.— We are now ready to make the prediction for the ^{100}Sn GT decay. Phrased in terms of the GT strength \mathcal{B}_{GT} defined in [5, 23], the Landau Fermi-liquid fixed point GT strength can be translated into the GT strength given in terms of the ESPM quantities

$$\mathcal{B}_{\text{GT}}^{\text{theory}} = \mathcal{B}_{\text{GT}}^{\text{EPSM}}(q_{\text{ssb}}q_{\text{snc}}^{\text{Landau}})^2 \approx 10.8q_{\text{ssb}}^2 \quad (11)$$

with $\mathcal{B}_{\text{GT}}^{\text{EPSM}} = 160/9$ and the full mixing factor given in the LFL fixed-point theory (5). In the absence of exact calculations, $\mathcal{B}_{\text{GT}}^{\text{theory}}$ of Eq. (11) should be replaced by a *reliable* experiment.

Let us see what Nature says.

Eschewing entering into detailed experimental data available involving nuclear model dependence, an adequate assessment of which is beyond the author’s capability, we will limit to two extreme cases: One referred to as GSI [23] signaling no *AIQ* and the other RIKEN [5] signaling an important *AIQ*.

For the GSI result, among various possibilities clouded by not necessarily trustful nuclear models, we limit to the analysis that zeroes-in on the daughter state of a pure ($\nu g_{7/2}$) particle-($\pi g_{9/2}$) hole configuration. The cited measured \mathcal{B}_{GT} is

$$\mathcal{B}_{\text{GT}}^{\text{GSI}} \approx 10. \quad (12)$$

From (11), this leads to $q_{\text{ssb}} \approx 1$. This would imply that the *AIQ* effect should be negligible, supporting $g_{Ae}/g_A = 1$ in (1).

We should stress that (12) corresponds to the β decay to a pure single daughter state⁶. Of course to what extent the daughter state is pure could very well be subject to debate, both theoretical and experimental – and there are discussions on the matter in [23] – but what seems unquestionable is that this result could be taken, at least, semi-quantitatively robust.

Now let’s turn to the more recent, what’s claimed to be “improved,” RIKEN result [5], which is in serious tension with the GSI result, (12)

$$\mathcal{B}_{\text{GT}}^{\text{RIKEN}} = 4.4_{-0.7}^{+0.9}. \quad (13)$$

Within the range of values involved, this implies

$$q_{\text{ssb}}^{\text{RIKEN}} \approx 0.58 - 0.69. \quad (14)$$

This result implies that the “fundamental” g_A in the axial current effective in nuclear medium defined by the chiral symmetry cutoff $\Lambda_{\text{chi}} \sim 4\pi f_\pi$ can be quenched from 1.276 to $\sim (0.74 - 0.88)$. This is a big *fundamental* quenching.

A legitimate question to ask at this point on such a drastic quenching is whether one cannot arrive at (13) by other mechanisms that do not require the *AIQ*. There may be several possibilities available in the literature but as far as the author is aware none seems viable as they stand. However one notable approach to be seriously considered is the *ab initio* calculation claimed to be a “first-principles resolution” of the g_A puzzle [25]. The strategy in this calculation – which is anchored on standard nuclear chiral EFT – is to tweak the resolution scale (or cut-off scale) in the EFT to shift the quenching effect from one-body to 2 or more-body currents to arrive at q_{snc} that requires no *AIQ*. It would be $q_{\text{snc}} \approx 0.76$ if (12) were the experimental value.⁷ As argued in [1], this strategy exploiting N^3LO chiral expansion terms is untenable unless there is justification to ignore the next-order (N^4LO) terms – and there is none. Furthermore N^4LO terms contain far too many unknown parameters to fix. This makes, we conclude, moot the assertion made in [25].

We now turn to another potentially serious indication for a possible *AIQ* quenching, this in the recent developments on the spectral shape of multifold forbidden β decay which is extremely sensitive to the g_A coupling. Comparing theoretical predictions to the experimental β spectrum, it has been argued [26] that the spectral shape of the multifold forbidden β decay of ^{115}In requires a quenching of g_A by a factor $q^{115\text{In}} \sim (0.65 - 0.75)$ with the range of the value accounting for the nuclear model dependence in calculating the matrix elements of the weak current involved in the spectrum shape. Suppose the axial matrix elements in the spectral shape were calculated with accuracy. Then one could take the quenching factor found in the measurement to be the *AIQ*, $q_{\text{ssb}}^{115\text{In}} \sim (0.65 - 0.75)$. This is comparable to the *AIQ* factor found in the RIKEN experiment.

There is however a caveat in arriving at this result. As noted by the authors of [26], the spectral shape and decay do not match. This is not surprising since the operators involved are of different structure. Unlike in the superallowed decay in ^{100}Sn where q_{snc} and q_{ssb} are disentangled from each other thanks to the hidden scale symmetry governing in the doubly magic shell structure, the one-body axial-current operator figuring in the spectral shape does not enjoy any known symmetry protection. Its matrix element could therefore be highly nuclear-model dependent [26].

Finally a few brief remarks on evidences against a large *AIQ* effect.

That g_{Ae}/g_A is near 1 in light nuclei but deviates from 1 in heavier nuclei is not impossible, but it is more or less ruled out in the formula for *AIQ* (9) unless $\beta' > 2$ – which appears most unlikely. Although q_{ssb} cannot be

⁶ The authors estimate that the decay takes place $\sim 95\%$ to this single dominant daughter state. We are taking this to be 100%.

⁷ It is not obvious that this strategy would succeed to arrive at the RIKEN result (14) without an *AIQ* factor. This matter was not unaddressed.

calculated at present, there is nothing to suggest a big density dependence. In fact $g_A^L \simeq 1$ seems to permeate from $\sim n_0$ to the dilaton-limit fixed point $\gtrsim 25n_0$ [11].

Another indication that an AIQ factor of the magnitude (14) is at odds with Nature comes from a precision calculation of ϵ_{MEC} . This quantity which determines the contribution from the soft-pion exchange 2-body contribution to the matrix element of the axial charge operator $J_{\gamma_5 0}^i$ has been measured and shown to be enhanced by a factor of ~ 2 in Pb nuclei [27]. This result is precisely reproduced in the framework of $GnEFT$ without the AIQ factor [28]. This Pb result is given a quantitative support in $A = 12$ nuclei where the density dependence in Φ is consistent with the BR scaling [29]. In fact this “precision calculation” would be ruined if g_A were quenched by the fundamental renormalization (14).

The enhancement factor ϵ_{MEC} obtained in [27, 29] has the merit of being nuclear-model independent. First being protected by the soft-pion theorems, the exchange current is totally model-independent. Furthermore the ratio of the 2-body matrix element to the leading 1-body term is also nuclear model-independent. The question could however remain as to the precision with which the one-body matrix element has been calculated. This could surely be checked in modern high-powered *ab initio* calculations.

The idea to exploit this enhanced $J_{\gamma_5 0}^i$ in first-forbidden transitions - in a stark contrast to the highly suppressed 2-body current for superallowed Gamow-Teller transitions - was studied in combination with the effect on the spectral shape in [30]. There seems to appear a hint for a quenched g_A but the disentanglement issue

discussed above may still have to be addressed.

Concluding remarks.— In this note we have shown by mapping the RG-based Landau Fermi-liquid fixed point approximation to the ESPM in heavy doubly-magic nuclei that what’s referred to as “quenched g_A ,” i.e., g_A^Q , in the literature consists of two factors: one q_{snc} given entirely by “snc” (strong nuclear correlations) and the other, q_{ssb} induced by the trace anomaly of QCD – which is invisible in the vacuum but exposed by density in nuclear medium. It was also shown that $q_{\text{snc}}g_A \rightarrow 1$ independent of density. While there is no clear evidence in nuclear processes for the presence of a fundamental quenching $q_{\text{ssb}} \neq 1$, there appear in the literature several experiments that show an indication for as much as $q_{\text{ssb}} \sim 0.6$. Such a big quenching in the fundamental coupling constant would make a game change not only in nuclear theory in general but also in the effort to go beyond the SM, such as $0\nu\beta\beta$ decay, in particular. A possible resolution to this problem is proposed: First establish q_{ssb} in the superallowed GT matrix element in doubly-magic nuclei, insert it in the analysis of the spectral shape of highly forbidden β decays to fix the nuclear model dependence (by fitting the spectral shape) and then employ the resulting nuclear model for the Gamow-Teller matrix elements in the $2\nu\beta\beta$ and $0\nu\beta\beta$ processes where non-negligible momentum transfer could be involved.

Acknowledgments

The author is grateful for useful and informative comments from Jouni Suhonen.

-
- [1] Y. L. Ma and M. Rho, Phys. Rev. Lett. **125**, 142501 (2020).
- [2] J.T. Suhonen, Rept. Prog. Phys. **80**, 046301 (2017).
- [3] D.H. Wilkinson, “Symmetries and Nuclei,” in *Nuclear Physics with Heavy Ions and Mesons* (North-Holland, Amsterdam, 1978) eds. by R. Balian, M Rho and G. Ripka).
- [4] B. King *et al.*, Phys. Rev. C **102**, 025501 (2020).
- [5] D. Lubos *et al.*, Phys. Rev. Lett. **122**, 222502 (2019).
- [6] R.J. Crewther, Universe **6**, 96 (2020).
- [7] R.J. Crewther and L.C. Tunstall, Phys. Rev. D **91**, 034016 (2015).
- [8] L. Del Debbio and R. Zwicky, JHEP **08**, 007 (2022).
- [9] M. Harada and K. Yamawaki, Phys. Rept. **381**, 1 (2003).
- [10] Z. Komargodski, JHEP **02**, 019 (2011).
- [11] Y. L. Ma and M. Rho, Prog. Part. Nucl. Phys. **113**, 103791 (2020).
- [12] O. Catà and C. Müller, Nucl. Phys. B **952**, 114938 (2020).
- [13] Y. L. Li, Y. L. Ma and M. Rho, Phys. Rev. D **95**, no.11, 114011 (2017).
- [14] H. W. Hammer, S. König and U. van Kolck, Rev. Mod. Phys. **92**, 025004 (2020).
- [15] R. Shankar, Rev. Mod. Phys. **66**, 129-192 (1994).
- [16] B. Friman and M. Rho, Nucl. Phys. A **606**, 303 (1996).
- [17] B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. **16**, 1 (1986).
- [18] G. E. Brown and M. Rho, Phys. Rev. Lett. **66**, 2720 (1991).
- [19] T. Matsui, Nucl. Phys. A **370**, 365 (1981).
- [20] W. Kohn, Phys. Rev. **123**, 1242 (1961).
- [21] A.B. Migdal, *Theory of Finite Systems and Applications to Finite Nuclei* (Interscience, London, 1967).
- [22] J. W. Holt, M. Rho and W. Weise, Phys. Rept. **621**, 2 (2016).
- [23] C.B. Hinke *et al.*, Nature **486**, 341 (2012).
- [24] T. Faestermann, M. Gorska and H. Grawe, Prog. Part. Nucl. Phys. **69**, 85 (2013).
- [25] P. Gysbers *et al.*, Nature Phys. **15**, 428 (2019).
- [26] A. F. Leder *et al.*, Phys. Rev. Lett. **129**, 232502 (2022).
- [27] E. K. Warburton, Phys. Rev. C **44**, 233 (1991).
- [28] K. Kubodera and M. Rho, Phys. Rev. Lett. **67**, 3479 (1991).
- [29] K. Minamisono *et al.*, Phys. Rev. C **65**, 015209 (2002).
- [30] J. Kostensalo and J. Suhonen, Phys. Lett. B **781**, 480 (2018).