

# HEAT-WAVE-SCHRÖDINGER TYPE EQUATIONS ON LOCALLY COMPACT GROUPS

SANTIAGO GÓMEZ COBOS, JOEL E. RESTREPO, AND MICHAEL RUZHANSKY

**ABSTRACT.** We prove that the noncommutative Lorentz norm (associated to a semifinite von Neumann algebra) of a propagator of the form  $\varphi(|\mathcal{L}|)$  can be estimated if the Borel function  $\varphi$  is bounded by a positive monotonically decreasing vanishing at infinity continuous function  $\psi$ . As a consequence we obtain the  $L^p - L^q$  ( $1 < p \leq 2 \leq q < +\infty$ ) norm estimates for the solutions of heat, wave and Schrödinger type equations (new in this setting) on a locally compact separable unimodular group  $G$  by using a non-local integro-differential operator in time and any positive left invariant operator (maybe unbounded and either with discrete or continuous spectrum) on  $G$ . We also provide asymptotic estimates (large-time behavior) for the solutions which in some cases can be claimed to be sharp. Illustrative examples are given for several groups.

## 1. INTRODUCTION

Regularity of integro-differential equations has been in the sights of mathematicians since long time ago not only for its wide range of applications, but also for its abstract relevance [21, 27, 34]. From the abstract point of view one can quickly identify that there are two promising ways to extend this kind of studies, on one hand one can try to study abstract equations on simpler spaces (the basic model is  $\mathbb{R}^n$ , see e.g. [6, 17, 22, 31]); or on the other hand one can work with some simpler equations on very general spaces ( $C^*$ -algebras, symmetric spaces, groups, etc, see e.g. [1, 11, 14]). In [2], Akylzhanov and the third author went one step further in the direction of the second option mentioned above by obtaining regularity in a quite general set up, considering linear evolution equations on locally compact unimodular groups; to describe the generality, the type of groups which can be considered is very wide. Some of them are compact, semisimple, exponential, nilpotent, some solvable ones, real algebraic, and many more. Moreover, the operators which can be involved are very general as well, and in particular one can use the following relevant ones: Laplacians, sublaplacians, (non)-Rockland operators, weighted subcoercive operators, etc. This was done through an application of a spectral multipliers boundedness theorem, which was obtained mainly due to the application of von Neumann algebras theory [13, 23] and the generalized non-commutative Lorentz spaces [20]. Using their result [2, Theorem 6.1] one can translate the regularity problem to the study of boundedness of propagators of the form  $\varphi(|\mathcal{L}|^1)$  for  $\varphi$  a positive monotonically decreasing vanishing at infinity continuous function, and if we are provided good

---

2010 *Mathematics Subject Classification.* 43A15, 43A85, 45K05, 35Q41.

*Key words and phrases.* Locally compact groups, heat type equations, wave type equations, Schrödinger type equations, asymptotic estimates, non-local operators.

<sup>1</sup>We denote by  $L = U|L|$  its polar decomposition.

asymptotic behaviour of the traces of the spectral projections  $E_{(s,+\infty)}(|\mathcal{L}|)$  we end up just imposing properties of the function  $\varphi$ . In [2], the main examples were functions  $e^{-tx}$  and  $1/(1+x)^\gamma$  for  $t, x, \gamma > 0$ , i.e. boundedness of solutions of the  $\mathcal{L}$ -heat equation and Sobolev embeddings, respectively. Obviously, one can not expect to be able to use any function  $\varphi$ , and in particular it does not work for  $\varphi(x) = e^{it\sqrt{x}}$  with  $t, x > 0$ , which is representing the propagator for the wave equation, hence it is not covered by these results as well as the Schrödinger one. Nevertheless, on a compact/graded Lie group, or a Hilbert space, in [10, 12], it was shown that for the time fractional  $\mathcal{L}$ -heat equation ( $\mathcal{L}$ -heat type) one can apply the previous reasoning since on those cases the propagator is related to the well-known Mittag-Leffler function (completely monotonic function).

Having this on mind the authors were motivated to extend the class of equations which can be studied in this way, and we managed to obtain the following refinement on the estimation of Lorentz norms of a general propagator:

**Theorem 1.1.** *Let  $L$  be a closed (maybe unbounded) operator affiliated<sup>2</sup> with a semifinite von Neuman algebra  $M$ . Let  $\phi$  be a Borel measurable function on  $[0, +\infty)$ . Suppose also that  $\psi$  is a monotonically decreasing continuous function on  $[0, +\infty)$  such that  $\psi(0) = 1$ ,  $\lim_{v \rightarrow +\infty} \psi(v) = 0$  and  $\phi(v) \leq \psi(v)$  for all  $v \in [0, +\infty)$ . Then for every  $1 \leq r < +\infty$  we have the inequality*

$$\|\phi(|L|)\|_{L^{r,\infty}(M)} \leq \sup_{v>0} \psi(v) [\tau(E_{(0,v)}(|L|))]^{\frac{1}{r}}.$$

This improvement allows us, at least for now, to treat the  $\mathcal{L}$ -wave type equation, by this we mean the wave equation but replacing the classical time derivative by the non-local operator  ${}^C\partial_t^\beta w(t, x)$ , the so-called Dzhrbashyan–Caputo fractional derivative, defined by  ${}^C\partial_t^\beta w(t, x) = {}^{RL}_0 I^{2-\beta} \partial_t^{(2)} w(t, x)$ , where  $\beta \in (1, 2)$  and  ${}^{RL}_0 I^\beta w(t, x) = \frac{1}{\Gamma(\beta)} \int_0^t (t-s)^{\beta-1} w(s, x) ds$ , is the Riemann-Liouville fractional integral of order  $\beta > 0$ . For  $\beta = 2$  we have the classical partial derivative in time, i.e.  ${}^C\partial_t^\beta w(t, x) = {}^{RL}_0 I^0 \partial_t^{(2)} w(t, x) = \partial_t^{(2)} w(t, x)$  since  ${}^{RL}_0 I^0$  acts like the identity operator. Notice also that for the range  $0 < \beta < 1$  we have  ${}^C\partial_t^\beta w(t, x) = {}^{RL}_0 I^{1-\beta} \partial_t w(t, x)$ , and  $\beta = 1$  coincides with the classical partial derivative in time  ${}^C\partial_t^\beta w(t, x) = \partial_t w(t, x)$ . Also, the theorem above can be applied for Schrödinger type equations, specifically

$$i {}^C\partial_t^\beta w(t, x) + \mathcal{L}w(t, x) = 0, \quad \text{for } 0 < \beta < 1.$$

These applications are complemented by the study of the  $\mathcal{L}$ -heat type equation, i.e.

$${}^C\partial_t^\beta w(t, x) + \mathcal{L}w(t, x) = 0, \quad \text{for } 0 < \beta < 1.$$

Notice that for the classical heat equation, the result was already obtained in [2].

For the reader's convenience we now plot the propagators (functions) for the wave type equation, which are given in terms of the two parametric Mittag-Leffler function  $E_{\alpha,\delta}(-x)$  for  $x > 0$  (see formula (2.7)), so that one can visualize the intrinsic idea of our main result. The functions of the propagators are of the form  $E_{\alpha,1}(-x)$  and  $E_{\alpha,2}(-x)$  for  $1 < \alpha < 2$ , see Section 3 for full details. The idea is to show a clear picture of the oscillating behaviour, for a particular  $\alpha$ , of these type of functions

---

<sup>2</sup>See the preliminaries for the definition.

which are always bounded uniformly by  $C/(1+x)$  for some positive constant  $C$  (see formula (2.8)):




FIGURE 1. Wave propagators functions for  $\alpha = 1.95$  bounded uniformly.

In Figure 2, one can also realize that for  $\alpha = 2$  we have an oscillating behaviour like cosine or sine, and does not vanish at infinity:




FIGURE 2. Classical wave propagator function (green) is not uniformly bounded by a decreasing vanishing at infinity function.

Also, we give a complementary perspective of the orders and behaviour of some other type of propagators associated with Mittag-Leffler functions which can be bounded uniformly:




FIGURE 3. Mittag-Leffler functions with one small parameter.

Let us now summarize the applications concerning the  $L^p - L^q$  ( $1 < p \leq 2 \leq q < +\infty$ ) norm estimates for  $\mathcal{L}$ -heat-wave-Schrödinger type equations. These results are new in this context and open a door to studying different estimates of other type of equations which, in principle, can not be calculated in the classical case but perhaps could be carried out in the set up of non-local operators. Also, we discuss existence, uniqueness and time-decay behaviour of such equations. Full details are given in Section 3.

Now we write explicitly the type of equations for which one can apply the main theorem above.

**$\mathcal{L}$ -heat-wave type equation:**

$$\int_0^t \frac{(t-s)^{n-\beta-1}}{\Gamma(n-\beta)} \partial_s^{(n)} w(s, x) ds - \mathcal{L}w(t, x) = 0, \quad t > 0, \quad x \in G, \quad (1.1)$$

$$w(t, x)|_{t=0} = w_0(x), \quad w_0 \in L^p(G), \quad (1.2)$$

$$\partial_t w(t, x)|_{t=0} = w_1(x), \quad w_1 \in L^p(G), \quad (1.3)$$

where  $0 < \beta < 2$  ( $\beta \neq 1$ ),  $n = \lfloor \beta \rfloor + 1$  ( $\lfloor \cdot \rfloor$  is the floor function),  $1 < p \leq 2$ ,  $\mathcal{L}$  is any positive linear left invariant operator on  $G$  (a locally compact separable unimodular group). If  $0 < \beta < 1$  we just consider the first initial condition  $w_0$ . While, for  $1 < \beta < 2$ , we use both initial conditions  $w_0, w_1$ . We exclude the value  $\beta = 1$  from consideration since this case is already known ([2]).

We also study the  **$\mathcal{L}$ -Schrödinger type equation:**

$$i \int_0^t \frac{(t-s)^{-\beta}}{\Gamma(1-\beta)} \partial_s w(s, x) ds - \mathcal{L}w(t, x) = 0, \quad t > 0, \quad x \in G, \quad 0 < \beta < 1, \quad (1.4)$$

$$w(t, x)|_{t=0} = w_0(x), \quad w_0 \in L^p(G). \quad (1.5)$$

**Theorem 1.2.** *Let  $G$  be a locally compact separable unimodular group and let  $1 < p \leq 2 \leq q < +\infty$ . Let  $\mathcal{L}$  be any positive left invariant operator on  $G$  (maybe unbounded).*

- If  $0 < \beta < 1$ ,  $w_0 \in L^p(G)$  and

$$\sup_{t>0} \sup_{s>0} [\tau(E_{(0,s)}(\mathcal{L}))]^{\frac{1}{p}-\frac{1}{q}} E_\beta(-t^\beta s) < +\infty,$$

*then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -heat type equation (1.1)-(1.2) given by*

$$w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x), \quad t > 0, \quad x \in G.$$

- If  $1 < \beta < 2$ ,  $w_0, w_1 \in L^p(G)$  and

$$\sup_{t>0} \sup_{s>0} \frac{[\tau(E_{(0,s)}(\mathcal{L}))]^{\frac{1}{p}-\frac{1}{q}}}{1+t^\beta s} < +\infty, \quad (1.6)$$

*then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -wave type equation (1.1)-(1.3) given by*

$$w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x) + {}^{RL}I_t^1 E_\beta(-t^\beta \mathcal{L})w_1(x), \quad t > 0, \quad x \in G.$$

- If  $w_0 \in L^p(G)$  and the condition (1.6) holds with  $0 < \beta < 1$ , then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -Schrödinger type equation (1.4)-(1.5) given by

$$w(t, x) = E_\beta(it^\beta \mathcal{L})w_0(x), \quad t > 0, x \in G.$$

In particular, if

$$\tau(E_{(0,s)}(\mathcal{L})) \lesssim s^\lambda, \quad s \rightarrow +\infty, \quad \text{for some } \lambda > 0,$$

we obtain the following time decay rate for the solutions of  $\mathcal{L}$ -heat type equation (1.1)-(1.2) and  $\mathcal{L}$ -Schrödinger type equation (1.4)-(1.5):

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\beta, \lambda, p, q} t^{-\beta\lambda(\frac{1}{p} - \frac{1}{q})} \|w_0\|_{L^p(G)},$$

and also for the  $\mathcal{L}$ -wave type equation (1.1)-(1.3):

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\beta, \lambda, p, q} t^{-\beta\lambda(\frac{1}{p} - \frac{1}{q})} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}),$$

whenever  $\frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}$ .

Preceding results are used to give different examples by choosing concrete operators and groups. Moreover, in some occasions, we are able to claim the sharpness of the time-decay rate. See Section 4.

## 2. PRELIMINARY RESULTS

In this section we recall definitions and some necessary results on integro-differential operators, abstract fractional differential equations and von Neumann algebras, which will appear frequently in this paper.

**2.1. Abstract integro-differential equations.** Here we recall some concepts with respect to the solution of an evolution integral equation. General and fundamental aspects of this theory can be found in e.g. the books [27, 34]. For a more specialized treatment of abstract fractional differential equations in this framework we can refer the works [4, 8].

**2.1.1. Integro-differential operators.** As usual, the Riemann–Liouville fractional integral of order  $\beta > 0$  ([30]) is defined by

$${}^{RL}_a I^\beta f(t) = \frac{1}{\Gamma(\beta)} \int_a^t (t-s)^{\beta-1} f(s) ds, \quad f \in L^1(a, T),$$

where  $L^1(a, T)$  is the Lebesgue integrable space on  $(a, T)$ .

Now we recall the well-known Dzhrbashyan–Caputo fractional derivative:

$${}_a^C D^\beta f(t) = {}^{RL}_a I^{n-\beta} f^{(n)}(t), \quad f \in AC^n[a, T], \quad n = \lfloor \beta \rfloor + 1, \quad (2.1)$$

where  $AC^n[a, T]$  is the set of functions such that  $f^{(n-1)}$  exists and is absolutely continuous on  $[a, T]$ .

Below we use another equivalent representation for operator (2.1). The new form of this operator is useful in applications since it can be rewritten utilizing the initial

conditions. For this reason, we first need to recall some suitable Sobolev type spaces [7, Appendix]:

$$W^{m,p}(I; X) := \left\{ g \middle/ \exists \phi \in L^p(I; X); \quad g(t) = \sum_{j=0}^{m-1} a_j \frac{t^j}{j!} + \frac{t^{m-1}}{(m-1)!} * \phi(t), \quad t \in I \right\},$$

where  $m \in \mathbb{N}$ ,  $1 \leq p < +\infty$  and  $I$  is an interval in  $\mathbb{R}$ . One can see that  $a_j = g^{(j)}(0)$  and  $\phi(t) = g^{(j)}(t)$ . So, if  $f \in C^{n-1}(I)$  and  $h_{n-\beta} * f \in W^{n,1}(I)$ , where  $n = \lfloor \beta \rfloor + 1$  and

$$h_\beta(t) := \begin{cases} \frac{t^{\beta-1}}{\Gamma(\beta)}, & t > 0, \\ 0, & t \leq 0, \end{cases}$$

then we get

$${}_a^C D^\beta f(t) = {}^{RL}_a D^\beta \left( f(t) - \sum_{k=0}^{n-1} \frac{f^{(k)}(a)}{k!} (t-a)^k \right), \quad (2.2)$$

where  ${}^{RL}_a D^\beta f(t) = D^n {}^{RL}_a I^{n-\beta} f(t)$  is the Riemann–Liouville fractional derivative. For more details, see e.g. [4, P. 10].

**2.1.2. Integro-differential equations.** Below we always consider the linear operator  $\mathcal{L}$  to be closed and densely defined in a Banach space  $X$ . We study the following Cauchy problem by means of the Dzhrbashyan–Caputo fractional derivative:

$${}_0^C \partial_t^\beta w(t) = \mathcal{L}w(t), \quad t > 0, \quad (2.3)$$

under the initial conditions

$$\left. \frac{\partial^j w(t)}{\partial t^j} \right|_{t=0+} = w_j \in X, \quad j = 0, 1, \dots, n-1, \quad (2.4)$$

where  $\beta > 0$  and  $n = 1 + \lfloor \beta \rfloor$ .

**Definition 2.1.** Let  $w$  be a function of the space  $C(\mathbb{R}^+, X)$ . It is said that  $w$  is a strong solution of equation (2.3)–(2.4) if  $w \in C(\mathbb{R}^+, X) \cap C^{n-1}(\mathbb{R}^+, X)$ ,

$${}^{RL}_0 I^{n-\beta} \left( w(t) - \sum_{j=0}^{n-1} \frac{\partial^j w(t)}{\partial t^j} \bigg|_{t=0+} \frac{t^j}{j!} \right) \in C^{n-1}(\mathbb{R}^+, X)$$

and equation (2.3)–(2.4) is satisfied on  $(\mathbb{R}^+, X)$ .

**Definition 2.2.** The fractional Cauchy problem (2.3)–(2.4) is said to be well-posed if the following two conditions are satisfied:

- (1) For any  $w_j \in \mathcal{D}(\mathcal{L})$ ,  $j = 0, \dots, n-1$ , there exists a unique strong solution  $w$  of equation (2.3)–(2.4).
- (2) For all sequences  $\{w_{k,m}\}_{m \geq 1} \subset \mathcal{D}(\mathcal{L})$  ( $k = 0, \dots, n-1$ ), such that  $w_{k,m} \rightarrow 0$  as  $m \rightarrow +\infty$ , imply  $w_m \rightarrow 0$  as  $m \rightarrow +\infty$  in  $X$ , uniformly on compact intervals, where  $w_m$  is the unique strong solution of equation (2.3)–(2.4) with initial conditions  $w_{0,m}, w_{1,m}, \dots, w_{n-1,m}$ .

We now show that the study of the solution of equation (2.3)-(2.4) can be reduced to the investigation of equation (2.3) under the following initial conditions:

$$w_0(t) \Big|_{t=0+} = w_0, \quad \frac{\partial^j w(t)}{\partial t^j} \Big|_{t=0+} = 0, \quad j = 1, \dots, n-1. \quad (2.5)$$

First of all, by [4, Formula (1.21)], one can see that equation (2.3),(2.5) is well-posed if and only if the following Volterra integral equation

$$w(t) = w_0(t) + {}^{RL}I^\beta(\mathcal{L}w(t)), \quad (2.6)$$

is well-posed in the sense of [27, Def. 1.2]. Hence, we are prepared to define the notion of a solution operator (we always assume exponentially bounded) of equation (2.3),(2.5) by means of its equivalent integral equation (2.6).

**Definition 2.3.** A family  $\{E_\beta(-t^\beta \mathcal{L})\}_{t \geq 0} \subset \mathcal{B}(X)$  is called a solution operator of equation (2.3),(2.5) if the following conditions are satisfied:

- (1)  $E_\beta(-t^\beta \mathcal{L})$  is strongly continuous for  $t \geq 0$  and  $E_\beta(0) = I$ ;
- (2)  $E_\beta(-t^\beta \mathcal{L})\mathcal{D}(\mathcal{L}) \subset \mathcal{D}(\mathcal{L})$  and  $\mathcal{L}E_\beta(-t^\beta \mathcal{L})w = E_\beta(-t^\beta \mathcal{L})\mathcal{L}w$  for any  $w \in \mathcal{D}(\mathcal{L})$ ,  $t \geq 0$ ;
- (3)  $E_\beta(-t^\beta \mathcal{L})w$  is a solution of equation (2.6) for any  $w \in \mathcal{D}(\mathcal{L})$ ,  $t \geq 0$ .

By using the same terminology of [27], equation (2.3),(2.5) is well-posed if and only if it has a solution operator. Therefore, if  $E_\beta(-t^\beta \mathcal{L})$  is a solution operator of equation (2.3),(2.5) then the general equation (2.3)-(2.4) is uniquely solvable by

$$w(t) = \sum_{j=0}^{n-1} {}^{RL}I^j(E_\beta(-t^\beta \mathcal{L}))w_j,$$

where  $w_j \in \mathcal{D}(\mathcal{L})$ ,  $j = 0, 1, \dots, n-1$ . So, it is well-posed. This means that is enough to study equation (2.3),(2.5) to know the solution of the general case (2.3)-(2.4).

The solution operator can be written in an integral form as [4, Chapters 2 and 3] (see also [27, Chapter 2] or [8, Theorem 2.41])

$$E_\beta(-t^\beta \mathcal{L}) = \frac{1}{2\pi i} \int_H e^{\gamma t} \gamma^{\beta-1} (\gamma^\beta + \mathcal{L})^{-1} d\gamma, \quad t \geq 0, \quad \beta \in (0, 2),$$

where  $H$  is the Hankel path of [8, Formula (2.5)] and it is contained in the resolvent set  $\rho(-\mathcal{L})$ . The above notation is consistent (convenient) since the integral representation of the Mittag-Leffler function  $E_\alpha(z) = \sum_{k=0}^{+\infty} \frac{z^k}{\Gamma(\alpha k + 1)}$  for  $z \in \mathbb{C}$  and  $\Re(\alpha) > 0$  is given by

$$E_\alpha(z) = \frac{1}{2\pi i} \int_{\mathcal{H}} e^{\gamma} \gamma^{\alpha-1} (\gamma^\alpha - z)^{-1} d\gamma,$$

where  $\mathcal{H}$  is a suitable Hankel path [15, Formula (3.4.12)].

In this paper, the two-parametric Mittag-Leffler function will appear frequently:

$$E_{\alpha,\delta}(z) = \sum_{k=0}^{+\infty} \frac{z^k}{\Gamma(\alpha k + \delta)}, \quad z, \delta \in \mathbb{C}, \quad \Re(\alpha) > 0, \quad (2.7)$$

which is absolutely and locally uniformly convergent for the given parameters ([15]).



Also, we need the following particular estimate from [26, Theorem 1.6]:

$$|E_{\alpha,\delta}(z)| \leq \frac{C}{1+|z|}, \quad z \in \mathbb{C}, \delta \in \mathbb{R}, \alpha < 2, \quad (2.8)$$

where  $\mu \leq |\arg(z)| \leq \pi$ ,  $\pi\alpha/2 < \mu < \min\{\pi, \pi\alpha\}$  and  $C$  is a positive constant.

**2.2. Von Neumann algebras.** The starting point of this algebras can be found in the well-known papers of von Neumann in [23, 24], where one builds the mathematical foundations for studying quantum mechanics. Let  $\mathfrak{L}(\mathcal{H})$  be the set of linear operators defined on a Hilbert space  $\mathcal{H}$ . Notice that the concept of  $\tau$ -measurability on a von Neumann algebra  $M$  and the technique of spectral projections allow us to approximate unbounded operators by bounded ones.

For our study we just need to set  $M$  to be the right group von Neumann algebra  $VN_R(G)$  with  $G$  being a locally compact separable unimodular group. The latter assumption gives the possibility to use the notion of noncommutative Lorentz spaces on  $M$  as given in [20]. For more specific details on von Neumann algebras we refer [13, 32].

It is important to recall that the group von Neumann algebra  $VN_R(G)$  is generated by all the right actions of  $G$  on  $L^2(G)$  ( $\pi_R(g)f(x) = f(xg)$  with  $g \in G$ ), which means that  $VN_R(G) = \{\pi_R(g)\}_{g \in G}^{!!}$ , where  $!!$  is the bicommutant of the subalgebra  $\{\pi_R(g)\}_{g \in G} \subset \mathfrak{L}(L^2(G))$ . The latter result is a consequence of the fact [13]:  $VN_R(G)^! = VN_L(G)$  and  $VN_L(G)^! = VN_R(G)$ , where the symbol  $!$  represents the commutant of the group.

Before recalling the noncommutative Lorentz spaces we need the following definitions.

**Definition 2.4.** Let  $M \subset \mathfrak{L}(\mathcal{H})$  be a semifinite von Neumann algebra acting over the Hilbert space  $\mathcal{H}$  with a trace  $\tau$ . A linear operator  $L$  (maybe unbounded) is said to be affiliated with  $M$ , if it commutes with the elements of the commutant  $M^!$  of  $M$ , which means  $LV = VL$  for all  $V \in M^!$ .

We highlight that in this paper we will consider operators affiliated with  $VN_R(G)$  (see [2, Def. 2.1]), which are precisely those ones who are left invariant on  $G$  [2, Remark 2.17].

**Remark 2.5.** If  $L$  is an affiliated bounded operator with  $M$ , then  $L \in M$  by the double commutant theorem.

So, the next definitions are given for the above type of operators.

**Definition 2.6.** Let  $M$  be a von Neumann algebra. A trace of the positive part  $M_+ = \{L \in M : L^* = L > 0\}$  of  $M$  is a functional defined on  $M_+$ , taking non-negative, possibly infinite, real values, with the following properties:

- (1) If  $L \in M_+$  and  $T \in M_+$  then  $\tau(L + T) = \tau(L) + \tau(T)$ ;
- (2) If  $L \in M_+$  and  $\gamma \in \mathbb{R}^+$  then  $\tau(\gamma L) = \gamma\tau(L)$  (with  $0 \cdot +\infty = 0$ );
- (3) If  $L \in M_+$  and  $U$  is an unitary operator on  $M$  then  $\tau(ULU^{-1}) = \tau(L)$ .

A trace  $\tau$  is faithful (or exact) if  $\tau(L) = 0$  ( $L \in M_+$ ) implies that  $L = 0$ . We have that  $\tau$  is finite if  $\tau(L) < +\infty$  for all  $L \in M_+$ . We have that  $\tau$  is semifinite if, for each



$L \in M_+$ ,  $\tau(L)$  is the supremum of the numbers  $\tau(T)$  over those  $T \in M_+$  such that  $L \leq T$  and  $\tau(T) < +\infty$ .

**Definition 2.7.** A closeable operator  $L$  (maybe unbounded) is called  $\tau$ -measurable if for each  $\epsilon > 0$  there exists a projection  $P$  in  $M$  such that  $P(\mathcal{H}) \subset D(L)$  and  $\tau(I - P) \leq \epsilon$ , where  $D(L)$  is the domain of  $L$  in  $\mathcal{H}$  and  $M \subset \mathfrak{L}(\mathcal{H})$ . We denote by  $S(M)$  the set of all  $\tau$ -measurable operators.

**Definition 2.8.** Let us take an operator  $L \in S(M)$  and let  $L = U|L|$  be its polar decomposition. We define the distribution function by  $d_\gamma(L) := \tau(E_{(\gamma, +\infty)}(|L|))$  for  $\gamma \geq 0$ , where  $E_{(\gamma, +\infty)}(|L|)$  is the spectral projection of  $|L|$  over the interval  $(\gamma, +\infty)$ . Also, for any  $t > 0$ , we define the generalized  $t$ -th singular numbers as

$$\mu_t(L) := \inf\{\gamma \geq 0 : d_\gamma(L) \leq t\}.$$

For more details and properties of the distribution function and generalized singular numbers, we recommend [33].

Below we recall the noncommutative Lorentz spaces associated with a semifinite von Neumann algebra  $M$ , which are a noncommutative extension of the classical Lorentz spaces [20].

**Definition 2.9.** We denote by  $L^{p,q}(M)$  ( $1 \leq p < +\infty$ ,  $1 \leq q < +\infty$ ) the set of all operators  $L \in S(M)$  such that

$$\|L\|_{L^{p,q}(M)} = \left( \int_0^{+\infty} (t^{1/p} \mu_t(L))^q \frac{dt}{t} \right)^{1/q} < +\infty.$$

Notice that the  $L^p$ -spaces on  $M$  can be defined by

$$\|L\|_{L^p(M)} := \|L\|_{L^{p,p}(M)} = \left( \int_0^{+\infty} \mu_t(L)^p dt \right)^{1/p}.$$

In the case  $q = \infty$ ,  $L^{p,\infty}(M)$  is the set of all operators  $L \in S(M)$  such that

$$\|L\|_{L^{p,\infty}(M)} = \sup_{t>0} t^{1/p} \mu_t(L).$$

### 3. MAIN RESULT AND SOME APPLICATIONS

In this section, we prove the main result of this paper and give some applications regarding some well-known integral equations.

**Theorem 3.1.** *Let  $L$  be a closed (maybe unbounded) operator affiliated with a semifinite von Neumann algebra  $M$ . Let  $\phi$  be a Borel measurable function on  $[0, +\infty)$ . Suppose also that  $\psi$  is a monotonically decreasing continuous function on  $[0, +\infty)$  such that  $\psi(0) = 1$ ,  $\lim_{v \rightarrow +\infty} \psi(v) = 0$  and  $\phi(v) \leq \psi(v)$  for all  $v \in [0, +\infty)$ . Then for every  $1 \leq r < \infty$  we have the inequality*

$$\|\phi(|L|)\|_{L^{r,\infty}(M)} \leq \sup_{v>0} \psi(v) [\tau(E_{(0,v)}(|L|))]^{\frac{1}{r}}.$$

*Proof.* By definition of the Lorentz space and [2, Proposition 2.9] we have

$$\|\phi(|L|)\|_{L^{r,\infty}(M)} = \sup_{s>0} s [\tau(E_{(s,+\infty)}(\phi(|L|)))]^{\frac{1}{r}}. \quad (3.1)$$

The Borel functional calculus [3] allow us to express the spectral projections as follows

$$E_{(s,+\infty)}(|L|) = \int_0^{+\infty} \chi_{(s,+\infty)}(t) dE_t(|L|) = \chi_{(s,+\infty)}(|L|),$$

therefore by [18, Corollary 5.6.29, p. 363]

$$E_{(s,+\infty)}(\phi(|L|)) = \chi_{(s,+\infty)}(\phi(|L|)) = (\chi_{(s,+\infty)} \circ \phi)(|L|) = \int_0^{+\infty} \chi_{(s,+\infty)}(\phi(t)) dE_t(|L|).$$

Using [2, Proposition 2.6] (see also [33]) we are able to calculate the trace of this operator, namely

$$\tau(E_{(s,+\infty)}(\phi(|L|))) = \int_0^{+\infty} \chi_{(s,+\infty)}(\phi(t)) d\mu_t,$$

where  $\mu_t = \tau(E_t(|L|))$  is a well-defined measure. Since  $\phi(v) \leq \psi(v)$  for all  $v \in [0, +\infty)$ , we have that  $\chi_{(s,+\infty)}(\phi(v)) \leq \chi_{(s,+\infty)}(\psi(v))$  for all  $v \in [0, +\infty)$ . As a consequence of the monotonicity of  $\mu_t$  we get

$$\tau(E_{(s,+\infty)}(\phi(|L|))) \leq \int_0^{+\infty} \chi_{(s,+\infty)}(\psi(t)) d\mu_t = \tau(E_{(s,+\infty)}(\psi(|L|))). \quad (3.2)$$

Putting (3.1) and (3.2) together we obtain

$$\|\phi(|L|)\|_{L^{r,\infty}(M)} \leq \sup_{s>0} s [\tau(E_{(s,+\infty)}(\psi(|L|)))]^{\frac{1}{r}}.$$

From the hypothesis given on  $\psi$ , the proof can be completed by mimicking the proof of [2, Theorem 6.1].  $\square$

**Remark 3.2.** The condition  $\psi(0) = 1$  in Theorem 3.1 can be rescaled, thus we can prove the same result if one has  $\psi(0) = \kappa_0 > 0$ .

In the next subsections we establish for heat, wave and Schrödinger type equations the  $L^p - L^q$  norm estimates for  $1 < p \leq 2 \leq q < +\infty$ . Indeed, we will see that the norm estimates can be reduced to the time asymptotic of the propagator (some of them oscillatory ones) in the noncommutative Lorentz space norm. Also, we show that the latter norm mainly involves to calculate the trace of the spectral projections of the operator  $\mathcal{L}$ .

In each equation below we assume that  $G$  is a locally compact separable unimodular group,  $\mathcal{L}$  is any positive linear left invariant operator on  $G$  (maybe unbounded) and  ${}^C\partial_t^\beta$  is the Dzhrbashyan-Caputo fractional derivative of order  $\beta$  from (2.1).

**3.1.  $\mathcal{L}$ -heat type equation.** We consider the following heat type equation:

$$\begin{aligned} {}^C\partial_t^\beta w(t, x) + \mathcal{L}w(t, x) &= 0, \quad t > 0, \quad x \in G, \quad 0 < \beta < 1, \\ w(t, x)|_{t=0} &= w_0(x). \end{aligned} \quad (3.3)$$

Our next result also covers the case  $\beta = 1$ , i.e. the classical  $\mathcal{L}$ -heat equation, but it is already known [2, Section 7].

**Theorem 3.3.** *Let  $G$  be a locally compact separable unimodular group,  $0 < \beta < 1$  and  $1 < p \leq 2 \leq q < +\infty$ . Let  $\mathcal{L}$  be any positive left invariant operator on  $G$  (maybe unbounded) such that*

$$\sup_{t>0} \sup_{s>0} [\tau(E_{(0,s)}(\mathcal{L}))]^{\frac{1}{p}-\frac{1}{q}} E_\beta(-t^\beta s) < +\infty. \quad (3.4)$$

*If  $w_0 \in L^p(G)$  then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -heat type equation (3.3) represented by*

$$w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x), \quad t > 0, \quad x \in G.$$

*In the particular case that*

$$\tau(E_{(0,s)}(\mathcal{L})) \lesssim s^\lambda, \quad s \rightarrow +\infty, \quad \text{for some } \lambda > 0, \quad (3.5)$$

*we get the following time decay rate for the solution of equation (3.3) for all  $t > 0$ ,*

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\beta, \lambda, p, q} t^{-\beta \lambda (\frac{1}{p} - \frac{1}{q})} \|w_0\|_{L^p(G)}, \quad \frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}.$$

*Proof.* Since the operator  $\mathcal{L}$  is positive we have that the solution operator of equation (3.3) is given by  $w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x)$  [8, Chapter 3] (see also [4, Prop. 3.8 and Def. 2.3]). So, by [2, Theorem 5.1], we have

$$\|w(t, \cdot)\|_{L^q(G)} = \|E_\beta(-t^\beta \mathcal{L})w_0\|_{L^q(G)} \lesssim \|E_\beta(-t^\beta \mathcal{L})\|_{L^{r, \infty}(VN_R(G))} \|w_0\|_{L^p(G)}, \quad (3.6)$$

with  $\frac{1}{r} = \frac{1}{p} - \frac{1}{q}$ , where the above Lorentzian norm is calculated as [2, Theorem 6.1]:

$$\|E_\beta(-t^\beta \mathcal{L})\|_{L^{r, \infty}(VN_R(G))} = \sup_{s>0} [\tau(E_{(0,s)}(\mathcal{L}))]^{\frac{1}{r}} E_\beta(-t^\beta s).$$

In the above result we are using that  $E_\beta(-t^\beta s)$  ( $t, s > 0$ ) is completely monotonic for all  $0 < \beta \leq 1$  ([25]) such that  $E_\beta(0) = 1$  and  $\lim_{s \rightarrow +\infty} E_\beta(-t^\beta s) = 0$  by the uniform estimate (2.8). Thus, by (3.5) and again estimate (2.8) we get

$$\|E_\beta(-t^\beta \mathcal{L})\|_{L^{r, \infty}(VN_R(G))} \lesssim \sup_{s>0} \frac{s^{\frac{\lambda}{r}}}{1 + t^\beta s}.$$

The above supremum is attained at  $s = \frac{\lambda t^{-\beta}}{r - \lambda}$  whenever  $\frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}$ , hence

$$\|E_\beta(-t^\beta \mathcal{L})\|_{L^{r, \infty}(VN_R(G))} \leq C_{\beta, \lambda, p, q} t^{-\beta \lambda / r}, \quad (3.7)$$

and then the result follows by (3.6).  $\square$

**3.2.  $\mathcal{L}$ -wave type equation.** Here we investigate the solution of the following equation, which in a sense interpolates between wave (without being wave,  $\beta < 2$ ) and heat types:

$$\begin{aligned} {}^C \partial_t^\beta w(t, x) + \mathcal{L}w(t, x) &= 0, \quad t > 0, \quad x \in G, \quad 1 < \beta < 2, \\ w(t, x)|_{t=0} &= w_0(x), \\ \partial_t w(t, x)|_{t=0} &= w_1(x). \end{aligned} \quad (3.8)$$

**Theorem 3.4.** *Let  $G$  be a locally compact separable unimodular group,  $1 < \beta < 2$  and  $1 < p \leq 2 \leq q < +\infty$ . Let  $\mathcal{L}$  be any positive left invariant operator on  $G$  (maybe unbounded) satisfying the following condition*

$$\sup_{t>0} \sup_{s>0} \frac{[\tau(E_{(0,s)}(\mathcal{L}))]^{\frac{1}{p}-\frac{1}{q}}}{1+t^\beta s} < +\infty. \quad (3.9)$$

*If  $w_0, w_1 \in L^p(G)$  then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -wave type equation (3.8) given by*

$$w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x) + {}^{RL}_0 I_t^1 E_\beta(-t^\beta \mathcal{L})w_1(x), \quad t > 0, x \in G.$$

*In particular, if the condition (3.5) holds then for any  $1 < p \leq 2 \leq q < +\infty$  such that  $\frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}$  we get the following time decay rate for the solution of equation (3.8):*

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\beta, \lambda, p, q} t^{-\beta \lambda (\frac{1}{p} - \frac{1}{q})} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}).$$

*Proof.* We know that the solution operator of equation (3.8) can be expressed as (see Subsection 2.1.2):

$$w(t, x) = E_\beta(-t^\beta \mathcal{L})w_0(x) + {}^{RL}_0 I_t^1 E_\beta(-t^\beta \mathcal{L})w_1(x). \quad (3.10)$$

Notice now that

$${}^{RL}_0 I_t^1 E_\beta(-t^\beta s) = \sum_{k=0}^{+\infty} \frac{(-s)^k}{\Gamma(\beta k + 1)} \frac{t^{\beta k + 1}}{\beta k + 1} = t \sum_{k=0}^{+\infty} \frac{(-st^\beta)^k}{\Gamma(\beta k + 2)} = t E_{\beta, 2}(-t^\beta s). \quad (3.11)$$

Taking the propagators of equation (3.10), equality (3.11), the condition (3.9), estimate (2.8), Theorem 3.1 and [2, Theorem 5.1], we obtain

$$\begin{aligned} \|w(t, \cdot)\|_{L^q(G)} &\leq \|E_\beta(-t^\beta \mathcal{L})w_0\|_{L^q(G)} + \|{}^{RL}_0 I_t^1 E_\beta(-t^\beta \mathcal{L})w_1\|_{L^q(G)} \\ &\lesssim \sup_{v>0} \frac{1}{1+vt^\beta} (\tau(E_{(0,v)}(\mathcal{L})))^{\frac{1}{p}-\frac{1}{q}} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}) \\ &\lesssim \sup_{v>0} \frac{v^{\lambda(\frac{1}{p}-\frac{1}{q})}}{1+vt^\beta} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}) \lesssim t^{-\beta \lambda (\frac{1}{p}-\frac{1}{q})} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}), \end{aligned}$$

completing the proof.  $\square$

**3.3.  $\mathcal{L}$ -Schrödinger type equation.** Consider the  $\mathcal{L}$ -Schrödinger type equation:

$$\begin{aligned} i {}^C \partial_t^\beta w(t, x) + \mathcal{L}w(t, x) &= 0, \quad t > 0, x \in G, \quad 0 < \beta < 1, \\ w(t, x)|_{t=0} &= w_0(x). \end{aligned} \quad (3.12)$$

**Theorem 3.5.** *Let  $0 < \beta < 1$  and  $1 < p \leq 2 \leq q < +\infty$ . Suppose that the operator  $\mathcal{L}$  satisfies the condition (3.9) with  $0 < \beta < 1$ . If  $w_0 \in L^p(G)$  then there exists a unique solution  $w \in \mathcal{C}([0, +\infty); L^q(G))$  for the  $\mathcal{L}$ -Schrödinger type equation (3.12) represented by*

$$w(t, x) = E_\beta(it^\beta \mathcal{L})w_0(x), \quad t > 0, x \in G.$$

*In particular, if the condition (3.5) holds then for any  $1 < p \leq 2 \leq q < +\infty$  such that  $\frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}$  we get the following time decay rate for the solution of equation (3.12) for all  $t > 0$ ,*

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\beta, \lambda, p, q} t^{-\beta \lambda (\frac{1}{p} - \frac{1}{q})} \|w_0\|_{L^p(G)}.$$

*Proof.* Notice that the solution operator of equation (3.12) is given by

$$w(t, x) = E_\beta(it^\beta \mathcal{L})w_0(x), \quad 0 < \beta < 1.$$

Hence, by [2, Theorem 5.1] we have

$$\|w(t, \cdot)\|_{L^q(G)} = \|E_\beta(it^\beta \mathcal{L})w_0\|_{L^q(G)} \lesssim \|E_\beta(it^\beta \mathcal{L})\|_{L^{r,\infty}(VN_R(G))} \|w_0\|_{L^p(G)}, \quad (3.13)$$

with  $\frac{1}{r} = \frac{1}{p} - \frac{1}{q}$ . Since the argument of the Mittag-Leffler function is a pure imaginary number then we can apply inequality (2.8) and get

$$|E_\beta(it^\beta s)| \leq \frac{C}{1 + t^\beta s}, \quad s, t \geq 0, \quad 0 < \beta < 1.$$

In this way, applying Theorem 3.1 with  $\psi(s) = C/(1 + t^\beta s)$  we finally obtain that

$$\|E_\beta(it^\beta \mathcal{L})\|_{L^{r,\infty}(VN_R(G))} \lesssim \sup_{s>0} \frac{s^{\frac{\lambda}{r}}}{1 + t^\beta s} \leq C_{\beta,\lambda,p,q} t^{-\beta\lambda/r}, \quad \frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q},$$

and thus the result follows by (3.13).  $\square$

**Remark 3.6.** We emphasize that Theorems 3.4, 3.5 do not cover the classical  $\mathcal{L}$ -wave equation ( $\beta = 2$ ) and  $\mathcal{L}$ -Schrödinger equation ( $\beta = 1$ ), respectively. For this case one would need a more delicate analysis of the problem, especially focusing on understanding the behavior of the trace of the spectral projection of the Schrödinger (wave) propagator  $\tau(E_{(\lambda,+\infty)}(e^{it\mathcal{L}}))$ . The latter remains an open problem.

#### 4. EXAMPLES

In this section we provide some explicit examples to clarify and compare the wide class of groups and operators we can be treated with our results. We do it for the case of heat and wave type equations, but obviously the same examples work for the Schrödinger type equations.

Let us consider the following general  $\mathfrak{D}$ -heat-wave type equation:

$$\begin{aligned} {}^C\partial_t^\alpha w(t, x) - \mathfrak{D}w(t, x) &= 0, \quad t > 0, \quad x \in G, \\ w(t, x)|_{t=0} &= w_0(x), \quad w_0 \in L^p(G), \\ \partial_t w(t, x)|_{t=0} &= w_1(x), \quad w_1 \in L^p(G), \end{aligned} \quad (4.1)$$

where  $0 < \alpha < 2$  ( $\alpha \neq 1$ ),  $1 < p \leq 2$ ,  $\mathfrak{D}$  is any positive linear left invariant operator on a locally compact separable unimodular group  $G$ . We use the first initial condition  $w_0$  when  $0 < \alpha < 1$ . We consider both initial conditions  $w_0, w_1$  when  $1 < \alpha < 2$ .

By Theorem 3.4, we get for any  $1 < p \leq 2 \leq q < +\infty$  the following time decay rate for the solution of equation (4.1):

$$\|w(t, \cdot)\|_{L^q(G)} \leq C_{\alpha,\nu,Q,p,q} t^{-\alpha\lambda(\frac{1}{p}-\frac{1}{q})} (\|w_0\|_{L^p(G)} + t\|w_1\|_{L^p(G)}), \quad (4.2)$$

whenever

$$\tau(E_{(0,s)}(\mathfrak{D})) \lesssim s^\lambda, \quad s \rightarrow +\infty, \quad \text{for some } \lambda > 0, \quad \frac{1}{\lambda} > \frac{1}{p} - \frac{1}{q}.$$

Below we give several examples of equation (4.1) in different *Lie* groups and particular operators acting there, whose trace of the spectral projections behave like  $s^\lambda$  as

$s \rightarrow +\infty$ . Moreover, in some cases, we can claim the sharpness of the time-decay rate.

#	Group	Operator $\mathfrak{D}$	$\tau(\mathbf{E}_{(0,s)}(\mathfrak{D}))$	Time Decay
1	The Euclidean space $\mathbb{R}^n$	$\Delta_{\mathbb{R}^n}$	$s^{n/2}$	$t^{-\alpha n/2(\frac{1}{p}-\frac{1}{q})}$
2	Any compact Lie group $G$	$\Delta_{sub}$	$s^{Q/2}$	$t^{-\alpha Q/2(\frac{1}{p}-\frac{1}{q})}$
3	The Heisenberg group $\mathbb{H}^n$	$\mathcal{L}$	$s^{n+1}$	$t^{-\alpha(n+1)(\frac{1}{p}-\frac{1}{q})}$
4	Any graded Lie group $G$	$\mathcal{R}$	$s^{Q/\nu}$	$t^{-\alpha Q/\nu(\frac{1}{p}-\frac{1}{q})}$
5	Engel group $\mathfrak{B}_4$	$\mathfrak{D}_1$	$s^3$	$t^{-3\alpha(\frac{1}{p}-\frac{1}{q})}$
6	Cartan group $\mathfrak{B}_5$	$\mathfrak{D}_2$	$s^{9/2}$	$t^{-9\alpha/2(\frac{1}{p}-\frac{1}{q})}$
7	Any connected Lie group $G$	$\mathcal{L}$	$s^{Q_*/m}$	$t^{-\alpha Q_*/m(\frac{1}{p}-\frac{1}{q})}$

1. The Laplacian  $\Delta_{\mathbb{R}^n}$  on the Euclidean space  $\mathbb{R}^n$ . By [2, Example 7.3], it follows that

$$\tau(E_{(0,s)}(\Delta_{\mathbb{R}^n})) \lesssim s^{n/2}, \quad s \rightarrow +\infty.$$

Here we can recover the sharp estimate (time-decay) given in [19, Theorem 3.3, item (i)] whenever  $\frac{2}{n} > \frac{1}{p} - \frac{1}{q}$ .

2. The sub-Laplacian  $\Delta_{sub}$  on a compact Lie group. By [16], we have

$$\tau(E_{(0,s)}(-\Delta_{sub})) \lesssim s^{Q/2}, \quad s \rightarrow +\infty,$$

where  $Q$  is the Hausdorff dimension of  $G$ .

3. The positive sub-Laplacian on the Heisenberg group  $\mathbb{H}^n$ . By [2, Formula (7.17)], we get

$$\tau(E_{(0,s)}(\mathcal{L})) \lesssim s^{n+1}, \quad s \rightarrow +\infty.$$

4. A positive Rockland operator  $\mathcal{R}$  on a graded Lie group. By [28, Theorem 8.2], we obtain

$$\tau(E_{(0,s)}(\mathcal{R})) \lesssim s^{Q/\nu}, \quad s \rightarrow +\infty,$$

where  $Q$  is the homogeneous dimension of  $G$ . In this case the estimate is sharp and we can then claim the sharpness of the time-decay rate.

5. The non-Rockland-type operator  $\mathfrak{D}_1 = -(X_1^2 + X_2^2 + X_3^2 + X_4^2 + X_4^{-2})$  on the Engel group  $\mathfrak{B}_4$ , where  $\{X_i\}$  are the vector fields that form the canonical basis of its Lie algebra. By [9, Example 2.2], we know

$$\tau(E_{(0,s)}(\mathfrak{D}_1)) \lesssim s^3, \quad s \rightarrow +\infty.$$

6. The non-Rockland-type operator  $\mathfrak{D}_2 = -(X_1^2 + X_2^2 + X_3^2 + X_4^2 + X_5^2 + X_4^{-2} + X_5^{-2})$  on the Cartan group  $\mathfrak{B}_5$ , where  $\{X_i\}$  are the vector fields that form the canonical basis of its Lie algebra. By [9, Example 3.2], one has

$$\tau(E_{(0,s)}(\mathfrak{D}_2)) \lesssim s^{9/2}, \quad s \rightarrow +\infty.$$

7. An  $m$ -th order weighted subcoercive positive operator on a connected unimodular Lie group. By [29, Proposition 0.3], we arrive at

$$\tau(E_{(0,s)}(\mathcal{L})) \lesssim s^{\frac{Q_*}{m}}, \quad s \rightarrow \infty,$$

where  $Q_*$  is the local dimension of  $G$  relative to the chosen weighted structure on its Lie algebra.

**Example of an abelian locally compact group of non-Lie type.** We conclude this section by presenting in more detail the case of the group of  $\rho$ -adic numbers, denoted by  $\mathbb{Q}_\rho$  ( $\rho$  is a prime number). Since there is no smooth structure in  $\mathbb{Q}_\rho$  we can not define a Laplacian-type operator using derivatives, but we can do it as a Fourier multiplier. In fact, we consider the Vladimirov operator [35], an analogous operator to the fractional Laplacian, defined as follows:

$$\mathfrak{D}^\mu f(x) = \int_{\mathbb{Q}_\rho} e^{2\pi i \{x\xi\}_\rho} |\xi|_\rho^\mu \widehat{f}(\xi) d\xi, \quad \mu > 0, \quad x \in \mathbb{Q}_\rho, \quad (4.3)$$

where  $\widehat{f}$  is the Fourier transform of  $f$ ,  $|\cdot|_\rho$  is the  $\rho$ -adic norm and  $\{\cdot\}_\rho$  is the fractional part of a  $\rho$ -adic number. This means that  $\widehat{\mathfrak{D}^\mu f}(\xi) = |\xi|_\rho^\mu \widehat{f}(\xi)$ , so that its symbol is given by  $\sigma_{\mathfrak{D}^\mu}(\xi) = |\xi|_\rho^\mu$ . From the general theory (see [18, Theorem 5.6.26, p. 360] and [18, Corollary 5.6.29, p. 363]) we get that  $E_{(0,s)}(\mathfrak{D}^\mu) = \chi_{(0,s)}(|\sigma_{\mathfrak{D}^\mu}(\xi)|)$ . Therefore the trace of this spectral projection can be directly computed as

$$\tau(E_{(0,s)}(\mathfrak{D}^\mu)) = \int_{\substack{\mathbb{Q}_\rho \\ |\sigma_{\mathfrak{D}^\mu}(\xi)| \leq s}} d\xi = \int_{\substack{\mathbb{Q}_\rho \\ |\xi|_\rho^\mu \leq s}} d\xi = \text{Vol}(B_{s^{1/\mu}}(0)),$$

where as usual  $B_r(y)$  denotes the  $\rho$ -adic ball of radius  $r$  and center  $y$ . Let  $v$  be the smallest integer such that  $1/\rho^v \sim s^{1/\mu}$ , thus by the topology given in  $\mathbb{Q}_\rho$  we have

$$\text{Vol}(B_{s^{1/\mu}}(0)) = \text{Vol}(B_{\rho^{-v}}(0)) = 1/\rho^v,$$

since this is the normalization of the Haar measure in  $\mathbb{Q}_\rho$  ([5]). Hence

$$\tau(E_{(0,s)}(\mathfrak{D}^\mu)) = 1/\rho^v \lesssim s^{1/\mu}, \quad s \rightarrow \infty.$$

Finally, we obtain time decay rate  $t^{-\alpha/\mu(\frac{1}{p}-\frac{1}{q})}$  for the solution of  $\mathfrak{D}^\mu$ -equation (4.1), with  $\lambda = 1/\mu$  in (4.2).

## 5. ACKNOWLEDGEMENTS

The authors were supported by the FWO Odysseus 1 grant G.0H94.18N: Analysis and Partial Differential Equations, the Methusalem programme of the Ghent University Special Research Fund (BOF) (Grant number 01M01021). MR is also supported by EPSRC grant EP/R003025/2 and FWO Senior Research Grant G011522N.

## 6. CONFLICTS OF INTEREST

On behalf of all authors, the corresponding author states that there is no conflict of interest.



## REFERENCES

- [1] Akylzhanov R., Majid S., Ruzhansky M. Smooth dense subalgebras and Fourier multipliers on compact quantum groups. *Comm. Math. Phys.*, 362, 761–799, (2018).
- [2] Akylzhanov, R., Ruzhansky, M.  $L^p - L^q$  multipliers on locally compact groups. *J. Funct. Anal.*, 278(3), #108324, (2020).
- [3] Arveson, W. A. Short Course on Spectral Theory, vol. 209, Springer Science & Business Media, 2006.
- [4] Bajlekova, E. G. Fractional Evolution Equations in Banach Spaces. Ph.D. Thesis, Eindhoven University of Technology, 2001.
- [5] Bourbaki, N. Elements of Mathematics. Integration II, Springer, Berlin, 2004.
- [6] Brenner, P.  $L^p - L^{p'}$ -Estimates for Fourier integral operators related to hyperbolic equations. *Math. Z.*, 152, 273–286, (1977).
- [7] Brezis, H. Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert. *Math. Studies* 5, North-Holland, Amsterdam, (1973).
- [8] Carvalho-Neto, P. M. Fractional differential equations: a novel study of local and global solutions in Banach spaces. PhD thesis, Universidade de São Paulo, São Carlos, 2013.
- [9] Chatzakou, M. A note on spectral multipliers on Engel and Cartan groups. *Proc. Amer. Math. Soc.*, 150(5), 2259–2270, (2022).
- [10] Chatzakou, M., Restrepo, J. E., Ruzhansky, M. Heat and wave type equations with non-local operators, II. Hilbert spaces and graded Lie groups. *arXiv:2301.12256*, (2023).
- [11] Cowling, M., Giulini, S., Meda, S.  $L^p - L^q$  estimates for functions of the Laplace-Beltrami operator on noncompact symmetric spaces I. *Duke Math. J.*, 72(1), 109–150, (1993).
- [12] de Moraes, W.A.A., Restrepo, J.E., Ruzhansky, M. Heat and wave type equations with non-local operators, I. Compact Lie groups. *Int. Math. Res. Not. IMRN.*, (2023), DOI: 10.1093/imrn/rnad017.
- [13] Dixmier, J. Von Neumann Algebras. North-Holland, Amsterdam, 1981.
- [14] Giga, Y. Solutions for semilinear parabolic equations in  $L^p$  and regularity of weak solutions of the Navier-Stokes system. *J. Differ. Equ.*, 62(2), 186–212, (1986).
- [15] Gorenflo, R., Kilbas, A. A., Mainardi, F., Rogosin, S. V. Mittag-Leffler Functions, Related Topics and Applications, 2nd ed. Springer Monographs in Mathematics, Springer, New York, 2020.
- [16] Hassannezhad, A., Kokarev, G. Sub-Laplacian eigenvalue bounds on sub-Riemannian manifolds. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, 16(4), 1049–1092, (2016).
- [17] Hörmander, L. Estimates for translation invariant operators in  $L^p$  spaces. *Acta Math.*, 104(1–2), 93–140, (1960).
- [18] Kadison, R., Ringrose, J.: Fundamentals of the Theory of Operator Algebras. Volume I: Elementary Theory. Graduate Studies in Mathematics, 1997.
- [19] Kemppainen, J., Siljander, J., Vergara, V., Zacher, R. Decay estimates for time-fractional and other non-local in time subdiffusion equations in  $\mathbb{R}^d$ . *Math. Ann.*, 366(3), 941–979, (2016).
- [20] Kosaki, H. Non-commutative Lorentz spaces associated with a semi-finite von Neumann algebra and applications. *Proc. Japan Acad. Ser. A Math. Sci.*, 57(6), 303–306, (1981).
- [21] Krein, S. G. Linear differential equations in Banach space. American Mathematical Society, 1972.
- [22] Littman, W.  $L^p - L^q$ -estimates for singular integral operators arising from hyperbolic equations. In Partial differential equations (Proc. Sympos. Pure Math., Vol. XXIII, Univ. California, Berkeley, Calif., 1971), 479–481, (1973).
- [23] Murray, F. J., von Neumann, J. On rings of operators. *Ann. of Math.*, (2), 37(1), 116–229, (1936).
- [24] Murray, F. J., von Neumann, J. On rings of operators II. *Trans. Amer. Math. Soc.*, 41(2), 208–248, (1937).
- [25] Pollard, H. The completely monotonic character of the Mittag-Leffler function  $E_a(-x)$ . *Bull. Amer. Math. Soc.* 54, 1115–1116, (1948).
- [26] Podlubny, I. Fractional Differential Equations. Academic Press, San Diego, 1999.

- [27] Prüss, J. Evolutionary integral equations and applications. Birkhäuser, Basel, Boston, Berlin, 1993.
- [28] Rottensteiner, D., Ruzhansky, M. Harmonic and anharmonic oscillators on the Heisenberg group. J. Math. Phys., 63, #111509, (2022).
- [29] Rottensteiner, D., Ruzhansky, M. An update on the  $L^p - L^q$  norms of spectral multipliers on unimodular Lie groups. Arch. Math., 120, 507-520, (2023).
- [30] Samko, S. G., Kilbas, A. A., Marichev, O. I. Fractional integrals and derivatives, translated from the 1987 Russian original, Gordon and Breach, Yverdon, 1993.
- [31] Strichartz, R. S. A priori estimates for the wave equation and some applications. J. Funct. Anal., 5, 218-235, (1970).
- [32] Terp, M.  $L^p$  Spaces Associated with Von Neumann Algebras. Copenhagen University, 1981.
- [33] Thierry, F., Kosaki, H. Generalized  $s$ -numbers of  $\tau$ -measurable operators. Pacific J. Math. 123(2), 269-300, (1986).
- [34] Tricomi, F. G. Integral equations (Pure and Applied Mathematics, Vol. V). Interscience Publishers, Inc., New York; Interscience Publishers Ltd., London 1957 viii+238 pp.
- [35] Vladimirov, V. S., Volovich, I., Zelenov, E.  $p$ -adic analysis and mathematical physics. Series on Soviet and East European mathematics, V1. World Scientific, 1994.

SANTIAGO GÓMEZ COBOS:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, KRIJGSLAAN 281, BUILDING S8, B 9000 GHENT  
BELGIUM

*E-mail address* [davidsantiago.gomezcobos@ugent.be](mailto:davidsantiago.gomezcobos@ugent.be)

JOEL E. RESTREPO:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, KRIJGSLAAN 281, BUILDING S8, B 9000 GHENT  
BELGIUM

*E-mail address* [joel.restrepo@ugent.be](mailto:joel.restrepo@ugent.be); [cocojoel89@yahoo.es](mailto:cocojoel89@yahoo.es)

MICHAEL RUZHANSKY:

DEPARTMENT OF MATHEMATICS: ANALYSIS, LOGIC AND DISCRETE MATHEMATICS  
GHENT UNIVERSITY, KRIJGSLAAN 281, BUILDING S8, B 9000 GHENT  
BELGIUM

AND

SCHOOL OF MATHEMATICAL SCIENCES  
QUEEN MARY UNIVERSITY OF LONDON  
UNITED KINGDOM

*E-mail address* [michael.ruzhansky@ugent.be](mailto:michael.ruzhansky@ugent.be)