

A NOTE ON COMMUTATORS OF SINGULAR INTEGRALS WITH BMO AND VMO FUNCTIONS IN THE DUNKL SETTING

JACEK DZIUBAŃSKI AND AGNIESZKA HEJNA

ABSTRACT. On \mathbb{R}^N equipped with a root system R , multiplicity function $k \geq 0$, and the associated measure $dw(\mathbf{x}) = \prod_{\alpha \in R} |\langle \mathbf{x}, \alpha \rangle|^{k(\alpha)} d\mathbf{x}$, we consider a (non-radial) kernel $K(\mathbf{x})$ which has properties similar to those from the classical theory of singular integrals and the Dunkl convolution operator $\mathbf{T}f = f * K$ associated with K . Assuming that b belongs to the BMO space on the space of homogeneous type $X = (\mathbb{R}^N, \|\cdot\|, dw)$, we prove that the commutator $[b, \mathbf{T}]f(\mathbf{x}) = b(\mathbf{x})\mathbf{T}f(\mathbf{x}) - \mathbf{T}(bf)(\mathbf{x})$ is a bounded operator on $L^p(dw)$ for all $1 < p < \infty$. Moreover, $[b, \mathbf{T}]$ is compact on $L^p(dw)$, provided $b \in \text{VMO}(X)$. The paper extends results of Han, Lee, Li and Wick.

1. INTRODUCTION

1.1. **Introduction.** Consider \mathbb{R}^N equipped with a root system R and a non-negative multiplicity function $k \geq 0$. Let $dw(\mathbf{x}) = \prod_{\alpha \in R} |\langle \alpha, \mathbf{x} \rangle|^{k(\alpha)} d\mathbf{x}$ be the associated measure. For $f \in L^1_{\text{loc}}(dw)$ and a measurable bounded set $E \subset \mathbb{R}^N$, we denote

$$(1.1) \quad f_E := \frac{1}{w(E)} \int_E f(\mathbf{x}) dw(\mathbf{x}).$$

Let G be the Coxeter group generated by the reflections σ_α , $\alpha \in R$. For $E \subset \mathbb{R}^N$, we set

$$\mathcal{O}(E) = \{\sigma(\mathbf{x}) : \sigma \in G, \mathbf{x} \in E\}.$$

In Han, Lee, Li, and Wick [14] the authors investigated two types of BMO and VMO spaces in the Dunkl setting which are connected with two distances: the Euclidean distance $\|\mathbf{x} - \mathbf{y}\|$ and the orbit distance $d(\mathbf{x}, \mathbf{y}) = \min_{\sigma \in G} \|\mathbf{x} - \sigma(\mathbf{y})\|$. The spaces BMO and BMO_d are defined as

$$\text{BMO} = \{b \in L^1_{\text{loc}}(dw) : \|b\|_{\text{BMO}} < \infty\}, \quad \|b\|_{\text{BMO}} := \sup_B \frac{1}{w(B)} \int_B |b(\mathbf{x}) - b_B| dw(\mathbf{x}),$$

$$\text{BMO}_d = \{b \in L^1_{\text{loc}}(dw) : \|b\|_d < \infty\}, \quad \|b\|_d := \sup_B \frac{1}{w(\mathcal{O}(B))} \int_{\mathcal{O}(B)} |b(\mathbf{x}) - b_{\mathcal{O}(B)}| dw(\mathbf{x}),$$

where the supremum is taken over all Euclidean balls $B = B(\mathbf{y}, r) = \{\mathbf{z} \in \mathbb{R}^N : \|\mathbf{y} - \mathbf{z}\| < r\}$. The space BMO_d is a proper subspace of BMO (see [16]) and

$$\|b\|_{\text{BMO}} \leq C \|b\|_{\text{BMO}_d} \quad \text{for } b \in \text{BMO}_d.$$

In [14] commutators of BMO and BMO_d functions with the Dunkl-Riesz transforms R_j are studied. The Dunkl-Riesz transforms are Calderón-Zygmund type operators which are formally defined by $R_j = T_{e_j}(-\Delta_k)^{-1/2}$, where T_{e_j} are the Dunkl operators (see (2.6)) and

2020 *Mathematics Subject Classification.* primary: 44A20, 42B20, 42B25, 47B38, 35K08, 33C52, 39A70.

Key words and phrases. rational Dunkl theory, root systems, generalized translations, singular integrals, commutators, bounded mean oscillation spaces, Riesz transforms.

$\Delta_k = \sum_{j=1}^N T_{e_j}^2$ is the Dunkl Laplacian. They were studied by Thangavelu and Xu [24] (in dimension 1 and in the product case) and by Amri and Sifi [1] (in higher dimensions) who proved their bounds on $L^p(dw)$ spaces. One of the main results of [14] asserts that if $b \in \text{BMO}_d$, then the commutator $[b, R_j]f(\mathbf{x}) = b(\mathbf{x})R_j f(\mathbf{x}) - R_j(b(\cdot)f(\cdot))(\mathbf{x})$ is a bounded operator on $L^p(dw)$ for $1 < p < \infty$ and

$$(1.2) \quad \|[b, R_j]\|_{L^p(dw) \rightarrow L^p(dw)} \lesssim \|b\|_{\text{BMO}_d}.$$

Conversely, if for $b \in L^1_{\text{loc}}(dw)$, the commutator $[b, R_j]$ is bounded on $L^p(dw)$ for some $1 < p < \infty$, then $b \in \text{BMO}$ and

$$(1.3) \quad \|b\|_{\text{BMO}} \lesssim \|[b, R_j]\|_{L^p(dw) \rightarrow L^p(dw)}.$$

The authors of [14] raised the question if the possible lower bound $\|b\|_{\text{BMO}_d} \lesssim \|[b, R_j]\|_{L^p(dw) \rightarrow L^p(dw)}$ holds true.

Our first goal in this note is to improve (1.2) by showing that it holds for $b \in \text{BMO}$, that is, there is a constant $C_p > 0$ such that

$$(1.4) \quad \|[b, R_j]\|_{L^p(dw) \rightarrow L^p(dw)} \leq C_p \|b\|_{\text{BMO}}$$

(see Theorem 3.1).

Let us point out that (1.4) gives a negative answer to the question formulated above, because otherwise we would get $\|b\|_{\text{BMO}_d} \lesssim \|[b, R_j]\|_{L^p(dw) \rightarrow L^p(dw)} \lesssim \|b\|_{\text{BMO}}$, which is impossible (see [16]).

An essential part of [14] is devoted for studying compactness of the commutators of VMO functions with the Dunkl-Riesz transforms. The VMO and VMO_d spaces are defined as the closures of the sets of compactly supported functions from the Lipschitz spaces Λ and Λ_d in the norms $\|\cdot\|_{\text{BMO}}$ and $\|\cdot\|_{\text{BMO}_d}$ respectively. Then $\text{VMO}_d \subset \text{VMO}$ and $\|b\|_{\text{VMO}} \lesssim \|b\|_{\text{VMO}_d}$ and, thanks to [5, Theorem 4.1], the dual space to VMO is the Hardy space H^1 considered in [2] and [10]. Theorem 1.5 of [14] states that if $b \in \text{VMO}_d$, then the commutator $[b, R_j]$ is a compact operator on $L^p(dw)$ for all $1 < p < \infty$. Our second aim is to extend this result for all $b \in \text{VMO}$ (see Theorem 3.2). Actually we will prove (1.2) and the compactness result for commutators $[b, \mathbf{T}]$ of BMO and VMO functions with Dunkl singular integral operators \mathbf{T} of convolution type (under certain regularity for the associated kernels $K(\mathbf{x})$). The Dunkl-Riesz transforms are the basic examples of such operators.

Let us remark that for the Riesz transforms, the lower bounds (1.3) proved in [14] together with the upper bounds (1.4) (see Theorem 3.1) generalize (to the Dunkl setting) the classical results of Coifman, Rochberg and Weiss [4] and Janson [15] obtained on the Euclidean spaces $(\mathbb{R}^N, d\mathbf{x})$. As far as the compactness is concerned, the necessity result [14, Theorem 1.5] together with the sufficiency result (see Theorem 3.2) extend to the Dunkl theory the classical theorems of Uchiyama [26] about the characterization of the VMO functions by commutators with the Riesz transforms.

2. PRELIMINARIES

2.1. Dunkl theory. In this section we present basic facts concerning the theory of the Dunkl operators. For more details we refer the reader to [7], [18], [19], and [20].

We consider the Euclidean space \mathbb{R}^N with the scalar product $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{j=1}^N x_j y_j$, where $\mathbf{x} = (x_1, \dots, x_N)$, $\mathbf{y} = (y_1, \dots, y_N)$, and the norm $\|\mathbf{x}\|^2 = \langle \mathbf{x}, \mathbf{x} \rangle$.

A *normalized root system* in \mathbb{R}^N is a finite set $R \subset \mathbb{R}^N \setminus \{0\}$ such that $R \cap \alpha\mathbb{R} = \{\pm\alpha\}$, $\sigma_\alpha(R) = R$, and $\|\alpha\| = \sqrt{2}$ for all $\alpha \in R$, where σ_α are defined by

$$(2.1) \quad \sigma_\alpha(\mathbf{x}) = \mathbf{x} - 2\frac{\langle \mathbf{x}, \alpha \rangle}{\|\alpha\|^2}\alpha.$$

The finite group G generated by the reflections σ_α , $\alpha \in R$, is called the *Coxeter group* (*reflection group*) of the root system.

A *multiplicity function* is a G -invariant function $k : R \rightarrow \mathbb{C}$ which will be fixed and ≥ 0 throughout this paper.

The associated measure dw is defined by $dw(\mathbf{x}) = w(\mathbf{x}) d\mathbf{x}$, where

$$(2.2) \quad w(\mathbf{x}) = \prod_{\alpha \in R} |\langle \mathbf{x}, \alpha \rangle|^{k(\alpha)}.$$

Let $\mathbf{N} = N + \sum_{\alpha \in R} k(\alpha)$. Then,

$$(2.3) \quad w(B(t\mathbf{x}, tr)) = t^{\mathbf{N}} w(B(\mathbf{x}, r)) \quad \text{for all } \mathbf{x} \in \mathbb{R}^N, t, r > 0.$$

Observe that there is a constant $C > 0$ such that for all $\mathbf{x} \in \mathbb{R}^N$ and $r > 0$ we have

$$(2.4) \quad C^{-1}w(B(\mathbf{x}, r)) \leq r^{\mathbf{N}} \prod_{\alpha \in R} (|\langle \mathbf{x}, \alpha \rangle| + r)^{k(\alpha)} \leq Cw(B(\mathbf{x}, r)),$$

so $dw(\mathbf{x})$ is doubling. Moreover, there exists a constant $C \geq 1$ such that, for every $\mathbf{x} \in \mathbb{R}^N$ and for all $r_2 \geq r_1 > 0$,

$$(2.5) \quad C^{-1}\left(\frac{r_2}{r_1}\right)^{\mathbf{N}} \leq \frac{w(B(\mathbf{x}, r_2))}{w(B(\mathbf{x}, r_1))} \leq C\left(\frac{r_2}{r_1}\right)^{\mathbf{N}}.$$

For $\xi \in \mathbb{R}^N$, the *Dunkl operators* T_ξ are the following k -deformations of the directional derivatives ∂_ξ by difference operators:

$$(2.6) \quad T_\xi f(\mathbf{x}) = \partial_\xi f(\mathbf{x}) + \sum_{\alpha \in R} \frac{k(\alpha)}{2} \langle \alpha, \xi \rangle \frac{f(\mathbf{x}) - f(\sigma_\alpha(\mathbf{x}))}{\langle \alpha, \mathbf{x} \rangle}.$$

The Dunkl operators T_ξ , which were introduced in [7], commute and are skew-symmetric with respect to the G -invariant measure dw .

The closures of connected components of

$$\{\mathbf{x} \in \mathbb{R}^N : \langle \mathbf{x}, \alpha \rangle \neq 0 \text{ for all } \alpha \in R\}$$

are called (closed) *Weyl chambers*. We remark that $\|\mathbf{x} - \mathbf{y}\| = d(\mathbf{x}, \mathbf{y})$ if and only if $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$ belong to the same closed Weyl chamber.

Let

$$(2.7) \quad Mf(\mathbf{x}) = \sup_{B \ni \mathbf{x}} \frac{1}{w(B)} \int_B |f(\mathbf{y})| dw(\mathbf{y}),$$

denote the (uncentered) Hardy–Littlewood maximal function on $(\mathbb{R}^N, \|\cdot\|, dw)$.

2.2. Dunkl transform. For fixed $\mathbf{y} \in \mathbb{R}^N$, the *Dunkl kernel* $\mathbf{x} \mapsto E(\mathbf{x}, \mathbf{y})$ is a unique solution to the system

$$T_\xi f = \langle \xi, \mathbf{y} \rangle f, \quad f(0) = 1.$$

The function $E(\mathbf{x}, \mathbf{y})$, which generalizes the exponential function $e^{\langle \mathbf{x}, \mathbf{y} \rangle}$, has a unique extension to a holomorphic function $E(\mathbf{z}, \mathbf{w})$ on $\mathbb{C}^N \times \mathbb{C}^N$. Let $f \in L^1(dw)$. We define the *Dunkl transform* $\mathcal{F}f$ of f by

$$(2.8) \quad \mathcal{F}f(\xi) = \mathbf{c}_k^{-1} \int_{\mathbb{R}^N} f(\mathbf{x}) E(\mathbf{x}, -i\xi) dw(\mathbf{x}), \quad \mathbf{c}_k = \int_{\mathbb{R}^N} e^{-\frac{\|\mathbf{x}\|^2}{2}} dw(\mathbf{x}) > 0$$

The Dunkl transform is a generalization of the Fourier transform on \mathbb{R}^N . It was introduced in [8] for $k \geq 0$ and further studied in [6] in the more general context. It was proved in [8, Corollary 2.7] (see also [6, Theorem 4.26]) that it extends uniquely to an isometry on $L^2(dw)$. We have also the following inversion theorem ([6, Theorem 4.20]): for all $f \in L^1(dw)$ such that $\mathcal{F}f \in L^1(dw)$ we have $f(\mathbf{x}) = (\mathcal{F})^2 f(-\mathbf{x})$ for almost all $\mathbf{x} \in \mathbb{R}^N$. The inverse \mathcal{F}^{-1} of \mathcal{F} has the form

$$(2.9) \quad \mathcal{F}^{-1}f(\mathbf{x}) = \mathbf{c}_k^{-1} \int_{\mathbb{R}^N} f(\xi) E(i\xi, \mathbf{x}) dw(\xi) = \mathcal{F}f(-\mathbf{x}) \quad \text{for } f \in L^1(dw).$$

2.3. Dunkl translations. Suppose that $f \in \mathcal{S}(\mathbb{R}^N)$ (the Schwartz class of functions on \mathbb{R}^N) and $\mathbf{x} \in \mathbb{R}^N$. We define the *Dunkl translation* $\tau_{\mathbf{x}}f$ of f to be

$$(2.10) \quad \tau_{\mathbf{x}}f(-\mathbf{y}) = \mathbf{c}_k^{-1} \int_{\mathbb{R}^N} E(i\xi, \mathbf{x}) E(-i\xi, \mathbf{y}) \mathcal{F}f(\xi) dw(\xi) = \mathcal{F}^{-1}(E(i\cdot, \mathbf{x})\mathcal{F}f)(-\mathbf{y}).$$

The Dunkl translation was introduced in [17]. The definition can be extended to functions which are not necessary in $\mathcal{S}(\mathbb{R}^N)$. For instance, thanks to the Plancherel's theorem, one can define the Dunkl translation of $L^2(dw)$ function f by

$$(2.11) \quad \tau_{\mathbf{x}}f(-\mathbf{y}) = \mathcal{F}^{-1}(E(i\cdot, \mathbf{x})\mathcal{F}f(\cdot))(-\mathbf{y})$$

(see [17] of [23, Definition 3.1]). In particular, the operators $f \mapsto \tau_{\mathbf{x}}f$ are contractions on $L^2(dw)$. Here and subsequently, for a reasonable function $g(\mathbf{x})$, we write $g(\mathbf{x}, \mathbf{y}) := \tau_{\mathbf{x}}g(-\mathbf{y})$.

We will need the following result concerning the support of the Dunkl translation of a compactly supported function.

Theorem 2.1 ([9, Theorem 1.7]). *Let $f \in L^2(dw)$, $\text{supp } f \subseteq B(0, r)$, and $\mathbf{x} \in \mathbb{R}^N$. Then*

$$(2.12) \quad \text{supp } \tau_{\mathbf{x}}f(-\cdot) \subseteq \mathcal{O}(B(\mathbf{x}, r)).$$

2.4. Dunkl convolution. Assume that $f, g \in L^2(dw)$. The *generalized convolution* (or *Dunkl convolution*) $f * g$ is defined by the formula

$$(2.13) \quad f * g(\mathbf{x}) = \mathbf{c}_k \mathcal{F}^{-1}((\mathcal{F}f)(\mathcal{F}g))(\mathbf{x}),$$

equivalently, by

$$(2.14) \quad (f * g)(\mathbf{x}) = \int_{\mathbb{R}^N} f(\mathbf{y}) \tau_{\mathbf{x}}g(-\mathbf{y}) dw(\mathbf{y}) = \int_{\mathbb{R}^N} g(\mathbf{y}) \tau_{\mathbf{x}}f(-\mathbf{y}) dw(\mathbf{y}).$$

Generalized convolution of $f, g \in \mathcal{S}(\mathbb{R}^N)$ was considered in [17] and [25], the definition was extended to $f, g \in L^2(dw)$ in [23].

2.5. Singular integral kernels. Let us consider a (non-radial) kernel $K(\mathbf{x})$ which has properties similar to those from the classical theory. Namely, let s_0 be an even positive integer larger than \mathbf{N} , which will be fixed in the whole paper. Consider a function $K \in C^{s_0}(\mathbb{R}^N \setminus \{0\})$ such that

$$(A) \quad \sup_{0 < a < b < \infty} \left| \int_{a < \|\mathbf{x}\| < b} K(\mathbf{x}) \, dw(\mathbf{x}) \right| < \infty,$$

$$(D) \quad \left| \frac{\partial^\beta}{\partial \mathbf{x}^\beta} K(\mathbf{x}) \right| \leq C_\beta \|\mathbf{x}\|^{-\mathbf{N}-|\beta|} \quad \text{for all } |\beta| \leq s_0,$$

$$(L) \quad \lim_{\varepsilon \rightarrow 0} \int_{\varepsilon < \|\mathbf{x}\| < 1} K(\mathbf{x}) \, dw(\mathbf{x}) = L \text{ for some } L \in \mathbb{C}.$$

Set

$$K^{\{t\}}(\mathbf{x}) = K(\mathbf{x})(1 - \phi(t^{-1}\mathbf{x})),$$

where ϕ is a fixed radial C^∞ -function supported by the unit ball $B(0, 1)$ such that $\phi(\mathbf{x}) = 1$ for $\|\mathbf{x}\| < 1/2$. The following theorem was proved in [11].

Theorem 2.2. *Assume that (A) and (D) are satisfied, then*

- (i) *(Theorems 4.1 and 4.2 of [11]) the operators $f \mapsto f * K^{\{t\}}$ are bounded on $L^p(dw)$ for $1 < p < \infty$ and they are of weak-type $(1, 1)$ with the bounds independent of $t > 0$;*
- (ii) *(Theorems 3.7 and 4.3 of [11]) assuming additionally (L), the limit $\lim_{t \rightarrow 0} f * K^{\{t\}}(\mathbf{x})$ exists and defines a bounded operator \mathbf{T} on $L^p(dw)$ for $1 < p < \infty$, which is of weak-type $(1, 1)$ as well.*

3. STATEMENT OF THE RESULTS

3.1. Commutators. In order to define the commutator operator, we come back to the definition of the limit operator \mathbf{T} . Let $0 < \varepsilon < \min(1, s_0 - \mathbf{N})$. For any $t > 0$ let us denote $K^{\{t/2, t\}} := K^{\{t/2\}} - K^{\{t\}}$. Then $K^{\{t/2, t\}}$ is $C^{s_0}(\mathbb{R}^N)$ -function supported by $B(0, t) \setminus B(0, t/4)$ (cf. [11, (3.1)]). In order to simplify the notation, we write $K_\ell = K^{\{2^{\ell-1}, 2^\ell\}}$. It is proved in [12, (4.24), (4.25)] that for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$ and $\ell \in \mathbb{Z}$, we have

$$(3.1) \quad |K_\ell(\mathbf{x}, \mathbf{y})| \leq C \left(1 + \frac{\|\mathbf{x} - \mathbf{y}\|}{2^\ell} \right)^{-\varepsilon} w(B(\mathbf{x}, 2^\ell))^{-1/2} w(B(\mathbf{y}, 2^\ell))^{-1/2},$$

$$(3.2) \quad \begin{aligned} & |K_\ell(\mathbf{x}, \mathbf{y}) - K_\ell(\mathbf{x}, \mathbf{y}')| \\ & \leq C \frac{\|\mathbf{y} - \mathbf{y}'\|^\varepsilon}{2^{\varepsilon\ell}} \left(1 + \frac{\|\mathbf{x} - \mathbf{y}\|}{2^\ell} \right)^{-\varepsilon} w(B(\mathbf{x}, 2^\ell))^{-1/2} (w(B(\mathbf{y}, 2^\ell))^{-1/2} + w(B(\mathbf{y}', 2^\ell))^{-1/2}). \end{aligned}$$

Moreover, $K_\ell(\mathbf{x}, \mathbf{y}) = K_\ell(-\mathbf{y}, -\mathbf{x})$ and by [12, proof of Theorem 4.6],

$$(3.3) \quad \sum_{\ell=-\infty}^{\infty} |K_\ell(\mathbf{x}, \mathbf{y})| \leq C w(B(\mathbf{x}, d(\mathbf{x}, \mathbf{y})))^{-1} \frac{d(\mathbf{x}, \mathbf{y})^\varepsilon}{\|\mathbf{x} - \mathbf{y}\|^\varepsilon}$$

for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$ such that $d(\mathbf{x}, \mathbf{y}) \neq 0$ and

$$(3.4) \quad \sum_{\ell=-\infty}^{\infty} |K_{\ell}(\mathbf{x}, \mathbf{y}) - K_{\ell}(\mathbf{x}, \mathbf{y}')| \leq C \frac{\|\mathbf{y} - \mathbf{y}'\|^{\varepsilon}}{\|\mathbf{x} - \mathbf{y}\|^{\varepsilon}} w(B(\mathbf{x}, d(\mathbf{x}, \mathbf{y})))^{-1}$$

for all $\mathbf{x}, \mathbf{y}, \mathbf{y}' \in \mathbb{R}^N$ such that $\|\mathbf{y} - \mathbf{y}'\| < \frac{d(\mathbf{x}, \mathbf{y})}{2}$. The number $\varepsilon > 0$ will be fixed in the whole paper. Thanks to (3.3) the function

$$(3.5) \quad K(\mathbf{x}, \mathbf{y}) = \sum_{\ell=-\infty}^{\infty} K_{\ell}(\mathbf{x}, \mathbf{y})$$

is well defined for $d(\mathbf{x}, \mathbf{y}) > 0$ and, by Theorem 2.2, it is the associated kernel to the operator \mathbf{T} , that is,

$$(3.6) \quad \mathbf{T}f(\mathbf{x}) = \int_{\mathbb{R}^N} K(\mathbf{x}, \mathbf{y})f(\mathbf{y}) dw(\mathbf{y})$$

for $f \in L^p(dw)$ and $\mathbf{x} \notin \text{supp } f$.

Let us emphasize that the estimate for $K(\mathbf{x}, \mathbf{y})$ which are consequences of (3.3) and (3.4) turn out to be very useful in handling some harmonic analysis problems in the Dunkl setting (see [22]).

From now on we fix a kernel $K \in C^{s_0}(\mathbb{R}^N \setminus \{0\})$ satisfying (A), (L), and (D) for some $s_0 > \mathbf{N}$. Let $b \in \text{BMO}$. For any compactly supported $f \in L^p(dw)$ for some $p > 1$, we define

$$(3.7) \quad \mathcal{C}f(\mathbf{x}) = [b, \mathbf{T}]f(\mathbf{x}) = \lim_{m \rightarrow \infty} \int_{\mathbb{R}^N} (b(\mathbf{x}) - b(\mathbf{y})) \sum_{\ell=-m}^m K_{\ell}(\mathbf{x}, \mathbf{y})f(\mathbf{y}) dw(\mathbf{y}) = \lim_{m \rightarrow \infty} \mathcal{C}_m f(\mathbf{x})$$

The existence of the limit in (3.7) in any $L^{p_0}(dw)$ -norm, provided $1 < p_0 < p$, is proved in Lemma 4.2. Then, Theorem 3.1 and its proof allows one to extend the definition for all $f \in L^p(dw)$.

3.2. Statements of main theorems. Our main results are the following two theorems.

Theorem 3.1. *Let $p > 1$. Assume that a kernel $K \in C^{s_0}(\mathbb{R}^N \setminus \{0\})$ satisfies (A), (L), (D) for a certain even integer $s_0 > \mathbf{N}$, and $b \in \text{BMO}$. Then there is a constant $C > 0$ independent of b such that for all $f \in L^p(dw)$ we have*

$$(3.8) \quad \|\mathcal{C}f\|_{L^p(dw)} \leq C \|b\|_{\text{BMO}} \|f\|_{L^p(dw)}.$$

In order to formulate our second theorem, recall that VMO is the closure in BMO of compactly supported Lipschitz functions, i.e., functions f satisfying $\sup_{\mathbf{x} \neq \mathbf{y}} \frac{|f(\mathbf{x}) - f(\mathbf{y})|}{\|\mathbf{x} - \mathbf{y}\|} < \infty$.

Theorem 3.2. *Assume $b \in \text{VMO}$. For $1 < p < \infty$, the commutator \mathcal{C} is a compact operator on $L^p(dw)$.*

To prove the theorems, we adapt the ideas of classical proofs (cf. e.g. [13]) to apply estimates for integral kernels of operators which are expressed in terms of the orbit distance $d(\mathbf{x}, \mathbf{y})$ and the Euclidean metric, see (3.3) and (3.4) (cf. also [14]). These require in some places much careful analysis. For example, we use the Coxeter group for decomposing $L^p(dw)$ -functions (see (4.9)) or we split integration over \mathbb{R}^N onto the Weyl chambers (see (4.10)).

4. PROOF OF THEOREM 3.1

Let us begin with the following lemma. Recall that $b_{B(\mathbf{x},r)}$ is defined by (1.1).

Lemma 4.1. *There is a constant $C > 0$ such that for all $b \in L^1_{\text{loc}}(dw)$, $r_1 > r > 0$, $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$, and $\sigma \in G$, we have:*

$$(4.1) \quad |b_{B(\mathbf{x},r)} - b_{B(\mathbf{x},r_1)}| \leq C \log(r_1/r) \|b\|_{\text{BMO}},$$

$$(4.2) \quad |b_{B(\mathbf{x},r)} - b_{B(\mathbf{y},r)}| \leq C \|b\|_{\text{BMO}} \quad \text{provided } \|\mathbf{x} - \mathbf{y}\| \leq 2r,$$

$$(4.3) \quad |b_{B(\mathbf{x},r)} - b_{B(\sigma(\mathbf{x}),r)}| \leq C \log \left(\frac{\|\sigma(\mathbf{x}) - \mathbf{x}\|}{r} + 4 \right) \|b\|_{\text{BMO}},$$

(John–Nirenberg inequality) for any $1 \leq s < \infty$ there is $C_s > 0$ such that for all positive integers j we have

$$(4.4) \quad \left(\frac{1}{w(B(\mathbf{x}, 2^j r))} \int_{B(\mathbf{x}, 2^j r)} |b(\mathbf{y}) - b_{B(\mathbf{x},r)}|^s dw(\mathbf{y}) \right)^{1/s} \leq C_s j \|b\|_{\text{BMO}}.$$

Proof. Inequalities (4.1), (4.2), and (4.4) are consequences of the well-known results in metric spaces with doubling measures (see e.g. [3], [5], or [16, Section 4] in the Dunkl setting). In order to prove (4.3), note that if $\|\mathbf{x} - \sigma(\mathbf{x})\| \leq 2r$, then (4.3) follows by (4.2). Assume that $\|\mathbf{x} - \sigma(\mathbf{x})\| > 2r$ and let j be the smallest positive integer such that $\|\mathbf{x} - \sigma(\mathbf{x})\| \leq 2^j r$. Then $B(\mathbf{x}, 2^{j+2}r) \cap B(\sigma(\mathbf{x}), 2^{j+2}r) \neq \emptyset$. So, applying (4.1) and (4.2), we get

$$\begin{aligned} & |b_{B(\mathbf{x},r)} - b_{B(\sigma(\mathbf{x}),r)}| \\ & \leq |b_{B(\mathbf{x},r)} - b_{B(\mathbf{x},2^{j+2}r)}| + |b_{B(\mathbf{x},2^{j+2}r)} - b_{B(\sigma(\mathbf{x}),2^{j+2}r)}| + |b_{B(\sigma(\mathbf{x}),2^{j+2}r)} - b_{B(\sigma(\mathbf{x}),r)}| \\ & \leq C j \|b\|_{\text{BMO}} \leq C' \log \left(\frac{\|\mathbf{x} - \sigma(\mathbf{x})\|}{r} + 4 \right) \|b\|_{\text{BMO}}. \end{aligned}$$

□

Lemma 4.2. *Let $p > p_0 > 1$. If $f \in L^p(dw)$ is compactly supported, then $\mathcal{C}_m f \in L^{p_0}(dw)$ for all $m \in \mathbb{Z}$ and*

$$\lim_{m \rightarrow \infty} \mathcal{C}_m f = \mathcal{C} f \text{ in } L^{p_0}(dw).$$

Proof. Let $B = B(0, r)$ be any ball such that $\text{supp } f \subseteq B$. We write

$$\mathcal{C}_m f(\mathbf{x}) = (b(\mathbf{x}) - b_B) \sum_{\ell=-m}^m K_\ell * f(\mathbf{x}) + \sum_{\ell=-m}^m K_\ell * ((b - b_B)f)(\mathbf{x}).$$

By Hölder's inequality with $s = p/p_0$ and the John–Nirenberg inequality (4.4) we obtain

$$\left(\int_B |b(\mathbf{x}) - b_B|^{p_0} |f(\mathbf{x})|^{p_0} dw(\mathbf{x}) \right)^{1/p_0} \leq C w(B)^{1/(p_0 s')} \|b\|_{\text{BMO}} \|f\|_{L^p(dw)},$$

where $s' > 1$ is such that $s + s' = ss'$, so $(b - b_B)f \in L^{p_0}(dw)$. Hence, by Theorem 2.2 (ii),

$$\lim_{m \rightarrow \infty} \sum_{\ell=-m}^m K_\ell * ((b - b_B)f)$$

exists in the $L^{p_0}(dw)$ -norm and it is equal to $\mathbf{T}((b - b_B)f) \in L^{p_0}(dw)$. Again, by Theorem 2.2 (ii), the limit

$$\lim_{m \rightarrow \infty} \sum_{\ell=-m}^m K_\ell * f$$

exists in the $L^{p_0}(dw)$ and $L^p(dw)$ -norms and it is equal to $\mathbf{T}f$. We write

$$(b(\mathbf{x}) - b_B)\mathbf{T}f(\mathbf{x}) = (b(\mathbf{x}) - b_B)\mathbf{T}f(\mathbf{x})\chi_{5B}(\mathbf{x}) + (b(\mathbf{x}) - b_B)\mathbf{T}f(\mathbf{x})\chi_{(5B)^c}(\mathbf{x}) =: g_1(\mathbf{x}) + g_2(\mathbf{x}).$$

We will show that $g_1, g_2 \in L^{p_0}(dw)$. For g_1 , by Hölder's inequality with $s = p/p_0$, the John-Nirenberg inequality (4.4), and $L^p(dw)$ -boundedness of \mathbf{T} (Theorem 2.2 (ii)), we get

$$\begin{aligned} & \left(\int_{5B} |b(\mathbf{x}) - b_B|^{p_0} |\mathbf{T}f(\mathbf{x})|^{p_0} dw(\mathbf{x}) \right)^{1/p_0} \\ & \leq C \left(\int_{5B} |b(\mathbf{x}) - b_B|^{p_0 s'} dw(\mathbf{x}) \right)^{1/(p_0 s')} \left(\int_{5B} |\mathbf{T}f(\mathbf{x})|^p dw(\mathbf{x}) \right)^{1/p} \\ & \leq C w(B)^{1/(p_0 s')} \|b\|_{\text{BMO}} \|f\|_{L^p(dw)}. \end{aligned}$$

In order to prove that $g_2 \in L^{p_0}(dw)$, we observe that, by (2.3) and (2.4), $w(B(\mathbf{x}, d(\mathbf{x}, \mathbf{y}))) \sim w(B(0, \|\mathbf{x}\|)) = c\|\mathbf{x}\|^{\mathbf{N}}$ for $\mathbf{y} \in B$ and $\mathbf{x} \in (5B)^c$. Using (3.3) we have

$$\begin{aligned} (4.5) \quad \chi_{(5B)^c}(\mathbf{x}) |\mathbf{T}f(\mathbf{x})| & \leq C \chi_{(5B)^c}(\mathbf{x}) \int_B \frac{|f(\mathbf{y})|}{w(B(\mathbf{x}, d(\mathbf{x}, \mathbf{y})))} dw(\mathbf{y}) \\ & \leq C \int_B \frac{|f(\mathbf{y})|}{w(B(0, \|\mathbf{x}\|))} dw(\mathbf{y}) \leq C \|f\|_{L^p(dw)} \frac{w(B)^{1/p'}}{w(B(0, \|\mathbf{x}\|))}. \end{aligned}$$

Consequently,

$$\begin{aligned} (4.6) \quad \|g_2\|_{L^{p_0}(dw)}^{p_0} & \leq C \|f\|_{L^p(dw)}^{p_0} w(B)^{p_0/p'} \int_{\|\mathbf{x}\| > 5r} \frac{|b(\mathbf{x}) - b_B|^{p_0}}{w(B(0, \|\mathbf{x}\|))^{p_0}} dw(\mathbf{x}) \\ & \leq C \|f\|_{L^p(dw)}^{p_0} w(B)^{p_0/p'} \sum_{j=1}^{\infty} \int_{\|\mathbf{x}\| \leq 2^j r} \frac{|b(\mathbf{x}) - b_B|^{p_0}}{w(B(0, 2^j r))^{p_0}} dw(\mathbf{x}). \end{aligned}$$

Now, applying the John-Nirenberg inequality (4.4) we get

$$\begin{aligned} (4.7) \quad \|g_2\|_{L^{p_0}(dw)}^{p_0} & \leq C \|f\|_{L^p(dw)}^{p_0} w(B)^{p_0/p'} \sum_{j=1}^{\infty} \|b\|_{\text{BMO}}^{p_0} \frac{j^{p_0}}{(2^j r)^{(p_0-1)\mathbf{N}}} \\ & \leq C' \|f\|_{L^p(dw)}^{p_0} w(B)^{p_0/p'} \|b\|_{\text{BMO}}^{p_0} r^{-(p_0-1)\mathbf{N}}. \end{aligned}$$

□

Proof of Theorem 3.1. We shall prove the inequality (3.8) for compactly supported functions $f \in L^p(dw)$ which form a dense subspace in $L^p(dw)$. To this end, thanks to Lemma 4.2 and [21, page 148, Theorem 2 and the remark below it], it suffices to verify that

$$(4.8) \quad \|(\mathcal{C}f)^\# \|_{L^p(dw)} \leq C \|b\|_{\text{BMO}} \|f\|_{L^p(dw)},$$

where $g^\#$ denotes the sharp maximal function

$$g^\#(\mathbf{x}) = \sup_{B \ni \mathbf{x}} \inf_{c \in \mathbb{C}} \frac{1}{w(B)} \int_B |g(\mathbf{y}) - c| dw(\mathbf{y}).$$

Let $\mathbf{x} \in \mathbb{R}^N$ and let $B = B(\mathbf{x}_0, r)$ be any ball which contains \mathbf{x} . We write all the elements of $G \setminus \{\text{id}\}$ in a sequence $\sigma_1, \sigma_2, \dots, \sigma_{|G|-1}$. We define the sets $U_j \subseteq \mathbb{R}^N$, $j = 1, 2, \dots, |G| - 1$, inductively:

$$U_1 := \{\mathbf{z} \in \mathbb{R}^N : \|\mathbf{z} - \mathbf{x}_0\| > 5r, \|\mathbf{z} - \sigma_1(\mathbf{x}_0)\| \leq 5r\},$$

$$U_{j+1} := \{\mathbf{z} \in \mathbb{R}^N : \|\mathbf{z} - \mathbf{x}_0\| > 5r, \|\mathbf{z} - \sigma_{j+1}(\mathbf{x}_0)\| \leq 5r\} \setminus \left(\bigcup_{j_1=1}^j U_{j_1} \right) \text{ for } j \geq 1.$$

For a compactly supported function $f \in L^p(dw)$, we decompose

$$(4.9) \quad f = f_1 + f_2 + \sum_{j=1}^{|G|-1} f_{\sigma_j}, \text{ where } f_1 := f \cdot \chi_{5B}, \quad f_2 := f \cdot \chi_{(\mathcal{O}(5B))^c}, \quad f_{\sigma_j} := f \cdot \chi_{U_j}.$$

For $\mathbf{y} \in B$ we set

$$g_1(\mathbf{y}) := \mathcal{C}f_1(\mathbf{y}) = (b(\mathbf{y}) - b_B)\mathbf{T}f_1(\mathbf{y}) + \mathbf{T}((b_B - b)f_1)(\mathbf{y}) =: g_{11}(\mathbf{y}) + g_{12}(\mathbf{y}),$$

$$g_2(\mathbf{y}) := \mathcal{C}f_2(\mathbf{y}) = (b(\mathbf{y}) - b_B)\mathbf{T}f_2(\mathbf{y}) + \mathbf{T}((b_B - b)f_2)(\mathbf{y}) =: g_{21}(\mathbf{y}) + g_{22}(\mathbf{y}),$$

$$g_{\sigma_j}(\mathbf{y}) := \mathcal{C}f_{\sigma_j}(\mathbf{y}) = (b(\mathbf{y}) - b_B)\mathbf{T}f_{\sigma_j}(\mathbf{y}) + \mathbf{T}((b_B - b)f_{\sigma_j})(\mathbf{y}) =: g_{\sigma_j 1}(\mathbf{y}) + g_{\sigma_j 2}(\mathbf{y}).$$

Fix $1 < s < p$. Further, by the fact that $|(g_{11})_B| \leq \frac{1}{w(B)} \int_B |g_{11}(\mathbf{y})| dw(\mathbf{y})$, by the definition of g_{11} , and Hölder's inequality,

$$\begin{aligned} \frac{1}{w(B)} \int_B |g_{11}(\mathbf{y}) - (g_{11})_B| dw(\mathbf{y}) &\leq \frac{2}{w(B)} \int_B |g_{11}(\mathbf{y})| dw(\mathbf{y}) \\ &\leq 2 \frac{1}{w(B)} \int_B |b(\mathbf{y}) - b_B| \cdot |\mathbf{T}f_1(\mathbf{y})| dw(\mathbf{y}) \\ &\leq 2 \left(\frac{1}{w(B)} \int_B |b(\mathbf{y}) - b_B|^{s'} dw(\mathbf{y}) \right)^{1/s'} \cdot \left(\frac{1}{w(B)} \int_B |\mathbf{T}f_1(\mathbf{y})|^s dw(\mathbf{y}) \right)^{1/s}, \end{aligned}$$

where $s' > 1$ is such that $\frac{1}{s} + \frac{1}{s'} = 1$. Applying inequality (4.4) we conclude that

$$g_{11}^\#(\mathbf{x}) \leq C \|b\|_{\text{BMO}} (M(|\mathbf{T}f_1|^s)(\mathbf{x}))^{1/s}.$$

The same analysis gives

$$g_{21}^\#(\mathbf{x}) \leq C \|b\|_{\text{BMO}} (M(|\mathbf{T}f_2|^s)(\mathbf{x}))^{1/s}, \quad g_{\sigma_j 1}^\#(\mathbf{x}) \leq C \|b\|_{\text{BMO}} (M(|\mathbf{T}f_{\sigma_j}|^s)(\mathbf{x}))^{1/s}.$$

To deal with g_{12} , we choose $q, v \in (1, \infty)$ such that $1 < qv < p < \infty$ and $s = qv$. Then, by Hölder's inequality and $L^q(dw)$ -boundedness of \mathbf{T} ,

$$\begin{aligned} \frac{1}{w(B)} \int_B |g_{12}(\mathbf{y}) - (g_{12})_B| dw(\mathbf{y}) &\leq \frac{2}{w(B)} \int_B |g_{12}(\mathbf{y})| dw(\mathbf{y}) \\ &\leq 2 \frac{1}{w(B)} \int_B |\mathbf{T}((b(\cdot) - b_B) \cdot f_1)(\mathbf{y})| dw(\mathbf{y}) \\ &\leq 2 \left(\frac{1}{w(B)} \int_B \mathbf{T}((b(\cdot) - b_B) \cdot f_1)(\mathbf{y})^q dw(\mathbf{y}) \right)^{1/q} \\ &\leq C \left(\frac{1}{w(5B)} \int_{5B} |(b(\mathbf{y}) - b_B) \cdot f_1(\mathbf{y})|^q dw(\mathbf{y}) \right)^{1/q} \\ &\leq C \left(\frac{1}{w(5B)} \int_{5B} |b(\mathbf{y}) - b_B|^{qv'} dw(\mathbf{y}) \right)^{1/(qv')} \cdot \left(\frac{1}{w(5B)} \int_{5B} |f_1(\mathbf{y})|^{qv} dw(\mathbf{y}) \right)^{1/(qv)}. \end{aligned}$$

Hence, applying the John-Nirenberg inequality (4.4), we get

$$g_{12}^\#(\mathbf{x}) \leq C \|b\|_{\text{BMO}} (M(|f_1|^s)(\mathbf{x}))^{1/s}.$$

We turn to analyse g_{22} . Observe that for $\mathbf{z} \notin \mathcal{O}(5B)$ and $\mathbf{y} \in B$ we have $\|\mathbf{x}_0 - \mathbf{y}\| \leq d(\mathbf{x}_0, \mathbf{z})$. Let Γ be a fixed closed Weyl chamber such that $\mathbf{x}_0 \in \Gamma$, then by the estimates (3.4),

(4.10)

$$\begin{aligned} |g_{22}(\mathbf{y}) - g_{22}(\mathbf{x}_0)| &\leq \int_{\mathbb{R}^N} \sum_{\ell=-\infty}^{\infty} |K_\ell(\mathbf{z}, \mathbf{y}) - K_\ell(\mathbf{z}, \mathbf{x}_0)| |b_B - b(\mathbf{z})| |f_2(\mathbf{z})| dw(\mathbf{z}) \\ &= \sum_{\sigma \in G} \int_{\mathbf{z} \in \sigma(\Gamma)} \sum_{\ell=-\infty}^{\infty} |K_\ell(\mathbf{z}, \mathbf{y}) - K_\ell(\mathbf{z}, \mathbf{x}_0)| |b_B - b(\mathbf{z})| |f_2(\mathbf{z})| dw(\mathbf{z}) \\ &\leq C \sum_{\sigma \in G} \int_{\mathbf{z} \in \sigma(\Gamma)} \frac{\|\mathbf{y} - \mathbf{x}_0\|^\varepsilon}{\|\mathbf{x}_0 - \mathbf{z}\|^\varepsilon} \frac{1}{w(B(\mathbf{x}_0, d(\mathbf{x}_0, \mathbf{z})))} |b_B - b(\mathbf{z})| |f_2(\mathbf{z})| dw(\mathbf{z}) \\ &=: \sum_{\sigma \in G} J_\sigma(\mathbf{y}). \end{aligned}$$

In dealing with $J_\sigma(\mathbf{y})$ we shall use the inequalities:

$$\begin{aligned} \|\mathbf{x}_0 - \mathbf{z}\| &\geq \max(\|\mathbf{x}_0 - \sigma(\mathbf{x}_0)\|/2, r) \quad \text{for } \mathbf{z} \in \sigma(\Gamma), \\ \|\sigma(\mathbf{x}_0) - \sigma(\mathbf{x})\| &\leq r < 5r \leq \|\sigma(\mathbf{x}_0) - \mathbf{z}\| = d(\mathbf{x}_0, \mathbf{z}) \leq \|\mathbf{x}_0 - \mathbf{z}\| \quad \text{for } \mathbf{z} \in \sigma(\Gamma), \mathbf{z} \notin \mathcal{O}(5B). \end{aligned}$$

So,

$$\begin{aligned} J_\sigma(\mathbf{y}) &\leq C \int_{\mathbf{z} \in \sigma(\Gamma)} \frac{r^\varepsilon}{\|\mathbf{x}_0 - \mathbf{z}\|^\varepsilon} \frac{1}{w(B(\sigma(\mathbf{x}_0), \|\sigma(\mathbf{x}_0) - \mathbf{z}\|))} |b_B - b_{\sigma(B)}| |f_2(\mathbf{z})| dw(\mathbf{z}) \\ &\quad + C \int_{\mathbf{z} \in \sigma(\Gamma)} \frac{r^\varepsilon}{\|\mathbf{x}_0 - \mathbf{z}\|^\varepsilon} \frac{1}{w(B(\sigma(\mathbf{x}_0), \|\sigma(\mathbf{x}_0) - \mathbf{z}\|))} |b_{\sigma(B)} - b(\mathbf{z})| |f_2(\mathbf{z})| dw(\mathbf{z}) =: J_{\sigma,1}(\mathbf{y}) + J_{\sigma,2}(\mathbf{y}). \end{aligned}$$

Further, by (4.3),

(4.11)

$$\begin{aligned} J_{\sigma,1}(\mathbf{y}) &\leq C \int_{\mathbf{z} \in \sigma(\Gamma)} \frac{r^{\varepsilon/2}}{\|\mathbf{x}_0 - \mathbf{z}\|^{\varepsilon/2}} \frac{r^{\varepsilon/2}}{r^{\varepsilon/2} + \|\mathbf{x}_0 - \sigma(\mathbf{x}_0)\|^{\varepsilon/2}} \log\left(\frac{\|\mathbf{x}_0 - \sigma(\mathbf{x}_0)\|}{r} + 4\right) \|b\|_{\text{BMO}} \\ &\quad \times \frac{|f_2(\mathbf{z})|}{w(B(\sigma(\mathbf{x}_0), \|\sigma(\mathbf{x}_0) - \mathbf{z}\|))} dw(\mathbf{z}) \\ &\leq C \|b\|_{\text{BMO}} \int_{\mathbf{z} \in \sigma(\Gamma)} \frac{r^{\varepsilon/2}}{\|\mathbf{x}_0 - \mathbf{z}\|^{\varepsilon/2}} \cdot \frac{|f_2(\mathbf{z})|}{w(B(\sigma(\mathbf{x}_0), \|\sigma(\mathbf{x}_0) - \mathbf{z}\|))} dw(\mathbf{z}) \\ &\leq C \|b\|_{\text{BMO}} \sum_{j=2}^{\infty} \int_{\mathbf{z} \in \sigma(\Gamma), \|\sigma(\mathbf{x}_0) - \mathbf{z}\| \sim 2^j r} \frac{r^{\varepsilon/2}}{\|\mathbf{x}_0 - \mathbf{z}\|^{\varepsilon/2}} \cdot \frac{|f_2(\mathbf{z})|}{w(B(\sigma(\mathbf{x}_0), \|\sigma(\mathbf{x}_0) - \mathbf{z}\|))} dw(\mathbf{z}) \\ &\leq C \|b\|_{\text{BMO}} M f_2(\sigma(\mathbf{x})). \end{aligned}$$

We turn to considering $J_{\sigma,2}(\mathbf{y})$. Applying Hölder's inequality and then the John-Nirenberg inequality (4.4) we obtain

$$\begin{aligned}
(4.12) \quad J_{\sigma,2}(\mathbf{y}) &\leq C \sum_{j=2}^{\infty} \int_{\mathbf{z} \in \sigma(\Gamma), \|\mathbf{z} - \sigma(\mathbf{x}_0)\| \sim 2^j r} \frac{r^\varepsilon}{2^{\varepsilon j} r^\varepsilon} |b_{\sigma(B)} - b(\mathbf{z})| \frac{|f_2(\mathbf{z})|}{w(B(\sigma(\mathbf{x}_0), 2^j r))} dw(\mathbf{z}) \\
&\leq C \sum_{j=2}^{\infty} 2^{-\varepsilon j} \left(\int_{\|\mathbf{z} - \sigma(\mathbf{x}_0)\| \sim 2^j r} |b_{\sigma(B)} - b(\mathbf{z})|^{s'} \frac{dw(\mathbf{z})}{w(B(\sigma(\mathbf{x}_0), 2^j r))} \right)^{1/s'} \\
&\quad \times \left(\int_{\|\mathbf{z} - \sigma(\mathbf{x}_0)\| \leq 2^j r} |f_2(\mathbf{z})|^s \frac{dw(\mathbf{z})}{w(B(\sigma(\mathbf{x}_0), 2^j r))} \right)^{1/s} \\
&\leq C \sum_{j=2}^{\infty} 2^{-\varepsilon j} \|b\|_{\text{BMO}} (M(|f_2|^s)(\sigma(\mathbf{x})))^{1/s} \leq C \|b\|_{\text{BMO}} (M(|f_2|^s)(\sigma(\mathbf{x})))^{1/s}.
\end{aligned}$$

Thus, by (4.11) and (4.12) we have got

$$g_{22}^\#(\mathbf{x}) \leq C \sum_{\sigma \in G} \left(M f_2(\sigma(\mathbf{x})) + (M(|f_2|^s)(\sigma(\mathbf{x})))^{1/s} \right).$$

Finally we turn to estimate $g_{\sigma_j,2}$. To this end we note that for $\mathbf{z} \in U_j$ and $\mathbf{y} \in B$ we have

$$\|\mathbf{z} - \mathbf{y}\| \geq \|\mathbf{z} - \mathbf{x}_0\| - \|\mathbf{x}_0 - \mathbf{y}\| \geq 5r - r = 4r,$$

$$\|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\| \leq \|\mathbf{x}_0 - \mathbf{y}\| + \|\mathbf{z} - \mathbf{y}\| + \|\mathbf{z} - \sigma_j(\mathbf{x}_0)\| \leq 6r + \|\mathbf{z} - \mathbf{y}\| \leq \frac{5}{2} \|\mathbf{z} - \mathbf{y}\|,$$

Consequently, by (3.3),

$$\begin{aligned}
(4.13) \quad &\int_B \sum_{\ell=-\infty}^{\infty} |K_\ell(\mathbf{z}, \mathbf{y})| dw(\mathbf{y}) \leq C \int_B \frac{d(\mathbf{z}, \mathbf{y})^\varepsilon}{\|\mathbf{z} - \mathbf{y}\|^\varepsilon} \frac{1}{w(B(\mathbf{z}, d(\mathbf{z}, \mathbf{y})))} dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\|)^\varepsilon} \int_B \frac{d(\mathbf{z}, \mathbf{y})^\varepsilon}{r^\varepsilon} \frac{1}{w(B(\mathbf{z}, d(\mathbf{z}, \mathbf{y})))} dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\|)^\varepsilon} \int_{\mathcal{O}(B(\mathbf{z}, 16r))} \frac{d(\mathbf{z}, \mathbf{y})^\varepsilon}{r^\varepsilon} \frac{1}{w(B(\mathbf{z}, d(\mathbf{z}, \mathbf{y})))} dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\|)^\varepsilon} \sum_{j=-4}^{\infty} \int_{d(\mathbf{z}, \mathbf{y}) \sim 2^j r} \frac{d(\mathbf{z}, \mathbf{y})^\varepsilon}{r^\varepsilon} \frac{1}{w(B(\mathbf{z}, d(\mathbf{z}, \mathbf{y})))} dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\|)^\varepsilon} \sum_{j=-4}^{\infty} 2^{-\varepsilon j} \int_{d(\mathbf{z}, \mathbf{y}) \sim 2^{-j} r} \frac{1}{w(B(\mathbf{z}, 2^{-j} r))} dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\mathbf{x}_0 - \sigma_j(\mathbf{x}_0)\|)^\varepsilon}.
\end{aligned}$$

Hence, by (4.13) and (4.3),

$$\begin{aligned}
\frac{1}{w(B)} \int_B |g_{\sigma_j 2}(\mathbf{y}) - (g_{\sigma_j 2})_B| dw(\mathbf{y}) &\leq \frac{2}{w(B)} \int_B |g_{\sigma_j 2}(\mathbf{y})| dw(\mathbf{y}) \\
&\leq \frac{2}{w(B)} \int_B \int_{U_j} \sum_{\ell=-\infty}^{\infty} |K_\ell(\mathbf{y}, \mathbf{z})| |b_B - b(\mathbf{z})| \cdot |f_{\sigma_j}(\mathbf{z})| dw(\mathbf{z}) dw(\mathbf{y}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\sigma_j(\mathbf{x}_0) - \mathbf{x}_0\|)^\varepsilon} \frac{1}{w(B)} \int_{U_j} |b_B - b(\mathbf{z})| \cdot |f_{\sigma_j}(\mathbf{z})| dw(\mathbf{z}) \\
&\leq C \frac{r^\varepsilon}{(r + \|\sigma_j(\mathbf{x}_0) - \mathbf{x}_0\|)^\varepsilon} \left(\frac{1}{w(B)} \int_{U_j} |b_B - b(\mathbf{z})|^{s'} dw(\mathbf{z}) \right)^{1/s'} \left(\frac{1}{w(B)} \int_{U_j} |f_{\sigma_j}(\mathbf{z})|^s dw(\mathbf{z}) \right)^{1/s} \\
&\leq C \frac{r^\varepsilon}{(r + \|\sigma_j(\mathbf{x}_0) - \mathbf{x}_0\|)^\varepsilon} \log \left(\frac{\|\sigma_j(\mathbf{x}_0) - \mathbf{x}_0\|}{r} + 4 \right) \|b\|_{\text{BMO}} (M|f_{\sigma_j}|^s(\sigma(\mathbf{x})))^{1/s},
\end{aligned}$$

so

$$|g_{\sigma_j 2}^\#(\mathbf{x})| \leq C \|b\|_{\text{BMO}} (M|f_{\sigma_j}|^s(\sigma(\mathbf{x})))^{1/s}.$$

Finally we end up with the estimate

$$\begin{aligned}
(\mathcal{C}f)^\#(\mathbf{x}) &\leq C \|b\|_{\text{BMO}} \left((M(|\mathbf{T}f_1|^s)(\mathbf{x}))^{1/s} + M(|\mathbf{T}f_2|^s(\mathbf{x}))^{1/s} \right. \\
(4.14) \quad &\left. + \sum_{j=1}^{|G|-1} M(|\mathbf{T}f_{\sigma_j}|^s(\mathbf{x}))^{1/s} + \sum_{\sigma \in G} (Mf(\sigma(\mathbf{x})) + (M|f|^s(\sigma(\mathbf{x})))^{1/s}) \right).
\end{aligned}$$

Hence, using the $L^{p_1}(dw)$ -boundedness of the Hardy–Littlewood maximal M function for all $1 < p_1 < \infty$ and the fact that the measure dw is G -invariant, we conclude (4.8) from (4.14) and (4.9). \square

5. PROOF OF THEOREM 3.2

Lemma 5.1. *Let $1 < p < \infty$. Assume that b is a compactly supported Lipschitz function. Then*

$$(5.1) \quad \lim_{m \rightarrow \infty} \|\mathcal{C} - \mathcal{C}_m\|_{L^p(dw) \rightarrow L^p(dw)} = 0.$$

Proof. Let $r_b > 1$ be such that $\text{supp } b \subset B(0, r_b)$ and let $L_b > 0$ be such that $|b(\mathbf{x}) - b(\mathbf{y})| \leq L_b \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^N$. By (3.7), it is enough to prove that there is a constant $C > 0$ such that for all positive integers m such that $2^m \geq 2r_b$, $f \in L^p(dw)$, and $\mathbf{x} \in \mathbb{R}^N$, we have

$$(5.2) \quad \sum_{\ell=-\infty}^{-m-1} \left| \int_{\mathbb{R}^N} (b(\mathbf{x}) - b(\mathbf{y})) K_\ell(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \leq CL_b 2^{-\varepsilon m} \sum_{\sigma \in G} Mf(\sigma(\mathbf{x})),$$

$$(5.3) \quad \sum_{\ell=m+1}^{\infty} \left| b(\mathbf{x}) \int_{\mathbb{R}^N} K_\ell(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \leq C \|b\|_{L^\infty} 2^{-mN/p} \chi_{B(0, r_b)}(\mathbf{x}) \|f\|_{L^p(dw)},$$

$$(5.4) \quad \sum_{\ell=m+1}^{\infty} \left| \int_{\mathbb{R}^N} K_\ell(\mathbf{x}, \mathbf{y}) b(\mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \leq C \|b\|_{L^\infty} (\chi_{B(0, 2^m)}(\mathbf{x}) 2^{-mN} + \chi_{B(0, 2^m)^c}(\mathbf{x}) \|\mathbf{x}\|^{-N}) \|f\|_{L^p(dw)}.$$

Let us note that by Theorem 2.1, $\text{supp } K_\ell(\mathbf{x}, \cdot) \subseteq \mathcal{O}(B(\mathbf{x}, 2^\ell))$. Hence, by (3.1) and the doubling property of the measure dw (see (2.5)) we have

$$(5.5) \quad |K_\ell(\mathbf{x}, \mathbf{y})| \leq C \left(1 + \frac{\|\mathbf{x} - \mathbf{y}\|}{2^\ell}\right)^{-\varepsilon} w(B(\mathbf{x}, 2^\ell))^{-1}.$$

In order to prove (5.2), we use now the Lipschitz condition for b and get

$$\begin{aligned} & \sum_{\ell=-\infty}^{-m-1} \left| \int_{\mathbb{R}^N} (b(\mathbf{x}) - b(\mathbf{y})) K_\ell(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \\ & \leq C \sum_{\ell=-\infty}^{-m-1} \frac{L_b}{w(B(\mathbf{x}, 2^\ell))} \int_{\mathcal{O}(B(\mathbf{x}, 2^\ell))} \|\mathbf{x} - \mathbf{y}\|^\varepsilon \frac{2^{\varepsilon\ell}}{\|\mathbf{x} - \mathbf{y}\|^\varepsilon} |f(\mathbf{y})| dw(\mathbf{y}) \leq CL_b 2^{-\varepsilon m} \sum_{\sigma \in G} Mf(\sigma(\mathbf{x})). \end{aligned}$$

In order to prove (5.3), let us note that by (2.4), $w(B(\mathbf{x}, 2^\ell)) \geq c2^{\ell N}$ for all $\ell \geq 0$. Hence, from (5.5), Hölder's inequality, and the fact that $\text{supp } b \subseteq B(0, r_b)$ we conclude that

$$(5.6) \quad \begin{aligned} & \sum_{\ell=m+1}^{\infty} \left| b(\mathbf{x}) \int_{\mathbb{R}^N} K_\ell(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \\ & \leq C \chi_{B(0, r_b)}(\mathbf{x}) \|b\|_{L^\infty} \sum_{\ell=m+1}^{\infty} \left(\int_{\mathcal{O}(B(\mathbf{x}, 2^\ell))} \frac{1}{w(B(\mathbf{x}, 2^\ell))^{p'}} dw(\mathbf{y}) \right)^{1/p'} \|f\|_{L^p(dw)} \\ & \leq C \chi_{B(0, r_b)}(\mathbf{x}) \|b\|_{L^\infty} 2^{-mN/p} \|f\|_{L^p(dw)}. \end{aligned}$$

Finally, for (5.4), we consider two cases. If $\|\mathbf{x}\| \leq 2^m$, then similarly to (5.6) we get

$$(5.7) \quad \begin{aligned} & \sum_{\ell=m+1}^{\infty} \left| \int_{\mathbb{R}^N} K_\ell(\mathbf{x}, \mathbf{y}) b(\mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \\ & \leq C \chi_{B(0, 2^m)}(\mathbf{x}) \|b\|_{L^\infty} \sum_{\ell=m+1}^{\infty} \left(\int_{B(0, r_b)} \frac{1}{w(B(\mathbf{y}, 2^\ell))^{p'}} dw(\mathbf{y}) \right)^{1/p'} \|f\|_{L^p(dw)} \\ & \leq C \chi_{B(0, 2^m)}(\mathbf{x}) \|b\|_{L^\infty} 2^{-mN} \|f\|_{L^p(dw)}. \end{aligned}$$

Now we assume that $\|\mathbf{x}\| > 2^m$. Recall that $2^m \geq 2r_b$. From Theorem 2.1 and the fact that $\text{supp } K_\ell \subseteq B(0, 2^\ell)$, we conclude that

$$K_\ell(\mathbf{x}, \mathbf{y}) = 0 \quad \text{for all } \mathbf{y} \in B(0, r_b) \text{ and } \ell \geq m \text{ such that } 2^{\ell+1} < \|\mathbf{x}\|.$$

Further, by (5.5) and the doubling property (2.5) of dw ,

$$|K_\ell(\mathbf{x}, \mathbf{y})| \leq \frac{C}{w(B(0, 2^\ell))} \quad \text{for all } \mathbf{y} \in B(0, r_b) \text{ and } \ell \geq m \text{ such that } 2^{\ell+1} \geq \|\mathbf{x}\|.$$

Hence, by the fact that b is supported by $B(0, r_b)$, we obtain

$$\begin{aligned} & \sum_{\ell=m+1}^{\infty} \left| \int_{\mathbb{R}^N} K_\ell(\mathbf{x}, \mathbf{y}) b(\mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) \right| \leq C \sum_{\ell=\lfloor \log(\|\mathbf{x}\|) \rfloor - 2}^{\infty} \int_{B(0, r_b)} \frac{\|b\|_{L^\infty}}{w(B(0, 2^\ell))} |f(\mathbf{y})| dw(\mathbf{y}) \\ & \leq C \frac{\|b\|_{L^\infty}}{w(B(0, \|\mathbf{x}\|))} \int_{B(0, r_b)} |f(\mathbf{y})| dw(\mathbf{y}) \leq Cr_b^{N/p'} \|b\|_{L^\infty} \|\mathbf{x}\|^{-N} \|f\|_{L^p(dw)}. \end{aligned}$$

□

Proof of Theorem 3.2. Let us recall that the compactly supported Lipschitz functions form a dense subspace in VMO. Hence, by Theorem 3.1, it suffices to show that \mathcal{C} is a compact operator on $L^p(dw)$ for any compactly supported Lipschitz function b . Further, thanks to Lemma 5.1, it is enough to prove that for any compactly supported Lipschitz function b , the commutators \mathcal{C}_m are compact operators on $L^p(dw)$ for large enough positive integers m . To this end, let $r_b > 1$ be such that $\text{supp } b \subset B(0, r_b)$. First we note that if f is supported by $B(0, r_b + 2^m)^c$, then $b \cdot f \equiv 0$. Moreover, since $\text{supp } \sum_{\ell=-m}^m K_\ell(\mathbf{x}, \cdot) \subseteq \mathcal{O}(B(\mathbf{x}, 2^m))$ (cf. Theorem 2.1), we have $\sum_{\ell=-m}^m K_\ell(\mathbf{x}, \cdot) \cdot f(\cdot) \equiv 0$ for all $\mathbf{x} \in B(0, r_b)$. Consequently, for all $f \in L^p(dw)$ we have

$$\mathcal{C}_m f(\mathbf{x}) = \int_{\mathbb{R}^N} (b(\mathbf{x}) - b(\mathbf{y})) \sum_{\ell=-m}^m K_\ell(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) dw(\mathbf{y}) = \mathcal{C}_m(f \cdot \chi_{B(0, r_b + 2^m)})(\mathbf{x}).$$

Moreover, since $\text{supp } \sum_{\ell=-m}^m K_\ell(\mathbf{x}, \cdot) \subseteq \mathcal{O}(B(\mathbf{x}, 2^m))$,

$$\text{supp } \mathcal{C}_m(f) = \text{supp } \mathcal{C}_m(f \cdot \chi_{B(0, r_b + 2^m)}) \subseteq B(0, r_b + 2^{m+1}).$$

Let $\Omega = B(0, r_b + 2^{m+1})$. The proof now is reduced to showing that \mathcal{C}_m is a compact operator $L^p(\Omega, dw) \mapsto L^p(\Omega, dw)$. The estimates (3.1) and (3.2) imply

$$|\mathcal{C}_m f(\mathbf{x})| \leq C_{\Omega, m} \|f\|_{L^p(\Omega, dw)},$$

$$|\mathcal{C}_m f(\mathbf{x}) - \mathcal{C}_m f(\mathbf{x}')| \leq C_{\Omega, m} \|\mathbf{x} - \mathbf{x}'\|^\varepsilon \cdot \|f\|_{L^p(\Omega, dw)}$$

for all $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^N$. By the Arzelá-Ascoli theorem the set

$$\{\mathcal{C}_m f : f \in L^p(\Omega, dw), \|f\|_{L^p(\Omega, dw)} \leq 1\}$$

is relatively compact in $C(\Omega)$ with the sup-norm. Hence it is relatively compact in $L^p(\Omega, dw)$, since $w(\Omega)$ is finite. \square

REFERENCES

- [1] B. Amri, M. Sifi, *Riesz transforms for Dunkl transform*, Ann. Math. Blaise Pascal 19 (2012), no. 1, 247–262.
- [2] J.-Ph. Anker, J. Dziubański, A. Hejna, *Harmonic functions, conjugate harmonic functions and the Hardy space H^1 in the rational Dunkl setting*, J. Fourier Anal. Appl. 25 (2019), 2356–2418.
- [3] S. M. Buckley, *Inequalities of John-Nirenberg type in doubling spaces*, J. Anal. Math. 79 (1999), 215–240.
- [4] R.R. Coifman, R. Rochberg G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. 103 (1976), no. 4, 511–635.
- [5] R.R. Coifman, G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. 83 (1977), no. 4, 569–615.
- [6] M.F.E. de Jeu, *The Dunkl transform*, Invent. Math. 113 (1993), 147–162.
- [7] C.F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. 311 (1989), no. 1, 167–183.
- [8] C.F. Dunkl, *Hankel transforms associated to finite reflection groups*, in: *Proc. of the special session on hypergeometric functions on domains of positivity, Jack polynomials and applications*, Proceedings, Tampa 1991, Contemp. Math. 138 (1989), 123–138.
- [9] J. Dziubański and A. Hejna, *Hörmander’s multiplier theorem for the Dunkl transform*, Journal of Functional Analysis 277 (2019), 2133–2159.
- [10] J. Dziubański, A. Hejna, *Remark on atomic decompositions for the Hardy space H^1 in the rational Dunkl setting*, Studia Math. 251 (2020), no. 1, 89–110.
- [11] J. Dziubański, A. Hejna, *Singular integrals in the rational Dunkl setting*, Revista Matemática Complutense, (2021).
- [12] J. Dziubański, A. Hejna, *Remarks on Dunkl translations of non-radial kernels*, arXiv:2211.02518.

- [13] L. Grafakos, *Modern Fourier Analysis*, 3rd edition, Graduate Texts in Mathematics, 250. Springer, New York, 2014.
- [14] Y. Han, M-Y. Lee, J. Li, B. D. Wick, *Riesz transform and commutators in the Dunkl setting*, preprint [arXiv:2105.11275](https://arxiv.org/abs/2105.11275).
- [15] S. Janson, *Mean oscillation and commutators of singular integral operators*, ARK. Math. 16 (1978), 263–270. Volume 412, (2023).
- [16] J. Jiu and Z. Li, *The dual of the Hardy space associated with the Dunkl operators*, Advances in Mathematics, Volume 412, (2023).
- [17] M. Rösler, *Generalized Hermite polynomials and the heat equation for Dunkl operators*, Comm. Math. Phys. 192 (1998), 519–542.
- [18] M. Rösler, *Positivity of Dunkl’s intertwining operator*, Duke Math. J. 98 (1999), no. 3, 445–463.
- [19] M. Rösler: *Dunkl operators (theory and applications)*. In: Koelink, E., Van Assche, W. (eds.) *Orthogonal polynomials and special functions* (Leuven, 2002), 93–135. Lect. Notes Math. 1817, Springer-Verlag (2003).
- [20] M. Rösler, M. Voit, *Dunkl theory, convolution algebras, and related Markov processes*, in *Harmonic and stochastic analysis of Dunkl processes*, P. Graczyk, M. Rösler, M. Yor (eds.), 1–112, Travaux en cours 71, Hermann, Paris, 2008.
- [21] E.M. Stein, *Harmonic analysis (real variable methods, orthogonality and oscillatory integrals)*, Princeton Math. Series 43, Princeton Univ. Press, 1993.
- [22] Ch. Tan, Y. Han, Y. Han, M-Y. Lee, J. Li, *Singular integral operators, T1 theorem, Littlewood-Paley theory and Hardy spaces in Dunkl Setting*, [arXiv:2204.01886](https://arxiv.org/abs/2204.01886).
- [23] S. Thangavelu, Y. Xu, *Convolution operator and maximal function for the Dunkl transform*, J. Anal. Math. 97 (2005), 25–55.
- [24] S. Thangavelu, Y. Xu, *Riesz transform and Riesz potentials for Dunkl transform*, J. Comput. Appl. Math. 199 (2007), no. 1, 181–195, J. Anal. Math. 97 (2005), 25–55.
- [25] K. Trimèche, *Paley-Wiener theorems for the Dunkl transform and Dunkl translation operators*, Integral Transforms Spec. Funct. 13 (2002), no. 1, 17–38.
- [26] A. Uchiyama, *On compactness operators of Hankel type*, Tohoku Math. J. 30 (1978), 163–171.

JACEK DZIUBAŃSKI, UNIWERSYTET WROCŁAWSKI, INSTYTUT MATEMATYCZNY, PL. GRUNWALDZKI 2, 50-384 WROCŁAW, POLAND

Email address: jdziuban@math.uni.wroc.pl

AGNIESZKA HEJNA, UNIWERSYTET WROCŁAWSKI, INSTYTUT MATEMATYCZNY, PL. GRUNWALDZKI 2, 50-384 WROCŁAW, POLAND & DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854-8019, USA

Email address: hejna@math.uni.wroc.pl