

# Universality of Koba-Nielsen-Olesen scaling in QCD at high energy and entanglement

Yizhuang Liu

*Institute of Theoretical Physics, Jagiellonian University, 30-348 Kraków, Poland.\**

Maciej A. Nowak<sup>†</sup>

*Institute of Theoretical Physics and Mark Kac Center for Complex Systems Research,  
Jagiellonian University, 30-348 Kraków, Poland*

Ismail Zahed<sup>‡</sup>

*Center for Nuclear Theory, Department of Physics and Astronomy,  
Stony Brook University, Stony Brook, New York 11794-3800, USA*

(Dated: February 6, 2023)

Using Mueller's dipole formalism for deep inelastic scattering in QCD, we formulate and solve the evolution for the generating function for the multiplicities of the produced particles, in hadronic processes at high energy. The solution for the multiplicities satisfies Koba-Nielsen-Olesen (KNO) scaling, with good agreement with the recently re-analyzed data from the H1 experiment at HERA (DESY), and the old ALEPH data for hadronic  $Z$  decay at LEP (CERN). The same scaling function with KNO scaling, carries to the hadronic multiplicities from jets in electron-positron annihilation. This agreement is *a priori* puzzling, since in Mueller's dipole evolution, one accounts for virtual dipoles in a wave function, whereas in electron-positron annihilation, one describes cross-sections of real particles. We explain the origin of this similarity, pointing at a particular duality between the two processes. Finally, we interpret our results from the point of view of quantum entanglement between slow and fast degrees of freedom in QCD, and derive the entanglement entropy pertinent to electron-positron annihilation into hadronic jets.

1. Universality is a powerful concept permeating several branches of physics, whereby different physical systems can exhibit similar behavior. This is usually captured by universal exponents, given general assumptions. Perhaps, the best example is the universality of the critical exponents in scaling laws in the vicinity of phase changes. Scaling laws, *per se*, form an important theoretical corpus in physics. In general, they describe the functional relationship between two physical quantities, that scale with each other over a significant interval.

In the context of high energy particle physics, the so-called Koba-Nielsen-Olesen scaling (named KNO scaling hereafter), formulated half a century ago, is of paramount importance

in the empirical analysis of many high energy hadronic multiplicities. Yet, it is usually challenging to derive from first principles in QCD. Historically, KNO scaling was first formulated in two, independent theoretical works [1] and [2], which suggested that at high energies with large Mandelstam  $s$  (squared invariant mass), the probability distribution of producing  $n$  particles in a specific collision process, scales as

$$\bar{n}(s)p_n(s) = f(z) \quad (1)$$

where  $\bar{n}(s)$  is the average multiplicity at large  $s$ , and  $z \equiv \frac{n}{\bar{n}(s)}$  is the argument of the *scaling function*  $f(z)$ . Remarkably, the KNO scaling hypothesis precedes the emergence of Quantum Chromodynamics (QCD), and the advent of high energy and luminosity data currently available at colliders.

This letter is motivated by the recent work in [3], where the DIS data from the H1 experiment at DESY were re-analyzed, with inter-

---

\* yizhuang.liu@uj.edu.pl

† maciej.a.nowak@uj.edu.pl

‡ ismail.zahed@stonybrook.edu

est in an assessment of the quantum entanglement in high energy particle physics. Clearly, the data analysed, especially for the highest energy range, shows KNO scaling (see Fig. 1). Also, the Shannon entropy of the multiplicities presented, bears some similarity to the entanglement entropy. However, the explicit form of the scaling function was unknown, and the QCD understanding of the hypothetical entanglement was not specified.

In the first part of this letter, we discuss the unexpected *a priori* fact, that identical differential equations, such as the ones we derived recently in [4], yield a scaling function that applies to different settings at high energy, e.g. deep inelastic scattering (DIS) and jets in electron-positron or  $e^+e^-$  annihilation. Exploiting the formalism based on the Banfi-Marchesini-Smye (BMS) construction [5], we point to the fact that the pertinent generating functions for the multiplicity probabilities in the case of DIS and jets, respectively, are mathematically *identical*, leading to similar equations. Furthermore, in the double logarithm approximation (DLA), we arrive at the final differential equation for the KNO function, which we then solve using methods based on analyticity. The resulting, parameter-free curve, does not only agree well with DIS data from the H1 experiment at DESY, and the Z-decay data from ALEPH at CERN, but also represents an exact prediction for future DIS experiments, alike EIC or EicC.

In sum, this letter consists of three new results: 1/ The derivation of the KNO scaling function in QCD for both DIS and jets in the DLA; 2/ The use of the KNO scaling function in the DLA, to show the universality of the hadronic multiplicities from current colliders, for both DIS and jets; 3/ The explicit derivation of the entanglement entropy for  $e^+e^-$  annihilation into hadronic jets, to be measured at collider energies.

**2.** The BMS equation [5–7], describes the “non-global” logarithms in the  $e^+e^-$  annihilation process, and is based on the

universal features of the soft divergences in the  $n$ -gluon contribution to the total cross section  $\sigma_n$ . To leading logarithm accuracy, the “most singular” part of  $\sigma_n$ , can be effectively generated through a Markov process. Defining the directions of the quark-antiquark pair as  $p$  and  $n$ , soft gluons ( $k$ ) are emitted from harder ones ( $p$ ) through a universal eikonal current  $\frac{gp^\mu}{p \cdot k}$ , and the emissions are strongly ordered in time and energy. As a result, the emission depends only on the color-charges that are already present in the final state, but not on the history of how they are emitted. In the large number of colors  $N_c$ , the generating functional for  $\sigma_n$ ,

$$Z\left(\frac{E}{E_0}; n, p; \lambda\right) = \sum_{n=0}^{\infty} \lambda^n \sigma_n \quad (2)$$

satisfies a closed integral equation [4]

$$\begin{aligned} Z\left(\frac{E}{E_0}; n, p; \lambda\right) &= e^{-\bar{\alpha}_s \ln \frac{E}{E_0}} \int d\Omega_k K(\hat{k}; \hat{p}, \hat{n}) \\ &+ \bar{\alpha}_s \lambda \int_{E_0}^E \frac{d\omega}{\omega} e^{-\bar{\alpha}_s \ln \frac{E}{\omega}} \int d\Omega_k K(\hat{k}; \hat{p}, \hat{n}) \int d\Omega_k \\ &\times K(\hat{k}; \hat{p}, \hat{n}) Z\left(\frac{\omega}{E_0}; n, \hat{k}; \lambda\right) Z\left(\frac{\omega}{E_0}; p, \hat{k}; \lambda\right), \quad (3) \end{aligned}$$

with the eikonalized gluonic emission kernel  $K(\hat{k}; \hat{p}, \hat{n}) = \frac{1}{4\pi} \frac{p \cdot n}{\hat{k} \cdot p \hat{k} \cdot n}$  and  $\bar{\alpha}_s = N_c \alpha_s / \pi$ . The first term is the Sudakov contribution where all the soft gluons are virtual, and the second term is the contribution where at least one soft gluon is real. (3) is the integral form of the BMS equation, which can be brought to the standard form discussed in [5–7], by taking a derivative with respect to  $\ln \frac{E}{E_0}$  and some re-arrangements.

**3.** Mueller [8] has shown that the small- $x$  evolution equations such as BFKL [9–11], and BK [12, 13], are also based on a very similar branching process, where small- $x$  virtual gluons are released into the light-front wave functions (LFWFs). The same reasoning yields an evolution equation, this time for the generating function for the distribution of the virtual dipoles or  $Z(b, \frac{x}{x_{\min}}, \lambda)$ , which

is exactly of the form (3), with the substitution  $\int d\Omega_k \rightarrow \int d^2b_2$ . Here  $b_2$  denotes the transverse position of the emitted soft gluon, with a different eikonalized emission kernel  $K(b_2; b_0, b_1)$  that on the transverse dipole positions.

In fact, one can show that in the leading order, the BMS and BK equations map onto each through a pertinent conformal transformation [14–16], where the asymptotic real soft gluons at  $t = \infty$ , map onto the virtual gluons present at  $x^+ = 0$  [4]. In this sense, the mapping is a “virtual-real” duality, in addition to the standard interpretation that it maps rapidity divergence to UV divergence [15][17]. In Table I we have highlighted this duality through a parallel between the two constructions.

**4.** The BK (BMS) equation resums *single* logarithms in rapidity (energy). However, in both cases there are two types of divergences instead of one: in the Mueller’s dipole construction for the LFWFs [8], there are UV divergences when  $k_\perp$  becomes large, while in the BMS construction with Wilson-line cusps, there are rapidity-divergences when the emissions become collinear to the Wilson-lines. It is natural to resum the double logarithms for both. One can then show that the double logarithm approximation (DLA) for BK (BMS) is generated from the same branching process, with one more strong ordering in dipole sizes (emitting angles) (see Table I). The strong ordering in virtuality is preserved by the DLA limit. Clearly, the two DLA and the underlying size (angle) orderings, simply map onto each other under a conformal transformation. The similarity of the branching process follows from the non-Abelian three-gluon coupling and large  $N_c$  in QCD, giving rise to a binary tree branching.

**5.** A unique feature of the DLA is that the distribution of dipoles (soft gluons) has a nontrivial KNO scaling function  $f(z)$ , which coincides with that suggested many years ago

for jets in [18–20]. This is due to the same strong ordering in emitting angles in both cases, with Table I making now this plausible.

In the DLA, the equation for the generating function (3) simplifies

$$Z(\rho) = e^{-\rho} + \rho \int_0^1 dx \int_0^1 dy e^{-\rho(1-y)} Z(\rho xy) Z(\rho y)$$

where

$$\rho = \frac{2C_F}{\pi\beta_0} y \ln \ln \frac{Q^2}{M^2}$$

with  $2C_F \sim N_c$  and  $\beta_0 = \frac{11N_c}{12\pi}$  for large  $N_c$ . Defining  $Z = \exp W$ , and introducing  $u = 2\sqrt{\rho}$ , we arrive at the final equation  $\Delta_2 W = e^W - 1$ , where  $\Delta_2$  is a radial part of the 2-dimensional Laplacian. This equation is reminiscent of the Poisson-Boltzmann equation, if  $\Delta_2$  was the full Laplacian. For large energies, it reduces to

$$\frac{d^2 W}{du^2} = e^W - 1. \quad (4)$$

The solution of this equation, encodes the shape of the KNO scaling function  $f(z)$ , through a Fourier-Laplace transform

$$Z(t = -e^u) = \int_0^\infty dz e^{-tz} f(z) \quad (5)$$

A detailed investigation of (4-5) can be found in our recent analysis [4]. With the help of complex-analytic method, we are able to unravel the scaling function  $f(z)$  both in the asymptotic region analytically, and throughout numerically. For  $z < 0.1$ , the scaling function can be made explicit

$$f(z) \sim \frac{\alpha}{z^2} \ln \frac{\alpha}{z} \exp \left( -\frac{1}{2} \ln^2 \frac{\alpha}{z} \ln \frac{\alpha}{z} + \mathcal{O}(1) \right). \quad (6)$$

where  $\alpha = 1.50972$ . For  $z > 2.0$ , the scaling function behaves asymptotically as

$$f(z) = 2r \left( rz - 1 + \mathcal{O} \left( \frac{\ln z}{z} \right) \right) e^{-rz}, \quad z \rightarrow \infty, \quad (7)$$

where  $r = 2.55297$ .

TABLE I. Mueller hierarchy and BMS hierarchy

	Dipole	Cusp
distribution in	<i>virtual</i> gluon in wave function	<i>real</i> gluon in asymptotic state
large $N_c$	yes	yes
kernel	$\frac{b_{10}^2}{b_{11}^2, b_{20}^2}$	$\frac{n \cdot p}{k \cdot n k \cdot p}$
virtual part	TMD soft factor	Sudakov form factor
time ordering	in LF time	in CM time
momentum ordering	decreasing $k^+$	decreasing energy $\omega$
virtuality ordering	increasing	decreasing
Markov process	yes	yes
DLA	$b_{10} \gg b_{12} \gg \dots$	$\theta_{01} \gg \theta_{12} \gg \dots$

Since this scaling function represents a parameter-free QCD prediction for future experiments, including the EIC and EicC, we record our numerical solution in Table II, following from our analysis in [4]. For completeness, we note that a moment reconstruction of  $f(z)$  using different arguments, was used in the context of jets in [18, 20, 21].

In Figure 1, we compare our results recorded in Table II (black-solid curve) with the H1 data for DIS [3] (red data) and the old ALEPH data for  $Z$  decay [22] (grey data). The agreement is very good for both data sets, supporting the universality of our results. For comparison we also show the exact asymptotic (7) (blue-solid curve), and the KNO particle multiplicity  $e^{-z}$  [8], following from the dimensional reduction (diffusive approximation) of Mueller's dipole wave-function evolution (dashed-green curve).

**6.** The knowledge of the effective reduced density matrix for the virtual dipoles in the LFWF [23], and the shape of the KNO function, allow for the evaluation of the entanglement entropy between fast and slow degrees of freedom in DIS [23–26]. For DIS in the DLA with KNO scaling, the result is [4]

$$S_{DIS}(y, Q^2) \rightarrow \ln \bar{n} \equiv 2 \left( \frac{2C_F}{\pi\beta_0} y \ln \ln \frac{Q^2}{M^2} \right)^{\frac{1}{2}} \quad (8)$$

a measure of the Sudakov contribution. (8) is measurable in the DGLAP regime of DIS. We note that the KNO scaling function in the




FIG. 1. The exact (black-solid curve) and asymptotic (blue-solid curve) scaling of the KNO particle multiplicity  $f(z)$  based on Table II, compared to the recent data for DIS [3] (red) and the  $Z$ -decay data at  $\sqrt{s} = M_Z$  by ALEPH [22] (grey).

diffusive (BFKL) regime with  $f(z) = e^{-z}$  [8], leads to maximal decoherence in the entanglement entropy  $S_{BFKL} \sim y$  [23–26]. In contrast, the unimodal character of the scaling function in the DLA, leads to a smaller entanglement entropy  $S_{DIS} \sim \sqrt{y}$ .

Using the BMS-BK duality, we can readily formulate the entanglement entropy between soft and hard degrees of freedom in the final state of  $e^+e^-$  annihilation into hadronic jets

$$S_{e^+e^-}(Q^2) \equiv 2 \left( \frac{C_F}{\pi\beta_0} \ln \frac{Q^2}{M^2} \ln \ln \frac{Q^2}{M^2} \right)^{\frac{1}{2}}. \quad (9)$$

also a measure of the Sudakov contribution (with no extra 2 in the bracket). The rapidity gap between the quark-antiquark pair

TABLE II. Table of the scaling function  $f(z)$ 

$z$	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0
$f(z)$	0.01	0.21	0.45	0.65	0.77	0.82	0.82	0.78	0.72	0.64	0.56	0.49	0.42	0.35	0.29	0.24	0.20	0.15	0.12	0.1

$y \rightarrow \ln \frac{Q^2}{M^2}$ , produces *another logarithm* in  $Q^2$ . Note that for  $e^+e^-$ , this contribution dominates the single logarithm resummation in the BFKL contribution at large  $Q^2$ , which is

$$\ln \bar{n}_{\text{BFKL}} \sim \frac{2C_F \ln 2}{\pi \beta_0} \ln \ln \frac{Q^2}{M^2} \quad (10)$$

The prediction (9) is amenable to experimental verification in high energy hadronic jet physics.

#### Acknowledgements:

We are grateful to Jacek Wosiek for bringing [19] to our attention. This work is supported by the Office of Science, U.S. Department of Energy under Contract No. DE-FG-88ER40388, and by the Priority Research Areas SciMat and DigiWorld under program Excellence Initiative - Research University at the Jagiellonian University.

- 
- [1] A. M. Polyakov, Zh. Eksp. Teor. Fiz. **59**, 542 (1970).
  - [2] Z. Koba, H. B. Nielsen, and P. Olesen, Nucl. Phys. B **40**, 317 (1972).
  - [3] V. Andreev *et al.* (H1), Eur. Phys. J. C **81**, 212 (2021), arXiv:2011.01812 [hep-ex].
  - [4] Y. Liu, M. A. Nowak, and I. Zahed, “Mueller’s dipole wave function in QCD: emergent KNO scaling in the double logarithm limit,” (2022), arXiv:2211.05169 [hep-ph].
  - [5] A. Banfi, G. Marchesini, and G. Smye, JHEP **08**, 006 (2002), arXiv:hep-ph/0206076.
  - [6] G. Marchesini and A. H. Mueller, Phys. Lett. B **575**, 37 (2003), arXiv:hep-ph/0308284.
  - [7] S. Caron-Huot, JHEP **03**, 036 (2018), arXiv:1501.03754 [hep-ph].
  - [8] A. H. Mueller, Nucl. Phys. B **415**, 373 (1994).
  - [9] V. S. Fadin, E. A. Kuraev, and L. N. Lipatov, Phys. Lett. B **60**, 50 (1975).
  - [10] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, Sov. Phys. JETP **44**, 443 (1976).
  - [11] E. A. Kuraev, L. N. Lipatov, and V. S. Fadin, Sov. Phys. JETP **45**, 199 (1977).
  - [12] I. Balitsky, Nucl. Phys. B **463**, 99 (1996), arXiv:hep-ph/9509348.
  - [13] Y. V. Kovchegov, Phys. Rev. D **61**, 074018 (2000), arXiv:hep-ph/9905214.
  - [14] L. Cornalba, “Eikonal methods in AdS/CFT: Regge theory and multi-reggeon exchange,” (2007), arXiv:0710.5480 [hep-th].
  - [15] A. A. Vladimirov, Phys. Rev. Lett. **118**, 062001 (2017), arXiv:1610.05791 [hep-ph].
  - [16] Y. Hatta, E. Iancu, A. H. Mueller, and D. N. Triantafyllopoulos, JHEP **02**, 075 (2018), arXiv:1710.06722 [hep-ph].
  - [17] In non-conformal theory, the exact mapping breaks at two-loop already. But for the virtual part, it can be generalized to all orders.
  - [18] Y. L. Dokshitzer, V. S. Fadin, and V. A. Khoze, Z. Phys. C **15**, 325 (1982).
  - [19] Y. L. Dokshitzer, V. S. Fadin, and V. A. Khoze, Z. Phys. C **18**, 37 (1983).
  - [20] A. Bassetto, M. Ciafaloni, and G. Marchesini, Phys. Rept. **100**, 201 (1983).
  - [21] A. Bassetto, Nucl. Phys. B **303**, 703 (1988).
  - [22] D. Buskulic *et al.* (ALEPH), Z. Phys. C **69**, 15 (1995).
  - [23] Y. Liu, M. A. Nowak, and I. Zahed, accepted by Physical Review D (2022), arXiv:2203.00739 [hep-ph].
  - [24] A. Stoffers and I. Zahed, Phys. Rev. D **88**,

- 025038 (2013), [arXiv:1211.3077 \[nucl-th\]](#).
- [25] D. E. Kharzeev and E. M. Levin, *Phys. Rev. D* **95**, 114008 (2017), [arXiv:1702.03489 \[hep-ph\]](#).
- [26] D. E. Kharzeev and E. Levin, *Phys. Rev. D* **104**, L031503 (2021), [arXiv:2102.09773 \[hep-ph\]](#).
- [27] G. Marchesini and A. H. Mueller, *JHEP* **02**, 010 (2016), [arXiv:1510.08763 \[hep-ph\]](#).
- [28] T. Liou, A. H. Mueller, and S. Munier, *Phys. Rev. D* **95**, 014001 (2017), [arXiv:1608.00852 \[hep-ph\]](#).