

ANALYSIS OF THE $T_{c\bar{s}}(2900)$ AND RELATED TETRAQUARK STATES WITH THE QCD SUM RULES

Xiao-Song Yang^{*†}, Qi Xin^{*†}, Zhi-Gang Wang^{* 1}

Department of Physics, North China Electric Power University, Baoding 071003, P. R. China*
School of Nuclear Science and Engineering, North China Electric Power University, Beijing 102206, P. R. China[†]

Abstract

In this research, we tentatively assign the $T_{c\bar{s}}(2900)$ as the $A\bar{A}$ -type tetraquark state, and study the mass spectrum of the tetraquark states with strange and doubly strange, which have the spin-parity $J^P = 0^+, 1^+$ and 2^+ , in the framework of the QCD sum rules in details, where the A denotes the axialvector diquark state. The predicted mass $M = 2.92 \pm 0.12$ GeV is in consistent with the experimental values $M = 2.892 \pm 0.014 \pm 0.015$ GeV and $2.921 \pm 0.017 \pm 0.020$ GeV from the LHCb collaboration and supports assigning the $T_{c\bar{s}}(2900)$ to be the $A\bar{A}$ -type scalar $c\bar{s}q\bar{q}$ tetraquark state. The predictions for other tetraquark states can be confronted to the experimental data in the future to diagnose the nature of the fully open flavor exotic states.

PACS number: 12.39.Mk, 12.38.Lg

Key words: Tetraquark state, QCD sum rules

1 Introduction

In 2020, the LHCb collaboration carried out the amplitude analysis of the process $B^+ \rightarrow D^+ D^- K^+$ using the LHCb pp collision data taken at $\sqrt{s} = 7, 8$, and 13 TeV corresponding to a total integrated luminosity of 9 fb^{-1} and observed a narrow peak in the $D^- K^+$ invariant mass spectrum [1, 2]. The peak can be parameterized in terms of two Breit-Wigner resonances with the spin-0 and spin-1, the masses and widths are determined to be,

$$\begin{aligned} X_0(2900) &: M = 2.866 \pm 0.007 \pm 0.002 \text{ GeV}, \Gamma = 57 \pm 12 \pm 4 \text{ MeV}, \\ X_1(2900) &: M = 2.904 \pm 0.005 \pm 0.001 \text{ GeV}, \Gamma = 110 \pm 11 \pm 4 \text{ MeV}, \end{aligned} \quad (1)$$

respectively. It is the first exotic structure with fully open flavor and leads to several research works to explore its inner structures, such as the tetraquark state [3, 4, 5, 6, 7, 8], molecular state [9, 10, 11, 12], triangle singularity and cusp effect [13, 14]. Among those interpretations, we adopted the tetraquark state scenario, and studied the axialvector-diquark-axialvector-antidiquark ($A\bar{A}$) type and scalar-diquark-scalar-antidiquark ($S\bar{S}$) type fully open flavor $c\bar{s}u\bar{d}$ tetraquark states with the spin-parity $J^P = 0^+$ via the QCD sum rules, and got the conclusion that it is reasonable to assign the $X_0(2900)$ to be the $A\bar{A}$ -type $c\bar{s}u\bar{d}$ tetraquark state.

Very recently, the LHCb collaboration observed the tetraquark candidates $T_{c\bar{s}0}^a(2900)^{0/++}$ with the spin-parity $J^P = 0^+$ in the processes $B^+ \rightarrow D^- D_s^+ \pi^+$ and $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ with the significance larger than 9σ [15, 16]. The measured Breit-Wigner masses and widths are

$$\begin{aligned} T_{c\bar{s}0}^a(2900)^0 &: M = 2.892 \pm 0.014 \pm 0.015 \text{ GeV}, \Gamma = 0.119 \pm 0.026 \pm 0.013 \text{ GeV}, \\ T_{c\bar{s}0}^a(2900)^{++} &: M = 2.921 \pm 0.017 \pm 0.020 \text{ GeV}, \Gamma = 0.137 \pm 0.032 \pm 0.017 \text{ GeV}, \end{aligned} \quad (2)$$

respectively, such tetraquark candidates maybe exist in the diquark-antidiquark picture [3, 5, 8], due to the light-flavor $SU(3)$ symmetry and the flavor-blinded strong interaction.

They may be the isospin partners of the tetraquark candidates composed of the $c\bar{s}u\bar{d}$ and $c\bar{s}u\bar{d}$, and hence have attracted a lot of research works to comprehend their nature. In the scenario of tetraquark states, the masses of the $T_{c\bar{s}0}^a(2900)^{0/++}$ can be reproduced in the framework of

¹E-mail: zgwang@aliyun.com.

the nonrealistic potential quark model [17], and the color flux-tube model [18]. The $T_{c\bar{s}0}(2900)^0$, $T_{c\bar{s}0}(2900)^{++}$ and $X_0(2900)$ can be accommodated in the light flavor $SU(3)_F$ symmetry sextet [19]. In the scenario of molecular states, the $T_{c\bar{s}0}^a(2900)^{0/++}$ can (cannot) be assigned as the D^*K^* molecular states in the framework of the QCD sum rules [20, 21], the one-boson-exchange model [22], and the effective Lagrangian approach [23] (Bethe-Salpeter equation [24]). On the other hand, we cannot exclude that they are not real resonances and just threshold effects [25, 26]. More works on analogous structures are still needed to diagnose their nature. (After the present work is finished, Ref.[27] appears, where the strong decays of the $T_{c\bar{s}0}^a(2900)^{0/++}$ in the scenario of tetraquark states are studied with the QCD sum rules.)

In our previous works, we have investigated the hidden-charm, doubly-charm, hidden-bottom tetraquark (molecular) states in our special scheme via the QCD sum rules in a comprehensive way, and reach satisfactory assignments of the exotic X , Y , Z and T states and made many predictions [28, 29, 30, 31, 32, 33]. In the present work, we extend our previous works to make possible assignments of the $T_{c\bar{s}0}^a(2900)^{0/++}$, just like what we have done in Ref.[4]. Due to the similar masses, spin-parity and quark constituents of the $T_{c\bar{s}0}^a(2900)^{0/++}$ and $X_0(2900)$, we consider to assign the $T_{c\bar{s}0}^a(2900)^{0/++}$ and $X_0(2900)$ in the same picture as the compact $A\bar{A}$ -type tetraquark states tentatively, and construct the $A\bar{A}$ -type currents to study the ground state mass spectrum of the tetraquark states with strange and doubly strange (and have the spin-parity $J^P = 0^+, 1^+$ and 2^+) via the QCD sum rules to verify the inner structures of the $T_{c\bar{s}0}^a(2900)^{0/++}$.

The article is organized as follows: we get the QCD sum rules for the tetraquark states with strange and doubly strange in Section 2; in Section 3, we present the numerical results and discussions; finally, Section 4 is reserved for our conclusion.

2 QCD sum rules for the tetraquark states

Let us write down the two-point correlation functions at first,

$$\begin{aligned}\Pi(p) &= i \int d^4x e^{ipx} \langle 0 | T \{ J(x) J^\dagger(0) \} | 0 \rangle, \\ \Pi_{\mu\nu\alpha\beta}(p) &= i \int d^4x e^{ipx} \langle 0 | T \{ J_{\mu\nu}(x) J_{\alpha\beta}^\dagger(0) \} | 0 \rangle,\end{aligned}\quad (3)$$

where the currents $J(x) = J_s^0(x)$, $J_{ss}^0(x)$ and $J_{\mu\nu}(x) = J_{s,\mu\nu}^1(x)$, $J_{ss,\mu\nu}^1(x)$, $J_{s,\mu\nu}^2(x)$, $J_{ss,\mu\nu}^2(x)$,

$$\begin{aligned}J_s^0(x) &= \varepsilon^{ijk} \varepsilon^{imn} u_j^T(x) C \gamma_\mu c_k(x) \bar{d}_m(x) \gamma^\mu C \bar{s}_n^T(x), \\ J_{ss}^0(x) &= \varepsilon^{ijk} \varepsilon^{imn} u_j^T(x) C \gamma_\mu c_k(x) \bar{s}_m(x) \gamma^\mu C \bar{s}_n^T(x), \\ J_{s,\mu\nu}^1(x) &= \varepsilon^{ijk} \varepsilon^{imn} [u_j^T(x) C \gamma_\mu c_k(x) \bar{d}_m(x) \gamma_\nu C \bar{s}_n^T(x) - u_j^T(x) C \gamma_\nu c_k(x) \bar{d}_m(x) \gamma_\mu C \bar{s}_n^T(x)], \\ J_{ss,\mu\nu}^1(x) &= \varepsilon^{ijk} \varepsilon^{imn} [u_j^T(x) C \gamma_\mu c_k(x) \bar{s}_m(x) \gamma_\nu C \bar{s}_n^T(x) - u_j^T(x) C \gamma_\nu c_k(x) \bar{s}_m(x) \gamma_\mu C \bar{s}_n^T(x)], \\ J_{s,\mu\nu}^2(x) &= \varepsilon^{ijk} \varepsilon^{imn} [u_j^T(x) C \gamma_\mu c_k(x) \bar{d}_m(x) \gamma_\nu C \bar{s}_n^T(x) + u_j^T(x) C \gamma_\nu c_k(x) \bar{d}_m(x) \gamma_\mu C \bar{s}_n^T(x)], \\ J_{ss,\mu\nu}^2(x) &= \varepsilon^{ijk} \varepsilon^{imn} [u_j^T(x) C \gamma_\mu c_k(x) \bar{s}_m(x) \gamma_\nu C \bar{s}_n^T(x) + u_j^T(x) C \gamma_\nu c_k(x) \bar{s}_m(x) \gamma_\mu C \bar{s}_n^T(x)],\end{aligned}\quad (4)$$

the i, j, k, m and n are color indexes, (thereafter) the superscripts 0, 1 and 2 denote the spins of the currents, the μ and ν are the Lorentz indices, and the subscripts s and ss denote the currents with strange and doubly strange, respectively. With a simple replacement $u \leftrightarrow d$, we obtain the corresponding currents in the same isospin multiplets. In the isospin limit, the tetraquark states in the same multiplets have the same masses.

On the hadron side, we insert a complete set of intermediate hadronic states with the same quantum numbers as the currents into the correlation functions [34, 35], and isolate the ground

state contributions,

$$\begin{aligned}
\Pi^0(p) &= \frac{\lambda_T^2}{M_T^2 - p^2} + \dots, \\
&= \Pi_T(p^2), \\
\Pi_{\mu\nu\alpha\beta}^1(p) &= \frac{\lambda_T^2}{M_T^2 (M_T^2 - p^2)} (p^2 g_{\mu\alpha} g_{\nu\beta} - p^2 g_{\mu\beta} g_{\nu\alpha} - g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta) \\
&\quad + \frac{\lambda_Y^2}{M_Y^2 (M_Y^2 - p^2)} (-g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta) + \dots, \\
&= \tilde{\Pi}_T(p^2) (p^2 g_{\mu\alpha} g_{\nu\beta} - p^2 g_{\mu\beta} g_{\nu\alpha} - g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta) \\
&\quad + \tilde{\Pi}_Y(p^2) (-g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta), \\
\Pi_{\mu\nu\alpha\beta}^2(p) &= \frac{\lambda_T^2}{M_T^2 - p^2} \left(\frac{\tilde{g}_{\mu\alpha} \tilde{g}_{\nu\beta} + \tilde{g}_{\mu\beta} \tilde{g}_{\nu\alpha}}{2} - \frac{\tilde{g}_{\mu\nu} \tilde{g}_{\alpha\beta}}{3} \right) + \dots, \\
&= \Pi_T(p^2) \left(\frac{\tilde{g}_{\mu\alpha} \tilde{g}_{\nu\beta} + \tilde{g}_{\mu\beta} \tilde{g}_{\nu\alpha}}{2} - \frac{\tilde{g}_{\mu\nu} \tilde{g}_{\alpha\beta}}{3} \right) + \dots,
\end{aligned} \tag{5}$$

where $\tilde{g}_{\mu\nu} = g_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}$, the pole residues are defined by

$$\begin{aligned}
\langle 0 | J^0(0) | T(p) \rangle &= \lambda_T, \\
\langle 0 | J_{\mu\nu}^1(0) | T(p) \rangle &= \frac{\lambda_T}{M_T} \varepsilon_{\mu\nu\alpha\beta} \varepsilon^\alpha p^\beta, \\
\langle 0 | J_{\mu\nu}^1(0) | Y(p) \rangle &= \frac{\lambda_Y}{M_Y} (\varepsilon_\mu p_\nu - \varepsilon_\nu p_\mu), \\
\langle 0 | J_{\mu\nu}^2(0) | T(p) \rangle &= \lambda_T \varepsilon_{\mu\nu},
\end{aligned} \tag{6}$$

the current $J_{\mu\nu}^1(x)$ couples potentially to both the $J^P = 1^+$ and 1^- tetraquark states T and Y , respectively, and the ε_μ and $\varepsilon_{\mu\nu}$ are polarization vectors of the tetraquark states. We choose the components $\Pi_T(p^2)$ to explore the properties of the T -states.

We can project out the components $\Pi_T(p^2)$ and $\Pi_Y(p^2)$ via the projective operators $P_T^{\mu\nu\alpha\beta}$ and $P_Y^{\mu\nu\alpha\beta}$,

$$\begin{aligned}
\Pi_T(p^2) &= p^2 \tilde{\Pi}_T(p^2) = P_T^{\mu\nu\alpha\beta} \Pi_{\mu\nu\alpha\beta}^T(p), \\
\Pi_Y(p^2) &= p^2 \tilde{\Pi}_Y(p^2) = P_Y^{\mu\nu\alpha\beta} \Pi_{\mu\nu\alpha\beta}^Y(p),
\end{aligned} \tag{7}$$

where

$$\begin{aligned}
P_T^{\mu\nu\alpha\beta} &= \frac{1}{6} \left(g^{\mu\alpha} - \frac{p^\mu p^\alpha}{p^2} \right) \left(g^{\nu\beta} - \frac{p^\nu p^\beta}{p^2} \right), \\
P_Y^{\mu\nu\alpha\beta} &= \frac{1}{6} \left(g^{\mu\alpha} - \frac{p^\mu p^\alpha}{p^2} \right) \left(g^{\nu\beta} - \frac{p^\nu p^\beta}{p^2} \right) - \frac{1}{6} g^{\mu\alpha} g^{\nu\beta}.
\end{aligned} \tag{8}$$

On the QCD side, we firstly contract the u , d , s and c quark fields in the correlation functions with the Wick's theorem, and obtain one heavy full quark propagator and three light full quark propagators,

$$\begin{aligned}
U/D_{ij}(x) &= \frac{i\delta_{ij} \not{x}}{2\pi^2 x^4} - \frac{\delta_{ij} \langle \bar{q}q \rangle}{12} - \frac{\delta_{ij} x^2 \langle \bar{q}g_s \sigma G q \rangle}{192} - \frac{ig_s G_{\alpha\beta}^a t_{ij}^a (\not{x} \sigma^{\alpha\beta} + \sigma^{\alpha\beta} \not{x})}{32\pi^2 x^2} \\
&\quad - \frac{\delta_{ij} x^4 \langle \bar{q}q \rangle \langle g_s^2 GG \rangle}{27648} - \frac{1}{8} \langle \bar{q}_j \sigma^{\mu\nu} q_i \rangle \sigma_{\mu\nu} + \dots,
\end{aligned} \tag{9}$$

$$S_{ij}(x) = \frac{i\delta_{ij}\not{x}}{2\pi^2x^4} - \frac{\delta_{ij}m_s}{4\pi^2x^2} - \frac{\delta_{ij}\langle\bar{s}s\rangle}{12} + \frac{i\delta_{ij}\not{x}m_s\langle\bar{s}s\rangle}{48} - \frac{\delta_{ij}x^2\langle\bar{s}g_s\sigma Gs\rangle}{192} + \frac{i\delta_{ij}x^2\not{x}m_s\langle\bar{s}g_s\sigma Gs\rangle}{1152} - \frac{ig_sG_{\alpha\beta}^at_{ij}^a(\not{x}\sigma^{\alpha\beta} + \sigma^{\alpha\beta}\not{x})}{32\pi^2x^2} - \frac{\delta_{ij}x^4\langle\bar{s}s\rangle\langle g_s^2GG\rangle}{27648} - \frac{1}{8}\langle\bar{s}_j\sigma^{\mu\nu}s_i\rangle\sigma_{\mu\nu} + \dots, \quad (10)$$

$$C_{ij}(x) = \frac{i}{(2\pi)^4} \int d^4k e^{-ik\cdot x} \left\{ \frac{\delta_{ij}}{\not{k} - m_c} - \frac{g_s G_{\alpha\beta}^n t_{ij}^n}{4} \frac{\sigma^{\alpha\beta}(\not{k} + m_c) + (\not{k} + m_c)\sigma^{\alpha\beta}}{(k^2 - m_c^2)^2} - \frac{g_s^2(t^a t^b)_{ij} G_{\alpha\beta}^a G_{\mu\nu}^b (f^{\alpha\beta\mu\nu} + f^{\alpha\mu\beta\nu} + f^{\alpha\nu\mu\beta})}{4(k^2 - m_c^2)^5} + \dots \right\},$$

$$f^{\alpha\beta\mu\nu} = (\not{k} + m_c)\gamma^\alpha(\not{k} + m_c)\gamma^\beta(\not{k} + m_c)\gamma^\mu(\not{k} + m_c)\gamma^\nu(\not{k} + m_c), \quad (11)$$

where $t^n = \frac{\lambda^n}{2}$, the λ^n is the Gell-Mann matrix [35, 36, 37], then we accomplish all the integrals to obtain the QCD representations. For technical details, the interested readers can consult Refs.[37, 38, 39, 40]. If the heavy quark line emits a gluon and every light quark line contributes a quark-antiquark pair, we can get a quark-gluon operator $g_s G \bar{q} q \bar{q} q$ of dimension 11, its vacuum expectation can be factorized into the form $\langle \bar{q} q \rangle \langle q \bar{q} \rangle \langle q \bar{q} \sigma G q \rangle$ by assuming the vacuum saturation, where $q = u, d$ or s . Therefore, we should calculate the vacuum condensates up to dimension 11. In details, we take account of the vacuum condensates $\langle \bar{q} q \rangle$, $\langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q} g_s \sigma G q \rangle$, $\langle \bar{q} q \rangle^2$, $\langle \bar{q} q \rangle \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q} q \rangle \langle \bar{q} g_s \sigma G q \rangle$, $\langle \bar{q} q \rangle^3$, $\langle \bar{q} g_s \sigma G q \rangle^2$, $\langle \bar{q} q \rangle^2 \langle \frac{\alpha_s GG}{\pi} \rangle$, $\langle \bar{q} q \rangle^2 \langle \bar{q} g_s \sigma G q \rangle$, with the truncations of the operators of the orders $\mathcal{O}(\alpha_s^k)$ with $k \leq 1$ consistently, just like in our previous studies [37, 38, 39, 40].

We obtain the QCD spectral densities $\rho_{QCD}(s)$ through dispersion relation, and the explicit spectral densities $\rho_{QCD}(s)$ are given in the Appendix. Then, we match the hadron spectral densities with QCD spectral densities bellow the continuum thresholds with the assumption of quark-hadron duality, and perform Borel transform in regard to the variable $P^2 = -p^2$ to get the QCD sum rules,

$$\lambda_T^2 \exp\left(-\frac{M_T^2}{T^2}\right) = \int_{m_c^2}^{s_0} ds \rho_{QCD}(s) \exp\left(-\frac{s}{T^2}\right), \quad (12)$$

where the T^2 is the Borel parameter.

Finally, we differentiate Eq.(12) with respect to $\tau = \frac{1}{T^2}$ and obtain the masses of the tetraquark states,

$$M_T^2 = -\frac{\frac{d}{d\tau} \int_{m_c^2}^{s_0} ds \rho_{QCD}(s) \exp(-\tau s)}{\int_{m_c^2}^{s_0} ds \rho_{QCD}(s) \exp(-\tau s)}. \quad (13)$$

3 Numerical results and discussions

We adopt the standard values of the input parameters on the QCD side $\langle \bar{q} q \rangle = -(0.24 \pm 0.01 \text{ GeV})^3$, $\langle \bar{s} s \rangle = (0.8 \pm 0.1) \langle \bar{q} q \rangle$, $\langle \bar{q} g_s \sigma G q \rangle = m_0^2 \langle \bar{q} q \rangle$, $\langle \bar{s} g_s \sigma G s \rangle = m_0^2 \langle \bar{s} s \rangle$, $m_0^2 = (0.8 \pm 0.1) \text{ GeV}^2$, $\langle \frac{\alpha_s GG}{\pi} \rangle = (0.33 \text{ GeV})^4$ at the energy scale $\mu = 1 \text{ GeV}$ [34, 35, 41], and take the modified minimal subtraction masses of the charm quark and strange quark $m_c(m_c) = (1.275 \pm 0.025) \text{ GeV}$ and $m_s(\mu = 2 \text{ GeV}) = (0.095 \pm 0.005) \text{ GeV}$ from the Particle Data Group [42]. Furthermore, we set $m_u = m_d = 0$ because of the small current quark masses and take account of the energy-scale dependence of the input

parameters from the renormalization group equation [43],

$$\begin{aligned}
\langle \bar{q}q \rangle(\mu) &= \langle \bar{q}q \rangle(1 \text{ GeV}) \left[\frac{\alpha_s(1 \text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{12}{33-2n_f}}, \\
\langle \bar{s}s \rangle(\mu) &= \langle \bar{s}s \rangle(1 \text{ GeV}) \left[\frac{\alpha_s(1 \text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{12}{33-2n_f}}, \\
\langle \bar{q}g_s\sigma Gq \rangle(\mu) &= \langle \bar{q}g_s\sigma Gq \rangle(1 \text{ GeV}) \left[\frac{\alpha_s(1 \text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{2}{33-2n_f}}, \\
\langle \bar{s}g_s\sigma Gs \rangle(\mu) &= \langle \bar{s}g_s\sigma Gs \rangle(1 \text{ GeV}) \left[\frac{\alpha_s(1 \text{ GeV})}{\alpha_s(\mu)} \right]^{\frac{2}{33-2n_f}}, \\
m_c(\mu) &= m_c(m_c) \left[\frac{\alpha_s(\mu)}{\alpha_s(m_c)} \right]^{\frac{12}{33-2n_f}}, \\
m_s(\mu) &= m_s(2 \text{ GeV}) \left[\frac{\alpha_s(\mu)}{\alpha_s(2 \text{ GeV})} \right]^{\frac{12}{33-2n_f}}, \\
\alpha_s(\mu) &= \frac{1}{b_0 t} \left[1 - \frac{b_1}{b_0^2} \frac{\log t}{t} + \frac{b_1^2(\log^2 t - \log t - 1) + b_0 b_2}{b_0^4 t^2} \right], \tag{14}
\end{aligned}$$

where $t = \log \frac{\mu^2}{\Lambda^2}$, $b_0 = \frac{33-2n_f}{12\pi}$, $b_1 = \frac{153-19n_f}{24\pi^2}$, $b_2 = \frac{2857-\frac{5033}{9}n_f+\frac{325}{27}n_f^2}{128\pi^3}$, $\Lambda = 213 \text{ MeV}$, 296 MeV and 339 MeV for the flavors $n_f = 5, 4$ and 3 , respectively [42]. We take the flavor numbers $n_f = 4$ (i.e. u, d, s and c) and the typical energy scale $\mu = 1.0 \text{ GeV}$ in the present study as in our previous works [4].

The energy gaps from the ground states to the first radial excited states are $0.5 \sim 0.7 \text{ GeV}$ approximately for the charmed mesons. We tentatively adjust the continuum threshold parameters $\sqrt{s_0} = M_T + 0.5 \sim 0.7 \text{ GeV}$ and Borel parameters T^2 , and search for the best values via trial and error to avoid the contaminations from the higher resonances and continuum states, and to satisfy the two fundamental criteria of the QCD sum rules, which are pole dominance at the hadron side and convergence of the operator product expansion at the QCD side. The pole contributions are defined by

$$\text{pole} = \frac{\int_{m_c^2}^{s_0} ds \rho_{QCD}(s) \exp\left(-\frac{s}{T^2}\right)}{\int_{m_c^2}^{\infty} ds \rho_{QCD}(s) \exp\left(-\frac{s}{T^2}\right)}, \tag{15}$$

and are required to reach the values about 40% – 60% consistently, just like in our previous works [4, 28, 29, 30, 31, 32, 33].

In calculations, we determine the continuum threshold parameters and Borel parameters by the pole contributions alone, then we check whether or not the operator product expansion is convergent automatically. In Table 1, we can see clearly that the maximum ranges of the pole contributions are 38% – 66% and all the central values for the six states are larger than 50%, which implies that the criterion of pole dominance is satisfied pretty well.

In addition, we calculate the contributions of the vacuum condensates in regard to dimension, which is defined by

$$D(n) = \frac{\int_{m_c^2}^{s_0} ds \rho_{n,QCD}(s) \exp\left(-\frac{s}{T^2}\right)}{\int_{m_c^2}^{s_0} ds \rho_{QCD}(s) \exp\left(-\frac{s}{T^2}\right)}, \tag{16}$$

to judge the convergence of the operator product expansion, where the n presents the spectral densities concerning the condensates of dimension n . We plot the results of the $c\bar{s}q\bar{q}$ type tetraquark states as an example in Fig.1, and find that the contributions $D(0)$, $D(3)$, $D(5)$, $D(6)$ and $D(8)$ play




Figure 1: The contributions $D(n)$ for the $c\bar{s}q\bar{q}$ tetraquark states, where A , B and C denote the tetraquark states with $J^P = 0^+$, 1^+ and 2^+ , respectively.

a major role, while the contributions $D(4)$ and $D(7)$ are rather small due to the tiny contributions of the gluon condensate $\langle \frac{\alpha_s G G}{\pi} \rangle$. The absolute values show a descending trend $|D(6)| > |D(8)| > |D(9)| > |D(10)| \sim |D(11)| \sim 0$. The criterion of convergence of the operator product expansion is satisfied very well.

Then we consider all the uncertainties of the input parameters to get the values of the masses and pole residues, and plot the masses of the six tetraquark states with respect to the Borel parameters T^2 in Fig.2. We can see clearly that there appear flat platforms in the Borel windows, which are the regions between the two short perpendicular lines, for all the six states, and thus it declares the reliability of our predictions.

In Table 1, we display the spin-parity, Borel parameters, continuum threshold parameters, pole contributions, masses and pole residues of the tetraquark states with strange and doubly strange. The predicted mass of the $J^P = 0^+$ state $c\bar{u}\bar{d}\bar{s}$, $M = 2.92 \pm 0.12$ GeV, is in very good agreement with the experimental values $M = 2.892 \pm 0.014 \pm 0.015$ GeV and $2.921 \pm 0.017 \pm 0.020$ GeV from the LHCb collaboration [15, 16], and supports assigning the $T_{c\bar{s}}(2900)$ to be the $A\bar{A}$ -type $c\bar{s}q\bar{q}$ tetraquark states with the spin-parity $J^P = 0^+$. The pole residues can be used to calculate the decay widths of the corresponding tetraquark states in the following works. As the experimental data for the fully open flavor exotic states are still far from complete at the present time, we expect for further experimental data to confront with our predictions for the other related tetraquark states to illuminate the nature of the fully open flavor exotic hadrons.

4 Conclusion

In this article, we construct the $A\bar{A}$ -type currents to study the fully open heavy flavor tetraquark states with strange and doubly strange which have the spin-parity $J^P = 0^+$, 1^+ and 2^+ via the QCD sum rules approach in details. We perform the operator product expansion up to the vacuum condensates of dimension 11 in a consistent way, and obtain the masses and pole residues of those tetraquark states. The predicted mass of the $c\bar{u}\bar{d}\bar{s}$ state, $M = 2.92 \pm 0.12$ GeV, is in very good agreement with the experimental data $M = 2.892 \pm 0.014 \pm 0.015$ GeV and $2.921 \pm 0.017 \pm 0.020$ GeV from the LHCb collaboration, as a consequence, it is reasonable to interpret the $T_{c\bar{s}}(2900)$ as the $A\bar{A}$ -type scalar $c\bar{u}\bar{d}\bar{s}$ and $c\bar{d}\bar{u}\bar{s}$ tetraquark states. Besides, we expect more experimental data to

	J^P	$T^2(\text{GeV}^2)$	$\sqrt{s_0}(\text{GeV})$	pole	$M_T(\text{GeV})$	$\lambda_T(10^{-2}\text{GeV}^5)$
$cu\bar{d}\bar{s}$	0^+	$1.9 - 2.3$	3.50 ± 0.10	$(40 - 65)\%$	2.92 ± 0.12	1.62 ± 0.33
$cu\bar{d}\bar{s}$	1^+	$2.2 - 2.6$	3.65 ± 0.10	$(40 - 62)\%$	3.10 ± 0.10	1.59 ± 0.27
$cu\bar{d}\bar{s}$	2^+	$2.5 - 2.9$	3.95 ± 0.10	$(42 - 61)\%$	3.40 ± 0.10	3.51 ± 0.55
$cu\bar{s}\bar{s}$	0^+	$2.0 - 2.4$	3.60 ± 0.10	$(38 - 66)\%$	3.02 ± 0.12	2.72 ± 0.43
$cu\bar{s}\bar{s}$	1^+	$2.3 - 2.7$	3.75 ± 0.10	$(40 - 65)\%$	3.20 ± 0.10	2.64 ± 0.33
$cu\bar{s}\bar{s}$	2^+	$2.6 - 3.0$	4.05 ± 0.10	$(41 - 64)\%$	3.49 ± 0.10	5.70 ± 0.65

Table 1: The spin-parity, Borel parameters, continuum threshold parameters, pole contributions, masses and pole residues of the tetraquark states.




Figure 2: The masses with variations of the Borel parameters T^2 , where the A, B and C (D, E and F) denote the $J^P = 0^+$, 1^+ and 2^+ tetraquark states with strange (doubly strange), respectively.

examine our predictions for the other tetraquark states in the future to illustrate the nature of the T -states.

Appendix

The detailed QCD spectral densities,

$$\rho_{QCD}(s) = \rho_{s/ss}^{0/1/2}(n), \quad (17)$$

where the superscripts 0, 1 and 2 stand for the spins of the tetraquark states, the subscripts s and ss stand for the valence s -quarks, and the n stands for the dimensions of the condensates,

$$\rho_s^0(0) = \frac{1}{1536\pi^6} \int_{x_i}^1 dx x (1-x)^4 (3s - \tilde{m}_c^2) (s - \tilde{m}_c^2)^3,$$

$$\begin{aligned} \rho_s^0(3) = & -\frac{m_s(\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{16\pi^4} \int_{x_i}^1 dx x (1-x)^2 (2s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \\ & -\frac{m_c \langle \bar{q}q \rangle}{32\pi^4} \int_{x_i}^1 dx (1-x)^2 (s - \tilde{m}_c^2)^2, \end{aligned}$$

$$\begin{aligned} \rho_s^0(4) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{1}{768\pi^4} \int_{x_i}^1 dx (1+2x) (1-x)^2 (2s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\ & \left. - \frac{m_c^2}{2304\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} (3s - 2\tilde{m}_c^2) \right\}, \end{aligned}$$

$$\begin{aligned} \rho_s^0(5) = & \frac{m_s(9\langle \bar{q}g_s\sigma Gq \rangle - 8\langle \bar{s}g_s\sigma Gs \rangle)}{384\pi^4} \int_{x_i}^1 dx x (1-x) (3s - 2\tilde{m}_c^2) \\ & - \frac{m_s \langle \bar{q}g_s\sigma Gq \rangle}{256\pi^4} \int_{x_i}^1 dx (1-x)^2 (3s - 2\tilde{m}_c^2) + \frac{m_c \langle \bar{q}g_s\sigma Gq \rangle}{64\pi^4} \int_{x_i}^1 dx (1-x) (s - \tilde{m}_c^2), \end{aligned}$$

$$\rho_s^0(6) = \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{6\pi^2} \int_{x_i}^1 dx x (1-x) (3s - 2\tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle (4\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{12\pi^2} \int_{x_i}^1 dx,$$

$$\begin{aligned} \rho_s^0(7) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{m_s m_c^2 (\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{288\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^2} \left(1 + \frac{s}{T^2}\right) \delta(s - \tilde{m}_c^2) \right. \\ & + \frac{m_c^3 \langle \bar{q}q \rangle}{288\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^3} \delta(s - \tilde{m}_c^2) + \frac{m_c \langle \bar{q}q \rangle}{288\pi^2} \int_{x_i}^1 dx \left(2 - \frac{3(1-x)}{x^2}\right) \\ & + \frac{m_s(4\langle \bar{q}q \rangle + \langle \bar{s}s \rangle)}{384\pi^2} \int_{x_i}^1 dx (1-x) [2 + s\delta(s - \tilde{m}_c^2)] \\ & \left. - \frac{m_s(4\langle \bar{q}q \rangle - 3\langle \bar{s}s \rangle)}{1152\pi^2} \int_{x_i}^1 dx x [2 + s\delta(s - \tilde{m}_c^2)] \right\}, \end{aligned}$$

$$\begin{aligned}
\rho_s^0(8) &= \frac{\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{96\pi^2} \int_{x_i}^1 dx (1-4x) [2 + s\delta(s - \tilde{m}_c^2)] \\
&+ \frac{m_s m_c (8\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 9\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 96\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{576\pi^2} \delta(s - m_c^2) \\
&+ \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle}{48\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s - \tilde{m}_c^2) ,
\end{aligned}$$

$$\rho_s^0(9) = -\frac{4m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{9} \delta(s - m_c^2) ,$$

$$\begin{aligned}
\rho_s^0(10) &= \frac{\langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{1536\pi^2} \left\{ 8 \left(1 + \frac{s}{T^2}\right) \delta(s - m_c^2) + 3 \int_{x_i}^1 dx \left(1 + \frac{s}{T^2}\right) \delta(s - \tilde{m}_c^2) \right\} \\
&+ \frac{m_s m_c^3 \langle \bar{q}g_s \sigma Gq \rangle (8\langle \bar{q}g_s \sigma Gq \rangle - \langle \bar{s}g_s \sigma Gs \rangle)}{384\pi^2 T^4} \delta(s - m_c^2) - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle^2}{192\pi^2 T^2} \delta(s - m_c^2) \\
&+ \langle \frac{\alpha_s GG}{\pi} \rangle \left\{ \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{216} \left[-3 \int_{x_i}^1 dx \left(1 + \frac{s}{T^2}\right) \delta(s - \tilde{m}_c^2) + 2 \left(1 + \frac{s}{T^2}\right) \delta(s - m_c^2) \right] \right. \\
&- \frac{m_s m_c^3 \langle \bar{q}q \rangle (4\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{216T^4} \int_{x_i}^1 dx \frac{1}{x^3} \delta(s - \tilde{m}_c^2) + \frac{m_s m_c^3 \langle \bar{q}q \rangle (8\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{432T^4} \delta(s - m_c^2) \\
&- \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{288T^2} \delta(s - m_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle (4\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{72T^2} \int_{x_i}^1 dx \frac{1}{x^2} \delta(s - \tilde{m}_c^2) \\
&\left. - \frac{m_c^2 \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{108T^4} \int_{x_i}^1 dx \frac{s(1-x)}{x^2} \delta(s - \tilde{m}_c^2) \right\} ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^0(11) &= \frac{m_c^3 \langle \bar{q}q \rangle (\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 2\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{9T^4} \delta(s - m_c^2) \\
&- \frac{m_c \langle \bar{q}q \rangle (\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{36T^2} \delta(s - m_c^2) ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(0) &= \frac{1}{30720\pi^6} \int_{x_i}^1 dx x (1-x)^5 \left[(s - \tilde{m}_c^2)^4 + 12s (s - \tilde{m}_c^2)^3 + 8s^2 (s - \tilde{m}_c^2)^2 \right] \\
&+ \frac{1}{18432\pi^6} \int_{x_i}^1 dx x (1-x)^4 (3s + \tilde{m}_c^2) (s - \tilde{m}_c^2)^3 ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(3) &= -\frac{m_c \langle \bar{q}q \rangle}{288\pi^4} \int_{x_i}^1 dx \left[3(1-x)^2 (s - \tilde{m}_c^2)^2 + (1-x)^3 (5s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \right] \\
&- \frac{m_s \langle \bar{q}q \rangle}{32\pi^4} \int_{x_i}^1 dx x (1-x)^2 (3s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \\
&+ \frac{m_s \langle \bar{s}s \rangle}{576\pi^4} \int_{x_i}^1 dx \left\{ 3x(1-x)^2 (s^2 - \tilde{m}_c^4) + x(1-x)^3 \left[3(s - \tilde{m}_c^2)^2 + 18s(s - \tilde{m}_c^2) + 4s^2 \right] \right\} ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(4) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ -\frac{1}{1152\pi^4} \int_{x_i}^1 dx x (1-x)^2 (2s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\
& + \frac{1}{13824\pi^4} \int_{x_i}^1 dx x (1-x)^3 \left[3 (s - \tilde{m}_c^2)^2 + 18s (s - \tilde{m}_c^2) + 4s^2 \right] \\
& - \frac{1}{55296\pi^4} \int_{x_i}^1 dx (1-x)^3 \left[13 (s - \tilde{m}_c^2)^2 + 4s (s - \tilde{m}_c^2) \right] \\
& - \frac{1}{221184\pi^4} \int_{x_i}^1 dx (1-x)^4 \left[9 (s - \tilde{m}_c^2)^2 + 48s (s - \tilde{m}_c^2) + 8s^2 \right] \\
& - \frac{m_c^2}{69120\pi^4} \int_{x_i}^1 dx \frac{(1-x)^5}{x^2} \left[3 (4s - \tilde{m}_c^2) + 2s^2 \delta(s - \tilde{m}_c^2) \right] \\
& \left. - \frac{m_c^2}{13824\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} \tilde{m}_c^2 \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(5) = & -\frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{576\pi^4} \int_{x_i}^1 dx \left\{ 2x (1-x) \tilde{m}_c^2 + x (1-x)^2 [12s - 3\tilde{m}_c^2 + 2s^2 \delta(s - \tilde{m}_c^2)] \right\} \\
& + \frac{5m_c \langle \bar{q} g_s \sigma G q \rangle}{1152\pi^4} \int_{x_i}^1 dx \left[2 (1-x) (s - \tilde{m}_c^2) + (1-x)^2 (3s - \tilde{m}_c^2) \right] \\
& + \frac{m_s \langle \bar{q} g_s \sigma G q \rangle}{768\pi^4} \int_{x_i}^1 dx (23x - 1) (1-x) (2s - \tilde{m}_c^2),
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(6) = & \frac{\langle \bar{q} q \rangle \langle \bar{s} s \rangle}{6\pi^2} \int_{x_i}^1 dx x (1-x) (2s - \tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q} q \rangle (6\langle \bar{q} q \rangle - \langle \bar{s} s \rangle)}{36\pi^2} \int_{x_i}^1 dx \\
& - \frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle}{36\pi^2} \int_{x_i}^1 dx (1-x) [1 + 2s \delta(s - \tilde{m}_c^2)],
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(7) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ -\frac{m_s \langle \bar{s} s \rangle}{13824\pi^2} \int_{x_i}^1 dx \left\{ 2 (1-x) [17 + 4s \delta(s - \tilde{m}_c^2)] \right. \right. \\
& + (1-x)^2 \left[15 + 2 \left(13s + \frac{4s^2}{T^2} \right) \delta(s - \tilde{m}_c^2) \right] \left. \right\} \\
& + \frac{m_c \langle \bar{q} q \rangle}{6912\pi^2} \int_{x_i}^1 dx \left[32 - \frac{(21x^2 - 31x + 32)(1-x)}{x^2} - \frac{2s(5x^2 - 13x + 24)(1-x)}{x^2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c^2 \langle \bar{s} s \rangle}{5184\pi^2} \int_{x_i}^1 dx \left[\frac{(1-x)^2}{x^2} \left(6 - \frac{3s}{T^2} \right) + \frac{(1-x)^3}{x^2} \left(2 - \frac{s}{T^2} + \frac{2s^2}{T^4} \right) \right] \delta(s - \tilde{m}_c^2) \\
& + \frac{m_c^3 \langle \bar{q} q \rangle}{2592\pi^2} \int_{x_i}^1 dx \left[\frac{(2+x)(1-x)^2}{x^3} + \frac{2s(1-x)^3}{x^3 T^2} \right] \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s (4\langle \bar{q} q \rangle - \langle \bar{s} s \rangle)}{1152\pi^2} \int_{x_i}^1 dx x [1 + s \delta(s - \tilde{m}_c^2)] + \frac{m_s m_c^2 \langle \bar{q} q \rangle}{288\pi^2 T^2} \int_{x_i}^1 dx \frac{s(1-x)^2}{x^2} \delta(s - \tilde{m}_c^2) \\
& \left. + \frac{5m_s \langle \bar{q} q \rangle}{1152\pi^2} \int_{x_i}^1 dx (1-x) [3 + s \delta(s - \tilde{m}_c^2)] \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(8) = & \frac{\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle}{288\pi^2} \int_{x_i}^1 dx (1-12x) [1 + s\delta(s - \tilde{m}_c^2)] \\
& + \frac{m_s m_c (8\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 11\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle - 144\langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{1728\pi^2} \delta(s - m_c^2) \\
& + \frac{m_s m_c (8\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 11\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{1728\pi^2} \int_{x_i}^1 dx \left(-1 + \frac{2s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma Gq \rangle}{288\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s - \tilde{m}_c^2) ,
\end{aligned}$$

$$\rho_s^1(9) = -\frac{2m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{9} \delta(s - m_c^2) ,$$

$$\begin{aligned}
\rho_s^1(10) = & \frac{\langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{4608\pi^2 T^2} \left[40s\delta(s - m_c^2) - 19 \int_{x_i}^1 dx s \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c^3 \langle \bar{q}g_s \sigma Gq \rangle (36\langle \bar{q}g_s \sigma Gq \rangle - 11\langle \bar{s}g_s \sigma Gs \rangle)}{3456\pi^2 T^4} \delta(s - m_c^2) \\
& - \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle (3\langle \bar{q}g_s \sigma Gq \rangle - 11\langle \bar{s}g_s \sigma Gs \rangle)}{3456\pi^2 T^2} \delta(s - m_c^2) \\
& + \langle \frac{\alpha_s G}{\pi} \rangle \left\{ \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{864} \left[-5 \int_{x_i}^1 dx \left(2 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) + \frac{8s}{T^2} \delta(s - m_c^2) \right] \right. \\
& + \frac{m_s m_c^3 \langle \bar{q}q \rangle (4\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{432T^4} \delta(s - m_c^2) + \frac{m_s m_c^3 \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{648} \int_{x_i}^1 dx \frac{1-x}{x^3} \left(-\frac{5}{T^4} + \frac{2s}{T^6} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c^3 \langle \bar{q}q \rangle (6\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{648T^4} \int_{x_i}^1 dx \frac{1}{x^3} \delta(s - \tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle (6\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{216T^2} \int_{x_i}^1 dx \frac{1}{x^2} \delta(s - \tilde{m}_c^2) \\
& + \frac{11m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{5184T^2} \delta(s - m_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{5184} \int_{x_i}^1 dx \frac{24-23x}{x^2} \left(\frac{3}{T^2} - \frac{2s}{T^4} \right) \delta(s - \tilde{m}_c^2) \\
& \left. + \frac{m_c^2 \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{108} \int_{x_i}^1 dx \frac{1-x}{x^2} \left(\frac{1}{T^2} - \frac{s}{T^4} \right) \delta(s - \tilde{m}_c^2) \right\} ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^1(11) = & \frac{m_c^3 \langle \bar{q}q \rangle (\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 2\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{18T^4} \delta(s - m_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle (\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{216T^2} \delta(s - m_c^2) ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(0) = & \frac{1}{46080\pi^6} \int_{x_i}^1 dx x (1-x)^4 (27s - 7\tilde{m}_c^2) (s - \tilde{m}_c^2)^3 \\
& + \frac{1}{76800\pi^6} \int_{x_i}^1 dx x (1-x)^5 \left[3(s - \tilde{m}_c^2)^4 + 44s(s - \tilde{m}_c^2)^3 + 40s^2(s - \tilde{m}_c^2)^2 \right] ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(3) = & -\frac{m_c \langle \bar{q}q \rangle}{144\pi^4} \int_{x_i}^1 dx \left[3(1-x)^2 (s - \tilde{m}_c^2)^2 + (1-x)^3 (5s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \right] \\
& + \frac{m_s \langle \bar{s}s \rangle}{1440\pi^4} \int_{x_i}^1 dx \left\{ 3x(1-x)^2 (17s - 7\tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\
& \left. + x(1-x)^3 \left[9(s - \tilde{m}_c^2)^2 + 66s(s - \tilde{m}_c^2) + 20s^2 \right] \right\} \\
& - \frac{m_s \langle \bar{q}q \rangle}{16\pi^4} \int_{x_i}^1 dx x (1-x)^2 (3s - \tilde{m}_c^2) (s - \tilde{m}_c^2) ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(4) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ -\frac{1}{5760\pi^4} \int_{x_i}^1 dx x (1-x)^2 (59s - 19\tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\
& + \frac{1}{34560\pi^4} \int_{x_i}^1 dx x (1-x)^3 \left[9(s - \tilde{m}_c^2)^2 + 66s(s - \tilde{m}_c^2) + 20s^2 \right] \\
& - \frac{1}{34560\pi^4} \int_{x_i}^1 dx (1-x)^3 (43s - 23\tilde{m}_c^2) (s - \tilde{m}_c^2) \\
& - \frac{1}{34560\pi^4} \int_{x_i}^1 dx (1-x)^4 \left[9(s - \tilde{m}_c^2)^2 + 51s(s - \tilde{m}_c^2) + 10s^2 \right] \\
& - \frac{m_c^2}{172800\pi^4} \int_{x_i}^1 dx \frac{(1-x)^5}{x^2} \left[3(14s - 3\tilde{m}_c^2) + 10s^2 \delta(s - \tilde{m}_c^2) \right] \\
& \left. - \frac{m_c^2}{34560\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} (12s - 7\tilde{m}_c^2) \right\} ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(5) = & \frac{7m_c \langle \bar{q}g_s \sigma Gq \rangle}{576\pi^4} \int_{x_i}^1 dx \left[2(1-x) (s - \tilde{m}_c^2) + (1-x)^2 (3s - \tilde{m}_c^2) \right] \\
& - \frac{m_s \langle \bar{s}g_s \sigma Gs \rangle}{1440\pi^4} \int_{x_i}^1 dx \left\{ 2x(1-x) (12s - 7\tilde{m}_c^2) + x(1-x)^2 [42s - 9\tilde{m}_c^2 + 10s^2 \delta(s - \tilde{m}_c^2)] \right\} \\
& + \frac{m_s \langle \bar{q}g_s \sigma Gq \rangle}{384\pi^4} \int_{x_i}^1 dx (25x + 1) (1-x) (2s - \tilde{m}_c^2) ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(6) = & \frac{\langle \bar{q}q \rangle \langle \bar{s}s \rangle}{3\pi^2} \int_{x_i}^1 dx x (1-x) (2s - \tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle (6\langle \bar{q}q \rangle - \langle \bar{s}s \rangle)}{18\pi^2} \int_{x_i}^1 dx \\
& - \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{18\pi^2} \int_{x_i}^1 dx (1-x) [1 + 2s \delta(s - \tilde{m}_c^2)] ,
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(7) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{m_c^3 \langle \bar{q}q \rangle}{1296\pi^2} \int_{x_i}^1 dx \left[\frac{(2+x)(1-x)^2}{x^3} + \frac{2s(1-x)^3}{x^3 T^2} \right] \delta(s - \tilde{m}_c^2) \right. \\
& + \frac{m_c \langle \bar{q}q \rangle}{864\pi^2} \int_{x_i}^1 dx \left[8 - \frac{(7x^2 - 13x + 8)(1-x)}{x^2} - \frac{2s(7x^2 - 5x + 2)(1-x)}{x^2} \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s \langle \bar{s}s \rangle}{5760\pi^2} \int_{x_i}^1 dx \left\{ 19x^2 - 52x + 43 + \left\{ \left[10 + 33(1-x)^2 \right] s + \frac{10s^2}{T^2} \right\} \delta(s - \tilde{m}_c^2) \right\} \\
& + \frac{m_s \langle \bar{q}q \rangle}{144\pi^2} \int_{x_i}^1 dx \left[3 - 4x + (1-2x)s \delta(s - \tilde{m}_c^2) \right] \\
& - \frac{m_s m_c^2 \langle \bar{s}s \rangle}{12960\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^2} \left[2(1+2x) + \frac{s(8+7x)}{T^2} + \frac{10s^2(1-x)}{T^4} \right] \delta(s - \tilde{m}_c^2) \\
& \left. + \frac{m_s m_c^2 \langle \bar{q}q \rangle}{144\pi^2 T^2} \int_{x_i}^1 dx \frac{s(1-x)^2}{x^2} \delta(s - \tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(8) = & - \frac{\langle \bar{q}q \rangle \langle \bar{s}s \sigma G s \rangle + \langle \bar{s}s \rangle \langle \bar{q}q \sigma G q \rangle}{144\pi^2} \int_{x_i}^1 dx (1+12x) [1 + s \delta(s - \tilde{m}_c^2)] \\
& + \frac{m_s m_c (8 \langle \bar{q}q \rangle \langle \bar{s}s \sigma G s \rangle + 13 \langle \bar{s}s \rangle \langle \bar{q}q \sigma G q \rangle - 144 \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma G q \rangle)}{864\pi^2} \delta(s - m_c^2) \\
& + \frac{m_s m_c (8 \langle \bar{q}q \rangle \langle \bar{s}s \sigma G s \rangle + 13 \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma G q \rangle)}{864\pi^2} \int_{x_i}^1 dx \left(-1 + \frac{2s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{q}g_s \sigma G q \rangle}{144\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s - \tilde{m}_c^2),
\end{aligned}$$

$$\rho_s^2(9) = - \frac{4m_c \langle \bar{q}q \rangle^2 \langle \bar{s}s \rangle}{9} \delta(s - m_c^2),$$

$$\begin{aligned}
\rho_s^2(10) = & \frac{7\langle\bar{q}g_s\sigma Gq\rangle\langle\bar{s}g_s\sigma Gs\rangle}{6912\pi^2T^2} \left[24s\delta(s-m_c^2) + 5\int_{x_i}^1 dxs\delta(s-\tilde{m}_c^2) \right] \\
& + \frac{m_sm_c^3\langle\bar{q}g_s\sigma Gq\rangle(36\langle\bar{q}g_s\sigma Gq\rangle-13\langle\bar{s}g_s\sigma Gs\rangle)}{1728\pi^2T^4}\delta(s-m_c^2) \\
& + \frac{m_sm_c\langle\bar{q}g_s\sigma Gq\rangle(3\langle\bar{q}g_s\sigma Gq\rangle+13\langle\bar{s}g_s\sigma Gs\rangle)}{1728\pi^2T^2}\delta(s-m_c^2) \\
& + \langle\frac{\alpha_s GG}{\pi}\rangle \left\{ \frac{\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{108} \left[-\int_{x_i}^1 dx \left(2 + \frac{s}{T^2} \right) \delta(s-\tilde{m}_c^2) + \frac{2s}{T^2}\delta(s-m_c^2) \right] \right. \\
& + \frac{7m_sm_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{1296T^2}\delta(s-m_c^2) + \frac{m_sm_c^3\langle\bar{q}q\rangle(4\langle\bar{q}q\rangle-\langle\bar{s}s\rangle)}{216T^4}\delta(s-m_c^2) \\
& + \frac{m_sm_c\langle\bar{q}q\rangle(6\langle\bar{q}q\rangle-\langle\bar{s}s\rangle)}{108T^2} \int_{x_i}^1 dx \frac{1}{x^2} \delta(s-\tilde{m}_c^2) \\
& - \frac{m_sm_c^3\langle\bar{q}q\rangle(6\langle\bar{q}q\rangle-\langle\bar{s}s\rangle)}{324T^4} \int_{x_i}^1 dx \frac{1}{x^3} \delta(s-\tilde{m}_c^2) \\
& + \frac{m_sm_c^3\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{324} \int_{x_i}^1 dx \frac{1-x}{x^3} \left(-\frac{5}{T^4} + \frac{2s}{T^6} \right) \delta(s-\tilde{m}_c^2) \\
& + \frac{m_sm_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{1296} \int_{x_i}^1 dx \frac{12-13x}{x^2} \left(\frac{3}{T^2} - \frac{2s}{T^4} \right) \delta(s-\tilde{m}_c^2) \\
& \left. + \frac{m_c^2\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{54} \int_{x_i}^1 dx \frac{1-x}{x^2} \left(\frac{1}{T^2} - \frac{s}{T^4} \right) \delta(s-\tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_s^2(11) = & \frac{m_c\langle\bar{q}q\rangle(\langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle+\langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle)}{108T^2}\delta(s-m_c^2) \\
& + \frac{m_c^3\langle\bar{q}q\rangle(\langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle+2\langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle)}{9T^4}\delta(s-m_c^2),
\end{aligned}$$

$$\rho_{ss}^0(0) = \frac{1}{1536\pi^6} \int_{x_i}^1 dx x (1-x)^4 (3s-\tilde{m}_c^2) (s-\tilde{m}_c^2)^3,$$

$$\rho_{ss}^0(3) = -\frac{m_c\langle\bar{q}q\rangle}{32\pi^4} \int_{x_i}^1 dx (1-x)^2 (s-\tilde{m}_c^2)^2,$$

$$\begin{aligned}
\rho_{ss}^0(4) = & \langle\frac{\alpha_s GG}{\pi}\rangle \left\{ \frac{1}{768\pi^4} \int_{x_i}^1 dx (1+2x)(1-x)^2 (2s-\tilde{m}_c^2) (s-\tilde{m}_c^2) \right. \\
& \left. - \frac{m_c^2}{2304\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} (3s-2\tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^0(5) = & \frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{192\pi^4} \int_{x_i}^1 dx x (1-x) (3s-2\tilde{m}_c^2) - \frac{m_s\langle\bar{s}g_s\sigma Gs\rangle}{128\pi^4} \int_{x_i}^1 dx (1-x)^2 (3s-2\tilde{m}_c^2) \\
& + \frac{m_c\langle\bar{q}g_s\sigma Gq\rangle}{64\pi^4} \int_{x_i}^1 dx (1-x) (s-\tilde{m}_c^2),
\end{aligned}$$

$$\rho_{ss}^0(6) = \frac{\langle \bar{s}s \rangle^2}{6\pi^2} \int_{x_i}^1 dx x (1-x) (3s - 2\tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{2\pi^2} \int_{x_i}^1 dx,$$

$$\begin{aligned} \rho_{ss}^0(7) = & \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{m_c^3 \langle \bar{q}q \rangle}{288\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^3} \delta(s - \tilde{m}_c^2) + \frac{m_c \langle \bar{q}q \rangle}{288\pi^2} \int_{x_i}^1 dx \left(2 - \frac{3(1-x)}{x^2} \right) \right. \\ & \left. + \frac{5m_s \langle \bar{s}s \rangle}{192\pi^2} \int_{x_i}^1 dx (1-x) [2 + s\delta(s - \tilde{m}_c^2)] - \frac{m_s \langle \bar{s}s \rangle}{576\pi^2} \int_{x_i}^1 dx x [2 + s\delta(s - \tilde{m}_c^2)] \right\}, \end{aligned}$$

$$\begin{aligned} \rho_{ss}^0(8) = & \frac{\langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{48\pi^2} \int_{x_i}^1 dx (1-4x) [2 + s\delta(s - \tilde{m}_c^2)] \\ & - \frac{m_s m_c (40\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + 39\langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{288\pi^2} \delta(s - m_c^2) \\ & + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle}{24\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s - \tilde{m}_c^2), \end{aligned}$$

$$\rho_{ss}^0(9) = -\frac{4m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle^2}{9} \delta(s - m_c^2),$$

$$\begin{aligned} \rho_{ss}^0(10) = & \frac{\langle \bar{s}g_s \sigma Gs \rangle^2}{1536\pi^2} \left\{ 8 \left(1 + \frac{s}{T^2} \right) \delta(s - m_c^2) + 3 \int_{x_i}^1 dx \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) \right\} \\ & + \frac{m_s m_c \langle \bar{q}g_s \sigma Gq \rangle \langle \bar{s}g_s \sigma Gs \rangle}{192\pi^2} \left(-\frac{2}{T^2} + \frac{7s}{T^4} \right) \delta(s - m_c^2) \\ & + \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{\langle \bar{s}s \rangle^2}{216} \left[-3 \int_{x_i}^1 dx \left(1 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) + 2 \left(1 + \frac{s}{T^2} \right) \delta(s - m_c^2) \right] \right. \\ & - \frac{m_s m_c^3 \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{36T^4} \int_{x_i}^1 dx \frac{1}{x^3} \delta(s - \tilde{m}_c^2) - \frac{m_c^2 \langle \bar{s}s \rangle^2}{108T^4} \int_{x_i}^1 dx \frac{s(1-x)}{x^2} \delta(s - \tilde{m}_c^2) \\ & \left. + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{12T^2} \int_{x_i}^1 dx \frac{1}{x^2} \delta(s - \tilde{m}_c^2) + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{432} \left(-\frac{3}{T^2} + \frac{14s}{T^4} \right) \delta(s - m_c^2) \right\}, \end{aligned}$$

$$\begin{aligned} \rho_{ss}^0(11) = & \frac{m_c^3 \langle \bar{s}s \rangle (2\langle \bar{q}q \rangle \langle \bar{s}g_s \sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s \sigma Gq \rangle)}{9T^4} \delta(s - m_c^2) \\ & - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle \langle \bar{s}g_s \sigma Gs \rangle}{18T^2} \delta(s - m_c^2), \end{aligned}$$

$$\rho_{ss}^1(0) = \frac{1}{1536\pi^6} \int_{x_i}^1 dx x (1-x)^4 s (s - \tilde{m}_c^2)^3,$$

$$\begin{aligned} \rho_{ss}^1(3) = & -\frac{m_c \langle \bar{q}q \rangle}{288\pi^4} \int_{x_i}^1 dx \left[(4-x)(1-x)^2 (s - \tilde{m}_c^2)^2 + 4(1-x)^3 s (s - \tilde{m}_c^2) \right] \\ & - \frac{m_s \langle \bar{s}s \rangle}{16\pi^4} \int_{x_i}^1 dx x (1-x)^2 (2s - \tilde{m}_c^2) (s - \tilde{m}_c^2), \end{aligned}$$

$$\rho_{ss}^1(4) = -\langle \frac{\alpha_s GG}{\pi} \rangle \left\{ \frac{1}{4608\pi^4} \int_{x_i}^1 dx (1+2x)(1-x)^2 (s-\tilde{m}_c^2)^2 + \frac{m_c^2}{4608\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} s \right\},$$

$$\begin{aligned} \rho_{ss}^1(5) = & -\frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{384\pi^4} \int_{x_i}^1 dx [2(1-19x)(1-x)s - (1-23x)(1-x)\tilde{m}_c^2] \\ & -\frac{5m_c \langle \bar{q} g_s \sigma G q \rangle}{1152\pi^4} \int_{x_i}^1 dx [(3x-5)(1-x)s - (x-3)(1-x)\tilde{m}_c^2], \end{aligned}$$

$$\begin{aligned} \rho_{ss}^1(6) = & \frac{\langle \bar{s} s \rangle^2}{6\pi^2} \int_{x_i}^1 dx x (1-x) (2s - \tilde{m}_c^2) \\ & -\frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle}{18\pi^2} \int_{x_i}^1 dx \{ (1-x) [1 + 2s\delta(s - \tilde{m}_c^2)] - 5 \}, \end{aligned}$$

$$\begin{aligned} \rho_{ss}^1(7) = & \langle \frac{\alpha_s GG}{\pi} \rangle \left\{ -\frac{m_c \langle \bar{q} q \rangle}{864\pi^2} \int_{x_i}^1 dx \left(-4 + \frac{(3-2x)(1-x)}{x^2} \right) \right. \\ & -\frac{3m_s \langle \bar{s} s \rangle}{576\pi^2} \int_{x_i}^1 dx [4x-3+s(2x-1)\delta(s-\tilde{m}_c^2)] \\ & +\frac{m_c^3 \langle \bar{q} q \rangle}{2592\pi^2} \int_{x_i}^1 dx \left[\frac{(2+x)(1-x)^2}{x^3} + \frac{2s(1-x)^3}{x^3 T^2} \right] \delta(s-\tilde{m}_c^2) \\ & +\frac{m_s m_c^2 \langle \bar{s} s \rangle}{288\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^2} \left(1 + \frac{s}{T^2} \right) \delta(s-\tilde{m}_c^2) \\ & \left. -\frac{m_c \langle \bar{q} q \rangle}{1728\pi^2} \int_{x_i}^1 dx \frac{(5x^2-3x+2)(1-x)}{x^2} [1 + 2s\delta(s-\tilde{m}_c^2)] \right\}, \end{aligned}$$

$$\begin{aligned} \rho_{ss}^1(8) = & -\frac{\langle \bar{s} s \rangle \langle \bar{s} g_s \sigma G s \rangle}{144\pi^2} \int_{x_i}^1 dx (12x-1) [1 + s\delta(s-\tilde{m}_c^2)] \\ & -\frac{m_s m_c (64\langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + 61\langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle)}{864\pi^2} \delta(s-\tilde{m}_c^2) \\ & -\frac{m_s m_c (8\langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + 11\langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle)}{864\pi^2} \int_{x_i}^1 dx \left(1 - \frac{2s}{T^2} \right) \delta(s-\tilde{m}_c^2) \\ & +\frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle}{144\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s-\tilde{m}_c^2), \end{aligned}$$

$$\rho_{ss}^1(9) = -\frac{2m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle^2}{9} \delta(s-\tilde{m}_c^2),$$

$$\begin{aligned}
\rho_{ss}^1(10) = & -\frac{\langle \bar{s}g_s\sigma Gs \rangle^2}{4608\pi^2} \left[-\frac{40s}{T^2} \delta(s - m_c^2) + 19 \int_{x_i}^1 dx \frac{s}{T^2} \delta(s - \tilde{m}_c^2) \right] \\
& + \frac{m_s m_c \langle \bar{q}g_s\sigma Gq \rangle \langle \bar{s}g_s\sigma Gs \rangle}{1728\pi^2} \left(\frac{8}{T^2} + \frac{25s}{T^4} \right) \delta(s - m_c^2) \\
& + \langle \frac{\alpha_s GG}{\pi} \rangle \left\{ -\frac{\langle \bar{s}s \rangle^2}{216} \left[\int_{x_i}^1 dx \left(2 + \frac{s}{T^2} \right) \delta(s - \tilde{m}_c^2) - \frac{2s}{T^2} \delta(s - m_c^2) \right] \right. \\
& + \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{1296} \left(\frac{5}{T^2} + \frac{18s}{T^4} \right) \delta(s - m_c^2) \\
& - \frac{m_s m_c^3 \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{324} \int_{x_i}^1 dx \left(\frac{5(2-x)}{x^3 T^4} - \frac{2s(1-x)}{x^3 T^6} \right) \delta(s - \tilde{m}_c^2) \\
& - \frac{m_s m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle}{1296} \int_{x_i}^1 dx \left(-\frac{3(32-11x)}{x^2 T^2} + \frac{2s(12-11x)}{x^2 T^4} \right) \delta(s - \tilde{m}_c^2) \\
& \left. - \frac{m_c^2 \langle \bar{s}s \rangle^2}{108} \int_{x_i}^1 dx \frac{1-x}{x^2} \left(-\frac{1}{T^2} + \frac{s}{T^4} \right) \delta(s - \tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^1(11) = & \frac{m_c^3 \langle \bar{s}s \rangle (2 \langle \bar{q}q \rangle \langle \bar{s}g_s\sigma Gs \rangle + \langle \bar{s}s \rangle \langle \bar{q}g_s\sigma Gq \rangle)}{18T^4} \delta(s - m_c^2) \\
& - \frac{m_c \langle \bar{q}q \rangle \langle \bar{s}s \rangle \langle \bar{s}g_s\sigma Gs \rangle}{108T^2} \delta(s - m_c^2),
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(0) = & \frac{1}{38400\pi^6} \int_{x_i}^1 dx x (1-x)^5 \left[(s - \tilde{m}_c^2)^4 + 16s (s - \tilde{m}_c^2)^3 + 16s^2 (s - \tilde{m}_c^2)^2 \right] \\
& + \frac{1}{11520\pi^6} \int_{x_i}^1 dx x (1-x)^4 (9s - 2\tilde{m}_c^2) (s - \tilde{m}_c^2)^3,
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(3) = & -\frac{m_s \langle \bar{s}s \rangle}{360\pi^4} \int_{x_i}^1 dx \left\{ 3x(1-x)^2 (34s - 11\tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\
& - x(1-x)^3 \left[3(s - \tilde{m}_c^2)^2 + 24s(s - \tilde{m}_c^2) + 8s^2 \right] \left. \right\} \\
& - \frac{m_c \langle \bar{q}q \rangle}{144\pi^4} \int_{x_i}^1 dx \left[3(1-x)^2 (s - \tilde{m}_c^2)^2 + (1-x)^3 (5s - \tilde{m}_c^2) (s - \tilde{m}_c^2) \right],
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(4) = & \langle \frac{\alpha_s GG}{\pi} \rangle \left\{ -\frac{1}{34560\pi^4} \int_{x_i}^1 dx (1+2x)(1-x)^2 (113s - 37\tilde{m}_c^2) (s - \tilde{m}_c^2) \right. \\
& + \frac{1}{69120\pi^4} \int_{x_i}^1 dx (1+3x)(1-x)^3 \left[3(s - \tilde{m}_c^2)^2 + 24s(s - \tilde{m}_c^2) + 8s^2 \right] \\
& - \frac{m_c^2}{43200\pi^4} \int_{x_i}^1 dx \frac{(1-x)^5}{x^2} \left[3(7s - 5\tilde{m}_c^2) + 2s^2 \delta(s - \tilde{m}_c^2) \right] \\
& \left. - \frac{m_c^2}{34560\pi^4} \int_{x_i}^1 dx \frac{(1-x)^4}{x^2} (15s - 8\tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(5) &= \frac{7m_c \langle \bar{q} g_s \sigma G q \rangle}{288\pi^4} \int_{x_i}^1 dx \left[(1-x)(s - \tilde{m}_c^2) + (1-x)^2(3s - \tilde{m}_c^2) \right] \\
&+ \frac{m_s \langle \bar{s} g_s \sigma G s \rangle}{5760\pi^4} \int_{x_i}^1 dx \left\{ x(1-x)(1320s - 652\tilde{m}_c^2) \right. \\
&\left. - 16x(1-x)^2[3(5s - \tilde{m}_c^2) + 4s^2\delta(s - \tilde{m}_c^2)] + 30(1-x)^2(2s - \tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(6) &= -\frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle}{9\pi^2} \int_{x_i}^1 dx \left\{ (1-x)[1 + 2s\delta(s - \tilde{m}_c^2)] - 5 \right\} \\
&+ \frac{\langle \bar{s} s \rangle^2}{3\pi^2} \int_{x_i}^1 dx x(1-x)(2s - \tilde{m}_c^2),
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(7) &= \left\langle \frac{\alpha_s GG}{\pi} \right\rangle \left\{ \frac{m_c^3 \langle \bar{q} q \rangle}{1296\pi^2} \int_{x_i}^1 dx \left[\frac{(2+x)(1-x)^2}{x^3} + \frac{2s(1-x)^3}{x^3 T^2} \right] \delta(s - \tilde{m}_c^2) \right. \\
&- \frac{m_c \langle \bar{q} q \rangle}{864\pi^2} \int_{x_i}^1 dx \left[-8 + \frac{(7x^2 - 13x + 8)(1-x)}{x^2} + \frac{2s(7x^2 - 5x + 2)(1-x)}{x^2} \delta(s - \tilde{m}_c^2) \right] \\
&- \frac{m_s \langle \bar{s} s \rangle}{8640\pi^2} \int_{x_i}^1 dx \left\{ 150x[1 + s\delta(s - \tilde{m}_c^2)] + 2(1-x)[-121 + s\delta(s - \tilde{m}_c^2)] \right. \\
&- 4(1-x)^2 \left[3 + 4 \left(s + \frac{s^2}{T^2} \right) \delta(s - \tilde{m}_c^2) \right] \left. \right\} \\
&- \frac{m_s m_c^2 \langle \bar{s} s \rangle}{6480\pi^2} \int_{x_i}^1 dx \frac{(1-x)^2}{x^2} \left[(1+2x) + \frac{s(8x-77)}{T^2} + \frac{8s^2(1-x)}{T^4} \right] \delta(s - \tilde{m}_c^2) \left. \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(8) &= -\frac{\langle \bar{s} s \rangle \langle \bar{s} g_s \sigma G s \rangle}{72\pi^2} \int_{x_i}^1 dx (12x + 1)[1 + s\delta(s - \tilde{m}_c^2)] \\
&- \frac{m_s m_c (64 \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + 59 \langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle)}{432\pi^2} \delta(s - m_c^2) \\
&- \frac{m_s m_c (8 \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle + 13 \langle \bar{s} s \rangle \langle \bar{q} g_s \sigma G q \rangle)}{432\pi^2} \int_{x_i}^1 dx \left(1 - \frac{2s}{T^2} \right) \delta(s - \tilde{m}_c^2) \\
&- \frac{m_s m_c \langle \bar{q} q \rangle \langle \bar{s} g_s \sigma G s \rangle}{72\pi^2} \int_{x_i}^1 dx \frac{1}{x} \delta(s - \tilde{m}_c^2),
\end{aligned}$$

$$\rho_{ss}^2(9) = -\frac{4m_c \langle \bar{q} q \rangle \langle \bar{s} s \rangle^2}{9} \delta(s - m_c^2),$$

$$\begin{aligned}
\rho_{ss}^2(10) = & \frac{7\langle\bar{s}g_s\sigma Gs\rangle^2}{2304\pi^2} \left[\frac{8s}{T^2}\delta(s-m_c^2) + \frac{5}{3}\int_{x_i}^1 dx \frac{s}{T^2}\delta(s-\tilde{m}_c^2) \right] \\
& + \frac{m_sm_c\langle\bar{q}g_s\sigma Gq\rangle\langle\bar{s}g_s\sigma Gs\rangle}{864\pi^2} \left(\frac{16}{T^2} + \frac{23s}{T^4} \right) \delta(s-m_c^2) \\
& + \langle\frac{\alpha_s GG}{\pi}\rangle \left\{ -\frac{\langle\bar{s}s\rangle^2}{108} \left[\int_{x_i}^1 dx \left(2 + \frac{s}{T^2} \right) \delta(s-\tilde{m}_c^2) - \frac{2s}{T^2}\delta(s-m_c^2) \right] \right. \\
& + \frac{m_sm_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{648} \left(\frac{7}{T^2} + \frac{18s}{T^4} \right) \delta(s-m_c^2) \\
& - \frac{m_sm_c^3\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{162} \int_{x_i}^1 dx \left(\frac{5(2-x)}{x^3T^4} - \frac{2s(1-x)}{x^3T^6} \right) \delta(s-\tilde{m}_c^2) \\
& - \frac{m_sm_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle}{648} \int_{x_i}^1 dx \left(\frac{3(13x-32)}{x^2T^2} + \frac{2s(12-13x)}{x^2T^4} \right) \delta(s-\tilde{m}_c^2) \\
& \left. - \frac{m_c^2\langle\bar{s}s\rangle^2}{54} \int_{x_i}^1 dx \frac{1-x}{x^2} \left(-\frac{1}{T^2} + \frac{s}{T^4} \right) \delta(s-\tilde{m}_c^2) \right\},
\end{aligned}$$

$$\begin{aligned}
\rho_{ss}^2(11) = & \frac{m_c^3\langle\bar{s}s\rangle(2\langle\bar{q}q\rangle\langle\bar{s}g_s\sigma Gs\rangle + \langle\bar{s}s\rangle\langle\bar{q}g_s\sigma Gq\rangle)}{9T^4} \delta(s-m_c^2) \\
& + \frac{m_c\langle\bar{q}q\rangle\langle\bar{s}s\rangle\langle\bar{s}g_s\sigma Gs\rangle}{54T^2} \delta(s-m_c^2),
\end{aligned}$$

where $\tilde{m}_c^2 = \frac{m_c^2}{x}$ and $x_i = \frac{m_c^2}{s}$.

Acknowledgements

This work is supported by National Natural Science Foundation, Grant Number 12175068.

References

- [1] R. Aaij et al., Phys. Rev. **D102** (2020) 112003.
- [2] R. Aaij et al., Phys. Rev. Lett. **125** (2020) 242001.
- [3] Q. F. Lu, D. Y. Chen and Y. B. Dong, Phys. Rev. **D102** (2020) 074021.
- [4] Z. G. Wang, Int. J. Mod. Phys. **A35** (2020) 2050187.
- [5] X. G. He, W. Wang and R. L. Zhu, Eur. Phys. J. **C80** (2020) 1026.
- [6] M. Karliner and J. L. Rosner, Phys. Rev. **D102** (2020) 094016.
- [7] J. R. Zhang, Phys. Rev. **D103** (2021) 054019.
- [8] S. S. Agaev, K. Azizi and H. Sundu, Phys. Lett. **B820** (2021) 136530.
- [9] M. W. Hu, X. Y. Lao, P. Ling and Q. Wang, Chin. Phys. **C45** (2021) 021003.
- [10] M. Z. Liu, J. J. Xie and L. S. Geng, Phys. Rev. **D102** (2020) 091502.
- [11] H. X. Chen, W. Chen, R. R. Dong and N. Su, Chin. Phys. Lett. **37** (2020) 101201.

- [12] Y. Huang, J. X. Lu, J. J. Xie and L. S. Geng, Eur. Phys. J. **C80** (2020) 973.
- [13] X. H. Liu, M. J. Yan, H. W. Ke, G. Li and J. J. Xie, Eur. Phys. J. **C80** (2020) 1178.
- [14] T. J. Burns and E. S. Swanson, Phys. Lett. **B813** (2021) 136057.
- [15] R. Aaij et al, arXiv:2212.02716 [hep-ex].
- [16] R. Aaij et al, arXiv:2212.02717 [hep-ex].
- [17] F. X. Liu, R. H. Ni, X. H. Zhong and Q. Zhao, arXiv:2211.01711 [hep-ph].
- [18] J. Wei, Y. H. Wang, C. S. An and C. R. Deng, Phys. Rev. **D106** (2022) 096023.
- [19] V. Dmitrasinovic, arXiv:2301.05471 [hep-ph].
- [20] S. S. Agaev, K. Azizi and H. Sundu, arXiv:2207.02648 [hep-ph].
- [21] S. S. Agaev, K. Azizi and H. Sundu, arXiv:2212.12001 [hep-ph].
- [22] R. Chen and Q. Huang, arXiv:2208.10196 [hep-ph].
- [23] Z. L. Yue, C. J. Xiao, D. Y. Chen, arXiv:2212.03018 [hep-ph].
- [24] H. W. Ke, Y. F. Shi, X. H. Liu and X. Q. Li, Phys. Rev. **D106** (2022) 114032.
- [25] Y. H. Ge, X. H. Liu and H. W. Ke, Eur. Phys. J. **C82** (2022) 955.
- [26] R. Molina and E. Oset, arXiv:2211.01302 [hep-ph].
- [27] D. K. Lian, W. Chen, H. X. Chen, L. Y. Dai and T. G. Steele, arXiv:2302.01167 [hep-ph].
- [28] Z. G. Wang, Chin. Phys. **C45** (2021) 073107.
- [29] Z. G. Wang and Z. H. Yan, Eur. Phys. J. **C78** (2018) 19.
- [30] Q. Xin and Z. G. Wang, Eur. Phys. J. **A58** (2022) 110.
- [31] Z. G. Wang, Phys. Rev. **D102** (2020) 014018.
- [32] Z. G. Wang, Eur. Phys. J. **C79** (2019) 489.
- [33] Z. G. Wang, Int. J. Mod. Phys. **A36** (2021) 2150107.
- [34] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385; Nucl. Phys. **B147** (1979) 448.
- [35] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.
- [36] P. Pascual and R. Tarrach, "QCD: Renormalization for the practitioner", Springer Berlin Heidelberg (1984).
- [37] Z. G. Wang and T. Huang, Phys. Rev. **D89** (2014) 054019.
- [38] Z. G. Wang and T. Huang, Eur. Phys. J. **C74** (2014) 2891.
- [39] Z. G. Wang, Eur. Phys. J. **C74** (2014) 2874.
- [40] Z. G. Wang, Int. J. Mod. Phys. **A35** (2020) 2050003.
- [41] P. Colangelo and A. Khodjamirian, hep-ph/0010175.
- [42] P. A. Zyla et al., Prog. Theor. Exp. Phys. **2020** (2020) 083C01.
- [43] S. Narison and R. Tarrach, Phys. Lett. **125B** (1983) 217.