

ORIENTED AND STANDARD SHADOWING PROPERTIES ON CLOSED SURFACES

SOGO MURAKAMI

ABSTRACT. We prove that oriented and standard shadowing properties are equivalent for topological flows on closed surfaces with the nonwandering set consisting of the finite number of critical elements (i.e., singularities or closed orbits). Moreover, we prove that each isolated singularity of a topological flow on a closed surface with the oriented shadowing property is either asymptotically stable, backward asymptotically stable, or admits a neighborhood which splits into two or four hyperbolic sectors.

1. INTRODUCTION

In the qualitative study of dynamical systems, the shadowing property has been a key property in the hyperbolic theory (see [9] or [12] for instance), which is useful in the proof of the structural stability of hyperbolic systems. One of the highlights of the studies on the relation between various shadowing properties and the structural stability or the Ω -stability is due to Pilyugin and Tikhomirov, who proved that the Lipschitz shadowing property and the structural stability are equivalent for C^1 vector fields on a compact manifold [10]. The Lipschitz shadowing property is stronger than the standard shadowing property, while there are several shadowing properties weaker than the standard one. Among them, we can find the oriented shadowing property. This concerns only time-orientations of orbits, allowing orientation preserving time-reparametrizations for true orbits in order to shadow pseudotrajectories. It is an interesting problem to know when the oriented shadowing property implies the standard one.

As a classical work, Komuro [6] proved the equivalence of oriented and standard shadowing properties for topological flows on a compact metric space without singularities. Recently, this has been extended to those with finitely many Lyapunov stable or backward Lyapunov stable singularities [8]. When the compact metric space is changed to a closed surface, it is expected that a stronger property holds. However, we have an important example by Tikhomirov [13] for which the equivalence does not hold. The example is a C^1 vector field on a four-dimensional closed manifold, whose nonwandering set consists of finitely many singularities including two hyperbolic saddles with index two. In this paper, we prove the equivalence for topological flows on a closed surface whose nonwandering set consists of a finite number of critical elements. Note that all hyperbolic singularities of saddle type have index one here, so the situation that Tikhomirov constructed in his example cannot occur in this case.

2010 *Mathematics Subject Classification.* Primary 37C50.

Key words and phrases. topological flows; shadowing; hyperbolic sectors.

First we introduce some notation. Let (M, dist) be a C^0 closed surface with a distance function on M . Take a topological flow ϕ on M (i.e., $\phi : \mathbb{R} \times M \rightarrow M$ is a continuous map satisfying $\phi(0, x) = x$ and $\phi(s + t, x) = \phi(s, \phi(t, x))$ for all $s, t \in \mathbb{R}$ and $x \in M$).

We say that $\xi : \mathbb{R} \rightarrow M$ is a d -pseudotrajectory of ϕ if

$$\text{dist}(\xi(t + s), \phi(\xi(t), s)) < d$$

for all $t \in \mathbb{R}$ and $s \in [0, 1]$. Let $\text{Ps}(d)$ be the set of all d -pseudotrajectories of ϕ . Denote by Rep the set of all homeomorphisms from \mathbb{R} to \mathbb{R} which preserves the orientation. For $\varepsilon > 0$, let

$$\text{Rep}(\varepsilon) = \left\{ f \in \text{Rep}; \left| \frac{f(a) - f(b)}{a - b} - 1 \right| < \varepsilon, \forall a, b \in \mathbb{R}, a > b \right\}.$$

We say that a topological flow ϕ has the *standard shadowing property* if for every $\varepsilon > 0$ there exists $d > 0$ such that if $\xi \in \text{Ps}(d)$ then

$$\text{dist}(\xi(t), \phi(h(t), x)) < \varepsilon, \quad t \in \mathbb{R}$$

for some $x \in M$ and $h \in \text{Rep}(\varepsilon)$. Denote the set of all topological flows with the standard shadowing property by $\text{StSh}(M)$. As a weaker form of shadowing properties, we say that a topological flow ϕ has the *oriented shadowing property* if for every $\varepsilon > 0$ there exists $d > 0$ such that if $\xi \in \text{Ps}(d)$ then

$$\text{dist}(\xi(t), \phi(h(t), x)) < \varepsilon, \quad t \in \mathbb{R}$$

for some $x \in M$ and $h \in \text{Rep}$. Denote the set of all topological flows with the oriented shadowing property by $\text{OrientSh}(M)$. Singularities (i.e., fixed points for topological flows) and closed orbits of ϕ are called *critical elements*. Let $\text{Sing}(\phi)$ be the set of all singularities of ϕ , and let $\text{Crit}(\phi)$ be that of all critical elements of ϕ . We say that a ϕ -invariant compact set K is *asymptotically stable* if for any neighborhood V of K , there exists a neighborhood U of K such that

$$\phi(t, x) \in V, \quad t \geq 0$$

and

$$\phi(t, x) \rightarrow p, \quad t \rightarrow \infty$$

for all $x \in U$. Similarly, a ϕ -invariant compact set K is said to be *backward asymptotically stable* if for any neighborhood V of K , there exists a neighborhood U of K such that


$$\phi(t, x) \in V, \quad t \leq 0$$

and


$$\phi(t, x) \rightarrow p, \quad t \rightarrow -\infty$$

for all $x \in U$. For $r > 0$ and $x \in M$, let $B(r, x)$ be an open ball centered at x with radius r . Denote the α -limit set (resp. ω -limit set) of x by $\alpha(x)$ (resp. $\omega(x)$), and denote the nonwandering set of ϕ by $\Omega(\phi)$.


Our first theorem is a classification result on singularities of $\phi \in \text{OrientSh}(M)$. There are only four types of singularities, which are simple enough to understand local dynamical behavior around singularities. A similar result can be found in [7] but ours is more precise and based on the theory of sectors [5] that was not used in [7]. So, we provide a full proof of it for completeness.




(a) Asymptotically stable singularity.



(b) Backward asymptotically stable singularity.



(c) Singularity with four hyperbolic sectors.



(d) Singularity with two hyperbolic sectors.

Figure 1. Four types of singularities in Theorem 1.1.

Theorem 1.1. *Let M be a C^0 closed surface and let $\phi \in \text{OrientSh}(M)$ with $p \in \text{Sing}(\phi)$. Suppose that p has a neighborhood D homeomorphic to a closed disk in \mathbb{R}^2 without containing any α -limit point or ω -limit point except p . Then one of the following properties hold:*

- (a) p is asymptotically stable.
- (b) p is backward asymptotically stable.
- (c) D has an open dense subset consisting of four hyperbolic sectors associated with ∂D .
- (d) D has an open dense subset consisting of two hyperbolic sectors associated with ∂D .

We give definitions of hyperbolic sectors in Section 2 (see also Figure 1). Using Theorem 1.1, we prove the equivalence between oriented and standard shadowing properties.

Theorem 1.2. *Let M be a C^0 closed surface, and let ϕ be a topological flow on M . Suppose that the nonwandering set $\Omega(\phi)$ consists of a finite number of critical elements. Then the following conditions are equivalent:*

- $\phi \in \text{StSh}(M)$.
- $\phi \in \text{OrientSh}(M)$.

It is trivial that $\text{StSh}(M) \subset \text{OrientSh}(M)$. For the proof of Theorem 1.2, we need to show that $\text{OrientSh}(M) \subset \text{StSh}(M)$. The following corollary is immediate from Theorem 1.2.

Corollary 1.3. *Let M be a smooth closed surface, and let ϕ be a C^1 Axiom A flow on M . Then the following conditions are equivalent:*

- $\phi \in \text{StSh}(M)$.
- $\phi \in \text{OrientSh}(M)$.

In Section 2, several propositions and lemmas are given, which play important roles in the proofs of the main theorems. In Section 3, Theorem 1.1 is proved. In Section 4, we reduce Theorem 1.2 to a proposition, and introduce some notations and lemmas. In Sections 5 and 6, the behavior of pseudotrajectories near critical elements is investigated in order to prove Theorem 1.2. In Section 7, we finish the proof of Theorem 1.2.

2. PRELIMINARIES

In this section, we give several lemmas and propositions that were proved elsewhere, which are basic to the proof of Theorems 1.1 and 1.2. The first lemma is essentially proved in [2].

Lemma 2.1 ([2]). *Let $x_0 \in M \setminus \text{Sing}(\phi)$ and let U be a neighborhood of x_0 . Then there exist a set S containing x_0 and $\tau_0 > 0$ such that the map*

$$\varphi : (-\tau_0, \tau_0) \times S \ni (x, t) \mapsto \phi(x, t) \in U$$

is a homeomorphism onto its image that is an open neighborhood of x_0 .

Remark. The fact that $\text{Im}\varphi$ contains a neighborhood of x_0 follows from an easy additional argument of the proof of [2, Theorem 2.9], which is omitted here.

The following lemma is immediate from the Jordan-Schoenflies theorem [3].

Lemma 2.2 ([3]). *Let C be a simple closed curve in \mathbb{R}^2 . Then for all $p \notin C$ and $q \in C$, there exists a continuous map $i : [0, 1] \rightarrow \mathbb{R}^2$ such that $i(0) = p$, $i(1) = q$ and*

$$i([0, 1]) \cap C = \emptyset.$$

The following lemma is [14, Theorem 31.2, Theorem 31.5].

Lemma 2.3 ([14]). *Let X be a Hausdorff topological space, and let $G : [-1, 1] \rightarrow X$ be a surjective continuous map with $G(-1) \neq G(1)$. Then there exists an injective continuous map $j : [-1, 1] \rightarrow X$ such that $j(-1) = G(-1)$ and $j(1) = G(1)$.*

Using these lemmas, we can prove the following proposition, which claims that the flow box theorem is valid for topological flows on a closed surface. This fact has been used in many papers (see [4] or [1] for instance), but as far as the author knows the proof has not been given explicitly. We give a proof here for completeness.

Proposition 2.4. *Let $x_0 \in M \setminus \text{Sing}(\phi)$ and let U be a neighborhood of x_0 disjoint from the limit set of ϕ . Then there exist an embedded closed interval I with x_0 and $\tau_0 > 0$ such that the map*

$$\psi : (-\tau_0, \tau_0) \times I \ni (t, x) \mapsto \phi(t, x) \in U$$

is a homeomorphism onto its image that is a neighborhood of x_0 .

Proof. By Lemma 2.1, there exist a set S with x_0 and $\tau_0 > 0$ such that the map

$$\varphi : (-\tau_0, \tau_0) \times S \ni (t, x) \mapsto \phi(t, x) \in M$$

is a homeomorphism onto its image that is an open neighborhood of x_0 . Making U smaller if necessary, we may assume that U is homeomorphic to an open disk in \mathbb{R}^2 and $U \subset \text{Im}\varphi$. Let $C \subset U$ be a simple closed curve surrounding x_0 . Since U and the limit set of ϕ are disjoint, we can take

$$\begin{aligned} t_1 &= \sup\{t < 0; \phi(t, x_0) \in C\}, \\ t_2 &= \inf\{t > 0; \phi(t, x_0) \in C\}. \end{aligned}$$

Set $y_i = \phi(t_i, x_0)$ for $i = 1, 2$. Then C is divided into two subarcs C_1 and C_2 whose end points are y_1 and y_2 , and define another subarc with the same end points:

$$C_0 = \{\phi(t, x_0); t_1 < t < t_2\}.$$

Let U_1 and U_2 be the interiors of the simple closed curves $C_0 \sqcup C_1$ and $C_0 \sqcup C_2$, respectively. Applying Lemma 2.2 to $C_0 \sqcup C_1$ and $C_0 \sqcup C_2$, we have a continuous map $i : [-1, 1] \rightarrow U$ satisfying $i(0) = x_0$, $i(-1) \in U_1$, $i(1) \in U_2$ and

$$i([-1, 0) \cup (0, 1]) \cap (C_0 \sqcup C) = \emptyset.$$

Define $F : [-1, 1] \rightarrow S$ by

$$F(s) = \mathbf{pr}_2 \circ \varphi^{-1}(i(s))$$

for all $s \in [-1, 1]$, where \mathbf{pr}_2 is the projection to the second component. Notice that $\mathbf{pr}_1 \circ \varphi^{-1}(x) \rightarrow 0$ as $x \rightarrow x_0$, where \mathbf{pr}_1 is the projection to the first component. For sufficiently small $\eta \in (0, 1)$, we have $F(-\eta) \in U_1$, $F(\eta) \in U_2$ and

$$F([-\eta, 0) \cup (0, \eta]) \cap (C_0 \sqcup C) = \emptyset.$$

Define $G : [-1, 1] \rightarrow S$ by $G(s) = F(\eta s)$. Then apply Lemma 2.3 to $G : [-1, 1] \rightarrow G([-1, 1])$ in order to have an injective continuous map $j : [-1, 1] \rightarrow G([-1, 1])$ such that $j(-1) = G(-1)$ and $j(1) = G(1)$. Since $G([-1, 1]) \subset U_1 \sqcup C_0 \sqcup U_2$ and $j([-1, 1])$ is connected, there exists $t_0 \in (-1, 1)$ such that $j(t_0) \in C_0$. Since C_0 is a connected subset of an orbit and $U \subset \text{Im}\varphi$, we have $C_0 \cap S = \{x_0\}$. Thus $j(t_0) = x_0$ is obtained. Let $I = j([-1, 1])$ and define $\psi : (-\tau_0, \tau_0) \times I \rightarrow \text{Im}\varphi$ by

$$\psi(t, x) = \phi(t, x).$$

Then, by the choice of S , ψ is injective, and the Invariance of Domain Theorem shows that ψ is a homeomorphism onto its image. Finally, x_0 is in the interior of $\text{Im}\psi$, since x_0 is in an open set $\psi((-\tau_0, \tau_0) \times j((-1, 1)))$ of $\text{Im}\psi$. \square


Now we introduce some notations in order to give the next proposition. Let p be a singularity of ϕ , and let U be a neighborhood of p homeomorphic to an open disk in \mathbb{R}^2 . Let $C \subset U$ be a simple closed curve surrounding p and let U_C be the interior of C . For $x \in C$, we say that $\text{Orb}^+(x)$ (resp. $\text{Orb}^-(x)$) is a *positive* (resp. *negative*) *base orbit from* x if

$$\phi(t, x) \in U_C, \quad t > 0 \text{ (resp. } t < 0)$$

and

$$\phi(t, x) \rightarrow p, \quad t \rightarrow \infty \text{ (resp. } t \rightarrow -\infty).$$

Let c_i be a positive or negative base orbit from $x_i \in C$ for $i = 1, 2$. Open subset S of U_C is called *sector* determined by c_1 and c_2 if the boundary of S consists of c_1 , c_2 , p and a subarc C_{12} from x_1 to x_2 . A sector S determined by c_1 and c_2 is called *elliptic* if $\text{Orb}(x_1) = \text{Orb}(x_2) \subset \bar{S}$, and *hyperbolic* if S is not elliptic and $C_{12} \cup S$ contains neither positive nor negative base orbit. A sector S determined by c_1 and



(a) Elliptic sector.

(b) Hyperbolic sector.

(c) Parabolic sector.

Figure 2. Three types of sectors.

c_2 is called *positive* (resp. *negative*) *parabolic* sector if c_1 and c_2 are positive (resp. negative) base orbits and \bar{S} contains no negative (resp. positive) base orbit. As argued in Proposition 3.2 below, we may easily see that these three types of sectors do not overlap.

The following proposition is the Local Sector Classification Theorem [5, Lemmas 8.2 and 8.3]. This was proved in the C^1 setting, but the proof is still valid for topological flows.

Proposition 2.5 ([5]). *Let p be a singularity of ϕ , and let D be a closed neighborhood of p homeomorphic to a closed disk in \mathbb{R}^2 . If D contains no α -limit point or ω -limit point except p , then D has a dense subset consisting of finite number of elliptic, parabolic and hyperbolic sectors associated with ∂D .*

The three lemmas below will be used for the proof of Theorem 1.2. The following lemma has been already well-known (see [11] for instance).

Lemma 2.6 (Conley's Fundamental Theorem of Dynamical Systems). *There exists a continuous function $L_0 : M \rightarrow \mathbb{R}$ satisfying the following properties:*

- (1) *For any $x \notin \Omega(\phi)$,*

$$\mathbb{R} \ni t \mapsto L_0(\phi(t, x)) \in \mathbb{R}$$

is strictly increasing.

- (2) *For $x, y \in \Omega(\phi)$, x and y are in the same critical element iff $L_0(x) = L_0(y)$.*

For $\Lambda \in \text{Crit}(\phi)$, define

$$\begin{aligned} W^s(\Lambda) &= \{x \in M; \phi(t, x) \rightarrow \Lambda, \quad t \rightarrow \infty\}, \\ W^u(\Lambda) &= \{x \in M; \phi(t, x) \rightarrow \Lambda, \quad t \rightarrow -\infty\}. \end{aligned}$$

For $\Lambda_1, \Lambda_2 \in \text{Crit}(\phi)$, we write $\Lambda_1 \rightarrow \Lambda_2$ if

$$W^u(\Lambda_1) \cap W^s(\Lambda_2) \setminus \Omega(\phi) \neq \emptyset.$$

The proof of the following lemma is similar to that of [9, Lemma 2.4.6], which is omitted here.

Lemma 2.7. *If $\Omega(\phi)$ consists of finitely many critical elements, then $\phi \in \text{OrientSh}(M)$ has the no-cycles condition.*

The following lemma is the flows version of the Birkhoff Constant Theorem [9, Theorem 2.2.5]. The proof of this lemma is similar to [9, Theorem 2.2.5].

Lemma 2.8 ([9]). *Let U be an open set including $\Omega(\phi)$. Then there exist $R > 0$ and $d > 0$ such that if $x \in M$ satisfies*

$$\phi(t, x) \notin U, \quad t \in (0, \ell),$$

then $\ell < R$.

3. CLASSIFICATION OF SINGULARITIES

In this section, we prove Theorem 1.1, which classifies topological flows on a neighborhood of each $p \in \text{Sing}(\phi)$ with a neighborhood D satisfying the condition of Theorem 1.1.

Denote by $\text{Sing}_s(\phi)$ the set of all asymptotically stable singularities of ϕ , and by $\text{Sing}_u(\phi)$ that of all asymptotically unstable ones of ϕ . Denote by $\text{Sing}_4(\phi)$ the set of all singularities of ϕ with property (c) of Theorem 1.1, and by $\text{Sing}_2(\phi)$ that of those with property (d) of Theorem 1.1. Take $p \in \text{Sing}(\phi)$ and its neighborhood D satisfying the condition of Theorem 1.1.

Lemma 3.1. *For all $x \in D \setminus \{p\}$, we have $\text{Orb}(x) \not\subset D$. In particular, there is no elliptic sectors associated with ∂D .*

Proof. To get a contradiction, suppose there exists a point $x_0 \in D \setminus \{p\}$ such that $\text{Orb}(x_0) \subset D$. Then, by hypothesis, we have $\omega(x_0) = \alpha(x_0) = \{p\}$. Denote by U the open region in the interior of the simple closed curve $\text{Orb}(x_0) \cup \{p\}$. Since $U \subset D$, we have

$$\text{Orb}(x) \subset U \subset D \tag{3.1}$$

for all $x \in U$. Let x_1 be a point in U and take $\varepsilon_1 > 0$ so small that $B(2\varepsilon_1, x_1)$ is contained in U . Then (3.1) implies $\omega(x_1) = \alpha(x_1) = \{p\}$. For sufficiently small $d > 0$, all $\xi \in \text{Ps}(d)$ are ε_1 -shadowed. There exists $T_d > 0$ such that

$$\phi(T_d + t, x_1), \quad \phi(-T_d + t, x_1) \in B(d/2, p), \quad t \in [0, 1].$$

Consider a d -pseudotrajectory $\xi : \mathbb{R} \rightarrow U$ such that

$$\xi(t) = \phi(t - 2T_d n, x_1), \quad t \in [(2n - 1)T_d, (2n + 1)T_d]$$

for all $n \in \mathbb{Z}$. Then there exist $x_2 \in B(\varepsilon_1, x_1)$ and $h \in \text{Rep}$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t), \phi(h(t), x_2)) < \varepsilon_1, \quad t \in \mathbb{R}.$$

From the choice of x_1 and this inequality, we have

$$\begin{aligned} \text{dist}(p, \phi(h(2T_d n), x_2)) &\geq \text{dist}(p, \xi(2T_d n)) - \text{dist}(\xi(2T_d n), \phi(h(2T_d n), x_2)) \\ &\geq 2\varepsilon_1 - \varepsilon_1 = \varepsilon_1 \end{aligned}$$

for all $n \in \mathbb{Z}$. However, it follows from (3.1) that $\omega(x_2) = \alpha(x_2) = \{p\}$. This is a contradiction. \square

Claim 1. *If $p \notin \text{Sing}_s(\phi)$ (resp. $p \notin \text{Sing}_u(\phi)$), then there exists $x \in \partial D$ such that $\text{Orb}^-(x) \subset D$ (resp. $\text{Orb}^+(x) \subset D$) and*

$$\phi(t, x) \rightarrow p, \quad t \rightarrow -\infty, \quad (\text{resp. } \phi(t, x) \rightarrow p, \quad t \rightarrow \infty).$$

Proof. Assume $p \notin \text{Sing}_s(\phi)$. Then we can take $\varepsilon > 0$ such that $B(\varepsilon, p) \subset D$ and for any $n \in \mathbb{Z}_{>0}$, there exists $z_n \in B(1/n, p)$ satisfying

$$\text{Orb}^+(z_n) \not\subset B(\varepsilon, p)$$

or $\omega(z_n) \neq \{p\}$. Notice that if $\text{Orb}^+(z_n) \subset B(\varepsilon, p)$, then $\omega(z_n) = \{p\}$ by the hypothesis on D . Therefore, for all $n \geq 1$,

$$\text{Orb}^+(z_n) \not\subset B(\varepsilon, p).$$

Let

$$t_n = \inf\{t \geq 0; \phi(t, z_n) \in \partial B(\varepsilon, p)\}$$

and $w_n = \phi(t_n, z_n) \in \partial B(\varepsilon, p)$. Taking a subsequence $\{w_{n_k}\}_{k \geq 1}$ if necessary, we may assume that w_{n_k} converges to some $w \in \partial B(\varepsilon, p)$. We have $\text{Orb}^-(w) \subset D$ since $t_{n_k} \rightarrow \infty$ as $k \rightarrow \infty$. Thus by the hypothesis on D again,

$$\phi(t, w) \rightarrow p, \quad t \rightarrow -\infty.$$

Moreover, by Lemma 3.1 we have $\text{Orb}^+(w) \not\subset D$. Thus we may take

$$t_0 = \inf\{t > 0; \phi(t, w) \notin D\}$$

and define $x = \phi(t_0, w)$. Then it is easy to see that x satisfies the required condition. The proof for $p \notin \text{Sing}_u(\phi)$ is similar reversing the time direction. \square

Proposition 3.2. *If $p \notin \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)$, then p does not have any parabolic sector associated with ∂D .*

Proof. Assume that p has a positive parabolic sector S determined by $\text{Orb}^+(x_1)$ and $\text{Orb}^+(x_2)$. Let $C_{1,2}$ be the subarc of ∂D from x_1 to x_2 such that S is the subset of D with boundary consisting of

$$p, \text{Orb}^+(x_1), \text{Orb}^+(x_2) \text{ and } C_{1,2}.$$

Using Proposition 2.4, we can take an embedding $i : [-1, 1] \rightarrow D$ with $i(0) = \phi(1, x_1)$ such that an embedded closed interval $i([-1, 1])$ satisfies the condition of Proposition 2.4 with $x_0 = i(0)$. In addition, we may assume that $i((0, 1]) \subset S$. Then

$$\text{Orb}^+(i(s)) \subset S, \quad s \in (0, s_0] \tag{3.2}$$

for some small $s_0 \in (0, 1]$. In fact, suppose by contradiction that there exists a sequence $\{s_n\}_{n \geq 1}$ in $(0, 1]$ such that $s_n \rightarrow 0$ as $n \rightarrow \infty$ and

$$\text{Orb}^+(i(s_n)) \not\subset S.$$

Then for all $n \geq 1$, there exists $t_n > 0$ such that

$$t_n = \inf\{t > 0; \phi(t, i(s_n)) \in C_{1,2}\}.$$

Taking a subsequence if necessary, we may assume $\phi(t_n, i(s_n)) \rightarrow y_0$ for some $y_0 \in C_{1,2}$. Since $t_n \rightarrow \infty$, by continuity, $\phi(t, y_0) \in \overline{S}$ for all $t < 0$, contradicting that S is parabolic. Thus (3.2) holds. Making s_0 smaller if necessary, we may assume that $\text{Orb}^+(i(s_0)) \cap i([0, s_0]) = \{i(s_0)\}$. Denote by S_0 the region in the interior of the closed curve consisting of

$$i([0, s_0]), p, \text{Orb}^+(i(0)) \text{ and } \text{Orb}^+(i(s_0)),$$

where note that $\omega(i(s_0)) = \{p\}$ by (3.2). Let $x_0 = \phi(1, i(s_0/2)) \in S_0$. Take $\varepsilon_1 > 0$ satisfying the following properties:

$$B(\varepsilon_1, x_0) \subset S_0 \tag{3.3}$$

and

$$\text{dist}(\overline{S_0}, \partial D) > \varepsilon_1. \quad (3.4)$$

Since $p \notin \text{Sing}_s(\phi)$, using Claim 1, we have $w \in \partial D$ with $\text{Orb}^-(w) \subset D$ and

$$\phi(-t, w) \rightarrow p, \quad t \rightarrow \infty.$$

Our hypothesis $\phi \in \text{OrientSh}(M)$ implies that there exists $d > 0$ such that every d -pseudotrajectory is ε_1 -oriented shadowed. Take $T > 0$ so large that the function $\xi : \mathbb{R} \rightarrow M$ defined by

$$\xi(t) = \begin{cases} \phi(t, x_0), & t \leq T, \\ \phi(t - 2T, w), & t > T \end{cases}$$

is a d -pseudotrajectory. Then there exist $h \in \text{Rep}$ with $h(0) = 0$ and $z_0 \in S$ such that

$$\text{dist}(\xi(t), \phi(h(t), z_0)) < \varepsilon_1, \quad t \in \mathbb{R}.$$

From this inequality, we have $\text{dist}(x_0, z_0) < \varepsilon_1$. Thus, by (3.2) and (3.3), $\phi(h(t), z_0) \in S_0$ for all $t \geq 0$. We also have

$$\text{dist}(w, \phi(h(2T), z_0)) < \varepsilon_1.$$

This contradicts (3.4) and proves this proposition when p has a positive parabolic sector. The proof for the case where p has a negative parabolic sector is similar. \square

Assume that $p \notin \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)$. By Proposition 2.5, there is an open dense subset of D consisting of finite number of elliptic, hyperbolic and parabolic sectors. Then, combining Lemma 3.1 and Proposition 3.2, we can observe that D has an open dense subset consisting of even number of hyperbolic sectors.

Lemma 3.3. *Let S be a hyperbolic sector of p determined by $\text{Orb}^+(x_1)$ and $\text{Orb}^-(x_2)$ for some $x_1, x_2 \in \partial D$. Let $x'_1 \in \text{Orb}^+(x_1) \cup \{p\} \setminus \{x_1\}$ and $x'_2 \in \text{Orb}^-(x_2)$. For any neighborhood U_2 of x'_2 , there exists a neighborhood U_1 of x'_1 such that for any $x \in U_1 \cap S$,*

$$\phi(t, x) \in S, \quad t \in [0, t_x]$$

and

$$\phi(t_x, x) \in U_2$$

for some $t_x > 0$.

Proof. Since S is a hyperbolic sector, for all $x \in S$,

$$\text{Orb}^+(x) \not\subset S. \quad (3.5)$$

In fact, S has no positive base orbit, so if (3.5) is false then we would have $\text{Orb}^-(x) \subset S$ (note that $\text{Orb}^+(x) \subset S$ shows that $\omega(x) = \{p\}$). This contradicts Lemma 3.1.

Suppose to the contrary that there exists a sequence $\{y_n\}_{n \geq 1}$ in S such that

$$y_n \rightarrow x'_1, \quad n \rightarrow \infty$$

and for all $n \geq 1$ and $s > 0$, either

$$\phi([0, s], y_n) \not\subset S \quad (3.6)$$

or

$$\phi(s, y_n) \notin U_2 \quad (3.7)$$

hold. By (3.5), there exists $t_n = \inf\{t > 0; \phi(t, y_n) \in \partial S\} < \infty$. If $\phi(s, y_n) \in U_2$ for some $s \in [0, t_n]$, then (3.6) and (3.7) do not hold for s . Thus we get

$$\phi([0, t_n], y_n) \cap U_2 = \emptyset \quad (3.8)$$

for all $n \geq 1$. Taking a subsequence if necessary, we may assume that $\phi(t_n, y_n) \rightarrow y_0$ for some $y_0 \in \partial D$, where $\phi(t, y_0) \in \bar{S}$ for all $t \leq 0$ since $t_n \rightarrow \infty$. Combine this and the fact that S is a hyperbolic sector to have $y_0 = x_2$. This contradicts (3.8). \square

Proposition 3.4. *If $p \notin \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)$, then the number of hyperbolic sectors is either 2 or 4.*

Proof. Suppose to the contrary that there exist more than four hyperbolic sectors. Let S_1 and S_2 be adjacent sectors associated with ∂D , where S_i is determined by $\text{Orb}^+(x_0)$ and $\text{Orb}^-(y_i)$, $i = 1, 2$ for some $x_0, y_1, y_2 \in \partial D$. Since there are more than four hyperbolic sectors, we have $y_3 \in \partial D \setminus \{y_1, y_2\}$ with $\text{Orb}^-(y_3) \subset D$. Let $\varepsilon_1 > 0$ be such that

$$\varepsilon_1 < \text{dist}(\{\phi(-1, y_1), \phi(-1, y_2)\}, \text{Orb}^+(x_0) \cup \{p\} \cup \text{Orb}^-(y_3)) \quad (3.9)$$

and

$$\varepsilon_1 < \text{dist}(y_3, \overline{S_1} \cup \overline{S_2}). \quad (3.10)$$

Using Lemma 3.3 with respect to S_1 and S_2 , we can choose $\varepsilon_2 \in (0, \varepsilon_1)$ so small that if $x \in B(\varepsilon_2, \phi(1, x_0)) \setminus \text{Orb}^+(x_0)$, then there exists $t_0 > 0$ with

$$\phi(t, x) \in S_1 \cup S_2, \quad t \in [0, t_0]$$

and

$$\phi(t_0, x) \in B(\varepsilon_1, y_1) \cup B(\varepsilon_1, y_2).$$

Let $d > 0$ be such that every $\xi \in \text{Ps}(d)$ is ε_2 -oriented shadowed. The function $\xi : \mathbb{R} \rightarrow M$ defined by

$$\xi(t) = \begin{cases} \phi(1+t, x_0), & t \leq T_d, \\ \phi(t-2T_d, y_3), & t > T_d \end{cases}$$

is a d -pseudotrajectory for some $T_d > 0$ and hence ξ is ε_2 -shadowed. Take $h \in \text{Rep}$ with $h(0) = 0$ and $z_0 \in M$ satisfying

$$\text{dist}(\xi(t), \phi(h(t), z_0)) < \varepsilon_2, \quad t \in \mathbb{R}. \quad (3.11)$$

Then we have $z_0 \in B(\varepsilon_2, \phi(1, x_0))$.

Now let us consider two possible cases.

Case 1: $z_0 \in B(\varepsilon_2, \phi(1, x_0)) \setminus \text{Orb}^+(x_0)$.

By the choice of ε_2 , there exists $t_0 > 0$ such that

$$\phi(t, z_0) \in S_1 \cup S_2, \quad t \in [0, t_0]$$

and $\phi(t_0, z_0) \in B(\varepsilon_1, y_1) \cup B(\varepsilon_1, y_2)$. Let $t_1 = \min\{h^{-1}(t_0), 2T_d\}$. Then

$$\phi(h(t_1), z_0) \in \overline{S_1} \cup \overline{S_2}$$

and

$$\xi(t_1) \in \text{Orb}^+(x_0) \cup \{p\} \cup \text{Orb}^-(y_3).$$

In addition, we have either

$$\xi(t_1) = y_3 \text{ when } t_1 = 2T_d$$

or

$$\phi(h(t_1), z_0) \in B(\varepsilon_1, y_1) \cup B(\varepsilon_1, y_2) \text{ when } t_1 = h^{-1}(t_0).$$

By these facts together with (3.9) and (3.10),

$$\text{dist}(\xi(t_1), \phi(h(t_1), z_0)) > \varepsilon_1.$$

This contradicts (3.11) with $\varepsilon_2 < \varepsilon_1$.

Case 2: $z_0 \in B(\varepsilon_2, \phi(1, x_0)) \cap \text{Orb}^+(x_0)$.

Since $\phi(h(2T_d), z_0) \in \overline{S_1} \cup \overline{S_2}$, (3.10) implies

$$\text{dist}(y_3, \phi(h(2T_d), z_0)) > \varepsilon_1.$$

This contradicts (3.11) with $\varepsilon_2 < \varepsilon_1$ again. \square

Now combining Proposition 3.2 with Proposition 3.4, we obtain Theorem 1.1.

4. REDUCTION OF THEOREM 1.2

In this section, we reduce Theorem 1.2 to a proposition which is easier to show, and introduce some notations and lemmas. These notations and lemmas plays an important role in the rest of this paper. Assume that $\phi \in \text{OrientSh}(M)$ has the nonwandering set consisting of finite number of critical elements.

In order to prove Theorem 1.2, it is enough to show that for every $\varepsilon_0 > 0$, there exists $\delta > 0$ such that for all δ -pseudotrajectory ξ we have

$$\text{dist}(\xi(t), \phi(h(t), x)) < 2\varepsilon_0, \quad t \in \mathbb{R}$$

for some $x \in M$ and $h \in \text{Rep}(2\varepsilon_0)$. We may assume that $\varepsilon_0 \in (0, 4/5)$ satisfies

$$3\varepsilon_0 < \min\{\text{dist}(\Lambda_1, \Lambda_2); \Lambda_1, \Lambda_2 \in \text{Crit}(\phi), \Lambda_1 \neq \Lambda_2\}. \quad (4.1)$$

This choice of ε_0 will be used in the following sections. Set

$$K = M \setminus \bigcup_{x \in \text{Sing}(\phi)} B(\varepsilon_0/2, x).$$

Then we can take $T_0 > 0$ such that

$$\phi(t, x) \neq x, \quad t \in [0, 2T_0]$$

for all $x \in K$ because $K \cap \text{Sing}(\phi) = \emptyset$ and K is compact. Define

$$\tilde{K} = \bigcap_{t \in [-2T_0, 2T_0]} \phi(t, K).$$

Making $T_0 > 0$ smaller if necessary, we may assume that

$$B(\varepsilon_0, \Lambda) \subset \tilde{K} \quad (4.2)$$

for every closed orbit Λ . Denote by Pt_{T_0} the set of all functions $\xi : \mathbb{R} \rightarrow M$ satisfying

$$\xi(t + nT_0) = \phi(t, \xi(nT_0))$$

for all $t \in [0, T_0)$ and $n \in \mathbb{Z}$. For $d > 0$, let

$$\text{Pt}_{T_0}(d) = \{\xi \in \text{Pt}_{T_0}; \text{dist}(\phi(T_0, \xi(nT_0)), \xi((n+1)T_0)) < d, \quad n \in \mathbb{Z}\}.$$

The proof of Theorem 1.2 is reduced to showing the following proposition:

Proposition 4.1. *For $\varepsilon_0 > 0$ given by (4.1), there exists $d > 0$ such that if $\xi \in \text{Pt}_{T_0}(d)$ then*

$$\text{dist}(\xi(t), \phi(h(t), x)) < \varepsilon_0, \quad t \in \mathbb{R}$$

for some $x \in M$ and $h \in \text{Rep}(\varepsilon_0)$.

Let us show how Theorem 1.2 follows from Proposition 4.1. It is easy to see that there exists $\delta > 0$ such that if ξ is a δ -pseudotrajectory then

$$\text{dist}(\xi(t), \xi'(t)) < \varepsilon_0, \quad t \in \mathbb{R}$$

for some $\xi' \in \text{Pt}_{T_0}(d)$. Then, by Proposition 4.1, we have

$$\text{dist}(\xi'(t), \phi(h(t), x)) < \varepsilon_0, \quad t \in \mathbb{R}$$

for some $x \in M$ and $h \in \text{Rep}(\varepsilon_0)$. Therefore

$$\begin{aligned} \text{dist}(\xi(t), \phi(h(t), x)) &\leq \text{dist}(\xi(t), \xi'(t)) + \text{dist}(\xi'(t), \phi(h(t), x)) \\ &< \varepsilon_0 + \varepsilon_0 = 2\varepsilon_0, \quad t \in \mathbb{R}, \end{aligned}$$

proving (4.1) from which Theorem 1.2 follows.

The two lemmas below will be needed in the proof of Proposition 4.1, which involve the above choice of ε_0 . The following lemma is essentially [8, Proposition 2.1] where \tilde{K} was defined by given $T_0 \in (0, 1)$ and a compact set K , while we here fixed T_0 depending on ε_0 given above. In what follows, we write $\text{Pt}_{T_0} = \text{Pt}$ and $\text{Pt}_{T_0}(d) = \text{Pt}(d)$.

Lemma 4.2 ([8]). *For every $\varepsilon > 0$, there exists $\varepsilon' > 0$ satisfying the following property:*

If $y \in M$ satisfies

$$\text{dist}(\xi(t + t_0), \phi(g(t), y)) < \varepsilon', \quad t \in [0, T_1]$$

and

$$\xi(t + t_0), \phi(g(t), y) \in \tilde{K}, \quad t \in [0, T_1]$$

for some $\xi \in \text{Pt}$, $t_0 \in \mathbb{R}$, $T_1 \geq T_0$ and $g \in \text{Rep}$ with $g(0) = 0$, then there exists $\tilde{g} \in \text{Rep}(\varepsilon)$ with $\tilde{g}(0) = g(0)$ and $\tilde{g}(T_1) = g(T_1)$ such that

$$\text{dist}(\xi(t + t_0), \phi(\tilde{g}(t), y)) < \varepsilon, \quad t \in [0, T_1].$$

The following lemma is the closed orbits version of [8, Lemma 5.1].

Lemma 4.3 ([8]). *Let Λ be an asymptotically stable (resp. backward asymptotically stable) critical element of ϕ . Then there exist $d > 0$ and a neighborhood U of Λ such that if $\xi \in \text{Pt}(d)$ satisfies $\xi(t_0) \in U$ for some $t_0 \in \mathbb{R}$, then*

$$\xi(t) \in B(\varepsilon_0/2, \Lambda), \quad t \geq t_0 \text{ (resp. } t \leq t_0).$$

The proof of this lemma is similar to that of [8, Lemma 5.1], which is omitted here. In fact, the proof of [8, Lemma 5.1] is also valid for closed orbits just by replacing singularities by them.

5. NEIGHBORHOODS OF CLOSED ORBITS

In this section, we construct neighborhoods of closed orbits in order to investigate the behavior of pseudotrajectories.

Lemma 5.1. *Let Λ be a closed orbit of ϕ . Then Λ is asymptotically stable or backward asymptotically stable.*

Proof. Let ε_0 be the constant given by (4.1). Let $\varphi : [-1, 1] \rightarrow B(\varepsilon_0, \Lambda)$ be an embedding such that an embedded closed interval $\varphi([-1, 1])$ satisfies the condition of Proposition 2.4 with $x_0 = \varphi(0)$. For some $\delta_0 > 0$, we can take a Poincaré map $P : \varphi([- \delta_0, \delta_0]) \rightarrow \varphi([-1, 1])$. Define $\tilde{P} : [- \delta_0, \delta_0] \rightarrow [-1, 1]$ by $\tilde{P} = \varphi^{-1} \circ P \circ \varphi$. It is enough to show that 0 is attracting or repelling for \tilde{P} . Notice that P is a homeomorphism onto its image, and thus \tilde{P} is increasing or decreasing.

Let us consider the following two cases separately.

Case 1: \tilde{P} is increasing.

It follows from (4.1) that $\tilde{P}(t) \neq t$ hold for all $t \neq 0$. Since \tilde{P} is continuous, we may assume that $\tilde{P}(a) > a$ for all $a \in (0, \delta_0)$. The other case where $\tilde{P}(a) < a$ for all $a \in (0, \delta_0)$ is proved similarly. If $\tilde{P}(a) < a$ for all $a \in (-\delta_0, 0)$, then 0 is repelling for \tilde{P} .

To prove by contradiction, let us assume that $\tilde{P}(a) > a$ for some $a \in (-\delta_0, 0)$ which implies $\tilde{P}(a) > a$ for all $(-\delta_0, 0)$ because of (4.1). Let $p_0 = \varphi(-\delta_0/2)$ and let $p_1 = \varphi(\delta_0/2)$. Take $\varepsilon > 0$ so small that the closure of $B(\varepsilon, p_0)$ is disjoint from Λ . Then let Γ be the closure of $\text{Orb}^+(B(\varepsilon, p_0))$. Making ε smaller if necessary, we may assume that

$$B(\varepsilon, p_1) \cap \Gamma = \emptyset. \quad (5.1)$$

Since $\phi \in \text{OrientSh}(M)$, there exists $d > 0$ such that every d -pseudotrajectory is ε -oriented shadowed. Take $\tau_0, \tau_1 > 0$ so large that the function $\xi : \mathbb{R} \rightarrow M$ defined by

$$\xi(t) = \begin{cases} \phi(t + \tau_0, p_0), & t \leq 0, \\ \phi(t - \tau_1, p_1), & t > 0 \end{cases}$$

satisfies $\xi \in \text{Ps}(d)$. By the choice of d , ξ is ε -shadowed. Thus there exist $h \in \text{Rep}$ with $h(-\tau_0) = 0$ and $x_0 \in M$ such that

$$\text{dist}(\xi(t), \phi(h(t), x_0)) < \varepsilon, \quad t \in \mathbb{R}. \quad (5.2)$$

Then $\text{dist}(p_0, x_0) < \varepsilon$, which together with the definition of Γ implies that

$$\phi(h(t), x_0) \in \Gamma$$

for all $t \geq -\tau_0$. In particular, $\phi(h(\tau_1), x_0) \in \Gamma$ and then (5.1) contradicts (5.2) with $t = \tau_1$.

Case 2: \tilde{P} is decreasing.

Take $\delta_1 \in (0, \delta_0)$ satisfying $\tilde{P}((-\delta_1, \delta_1)) \subset (-\delta_0, \delta_0)$. Then it follows from (4.1) that $\tilde{P}^2(t) \neq t$ for all $t \neq 0$. Now assume that $\tilde{P}^2(a) > a$ for all $a \in (0, \delta_1)$. The proof for the other case where $\tilde{P}^2(a) < a$ for all $a \in (0, \delta_1)$ is similar. Since \tilde{P} is continuous and $\tilde{P}^2(\delta_1/2) > \delta_1/2$, there exists $a_0 \in (\tilde{P}(\delta_1/2), 0)$ such that $\tilde{P}(a_0) > \delta_1/2$. Then $(a_0, \delta_1/2)$ satisfies $\tilde{P}^{-1}(a_0, \delta_1/2) \subset (a_0, \delta_1/2)$, thus by (4.1), $(a_0, \delta_1/2)$ is a repelling neighborhood of 0 for \tilde{P} . \square

Denote by $\text{Clos}_s(\phi)$ the set of all asymptotically stable closed orbits, and denote by $\text{Clos}_u(\phi)$ that of all backward asymptotically stable ones.

Claim 2. *Let Λ be a closed orbit of ϕ and let ε_0 be the constant given by (4.1). There exists $r_0 \in (0, \varepsilon_0/2)$ such that if $x, y \in \Lambda$ satisfy $\text{dist}(x, y) < r_0$, then*

$$\text{dist}(\phi(t, x), \phi(t, y)) < \frac{\varepsilon_0}{8}, \quad t \in \mathbb{R}.$$

Proof. Choose $\varepsilon > 0$ so small that

$$\text{dist}(\phi(t, x), x) < \frac{\varepsilon_0}{8} \tag{5.3}$$

for all $t \in (-\varepsilon, \varepsilon)$ and $x \in \Lambda$. To prove by contradiction, suppose that there exist sequences $\{x_n\}_{n \geq 1}$ and $\{y_n\}_{n \geq 1}$ in Λ such that

$$\text{dist}(x_n, y_n) < \frac{1}{n} \tag{5.4}$$

and

$$\text{dist}(\phi(t_n, x_n), \phi(t_n, y_n)) \geq \frac{\varepsilon_0}{8}$$

for some $\{t_n\}_{n \geq 1}$. Then by (5.3),

$$\phi(t_n, y_n) \notin \phi((-\varepsilon, \varepsilon), \phi(t_n, x_n)),$$

implying

$$y_n \notin \phi((-\varepsilon, \varepsilon), x_n) \tag{5.5}$$

for $n \geq 1$. Taking a subsequence if necessary, we may assume that $x_n \rightarrow x_0$ for some $x_0 \in \Lambda$. Then, by (5.5), $y_n \notin \phi((-\varepsilon/2, \varepsilon/2), x_0)$ for sufficiently large n . This contradicts (5.4) for large n . \square

The following proposition is the main result of this section, whose proof is given after the proof of Lemma 5.3 below. The proposition shows that when Λ is asymptotically stable it has a neighborhood in which not only any forward pseudotrajectory with sufficiently small jump stays in the neighborhood, but also it is ε_0 -standard shadowed by the forward orbit of every nearby point.

Proposition 5.2. *Let $\Lambda \in \text{Clos}_s(\phi)$ (resp. $\Lambda \in \text{Clos}_u(\phi)$). Then there exist a neighborhood $U_\Lambda \subset B(\varepsilon_0, \Lambda)$ of Λ and $d_\Lambda, r_\Lambda > 0$ such that if $\xi \in \text{Pt}(d_\Lambda)$ satisfies $\xi(t_0) \in U_\Lambda$ for some $t_0 \in \mathbb{R}$ then the following properties hold:*

- (a) $\xi(t) \in B(\varepsilon_0/2, \Lambda)$ for all $t \geq t_0$ (resp. $t \leq t_0$).
- (b) For all $x \in B(r_\Lambda, \xi(t_0))$, there exists $h \in \text{Rep}(\varepsilon_0)$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t), \phi(h(t - t_0), x)) < \varepsilon_0, \quad t \geq t_0 \text{ (resp. } t \leq t_0).$$

In order to prove Proposition 5.2, we need the following lemma. This lemma claims that every forward orbit near a asymptotically stable closed orbit is standard shadowed by a sufficiently close forward orbit.

Lemma 5.3. *Let $\Lambda \in \text{Clos}_s(\phi)$ (resp. $\Lambda \in \text{Clos}_u(\phi)$). Then there exists $r > 0$ such that if $x_0 \in \Lambda$ and $x_1 \in B(r, x_0)$, then*

$$\text{dist}(\phi(t, x_0), \phi(g(t), x_1)) \leq \frac{\varepsilon_0}{4}, \quad t \geq 0 \text{ (resp. } t \leq 0)$$

for some $g \in \text{Rep}(\varepsilon_0/4)$ with $h(0) = 0$.

Proof. Without loss of generality, we may assume $\Lambda \in \text{Clos}_s(\phi)$, since the case for $\Lambda \in \text{Clos}_u(\phi)$ is similar. Let $r_0 \in (0, \varepsilon_0/2)$ be the constant given by Claim 2. Apply Lemma 4.2 with ε replaced by $\min\{\varepsilon_0/8, r_0/2\}$ and let $\varepsilon'_0 \in (0, r_0)$ be the constant given by the proposition for $\varepsilon = \min\{\varepsilon_0/8, r_0/2\}$. We may take $r \in (0, r_0/2)$ such that every $\xi \in \text{Pt}(r)$ is ε'_0 -oriented shadowed and

$$\text{Orb}^+(x) \subset B(\varepsilon_0/8, \Lambda)$$

for all $x \in B(r, \Lambda)$ since Λ is asymptotically stable. For every $x_0 \in \Lambda$ and $x_1 \in B(r, x_0)$, set

$$\xi(t) = \begin{cases} \phi(t, x_0), & t < 0, \\ \phi(t, x_1), & t \geq 0. \end{cases}$$

Since $\xi \in \text{Pt}(r)$, there exist $h \in \text{Rep}$ with $h(0) = 0$ and $x'_1 \in M$ such that

$$\text{dist}(\xi(t), \phi(h(t), x'_1)) < \varepsilon'_0, \quad t \in \mathbb{R}.$$

By the choice of r ,

$$\xi(t) \in B(\varepsilon_0/8, \Lambda), \quad t \in \mathbb{R}. \quad (5.6)$$

This fact and (4.2) enable us to apply Lemma 4.2. Thus we have

$$\text{dist}(\xi(t), \phi(g(t), x'_1)) < \min\{\varepsilon_0/8, r_0/2\}, \quad t \in \mathbb{R} \quad (5.7)$$

for some $g \in \text{Rep}(\min\{\varepsilon_0/8, r_0/2\})$ with $g(0) = 0$. This inequality and (5.6) imply that

$$\text{Orb}(x'_1) \subset B(\varepsilon_0/4, \Lambda).$$

Thus, by (4.1), we have $\alpha(x'_1) = \omega(x'_1) = \Lambda$. Then Lemma 2.7 shows $x'_1 \in \Lambda$. Since (5.7) implies

$$\text{dist}(x'_1, x_0) \leq \text{dist}(x'_1, x_1) + \text{dist}(x_1, x_0) < r_0/2 + r < r_0,$$

we have, $x'_1 \in \Lambda \cap B(r_0, x_0)$. Therefore, by Claim 2 and (5.7),

$$\begin{aligned} \text{dist}(\phi(t, x_0), \phi(g^{-1}(t), x_1)) &\leq \text{dist}(\phi(t, x_0), \phi(t, x'_1)) \\ &\quad + \text{dist}(\phi(t, x'_1), \phi(g^{-1}(t), x_1)) \\ &= \frac{\varepsilon_0}{8} + \text{dist}(\phi(g(g^{-1}(t)), x'_1), \xi(g^{-1}(t))) \\ &\leq \frac{\varepsilon_0}{8} + \frac{\varepsilon_0}{8} = \frac{\varepsilon_0}{4} \end{aligned}$$

for all $t \geq 0$. Combining this and $g^{-1} \in \text{Rep}(\varepsilon_0/4)$, we finish the proof. \square

Now, let us prove Proposition 5.2.

Proof of Proposition 5.2. Let us prove the proposition when $\Lambda \in \text{Clos}_s(\phi)$, since the case where $\Lambda \in \text{Clos}_u(\phi)$ can be proved similarly. Let $r_0, r > 0$ be the positive numbers given by Claim 2 and Lemma 5.3, respectively. Apply Lemma 4.2 with ε replaced by $\min\{\varepsilon_0/4, r/2\}$ and let $\varepsilon' \in (0, r_0)$ be the constant given by the proposition for $\varepsilon = \min\{\varepsilon_0/4, r/2\}$. Let d_Λ be such that every $\xi \in \text{Pt}(d_\Lambda)$ is ε' -oriented shadowed. Set $r_\Lambda = r/2$ and

$$U_\Lambda = B(r/2, \Lambda). \quad (5.8)$$

Lemma 4.3 implies that, by making d_Λ and U_Λ smaller if necessary, we may assume that if $\xi \in \text{Pt}(d_\Lambda)$ satisfies $\xi(t_0) \in U_\Lambda$ for some $t_0 \in \mathbb{R}$, then

$$\xi(t) \in B(\varepsilon_0/2, \Lambda) \quad (5.9)$$

for all $t \geq t_0$, proving (a). As for the proof of (b), as in the hypothesis, suppose that $\xi \in \text{Pt}(d_\Lambda)$ satisfies $\xi(t_0) \in U_\Lambda$ for some $t_0 \in \mathbb{R}$. Let $x_1 \in B(r_\Lambda, \xi(t_0))$. By (5.8), we can choose $x_0 \in \Lambda$ satisfying $\xi(t_0) \in B(r/2, x_0)$. On the other hand, by the choice of d_Λ , there exist x_2 and $h \in \text{Rep}$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t + t_0), \phi(h(t), x_2)) < \varepsilon', \quad t \in \mathbb{R}.$$

Since $\varepsilon' < \varepsilon_0/2$, together with (5.9) and (4.2) we may use Lemma 4.2 again. Then we have

$$\text{dist}(\xi(t + t_0), \phi(g(t), x_2)) < \min\{\varepsilon_0/4, r/2\}, \quad t \in \mathbb{R} \quad (5.10)$$

for some $g \in \text{Rep}(\min\{\varepsilon_0/4, r/2\})$ with $g(0) = 0$. Now apply Lemma 5.3 to x_1 and x_2 respectively, to have $h_1, h_2 \in \text{Rep}(\varepsilon_0/4)$ with $h_1(0) = h_2(0) = 0$ such that

$$\text{dist}(\phi(t, x_0), \phi(h_1(t), x_1)) < \varepsilon_0/4, \quad t \geq 0,$$

$$\text{dist}(\phi(t, x_0), \phi(h_2(t), x_2)) < \varepsilon_0/4, \quad t \geq 0.$$

Then these inequalities and (5.10) show that for all $t \geq 0$,

$$\begin{aligned} & \text{dist}(\xi(t + t_0), \phi(h_1 \circ h_2^{-1} \circ g(t), x_1)) \\ & \leq \text{dist}(\xi(t + t_0), \phi(g(t), x_2)) \\ & \quad + \text{dist}(\phi(g(t), x_2), \phi(h_2^{-1} \circ g(t), x_0)) \\ & \quad + \text{dist}(\phi(h_2^{-1} \circ g(t), x_0), \phi(h_1 \circ h_2^{-1} \circ g(t), x_1)) \\ & < \frac{\varepsilon_0}{4} + \frac{\varepsilon_0}{4} + \frac{\varepsilon_0}{4} = \frac{3\varepsilon_0}{4}. \end{aligned}$$

When $t \neq s$, we have

$$\begin{aligned} \left| \frac{h_1 \circ h_2^{-1} \circ g(t) - h_1 \circ h_2^{-1} \circ g(s)}{t - s} \right| &= \left| \frac{h_1 \circ h_2^{-1} \circ g(t) - h_1 \circ h_2^{-1} \circ g(s)}{h_2^{-1} \circ g(t) - h_2^{-1} \circ g(s)} \right| \\ & \quad \cdot \left| \frac{h_2^{-1} \circ g(t) - h_2^{-1} \circ g(s)}{g(t) - g(s)} \right| \\ & \quad \cdot \left| \frac{g(t) - g(s)}{t - s} \right|. \end{aligned}$$

From this and $\varepsilon_0 < 4/5$ it follows that for all $t \in \mathbb{R}$,

$$\begin{aligned} 1 - \varepsilon_0 &< \left(1 - \frac{\varepsilon_0}{4}\right) \left(1 + \frac{\varepsilon_0}{4}\right)^{-1} \left(1 - \frac{\varepsilon_0}{4}\right) \\ &< \left| \frac{h_2 \circ h_1^{-1} \circ g(t) - h_2 \circ h_1^{-1} \circ g(s)}{t - s} \right| \\ &< \left(1 + \frac{\varepsilon_0}{4}\right) \left(1 - \frac{\varepsilon_0}{4}\right)^{-1} \left(1 + \frac{\varepsilon_0}{4}\right) < 1 + \varepsilon_0. \end{aligned}$$

Thus we have $h_1 \circ h_2^{-1} \circ g \in \text{Rep}(\varepsilon_0)$, finishing the proof of Proposition 5.2.

6. NEIGHBORHOODS OF SINGULARITIES

In this section, we construct open sets near singularities in order to investigate the behavior of pseudotrajectories (see Figure 3).

Let $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$. Take a neighborhood $D \subset B(\varepsilon_0/2, p)$ of p homeomorphic to a closed disk in \mathbb{R}^2 . Then ∂D is a simple closed curve surrounding

p . If $p \in \text{Sing}_4(\phi)$, take $x_1, x_2, y_1, y_2 \in D$ such that the four hyperbolic sectors associated with ∂D of p is determined by basic orbits:

$$\text{Orb}^+(\phi(-1, x_i)) \text{ and } \text{Orb}^-(\phi(1, y_j))$$

with $i, j \in \{1, 2\}$. Similarly, if $p \in \text{Sing}_2(\phi)$, take $x_1, y_1 \in D$ such that the two hyperbolic sectors associated with ∂D of p is determined by basic orbits

$$\text{Orb}^+(\phi(-1, x_1)) \text{ and } \text{Orb}^-(\phi(1, y_1)).$$

Define

$$G^s(p) = \begin{cases} \{x_1, x_2\}, & p \in \text{Sing}_4(\phi), \\ \{x_1\}, & p \in \text{Sing}_2(\phi) \end{cases}$$

and

$$G^u(p) = \begin{cases} \{y_1, y_2\}, & p \in \text{Sing}_4(\phi), \\ \{y_1\}, & p \in \text{Sing}_2(\phi). \end{cases}$$

The following proposition claims that the α -limit set set of a point in $W^s(p) \setminus \{p\}$




Figure 3

or the ω -limit set set of a point in $W^u(p) \setminus \{p\}$ for $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ displays a simple dynamics. More precisely, the following proposition shows that there is no saddle connection.

Proposition 6.1. *Let $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$. If $z_0 \in W^s(p) \setminus \{p\}$, (resp. $z_0 \in W^u(p) \setminus \{p\}$) then $\alpha(z_0)$ (resp. $\omega(z_0)$) is a backward asymptotically stable (resp. asymptotically stable) critical element.*

Proof. Suppose that $z_0 \in W^s(p) \setminus \{p\}$. Let Λ_0 be the critical element containing $\alpha(z_0)$. Then we have that Λ_0 is not an asymptotically stable critical element. Arguing by contradiction we assume that $\Lambda_0 \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ (see Theorem 1.1 and Lemma 5.1). Let

$$d_1 \in (0, \inf\{L_0(\Lambda_0) - L_0(\Lambda); \Lambda \longrightarrow \Lambda_0\}).$$

Notice that $L_0(\Lambda_0) > L_0(\Lambda)$ for all Λ with $\Lambda \longrightarrow \Lambda_0$ (see Lemma 2.6). Making d_1 smaller if necessary, we may take a neighborhood $N^u(p)$ of $G^u(p)$ such that $L_0(y) > L_0(p) + d_1/2$ for all $y \in N^u(p)$. Then by Proposition 2.4, we may take an embedded closed interval Σ_0 satisfying the condition in Proposition 2.4 with $x_0 = \phi(1, z_0)$. By Lemma 3.3 we may assume that

$$\text{Orb}^+(x) \cap N^u(p) \neq \emptyset \quad (6.1)$$

for all $x \in \Sigma_0 \setminus \{\phi(1, z_0)\}$. Take $\varepsilon_1 > 0$ so small that

$$|L_0(x) - L_0(y)| \leq d_1/2 \quad (6.2)$$

for all $x, y \in M$ with $\text{dist}(x, y) < \varepsilon_1$ and

$$B(\varepsilon_1, z_0) \subset \{\phi(t, x); -2 < t < 0, x \in \Sigma_0\}.$$

Since $\phi \in \text{OrientSh}(M)$, there exists $d > 0$ such that every $\xi \in \text{Pt}(d)$ is ε_1 -oriented shadowed. Take $z_1 \in W^s(\Lambda_0) \setminus \Lambda_0$ and define

$$\xi(t) = \begin{cases} \phi(t, z_0), & t \geq -T_d, \\ \phi(t + 2T_0, z_1), & t < -T_d \end{cases}$$

for some $T_d > 0$. Then $\xi \in \text{Ps}(d)$ for sufficiently large T_d . Thus there exist $h \in \text{Rep}$ with $h(0) = 0$ and $w_0 \in M$ such that

$$\text{dist}(\xi(t), \phi(h(t), w_0)) < \varepsilon_1, \quad t \in \mathbb{R}.$$

In particular, (6.2) implies

$$|L_0(\xi(t)) - L_0(\phi(h(t), w_0))| \leq d_1/2, \quad t \in \mathbb{R}. \quad (6.3)$$

Since $w_0 \in B(\varepsilon_1, z_0)$, the choice of ε_1 implies that $\phi(s, w_0) \in \Sigma_0$ for some $s \in [0, 2]$. Now suppose that $\phi(s, w_0) \neq z_0$. Then, by (6.1), we may take $t' \geq 0$ satisfying $\phi(h(t'), w_0) \in N^u(p)$, which shows

$$L_0(\phi(h(t'), w_0)) > L_0(p) + d_1/2$$

by the choice of $N^u(p)$. On the other hand, it is obvious that

$$L_0(p) = \sup\{L_0(\phi(t, z_0)); t \in \mathbb{R}\} > L_0(\phi(t', z_0)) = L_0(\xi(t')),$$

contradicting inequality (6.3) with $t = t'$. This contradiction gives $\phi(s, w_0) = z_0$. From this and (6.3), it follows that

$$\begin{aligned} L_0(\Lambda_0) &= \inf\{L_0(\phi(t, z_0)); t \in \mathbb{R}\} \\ &= \inf\{L_0(\phi(t, w_0)); t \in \mathbb{R}\} \\ &\leq \inf\{L_0(\phi(t, z_1)); t \in \mathbb{R}\} + d_1/2, \end{aligned}$$

which contradicts $z_1 \in W^s(\Lambda_0) \setminus \Lambda_0$ and the choice of d_1 .

The proof for the ω -limit set is similar. \square

The following proposition is the singularities version of Proposition 5.2. However the proof is much simpler because $h \in \text{Rep}(\varepsilon_0)$ can be the identity.

Proposition 6.2. *Let $p \in \text{Sing}_s(\phi)$ (resp. $p \in \text{Sing}_u(\phi)$). Then there exist a neighborhood $U_p \subset B(\varepsilon_0, p)$ of p and $d_p, r_p > 0$ such that if $\xi \in \text{Pt}(d_p)$ satisfies $\xi(t_0) \in U_p$ for some $t_0 \in \mathbb{R}$ then the following properties hold:*

- (a) $\xi(t) \in B(\varepsilon_0/2, p)$ for all $t \geq t_0$ (resp. $t \leq t_0$).
- (b) For any $x \in B(r_p, \xi(t_0))$,

$$\text{dist}(\xi(t), \phi(t - t_0, x)) < \varepsilon_0, \quad t \geq t_0 \text{ (resp. } t \leq t_0).$$

Proof. It is enough to consider $p \in \text{Sing}_s(\phi)$ since the case for $p \in \text{Sing}_u(\phi)$ is similar. It follows from Lemma 4.3 that there exist $d > 0$ and a neighborhood V_p of p satisfying the following property:

- If $\xi \in \text{Ps}(d)$ satisfies $\xi(0) \in V_p$, then $\xi(t) \in B(\varepsilon_0/2, p)$ for $t \geq 0$.

Take $r_p > 0$ so small that $B(2r_p, p) \subset V_p$. Set $U_p = B(r_p, p)$ and let $d_p > 0$ be such that $\text{Pt}(d_p) \subset \text{Ps}(d)$. By hypothesis, we have $\xi \in \text{Pt}(d_p)$ with $\xi(t_0) \in U_p$ for some $t_0 \in \mathbb{R}$. Since $U_p \subset V_p$, the choice of V_p implies property (a). As for property (b), by the choice of r_p , we have $B(r_p, \xi(t_0)) \subset B(2r_p, p) \subset V_p$. Thus, using the above property twice (notice that $t \mapsto \phi(t, x)$ is a d -pseudotrajectory), we have

$$\text{dist}(\phi(t - t_0, x), \xi(t)) < \text{dist}(\phi(t - t_0, x), p) + \text{dist}(p, \xi(t)) < \frac{\varepsilon_0}{2} + \frac{\varepsilon_0}{2} = \varepsilon_0,$$

proving property (b). \square

For every $\Lambda \in \text{Clos}_s(\phi) \cup \text{Clos}_u(\phi) \cup \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)$, fix a neighborhood U_Λ of Λ given by Proposition 5.2 or Proposition 6.2. In these propositions, we also gave constants $d_\Lambda, r_\Lambda > 0$. Then let

$$d_0 = \min\{d_\Lambda; \Lambda \in \text{Clos}_s(\phi) \cup \text{Clos}_u(\phi) \cup \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)\} \quad (6.7)$$

and

$$r_0 = \min\{r_\Lambda; \Lambda \in \text{Clos}_s(\phi) \cup \text{Clos}_u(\phi) \cup \text{Sing}_s(\phi) \cup \text{Sing}_u(\phi)\}. \quad (6.8)$$

The following lemma claims that any backward (resp. forward) pseudotrajectory with a point in the fundamental domain of $W^s(p)$ (resp. $W^u(p)$) with $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ can be ε_0 -shadowed along the backward (resp. forward) pseudotrajectory.

Lemma 6.3. *Let $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$. There exist $d_p, \tau_p > 0$ satisfying the following property:*

- If $\xi \in \text{Pt}(d_p)$ and $x_0 \in M$ satisfies $\xi(t_0), x_0 \in B(\tau_p, x)$ for some $t_0 \in \mathbb{R}$ and $x \in G^s(p)$ (resp. $x \in G^u(p)$), then there exists $h \in \text{Rep}(\varepsilon_0)$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t + t_0), \phi(h(t), x_0)) < \varepsilon_0, \quad t \leq 0 \text{ (resp. } t \geq 0). \quad (6.9)$$

Proof. Let $x \in G^s(p)$. Then Proposition 6.1 shows that $\alpha(x) \in \text{Clos}_u(\phi) \cup \text{Sing}_u(\phi)$. Take $T > 0$ satisfying $\phi(-T, x) \in U_{\alpha(x)}$. Then choose $d_x \in (0, d_0)$ and $\tau_x > 0$ such that if $\xi \in \text{Pt}(d_x)$ with $\xi(t_0) \in B(\tau_x, x)$ and $y \in B(\tau_x, x)$ then

$$\xi(t_0 - T) \in U_{\alpha(x)} \quad (6.10)$$

and

$$\text{dist}(\xi(t_0 - t), \phi(-t, y)) < \min\{\varepsilon_0, r_{\alpha(x)}\}, \quad t \in [0, T]. \quad (6.11)$$

Suppose that $\xi \in \text{Pt}(d_x)$ and x_0 satisfies $\xi(t_0), x_0 \in B(\tau_x, x)$. By (6.10) and (6.11), Propositions 5.2 and 6.2 show that there exists $g \in \text{Rep}(\varepsilon_0)$ with $g(0) = 0$ such that

$$\text{dist}(\xi(t + t_0 - T), \phi(g(t), \phi(-T, x_0))) < \varepsilon_0, \quad t \leq 0. \quad (6.12)$$

Define $h \in \text{Rep}(\varepsilon_0)$ by

$$h(t) = \begin{cases} g(t + T) - T, & t \leq -T, \\ t, & t \geq -T. \end{cases}$$

Then, by (6.11),

$$\text{dist}(\xi(t + t_0), \phi(h(t), x_0)) < \varepsilon_0, \quad t \in [-T, 0].$$

This inequality and (6.12) yield (6.9). Similarly, we can choose d_y and τ_y for all $y \in G^s(p) \cup G^u(p)$. Taking the minimum over the finite set $G^s(p) \cup G^u(p)$, we complete the proof of the lemma. \square

For $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ and $x \in G^s(p) \cup G^u(p)$, fix an embedded closed interval

$$\Sigma_x \subset B(\tau_p, x) \quad (6.13)$$

satisfying the condition of Proposition 2.4 with $x_0 = x$. When $x \in G^s(p)$ and $y \in G^u(p)$, there exists a point of Σ_x which has the forward orbit intersecting Σ_y at an arbitrary large time.

Lemma 6.4. *Let $x_i, y_j \in D$ be the points given in the beginning of this section. Then there exists $T_p > 0$ such that for all $i, j \in \{1, 2\}$ and $T \geq T_p$,*

$$\phi(t, x) \in D, \quad t \in [0, T]$$

and $\phi(T, x) \in \Sigma_{y_j}$ for some $x \in \Sigma_{x_i}$.

Proof. Set $i, j \in \{1, 2\}$ and let S be the hyperbolic sector of p determined by $\phi(-1, x_i)$ and $\phi(1, y_j)$. By Lemma 3.3, making Σ_{x_i} smaller if necessary, we may assume that for any $x \in \Sigma_{x_i} \cap S$, there exists $t_x > 0$ such that

$$\phi(t, x) \in S, \quad t \in [0, t_x]$$

and $\phi(t_x, x) \in \Sigma_{y_j}$. Since the map $\Sigma_{x_i} \cap S \ni x \mapsto t_x \in (0, \infty)$ is continuous, and $t_x \rightarrow \infty$ when $x \rightarrow x_i$, this lemma has been proved. \square

The following lemma claims that some dynamical behavior around $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ is preserved for pseudotrajectories.

Lemma 6.5. *Let $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$ and let $d_p, \tau_p > 0$ and $T_p > 0$ be the constants given by Lemmas 6.3 and 6.4, respectively. Then there exist a neighborhood $U_p \subset D$ of p and $d'_p \in (0, d_p)$ satisfying the following properties:*

- (a) *Assume that $\xi \in \text{Pt}(d'_p)$ satisfies $\xi(t_0) \in U_p$ and $\xi(t_0 + t_1) \notin B(\varepsilon_0/2, p)$ for some $t_1 > 0$. Then there exists $t_2 > T_p/2$ such that*

$$\xi(t_0 + t_2) \in B(\tau_p, G^u(p)) \quad (6.14)$$

and

$$\xi(t_0 + t) \in B(\varepsilon_0/2, p), \quad t \in [0, t_2]. \quad (6.15)$$

- (b) Assume that $\xi \in \text{Pt}(d'_p)$ satisfies $\xi(t_0) \in U_p$ and $\xi(t_0 + t_1) \notin B(\varepsilon_0/2, p)$ for some $t_1 < 0$. Then there exists $t_2 < -T_p/2$ such that

$$\xi(t_0 + t_2) \in B(\tau_p, G^s(p))$$

and

$$\xi(t_0 + t) \in B(\varepsilon_0/2, p), \quad t \in [t_2, 0].$$

Proof. We prove this lemma for $p \in \text{Sing}_4(\phi)$; for the proof for $p \in \text{Sing}_2(\phi)$ is essentially the same. Let $\tau_p > 0$ be the constant given in Lemma 6.3. Taking $\tau'_p \in (0, \tau_p)$, we may assume that

$$\text{dist}(p, G^s(p) \cup G^u(p)) > \tau'_p.$$

If a neighborhood U of p and $d > 0$ are small enough,

$$\xi(t + t_0) \in B(\varepsilon_0/2, p) \setminus \overline{B(\tau'_p, G^s(p) \cup G^u(p))}, \quad t \in [-T_p/2, T_p/2] \quad (6.16)$$

for all $\xi \in \text{Pt}(d)$ with $\xi(t_0) \in U$ for some $t_0 \in \mathbb{R}$.

Choose $r_1 \in (0, \tau'_p/2)$ with $D \subset B(\varepsilon_0/2 - r_1, p)$. By Lemma 3.3, there exists $r_2 \in (0, r_1)$ such that if $x \in B(2r_2, p)$ and $\text{Orb}^+(x) \not\subset D$ then

$$\phi(t, x) \in D, \quad t \in [0, t_x] \quad (6.17)$$

and

$$\phi(t_x, x) \in B(r_1, G^u(p)) \quad (6.18)$$

for some $t_x > 0$. In addition, making r_2 smaller if necessary, we may assume that $B(r_2, p) \subset U$. Take $d'_p > 0$ such that every $\xi \in \text{Pt}(d'_p)$ is r_2 -oriented shadowed. Set $U_p = B(r_2, p)$.

As in the hypothesis of property (a), suppose that $\xi \in \text{Pt}(d'_p)$ satisfies $\xi(t_0) \in U_p = B(r_2, p)$ and $\xi(t_0 + t_1) \notin B(\varepsilon_0/2, p)$ for some $t_0 \in \mathbb{R}$ and $t_1 > 0$. Then there exist $x_0 \in B(2r_2, p)$ and $h \in \text{Rep}$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t_0 + t), \phi(h(t), x_0)) < r_2, \quad t \in \mathbb{R}, \quad (6.19)$$

where $x_0 \in B(2r_2, p)$ follows from

$$\text{dist}(x_0, p) \leq \text{dist}(x_0, \xi(t_0)) + \text{dist}(\xi(t_0), p) < r_2 + r_2.$$

Moreover, inequality

$$\text{dist}(p, \phi(h(t_1), x_0)) \geq \text{dist}(p, \xi(t_0 + t_1)) - \text{dist}(\xi(t_0 + t_1), \phi(h(t_1), x_0)) > \varepsilon_0/2 - r_2$$

shows that $\phi(h(t_1), x_0) \notin D$. Thus (6.17) and (6.18) with $x = x_0$ hold for some $t_{x_0} > 0$. Then (6.17), (6.19) and the choice of r_1, r_2 with $D \subset B(\varepsilon_0/2 - r_1, p) \subset B(\varepsilon_0/2 - r_2, p)$ imply that (6.15) for $t_2 = h^{-1}(t_{x_0})$. Now (6.19) and (6.18) give

$$\begin{aligned} \text{dist}(\xi(t_0 + h^{-1}(t_{x_0})), G^u(p)) &\leq \text{dist}(\xi(t_0 + h^{-1}(t_{x_0})), \phi(t_{x_0}, x_0)) \\ &\quad + \text{dist}(\phi(t_{x_0}, x_0), G^u(p)) \\ &< r_2 + r_1 < 2r_1 < \tau'_p < \tau_p. \end{aligned}$$

This and (6.16) show that $h^{-1}(t_{x_0}) > T_p/2$. In addition, this implies (6.14) with $t_2 = h^{-1}(t_{x_0})$. The proof for property (b) is similar. \square

7. PROOF OF PROPOSITION 4.1

In this section, let us finish the proof of Theorem 1.2 by showing Proposition 4.1 given in Section 4. We need Lemma 4.2 and Proposition 5.2 and two propositions and three lemmas in Section 6. To apply Lemma 4.2 later with $\varepsilon = \varepsilon_0$, let $\varepsilon'_0 > 0$ be the constant corresponding to ε' in the proposition. Let d_0 and d'_p be constants given by (6.7) and Lemma 6.5 respectively, and take

$$d_1 < \min\{d_0, d'_p; p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)\}$$

so that every $\xi \in \text{Pt}(d_1)$ is ε'_0 -oriented shadowed. Let

$$U = \bigcup_{\Lambda \in \text{Crit}(\phi)} U_\Lambda,$$

where U_Λ is the neighborhood of Λ given in Propositions 5.2, 6.2 and Lemma 6.5.

It is easy to see that the following claim follows from Lemma 2.8.

Claim 3. *There exist $R_0 > 0$ and $d_2 \in (0, d_1)$ such that if $\xi \in \text{Pt}(d_2)$ satisfies*

$$\xi(t) \notin U, \quad t \in (0, \ell),$$

then $\ell < R_0$.

In this claim, making $d_2 > 0$ smaller if necessary, we may assume that for all $\xi \in \text{Pt}(d_2)$ and $t_0 \in \mathbb{R}$,

$$\text{dist}(\xi(t_0 + t), \phi(t, \xi(t_0))) < \min\{\varepsilon_0, r_0\}, \quad t \in [0, R_0], \quad (7.1)$$

where ε_0 and r_0 were given in (4.1) and (6.8).

The hypothesis of the following proposition is used for classifying $\xi \in \text{Pt}(d_2)$ into several cases.

Proposition 7.1. *Suppose that $\xi(t_0) \in U_p$ for some $p \in \text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)$. If*

$$\sup\{t \in \mathbb{R}; \xi(t) \in U_p\} = \infty \text{ (resp. } \inf\{t \in \mathbb{R}; \xi(t) \in U_p\} = -\infty),$$

then $\xi(t) \in B(\varepsilon_0/2, p)$ for all $t \geq t_0$ (resp. $t \leq t_0$).

Proof. Let us consider the former case. Suppose to the contrary that there exists $t_1 > t_0$ with $\xi(t_1) \notin B(\varepsilon_0/2, p)$. Then Lemma 6.5 implies that $\xi(t_0 + t_2) \in B(\tau_p, G^u(p))$ for some $t_2 > T_p/2$. Then Proposition 6.1 for asymptotically stable critical elements and Lemma 6.3 for $t \geq 0$ show that $\xi(t) \in B(3\varepsilon_0/2, \Lambda)$ for some asymptotically stable critical element Λ and sufficiently large t , contradicting our hypothesis. The proof for the latter case is similar. \square

Let $\xi \in \text{Pt}(d_0)$ and define

$$S(\xi) = \{\Lambda \in \text{Crit}(\phi); \xi(t) \in U_\Lambda \text{ for some } t \in \mathbb{R}\}.$$

If $\xi(t_0) \in U_\Lambda$ for some $t_0 \in \mathbb{R}$ and some asymptotically stable critical element Λ , then by Propositions 6.2 and 5.2, we have $\xi(t) \in B(\varepsilon_0, \Lambda)$ for all $t \geq t_0$. Therefore $S(\xi)$ contains at most one asymptotically stable critical element. Similarly, $S(\xi)$ contains at most one backward asymptotically stable critical element. Thus it is enough to consider the following cases:

- (C1) $S(\xi) = \{\Lambda\}$ with $\Lambda \in \text{Sing}_s(\phi) \cup \text{Clos}_s(\phi)$.
- (C2) $S(\xi) = \{\Lambda\}$ with $\Lambda \in \text{Sing}_u(\phi) \cup \text{Clos}_u(\phi)$.
- (C3) $S(\xi) = \{\Lambda_1, \Lambda_2\}$ with $\Lambda_1 \in \text{Sing}_s(\phi) \cup \text{Clos}_s(\phi)$ and $\Lambda_2 \in \text{Sing}_u(\phi) \cup \text{Clos}_u(\phi)$.

(C4) There exists $p \in (\text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)) \cap S(\xi)$ such that

$$\inf\{t \in \mathbb{R}; \xi(t) \in U_p\} = -\infty \text{ and } \sup\{t \in \mathbb{R}; \xi(t) \in U_p\} = \infty.$$

(C5) There exists $p \in (\text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)) \cap S(\xi)$ such that

$$\inf\{t \in \mathbb{R}; \xi(t) \in U_p\} > -\infty \text{ and } \sup\{t \in \mathbb{R}; \xi(t) \in U_p\} = \infty.$$

(C6) There exists $p \in (\text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)) \cap S(\xi)$ such that

$$\inf\{t \in \mathbb{R}; \xi(t) \in U_p\} = -\infty \text{ and } \sup\{t \in \mathbb{R}; \xi(t) \in U_p\} < \infty.$$

(C7) There exists $p \in (\text{Sing}_4(\phi) \cup \text{Sing}_2(\phi)) \cap S(\xi)$ such that

$$\inf\{t \in \mathbb{R}; \xi(t) \in U_p\} > -\infty \text{ and } \sup\{t \in \mathbb{R}; \xi(t) \in U_p\} < \infty.$$

For the proof of Theorem 1.2, it suffices to show that any $\xi \in \text{Pt}(d_2)$ is ε_0 -standard shadowed in each case. Now let us prove it for the seven cases.

Proofs for cases from (C1) to (C7).

(C1): From Claim 3, it follows that $\{t \in \mathbb{R}; \xi(t) \in U_\Lambda\}$ does not have any lower bound. Then, by Propositions 6.2 and 5.2,

$$\xi(t) \in B(\varepsilon_0/2, \Lambda), \quad t \in \mathbb{R}. \quad (7.2)$$

In fact, otherwise there is $t' \in \mathbb{R}$ such that $\xi(t') \notin B(\varepsilon_0/2, \Lambda)$, but since there is no lower bounds, we can find some $t < t'$ such that $\xi(t) \in U_\Lambda$, which implies $\xi(t') \in B(\varepsilon_0/2, \Lambda)$ by these propositions, contradicting the above. From (7.2) it follows that ξ is ε_0 -shadowed by $\phi(t, \Lambda)$ for all $t \in \mathbb{R}$ when $\Lambda \in \text{Sing}_s(\phi)$. On the other hand, when $\Lambda \in \text{Clos}_s(\phi)$, by the choice of d_2 , there exist $x_0 \in M$ and $h \in \text{Rep}$ with $h(0) = 0$ such that

$$\text{dist}(\xi(t), \phi(h(t), x_0)) < \varepsilon'_0, \quad t \in \mathbb{R}.$$

Then, by (7.2) and (4.2), we can apply Lemma 4.2 to $[nT_0, (n+1)T_0]$, $n \in \mathbb{Z}$, (thinking of $[0, T_0]$ in the proposition) in order to show that ξ is ε_0 -standard shadowed.

(C2): The proof for this case is similar to that for case **(C1)**.

(C3): By Claim 3, we may take $t_1, t_2 \in \mathbb{R}$ such that

$$\xi(t_1) \in U_{\Lambda_1}, \quad \xi(t_2) \in U_{\Lambda_2}$$

and $t_1 - t_2 \in (0, R_0)$. By (7.1), we have

$$\text{dist}(\xi(t_2 + t), \phi(t, \xi(t_2))) < \min\{\varepsilon_0, r_0\}, \quad t \in [0, R_0], \quad (7.3)$$

implying

$$\text{dist}(\xi(t_1), \phi(t_1 - t_2, \xi(t_2))) < r_0.$$

Thus, Propositions 6.2 and 5.2 (b) show that there exist $h_1, h_2 \in \text{Rep}(\varepsilon_0)$ with $h_1(0) = h_2(0) = 0$ such that

$$\text{dist}(\xi(t), \phi(h_1(t - t_1), \phi(t_1 - t_2, \xi(t_2)))) < \varepsilon_0, \quad t \geq t_1 \quad (7.4)$$

and

$$\text{dist}(\xi(t), \phi(h_2(t - t_2), \xi(t_2))) < \varepsilon_0, \quad t \leq t_2. \quad (7.5)$$

Define $h \in \text{Rep}(\varepsilon_0)$ with $h(t_2) = 0$ by

$$h(t) = \begin{cases} h_1(t - t_1) + t_1 - t_2, & t \geq t_1, \\ t - t_2, & t \in [t_2, t_1], \\ h_2(t - t_2), & t \leq t_2. \end{cases}$$

Then it follows from (7.3), (7.4) and (7.5) that

$$\text{dist}(\xi(t), \phi(h(t), \xi(t_2))) < \varepsilon_0, \quad t \in \mathbb{R}.$$

(C4): By Proposition 7.1, $\xi(t) \in B(\varepsilon_0/2, p)$ for all $t \in \mathbb{R}$. Therefore ξ is ε_0 -shadowed by $\phi(t, p)$ for $t \in \mathbb{R}$.

(C5): Assume that $\xi(t_0) \in U_p$. By Claim 3 and $\inf\{t \in \mathbb{R}; \xi(t) \in U_p\} > -\infty$, we can find $t_1 < t_0$ with $\xi(t_1) \notin B(\varepsilon_0/2, p)$. Then Lemma 6.5 (b) implies that there exists $t_2 < -T_p/2$ such that

$$\xi(t_0 + t_2) \in B(\tau_p, G^s(p))$$

and

$$\xi(t_0 + t) \in B(\varepsilon_0/2, p), \quad t \in [t_2, 0]. \quad (7.6)$$

Let $x_0 \in G^s(p)$ with $\xi(t_0 + t_2) \in B(\tau_p, x_0)$. Then, by Lemma 6.3, there exists $h_0 \in \text{Rep}(\varepsilon_0)$ with $h_0(0) = 0$ such that

$$\text{dist}(\xi(t + t_0 + t_2), \phi(h_0(t), x_0)) < \varepsilon_0, \quad t \leq 0. \quad (7.7)$$

Define $h \in \text{Rep}(\varepsilon_0)$ with $h(0) = 0$ by

$$h(t) = \begin{cases} h_0(t), & t \leq 0, \\ t, & t \geq 0. \end{cases}$$

Then Proposition 7.1, (7.6) and (7.7) imply

$$\text{dist}(\xi(t + t_0 + t_2), \phi(h(t), x_0)) < \varepsilon_0, \quad t \in \mathbb{R}.$$

(C6): The proof for this case is similar to that for case **(C5)**.

(C7): Assume that $\xi(t_0) \in U_p$ for some $t_0 \in \mathbb{R}$. By Proposition 7.1 and the hypothesis, there exist $t_1 < t_0$ and $s_1 > t_0$ with $\xi(t_1), \xi(s_1) \notin B(\varepsilon_0/2, p)$. Then Lemma 6.5 shows that there are $t_2 < -T_p/2$ and $s_2 > T_p/2$ such that $\xi(t_0 + t_2) \in B(\tau_p, G^s(p))$, $\xi(t_0 + s_2) \in B(\tau_p, G^u(p))$ and

$$\xi(t_0 + t) \in B(\varepsilon_0/2, p), \quad t \in [t_2, s_2]. \quad (7.8)$$

Let $x_0 \in G^s(p)$ and $y_0 \in G^u(p)$ be such that $\xi(t_0 + t_2) \in B(\tau_p, x_0)$ and $\xi(t_0 + s_2) \in B(\tau_p, y_0)$, respectively. By Lemma 6.4, we can find x_1 in Σ_{x_0} satisfying

$$\phi(t, x_1) \in B \subset B(\varepsilon_0/2, p), \quad t \in [0, s_2 - t_2] \quad (7.9)$$

and $\phi(s_2 - t_2, x_1) \in \Sigma_{y_0}$, where Σ_{x_0} and Σ_{y_0} are the embedded closed intervals given by (6.13). Then it follows from Lemma 6.3 that there exist $h_0 \in \text{Rep}(\varepsilon_0)$ with $h_0(0) = 0$ such that

$$\text{dist}(\xi(t + t_0 + t_2), \phi(h_0(t), x_1)) < \varepsilon_0, \quad t \leq 0, \quad (7.10)$$

and $g_0 \in \text{Rep}(\varepsilon_0)$ with $g_0(0) = 0$ such that

$$\text{dist}(\xi(t + t_0 + s_2), \phi(g_0(t), \phi(s_2 - t_2, x_1))) < \varepsilon_0, \quad t \geq 0. \quad (7.11)$$

Define $h \in \text{Rep}(\varepsilon_0)$ with $h(0) = 0$ by

$$h(t) = \begin{cases} h_0(t), & t \leq 0, \\ t, & t \in [0, s_2 - t_2], \\ g_0(t - s_2 + t_2) + s_2 - t_2, & t \geq s_2 - t_2. \end{cases}$$

Thus, by (7.10), (7.11), (7.8) and (7.9), we obtain

$$\text{dist}(\xi(t + t_0 + t_2), \phi(h(t), x_1)) < \varepsilon_0, \quad t \in \mathbb{R},$$

completing the proof.

ACKNOWLEDGEMENTS

The author is grateful to my advisor S. Hayashi for his constructive suggestions and continuous support.

REFERENCES

1. Aranson S. Kh., Belitsky G. R., Zhuzhoma E. V. Introduction to the qualitative theory of dynamical systems on surfaces. Amer. Math. Soc., 1996.
2. Bhatia, N. P., Szegő, G. P. Stability theory of dynamical systems. Springer Science and Business Media, 2002.
3. Cairns, S. S. An elementary proof of the Jordan-Schoenflies theorem. *Proceedings of the American Mathematical Society*, **2.6**, 860-867 (1951).
4. Gutiérrez, C. Smoothing continuous flows on two-manifolds and recurrences. *Ergodic Theory and dynamical systems*, **6.1**, 17-44 (1986).
5. Hartman, P. Ordinary differential equations. Society for Industrial and Applied Mathematics, 2002.
6. Komuro, M. One-parameter flows with the pseudorbit tracing property, *MonatMath.*, **98**, 219-253 (1984).
7. Mai, J., Gu, R. Conditions of C^0 flows on closed surfaces having the pseudo-orbit tracing property. *Sci. China Ser. A-Math.* **40**, 1166-1175 (1997).
8. Murakami, S. Oriented and Standard shadowing Properties for Topological flows, *Tokyo J. Math.*, **46**, (2022).
9. Pilyugin, S., and Sakai, K. Shadowing and hyperbolicity. Vol. 2193. Berlin: Springer, 2017.
10. Pilyugin, S., and Tikhomirov, S. Lipschitz shadowing implies structural stability. *Nonlinearity* **23**(10), 2509-2515 (2010).
11. Robinson, C. Dynamical systems: stability, symbolic dynamics, and chaos. CRC press, 1998.
12. Robinson, C. Stability theorems and hyperbolicity in dynamical systems. *The Rocky Mountain Journal of Mathematics*, **7**(3), 425-437, (1977).
13. Tikhomirov, S. An Example of a Vector Field with the Oriented Shadowing Property, *J. Dyn. control syst.*, **21**, 643-654 (2015).
14. Willard, S. General topology. Courier Corporation, 2012.