

The Scholz conjecture on addition chains is true for $n = 2^m(23) + 7, m \in \mathbb{N}^*$

Amadou TALL¹

Departement de Mathématiques et Informatique
 Université Cheikh Anta Diop de Dakar
 amadou7.tall@ucad.edu.sn

Abstract. The Scholz conjecture on addition chains states that $\ell(2^n - 1) \leq \ell(n) + n - 1$ for all integers n where $\ell(n)$ stands for the minimal length of all addition chains for n . It is proven to hold for infinite sets of integers. In this paper, we will prove that the conjecture still holds for $n = 2^m(23) + 7$. It is the first set of integers given by Thurber [8] to prove that there are an infinity of integers satisfying $\ell(2n) = \ell(n)$. Later on, Thurber [4] give a second set of integers with the same properties ($n = 2^{2m+k+7} + 2^{2m+k+5} + 2^{m+k+4} + 2^{m+k+3} + 2^{m+2} + 2^{m+1} + 1$). We will prove that the conjecture holds for them as well.

Keywords: Addition chain, Scholz-Bauer conjecture, minimal length, factor method

Introduction

Exponentiation is a key operation in mathematics. It is often seen as a number of multiplications. Let x be an integer, finding x^n for a given n seems to be easy. We need to do n multiplications. But what if n is getting very large. It is a very important operation in many areas like cryptography. The best tool know to do fast exponentiation is addition chain.

Definition 1. An addition chain for an integer n is a set $\mathcal{C} = \{a_0 = 1, a_1, a_2, \dots, a_r\}$ such that the last element a_r is n and every element a_k is the sum of two previous elements a_i and a_j . The integer r is the length of the chain and is denoted $\ell(c)$.

One can easily see that there can be many addition chains of different length for the same integer n and finding the minimal length for all addition chains of n (denoted $\ell(n)$) is a NP-complete problem. Just to give an example, on can think that the fastest way to reach $2n$ is by reaching n and then doubling it, which means that $\ell(2n) = \ell(n) + 1$. It has been proven that there are infinitely many integers which don't follow that rule. We have $\ell(382) = \ell(191) = 11$. The most famous conjecture on addition chain is giving an upper bound on the length of addition chains for integers with only "1"s in their binary expansion ($n = 2^a - 1$ for some a).

Conjecture 2. The Scholz conjecture on addition chains states that

$$\ell(2^n - 1) \leq \ell(n) + n - 1, \forall n \in \mathbb{N}$$

Several decades of research and the conjecture has been proved to hold for $n < 5784689$. It has been proven to hold for several sets of integers. We know that if it is true for n and that $\ell(2n) = \ell(n) + 1$, then it is also true for $2n$. We would like to know if it holds when $\ell(2n) = \ell(n)$. An answer for the general case doesn't exist yet.

In this paper, we will investigate the two cases.

1. We will give a very simple proof that the conjecture holds for $2n$ if it is for n and $\ell(2n) = \ell(n) + 1$.
2. We will also prove that it holds for integers $n = 2^m(23) + 7$ which are known to be among the first infinite sets of integers verifying $\ell(2n) = \ell(n)$.

The main result from Thurber [8] is

Theorem 3. Let m be an integer greater than 5. Then

$$\ell(2^m(23) + 7) = m + 8 = \ell(2 \times (2^m(23) + 7))$$

Let us remind the factor method, which is a method to construct a chain for the product nm based on chains for n and m .

Definition 4. Let c_1 and c_2 be addition chains respectively for n_1 and n_2 . Then $c_1 \times c_2$ is an addition chain for $n_1 \times n_2$ of length $\ell(c_1) + \ell(c_2)$ where \times is defined as follows:

if $c_1 = \{a_0, a_1, \dots, a_r\}$ and $c_2 = \{b_0, b_1, \dots, b_l\}$, then

$$c_1 \times c_2 = \{a_0, a_1, \dots, a_r, a_r \times b_1, a_r \times b_2, \dots, a_r \times b_l\}.$$

The length of the new chain is the sum of the length of the chains, meaning that $\ell(mn) \leq \ell(n) + \ell(m)$.

1 Our contribution

Theorem 5. Let n be a positive integer satisfying $\ell(2n) = \ell(n) + 1$. If the Scholz conjecture holds for n , then it also holds for $2n$.

Proof. Let n be a positive integer which satisfies $\ell(2n) = \ell(n) + 1$ and $\ell(2^n - 1) \leq \ell(n) + n - 1$. Let $n_0 = 2n$ be another positive integer, we have

$$2^{n_0} - 1 = (2^n - 1)(2^n + 1),$$

using the factor method, we can deduce a chain for $2^{n_0} - 1$ of length

$$\ell(n) + n - 1 + n + 1 = \ell(n) + 2n = \ell(n_0) + n_0 - 1.$$

Theorem 6. The Scholz conjecture holds for all integers of the form $U_m = 2^m(23) + 7$.

It will be proven by induction on m

Proof. 1. Thanks to the computational results, the Scholz conjecture holds for U_m , $m \in \{1, 2, 3, 4, 5\}$

2. A chain for $X_m + 3$ (with $X_m = 2^m(23)$) of length $m + 7$ can be obtain as follows

$$\mathcal{C}_{X_{\frac{1}{2}}} = \{1, 2, 3, 5, 10, 20, 23, 2 \times 23, \dots, 2^m \times 23, 2^m \times 23 + 3\}$$

3. We know that

$$U_{m+1} = 2^{m+1}(23) + 7$$

so,

$$X_m = U_m - 7 = \frac{U_{m+1} - 1}{2} - 3$$

which leads to

$$\begin{aligned} 2^{U_{m+1}} - 1 &= 2^{2(X_m+3)+1} - 1 \\ &= 2(2^{2(X_m+3)} - 1) + 1 \\ &= 2(2^{X_m+3} - 1)(2^{X_m+3} + 1) + 1 \end{aligned}$$

From the chain of $(2^{X_m+3} - 1)$ and using the factor method, we can then deduce a chain for $2^{U_{m+1}} - 1$ of length

$$\begin{aligned} X_m + 3 + 1 + 1 + 1 + \ell(2^{X_m+3} - 1) &= X_m + 6 + \ell(X_m + 3) + X_m + 3 - 1 \\ &= 2X_m + 8 + m \\ &= (2x_m + 7) + (m + 9) - 1 \\ &= U_{m+1} + \ell(U_{m+1}) - 1 \end{aligned}$$

Two years later, Thurber has also stated that

Theorem 7. For each $m \geq 1$, the set of integers with $v(n) = 7$ and n of the binary form $n = 101 \dots m \dots 11 \dots k \dots 11 \dots m \dots 1$ (where $k \geq 3$) is an infinite class of integers which $\ell(2n) = \ell(n)$.

He prove that $\ell(n) = \lambda(n) + 4$. We will also prove that the Scholz conjecture holds for such n .

Theorem 8. *The Scholz conjecture on addition chain holds for $n = 2^{2m+k+7} + 2^{2m+k+5} + 2^{m+k+4} + 2^{m+k+3} + 2^{m+2} + 2^{m+1} + 1$.*

Proof. Let us rewrite

$$\begin{aligned} n &= 2^{2m+k+7} + 2^{2m+k+5} + 2^{m+k+4} + 2^{m+k+3} + 2^{m+2} + 2^{m+1} + 1 \\ &= 2^{m+k+3}(2^{m+4} + 2 + 2^{m+2} + 1) + (2^{m+2} + 1 + 2^{m+1}) \\ &= 2^{m+k+3}(2(2^{m+3} + 1) + (2^{m+2} + 1)) + (2^{m+2} + 1) + 2^{m+1} \end{aligned}$$

A minimal chain for n is

$$\begin{aligned} \mathcal{J} &= \{1, 2, \dots, 2^{m+1}, 2^{m+2}, 2^{m+2} + 1, 2^{m+3} + 1, 2^{m+1} + (2^{m+2} + 1) = \beta, \\ &\quad 2(2^{m+3} + 1), 2(2^{m+3} + 1) + (2^{m+2} + 1) = \alpha, \\ &\quad 2\alpha, 2^2\alpha, \dots, 2^{m+k+3}\alpha, n = 2^{m+k+3}\alpha + \beta\}. \end{aligned}$$

which is of length

$$m + 2 + 1 + 1 + 1 + 1 + 1 + m + k + 3 + 1 = 2m + k + 11 = \lambda(n) + 4.$$

Using the following rules:

1. If $A = 2B$, then we will get a chain for $2^A - 1 = (2^B - 1)(2^B + 1)$ by adding $B + 1$ steps to a chain for $2^B - 1$
2. If $A = B + 1$, then we will add two steps ($2^A - 1 = 2(2^B - 1) + 1$)
3. $2^{m+3} + 1 = 2^{m+2} + 2^{m+2} + 1$, so

$$2^{2^{m+3}+1} - 1 = 2^{2^{m+2}}(2^{2^{m+2}+1} - 1) + (2^{2^{m+2}} + 1)$$

4. $2^{m+2} + 2^{m+1} + 1 = (2^{m+2} + 1) + 2^{m+1}$, so

$$2^{2^{m+2}+2^{m+1}+1} - 1 = 2^{2^{m+1}}(2^{2^{m+2}+1} - 1) + (2^{2^{m+1}} - 1)$$

We can then construct a chain \mathcal{C} for $2^n - 1$ of length

value	step	additional steps	Comment
1	DBL		
2	DBL		
...	DBL		
2^{m+1}	DBL		
2^{m+2}	DBL	$2^{m+2} + m + 2 - 1$	
$2^{m+2} + 1$	+1	2	
$2^{m+3} + 1$	$+2^{m+2}$	$2^{m+2} + 1$	
$2^{m+2} + 2^{m+1} + 1 = \beta$	2^{m+1}	1	$2^{2^{m+1}}(2^{2^{m+2}+1} - 1) \in \mathcal{C}$
$2(2^{m+3} + 1)$	DBL	$2^{m+3} + 1 + 1$	
$2(2^{m+3} + 1) + (2^{m+2} + 1) = \alpha$	$+2^{m+2} + 1$	$2^{m+2} + 1 + 1$	
2α	DBL	$\alpha + 1$	
$2^2\alpha$	DBL	$2\alpha + 1$	
...	DBL	...	
$2^{m+k+3}\alpha$	DBL	$2^{m+k+3}\alpha + 1$	
$n = 2^{m+k+3}\alpha + \beta$	$+\beta$	$\beta + 1$	
TOTAL	4 small steps		$\ell(n) + n - 1 =$

$$\begin{aligned} \ell &= 2^{m+2} + m + 2 - 1 + 2 + 2^{m+2} + 1 + 2 + 2^{m+3} + \\ &\quad + 1 + 1 + 2^{m+2} + 1 + 1 + (m + k + 3) + \alpha(2^{m+k+3} - 1) + \beta \\ &= 2m + k + 13 + 2^{m+4} + 2^{m+2} + (2^{m+4} + 2^{m+2} + 2 + 1)(2^{m+k+3} - 1) + 2^{m+2} + 2^{m+1} + 1 \\ &= 2m + k + 14 + 2^{m+4} + 2^{m+3} + 2^{m+1} + 2^{2m+k+7} + 2^{2m+k+5}2^{m+k+4} + 2^{m+k+3} - 2^{m+4} - 2^{m+2} - 2 - 1 \\ &= (2m + k + 11) + (2^{2m+k+7} + 2^{2m+k+5}2^{m+k+4} + 2^{m+k+3} + 2^{m+2} + 2^{m+1}) = \ell(n) + n - 1. \end{aligned}$$

Conclusion

We don't know if the Scholz conjecture holds for all integers with $\ell(2n) = \ell(n)$. It must be more difficult to investigate the case where $\ell(2n) < \ell(n)$.

Acknowledgment

The author would like to thank IHES. The work was completed during his visiting fellowship.

References

1. D.E. Knuth, *The Art of Computer Programming*, Vol. 2 (Addison-Wesley, Read. Mass., 1969) 398-422.
2. Tall Amadou, *A generalization of Lucas addition chains*, Bulletin Mathématique de la Roumanie, Tome 55 (103) (1) (2012) 79 - 93.
3. Maurice Mignotte, Amadou Tall, *A note on addition chains*. International Journal of Algebra, Vol. 5, 2011, no. 6, 269 - 274
4. Thurber, Edward G., *The Scholz-Brauer problem on addition chains*, Pacific Journal of Mathematics, V.49 No.1, 1973 p.229-242
5. Schönhage, A. *A Lower Bound for the Length of Addition Chains*, Theoretical Computer Science, V. 1 1975 p. 1-12.
6. A. T. Brauer, *On addition chains*, Bull. Amer. Math. Soc. 45 (1939), 736-739.
7. W. R. Utz, *A note on the Scholz-Brauer problem in addition chains*, Proc. Amer. Math. Soc. 4 (1953), 462-463.
8. Thurber, Edward G., *On additions chains and solutions of $\ell(2n) = \ell(n)$ and $\ell(2^n - 1) = \ell(n) + n - 1$* , Discrete Mathematics 16 (1976) 279-289.
9. Cliff, N., *Calculating optimal addition chains*, Computing 2011, 265-284.