

Landau theory for ferro-paramagnetic phase transition in finitely-strained viscoelastic magnets

Tomáš Roubíček

*Mathematical Institute, Charles University,
Sokolovská 83, CZ-186 75 Praha 8,
email: tomas.roubicek@mff.cuni.cz*

*Institute of Thermomechanics, Czech Academy of Sciences,
Dolejšková 5, CZ-18200 Praha 8, Czech Republic*

Abstract.

The thermodynamic model of visco-elastic deformable magnetic materials at finite strains is formulated in a fully Eulerian way in rates. The Landau theory applies for ferro-to-paramagnetic phase transition, the gradient theory (leading exchange energy) for magnetization with general mechanically dependent coefficient, hysteresis in magnetization evolution by Landau-Lifshitz-Gilbert equation involving objective corotational time derivative of magnetization, and demagnetizing field are considered in the model. The Kelvin-Voigt viscoelastic rheology with a higher-order viscosity (exploiting the concept of multipolar materials) is used, allowing for physically relevant frame-indifferent stored energies and for local invertibility of deformation. The model complies with energy conservation and Clausius-Duhem entropy inequality. Existence and a certain regularity of weak solutions is proved by a Faedo-Galerkin semi-discretization and a suitable regularization.

Keywords: Elastodynamics, ferromagnetics, phase transition, micromagnetics, magnetostriiction, Kelvin-Voigt viscoelasticity, thermal coupling, large strains, multipolar continua, semi-Galerkin discretization, weak solutions.

AMS Subject Classification: 35Q74, 35Q79, 65M60, 74A30, 74F15, 74N30, 80A20.

1 Introduction – deforming magnetic continua

The magnetic materials which are not completely rigid represent interesting, important, and difficult multi-physical concatenation of mere (thermo)continuum mechanics and mere micromagnetism. Beside homogeneous visco-elastic magnets, it may concern elastically rather soft materials filled with magnetic particles, e.g. rocks (which can be considered soft on long time scales) and polymers (i.e. so-called magneto-rheological elastomers or ferrogels), which however needs to involve creep which is not consider in this paper rather for not making the

model too complicated. We will focus ourselves to general finite (also called large) strain mechanics in the Eulerian formulation.

This magneto-mechanical subject has been addressed in [14, Ch.6] or anisothermal but not with explicitly articulated equations [10] and also, in a thermodynamic context, [35, Ch.6]. Even in the purely mechanical isothermal cases, and a-fortiori in anisothermal situations, the visco-elastodynamics at finite strains has been articulated in [2, 3] as a difficult open problem as far as existence of weak solutions concerns. There is a certain agreement that, for analytical reasons, a certain enough strong dissipation mechanism is to be involved to make the dynamical problem parabolic, although some hyperbolic models exist, as mentioned below. The simplest variant is the *Kelvin-Voigt viscoelastic rheology*.

The mentioned *Eulerian approach* is standardly believed to be well fitted with fluids. It is particularly suitable in situations when there is no natural reference configuration or where a reference configuration becomes less and less relevant during long-time evolution, which may however apply also for solids. A formulation of equations in current deforming configuration needs rather velocity/strain than displacement to be involved in the momentum equation. The advantage is an easier possibility to involve interaction with outer spatial fields (here magnetic and gravity) and avoiding the pull-back and push-forward manipulation. On the other hand, there is a necessity to involve convective derivative and transport equations and also evolving the shape of the body is troublesome. In isothermal situations, such model was formulated and analyzed as incompressible in [30, 31] and as compressible in [24, 43]. The mentioned higher gradients that would allow for reasonable analysis can now be involved rather in the dissipative than conservative part, so that their influence manifests only in fast evolutions. In the isothermal situations it was used in quasistatic case in [47] and in dynamical case in [50] when considering the stored energy in the actual configuration, which then gives an energy pressure in the stress tensor. In anisothermal situations, such free-energy pressure would be directly added into stress tensor in a non-integrable way and likely would cause technical difficulties.

The main attributes of the devised model are:

- ⊕ Concept of *hyperelastic materials* (whose conservative-stress response comes from a free energy) combined with the *Kelvin-Voigt viscoelastic rheology* and also evolution of magnetization is driven by this free energy.
- ⊕ Inertial effects in fully compressible context (in particular with varying mass density) are considered.
- ⊕ The rate formulation in terms of velocity and deformation gradient is used while the deformation itself does not explicitly occur.
- ⊕ Magnetic phenomena covered by the model includes: ferro-to-para magnetic phase transition, hysteresis due to the pinning effects, exchange energy depending on deformation gradient (and in particular on compression/expansion), and demagnetizing field.
- ⊕ Mechanical consistency in the sense that *frame indifference* of the free energy (which is in particular *nonconvex* in terms of deformation gradient and in magnetization) and its *singularity* under infinite compression in relation with *local non-interpenetration* as well as objective corotational time derivative for magnetization transport.

- ⊕ Thermodynamic consistency of the thermally coupled system in the sense that the *total energy is conserved* in a closed system, the *Clausius-Duhem entropy inequality* holds, and temperature stays non-negative.
- ⊕ The nonconservative part of the stress in the Kelvin-Voigt model containing a higher-order component reflecting the concept of nonsimple *multipolar media* is exploited.
- ⊕ The model allows for rigorous mathematical analysis as far as existence and certain regularity of energy-conserving weak solutions concerns.

On the other hand, some simplifications are adopted:

- ⊖ Relatively slow evolution is implicitly assumed, which allows for reducing the full Maxwell electromagnetodynamics to magneto-statics.
- ⊖ Electric conductivity (and in particular eddy currents) is not considered.

As far as the non-negativity of temperature, below we will be able to prove only that at least some solutions enjoy this attribute, although there is an intuitive belief that all possible solutions will make it and a hope that more advanced analytical techniques would rigorously prove it.

The main notation used in this paper is summarized in the following table:

\mathbf{v} velocity (in m/s),	$\psi = \psi(\mathbf{F}, \mathbf{m}, \theta)$ referential free energy (in J/m ³ =Pa),
ϱ mass density (in kg/m ³),	$\varphi = \varphi(\mathbf{F}, \mathbf{m})$ referential stored energy (in J/m ³ =Pa),
ρ referential mass density,	$\zeta = \zeta(\mathbf{F}, \mathbf{m}, \theta)$ referential heat part of free energy,
\mathbf{F} deformation gradient,	$\mathbf{e}(\mathbf{v}) = \frac{1}{2}\nabla\mathbf{v}^\top + \frac{1}{2}\nabla\mathbf{v}$ small strain rate (in s ⁻¹),
\mathbf{m} magnetization (in A/m),	$h_c = h_c(\mathbf{F}, \theta)$ magnetic coercive force (in A/m),
θ temperature (in K),	w heat part of internal energy (enthalpy, in J/m ³),
\mathbf{T} Cauchy stress (symmetric, in Pa),	$(\bullet)^\cdot = \frac{\partial}{\partial t}\bullet + (\mathbf{v}\cdot\nabla)\bullet$ convective time derivative,
\mathbf{K} Korteweg stress (symmetric, in Pa),	$(\bullet)^\circ = (\bullet)^\cdot - \text{skw}(\nabla\mathbf{v})\bullet$ corotational time derivative,
\mathbf{S} skew-symmetric stress (in Pa),	$\text{skw } A = \text{a skew-symmetric part, i.e. } \frac{1}{2}A - \frac{1}{2}A^\top,$
\mathbf{D} dissipative stress (in Pa),	\mathcal{S} skew-symmetric magnetic hyperstress (in Pa m),
\mathcal{H} elastic hyperstress (in Pa m),	\cdot or $:$ scalar products of vectors or matrices,
\mathbf{j} heat flux (in W/m ²),	\vdots scalar products of 3rd-order tensors,
\mathbf{k} traction load,	$\mathcal{K} = \mathcal{K}(\mathbf{F}, \theta)$ thermal conductivity (in W/m ⁻² K ⁻¹),
\mathbf{h} (total) magnetic field (in A/m),	$\kappa = \kappa(\mathbf{F})$ exchange-energy coefficient (in kg·m/C ²),
\mathbf{h}_{ext} external magnetic field (in A/m),	$c = c(\mathbf{F}, \theta)$ heat capacity (in Pa/K),
u demagnetizing-field potential (in A),	\mathbf{t} magnetic “driving force” (in Jm ⁻² A ⁻¹),
ξ the return mapping,	$\gamma = \gamma(\mathbf{F}, \mathbf{m}, \theta)$ gyromagnetic ratio (in C/kg),
$\text{Cof}(\cdot)$ cofactor matrix,	τ “viscous” magnetic damping coefficient (in s),
$\nu_b > 0$ a boundary viscosity,	$\nu_1, \nu_2 > 0$ a bulk (hyper)viscosity coefficients,
$\det(\cdot)$ determinant of a matrix,	$\mathbf{h}_{\text{dem}} = -\nabla u$ demagnetizing field (in A/m),
$\mathbb{R}_{\text{sym}}^{d \times d} = \{A \in \mathbb{R}^{d \times d}; A^\top = A\},$	\mathbf{g} external bulk load (gravity acceleration in m/s ²),
$\mathbb{I} \in \mathbb{R}_{\text{sym}}^{d \times d}$ the unit matrix,	μ_0 vacuum permeability ($\sim 1.257 \times 10^{-6}$ H/m).

Table 1. Summary of the basic notation used through Sections 2 and 3.
Convention: upright = referential, slanted = actual.

In comparison with [47, 50], the novelty of this paper is to apply the Eulerian approach to solids in anisothermal situations, using the free energy in a reference configuration, which does not see the energy-pressure in the stress tensor and which is also more fitted with usually available experimental data. The analysis combines L^1 -theory for the heat equation adapted to the convective time derivatives and the techniques from compressible fluid dynamics adapted for solids.

For completeness, let us still mention a competitive, Lagrangian thermodynamic formulation (including also diffusion) [51] formulating the equations in a certain fixed “reference” configuration. This approach allows easily for deformation of the shape of the body and easier treatment inertial forces but a frame-indifferent viscosity and interaction with spatial gravity and magnetic forces is much more complicated.

The plan is as follows: formulation of the model in the actual Eulerian configuration and its energetics and thermodynamics is presented in Section 2, recalling first the micromagnetism and Landau transition in rigid magnets in Sect. 2.1 and finite-strain kinematics of deformable continua in Sect. 2.2 before formulating the model in Section 2.3 and showing its energetics in Section 2.4. Then, in Section 3, the rigorous analysis by a suitable regularization and a (semi) Faedo-Galerkin approximation is performed, combined with theory of transport by regular velocity fields.

2 The thermodynamic model and its energetics

It is important to distinguish carefully the referential and the actual time-evolving coordinates. Our aim is to formulate the model eventually in actual configurations, i.e. the Eulerian formulation, reflecting also the reality in many (or even most) situations (and a certain general agreement) that a reference configuration is only an artificial construction and, even if relevant in some situations, becomes successively more and more irrelevant during evolution at truly finite strains. Typical materials involve magnetic gels or elastomers or magnetic rocks which are viscoelastic on geological timescales. On the other hand, some experimental material data are related to some reference configuration – typically it concerns mass density and stored of free energies per mass (in J/kg) or per referential volume (in J/m³=Pa) as considered here.

We will present briefly the fundamental concepts and formulas which can mostly be found in the monographs, as e.g. [23, Part XI] or [32, Sect. 7.2].

2.1 Micromagnetism and ferro-paramagnetic transition

Let us briefly recall the micromagnetic model in rigid magnets and Landau’s phase-transition theory [27], cf. also [29, Sec.39] or the monographs [6, 15]. The basic ingredient governing static (and later also evolution) model is the free energy $\psi = \psi(\mathbf{m}, \theta)$ depending on magnetization \mathbf{m} and temperature θ .

In the micromagnetism, the free energy ψ is augmented by the exchange energy $\frac{\kappa}{2}|\nabla\mathbf{m}|^2$

with κ a coefficient determining an internal length-scale, responsible for a typical fine domain structure in ferromagnets.

The magnetization itself induces a magnetic field, called a self-induced *demagnetizing field* \mathbf{h}_{dem} . For many (or maybe most) applications, full Maxwell electro-magnetic system is considered simplified to *magnetostatics*, considering slow evolution and neglecting in particular eddy currents and even confining on electrically nonconductive media. The Maxwell system then reduces to the Ampère law $\text{curl} \mathbf{h}_{\text{dem}} = \mathbf{0}$ and the Gauss law $\text{div} \mathbf{b} = 0$ for the magnetic induction with is given by $\mathbf{b} = \mu_0 \mathbf{h}_{\text{dem}} + \mu_0 \mathbf{m}$ where μ_0 is the physical constant (vacuum permeability). The Ampère law ensures existence of a scalar-valued potential u such that $\mathbf{h}_{\text{dem}} = -\nabla u$. These equations are considered on the whole Universe \mathbb{R}^d while, of course, the magnetization \mathbf{m} is only in the body Ω while outside it is considered zero, which is articulated by introducing the characteristic function χ_Ω defined as $\chi_\Omega(\mathbf{x}) = 1$ if $\mathbf{x} \in \Omega$ and $\chi_\Omega(\mathbf{x}) = 0$ if $\mathbf{x} \in \mathbb{R}^d \setminus \Omega$. By substitution, we obtain the equation

$$\text{div}(\nabla u - \chi_\Omega \mathbf{m}) = 0 \quad \text{in } \mathbb{R}^d \quad (2.1)$$

to be considered in the sense of distributions. Under an external magnetic field \mathbf{h}_{ext} , the overall effective magnetic field \mathbf{h} is

$$\mathbf{h} = \mathbf{h}_{\text{ext}} - \mathbf{h}_{\text{dem}} = \mathbf{h}_{\text{ext}} + \nabla u. \quad (2.2)$$

Although not directly relevant in this paper, let us anyhow remind that, for a fixed temperature θ , the standard ferro-magnetostatic theory is based on the free energy $\psi(\mathbf{m}, \theta, \nabla \mathbf{m}) = \tilde{\psi}(\mathbf{m}, \theta) + \frac{\kappa}{2} |\nabla \mathbf{m}|^2$ leading to the overall energy

$$(\mathbf{m}, u) \mapsto \int_\Omega \underbrace{\tilde{\psi}(\mathbf{m}, \theta)}_{\text{free energy}} + \underbrace{\frac{\kappa}{2} |\nabla \mathbf{m}|^2}_{\text{exchange energy}} - \underbrace{\mu_0 (\mathbf{h}_{\text{ext}} + \nabla u) \cdot \mathbf{m}}_{\text{energy of } \mathbf{m} \text{ in the magnetic field } \mathbf{h}} \, d\mathbf{x} - \int_{\mathbb{R}^d} \underbrace{\frac{\mu_0}{2} |\nabla u|^2}_{\text{energy of demagnetizing field}} \, d\mathbf{x}. \quad (2.3)$$

Notably, this functional is concave with respect to u and has a saddle-point character. The static configurations (\mathbf{m}, u) are standardly considered as minimizing with respect to \mathbf{m} and maximizing with respect to u , i.e. a critical point of (2.3). The 1st-order optimality conditions then gives the system

$$\underbrace{\tilde{\psi}'_{\mathbf{m}}(\mathbf{m}, \theta) - \text{div}(\kappa \nabla \mathbf{m})}_{= \text{t magnetic "driving force"}} = \mu_0 \mathbf{h} \quad \text{and} \quad (2.1). \quad (2.4)$$

The mentioned saddle-point character can be eliminated by executing maximization with respect to u , i.e. in fact the partial Legendre transform. This gives, when testing (2.1) by $\mu_0 u$, which gives $\int_{\mathbb{R}^d} \mu_0 |\nabla u|^2 \, d\mathbf{x} = \int_\Omega \mu_0 \mathbf{m} \cdot \nabla u \, d\mathbf{x} = - \int_\Omega \mu_0 \mathbf{m} \cdot \mathbf{h}_{\text{dem}} \, d\mathbf{x}$. Substituting it into (2.3), the functional depending on \mathbf{m} which should be minimized by static configurations is:

$$\mathbf{m} \mapsto \int_\Omega \underbrace{\tilde{\psi}(\mathbf{m}, \theta)}_{\text{free energy}} + \underbrace{\frac{\kappa}{2} |\nabla \mathbf{m}|^2}_{\text{exchange energy}} - \underbrace{\mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m}}_{\text{Zeeman energy}} \, d\mathbf{x} + \int_{\mathbb{R}^d} \underbrace{\frac{\mu_0}{2} |\nabla u_{\mathbf{m}}|^2}_{\text{energy of demagnetizing field}} \, d\mathbf{x}. \quad (2.5)$$

In the rest of this paper, we will couple it with mechanical effects and a full thermodynamics, so that the minimization of energy will no longer be relevant.

In case of time-varying \mathbf{h}_{ext} , a dynamics of \mathbf{m} governed by the *Landau-Lifschitz-Gilbert equation* $\gamma^{-1} \frac{\partial \mathbf{m}}{\partial t} = \mathbf{m} \times \mathbf{h}_{\text{eff}}$ with $\mathbf{h}_{\text{eff}} = \mathbf{h} - \mathbf{t} - \mathbf{t}_{\text{dis}}$ an effective field composed from a conservative part \mathbf{t} arising from a free energy (2.3), cf. (2.4) or also (2.18c) below, while \mathbf{t}_{dis} is a magnetic field counting a dissipative-processes phenomenology, and \mathbf{h} is from (2.2). Equivalently [7], one can write it in the Gilbert form $\gamma^{-1} \mathbf{m} \times \frac{\partial \mathbf{m}}{\partial t} = \mathbf{h}_{\text{eff}}$. The basic choice of \mathbf{t}_{dis} is the magnetic “viscosity” $\tau \frac{\partial \mathbf{m}}{\partial t}$ with τ a phenomenological magnetic damping coefficient. To cover the (temperature dependent) *hysteresis* effects due to so-called pinning mechanism, we augment it by the dry-friction term $h_c(\theta) \text{Dir}(\frac{\partial \mathbf{m}}{\partial t})$ where “Dir” denotes the set-valued monotone “direction” mapping

$$\text{Dir}(\mathbf{r}) = \begin{cases} \{r \in \mathbb{R}^d; |r| \leq 1\} & \text{if } \mathbf{r} = \mathbf{0}, \\ \mathbf{r}/|\mathbf{r}| & \text{if } \mathbf{r} \neq \mathbf{0}, \end{cases} \quad (2.6)$$

cf. [52] and Remark 2.3 below. Note that $\mathbf{r} \cdot \text{Dir}(\mathbf{r}) = |\mathbf{r}|$. Here, having in mind an isotropic situation, $|\cdot|$ denotes the Euclidean norm, but in principle some other anisotropic norms on \mathbb{R}^d can be considered, too. Altogether, we consider the specific Gilbert equation as

$$\tau \frac{\partial \mathbf{m}}{\partial t} + h_c(\theta) \text{Dir}\left(\frac{\partial \mathbf{m}}{\partial t}\right) - \frac{\mathbf{m}}{\gamma(\theta)} \times \frac{\partial \mathbf{m}}{\partial t} = \mu_0 \mathbf{h} - \mathbf{t}. \quad (2.7)$$

The coercive force $h_c = h_c(\theta)$ determines the width of hysteresis loops within slowly time-varying oscillatory external field \mathbf{h}_{ext} . The gyromagnetic term should disappear under high temperatures, i.e. $1/\gamma(\cdot)$ going to 0 for temperatures around or above Curie temperature, as articulated in [34]. Let us note that (2.7) balances the terms in the physical units A/m, as standard.

Example 2.1 (*Ferro-to-para-magnetic transition*). A simplest example of free energy in rigid isotropic magnetic materials is

$$\tilde{\Psi}(\mathbf{m}, \theta) = a_0(\theta - \theta_c) |\mathbf{m}|^2 + b_0 |\mathbf{m}|^4 + c_0 \theta (1 - \ln \theta). \quad (2.8)$$

In static magnetically soft ferromagnetism, the magnetization minimizes the energy. Here the minimum of $\tilde{\Psi}(\bullet, \theta)$ is attained on the orbit $|\mathbf{m}| = m_s(\theta)$ with $m_s(\theta) = \sqrt{a_0(\theta_c - \theta)/(2b_0)}$ if $0 \leq \theta \leq \theta_c$ and at $\mathbf{m} = \mathbf{0}$ if $\theta \geq \theta_c$, cf. the solid line in Figure 1. Under an applied magnetic field \mathbf{h}_{ext} , the minimum of $\mathbf{m} \mapsto \tilde{\Psi}(\mathbf{m}, \theta) - \mathbf{h}_{\text{ext}} \cdot \mathbf{m}$ is at some magnetization whose magnitude is slightly bigger than $m_s(\theta)$, cf. the dashed line in Figure 1. This ansatz can be used for a ferro-para-magnetic transition for a mechanically rigid magnets as formulated (and analyzed) in [42]. This may be quite equally interpreted as ferri-antiferro-magnetic transition, too, cf. [16].




Fig. 1: Typical dependence of saturation magnetization m_s on absolute temperature under zero applied field \mathbf{h} (solid line) and under some applied field (dashed line), cf. e.g. [6].

2.2 Finite-strain kinematics and mass and momentum transport

In finite-strain continuum mechanics, the basic geometrical concept is the time-evolving deformation $\mathbf{y} : \Omega \rightarrow \mathbb{R}^d$ as a mapping from a reference configuration of the body $\Omega \subset \mathbb{R}^d$ into a physical space \mathbb{R}^d . The “Lagrangian” space variable in the reference configuration will be denoted as $\mathbf{X} \in \Omega$ while in the “Eulerian” physical-space variable by $\mathbf{x} \in \mathbb{R}^d$. The basic kinematic and geometrical objects are the Lagrangian velocity $\mathbf{v} = \frac{\partial}{\partial t} \mathbf{y}$ and the Lagrangian deformation gradient $\mathbf{F} = \nabla_{\mathbf{X}} \mathbf{y}$.

We will be interested in deformations $\mathbf{x} = \mathbf{y}(t, \mathbf{X})$ evolving in time, which are sometimes called “motions”. Further, assuming for a moment that $\mathbf{y}(t, \cdot)$ is invertible, we define the so-called *return* (sometimes called also a *reference*) *mapping* $\boldsymbol{\xi} : \mathbf{x} \mapsto \mathbf{y}^{-1}(t, \mathbf{x})$. The important quantities are the Eulerian velocity $\mathbf{v}(t, \mathbf{x}) = \mathbf{v}(t, \boldsymbol{\xi}(t, \mathbf{x}))$ and the Eulerian deformation gradient $\mathbf{F}(t, \mathbf{x}) = \mathbf{F}(t, \boldsymbol{\xi}(t, \mathbf{x}))$.

Here and thorough the whole article, having the Eulerian velocity at disposal, we use the dot-notation $(\cdot)^{\cdot} = \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}}$ for the *convective time derivative* applied to scalars or, component-wise, to vectors or tensors. Then the velocity gradient $\nabla \mathbf{v} = \nabla_{\mathbf{X}} \mathbf{v} \nabla_{\mathbf{x}} \mathbf{X} = \dot{\mathbf{F}} \mathbf{F}^{-1}$, where we used the chain-rule calculus and $\mathbf{F}^{-1} = (\nabla_{\mathbf{X}} \mathbf{x})^{-1} = \nabla_{\mathbf{x}} \mathbf{X}$. This gives the *transport equation-and-evolution for the deformation gradient* as

$$\dot{\mathbf{F}} = (\nabla \mathbf{v}) \mathbf{F}. \quad (2.9)$$

From this, we also obtain the evolution-and-transport equation for Jacobian $\det \mathbf{F}$ as $\overline{\dot{\det \mathbf{F}}} = (\det \mathbf{F}) \operatorname{div} \mathbf{v}$ and its inverse as

$$\overline{\left(\frac{1}{\det \mathbf{F}} \right)^{\cdot}} = - \frac{\operatorname{div} \mathbf{v}}{\det \mathbf{F}}. \quad (2.10)$$

The return mapping $\boldsymbol{\xi}$ satisfies the transport equation

$$\dot{\boldsymbol{\xi}} = \mathbf{0}; \quad (2.11)$$

note that, since we are confined on a spatially homogeneous material, actually $\boldsymbol{\xi}$ does not explicitly occur in the formulation of the problem.

As \mathbf{F} depends on \mathbf{x} , (2.9)–(2.11) are equalities which hold for a.a. \mathbf{x} . The same holds for (2.12)–(2.15) below. Here we will benefit from the boundary condition $\mathbf{v} \cdot \mathbf{n} = 0$ below, which causes that the shape of the actual domain Ω does not evolve in time, i.e. $\Omega = \Omega$. The same convention concerns temperature θ and thus also \mathbf{T} , η , and ν_1 in (2.18d) and (2.20) below, which will make the problem indeed fully Eulerian. Cf. the continuum-mechanics textbooks as e.g. [23, 32].

The mass density (in kg/m³) is an extensive variable, and its transport (expressing that the conservation of mass) writes as the *continuity equation* $\frac{\partial}{\partial t} \varrho + \operatorname{div}(\varrho \mathbf{v}) = 0$, or, equivalently, the *mass evolution-and-transport equation*

$$\dot{\varrho} = -\varrho \operatorname{div} \mathbf{v}. \quad (2.12)$$

Alternatively to (2.12), we will also use an evolution-and-transport equation for the “mass sparsity” as the inverse mass density $1/\varrho$:

$$\frac{\dot{1}}{1/\varrho} = (1/\varrho) \operatorname{div} \mathbf{v}. \quad (2.13)$$

The flow rule for the magnetization (2.7) is now to be considered in deforming medium, and then the partial time derivative in (2.7) should be replaced by an objective time derivative. Here we use the Zaremba-Jaumann (corotational) time derivative $\overset{\circ}{\mathbf{m}}$, defined as

$$\overset{\circ}{\mathbf{m}} = \dot{\mathbf{m}} - \operatorname{skw}(\nabla \mathbf{v}) \mathbf{m} = \frac{\partial \mathbf{m}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{m} - \operatorname{skw}(\nabla \mathbf{v}) \mathbf{m}, \quad (2.14)$$

where $\dot{\mathbf{m}} = \frac{\partial}{\partial t} \mathbf{m} + (\mathbf{v} \cdot \nabla) \mathbf{m}$ denotes the convective derivative of \mathbf{m} . Moreover, in deforming continuum, we can (and should) consider a more general $\gamma = \gamma(\mathbf{F}, \mathbf{m}, \theta)$ and $h_c = h_c(\mathbf{F}, \theta)$. Thus (2.7) turns into

$$\tau \overset{\circ}{\mathbf{m}} + h_c(\mathbf{F}, \theta) \operatorname{Dir}(\overset{\circ}{\mathbf{m}}) - \frac{\mathbf{m} \times \overset{\circ}{\mathbf{m}}}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} = \mu_0 \mathbf{h} - \mathbf{t}. \quad (2.15)$$

The convective derivative itself is not objective and would not be suitable in our context, except perhaps some laminar-like deformation as implicitly used in an incompressible isothermal variant in [5, 25, 53, 60] or in a nanoparticle transport in fluids [22]; for usage of $\overset{\circ}{\mathbf{m}}$ in (2.15) see Remark 2.4 below.

2.3 Magneto-viscoelasticity and its thermodynamics

The main ingredients of the model are the (volumetric) *free energy* ψ and the *dissipative stress*. The Helmholtz free energy $\psi = \psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta)$ is considered per the *referential volume*, while the free energy per actual deformed volume is $\psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta)/\det \mathbf{F}$. Considering the free energy per unit reference volume is more standard in continuum physics [23, 32] than the free energy per actual evolving volume and well corresponds to experimentally

available data. Here also the anisotropy (which is typical in ferromagnets on microscopical scale) in the stored energy needs rather large strains with referential stored energy. This last benefit is related to the fact that the referential free energy does not give an energy pressure contribution to the Cauchy stress (cf. the last term in (2.33) below or [47, Rem. 2]) and allows for more easy decoupling estimation strategy decoupling the magneto-mechanical part and the thermal part of the coupled system.

We will select out the temperature independent stored energy φ and consider the split:

$$\psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta) = \varphi(\mathbf{F}, \mathbf{m}) + \zeta(\mathbf{F}, \mathbf{m}, \theta) + \frac{\kappa(\mathbf{F})}{2} |\nabla \mathbf{m}|^2 \quad \text{with} \quad \zeta(\mathbf{F}, \mathbf{m}, 0) = 0. \quad (2.16)$$

The free energy considered per actual (not referential) volume extended by the Zeeman energy arising by an applied external actual (not referential) magnetic field \mathbf{h}_{ext} , i.e. the Gibbs-type *actual free energy* is thus

$$\psi_G(t; \mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta) = \underbrace{\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}}}_{\text{stored energy}} + \underbrace{\frac{\zeta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}}_{\text{coupling and heat energy}} - \underbrace{\mu_0 \mathbf{h}_{\text{ext}}(t) \cdot \mathbf{m}}_{\text{Zeeman energy}} + \underbrace{\frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}}}_{\text{exchange energy}}. \quad (2.17)$$

Thus the stored energy contain, beside the elastic stored energy, also the so-called anisotropy magnetic energy, which may distinguish directions of easy magnetization on the microscopical single-crystal level. Let us note that this rather general elasto-magnetic ansatz allows for modelling the *magnetic shape-memory materials* and *magnetostrictive effects*.

A particularly simple case $\kappa(\mathbf{F}) = k \det \mathbf{F}$ with some constant k would give the actual exchange energy $k |\nabla \mathbf{m}|^2 / 2$ independent of \mathbf{F} and the actual exchange coefficient as an intensive variable. Yet, as there does not seem any physical reason for exchange coefficient to be an intensive variable, we want to consider a general situation covering in particular also the “referential” case $\kappa(\mathbf{F})$ constant, which corresponds the *actual exchange coefficient* $\kappa(\mathbf{F}) / \det \mathbf{F}$ as an extensive variable.

From the free energy (2.16), we can read as partial (functional) derivatives of ψ with respect to \mathbf{F} , $\nabla \mathbf{m}$, \mathbf{m} , and θ respectively the *conservative part of the actual Cauchy stress* \mathbf{T} , a *capillarity-like stress* \mathbf{K} , the actual conservative *magnetic driving force* \mathbf{t} , and the *actual entropy* η as:

$$\mathbf{T} = \frac{\psi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} = \left(\varphi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}) + \zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) + \frac{\kappa'(\mathbf{F}) |\nabla \mathbf{m}|^2}{2} \right) \frac{\mathbf{F}^\top}{\det \mathbf{F}}, \quad (2.18a)$$

$$\mathbf{K} = - \frac{(\nabla \mathbf{m})^\top \psi'_{\nabla \mathbf{m}}(\mathbf{F}, \nabla \mathbf{m})}{\det \mathbf{F}} = - (\nabla \mathbf{m})^\top \frac{\mu_0 \kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} = - \mu_0 \kappa(\mathbf{F}) \frac{\nabla \mathbf{m} \otimes \nabla \mathbf{m}}{\det \mathbf{F}}, \quad (2.18b)$$

$$\begin{aligned} \mathbf{t} &= \frac{\psi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} - \operatorname{div} \frac{\psi'_{\nabla \mathbf{m}}(\mathbf{F}, \nabla \mathbf{m})}{\det \mathbf{F}} \\ &= \frac{\varphi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}) + \zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} - \operatorname{div} \left(\frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \right), \quad \text{and} \end{aligned} \quad (2.18c)$$

$$\eta = - \frac{\psi'_\theta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} = - \frac{\zeta'_\theta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}. \quad (2.18d)$$

The expected *symmetry* of such part \mathbf{T} of the Cauchy stress is granted by *frame indifference* of $\psi(\cdot, \cdot, \theta)$ and or $\kappa(\cdot)$. This means that

$$\forall(\mathbf{F}, \mathbf{m}, \theta) \in \text{GL}^+(d) \times \mathbb{R}^d \times \mathbb{R}, \quad Q \in \text{SO}(d) : \\ \varphi(\mathbf{F}, \mathbf{m}) = \varphi(Q\mathbf{F}, Q\mathbf{m}), \quad \zeta(\mathbf{F}, \mathbf{m}, \theta) = \zeta(Q\mathbf{F}, Q\mathbf{m}, \theta), \quad \text{and} \quad \kappa(\mathbf{F}) = \kappa(Q\mathbf{F}), \quad (2.19)$$

where $Q \in \text{SO}(d) = \{Q \in \mathbb{R}^{d \times d}; Q^\top Q = QQ^\top = \mathbb{I}\}$ is the special orthogonal group and $\text{GL}^+(d) = \{F \in \mathbb{R}^{d \times d}; \det F > 0\}$ denotes the orientation-preserving general linear group. This in particular implies that the stress \mathbf{T} is symmetric. The symmetry of the capillarity contribution \mathbf{K} to the Cauchy stress is automatic; actually, this contribution as $-(\nabla \mathbf{m})^\top [\psi_G]_{\nabla \mathbf{m}}(\mathbf{F}, \nabla \mathbf{m})$ was devised in [7, Formula (2.27)] or [13, Formula (5.16)].

Mainly for analytical reasons, we will use also a dissipative contribution to the Cauchy stress which, together with the conservative part \mathbf{T} , will realize the *Kelvin-Voigt rheological model* and make the system parabolic. To this goal, we consider a *dissipative* contribution to the *Cauchy stress* involving the standard dissipative stress depending (from the frame-invariancy reason) on the symmetric velocity gradient and also a higher-order elastic *hyperstress* \mathcal{H} , both isotropic for simplicity:

$$\mathbf{D} - \text{div } \mathcal{H} \quad \text{with} \quad \mathbf{D} = \mathbf{D}(\mathbf{e}(\mathbf{v})) \quad \text{and} \quad \mathcal{H} = \mathcal{H}(\nabla^2 \mathbf{v}) \\ \text{for} \quad \mathbf{D}(\mathbf{e}) = \nu_1 |\mathbf{e}|^{p-2} \mathbf{e} \quad \text{and} \quad \mathcal{H}(\mathbf{E}) = \nu_2 |\mathbf{E}|^{p-2} \mathbf{E}. \quad (2.20)$$

Actually, ν_1 and ν_2 may depend on $\det \mathbf{F}$ and θ without causing any structural and analytical problems, but we ignore it rather for notational simplicity.

The *momentum equilibrium* equation then balances the divergence of the total Cauchy stress with the inertial and gravity force:

$$\rho \dot{\mathbf{v}} - \text{div}(\mathbf{T} + \mathbf{D} + \mathbf{T}_{\text{mag}} - \text{div}(\mathcal{H} + \mathcal{S})) = \rho \mathbf{g} + \mathbf{f}_{\text{mag}} \quad (2.21)$$

with \mathbf{T} from (2.18d) and \mathbf{D} and \mathcal{H} from (2.20). Moreover, \mathbf{T}_{mag} and \mathbf{f}_{mag} are the magnetic stress and the magnetic force which balance the energetics, cf. $\mathbf{T}_{\text{mag}} := \mathbf{K} + \mathbf{S}$ and $\mu_0(\nabla \mathbf{h})^\top \mathbf{m} - \mu_0 \nabla(\mathbf{h} \cdot \mathbf{m}) =: \mathbf{f}_{\text{mag}}$ while $\mathcal{S} := \kappa(\mathbf{F}) \text{Skw}(\nabla \mathbf{m} \otimes \mathbf{m}) / \det \mathbf{F}$ will be a “magnetic exchange hyperstress” in (2.30b).

The driving magnetic force (2.18c) enters the Landau-Lifschitz-Gilbert equation (2.15) in the previous section.

The third ingredient, i.e. (2.18d), is subjected to the *entropy equation*:

$$\frac{\partial \eta}{\partial t} + \text{div}(\mathbf{v} \eta) = \frac{\xi - \text{div } \mathbf{j}}{\theta} \quad \text{with} \quad \mathbf{j} = -\mathcal{K}(\mathbf{F}, \theta) \nabla \theta \quad (2.22)$$

and with $\xi = \xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \dot{\mathbf{m}})$ denoting the heat production rate specified later in (2.30f). The latter equality in (2.22) is the *Fourier law* determining phenomenologically the heat flux \mathbf{j} proportional to the negative gradient of temperature θ through the heat conduction coefficient $\mathcal{K} = \mathcal{K}(\mathbf{F}, \theta)$. Assuming $\xi \geq 0$ and $\mathcal{K} \geq 0$ and integrating (2.22) over the domain Ω while imposing the non-penetrability of the boundary in the sense that the

normal velocity $\mathbf{v} \cdot \mathbf{n}$ vanishes across the boundary Γ of Ω , we obtain the *Clausius-Duhem inequality*:

$$\frac{d}{dt} \int_{\Omega} \eta \, d\mathbf{x} = \underbrace{\int_{\Omega} \left(\frac{\xi}{\theta} + \kappa \frac{|\nabla \theta|^2}{\theta^2} \right) d\mathbf{x}}_{\text{entropy production rate}} + \underbrace{\int_{\Gamma} \left(\kappa \frac{\nabla \theta}{\theta} - \eta \mathbf{v} \right) \cdot \mathbf{n} \, dS}_{\text{entropy flux}} \geq \int_{\Gamma} \kappa \frac{\nabla \theta \cdot \mathbf{n}}{\theta} \, dS. \quad (2.23)$$

If the system is thermally isolated in the sense that the normal heat flux $\mathbf{j} \cdot \mathbf{n}$ vanishes across the boundary Γ , we recover the *2nd law of thermodynamics*, i.e. the total entropy in isolated systems is nondecreasing in time.

Substituting η from (2.18d) into (2.22) written in the form $\theta \dot{\eta} = \xi - \operatorname{div} \mathbf{j} - \theta \eta \operatorname{div} \mathbf{v}$, we obtain

$$\begin{aligned} c(\mathbf{F}, \mathbf{m}, \theta) \dot{\theta} &= \xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \dot{\mathbf{m}}) + \theta \left(\frac{\zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right)'_{\mathbf{F}} : \dot{\mathbf{F}} \\ &\quad + \theta \left(\frac{\zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right)'_{\mathbf{m}} \cdot \dot{\mathbf{m}} + \theta \frac{\zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \operatorname{div} \mathbf{v} - \operatorname{div} \mathbf{j} \\ &= \xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \dot{\mathbf{m}}) + \theta \frac{\zeta''_{\mathbf{F}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} : \dot{\mathbf{F}} + \theta \frac{\zeta''_{\mathbf{m}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} - \operatorname{div} \mathbf{j} \\ &\quad \text{with the heat capacity } c(\mathbf{F}, \mathbf{m}, \theta) = -\theta \frac{\zeta''_{\theta\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}, \end{aligned} \quad (2.24)$$

which can be understood as the *heat equation* for the temperature θ as an intensive variable.

The referential *internal energy* is given by the *Gibbs relation* $\psi + \theta \eta$. In our Eulerian formulation, we will need rather the actual internal energy, which, in view of (2.18d), equals here to

$$\underbrace{\frac{\psi - \theta \psi'_{\theta}}{\det \mathbf{F}}}_{\text{actual internal energy}} = \underbrace{\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} + \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}}}_{\text{actual stored and exchange energy}} + \underbrace{\frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}}_{= w \text{ thermal part of the internal energy}}. \quad (2.25)$$

In terms of w , the heat equation (2.24) can be written in the so-called *enthalpy formulation*:

$$\begin{aligned} \frac{\partial w}{\partial t} + \operatorname{div}(\mathbf{v} w) &= \xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \dot{\mathbf{m}}) + \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} : \dot{\mathbf{F}} + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} - \operatorname{div} \mathbf{j} \\ &\quad \text{with } w = \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}. \end{aligned} \quad (2.26)$$

Note that w is an extensive variable so that the left-hand side of (2.26) is not just a convective derivative \dot{w} . For the passage from (2.24) to (2.26), we use the algebra $\mathbf{F}^{-1} = \operatorname{Cof} \mathbf{F}^{\top} / \det \mathbf{F}$ and the calculus $\det'(F) = \operatorname{Cof} F$ and (2.9) so that $(1/\det \mathbf{F})' : \dot{\mathbf{F}} = -(\operatorname{Cof} \mathbf{F} / \det \mathbf{F}^2) : (\nabla \mathbf{v}) \mathbf{F} = -(\mathbf{F}^{-\top} / \det \mathbf{F}) \mathbf{F}^{\top} : \nabla \mathbf{v} = (\operatorname{div} \mathbf{v}) / \det \mathbf{F}$, and thus we can calculate

$$\begin{aligned} \frac{\partial w}{\partial t} + \operatorname{div}(\mathbf{v} w) &= \dot{w} + w \operatorname{div} \mathbf{v} \\ &= \left(\frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right)' + \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \end{aligned}$$

$$\begin{aligned}
&= \left(\left(\frac{\zeta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right)'_{\mathbf{F}} - \theta \left(\frac{\zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right)'_{\mathbf{F}} \right) : \dot{\mathbf{F}} - \theta \frac{\zeta''_{\theta\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \dot{\theta} \\
&\quad + \left(\frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} - \theta \frac{\zeta''_{\mathbf{m}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right) : \dot{\mathbf{m}} + \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \\
&= \left(\frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} - \theta \frac{\zeta''_{\mathbf{F}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + (\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)) \left(\frac{1}{\det \mathbf{F}} \right)' \right) : \dot{\mathbf{F}} \\
&\quad + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta''_{\mathbf{m}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} : \dot{\mathbf{m}} + c(\mathbf{F}, \mathbf{m}, \theta) \dot{\theta} + \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_{\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \\
&= \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta''_{\mathbf{F}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} : \dot{\mathbf{F}} + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta''_{\mathbf{m}\theta}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} : \dot{\mathbf{m}} + c(\mathbf{F}, \mathbf{m}, \theta) \dot{\theta}.
\end{aligned}$$

This shows that (2.24) are (2.26) indeed (formally) equivalent to each other. Alternatively in (2.26), we could use (2.9) for writing $\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) : \dot{\mathbf{F}} = \zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^{\top} : (\nabla \mathbf{v}) = \zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^{\top} : \mathbf{e}(\mathbf{v})$; here the assumed frame indifference (2.19) of $\zeta(\cdot, \mathbf{m}, \theta)$ itself, leading to symmetry of $\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^{\top}$, was employed.

Remark 2.2 (*Gradient theories in rates*). So-called gradient theories in continuum mechanical models are nowadays very standard, referred as *nonsimple materials*, determining some internal length scales and often facilitating mathematical analysis. They can be applied to the conservative stress through the free energy or to the dissipation stress. Here, we have used the latter option in (2.20) which is better fitted to the rate formulation and which can make velocity field enough regular, as vitally needed for the transport of ϱ and \mathbf{F} in the Eulerian models. The higher-gradient hyper-stress as used below in (2.20) follows the theory by E. Fried and M. Gurtin [20], as already articulated in the general nonlinear context of *multipolar fluids* by J. Nečas et al. [37–39] or of solids [44, 55], inspired by R.A. Toupin [56] and R.D. Mindlin [36].

Remark 2.3 (*Dry friction in magnetization evolution*). Dry-friction-type rate-independent dissipation was proposed in [4, 59] as a device to model properly hysteresis in ferromagnets, modifying the Landau-Lifschitz equation by augmenting suitably the effective magnetic field. Although the original Gilbert's [21] and Landau-Lifschitz' [28] equations are equivalent with each other, the resulting augmented equations proposed in [4] and [59] are no longer mutually equivalent. This has been pointed out in [41], where the conceptual differences between the Gilbert and the Landau-Lifschitz formats have been elucidated.

Remark 2.4 (*Zaremba-Jaumann derivative $\dot{\mathbf{m}}$*). In deformable (and deforming) magnetic medium, the Jaumann corotational derivative for magnetization was suggested already by Maugin [33] to model situations when the magnetization can be “frozen” in hard-magnetic materials in their ferro- or ferri-magnetic state. Later it was used in [12, 13] in the linear viscosity (magnetic attenuation) term. It should be noted that the gyromagnetic term $\mathbf{m} \otimes \dot{\mathbf{m}} / \gamma(\mathbf{F}, \mathbf{m}, \theta)$ can be seen in literature; yet, it is the same as used in (2.15) since $\mathbf{m} \otimes \dot{\mathbf{m}} - \mathbf{m} \otimes \dot{\mathbf{m}} = \mathbf{m} \otimes (\operatorname{skw}(\nabla \mathbf{v}) \mathbf{m}) = \operatorname{skw}(\nabla \mathbf{v}) : (\mathbf{m} \otimes \mathbf{m}) = 0$. Instead of $\operatorname{skw}(\nabla \mathbf{v})$ in (2.14), an angular velocity either governed by a separate parabolic equation or approximated by $\operatorname{curl} \mathbf{v} / 2$ was used in [22, 40, 45, 54]. Usage of another (Lie) derivative was

proposed in [58, Formula (74)]. Mere convective derivative for magnetization has been used in [5, 25, 53] to model rather (incompressible isothermal) fluids containing magnetic particles.

Example 2.5 (*Neo-Hookean elastic magnets*). Modifying slightly the “rigid” model (2.8) and expanding it by standard neo-Hookean elastic ansatz, one obtains an example for elastic magnetic material amenable for ferro-to-paramagnetic transition and for complying with the (3.5b–e) below:

$$\begin{aligned} \psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta) = & \frac{1}{2}G \left(\frac{\text{tr}(\mathbf{F}\mathbf{F}^\top)}{(\det \mathbf{F})^{2/d}} - d \right) + v(\det \mathbf{F}) + \frac{\kappa(\mathbf{F})}{2} |\nabla \mathbf{m}|^2 \\ & + a_0(\det \mathbf{F}) \left(\frac{\theta}{1+\epsilon_1\theta} - \frac{\theta_c}{1+\epsilon_1\theta_c} \right) \frac{|\mathbf{m}|^2}{1+\epsilon_2|\mathbf{m}|^2} + b_0(\det \mathbf{F}) |\mathbf{m}|^4 + c_0\theta(1-\ln\theta) \end{aligned}$$

with some (referential) heat capacity $c_0 > 0$, some $\epsilon_1, \epsilon_2 > 0$, some shear modulus $G > 0$, and the non-negative volumetric energy $v \in C^1(\mathbb{R}^+)$ and $a_0(J) \geq \delta J$ and $b_0(J) \geq \delta J$ for some $\delta > 0$ so that φ fulfills (up to an irrelevant constant) the coercivity (3.5b) with $s = 4$. Note that $\psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \cdot)$ is concave for $c_0, c_1 > 0$ and $\epsilon_1, \epsilon_2 \geq 0$. Then, the split (2.16) uses

$$\zeta(\mathbf{F}, \mathbf{m}, \theta) = \frac{a_0(\det \mathbf{F})\theta|\mathbf{m}|^2}{(1+\epsilon_1\theta)(1+\epsilon_2|\mathbf{m}|^2)} + c_0\theta(1-\ln\theta) \quad (2.27)$$

so that

$$\omega(\mathbf{F}, \mathbf{m}, \theta) = \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_\theta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} = \frac{c_0\theta}{\det \mathbf{F}} + \frac{\epsilon_1 a_0(\det \mathbf{F})\theta^2|\mathbf{m}|^2}{(1+\epsilon_1\theta)^2(1+\epsilon_2|\mathbf{m}|^2)\det \mathbf{F}} \quad (2.28)$$

and the (actual) heat capacity $c(\mathbf{F}, \mathbf{m}, \theta) = -\theta \zeta''_{\theta\theta}(\mathbf{F}, \mathbf{m}, \theta)/\det \mathbf{F}$ is

$$c(\mathbf{F}, \mathbf{m}, \theta) = \omega'_\theta(\mathbf{F}, \mathbf{m}, \theta) = \frac{c_0}{\det \mathbf{F}} + \frac{2\epsilon_1 a_0(\det \mathbf{F})\theta|\mathbf{m}|^2}{(1+\epsilon_1\theta)^3(1+\epsilon_2|\mathbf{m}|^2)\det \mathbf{F}}. \quad (2.29)$$

Note that “thermo-coupling” stress $\zeta'_\theta(\mathbf{F}, \mathbf{m}, \theta)\mathbf{F}^\top/\det \mathbf{F} = a'_0(\det \mathbf{F})\theta|\mathbf{m}|^2\mathbb{I}/((1+\epsilon_1\theta)(1+\epsilon_2|\mathbf{m}|^2))$ is bounded provided a'_0 is bounded on $\text{GL}^+(d)$, so it surely complies with (3.5c) below. Also $|\zeta'_\theta(\mathbf{F}, \cdot, \cdot)/\det \mathbf{F}|$ is bounded for \mathbf{F} ranging over compact sets in $\text{GL}^+(d)$, so it surely complies with (3.5c); here we use $\epsilon_1 > 0$ and $\epsilon_2 > 0$. Also this ansatz satisfies (3.5d). Moreover, $|\omega'_\theta(\mathbf{F}, \mathbf{m}, \cdot)|$ has at most linear growth while $|\omega'_\theta(\mathbf{F}, \mathbf{m}, \cdot)|$ is even bounded as well as $|\omega''_{\theta\theta}|$ and $|\omega''_{\theta\mathbf{m}}|$ for \mathbf{F} ranging over compact sets in $\text{GL}^+(d)$, so that (3.5d) is satisfied, too.

2.4 The thermo-magneto-mechanical system and its energetics

Let us summarize the thermodynamically coupled system composed of six partial differential equations for ϱ , \mathbf{v} , \mathbf{F} , \mathbf{m} , u , and θ . More specifically, it is composed from the mass continuity equation for ϱ , the momentum equation written in terms of velocity \mathbf{v} , the evolution-and-transport of the deformation-gradient tensor \mathbf{F} , a flow rule (as an inclusion) for the magnetization \mathbf{m} , the Poisson equation for the demagnetizing-field potential u , and the heat-transfer equation for temperature θ .

Altogether, merging (2.1), (2.9), (2.12), (2.15), (2.21), and (2.26) with (2.18), we obtain a system of six equations for $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, u, w)$, respectively also for θ :

$$\frac{\partial \varrho}{\partial t} = -\operatorname{div}(\varrho \mathbf{v}), \quad (2.30a)$$

$$\frac{\partial}{\partial t}(\varrho \mathbf{v}) = \operatorname{div}\left(\mathbf{T} + \mathbf{K} + \mathbf{S} + \mathbf{D} - \operatorname{div}(\mathcal{H} + \mathcal{S}) - \varrho \mathbf{v} \otimes \mathbf{v}\right) + \mu_0(\nabla \mathbf{h})^\top \mathbf{m} - \mu_0 \nabla(\mathbf{h} \cdot \mathbf{m}) + \varrho \mathbf{g}$$

$$\text{with } \mathbf{T} = \left(\frac{\varphi'_F(\mathbf{F}, \mathbf{m}) + \zeta'_F(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F})}{2 \det \mathbf{F}} \right) \mathbf{F}^\top, \quad \mathbf{h} = \mathbf{h}_{\text{ext}} + \nabla u,$$

$$\mathbf{K} = \frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \nabla \mathbf{m} \otimes \nabla \mathbf{m}, \quad \mathbf{D} = \nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}), \quad \mathcal{H} = \nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v},$$

$$\mathbf{S} = \operatorname{skw}\left(\left(\mu_0 \mathbf{h} - \frac{\psi'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}\right) \otimes \mathbf{m}\right), \quad \text{and } \mathcal{S} = \frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \operatorname{Skw}(\mathbf{m} \otimes \nabla \mathbf{m}), \quad (2.30b)$$

$$\frac{\partial \mathbf{F}}{\partial t} = (\nabla \mathbf{v}) \mathbf{F} - (\mathbf{v} \cdot \nabla) \mathbf{F}, \quad (2.30c)$$

$$\tau \dot{\mathbf{m}} + h_c(\mathbf{F}, \theta) \operatorname{Dir}(\dot{\mathbf{m}}) - \frac{\mathbf{m} \times \dot{\mathbf{m}}}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} \ni \mu_0 \mathbf{h} - \frac{\psi'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + \operatorname{div}\left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \nabla \mathbf{m}\right), \quad (2.30d)$$

$$\Delta u = \operatorname{div}(\chi_\Omega \mathbf{m}) \quad \text{on } \mathbb{R}^d, \quad (2.30e)$$

$$\frac{\partial w}{\partial t} = \xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \dot{\mathbf{m}}) + \frac{\zeta'_F(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} : \mathbf{e}(\mathbf{v}) + \frac{\zeta'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} - \operatorname{div}(\mathbf{j} + w \mathbf{v})$$

$$\text{with } \xi(\mathbf{F}, \theta; \mathbf{e}, \mathbf{G}, \mathbf{r}) = \nu_1 |\mathbf{e}|^p + \nu_2 |\mathbf{G}|^p + \mu_0 \tau |\mathbf{r}|^2 + \mu_0 h_c(\mathbf{F}, \theta) |\mathbf{r}|$$

$$\text{and } \mathbf{j} = -\mathcal{K}(\mathbf{F}, \theta) \nabla \theta,$$

$$\text{and } w = \omega(\mathbf{F}, \mathbf{m}, \theta) = \frac{\zeta(\mathbf{F}, \mathbf{m}, \theta) - \theta \zeta'_\theta(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}}. \quad (2.30f)$$

The product $\nabla \mathbf{m} \otimes \nabla \mathbf{m}$ in (2.30b) is to be understood componentwise, specifically $[\nabla \mathbf{m} \otimes \nabla \mathbf{m}]_{ij} = \sum_{k=1}^d \frac{\partial}{\partial x_i} m_k \frac{\partial}{\partial x_j} m_k$ with $\mathbf{m} = (m_1, \dots, m_d)$ and $\mathbf{x} = (x_1, \dots, x_d)$, while the skew-symmetric part “Skw” of the 3rd-order tensor is defined as

$$[\operatorname{Skw}(\mathbf{m} \otimes \nabla \mathbf{m})]_{ijk} := \frac{1}{2} \left(m_i \frac{\partial m_j}{\partial x_k} - m_j \frac{\partial m_i}{\partial x_k} \right). \quad (2.31)$$

The equations (2.30a-d,f) are considered on the domain Ω while the static equation (2.30e) is to hold on the whole Universe in the distributional sense at all time instants. The symmetric Korteweg-like stress $\mathbf{K} = \kappa(\mathbf{F}) \nabla \mathbf{m} \otimes \nabla \mathbf{m} / \det \mathbf{F}$ in (2.30b) occur e.g. in [7, 17]. The skew-symmetric stress \mathbf{S} and couple-like hyperstress \mathcal{S} come from the calculation (2.39); for a similar skew-symmetric stress \mathbf{S} see [12, Formula (33)] or [13, Formula (5.37)] while the skew-symmetric hyperstress \mathcal{S} is like in the Cosserat theory in [57]. The pressures $\mu_0 \mathbf{h} \cdot \mathbf{m}$ in the momentum equation is related with that the Zeeman and the demagnetizing-field energies are actual (not referential). The magnetic, so-called Kelvin force $\mu_0(\nabla \mathbf{h})^\top \mathbf{m}$ comes from the magnetization \mathbf{m} exposed to the nonuniform magnetic field \mathbf{h} , and it arises from the calculus (2.46) and (2.48) below.

Denoting by \mathbf{n} the unit outward normal to the (fixed) boundary Γ of the domain Ω , we complete this system by suitable boundary conditions:

$$\mathbf{v} \cdot \mathbf{n} = 0, \quad \left[(\mathbf{T} + \mathbf{K} + \mathbf{S} + \mathbf{D} - \operatorname{div}(\mathcal{H} + \mathcal{S})) \mathbf{n} - \operatorname{div}_s(\mathcal{H} \mathbf{n} + \mathcal{S} \mathbf{n}) \right]_\top + \nu_b |\mathbf{v}|^{p-2} \mathbf{v} = \mathbf{k}, \quad (2.32a)$$

$$\nabla^2 \mathbf{v} : (\mathbf{n} \otimes \mathbf{n}) = \mathbf{0}, \quad (\mathbf{n} \cdot \nabla) \mathbf{m} = \mathbf{0}, \quad u(\infty) = 0, \quad \text{and} \quad \mathbf{j} \cdot \mathbf{n} = h(\theta) + \frac{\nu_b}{2} |\mathbf{v}|^p \quad (2.32b)$$

with $\nu_b > 0$ a boundary viscosity coefficient and with $[\cdot]_{\text{T}}$ a tangential part of a vector and with $\text{div}_s = \text{tr}(\nabla_s)$ denoting the $(d-1)$ -dimensional surface divergence with $\text{tr}(\cdot)$ being the trace of a $(d-1) \times (d-1)$ -matrix and $\nabla_s v = \nabla v - \frac{\partial v}{\partial \mathbf{n}} \mathbf{n}$ being the surface gradient of v . Naturally, $\mathbf{k} \cdot \mathbf{n} = 0$ is to be assumed if we want to recover the boundary conditions (2.32a) in the classical form, otherwise the weak form does not directly need it.

The first condition (i.e. normal velocity zero) expresses nonpenetrability of the boundary was used already for (2.23) and is most frequently adopted in literature for Eulerian formulation. This simplifying assumption fixes the shape of Ω in its referential configuration allows also for considering fixed boundary even for such time-evolving Eulerian description. The latter condition in (2.32a) involving a boundary viscosity comes from the Navier boundary condition largely used in fluid dynamics and is here connected with the technique used below, which is based on the total energy balance as the departing point and which, unfortunately, does not allow to cope with $\nu_b = 0$ and simultaneously $\mathbf{k} \neq \mathbf{0}$. This boundary viscosity naturally may contribute to the heat production on the boundary as well as to the outflow of the heat energy to the outer space. For notational simplicity, we consider that it is just equally distributed, one part remaining on the boundary of Ω and the other part leaving outside, which is related with the coefficient $1/2$ in the last condition in (2.32b). The condition $u(\infty) = 0$ in (2.32b) expresses shortly that $\lim_{|\mathbf{x}| \rightarrow 0} u(\mathbf{x}) = 0$.

The magnetization flow rule (2.30d) with the corotational derivative $\overset{\circ}{\mathbf{m}}$ in see also [7, 33] where it is articulated that the magnetization is “frozen” in the deforming medium if $\overset{\circ}{\mathbf{m}} = \mathbf{0}$ which then means that the magnetization is transported and rotates at the same local rate as the deforming medium; this is the situation below the blocking temperature θ_b and when the total driving magnetic field has small magnitude.

For the capillarity-like stress \mathbf{K} and the skew-symmetric stress \mathbf{S} see also [7, 12].

The magneto-mechanical energy balance of the model can be seen when testing the momentum equation (2.30b) by \mathbf{v} while using the continuity equation (2.30a) tested by $|\mathbf{v}|^2/2$ and the evolution-and-transport equation (2.30c) for \mathbf{F} tested by the stress $[\varphi(\mathbf{F}, \mathbf{m})/\det \mathbf{F}]'_{\mathbf{F}}$, the magnetic Landau-Lifshitz-Gilbert equation (2.30d) by $\overset{\circ}{\mathbf{m}}$, and the (rest from the) Maxwell system (2.30e) by $\mu_0 \frac{\partial u}{\partial t}$.

Let us first make the test of (2.30b) by \mathbf{v} . Using again the algebra $F^{-1} = \text{Cof } F^{\top} / \det F$ and the calculus $\det'(F) = \text{Cof } F$, we can write the part of the Cauchy stress arising from the stored energy as

$$\begin{aligned} \frac{\varphi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \mathbf{F}^{\top} &= \frac{\varphi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}) - \varphi(\mathbf{F}, \mathbf{m}) \mathbf{F}^{-\top}}{\det \mathbf{F}} \mathbf{F}^{\top} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \mathbb{I} \\ &= \left(\frac{\varphi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} - \frac{\varphi(\mathbf{F}, \mathbf{m}) \text{Cof } \mathbf{F}}{(\det \mathbf{F})^2} \right) \mathbf{F}^{\top} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \mathbb{I} \left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} \mathbf{F}^{\top} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \mathbb{I}. \end{aligned} \quad (2.33)$$

Let us recall that $\varphi(\mathbf{F}, \mathbf{m})/\det \mathbf{F}$ in (2.33) is the stored energy per actual (not referential) volume. Using the calculus (2.33), we obtain

$$\int_{\Omega} \text{div } \mathbf{T} \cdot \mathbf{v} \, d\mathbf{x} = \int_{\Gamma} (\mathbf{T} \mathbf{n}) \cdot \mathbf{v} \, dS - \int_{\Omega} \mathbf{T} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} \quad \text{with}$$

$$\begin{aligned}
\int_{\Omega} \mathbf{T} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} &= \int_{\Omega} \left(\frac{\varphi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}) + \zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F})}{2 \det \mathbf{F}} \right) \mathbf{F}^{\top} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} \\
&= \int_{\Omega} \left(\left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} \mathbf{F}^{\top} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \mathbb{I} + \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \mathbf{F}^{\top} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F})}{2 \det \mathbf{F}} \mathbf{F}^{\top} \right) : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} \\
&= \int_{\Omega} \left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} : (\nabla \mathbf{v}) \mathbf{F} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} + \left(\frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F})}{2 \det \mathbf{F}} \right) \mathbf{F}^{\top} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} \\
&= \frac{d}{dt} \int_{\Omega} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \, d\mathbf{x} - \int_{\Omega} \frac{\varphi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} - \left(\frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F})}{2 \det \mathbf{F}} \right) \mathbf{F}^{\top} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x}. \quad (2.34)
\end{aligned}$$

Here, we used the matrix algebra $A:(BC) = (B^{\top}A):C = (AC^{\top}):B$ for any square matrices A , B , and C and also we used (2.30c) together with the Green formula and the nonpenetrability boundary condition for

$$\begin{aligned}
&\int_{\Omega} \left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} : (\nabla \mathbf{v}) \mathbf{F} + \frac{\varphi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \, d\mathbf{x} \\
&\stackrel{(2.30c)}{=} \int_{\Omega} \left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} : \left(\frac{\partial \mathbf{F}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{F} \right) + \frac{\varphi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \cdot \left(\frac{\partial \mathbf{m}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{m} \right) + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \, d\mathbf{x} \\
&= \frac{d}{dt} \int_{\Omega} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \, d\mathbf{x} + \int_{\Omega} \nabla \left(\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right) \cdot \mathbf{v} + \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \, d\mathbf{x} \\
&= \frac{d}{dt} \int_{\Omega} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \, d\mathbf{x} + \int_{\Gamma} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \underbrace{(\mathbf{v} \cdot \mathbf{n})}_{=0} \, dS, \quad (2.35)
\end{aligned}$$

where we used

$$\begin{aligned}
&\int_{\Omega} \left[\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right]'_{\mathbf{F}} : (\mathbf{v} \cdot \nabla) \mathbf{F} + \frac{\varphi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \cdot (\mathbf{v} \cdot \nabla) \mathbf{m} \, d\mathbf{x} \\
&= \int_{\Omega} \nabla \left(\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right) \cdot \mathbf{v} \, d\mathbf{x} = \int_{\Gamma} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \underbrace{\mathbf{v} \cdot \mathbf{n}}_{=0} \, dS - \int_{\Omega} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} \, d\mathbf{x}. \quad (2.36)
\end{aligned}$$

The further contribution from the dissipative part of the Cauchy stress uses Green's formula over Ω twice and the surface Green formula over Γ . We abbreviate the elastic hyperstress $\mathcal{H} = \nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v}$. Then

$$\begin{aligned}
&\int_{\Omega} \operatorname{div} (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) \cdot \mathbf{v} \, d\mathbf{x} \\
&= \int_{\Gamma} \mathbf{v} \cdot (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) \mathbf{n} \, dS - \int_{\Omega} (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) : \nabla \mathbf{v} \, d\mathbf{x} \\
&= \int_{\Gamma} \mathbf{v} \cdot (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) \mathbf{n} - \mathbf{n} \cdot \mathcal{H} : \nabla \mathbf{v} \, dS - \int_{\Omega} \nu_1 |\mathbf{e}(\mathbf{v})|^p + \mathcal{H} : \nabla^2 \mathbf{v} \, d\mathbf{x} \\
&= \int_{\Gamma} \mathcal{H} : (\partial_{\mathbf{n}} \mathbf{v} \otimes \mathbf{n} \otimes \mathbf{n}) + \mathbf{n} \cdot \mathcal{H} : \nabla_{\mathbf{s}} \mathbf{v} + \mathbf{v} \cdot (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) \mathbf{n} \, dS \\
&\quad - \int_{\Omega} \nu_1 |\mathbf{e}(\mathbf{v})|^p + \nu_2 |\nabla^2 \mathbf{v}|^p \, d\mathbf{x} \\
&= \int_{\Gamma} \mathcal{H} : (\partial_{\mathbf{n}} \mathbf{v} \otimes \mathbf{n} \otimes \mathbf{n}) - (\operatorname{div}_{\mathbf{s}} (\mathbf{n} \cdot \mathcal{H}) + (\nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \operatorname{div} \mathcal{H}) \mathbf{n}) \cdot \mathbf{v} \, dS
\end{aligned}$$

$$- \int_{\Omega} \nu_1 |\mathbf{e}(\mathbf{v})|^p + \nu_2 |\nabla^2 \mathbf{v}|^p \, d\mathbf{x}, \quad (2.37)$$

where we used the decomposition of $\nabla \mathbf{v}$ into its normal component $\partial_{\mathbf{n}} \mathbf{v}$ and the tangential component, i.e. written componentwise $\nabla \mathbf{v}_i = (\mathbf{n} \cdot \nabla \mathbf{v}_i) \mathbf{n} + \nabla_s \mathbf{v}_i$.

Furthermore, the inertial force $\frac{\partial}{\partial t}(\varrho \mathbf{v}) + \operatorname{div}(\varrho \mathbf{v} \otimes \mathbf{v})$ in (2.30b) tested by \mathbf{v} gives the rate of kinetic energy $\varrho |\mathbf{v}|^2/2$ integrating over Ω . Here we use the continuity equation (2.12) tested by $|\mathbf{v}|^2/2$ and the Green formula with the boundary condition $\mathbf{v} \cdot \mathbf{n} = 0$:

$$\int_{\Omega} \left(\frac{\partial}{\partial t}(\varrho \mathbf{v}) + \operatorname{div}(\varrho \mathbf{v} \otimes \mathbf{v}) \right) \cdot \mathbf{v} \, d\mathbf{x} = \int_{\Omega} \varrho \dot{\mathbf{v}} \cdot \mathbf{v} \, d\mathbf{x} = \frac{d}{dt} \int_{\Omega} \frac{\varrho}{2} |\mathbf{v}|^2 \, d\mathbf{x} + \int_{\Gamma} \varrho |\mathbf{v}|^2 \underbrace{\mathbf{v} \cdot \mathbf{n}}_{=0} \, dS. \quad (2.38)$$

The test of (2.30d) by $\dot{\mathbf{m}}$ is quite technical. The exchange-energy term $\operatorname{div}(\kappa(\mathbf{F}) \nabla \mathbf{m})/\det \mathbf{F}$ tested by $(\mathbf{v} \cdot \nabla) \mathbf{m}$ is to be handled by using Green's formula twice. Namely,

$$\begin{aligned} \int_{\Omega} \operatorname{div} \left(\frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \right) \cdot (\mathbf{v} \cdot \nabla) \mathbf{m} \, d\mathbf{x} &= \int_{\Gamma} \frac{\kappa(\mathbf{F})(\mathbf{n} \cdot \nabla) \mathbf{m}}{\det \mathbf{F}} \cdot (\mathbf{v} \cdot \nabla) \mathbf{m} \, dS \\ &\quad - \int_{\Omega} \frac{\kappa(\mathbf{F}) \nabla^2 \mathbf{m}}{\det \mathbf{F}} : (\mathbf{v} \otimes \nabla \mathbf{m}) + \frac{\kappa(\mathbf{F})(\nabla \mathbf{m} \otimes \nabla \mathbf{m})}{\det \mathbf{F}} : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} \\ &= \int_{\Gamma} \frac{\kappa(\mathbf{F})(\mathbf{n} \cdot \nabla) \mathbf{m}}{\det \mathbf{F}} \cdot ((\mathbf{v} \cdot \nabla) \mathbf{m}) - \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}} \mathbf{v} \cdot \mathbf{n} \, dS \\ &\quad + \int_{\Omega} \frac{|\nabla \mathbf{m}|^2}{2} \nabla \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \right) \cdot \mathbf{v} + \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}} \operatorname{div} \mathbf{v} - \underbrace{\frac{\kappa(\mathbf{F})(\nabla \mathbf{m} \otimes \nabla \mathbf{m})}{\det \mathbf{F}} : \mathbf{e}(\mathbf{v})}_{= \mathbf{K} \text{ from (2.30b)}} \, d\mathbf{x}, \end{aligned} \quad (2.39)$$

where the boundary integral vanishes due to the boundary conditions $(\mathbf{n} \cdot \nabla) \mathbf{m} = \mathbf{0}$ and $\mathbf{v} \cdot \mathbf{n} = 0$. The latter equality in (2.39) follows by the calculus and the Green formula:

$$\begin{aligned} \int_{\Omega} \frac{\kappa(\mathbf{F}) \nabla^2 \mathbf{m}}{\det \mathbf{F}} : (\mathbf{v} \otimes \nabla \mathbf{m}) \, d\mathbf{x} &= \int_{\Gamma} \frac{\kappa(\mathbf{F})(\mathbf{n} \cdot \nabla) \mathbf{m}}{\det \mathbf{F}} \cdot ((\mathbf{v} \cdot \nabla) \mathbf{m}) \, dS \\ &\quad - \int_{\Omega} \frac{\kappa(\mathbf{F}) \nabla \mathbf{m} \otimes \mathbf{v}}{\det \mathbf{F}} : \nabla^2 \mathbf{m} + |\nabla \mathbf{m}|^2 \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \operatorname{div} \mathbf{v} + \nabla \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \right) \cdot \mathbf{v} \right) \, d\mathbf{x} \\ &= - \int_{\Omega} \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}} \operatorname{div} \mathbf{v} + \frac{|\nabla \mathbf{m}|^2}{2} \nabla \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \right) \cdot \mathbf{v} \, d\mathbf{x}, \end{aligned} \quad (2.40)$$

where again $(\mathbf{n} \cdot \nabla) \mathbf{m} = \mathbf{0}$ was used. For the last term, we can still use the calculus

$$\nabla \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \right) = \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \right)' : \nabla \mathbf{F} = \left(\frac{\kappa'(\mathbf{F})}{\det \mathbf{F}} - \frac{\kappa(\mathbf{F}) \operatorname{Cof} \mathbf{F}}{\det \mathbf{F}^2} \right) : \nabla \mathbf{F} = \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F}) \mathbf{F}^{-\top}}{\det \mathbf{F}} : \nabla \mathbf{F}, \quad (2.41)$$

where we again used the algebra $\mathbf{F}^{-\top} = \operatorname{Cof} \mathbf{F} / \det \mathbf{F}$ and the calculus $\det' = \operatorname{Cof}$. Thus (2.39) can be written as:

$$\begin{aligned} \int_{\Omega} \operatorname{div} \left(\frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \right) \cdot (\mathbf{v} \cdot \nabla) \mathbf{m} \, d\mathbf{x} &= \int_{\Omega} \left(|\nabla \mathbf{m}|^2 \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F}) \mathbf{F}^{-\top}}{2 \det \mathbf{F}} : (\mathbf{v} \cdot \nabla) \mathbf{F} \right. \\ &\quad \left. + \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}} \operatorname{div} \mathbf{v} - \mathbf{K} : \mathbf{e}(\mathbf{v}) \right) \, d\mathbf{x}, \end{aligned} \quad (2.42)$$

Similarly, this exchange-energy term $\text{div}(\kappa(\mathbf{F})\nabla\mathbf{m}/\det\mathbf{F})$ tested by $\frac{\partial}{\partial t}\mathbf{m}$ is to be handled by using Green's formula once:

$$\begin{aligned} \int_{\Omega} \text{div}\left(\frac{\kappa(\mathbf{F})\nabla\mathbf{m}}{\det\mathbf{F}}\right) \cdot \frac{\partial\mathbf{m}}{\partial t} d\mathbf{x} &= \int_{\Gamma} \frac{\kappa(\mathbf{F})(\mathbf{n}\cdot\nabla)\mathbf{m}}{\det\mathbf{F}} \cdot \frac{\partial\mathbf{m}}{\partial t} dS - \int_{\Omega} \frac{\kappa(\mathbf{F})\nabla\mathbf{m}}{\det\mathbf{F}} : \nabla \frac{\partial\mathbf{m}}{\partial t} d\mathbf{x} \\ &= -\frac{d}{dt} \int_{\Omega} \frac{\kappa(\mathbf{F})|\nabla\mathbf{m}|^2}{2\det\mathbf{F}} d\mathbf{x} + \int_{\Omega} \frac{|\nabla\mathbf{m}|^2}{2} \left(\frac{\kappa(\mathbf{F})}{\det\mathbf{F}}\right)' : \frac{\partial\mathbf{F}}{\partial t} d\mathbf{x} \\ &= -\frac{d}{dt} \int_{\Omega} \frac{\kappa(\mathbf{F})|\nabla\mathbf{m}|^2}{2\det\mathbf{F}} d\mathbf{x} + \int_{\Omega} |\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F})\mathbf{F}^{-\top}}{2\det\mathbf{F}} : \frac{\partial\mathbf{F}}{\partial t} d\mathbf{x}, \end{aligned} \quad (2.43)$$

where we again used $(\mathbf{n}\cdot\nabla)\mathbf{m} = \mathbf{0}$ on Γ and, where we again used, as in (2.41), that $(\kappa(\mathbf{F})/\det\mathbf{F})' = (\kappa'(\mathbf{F}) - \kappa(\mathbf{F})\mathbf{F}^{-\top})/\det\mathbf{F}$. To merge (2.42) and (2.43), we use (2.30c) and also the calculus

$$\begin{aligned} &|\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F})\mathbf{F}^{-\top}}{2\det\mathbf{F}} : \frac{\partial\mathbf{F}}{\partial t} + |\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F})\mathbf{F}^{-\top}}{2\det\mathbf{F}} : (\mathbf{v}\cdot\nabla)\mathbf{F} \\ &= |\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F}) - \kappa(\mathbf{F})\mathbf{F}^{-\top}}{2\det\mathbf{F}} : (\nabla\mathbf{v})\mathbf{F} = |\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F})\mathbf{F}^{\top} - \kappa(\mathbf{F})\mathbb{I}}{2\det\mathbf{F}} : \mathbf{e}(\mathbf{v}) \\ &= |\nabla\mathbf{m}|^2 \frac{\kappa'(\mathbf{F})\mathbf{F}^{\top}}{2\det\mathbf{F}} : \mathbf{e}(\mathbf{v}) - \frac{|\nabla\mathbf{m}|^2 \kappa(\mathbf{F})}{2\det\mathbf{F}} \text{div}\mathbf{v} \end{aligned} \quad (2.44)$$

when the frame indifference of κ is assumed. Noticing that the last term in (2.44) cancels with the same pressure term in (2.42), we obtain

$$\int_{\Omega} \text{div}\left(\frac{\kappa(\mathbf{F})\nabla\mathbf{m}}{\det\mathbf{F}}\right) \cdot \dot{\mathbf{m}} d\mathbf{x} = \int_{\Omega} \left(\frac{\kappa'(\mathbf{F})|\nabla\mathbf{m}|^2\mathbf{F}^{\top}}{2\det\mathbf{F}} - \mathbf{K}\right) : \mathbf{e}(\mathbf{v}) d\mathbf{x} - \frac{d}{dt} \int_{\Omega} \frac{\kappa(\mathbf{F})|\nabla\mathbf{m}|^2}{2\det\mathbf{F}} d\mathbf{x}. \quad (2.45)$$

Moreover, we use the Green theorem also for the driving magnetic field $\mathbf{h} = \mathbf{h}_{\text{ext}} - \mathbf{h}_{\text{dem}}$ with the demagnetizing field $\mathbf{h}_{\text{dem}} = -\nabla u$:

$$\begin{aligned} \int_{\Omega} \mu_0 \mathbf{h} \cdot \dot{\mathbf{m}} d\mathbf{x} &= \int_{\Omega} \mu_0 \mathbf{h}_{\text{ext}} \cdot \frac{\partial\mathbf{m}}{\partial t} + \mu_0 \mathbf{h} \cdot (\mathbf{v}\cdot\nabla)\mathbf{m} - \mu_0 \mathbf{h}_{\text{dem}} \cdot \frac{\partial\mathbf{m}}{\partial t} d\mathbf{x} \\ &= \int_{\Omega} \frac{\partial}{\partial t} (\mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m}) - \mu_0 \frac{\partial\mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m} - \mu_0 (\nabla\mathbf{h})^{\top} \mathbf{m} \cdot \mathbf{v} + \mu_0 \nabla(\mathbf{h} \cdot \mathbf{m}) \cdot \mathbf{v} - \mu_0 \mathbf{h}_{\text{dem}} \cdot \frac{\partial\mathbf{m}}{\partial t} d\mathbf{x}. \end{aligned}$$

Altogether, this is used to handle the right-hand side of (2.30d) tested by $\dot{\mathbf{m}}$:

$$\begin{aligned} &\int_{\Omega} \left(\mu_0 \mathbf{h} - \frac{\psi'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det\mathbf{F}} + \text{div}\left(\frac{\kappa(\mathbf{F})\nabla\mathbf{m}}{\det\mathbf{F}}\right)\right) \cdot \dot{\mathbf{m}} d\mathbf{x} \\ &= \int_{\Omega} \left(\mu_0 \mathbf{h} - \frac{\psi'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det\mathbf{F}} + \text{div}\left(\frac{\kappa(\mathbf{F})\nabla\mathbf{m}}{\det\mathbf{F}}\right)\right) \cdot \dot{\mathbf{m}} - (\mu_0 \mathbf{h} - \mathbf{t}) \cdot \text{skw}(\nabla\mathbf{v})\mathbf{m} d\mathbf{x} \\ &= \frac{d}{dt} \int_{\Omega} \mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m} - \frac{\kappa(\mathbf{F})|\nabla\mathbf{m}|^2}{2\det\mathbf{F}} d\mathbf{x} + \int_{\Omega} \left(\left(\frac{\kappa'(\mathbf{F})|\nabla\mathbf{m}|^2\mathbf{F}^{\top}}{2\det\mathbf{F}} - \mathbf{K}\right) : \mathbf{e}(\mathbf{v}) - \mu_0 \frac{\partial\mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m} \right. \\ &\quad \left. - \left(\frac{\varphi'_m(\mathbf{F}, \mathbf{m})}{\det\mathbf{F}} + \frac{\zeta'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det\mathbf{F}}\right) \cdot \dot{\mathbf{m}} - \underbrace{((\mu_0 (\nabla\mathbf{h})^{\top} \mathbf{m} - \mu_0 \nabla(\mathbf{h} \cdot \mathbf{m})) \cdot \mathbf{v})}_{\text{magnetic force in (2.30b)}} - \mu_0 \mathbf{h}_{\text{dem}} \cdot \frac{\partial\mathbf{m}}{\partial t} \right) d\mathbf{x} \end{aligned}$$

$$- \underbrace{\text{skw}((\mu_0 \mathbf{h} - \psi'_m(\mathbf{F}, \mathbf{m}, \theta)) \otimes \mathbf{m}) : \nabla \mathbf{v}}_{= \mathbf{S} \text{ from (2.30b)}} + \underbrace{\kappa(\mathbf{F}) \frac{\mathbf{m} \otimes \nabla \mathbf{m} - \nabla \mathbf{m} \otimes \mathbf{m}}{2 \det \mathbf{F}} : \nabla^2 \mathbf{v}}_{= \mathcal{S} \text{ from (2.30b)}} \, d\mathbf{x} \quad (2.46)$$

with \mathbf{t} and \mathbf{K} from (2.30b). Here the algebra $(\mu_0 \mathbf{h} - \psi'_m) \cdot \text{skw}(\nabla \mathbf{v}) \mathbf{m} = \text{skw}((\mu_0 \mathbf{h} - \psi'_m) \otimes \mathbf{m}) : \nabla \mathbf{v}$ has been used. Beside, to see the magnetic hyperstress \mathcal{S} , we used the calculus

$$\begin{aligned} & \int_{\Omega} \text{div} \left(\frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \nabla \mathbf{m} \right) \cdot \text{skw}(\nabla \mathbf{v}) \mathbf{m} \, d\mathbf{x} \\ &= \int_{\Gamma} \frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \underbrace{(\mathbf{n} \cdot \nabla) \mathbf{m}}_{= \mathbf{0}} \cdot \text{skw}(\nabla \mathbf{v}) \mathbf{m} \, dS - \int_{\Omega} \frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \nabla \mathbf{m} : \nabla (\text{skw}(\nabla \mathbf{v}) \mathbf{m}) \, d\mathbf{x} \\ &= - \int_{\Omega} \frac{\kappa(\mathbf{F})}{\det \mathbf{F}} \underbrace{(\nabla \mathbf{m} \otimes \nabla \mathbf{m}) : \text{skw}(\nabla \mathbf{v}) \mathbf{m}}_{= \mathbf{0}} + \kappa(\mathbf{F}) \frac{\nabla \mathbf{m} \otimes \mathbf{m} - \mathbf{m} \otimes \nabla \mathbf{m}}{2 \det \mathbf{F}} : \nabla^2 \mathbf{v} \, d\mathbf{x}. \end{aligned} \quad (2.47)$$

It remains to exploit (2.30e). Testing it by $\mu_0 \frac{\partial}{\partial t} u$, we use the calculus, including the Green theorem for the convective term, to obtain

$$\begin{aligned} 0 &= \int_{\mathbb{R}^d} \mu_0 \text{div}(\chi_{\Omega} \mathbf{m} - \nabla u) \frac{\partial u}{\partial t} \, d\mathbf{x} = \int_{\mathbb{R}^d} \mu_0 (\nabla u - \chi_{\Omega} \mathbf{m}) \cdot \nabla \frac{\partial u}{\partial t} \, d\mathbf{x} \\ &= \frac{d}{dt} \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u|^2 \, d\mathbf{x} + \int_{\Omega} \mu_0 \mathbf{m} \cdot \frac{\partial \mathbf{h}_{\text{dem}}}{\partial t} \, d\mathbf{x} \\ &= \frac{d}{dt} \left(\int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u|^2 + \int_{\Omega} \mu_0 \mathbf{h}_{\text{dem}} \cdot \mathbf{m} \, d\mathbf{x} \right) - \int_{\Omega} \mu_0 \frac{\partial \mathbf{m}}{\partial t} \cdot \mathbf{h}_{\text{dem}} \, d\mathbf{x}. \end{aligned} \quad (2.48)$$

This is to be subtracted from (2.46), giving cancellation of the terms $\pm \mu_0 \frac{\partial}{\partial t} \mathbf{m} \cdot \mathbf{h}_{\text{dem}}$ and merging $\mu_0 (\mathbf{h}_{\text{ext}} \cdot \mathbf{m}) - \mu_0 (\mathbf{h}_{\text{dem}} \cdot \mathbf{m}) = \mu_0 (\mathbf{h} \cdot \mathbf{m})$.

Summarizing the above calculations, we formulated:

Proposition 2.6 (Magneto-mechanical dissipation energy balance). *Any smooth solution of the system (2.30a–e) with the boundary conditions (2.32) satisfies the identity*

$$\begin{aligned} & \frac{d}{dt} \left(\underbrace{\int_{\Omega} \frac{\varrho}{2} |\mathbf{v}|^2}_{\text{kinetic energy}} + \underbrace{\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}}}_{\text{stored energy}} + \underbrace{\frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}}}_{\text{exchange energy}} - \underbrace{\mu_0 \mathbf{h} \cdot \mathbf{m}}_{\text{Zeeman energy}} \, d\mathbf{x} + \underbrace{\int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u|^2}_{\text{energy of de-magnetizing field}} \, d\mathbf{x} \right) \\ &+ \int_{\Omega} \underbrace{\xi(\mathbf{F}, \theta; e(\mathbf{v}), \dot{\mathbf{m}})}_{\text{dissipation rate from (2.30f)}} \, d\mathbf{x} + \int_{\Gamma} \underbrace{\nu_b |\mathbf{v}|^p}_{\text{boundary dissipation rate}} \, dS = \int_{\Omega} \left(\underbrace{\varrho \mathbf{g} \cdot \mathbf{v}}_{\text{power of gravity field}} - \underbrace{\mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}}_{\text{power of external field}} \right. \\ &\quad \left. - \underbrace{\frac{\zeta'_F(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} : e(\mathbf{v}) - \frac{\zeta'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \dot{\mathbf{m}}}_{\text{adiabatic effects}} \right) \, d\mathbf{x} + \int_{\Gamma} \underbrace{\mathbf{k} \cdot \mathbf{v}}_{\text{power of traction}} \, dS. \end{aligned} \quad (2.49)$$

When we add (2.30f) tested by 1, the adiabatic and the dissipative heat sources cancel with those in (2.49). Thus we obtain (at least formally):

Proposition 2.7 (*Total energy balance*). *Any smooth solution of the evolution boundary-value problem (2.30)–(2.32) satisfies the identity*

$$\begin{aligned}
& \frac{d}{dt} \left(\underbrace{\int_{\Omega} \frac{\rho}{2} |\mathbf{v}|^2}_{\text{kinetic energy}} + \underbrace{\frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}}}_{\text{stored energy}} + \underbrace{\frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{2 \det \mathbf{F}}}_{\text{exchange energy}} - \underbrace{\mu_0 \mathbf{h} \cdot \mathbf{m}}_{\text{Zeeman energy}} \, d\mathbf{x} + \underbrace{\omega(\mathbf{F}, \mathbf{m}, \theta)}_{\text{heat energy}} \, d\mathbf{x} \right. \\
& \left. + \int_{\mathbb{R}^d} \underbrace{\frac{\mu_0}{2} |\nabla u|^2}_{\text{energy of de-magnetizing field}} \, d\mathbf{x} \right) + \int_{\Gamma} \underbrace{\frac{\nu_b}{2} |\mathbf{v}|^p}_{\text{boundary heat production outflow}} \, dS = \int_{\Omega} \underbrace{\rho \mathbf{g} \cdot \mathbf{v}}_{\text{power of gravity field}} - \underbrace{\mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}}_{\text{power of external field}} \, d\mathbf{x} + \int_{\Gamma} \underbrace{\mathbf{k} \cdot \mathbf{v}}_{\text{power of traction}} + \underbrace{h(\theta)}_{\text{heat flux}} \, dS.
\end{aligned} \tag{2.50}$$

Another aspect important both thermodynamically and also for mathematical analysis is non-negativity of temperature, related with the 3rd law of thermodynamics. This will be demonstrated later when we will exploit some information about the quality of the velocity field extracted from (2.49), cf. (3.57) below.

Remark 2.8 (*Exchange hyper-stress*). In principle, to balance the energetics, the magnetic exchange driving force $\text{div}(\kappa(\mathbf{F}) \nabla \mathbf{m} / \det \mathbf{F})$ in (2.18c) may contribute either directly to the skew-symmetric magnetic stress \mathbf{S} by $\text{skw}(\text{div}(\frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}}) \otimes \mathbf{m})$, as considered in [48], or to the skew-symmetric hyperstress as \mathcal{S} . Physically it is rather questionable which option is more relevant. The former case would bring analytical troubles in the argumentation (3.52) below due to lack of compactness of $\nabla \mathbf{v}$ as $\frac{\partial}{\partial t} \mathbf{v}$ is not estimated, in contrast to [48] where the inertial force was handled in a simplified “semi-compressible” way. This have led us to adopt the latter option here, which seems also more physical and a similar skew-symmetric hyperstress can be found in [57].

Remark 2.9 (*Isotropic magnets*). Let us note that, when $\psi(\mathbf{F}, \cdot, \theta)$ is isotropic as in Example 2.5, the skew-symmetric magnetic stress \mathbf{S} simplifies to $\mu_0 \text{skw}(\mathbf{h} \otimes \mathbf{m})$ because $\psi'_m(\mathbf{F}, \mathbf{m}, \theta) = k(\mathbf{F}, \mathbf{m}, \theta) \mathbf{m}$ for some scalar-valued coefficient $k = k(\mathbf{F}, \mathbf{m}, \theta)$ so that $\text{skw}(\psi'_m(\mathbf{F}, \mathbf{m}, \theta) \otimes \mathbf{m}) = k(\mathbf{F}, \mathbf{m}, \theta) \text{skw}(\mathbf{m} \otimes \mathbf{m}) = \mathbf{0}$.

3 The analysis – weak solutions of (2.30)

We will provide a proof of existence and certain regularity of weak solutions. To this aim, the concept of multipolar viscosity is essential but, anyhow, still quite nontrivial and carefully ordered arguments will be needed. The peculiarities are that the inertial term in Eulerian setting involves varying mass density requiring sophisticated techniques from compressible fluid dynamics, the momentum equation is very geometrically nonlinear, and the heat equation has an L^1 -structure with \mathbf{F} -dependent heat capacity and with the convective time derivative and ever-troubling adiabatic effects due to necessarily general coupling of mechanical and thermal effect in the deforming configuration in compressible media.

Usual analysis is made by some approximation, a-priori estimates, and limit passage possibly in several steps. The mentioned strong nonlinearity makes time discretization problematic. On the other hand, the space discretization by a (conformal) Faedo-Galerkin method

is also not straightforward because of several “nonlinear” tests leading to the basic energy balances in Section 2.4, being confronted in particular with the Lavrentiev phenomenon as occurring already in static nonlinear elasticity [1, 2, 19]. Anyhow, careful suitably regularized “semi-Galerkin” discretization allowing estimation of the magneto-mechanical part separately from the thermal part and a successive limit passage will work.

3.1 Definition of weak solutions and the main results

We will use the standard notation concerning the Lebesgue and the Sobolev spaces, namely $L^p(\Omega; \mathbb{R}^n)$ for Lebesgue measurable functions $\Omega \rightarrow \mathbb{R}^n$ whose Euclidean norm is integrable with p -power, and $W^{k,p}(\Omega; \mathbb{R}^n)$ for functions from $L^p(\Omega; \mathbb{R}^n)$ whose all derivative up to the order k have their Euclidean norm integrable with p -power. We also write briefly $H^k = W^{k,2}$. The notation p^* will denote the exponent from the embedding $W^{1,p}(\Omega) \subset L^{p^*}(\Omega)$, i.e. $p^* = dp/(d-p)$ for $p < d$ while $p^* \geq 1$ arbitrary for $p = d$ or $p^* = +\infty$ for $p > d$. Moreover, for a Banach space X and for $I = [0, T]$, we will use the notation $L^p(I; X)$ for the Bochner space of Bochner measurable functions $I \rightarrow X$ whose norm is in $L^p(I)$ while $W^{1,p}(I; X)$ stands for functions $I \rightarrow X$ whose distributional derivative is in $L^p(I; X)$. Also, $C(\cdot)$ and $C^1(\cdot)$ will denote spaces of continuous and continuously differentiable functions.

Moreover, as usual, we will use C for a generic constant which may vary from estimate to estimate.

We will consider an initial-value problem, prescribing the initial conditions

$$\varrho(0) = \varrho_0, \quad \mathbf{v}(0) = \mathbf{v}_0, \quad \mathbf{F}(0) = \mathbf{F}_0, \quad \mathbf{m}(0) = \mathbf{m}_0, \quad \text{and} \quad \theta(0) = \theta_0; \quad (3.1)$$

here and in what follows, we will use the short-hand notation as $[\varrho(t)](\mathbf{x}) = \varrho(t, \mathbf{x})$. Referring to the referential mass density ρ , the initial conditions should satisfy $\varrho_0 = \rho/\det \mathbf{F}_0$. To devise a weak formulation of the initial-boundary-value problem (2.32) and (3.1) for the system (2.30), we use the by-part integration in time and the Green formula for the inertial force. The nonsmoothness of $\text{Dir}(\cdot)$ applied on $\mathring{\mathbf{m}}$ leads to a variational inequality, arising by a standard definition of the convex subdifferential of the convex potential of the monotone set-valued mapping $\mathbf{r} \mapsto \tau \mathbf{r} + h_c(\mathbf{F}, \theta) \text{Dir}(\mathbf{r})$, let us denote it as $D(\mathbf{F}, \theta; \mathbf{r}) = \tau |\mathbf{r}|^2/2 + h_c(\mathbf{F}, \theta) |\mathbf{r}|$. Then (2.30d) has the form $\partial_{\mathring{\mathbf{m}}} D(\mathbf{F}, \theta; \mathring{\mathbf{m}}) \ni \mu_0 \mathbf{h}_{\text{ext}} + \mu_0 \nabla u - \mathbf{t} + \mathbf{m} \times \mathring{\mathbf{m}}/\gamma(\mathbf{F}, \mathbf{m}, \theta)$ with \mathbf{t} from (2.18c), from which we obtain a variational inequality by taking into account the standard definition of the (partial) convex subdifferential $\partial_{\mathring{\mathbf{m}}}$. This involves $\mathbf{t} \cdot \mathring{\mathbf{m}}$ which contains the product of $\text{div}(\kappa'_{\nabla \mathbf{m}}(\mathbf{F}, \nabla \mathbf{m})/\det \mathbf{F})$ with $\mathring{\mathbf{m}}$. This product would cause troubles in convergence of approximate solutions, so we will better avoid it in the weak formulation by a substitution using (2.45) integrated over I , i.e.

$$\begin{aligned} \int_0^T \int_{\Omega} \text{div} \left(\frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \right) \cdot \mathring{\mathbf{m}} \, d\mathbf{x} dt &= \int_{\Omega} \frac{\kappa(\mathbf{F}_0) |\nabla \mathbf{m}_0|^2}{2 \det \mathbf{F}_0} - \frac{\kappa(\mathbf{F}(T)) |\nabla \mathbf{m}(T)|^2}{2 \det \mathbf{F}(T)} \, d\mathbf{x} \\ &+ \int_0^T \int_{\Omega} \left(\frac{\kappa'(\mathbf{F}) |\nabla \mathbf{m}|^2 \mathbf{F}^\top}{2 \det \mathbf{F}} - \mathbf{K} \right) : \mathbf{e}(\mathbf{v}) \, d\mathbf{x} dt. \end{aligned} \quad (3.2)$$

Also we use the orthogonality $(\mathbf{m} \times \mathring{\mathbf{m}}) \cdot \mathring{\mathbf{m}} = 0$, which eliminates this (otherwise not integrable) term and which altogether gives the variational inequality (3.4b) below.

Definition 3.1 (Weak solutions to (2.30)). *For $p \in [1, \infty)$, a six-tuple $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, u, \theta)$ with $\varrho \in H^1(I \times \Omega)$, $\mathbf{v} \in L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))$, $\mathbf{F} \in H^1(I \times \Omega; \mathbb{R}^{d \times d})$, $\mathbf{m} \in H^1(I; L^2(\Omega; \mathbb{R}^d)) \cap L^\infty(I; H^1(\Omega; \mathbb{R}^d))$, $u \in L^\infty(I; H^1(\mathbb{R}^d))$, and $\theta \in L^1(I; W^{1,1}(\Omega))$ will be called a weak solution to the system (2.30) with the boundary conditions (2.32) and the initial condition (3.1) if*

$$\frac{\psi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} \in L^1(I \times \Omega; \mathbb{R}^{d \times d}), \quad (3.3a)$$

$$\frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} \in L^{q'}(I \times \Omega; \mathbb{R}^{d \times d}), \quad (3.3b)$$

$$\operatorname{div} \left(\frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{\det \mathbf{F}} \right) \in L^2(I \times \Omega; \mathbb{R}^d) \quad (3.3c)$$

with $\det \mathbf{F} > 0$ a.e. such that the integral identities

$$\begin{aligned} & \int_0^T \int_\Omega \left(\left(\frac{\psi'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa'(\mathbf{F}) \mathbf{F}^\top}{2 \det \mathbf{F}} + \mathbf{K} + \nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \varrho \mathbf{v} \otimes \mathbf{v} \right) : \mathbf{e}(\tilde{\mathbf{v}}) \right. \\ & + \mathbf{S} : \operatorname{skw}(\nabla \tilde{\mathbf{v}}) + \mu_0 (\mathbf{h} \cdot \mathbf{m}) \operatorname{div} \tilde{\mathbf{v}} - \mu_0 (\nabla \mathbf{h})^\top : (\mathbf{m} \otimes \tilde{\mathbf{v}}) + (\nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v} + \mathcal{S}) : \nabla^2 \tilde{\mathbf{v}} \\ & \left. - \varrho \mathbf{v} \cdot \frac{\partial \tilde{\mathbf{v}}}{\partial t} \right) d\mathbf{x} dt = \int_0^T \int_\Omega \varrho \mathbf{g} \cdot \tilde{\mathbf{v}} d\mathbf{x} dt + \int_0^T \int_\Gamma (\mathbf{k} - \nu_b |\mathbf{v}|^{p-2} \mathbf{v}) \cdot \tilde{\mathbf{v}} dS dt + \int_\Omega \varrho_0 \mathbf{v}_0 \cdot \tilde{\mathbf{v}}(0) d\mathbf{x} \end{aligned} \quad (3.4a)$$

with $\mathbf{h} = \mathbf{h}_{\text{ext}} + \nabla u$, \mathbf{K} , \mathbf{S} , and \mathcal{S} from (2.30b) holds for any $\tilde{\mathbf{v}}$ smooth with $\tilde{\mathbf{v}} \cdot \mathbf{n} = \mathbf{0}$ and $\tilde{\mathbf{v}}(T) = \mathbf{0}$, and

$$\begin{aligned} & \int_0^T \int_\Omega \left(\frac{\tau}{2} |\tilde{\mathbf{r}}|^2 + h_c(\mathbf{F}, \theta) |\tilde{\mathbf{r}}| - \operatorname{div} \frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \cdot \operatorname{skw}(\nabla \mathbf{v}) \mathbf{m} - \left(\frac{\kappa'(\mathbf{F}) |\nabla \mathbf{m}|^2 \mathbf{F}^\top}{2 \det \mathbf{F}} - \mathbf{K} \right) : \mathbf{e}(\mathbf{v}) \right. \\ & \left. + \frac{\kappa(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} : \nabla \tilde{\mathbf{r}} - \left(\mu_0 \mathbf{h} - \frac{\psi'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right) \cdot (\tilde{\mathbf{r}} - \mathring{\mathbf{m}}) + \frac{\mathbf{m} \times \mathring{\mathbf{m}}}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} \cdot \tilde{\mathbf{r}} \right) d\mathbf{x} dt \\ & \geq \int_0^T \int_\Omega \frac{\tau}{2} |\mathring{\mathbf{m}}|^2 + h_c(\mathbf{F}, \theta) |\mathring{\mathbf{m}}| d\mathbf{x} dt + \int_\Omega \frac{\kappa(\mathbf{F}(T)) |\nabla \mathbf{m}(T)|^2}{2 \det \mathbf{F}(T)} - \frac{\kappa(\mathbf{F}_0) |\nabla \mathbf{m}_0|^2}{2 \det \mathbf{F}_0} d\mathbf{x} \end{aligned} \quad (3.4b)$$

for any $\tilde{\mathbf{r}} \in L^2(I; H^1(\Omega; \mathbb{R}^d))$, and further

$$\int_{\mathbb{R}^3} \nabla u(t) \cdot \nabla \tilde{u} d\mathbf{x} = \int_\Omega \mathbf{m}(t) \cdot \nabla \tilde{u} d\mathbf{x} \quad (3.4c)$$

holds for any $\tilde{u} \in H^1(\mathbb{R}^d)$ and for a.a. $t \in I$, and

$$\begin{aligned} & \int_0^T \int_\Omega \left(\omega(\mathbf{F}, \mathbf{m}, \theta) \frac{\partial \tilde{\theta}}{\partial t} + (\omega(\mathbf{F}, \mathbf{m}, \theta) \mathbf{v} - \mathcal{K}(\mathbf{F}, \theta) \nabla \theta) \cdot \nabla \tilde{\theta} \right. \\ & \left. + \left(\xi(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \mathring{\mathbf{m}}) + \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} : \mathbf{e}(\mathbf{v}) + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \mathring{\mathbf{m}} \right) \tilde{\theta} \right) d\mathbf{x} dt \\ & + \int_0^T \int_\Gamma \left(h(\theta) + \frac{\nu_b}{2} |\mathbf{v}|^p \right) \tilde{\theta} dS dt + \int_\Omega \omega(\mathbf{F}_0, \theta_0) \tilde{\theta}(0) d\mathbf{x} = 0 \end{aligned} \quad (3.4d)$$

with $\xi(\mathbf{F}, \theta; \cdot, \cdot)$ and $\omega(\cdot, \cdot, \cdot)$ from (2.30f) holds for any $\tilde{\theta}$ smooth with $\tilde{\theta}(T) = 0$, and the equations (2.30a) and (2.30c) hold a.e. on $I \times \Omega$ with $\mathbf{v}(0) = \mathbf{v}_0$ and $\mathbf{F}(0) = \mathbf{F}_0$ a.e. on Ω , and also $\mathbf{m}(0) = \mathbf{m}_0$ is to hold a.e. on Ω .

Before stating the main analytical result, let us summarize the data qualification which will be fitted to the motivating Example 2.5. For some $\delta > 0$ and $s > 0$, we assume:

$$\Omega \text{ a smooth bounded domain of } \mathbb{R}^d, \quad d = 2, 3, \quad (3.5a)$$

$$\varphi \in C^1(\text{GL}^+(d) \times \mathbb{R}^d), \quad \forall \mathbf{F} \in \text{GL}^+(d): \quad \varphi(\mathbf{F}, \mathbf{m}) \geq \delta(1 + |\mathbf{m}|^s \det \mathbf{F}),$$

$$\exists C \in C(\text{GL}^+(d)), \quad \forall \mathbf{F} \in \text{GL}^+(d), \quad \mathbf{m} \in \mathbb{R}^d: \quad |\varphi'_m(\mathbf{F}, \mathbf{m})| \leq C(\mathbf{F})(1 + |\mathbf{m}|^{1+2^*/2}), \quad (3.5b)$$

$$\zeta \in C^2(\text{GL}^+(d) \times \mathbb{R}^d \times \mathbb{R}^+), \quad \forall (\mathbf{F}, \mathbf{m}, \theta) \in \text{GL}^+(d) \times \mathbb{R}^d \times \mathbb{R}^+: \quad \zeta''_{\theta\theta}(\mathbf{F}, \mathbf{m}, \theta) \leq \frac{-\delta}{\det \mathbf{F}},$$

$$\left| \frac{\zeta'_F(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} \right| + \left| \frac{\zeta'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right|^2 \leq C \left(1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right), \quad (3.5c)$$

$$\forall K \subset \text{GL}^+(d) \text{ compact } \exists C_K < \infty \quad \forall (\mathbf{F}, \mathbf{m}, \theta) \in K \times \mathbb{R}^d \times \mathbb{R}^+: \quad |\omega'_F(\mathbf{F}, \mathbf{m}, \theta)| \leq C_K(1 + \theta),$$

$$\omega'_\theta(\mathbf{F}, \mathbf{m}, \theta) + |\omega'_m(\mathbf{F}, \mathbf{m}, \theta)| + |\omega''_{F\theta}(\mathbf{F}, \mathbf{m}, \theta)| + |\omega''_{m\theta}(\mathbf{F}, \mathbf{m}, \theta)| \leq C_K, \quad (3.5d)$$

$$\nu_1 > 0, \quad \nu_2 > 0, \quad \nu_b > 0, \quad (3.5e)$$

$$\gamma \in C(\text{GL}^+(d) \times \mathbb{R}^d \times \mathbb{R}^+) \text{ positive, } \forall K \subset \text{GL}^+(d) \text{ compact} \\ \exists C_K < \infty \quad \forall (\mathbf{F}, \mathbf{m}, \theta) \in K \times \mathbb{R}^d \times \mathbb{R}^+: \quad \frac{|\mathbf{m}|}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} \leq C_K, \quad (3.5f)$$

$$\kappa \in C^1(\text{GL}^+(d)), \quad \inf_{\mathbf{F} \in \text{GL}^+(d)} \kappa(\mathbf{F}) > 0, \quad (3.5g)$$

$$\mathcal{K} \in C(\text{GL}^+(d) \times \mathbb{R}^+) \text{ bounded, } \inf_{\mathbf{F} \in \text{GL}^+(d), \theta \in \mathbb{R}^+} \mathcal{K}(\mathbf{F}, \theta) > 0, \quad (3.5h)$$

$$h : I \times \Gamma \times \mathbb{R}^+ \rightarrow \mathbb{R} \text{ Carathéodory function, } 0 \leq \theta h(t, \mathbf{x}, \theta) \leq C(1 + \theta^2) \text{ and} \\ h(t, \mathbf{x}, \theta) \leq h_{\max}(t, \mathbf{x}) \text{ for some } h_{\max} \in L^1(I \times \Gamma), \quad (3.5i)$$

$$\mathbf{g} \in L^1(I; L^\infty(\Omega; \mathbb{R}^d)), \quad \mathbf{h}_{\text{ext}} \in W^{1,1}(I; L^{s'}(\Omega; \mathbb{R}^d)), \quad \mathbf{k} \in L^2(I \times \Gamma; \mathbb{R}^d), \quad \mathbf{k} \cdot \mathbf{n} = 0, \quad (3.5j)$$

$$\mathbf{v}_0 \in L^2(\Omega; \mathbb{R}^d), \quad \mathbf{F}_0 \in W^{1,r}(\Omega; \mathbb{R}^{d \times d}), \quad r > d, \quad \text{with } \min_{\bar{\Omega}} \det \mathbf{F}_0 > 0, \quad (3.5k)$$

$$\rho \in L^\infty(\Omega) \cap W^{1,r}(\Omega), \quad r > d, \quad \text{with } \min_{\bar{\Omega}} \rho > 0, \quad (3.5l)$$

$$\mathbf{m}_0 \in H^1(\Omega; \mathbb{R}^d), \quad \theta_0 \in L^1(\Omega), \quad \theta_0 \geq 0 \text{ a.e. on } \Omega, \quad (3.5m)$$

where ω in (3.5d) is from (2.30f). Let us note that the first condition in (3.5c) is just a condition on the heat capacity $c = c(\mathbf{F}, \mathbf{m}, \theta) = \omega'_\theta(\mathbf{F}, \mathbf{m}, \theta)$ and implies the coercivity $\omega(\mathbf{F}, \mathbf{m}, \theta) \geq \delta\theta/\det \mathbf{F}$ since $\omega(\mathbf{F}, \mathbf{m}, 0) = 0$. One should note that the referential stored energy (in contrast to the actual stored energy) enters the model only through its derivatives and can be modified without loss of generality by adding a constant, so that (3.5b) could be understood simply as coercivity $\varphi(\mathbf{F}, \mathbf{m})/\det \mathbf{F} \geq \delta|\mathbf{m}|^s$. Independently, the natural blow-up under compression, i.e. $\varphi(\mathbf{F}, \mathbf{m}) \rightarrow \infty$ if $\det \mathbf{F} \rightarrow 0+$, is allowed in (3.5b). The condition (3.5i) is well fitted with the standard situation that the boundary flux is $h(\theta) = f(\theta_{\text{ext}}) - f(\theta)$ with an increasing function f and with $\theta_{\text{ext}} \geq 0$ a prescribed external temperature, so that one can choose $h_{\max} = f(\theta_{\text{ext}})$ provided we prove that $\theta \geq 0$. Also the condition $\theta h(t, \mathbf{x}, \theta) \leq C(1 + \theta^2)$ is well compatible with this ansatz provided $f(0) = 0$ and $f(\theta_{\text{ext}}) \in L^2(I \times \Gamma)$.

Theorem 3.2 (Existence and regularity of weak solutions). *Let $p > d$ and $s \geq 2p/(p-2)$, and the assumptions (2.19) and (3.5) hold. Then:*

- (i) there exist a weak solution $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, u, \theta)$ according Definition 3.1 with a non-negative mass density $\varrho \in L^\infty(I; W^{1,r}(\Omega))$ such that $\frac{\partial}{\partial t} \varrho \in L^\sigma(I; L^{r\sigma/(r+\sigma)}(\Omega))$ with $3 \leq \sigma < p(pd+4p-2d)/(4p-2d)$, and a non-negative temperature $\theta \in L^\infty(I; L^1(\Omega)) \cap L^\mu(I; W^{1,\mu}(\Omega))$ with $1 \leq \mu < (d+2)/(d+1)$, and further $\frac{\partial}{\partial t} \mathbf{F} \in L^p(I; L^r(\Omega; \mathbb{R}^{d \times d}))$ and $\nabla \mathbf{F} \in L^\infty(I; L^r(\Omega; \mathbb{R}^{d \times d \times d}))$, and $\frac{\partial}{\partial t} \mathbf{m} \in L^2(I \times \Omega; \mathbb{R}^d)$ and $\nabla \mathbf{m} \in L^\infty(I; L^2(\Omega; \mathbb{R}^d))$, and $\nabla^2 \mathbf{m} \in L^2(I \times \Omega; \mathbb{R}^{d \times d \times d})$.
- (ii) Moreover, this solution complies with energetics in the sense that the energy dissipation balance (2.49) as well as the total energy balance (2.50) integrated over time interval $[0, t]$ with the initial conditions (3.1) hold.

3.2 Some auxiliary results and formal a-priori estimates

Let us first formulate two auxiliary assertions:

Lemma 3.3 (See [49]). *Given $\mathbf{v} \in L^1(I; W^{2,p}(\Omega; \mathbb{R}^d))$ with $p > d$ and $\mathbf{v} \cdot \mathbf{n} = 0$ on $I \times \Gamma$ and $\varrho_0 \in W^{1,r}(\Omega)$, (2.30a) has a unique weak solution $\varrho \in C_w(I; W^{1,r}(\Omega)) \cap W^{1,1}(I; L^r(\Omega))$ which satisfies it a.e. on $I \times \Omega$ and also the estimate holds:*

$$\|\varrho\|_{L^\infty(I; W^{1,r}(\Omega)) \cap W^{1,1}(I; L^r(\Omega))} \leq \mathfrak{C} \left(\|\nabla \mathbf{v}\|_{L^1(I; W^{1,p}(\Omega; \mathbb{R}^{d \times d}))}, \|\varrho_0\|_{W^{1,r}(\Omega)} \right) \quad (3.6)$$

holds with some $\mathfrak{C} \in C(\mathbb{R}^2)$. Moreover, the mapping

$$\mathbf{v} \mapsto \varrho : L^1(I; W^{2,p}(\Omega; \mathbb{R}^d)) \rightarrow L^\infty(I; W^{1,r}(\Omega)) \quad (3.7)$$

is (weak, weak*)-continuous. The analogous assertion holds for (2.10), assuming $1/\det \mathbf{F}_0 \in W^{1,r}(\Omega)$, and for (2.13), assuming $1/\varrho_0 \in W^{1,r}(\Omega)$. Eventually, it holds $\mathbb{R}^{d \times d}$ -valued also for (2.30c), assuming $\mathbf{F}_0 \in W^{1,r}(\Omega; \mathbb{R}^{d \times d})$.

For the approximation method in the proof below, we will still need a modification of Lemma 3.3 for a non-homogeneous evolution-and-transport equation (3.25e), whose proof is a straightforward modification (partly simplification) of [49, Sect.4]:

Lemma 3.4. *Given $\mathbf{v} \in L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))$ with $p > d$ and $\mathbf{r} \in L^1(I; L^2(\Omega; \mathbb{R}^d))$, the equation $\frac{\partial}{\partial t} \mathbf{m} + (\mathbf{v} \cdot \nabla) \mathbf{m} - \text{skw}(\nabla \mathbf{v}) \mathbf{m} = \mathbf{r}$ with the initial condition $\mathbf{m}_0 \in L^2(\Omega; \mathbb{R}^d)$ and has a unique weak solution $\mathbf{m} \in C_w(I; L^2(\Omega; \mathbb{R}^d)) \cap W^{1,1}(I; H^1(\Omega; \mathbb{R}^d)^*)$ and also the estimate holds:*

$$\|\mathbf{m}\|_{L^\infty(I; L^2(\Omega)) \cap W^{1,1}(I; H^1(\Omega; \mathbb{R}^d)^*)} \leq \mathfrak{C} \left(\|\nabla \mathbf{v}\|_{L^1(I; W^{1,p}(\Omega; \mathbb{R}^{d \times d}))}, \|\mathbf{m}_0\|_{L^2(\Omega; \mathbb{R}^d)}, \|\mathbf{r}\|_{L^1(I; L^2(\Omega; \mathbb{R}^d))} \right) \quad (3.8)$$

holds with some $\mathfrak{C} \in C(\mathbb{R}^3)$. Moreover, the mapping

$$(\mathbf{v}, \mathbf{r}) \mapsto \mathbf{m} : L^1(I; W^{2,p}(\Omega; \mathbb{R}^d)) \times L^1(I; L^2(\Omega; \mathbb{R}^d)) \rightarrow L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \quad (3.9)$$

is (weak, weak*)-continuous.

Formally, the assumptions (3.5) yield some a-priori bounds which can be obtained from the total energy balance (2.50) and the mechanical energy-dissipation balance (2.49) for any sufficiently regular solution $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, u, \theta)$ with $\theta \geq 0$ a.e. in $I \times \Omega$. Later, we will prove existence of such solutions, but unfortunately we are not able to claim that every weak solution has θ non-negative. For the approximation method used in the proof below, we assume the data ψ , \mathcal{K} , and h to be defined also for the negative temperature by extending them as

$$\begin{aligned}\psi(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}, \theta) &:= \varphi(\mathbf{F}, \mathbf{m}) + \theta(\ln(-\theta) - 1) + \kappa(\mathbf{F})|\nabla \mathbf{m}|^2/2, \\ \mathcal{K}(\mathbf{F}, \theta) &:= \mathcal{K}(\mathbf{F}, -\theta), \quad \text{and} \quad h(t, \mathbf{x}, \theta) := h(t, \mathbf{x}, -\theta) \quad \text{for } \theta < 0\end{aligned}\quad (3.10)$$

with φ and ζ from the split (2.17). This definition makes $\psi : \text{GL}^+(d) \times \mathbb{R}^d \times \mathbb{R}^{d \times d} \times \mathbb{R} \rightarrow \mathbb{R}$ continuous and implies that $\omega(\mathbf{F}, \mathbf{m}, \cdot)$ as well as $\zeta_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \cdot)$ and $\zeta_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \cdot)$ continuous; note that $\omega(\mathbf{F}, \mathbf{m}, \theta) = \theta/\det \mathbf{F}$, $\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) = \mathbf{0}$, and $\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) = \mathbf{0}$ for θ negative.

First, we use the total energy balance (2.50) integrated over a time interval $[0, t]$. At this point, we must now assume (while being later proved at least for some solution) that $\theta \geq 0$, and similarly we now assume $\det \mathbf{F} > 0$. In particular, we have also $\omega(\mathbf{F}, \mathbf{m}, \theta) \geq 0$ and thus we are “only” to estimate the right-hand side in (2.50) together with the Zeeman energy. For the bulk term $\varrho \mathbf{g} \cdot \mathbf{v}$ and the boundary terms $\mathbf{k} \cdot \mathbf{v} + h(\theta)$ we refer to [49]. The gravity force $\varrho \mathbf{g}$ tested by the velocity \mathbf{v} can be estimated by the Hölder/Young inequality as

$$\begin{aligned}\int_{\Omega} \varrho \mathbf{g} \cdot \mathbf{v} \, d\mathbf{x} &= \int_{\Omega} \sqrt{\frac{\varrho}{\det \mathbf{F}}} \sqrt{\varrho} \mathbf{v} \cdot \mathbf{g} \, d\mathbf{x} \leq \left\| \sqrt{\frac{\varrho}{\det \mathbf{F}}} \right\|_{L^2(\Omega)} \|\sqrt{\varrho} \mathbf{v}\|_{L^2(\Omega; \mathbb{R}^d)} \|\mathbf{g}\|_{L^\infty(\Omega; \mathbb{R}^d)} \\ &\leq \frac{1}{2} \left(\left\| \sqrt{\frac{\varrho}{\det \mathbf{F}}} \right\|_{L^2(\Omega)}^2 + \|\sqrt{\varrho} \mathbf{v}\|_{L^2(\Omega; \mathbb{R}^d)}^2 \right) \|\mathbf{g}\|_{L^\infty(\Omega; \mathbb{R}^d)} \\ &= \|\mathbf{g}\|_{L^\infty(\Omega; \mathbb{R}^d)} \int_{\Omega} \frac{\varrho}{2 \det \mathbf{F}} + \frac{\varrho}{2} |\mathbf{v}|^2 \, d\mathbf{x} \\ &\leq \|\mathbf{g}\|_{L^\infty(\Omega; \mathbb{R}^d)} \left(\frac{\max \varrho(\bar{\Omega})}{2 \inf \varphi(\text{GL}^+(d) \times \mathbb{R}^d)} \int_{\Omega} \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \, d\mathbf{x} + \int_{\Omega} \frac{\varrho}{2} |\mathbf{v}|^2 \, d\mathbf{x} \right). \quad (3.11)\end{aligned}$$

The integral on the right-hand side of (3.11) can then be treated by the Gronwall lemma. In order to apply the Gronwall lemma one needs the qualification (3.5j) for \mathbf{g} . The boundary terms in (2.50) can be estimated, at current time instant $t \in I$, as

$$\int_{\Gamma} \mathbf{k} \cdot \mathbf{v} + h(\theta) \, dS \leq \left(\frac{2}{\nu_b} \right)^{1/(p-1)} \|\mathbf{k}\|_{L^{p'}(\Gamma; \mathbb{R}^d)}^{p'} + \frac{\nu_b}{2p} \|\mathbf{v}\|_{L^p(\Gamma; \mathbb{R}^d)}^p + \|h_{\max}\|_{L^1(\Gamma)} \quad (3.12)$$

with the term $\nu_b \|\mathbf{v}\|_{L^p(\Gamma; \mathbb{R}^d)}^p / (2p)$ to be absorbed in the left hand side of (2.50); here we used the modelling assumption that part of the heat produced by the boundary viscosity leaves the system, otherwise we would have to confine ourselves to $\mathbf{f} = 0$. Then, integrating (2.50) in time, one can use the qualification of \mathbf{k} in (3.5j). The Zeeman energy $\mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m}$ and the power of external field $\mu_0 \frac{\partial}{\partial t} \mathbf{h}_{\text{ext}} \cdot \mathbf{m}$ are to be estimated as

$$\int_{\Omega} \mu_0 \mathbf{h}_{\text{ext}}(t) \cdot \mathbf{m}(t) - \mathbf{h}_{\text{ext}}(0) \cdot \mathbf{m}_0 \, d\mathbf{x} - \int_0^t \int_{\Omega} \mu_0 \frac{\partial}{\partial t} \mathbf{h}_{\text{ext}} \cdot \mathbf{m} \, d\mathbf{x} \, dt$$

$$\begin{aligned}
&\leq C_{\mu_0, \delta} \|\mathbf{h}_{\text{ext}}(t)\|_{L^{s'}(\Omega; \mathbb{R}^d)}^{s'} + \frac{\delta}{2} \|\mathbf{m}(t)\|_{L^s(\Omega; \mathbb{R}^d)}^s \\
&\quad + \mu_0 \int_0^t \left\| \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \right\|_{L^{s'}(\Omega; \mathbb{R}^d)} (1 + \|\mathbf{m}\|_{L^s(\Omega; \mathbb{R}^d)}^s) dt + \mu_0 \|\mathbf{h}_{\text{ext}}(0) \cdot \mathbf{m}_0\|_{L^1(\Omega)}
\end{aligned} \tag{3.13}$$

with some $C_{\mu_0, \delta}$ depending on μ_0 and δ chosen according to the assumption (3.5b). This assumption is then to be exploited for the stored energy on the left-hand side of (2.50) and, together with the qualification (3.5j) of \mathbf{h}_{ext} , used for the Gronwall inequality. As a result, since $\det \mathbf{F} > 0$, we obtain the (formal) a-priori estimates

$$\|\sqrt{\varrho} \mathbf{v}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d))} \leq C, \tag{3.14a}$$

$$\left\| \frac{\varphi(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} \right\|_{L^\infty(I; L^1(\Omega))} \leq C, \tag{3.14b}$$

$$\left\| \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{\det \mathbf{F}} \right\|_{L^\infty(I; L^1(\Omega))} \leq C, \tag{3.14c}$$

$$\|\nabla u\|_{L^\infty(I; L^2(\mathbb{R}^d; \mathbb{R}^d))} \leq C, \quad \text{and} \tag{3.14d}$$

$$\left\| \frac{\theta}{\det \mathbf{F}} \right\|_{L^\infty(I; L^1(\Omega))} \leq C. \tag{3.14e}$$

From (3.14b), using (3.5b), we also obtain

$$\|\mathbf{m}\|_{L^\infty(I; L^s(\Omega; \mathbb{R}^d))} \leq C. \tag{3.14f}$$

Now we come to (2.49); here we used the assumed frame indifference of $\zeta(\cdot, \mathbf{m}, \theta)$ so that $\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top$ is symmetric and thus $\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top : \nabla \mathbf{v} = \zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top : \mathbf{e}(\mathbf{v})$. The issue is now estimation of the adiabatic term in (2.49). Then, while relying on $p > d$, using the embedding

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega; \mathbb{R}^{d \times d})}^p \leq K \left(\|\nabla^2 \mathbf{v}\|_{L^p(\Omega; \mathbb{R}^{d \times d \times d})}^p + \|\mathbf{v}|_\Gamma\|_{L^p(\Gamma; \mathbb{R}^d)}^p \right), \tag{3.15}$$

we estimate

$$\begin{aligned}
&\int_\Omega \left| \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} : \mathbf{e}(\mathbf{v}) + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \cdot \dot{\mathbf{m}} \right| d\mathbf{x} \\
&\leq \left\| \frac{\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{\det \mathbf{F}} \right\|_{L^1(\Omega; \mathbb{R}^{d \times d})} \|\mathbf{e}(\mathbf{v})\|_{L^\infty(\Omega; \mathbb{R}^{d \times d})} + \left\| \frac{\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \right\|_{L^2(\Omega; \mathbb{R}^d)} \|\dot{\mathbf{m}}\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\stackrel{(3.5c)}{\leq} C \left\| 1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right\|_{L^1(\Omega)} \|\mathbf{e}(\mathbf{v})\|_{L^\infty(\Omega; \mathbb{R}^{d \times d})} + C \left\| 1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right\|_{L^1(\Omega)}^{1/2} \|\dot{\mathbf{m}}\|_{L^2(\Omega; \mathbb{R}^d)} \\
&\stackrel{(3.14b, e)}{\leq} \frac{C^{p'} K^{p'}}{\delta^{1/(p-1)}} \left\| 1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right\|_{L^1(\Omega)}^{p'} + \delta \|\nabla^2 \mathbf{v}\|_{L^p(\Omega; \mathbb{R}^{d \times d \times d})}^p + \delta \|\mathbf{v}|_\Gamma\|_{L^p(\Gamma; \mathbb{R}^d)}^p \\
&\quad + \frac{C^2}{\delta} \left\| 1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right\|_{L^1(\Omega)} + \delta \|\dot{\mathbf{m}}\|_{L^2(\Omega; \mathbb{R}^d)}^2 + \delta \|\text{skw}(\nabla \mathbf{v}) \mathbf{m}\|_{L^2(\Omega; \mathbb{R}^d)}^2 \\
&\leq \left(\frac{C^{p'} K^{p'}}{\delta^{1/(p-1)}} + \frac{C^{2p}}{\delta^p} \right) \left\| 1 + \frac{\varphi(\mathbf{F}, \mathbf{m}) + \theta}{\det \mathbf{F}} \right\|_{L^1(\Omega)}^{p'} + C_{p, s, K} (1 + \|\mathbf{m}\|_{L^s(\Omega; \mathbb{R}^d)}^s)
\end{aligned}$$

$$+ \delta \|\mathring{\mathbf{m}}\|_{L^2(\Omega; \mathbb{R}^d)}^2 + 2\delta \|\nabla^2 \mathbf{v}\|_{L^p(\Omega; \mathbb{R}^{d \times d \times d})}^p + 2\delta \|\mathbf{v}|_\Gamma\|_{L^p(\Gamma; \mathbb{R}^d)}^p. \quad (3.16)$$

The last inequality is due to the estimate, using (3.15), that

$$\begin{aligned} \|\text{skw}(\nabla \mathbf{v}) \mathbf{m}\|_{L^2(\Omega; \mathbb{R}^d)}^2 &\leq C_{p,s,K} (1 + \|\mathbf{m}\|_{L^s(\Omega; \mathbb{R}^d)}^s) + K^{-1} \|\nabla \mathbf{v}\|_{L^\infty(\Omega; \mathbb{R}^{d \times d})}^p \\ &\leq C_{p,s,K} (1 + \|\mathbf{m}\|_{L^s(\Omega; \mathbb{R}^d)}^s) + \|\nabla^2 \mathbf{v}\|_{L^p(\Omega; \mathbb{R}^{d \times d \times d})}^p + \|\mathbf{v}|_\Gamma\|_{L^p(\Gamma; \mathbb{R}^d)}^p \end{aligned} \quad (3.17)$$

which holds if $1/p + 1/s \leq 1/2$, as implied by the assumption $s \geq 2p/(p-2)$. Choosing $\delta > 0$ sufficiently small, the last three terms in (3.17) can be absorbed in the left-hand side of (2.49) while the other terms in (3.16) are already estimated by (3.14b,e,f). Thus we can exploit the dissipation rates in the magneto-mechanical energy dissipation balance (2.49) to obtain estimates

$$\|\mathbf{e}(\mathbf{v})\|_{L^p(I \times \Omega; \mathbb{R}^{d \times d})} \leq C, \quad \|\nabla^2 \mathbf{v}\|_{L^p(I \times \Omega; \mathbb{R}^{d \times d \times d})} \leq C, \quad \text{and} \quad \|\mathring{\mathbf{m}}\|_{L^2(I \times \Omega; \mathbb{R}^d)} \leq C. \quad (3.18)$$

When assuming $p > d$, the estimate (3.18) is essential by preventing evolution of singularities of the quantities transported by such a smooth velocity field. Here, due to qualification of \mathbf{F}_0 and $\varrho_0 = \rho / \det \mathbf{F}_0$ in (3.5k) and (3.5l) which implies also the qualification of the initial conditions

$$\begin{aligned} \nabla \left(\frac{1}{\det \mathbf{F}_0} \right) &= - \frac{\det'(\mathbf{F}_0) : \nabla \mathbf{F}_0}{(\det \mathbf{F}_0)^2} = - \frac{\mathbf{F}_0^{-\top} : \nabla \mathbf{F}_0}{\det \mathbf{F}_0} \in L^r(\Omega; \mathbb{R}^d), \\ \nabla \varrho_0 &= \frac{\nabla \rho}{\det \mathbf{F}_0} - \rho \frac{\mathbf{F}_0^{-\top} : \nabla \mathbf{F}_0}{\det \mathbf{F}_0} \in L^r(\Omega; \mathbb{R}^d), \quad \text{and} \\ \nabla \left(\frac{1}{\varrho_0} \right) &= \nabla \left(\frac{\det \mathbf{F}_0}{\rho} \right) = \frac{\text{Cof} \mathbf{F}_0 : \nabla \mathbf{F}_0}{\rho} - \frac{\det \mathbf{F}_0 \nabla \rho}{\rho^2} \in L^r(\Omega; \mathbb{R}^d), \end{aligned}$$

we can see that Lemma 3.3 yields the estimates

$$\|\mathbf{F}\|_{L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d}))} \leq C_r, \quad \left\| \frac{1}{\det \mathbf{F}} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r, \quad (3.19a)$$

$$\|\varrho\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r, \quad \text{and} \quad \left\| \frac{1}{\varrho} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r \quad \text{for any } 1 \leq r < +\infty. \quad (3.19b)$$

From (3.14a) and (3.19b) with $r > d$, we then have also

$$\|\mathbf{v}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d))} \leq \|\sqrt{\varrho} \mathbf{v}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d))} \left\| \frac{1}{\sqrt{\varrho}} \right\|_{L^\infty(I \times \Omega)} \leq C, \quad (3.19c)$$

and from (3.14c), using (3.5g), we obtain

$$\|\nabla \mathbf{m}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}))} \leq \left\| \frac{\det \mathbf{F}}{\kappa(\mathbf{F})} \right\|_{L^\infty(I \times \Omega)}^{1/2} \left\| \frac{\kappa(\mathbf{F}) |\nabla \mathbf{m}|^2}{\det \mathbf{F}} \right\|_{L^\infty(I; L^1(\Omega))}^{1/2} \leq C. \quad (3.19d)$$

Furthermore, having $\mathring{\mathbf{m}}$ estimated in (3.18) and by using the calculus $\text{div}(\kappa(\mathbf{F}) \nabla \mathbf{m} / \det \mathbf{F}) = \kappa(\mathbf{F}) \Delta \mathbf{m} / \det \mathbf{F} + (\kappa'(\mathbf{F}) / \det \mathbf{F} - \kappa(\mathbf{F}) \text{Cof} \mathbf{F} / \det \mathbf{F}^2) : (\nabla \mathbf{F} \otimes \nabla \mathbf{m})$, we can exploit (2.30d) in the form

$$\Delta \mathbf{m} \in \frac{\det \mathbf{F}}{\kappa(\mathbf{F})} \left(\tau \mathring{\mathbf{m}} + h_c(\mathbf{F}, \theta) \text{Dir}(\mathring{\mathbf{m}}) - \frac{\mathbf{m} \times \mathring{\mathbf{m}}}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} - \mathbf{h}_{\text{ext}} - \nabla u \right)$$

$$+ \frac{\varphi'_m(\mathbf{F}, \mathbf{m}) + \zeta'_m(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} - \left(\frac{\kappa'(\mathbf{F})}{\det \mathbf{F}} - \frac{\kappa(\mathbf{F}) \text{Cof} \mathbf{F}}{\det \mathbf{F}^2} \right) : (\nabla \mathbf{F} \otimes \nabla \mathbf{m}) \quad (3.20)$$

to estimate $\nabla^2 \mathbf{m}$ by the H^2 -regularity of the Laplacean with the homogeneous Neumann boundary conditions, as available on smooth or convex domains. Here we use (3.19a) with $r > d$ and (3.19d), we have $\nabla \mathbf{F} \otimes \nabla \mathbf{m}$ bounded in $L^\infty(I; L^{2r/(r+2)}(\Omega; \mathbb{R}^{d \times d \times d \times d}))$. By (3.5f) and by (3.18), we can still see that $\mathbf{m} \times \mathring{\mathbf{m}} / \gamma(\mathbf{F}, \mathbf{m}, \theta) \in L^2(I \times \Omega; \mathbb{R}^{d \times d})$. By (3.14d), $\nabla u|_\Omega \in L^2(I \times \Omega; \mathbb{R}^d)$. Moreover, by (3.5b), $\varphi'_m(\mathbf{F}, \mathbf{m}) \in L^\infty(I; L^{2^{*2}/(2^{*}+2)}(\Omega; \mathbb{R}^d))$ and by (3.5c) $\zeta'_m(\mathbf{F}, \mathbf{m}, \theta) \in L^\infty(I; L^2(\Omega; \mathbb{R}^d))$. By comparison and the mentioned H^2 -regularity, from (3.20) we obtain

$$\|\nabla^2 \mathbf{m}\|_{L^2(I \times \Omega; \mathbb{R}^{d \times d \times d})} \leq C. \quad (3.21)$$

Noting that the embedding $H^1(\Omega) \subset L^2(\Omega)$ is compact for $r > d$, (3.21) gives always a certain additional information about $\nabla \mathbf{m}$ in comparison with (3.19d).

3.3 Proof of Theorem 3.2

For clarity, we will divide the proof into ten steps. The inertial term and the continuity equation (2.30a) are treated as in [49] and we thus sketch the proof in these aspects.

Let us outline main technical difficulties: The time discretization (Rothe's method) standardly needs convexity of φ (which is not a realistic assumption in finite-strain mechanics) possibly weakened if there is some viscosity in \mathbf{F} (which is not directly considered here, however). Also the conformal space discretization (i.e. the Faedo-Galerkin method) is difficult since it cannot directly copy the energetics because the “nonlinear” test of (2.30c) by $[\varphi / \det]_{\mathbf{F}}'(\mathbf{F})$ needed in (2.34) is problematic in this approximation as $[\varphi / \det]_{\mathbf{F}}'(\mathbf{F})$ is not in the respective finite-dimensional space in general and similarly also the tests of (2.30a) by $|\mathbf{v}|^2$ and of (2.30d) by $\mathring{\mathbf{m}} = \frac{\partial}{\partial t} \mathbf{m} + (\mathbf{v} \cdot \nabla) \mathbf{m} - \text{skw}(\nabla \mathbf{v}) \mathbf{m}$ are problematic.

Step 1: a regularization. Referring to the formal estimates (3.19a), we can choose $\lambda > 0$ so small that, for any possible sufficiently regular solution, it holds

$$\det \mathbf{F} > \lambda \quad \text{and} \quad |\mathbf{F}| < \frac{1}{\lambda} \quad \text{a.e. on } I \times \Omega. \quad (3.22)$$

We first regularize the stress \mathbf{T} and the other nonlinearities in (2.30) by considering a smooth cut-off $\pi_\lambda \in C^1(\mathbb{R}^{d \times d})$ defined as

$$\pi_\lambda(\mathbf{F}) := \begin{cases} 1 & \text{for } \det \mathbf{F} \geq \lambda \text{ and } |\mathbf{F}| \leq 1/\lambda, \\ 0 & \text{for } \det \mathbf{F} \leq \lambda/2 \text{ or } |\mathbf{F}| \geq 2/\lambda, \\ \left(\frac{3}{\lambda^2} (2 \det \mathbf{F} - \lambda)^2 - \frac{2}{\lambda^3} (2 \det \mathbf{F} - \lambda)^3 \right) \times \\ \quad \times (3(\lambda |\mathbf{F}| - 1)^2 - 2(\lambda |\mathbf{F}| - 1)^3) & \text{otherwise.} \end{cases} \quad (3.23)$$

Here $|\cdot|$ stands for the Frobenius norm $|\mathbf{F}| = (\sum_{i,j=1}^d F_{ij}^2)^{1/2}$ for $\mathbf{F} = [F_{ij}]$, which makes π_λ frame indifferent. Thus we can regularize in a smooth way the singular nonlinearity $1/\det(\cdot)$

and also extend $\kappa(\cdot)/\det(\cdot)$ and \mathcal{K} and also ω :

$$\det_\lambda(\mathbf{F}) := \pi_\lambda(\mathbf{F}) \det \mathbf{F} + 1 - \pi_\lambda(\mathbf{F}), \quad (3.24a)$$

$$\kappa_\lambda(\mathbf{F}) := \pi_\lambda(\mathbf{F}) \kappa(\mathbf{F}) + (1 - \pi_\lambda(\mathbf{F})) \det \mathbf{F}, \quad \text{and} \quad (3.24b)$$

$$\mathcal{K}_\lambda(\mathbf{F}, \theta) := \pi_\lambda(\mathbf{F}) \mathcal{K}(\mathbf{F}, \theta) + 1 - \pi_\lambda(\mathbf{F}). \quad (3.24c)$$

Using the operator $\Delta^{-1} \text{div} : L^2(\Omega; \mathbb{R}^d) \rightarrow H^1(\mathbb{R}^d)$ defined by $u = [\Delta^{-1} \text{div}](\mathbf{m})$ as a unique weak solution to (2.1) with the “boundary” condition $u(\infty) = 0$, we can eliminate u ; actually, this is rather the scenario for $d = 3$ otherwise it is formally possible because only ∇u but not u itself occurs in the system and its energetics. Altogether, for the above chosen λ and for any $\varepsilon > 0$, we consider the regularized system

$$\frac{\partial \varrho}{\partial t} = -\text{div}(\varrho \mathbf{v}), \quad (3.25a)$$

$$\begin{aligned} \frac{\partial}{\partial t}(\varrho \mathbf{v}) = & \text{div} \left(\mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) + \mathbf{K}_\lambda(\mathbf{F}, \nabla \mathbf{m}) + \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) + \nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) - \frac{\varrho \mathbf{v} \otimes \mathbf{v}}{\det_\lambda(\mathbf{F})} \right. \\ & \left. - \text{div}(\mathcal{H} + \mathcal{S}_\lambda(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m})) \right) + \mu_0 (\nabla \mathbf{h})^\top \mathbf{m} - \mu_0 \nabla(\mathbf{h} \cdot \mathbf{m}) + \sqrt{\frac{\rho \varrho}{\det_\lambda(\mathbf{F})}} \mathbf{g} \end{aligned}$$

$$\text{with } \mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) = \left(\frac{[\pi_\lambda \varphi]_{\mathbf{F}}'(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} + \frac{\pi_\lambda(\mathbf{F}) \zeta_{\mathbf{F}}'(\mathbf{F}, \mathbf{m}, \theta)}{(1 + \varepsilon |\theta|) \det \mathbf{F}} + \frac{|\nabla \mathbf{m}|^2 \kappa_\lambda'(\mathbf{F})}{2 \det \mathbf{F}} \right) \mathbf{F}^\top,$$

$$\mathbf{h} = \mathbf{h}_{\text{ext}} + \nabla \Delta^{-1} \text{div}(\mathbf{m}), \quad \mathbf{K}_\lambda(\mathbf{F}, \nabla \mathbf{m}) = \frac{\kappa_\lambda(\mathbf{F})}{\det \mathbf{F}} \nabla \mathbf{m} \otimes \nabla \mathbf{m},$$

$$\mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) = \text{skw}((\mu_0 \mathbf{h} - \hat{\mathbf{t}}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta)) \otimes \mathbf{m}), \quad \mathcal{H} = \nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v},$$

$$\mathcal{S}_\lambda(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}) = \frac{\kappa_\lambda(\mathbf{F})}{\det \mathbf{F}} \text{Skw}(\nabla \mathbf{m} \otimes \mathbf{m}), \quad (3.25b)$$

$$\frac{\partial \mathbf{F}}{\partial t} = (\nabla \mathbf{v}) \mathbf{F} - (\mathbf{v} \cdot \nabla) \mathbf{F}, \quad (3.25c)$$

$$\begin{aligned} \tau \mathbf{r} + h_c(\mathbf{F}, \theta) \text{Dir}(\mathbf{r}) - \frac{\mathbf{m} \times \mathbf{r}}{\gamma(\mathbf{F}, \mathbf{m}, \theta)} \ni & \mu_0 \mathbf{h} - \hat{\mathbf{t}}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) - \text{div} \left(\frac{\kappa_\lambda(\mathbf{F}) \nabla \mathbf{m}}{\det \mathbf{F}} \right) \\ \text{with } \hat{\mathbf{t}}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) = & \frac{\pi_\lambda(\mathbf{F}) \varphi_{\mathbf{m}}'(\mathbf{F}, \mathbf{m})}{\det \mathbf{F}} + \frac{\pi_\lambda(\mathbf{F}) \zeta_{\mathbf{m}}'(\mathbf{F}, \mathbf{m}, \theta)}{(1 + \varepsilon |\theta|^{1/2}) \det \mathbf{F}}, \end{aligned} \quad (3.25d)$$

$$\frac{\partial \mathbf{m}}{\partial t} = \text{skw}(\nabla \mathbf{v}) \mathbf{m} - (\mathbf{v} \cdot \nabla) \mathbf{m} + \mathbf{r}, \quad (3.25e)$$

$$\begin{aligned} \frac{\partial w}{\partial t} = & \xi_\varepsilon(\mathbf{F}, \theta; \mathbf{e}(\mathbf{v}), \nabla^2 \mathbf{v}, \mathbf{r}) + \text{div}(\mathcal{K}(\mathbf{F}, \theta) \nabla \theta - w \mathbf{v}) + \frac{\pi_\lambda(\mathbf{F}) \zeta_{\mathbf{F}}'(\mathbf{F}, \mathbf{m}, \theta) \mathbf{F}^\top}{(1 + \varepsilon |\theta|) \det \mathbf{F}} : \mathbf{e}(\mathbf{v}) \\ & + \frac{\pi_\lambda(\mathbf{F}) \zeta_{\mathbf{m}}'(\mathbf{F}, \mathbf{m}, \theta)}{(1 + \varepsilon |\theta|^{1/2}) \det \mathbf{F}} \cdot (\mathbf{r} - \text{skw}(\nabla \mathbf{v}) \mathbf{m}) \quad \text{with } w = \omega(\mathbf{F}, \mathbf{m}, \theta) \end{aligned}$$

$$\text{and } \xi_\varepsilon(\mathbf{F}, \theta; \mathbf{e}, \mathbf{G}, \mathbf{r}) := \frac{\nu_1 |\mathbf{e}|^p + \tau |\mathbf{r}|^2 + h_c(\mathbf{F}, \theta) |\mathbf{r}| + \nu_2 |\mathbf{G}|^p}{1 + \varepsilon |\mathbf{e}|^p + \varepsilon |\mathbf{G}|^p + \varepsilon |\mathbf{r}|^2}, \quad (3.25f)$$

where $\omega(\cdot, \cdot)$ is from (2.30f). We complete this system with the correspondingly regularized boundary conditions on $I \times \Gamma$:

$$\begin{aligned} & \left[(\mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) + \mathbf{K}_\lambda(\mathbf{F}, \nabla \mathbf{m}) + \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}, \mathbf{m}, \theta) + \nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}) \right. \\ & \left. - \text{div}(\mathcal{H} + \mathcal{S}_\lambda(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m})) \right) \mathbf{n} - \text{div}_s(\mathcal{H} \mathbf{n} + \mathcal{S}_\lambda(\mathbf{F}, \mathbf{m}, \nabla \mathbf{m}) \mathbf{n}) \Big]_{\Gamma} + \nu_b |\mathbf{v}|^{p-2} \mathbf{v} = \mathbf{k}, \end{aligned} \quad (3.26a)$$

$$\mathbf{v} \cdot \mathbf{n} = 0, \quad \nabla^2 \mathbf{v} : (\mathbf{n} \otimes \mathbf{n}) = \mathbf{0}, \quad \frac{\kappa_\lambda(\mathbf{F})(\mathbf{n} \cdot \nabla) \mathbf{m}}{\det \mathbf{F}} = \mathbf{0}, \quad u(\infty) = 0, \quad \text{and} \quad (3.26b)$$

$$\mathcal{K}(\mathbf{F}, \theta) \nabla \theta \cdot \mathbf{n} - \frac{\nu_b |\mathbf{v}|^p}{2 + \varepsilon |\mathbf{v}|^p} = h_\varepsilon(\theta) := \frac{h(\theta)}{1 + \varepsilon |h(\theta)|} \quad (3.26c)$$

and initial conditions

$$\varrho(0) = \varrho_0, \quad \mathbf{v}(0) = \mathbf{v}_0, \quad \mathbf{F}(0) = \mathbf{F}_0, \quad \mathbf{m}(0) = \mathbf{m}_0, \quad \theta(0) = \theta_{0,\varepsilon} := \frac{\theta_0}{1 + \varepsilon \theta_0}. \quad (3.26d)$$

Note that $\pi_\lambda \varphi \in C^1(\mathbb{R}^{d \times d})$ if $\varphi \in C^1(\text{GL}^+(d))$ and that $[\pi_\lambda \varphi]'$ together with the regularized Cauchy stress $\mathbf{T}_{\lambda,\varepsilon}$ are bounded, continuous, and vanish if \mathbf{F} “substantially” violates the constraints (3.22). Altogether, also from (3.24b) and (3.24c) we have:

$$\left(\det \mathbf{F} \leq \frac{\lambda}{2} \quad \text{or} \quad |\mathbf{F}| \geq \frac{2}{\lambda} \right) \Rightarrow \mathbf{T}_{\lambda,\varepsilon} = \mathbf{0}, \quad \frac{\kappa_\lambda(\mathbf{F})}{\det \mathbf{F}} = 1, \quad \text{and} \quad \mathcal{K}(\mathbf{F}, \theta) = 1,$$

where a continuous (and smooth) extension of $[\kappa_\lambda/\det](\cdot)$ at 0 is considered. Also, due to (3.5c), $\mathbf{T}_{\lambda,\varepsilon} : \mathbb{R}^{d \times d} \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}_{\text{sym}}^{d \times d}$ is bounded. Thus the part (3.25a,b,d,e) of the system allows for bounds for $\varrho_\varepsilon \mathbf{v}_\varepsilon$, $\nabla \mathbf{v}_\varepsilon$, and \mathbf{m}_ε independently of $(\varrho_\varepsilon, \mathbf{F}_\varepsilon, \theta_\varepsilon)$ which can be estimated subsequently. Moreover, recalling the extension (3.10), note that it is defined also for negative temperatures.

The corresponding weak formulation of (3.25)–(3.26) a’la Definition 3.1 is quite straightforward and we will not explicitly write it, also because it will be obvious from its Galerkin version (3.27) below. The philosophy of the regularization (3.25) is that the estimation of the magneto-mechanical part (3.25a-e) and of the thermal part (3.25f) decouples since $\mathbf{T}_{\lambda,\varepsilon}$ and $\mathbf{S}_{\lambda,\varepsilon}$ are bounded and that heat sources in the heat equation are bounded for fixed $\varepsilon > 0$. Simultaneously, the heat equation has a non-negative solution and, for $\varepsilon \rightarrow 0$, the physical a priori estimates are the same as the formal estimates (3.14)–(3.19) and, when taking $\lambda > 0$ small to comply with (3.22), the λ -regularization becomes eventually inactive, cf. Step 9 below.

Step 2: a semi-discretization. For $\varepsilon > 0$ fixed, we use a spatial semi-discretization, keeping the transport equations (3.25a), (3.25c), and (3.25e) continuous (i.e. non-discretised) when exploiting Lemmas 3.3 and 3.4. More specifically, we make a conformal Galerkin approximation of (3.25b) by using a collection of nested finite-dimensional subspaces $\{V_k\}_{k \in \mathbb{N}}$ whose union is dense in $W^{2,p}(\Omega; \mathbb{R}^d)$ and a conformal Galerkin approximation of (3.25d) and of (3.25f) by using a collection of nested finite-dimensional subspaces $\{Z_k\}_{k \in \mathbb{N}}$ whose union is dense in $H^1(\Omega)$. Without loss of generality, we can assume $\mathbf{v}_0 \in V_1$ and $\theta_{0,\varepsilon} \in Z_1$.

The approximate solution of the regularized system (3.25) will be denoted by

$$(\varrho_{\varepsilon k}, \mathbf{v}_{\varepsilon k}, \mathbf{F}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) : I \rightarrow W^{1,r}(\Omega) \times V_k \times W^{1,r}(\Omega; \mathbb{R}^{d \times d}) \times Z_k^d \times H^1(\Omega; \mathbb{R}^d) \times Z_k.$$

Specifically, such a six-tuple should satisfy

$$\frac{\partial \varrho_{\varepsilon k}}{\partial t} = -\text{div}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}) \quad \text{in the } L^1(I \times \Omega)\text{-sense,} \quad (3.27a)$$

$$\frac{\partial \mathbf{F}_{\varepsilon k}}{\partial t} = (\nabla \mathbf{v}_{\varepsilon k}) \mathbf{F}_{\varepsilon k} - (\mathbf{v}_{\varepsilon k} \cdot \nabla) \mathbf{F}_{\varepsilon k} \quad \text{in the } L^1(I \times \Omega; \mathbb{R}^{d \times d})\text{-sense, and} \quad (3.27b)$$

$$\frac{\partial \mathbf{m}_{\varepsilon k}}{\partial t} = \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k} - (\mathbf{v}_{\varepsilon k} \cdot \nabla) \mathbf{m}_{\varepsilon k} + \mathbf{r}_{\varepsilon k} \quad \text{in the weak sense,} \quad (3.27c)$$

relying on $\varrho_{\varepsilon k} \in W^{1,1}(I \times \Omega)$ and $\mathbf{F}_{\varepsilon k} \in W^{1,1}(I \times \Omega; \mathbb{R}^{d \times d})$ which will be indeed proved later, together with the following integral identities

$$\begin{aligned} & \int_0^T \int_{\Omega} \left((\mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) + \mathbf{K}_{\lambda}(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) - \varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k} + \nu_1 |\mathbf{e}(\mathbf{v}_{\varepsilon k})|^{p-2} \mathbf{e}(\mathbf{v}_{\varepsilon k})) : \mathbf{e}(\tilde{\mathbf{v}}) \right. \\ & \quad + \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) : \text{skw}(\tilde{\mathbf{v}}) - \mu_0 (\nabla \mathbf{h}_{\varepsilon k}) : (\mathbf{m}_{\varepsilon k} \otimes \tilde{\mathbf{v}}) - \mu_0 \mathbf{h}_{\varepsilon k} \cdot \mathbf{m}_{\varepsilon k} (\text{div } \tilde{\mathbf{v}}) - \varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \cdot \frac{\partial \tilde{\mathbf{v}}}{\partial t} \\ & \quad + (\nu_2 |\nabla^2 \mathbf{v}_{\varepsilon k}|^{p-2} \nabla^2 \mathbf{v}_{\varepsilon k} + \mathcal{S}_{\lambda}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k})) : \nabla^2 \tilde{\mathbf{v}} \Big) d\mathbf{x} dt = \int_{\Omega} \varrho_0 \mathbf{v}_0 \cdot \tilde{\mathbf{v}}(0) d\mathbf{x} \\ & \quad + \int_0^T \int_{\Omega} \sqrt{\frac{\rho \varrho_{\varepsilon k}}{\det_{\lambda}(\mathbf{F}_{\varepsilon k})}} \mathbf{g} \cdot \tilde{\mathbf{v}} d\mathbf{x} dt + \int_0^T \int_{\Gamma} (\mathbf{k} - \nu_b |\mathbf{v}_{\varepsilon k}|^{p-2} \mathbf{v}_{\varepsilon k}) \cdot \tilde{\mathbf{v}} dS dt \end{aligned} \quad (3.27d)$$

with $\mathbf{T}_{\lambda, \varepsilon}$, \mathbf{K}_{λ} , $\mathbf{S}_{\lambda, \varepsilon}$, and \mathcal{S}_{λ} from (3.25b) and $\mathbf{h}_{\varepsilon k} = \mathbf{h}_{\text{ext}} + \nabla^2 \Delta^{-1} \text{div}(\mathbf{m}_{\varepsilon k})$ for any $\tilde{\mathbf{v}} \in L^{\infty}(I; V_k)$ with $\tilde{\mathbf{v}} \cdot \mathbf{n} = 0$ on $I \times \Gamma$ and $\tilde{\mathbf{v}}(T) = \mathbf{0}$, and

$$\begin{aligned} & \int_0^T \int_{\Omega} \left(\frac{\tau}{2} |\tilde{\mathbf{r}}|^2 + h_c(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}) |\tilde{\mathbf{r}}| - \text{div} \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon k}) \nabla \mathbf{m}_{\varepsilon k}}{\det \mathbf{F}_{\varepsilon k}} \cdot \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k} + \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon k}) \nabla \mathbf{m}_{\varepsilon k}}{\det \mathbf{F}_{\varepsilon k}} : \nabla \tilde{\mathbf{r}} \right. \\ & \quad - (\mu_0 \mathbf{h} - \hat{\mathbf{t}}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})) \cdot (\tilde{\mathbf{r}} - \mathbf{r}_{\varepsilon k}) + \frac{\mathbf{m}_{\varepsilon k} \times \mathbf{r}_{\varepsilon k}}{\gamma(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})} \cdot \tilde{\mathbf{r}} + \left(\mathbf{K}_{\lambda}(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) \right. \\ & \quad \left. - \frac{\kappa'_{\lambda}(\mathbf{F}_{\varepsilon k}) |\nabla \mathbf{m}_{\varepsilon k}|^2 \mathbf{F}_{\varepsilon k}^{\top}}{2 \det \mathbf{F}_{\varepsilon k}} \right) : \mathbf{e}(\mathbf{v}_{\varepsilon k}) \Big) d\mathbf{x} dt \geq \int_0^T \int_{\Omega} \frac{\tau}{2} |\mathbf{r}_{\varepsilon k}|^2 + h_c(\mathbf{F}, \theta) |\mathbf{r}_{\varepsilon k}| d\mathbf{x} dt \\ & \quad + \int_{\Omega} \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon k}(T)) |\nabla \mathbf{m}_{\varepsilon k}(T)|^2}{2 \det \mathbf{F}_{\varepsilon k}(T)} - \frac{\kappa_{\lambda}(\mathbf{F}_0) |\nabla \mathbf{m}_0|^2}{2 \det \mathbf{F}_0} d\mathbf{x} \end{aligned} \quad (3.27e)$$

holding for any $\tilde{\mathbf{r}} \in Z_k^d$ and for a.a. $t \in I$, and further

$$\begin{aligned} & \int_0^T \int_{\Omega} \left(w_{\varepsilon k} \frac{\partial \tilde{\theta}}{\partial t} + (w_{\varepsilon k} \mathbf{v}_{\varepsilon k} - \mathcal{K}(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}) \nabla \theta_{\varepsilon k}) \cdot \nabla \tilde{\theta} + \xi_{\varepsilon}(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) \tilde{\theta} \right. \\ & \quad + \pi_{\lambda}(\mathbf{F}_{\varepsilon k}) \left(\frac{\zeta'_{\mathbf{F}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^{\top}}{(1 + \varepsilon |\theta_{\varepsilon k}|) \det \mathbf{F}_{\varepsilon k}} : \mathbf{e}(\mathbf{v}_{\varepsilon k}) + \frac{\zeta'_{\mathbf{m}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k})}{(1 + \varepsilon |\theta_{\varepsilon k}|^{1/2}) \det \mathbf{F}_{\varepsilon k}} \right) \tilde{\theta} \Big) d\mathbf{x} dt \\ & \quad + \int_{\Omega} \omega(\mathbf{F}_0, \theta_{0, \varepsilon}) \tilde{\theta}(0) d\mathbf{x} + \int_0^T \int_{\Gamma} \left(h_{\varepsilon}(\theta_{\varepsilon k}) + \frac{\nu_b |\mathbf{v}_{\varepsilon k}|^p}{2 + \varepsilon |\mathbf{v}_{\varepsilon k}|^p} \right) \tilde{\theta} dS dt = 0 \end{aligned}$$

with $w_{\varepsilon k} = \omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \quad (3.27f)$

holds for any $\tilde{\theta} \in C^1(I; Z_k)$ with $\tilde{\theta}(T) = 0$.

Existence of this solution is based on the standard theory of systems of ordinary differential equations first locally in time combined here with the abstract $W^{1,r}(\Omega)$ - and $L^2(\Omega)$ valued differential equations based on Lemmas 3.3 and 3.4 for the scalar, the vector, and the tensor transport equations (3.27a–c) and then by successive prolongation on

the whole time interval based on the L^∞ -estimates below. Usage of Lemmas 3.3 and 3.4 with the fixed initial conditions ϱ_0 , \mathbf{F}_0 , and \mathbf{m}_0 defines the nonlinear operators $\mathfrak{R} : I \times L^p(I; W^{2,p}(\Omega; \mathbb{R}^d)) \rightarrow W^{1,r}(\Omega)$, $\mathfrak{F} : I \times L^p(I; W^{2,p}(\Omega; \mathbb{R}^d)) \rightarrow W^{1,r}(\Omega; \mathbb{R}^{d \times d})$, and $\mathfrak{M} : I \times L^p(I; W^{2,p}(\Omega; \mathbb{R}^d)) \times L^2(I \times \Omega; \mathbb{R}^d) \rightarrow L^2(\Omega; \mathbb{R}^d)$ by

$$\varrho_{\varepsilon k}(t) = \mathfrak{R}(t, \mathbf{v}_{\varepsilon k}), \quad \mathbf{F}_{\varepsilon k}(t) = \mathfrak{F}(t, \mathbf{v}_{\varepsilon k}), \quad \text{and} \quad \mathbf{m}_{\varepsilon k}(t) = \mathfrak{M}(t, \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}). \quad (3.28)$$

Step 3: first a priori estimates. In the Galerkin approximation, it is legitimate to use $\tilde{\mathbf{v}} = \mathbf{v}_{\varepsilon k}$ for (3.27d) and $\tilde{\theta} = \theta_{\varepsilon k}$ for (3.27f). We take the benefit from having the transport equations (3.27a,b) non-discretized and thus we can test them by the nonlinearities $|\mathbf{v}_{\varepsilon k}|^2/2$ and $[\pi_\lambda \varphi(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k})/\det \mathbf{F}_{\varepsilon k}]'_{\mathbf{F}}$, respectively. In particular, we can use the calculus (2.34) which holds also for $\pi_\lambda \varphi$ instead of φ and the calculus (2.38) also for the semi-Galerkin approximate solution. Also we can use the calculus (2.39)–(2.46) with $\pi_\lambda \zeta'_{\mathbf{F}}/(1+|\theta|)$ and $\pi_\lambda \zeta'_{\mathbf{m}}/(1+|\theta|^{1/2})$ and κ_λ instead of $\zeta'_{\mathbf{F}}$ and $\zeta'_{\mathbf{m}}$ and κ , respectively, using also that we have the nondiscretized equation (3.27c) at disposal. The philosophy of the regularization (3.25) is that, for this estimation procedure, the system decouples to the magneto-mechanical part and the thermal part which allows for basic estimates independent of $\mathbf{v}_{\varepsilon k}$, $\mathbf{m}_{\varepsilon k}$, and $\mathbf{r}_{\varepsilon k}$.

Specifically, from (3.27d) tested by $\mathbf{v}_{\varepsilon k}$ and (3.27e) tested by $\mathbf{0}$, like (2.49) we obtain the inequality

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \frac{\varrho_{\varepsilon k}}{2} |\mathbf{v}_{\varepsilon k}|^2 + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \varphi(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k})}{\det \mathbf{F}_{\varepsilon k}} + \frac{\kappa_\lambda(\mathbf{F}_{\varepsilon k})}{2 \det \mathbf{F}_{\varepsilon k}} |\nabla \mathbf{m}_{\varepsilon k}|^2 - \mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m}_{\varepsilon k} \, d\mathbf{x} \\ & + \int_{\Omega} \xi_\varepsilon(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) \, d\mathbf{x} + \int_{\Gamma} \nu_b |\mathbf{v}_{\varepsilon k}|^p \, dS + \frac{d}{dt} \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u_{\varepsilon k}|^2 \, d\mathbf{x} \\ & \leq \int_{\Gamma} \mathbf{k} \cdot \mathbf{v}_{\varepsilon k} \, dS + \int_{\Omega} \left(\sqrt{\frac{\rho \varrho_{\varepsilon k}}{\det_\lambda(\mathbf{F}_{\varepsilon k})}} \mathbf{g} \cdot \mathbf{v}_{\varepsilon k} - \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_{\mathbf{F}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^\top}{(1+\varepsilon|\theta_{\varepsilon k}|) \det \mathbf{F}_{\varepsilon k}} : \mathbf{e}(\mathbf{v}_{\varepsilon k}) \right. \\ & \quad \left. - \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_{\mathbf{m}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})}{(1+\varepsilon|\theta_{\varepsilon k}|^{1/2}) \det \mathbf{F}_{\varepsilon k}} \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k}) - \mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}_{\varepsilon k} \right) d\mathbf{x} \end{aligned} \quad (3.29)$$

with $u_{\varepsilon k} = \Delta^{-1} \text{div}(\mathbf{m}_{\varepsilon k})$.

The bulk-force term can be estimated similarly as (3.11):

$$\begin{aligned} & \int_{\Omega} \sqrt{\frac{\rho}{\det_\lambda(\mathbf{F}_{\varepsilon k})}} \sqrt{\varrho_{\varepsilon k}} \mathbf{v}_{\varepsilon k} \cdot \mathbf{g} \, d\mathbf{x} \leq \left\| \sqrt{\frac{\rho}{\det_\lambda(\mathbf{F}_{\varepsilon k})}} \right\|_{L^2(\Omega)} \left\| \sqrt{\varrho_{\varepsilon k}} \mathbf{v}_{\varepsilon k} \right\|_{L^2(\Omega; \mathbb{R}^d)} \left\| \mathbf{g} \right\|_{L^\infty(\Omega; \mathbb{R}^d)} \\ & \leq \left\| \mathbf{g} \right\|_{L^\infty(\Omega; \mathbb{R}^d)} \int_{\Omega} \frac{\rho}{2 \det_\lambda(\mathbf{F}_{\varepsilon k})} + \frac{\varrho_{\varepsilon k}}{2} |\mathbf{v}_{\varepsilon k}|^2 \, d\mathbf{x} \\ & = \left\| \mathbf{g} \right\|_{L^\infty(\Omega; \mathbb{R}^d)} \left(\frac{\text{meas}(\Omega)(1 + \max_{\Omega} \rho)}{2\lambda} + \int_{\Omega} \frac{\varrho_{\varepsilon k}}{2} |\mathbf{v}_{\varepsilon k}|^2 \, d\mathbf{x} \right) \end{aligned} \quad (3.30)$$

while the other terms in (3.29) can be estimated as done in (3.12), (3.13), (3.16), and (3.17).

By the Gronwall inequality, we obtain the estimates

$$\left\| \mathbf{v}_{\varepsilon k} \right\|_{L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))} \leq C_\varepsilon \quad \text{and} \quad \left\| \sqrt{\varrho_{\varepsilon k}} \mathbf{v}_{\varepsilon k} \right\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d))} \leq C_\varepsilon, \quad (3.31a)$$

$$\left\| \mathbf{r}_{\varepsilon k} \right\|_{L^2(I \times \Omega; \mathbb{R}^d)} \leq C_\varepsilon, \quad \text{and} \quad \left\| u_{\varepsilon k} \right\|_{L^\infty(I; H^1(\mathbb{R}^d))} \leq C_\varepsilon \quad (3.31b)$$

with C_ε depending on $\varepsilon > 0$ considered fixed in this Step. The former estimate in (3.31a) relies also on the Navier-type boundary conditions and allows us to use Lemma 3.3 to obtain the estimate

$$\|\mathbf{F}_{\varepsilon k}\|_{L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d}))} \leq C_{r,\varepsilon} \quad \text{with} \quad \left\| \frac{1}{\det \mathbf{F}_{\varepsilon k}} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_{r,\varepsilon}, \quad (3.31c)$$

$$\|\varrho_{\varepsilon k}\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_{r,\varepsilon} \quad \text{with} \quad \left\| \frac{1}{\varrho_{\varepsilon k}} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_{r,\varepsilon}, \quad (3.31d)$$

$$\|\mathbf{v}_{\varepsilon k}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d))} \leq C_\varepsilon, \quad \text{and} \quad (3.31e)$$

$$\|\mathbf{m}_{\varepsilon k}\|_{L^\infty(I; H^1(\Omega; \mathbb{R}^d))} \leq C_\varepsilon; \quad (3.31f)$$

for (3.31e) and (3.31f) we used argumentation like in (3.19c) and (3.31f), respectively. For λ and ε fixed, it is important that these estimates can be made independently of θ since $\mathbf{T}_{\lambda,\varepsilon}$ is a-priori bounded. It is also important that, due to the latter estimate in (3.31c), the singularity in $\zeta(\cdot, \mathbf{m}, \theta)$ is not active and $\omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})$ is well defined.

Also, we have the time derivatives estimated by comparison from (3.27) as

$$\left\| \frac{\partial \varrho_{\varepsilon k}}{\partial t} \right\|_{L^p(I; L^r(\Omega))} \leq C_\varepsilon, \quad \left\| \frac{\partial \mathbf{F}_{\varepsilon k}}{\partial t} \right\|_{L^p(I; L^r(\Omega; \mathbb{R}^{d \times d}))} \leq C_\varepsilon, \quad \left\| \frac{\partial \mathbf{m}_{\varepsilon k}}{\partial t} \right\|_{L^2(I \times \Omega; \mathbb{R}^d)} \leq C_\varepsilon. \quad (3.32)$$

The further estimates can be obtained by testing the Galerkin approximation of (3.27f) by $\tilde{\theta} = \theta_{\varepsilon k}$. This is to be made carefully not to see terms as $\theta_{\varepsilon k} \omega'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) : (\mathbf{v}_{\varepsilon k} \cdot \nabla) \mathbf{F}_{\varepsilon k}$ which is not integrable. To this goal, we consider the convective-derivative form $\frac{\partial}{\partial t} w + \operatorname{div}(w \mathbf{v}) = \dot{w} + w \operatorname{div} \mathbf{v}$ in (3.25f). We denote by $\widehat{\omega}(\mathbf{F}, \mathbf{m}, \theta)$ a primitive function to $\theta \mapsto \theta \omega'_\theta(\mathbf{F}, \mathbf{m}, \theta)$ depending smoothly on \mathbf{F} and on \mathbf{m} , specifically

$$\widehat{\omega}_\lambda(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 r \theta^2 \omega'_\theta(\mathbf{F}, \mathbf{m}, r\theta) \, dr. \quad (3.33)$$

For $w_{\varepsilon k} = \omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})$ and using (3.27b), the mentioned test by $\theta_{\varepsilon k}$ then gives

$$\begin{aligned} \theta_{\varepsilon k} \dot{w}_{\varepsilon k} &= \theta_{\varepsilon k} \omega'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) : \dot{\mathbf{F}}_{\varepsilon k} \\ &\quad + \theta_{\varepsilon k} \omega'_\mathbf{m}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \cdot \dot{\mathbf{m}}_{\varepsilon k} + \theta_{\varepsilon k} \omega'_\theta(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \dot{\theta}_{\varepsilon k} \\ &= \left(\theta_{\varepsilon k} \omega'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \widehat{\omega}'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) : (\nabla \mathbf{v}_{\varepsilon k}) \mathbf{F}_{\varepsilon k} \\ &\quad + \left(\theta_{\varepsilon k} \omega'_\mathbf{m}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \widehat{\omega}'_\mathbf{m}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) \cdot \dot{\mathbf{m}}_{\varepsilon k} + \overline{\widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})}. \end{aligned} \quad (3.34)$$

Integrating the last term over Ω gives, by the Green formula, $\int_\Omega \overline{\widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})} \, d\mathbf{x} = \frac{d}{dt} \int_\Omega \widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \, d\mathbf{x} - \int_\Omega \widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \operatorname{div} \mathbf{v}_{\varepsilon k} \, d\mathbf{x} + \int_\Gamma \widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{v}_{\varepsilon k} \cdot \mathbf{n} \, dS$. Thus, we obtain:

$$\begin{aligned} &\frac{d}{dt} \int_\Omega \widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \, d\mathbf{x} + \int_\Omega \mathcal{K}(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}) |\nabla \theta_{\varepsilon k}|^2 \, d\mathbf{x} \\ &= \int_\Omega \left(\left(\xi_\varepsilon(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^\top}{(1 + \varepsilon |\theta_{\varepsilon k}|) \det \mathbf{F}_{\varepsilon k}} : \mathbf{e}(\mathbf{v}_{\varepsilon k}) \right) \right. \end{aligned}$$

$$\begin{aligned}
& + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_m(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k})}{(1 + \varepsilon |\theta_{\varepsilon k}|^{1/2}) \det \mathbf{F}_{\varepsilon k}} - \omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \text{div } \mathbf{v}_{\varepsilon k} \Big) \theta_{\varepsilon k} \\
& - \left(\theta_{\varepsilon k} \omega'_F(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \widehat{\omega}'_F(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) : (\nabla \mathbf{v}_{\varepsilon k}) \mathbf{F}_{\varepsilon k} \\
& - \left(\theta_{\varepsilon k} \omega'_m(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \widehat{\omega}'_m(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k}) \\
& + \widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \text{div } \mathbf{v}_{\varepsilon k} \Big) \, \mathrm{d}\mathbf{x} + \int_\Gamma \left(h_\varepsilon(\theta_{\varepsilon k}) - \frac{\nu_b |\mathbf{v}_{\varepsilon k}|^p}{2 + \varepsilon |\mathbf{v}_{\varepsilon k}|^p} \right) \theta_{\varepsilon k} \, \mathrm{d}S. \tag{3.35}
\end{aligned}$$

We integrate (3.35) in time over an interval $[0, t]$ with $t \in I$. For the left-hand side, let us realize that $\widehat{\omega}(\mathbf{F}, \mathbf{m}, \theta) \geq c_K \theta^2$ due to (3.5c) with $c_K > 0$ depending on the fixed $\lambda > 0$ used in (3.22). Then the integrated right-hand side of (3.35) is to be estimated from above, in particular relying on (3.5c) and on (3.5d). Let us discuss the difficult terms. In view of (3.33), it holds that $\widehat{\omega}'_F(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 r \theta^2 \omega''_{F\theta}(\mathbf{F}, \mathbf{m}, r\theta) \, \mathrm{d}r$ and $\omega'_m(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 r \theta^2 \omega''_{m\theta}(\mathbf{F}, \mathbf{m}, r\theta) \, \mathrm{d}r$. Recalling (3.5d), we have $|\widehat{\omega}'_F(\mathbf{F}, \mathbf{m}, \theta) - \theta \omega'_F(\mathbf{F}, \mathbf{m}, \theta)| \leq C(1 + |\theta|^2)$ and $|\widehat{\omega}'_m(\mathbf{F}, \mathbf{m}, \theta) - \theta \omega'_m(\mathbf{F}, \mathbf{m}, \theta)| \leq C(1 + |\theta|)$. It allows for estimation

$$\begin{aligned}
& \left| \int_\Omega \left(\widehat{\omega}'_F(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \theta_{\varepsilon k} \omega'_F(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) : (\nabla \mathbf{v}_{\varepsilon k}) \mathbf{F}_{\varepsilon k} \, \mathrm{d}\mathbf{x} \right| \\
& \leq C(|\Omega| + \|\theta_{\varepsilon k}\|_{L^2(\Omega)}^2) \|(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{F}_{\varepsilon k}\|_{L^\infty(\Omega; \mathbb{R}^{d \times d})} \quad \text{and} \tag{3.36}
\end{aligned}$$

$$\begin{aligned}
& \left| \int_\Omega \left(\widehat{\omega}'_m(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \theta_{\varepsilon k} \omega'_m(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k}) \, \mathrm{d}\mathbf{x} \right| \\
& \leq 2C^2(|\Omega| + \|\theta_{\varepsilon k}\|_{L^2(\Omega)}^2) + \|(\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k})\|_{L^2(\Omega; \mathbb{R}^d)}^2. \tag{3.37}
\end{aligned}$$

Using (3.5d) together with $\omega(\mathbf{F}, \mathbf{m}, 0) = 0$ so that $|\omega(\mathbf{F}, \mathbf{m}, \theta)| \leq C_K |\theta|$, the convective terms $\omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) (\text{div } \mathbf{v}_{\varepsilon k}) \theta_{\varepsilon k}$ and $\widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \text{div } \mathbf{v}_{\varepsilon k}$ in (3.35) can be estimated as

$$\left| \int_\Omega \left(\widehat{\omega}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \theta_{\varepsilon k} \omega(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right) \text{div } \mathbf{v}_{\varepsilon k} \, \mathrm{d}\mathbf{x} \right| \leq \frac{C_K}{2} \|\theta_{\varepsilon k}\|_{L^2(\Omega)}^2 \|\text{div } \mathbf{v}_{\varepsilon k}\|_{L^\infty(\Omega)}. \tag{3.38}$$

The terms $\|\theta_{\varepsilon k}\|_{L^2(\Omega)}^2$ in (3.36) and in (3.37) and in (3.38) are to be treated by the Gronwall inequality. The boundary term in (3.35) can be estimated by (3.5i), taking also into the account the extension (3.10), as

$$\begin{aligned}
& \int_\Gamma \left(h_\varepsilon(\theta_{\varepsilon k}) - \frac{\nu_b |\mathbf{v}_{\varepsilon k}|^p}{2 + \varepsilon |\mathbf{v}_{\varepsilon k}|^p} \right) \theta_{\varepsilon k} \, \mathrm{d}S \leq C_{\varepsilon, \nu_b, a} + a \|\theta_{\varepsilon k}\|_{L^2(\Gamma)}^2 \\
& \leq C_{\varepsilon, \nu_b, a} + a N^2 (\|\theta_{\varepsilon k}\|_{L^2(\Omega)}^2 + \|\nabla \theta_{\varepsilon k}\|_{L^2(\Omega; \mathbb{R}^d)}^2), \tag{3.39}
\end{aligned}$$

where $C_{\varepsilon, \nu_b, \delta}$ depends also on C from (3.5i), N is the norm of the trace operator $H^1(\Omega) \rightarrow L^2(\Gamma)$. For $a > 0$ in (3.39) sufficiently small, the last term can be absorbed in the left-hand side of (3.35). Exploiting again the bound (3.31a,b), we eventually obtain the estimate

$$\|\theta_{\varepsilon k}\|_{L^\infty(I; L^2(\Omega)) \cap L^2(I; H^1(\Omega))} \leq C \quad \text{and also} \tag{3.40a}$$

$$\|w_{\varepsilon k}\|_{L^\infty(I; L^2(\Omega)) \cap L^2(I; H^1(\Omega))} \leq C. \tag{3.40b}$$

For (3.40b), we used the calculus

$$\begin{aligned} \nabla w_{\varepsilon k} &= [\omega_\lambda]_\theta'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \nabla \theta_{\varepsilon k} + [\omega_\lambda]_{\mathbf{F}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \nabla \mathbf{F}_{\varepsilon k} \\ &\quad + [\omega_\lambda]_{\mathbf{m}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \nabla \mathbf{m}_{\varepsilon k} \in L^2(I \times \Omega; \mathbb{R}^d) \end{aligned} \quad (3.41)$$

together with the already proved information that $|[\omega_\lambda]_{\mathbf{F}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})|$ is bounded in $L^2(I; L^{2^*}(\Omega))$ and $|[\omega_\lambda]_{\mathbf{m}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})|$ is bounded in $L^\infty(I \times \Omega)$ due to (3.5d), while $|\nabla \mathbf{F}_{\varepsilon k}|$ is bounded in $L^\infty(I; L^r(\Omega))$ and $|\nabla \mathbf{m}_{\varepsilon k}|$ is bounded in $L^\infty(I; L^2(\Omega))$.

Step 4: Limit passage in the magneto-mechanical part for $k \rightarrow \infty$. Using the Banach selection principle, we can extract some subsequence of $\{(\varrho_{\varepsilon k}, \mathbf{v}_{\varepsilon k}, \mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}, w_{\varepsilon k})\}_{k \in \mathbb{N}}$ and its limit $(\varrho_\varepsilon, \mathbf{v}_\varepsilon, \mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \mathbf{r}_\varepsilon, w_\varepsilon) : I \rightarrow W^{1,r}(\Omega) \times L^2(\Omega; \mathbb{R}^d) \times W^{1,r}(\Omega; \mathbb{R}^{d \times d}) \times H^1(\Omega; \mathbb{R}^d) \times L^2(\Omega; \mathbb{R}^d) \times L^2(\Omega)$ such that

$$\varrho_{\varepsilon k} \rightarrow \varrho_\varepsilon \quad \text{weakly}^* \text{ in } L^\infty(I; W^{1,r}(\Omega)), \quad (3.42a)$$

$$\mathbf{v}_{\varepsilon k} \rightarrow \mathbf{v}_\varepsilon \quad \text{weakly}^* \text{ in } L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d)), \quad (3.42b)$$

$$\mathbf{F}_{\varepsilon k} \rightarrow \mathbf{F}_\varepsilon \quad \text{weakly}^* \text{ in } L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d})), \quad (3.42c)$$

$$\mathbf{m}_{\varepsilon k} \rightarrow \mathbf{m}_\varepsilon \quad \text{weakly}^* \text{ in } L^\infty(I; H^1(\Omega; \mathbb{R}^d)), \quad (3.42d)$$

$$\mathbf{r}_{\varepsilon k} \rightarrow \mathbf{r}_\varepsilon \quad \text{weakly in } L^2(I \times \Omega; \mathbb{R}^d), \quad (3.42e)$$

$$w_{\varepsilon k} \rightarrow w_\varepsilon \quad \text{weakly}^* \text{ in } L^\infty(I; L^2(\Omega)) \cap L^2(I; H^1(\Omega)). \quad (3.42f)$$

Relying on the assumption $r > d$ and on estimates (3.32) on $\frac{\partial}{\partial t} \varrho_{\varepsilon k}$, $\frac{\partial}{\partial t} \mathbf{F}_{\varepsilon k}$, and $\frac{\partial}{\partial t} \mathbf{m}_{\varepsilon k}$, by the Aubin-Lions lemma we also have that

$$\varrho_{\varepsilon k} \rightarrow \varrho_\varepsilon \quad \text{strongly in } C(I \times \bar{\Omega}), \quad (3.43a)$$

$$\mathbf{F}_{\varepsilon k} \rightarrow \mathbf{F}_\varepsilon \quad \text{strongly in } C(I \times \bar{\Omega}; \mathbb{R}^{d \times d}), \quad \text{and} \quad (3.43b)$$

$$\mathbf{m}_{\varepsilon k} \rightarrow \mathbf{m}_\varepsilon \quad \text{strongly in } C(I \times \bar{\Omega}; \mathbb{R}^d). \quad (3.43c)$$

This already allows for the limit passage in the evolution-and-transport equations (3.27), cf. (3.7) and (3.9).

Further, by comparison in the equation (3.25f) with the boundary condition (3.26c) in its Galerkin approximation, we obtain a bound on $\frac{\partial}{\partial t} w_{\varepsilon k}$ in seminorms $|\cdot|_l$ on $L^2(I; H^1(\Omega)^*)$ arising from this Galerkin approximation, defined as $|f|_l := \sup\{\int_0^T \int_\Omega f \tilde{\theta} \, d\mathbf{x} dt; \|\tilde{\theta}\|_{L^2(I; H^1(\Omega))} \leq 1, \tilde{\theta}(t) \in Z_l \text{ for } t \in I\}$. More specifically, for any $k \geq l$, we can estimate

$$\begin{aligned} \left| \frac{\partial w_{\varepsilon k}}{\partial t} \right|_l &= \sup_{\substack{\tilde{\theta}(t) \in Z_l \text{ for } t \in I \\ \|\tilde{\theta}\|_{L^2(I; H^1(\Omega))} \leq 1}} \int_0^T \int_\Omega \left(-\mathcal{K}(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}) \nabla \theta_{\varepsilon k} \cdot \nabla \tilde{\theta} + \left(\xi_\varepsilon(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) \right. \right. \\ &\quad \left. \left. + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta_{\mathbf{F}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^\top : \mathbf{e}(\mathbf{v}_{\varepsilon k})}{(1 + \varepsilon |\theta_{\varepsilon k}|) \det \mathbf{F}_{\varepsilon k}} \right. \right. \\ &\quad \left. \left. + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta_{\mathbf{m}}'(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})}{(1 + \varepsilon |\theta_{\varepsilon k}|^{1/2}) \det \mathbf{F}_{\varepsilon k}} \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k}) \right) \tilde{\theta} \right) d\mathbf{x} dt \\ &\quad + \int_0^T \int_\Gamma \left(h_\varepsilon(\theta_{\varepsilon k}) + \frac{\nu_b |\mathbf{v}_{\varepsilon k}|^p}{2 + \varepsilon |\mathbf{v}_{\varepsilon k}|^p} \right) \tilde{\theta} \, dS dt \leq C \end{aligned} \quad (3.44)$$

with some C depending on the estimates (3.31a,b) and (3.40a) but independent on $l \in N$. Thus, by (3.42f) and by a generalized Aubin-Lions theorem [46, Ch.8], we obtain

$$w_{\varepsilon k} \rightarrow w_\varepsilon \quad \text{strongly in } L^s(I \times \Omega) \text{ for } 1 \leq s < 2 + 4/d. \quad (3.45a)$$

Since $\omega_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \cdot)$ is increasing, we can write $\theta_{\varepsilon k} = [\omega_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \cdot)]^{-1}(w_{\varepsilon k})$. Thanks to the continuity of $(\mathbf{F}, \mathbf{m}, w) \mapsto [\omega_\lambda(\mathbf{F}, \mathbf{m}, \cdot)]^{-1}(w) : \mathbb{R}^{d \times d} \times \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}$ and the at most linear growth with respect to w uniformly with respect to \mathbf{F} from any compact $K \subset \text{GL}^+(d)$, cf. (3.5c), we have also

$$\theta_{\varepsilon k} \rightarrow \theta_\varepsilon = [\omega_\lambda(\mathbf{F}_\varepsilon, \mathbf{m}_{\varepsilon k}, \cdot)]^{-1}(w_\varepsilon) \quad \text{strongly in } L^s(I \times \Omega) \text{ for } 1 \leq s < 2 + 4/d; \quad (3.45b)$$

actually, (3.45a,b) results from interpolation of (3.40). Note that we do not have any direct information about $\frac{\partial}{\partial t} \theta_{\varepsilon k}$ so that we could not use the Aubin-Lions arguments straight for $\{\theta_{\varepsilon k}\}_{k \in \mathbb{N}}$. Thus, by the continuity of the corresponding Nemytskiĭ (or here simply superposition) mappings, also the conservative part of the regularized Cauchy stress as well as the heat part of the internal energy, namely

$$\mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \rightarrow \mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \quad \text{strongly in } L^c(I \times \Omega; \mathbb{R}_{\text{sym}}^{d \times d}), \quad 1 \leq c < \infty, \quad (3.45c)$$

$$\begin{aligned} & \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_\mathbf{F}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^\top}{(1 + \varepsilon |\theta_{\varepsilon k}|) \det \mathbf{F}_{\varepsilon k}} \\ & \rightarrow \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_\mathbf{F}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top}{(1 + \varepsilon \theta_\varepsilon) \det \mathbf{F}_{\varepsilon k}} \quad \text{strongly in } L^c(I \times \Omega; \mathbb{R}_{\text{sym}}^{d \times d}), \quad 1 \leq c < \infty, \end{aligned} \quad (3.45d)$$

$$\mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \rightarrow \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \quad \text{strongly in } L^2(I; L^2(\Omega; \mathbb{R}_{\text{skw}}^{d \times d})), \quad 1 \leq c < \infty, \quad (3.45e)$$

$$\begin{aligned} & \mathcal{J}_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) \\ & \rightarrow \mathcal{J}_\lambda(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \nabla \mathbf{m}_\varepsilon) \quad \text{weakly* in } L^\infty(I; L^{2^*2/(2^*+2)}(\Omega; \mathbb{R}^{d \times d \times d})), \end{aligned} \quad (3.45f)$$

$$\hat{\mathbf{t}}_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \rightarrow \hat{\mathbf{t}}_\lambda(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \quad \text{strongly in } L^c(I \times \Omega; \mathbb{R}^d), \quad 1 \leq c < 2, \quad (3.45g)$$

$$\omega_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \rightarrow \omega_\lambda(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \quad \text{strongly in } L^c(I \times \Omega), \quad 1 \leq c < 2 + 4/d. \quad (3.45h)$$

It is important to notice that $\nabla(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}) = \nabla \varrho_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k} + \varrho_{\varepsilon k} \nabla \mathbf{v}_{\varepsilon k}$ is bounded in $L^\infty(I; L^r(\Omega; \mathbb{R}^{d \times d}))$ due to the already obtained bounds (3.31a,c,d). Therefore, $\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}$ converges weakly* in $L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^d))$. The limit of $\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}$ can be identified as $\varrho_\varepsilon \mathbf{v}_\varepsilon$ because we already showed that $\varrho_{\varepsilon k}$ converges strongly in (3.43a) and $\mathbf{v}_{\varepsilon k}$ converges weakly due to (3.42b).

By comparison (3.27d), we also obtain an information about $\frac{\partial}{\partial t}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})$. Specifically, for any $\tilde{\mathbf{v}} \in L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))$ with $\tilde{\mathbf{v}}(t) \in V_k$ for a.a. $t \in I$, we have

$$\begin{aligned} & \int_0^T \int_\Omega \frac{\partial}{\partial t}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}) \cdot \tilde{\mathbf{v}} \, d\mathbf{x} dt = \int_0^T \int_\Gamma (\mathbf{k} + \nu_b |\mathbf{v}_{\varepsilon k}|^{p-2} \mathbf{v}_{\varepsilon k}) \cdot \tilde{\mathbf{v}} \, dS dt + \int_0^T \int_\Omega \left(\sqrt{\frac{\rho \varrho_{\varepsilon k}}{\det_\lambda(\mathbf{F}_{\varepsilon k})}} \mathbf{g} \cdot \tilde{\mathbf{v}} \right. \\ & + \left(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k} - \mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \mathbf{K}_\lambda(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) - \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \right. \\ & \left. \left. - \nu_1 |\mathbf{e}(\mathbf{v}_{\varepsilon k})|^{p-2} \mathbf{e}(\mathbf{v}_{\varepsilon k}) \right) : \nabla \tilde{\mathbf{v}} - \left(\nu_2 |\nabla^2 \mathbf{v}_{\varepsilon k}|^{p-2} \nabla^2 \mathbf{v}_{\varepsilon k} + \mathcal{J}_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) \right) : \nabla \mathbf{e}(\tilde{\mathbf{v}}) \right. \\ & \left. + \mu_0 \nabla \mathbf{h}_{\varepsilon k} : (\mathbf{m}_{\varepsilon k} \otimes \tilde{\mathbf{v}}) + \mu_0 \mathbf{h}_{\varepsilon k} \cdot \mathbf{m}_{\varepsilon k} (\text{div } \tilde{\mathbf{v}}) \right) d\mathbf{x} dt \leq C \|\tilde{\mathbf{v}}\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))} \end{aligned} \quad (3.46)$$

with C independent of k . This yields a bound for $\frac{\partial}{\partial t}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})$ in a seminorm on $L^1(I; L^2(\Omega; \mathbb{R}^d)) + L^{p'}(I; W^{2,p}(\Omega; \mathbb{R}^d)^*)$ induced by the Galerkin discretization by V_k , and by any V_l with $l \leq k$ with C in (3.46) independent of k . Here we used in particular that $\mathbf{K}_\lambda(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k})$ and $\mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})$ are bounded in $L^\infty(I; L^1(\Omega; \mathbb{R}^{d \times d}))$ which is surely in duality with $L^p(I; W^{1,p}(\Omega; \mathbb{R}^{d \times d}))$ and that $\mathcal{S}_\lambda(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k})$ is bounded in $L^\infty(I; L^{2^*2/(2^+2)}(\Omega; \mathbb{R}^{d \times d \times d}))$ which is in duality with $L^p(I \times \Omega; \mathbb{R}^{d \times d \times d})$ if $p > d$. By a generalization of the Aubin-Lions compact-embedding theorem, cf. [46, Lemma 7.7], we then obtain

$$\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \rightarrow \varrho_k \mathbf{v}_\varepsilon \quad \text{strongly in } L^c(I \times \Omega; \mathbb{R}^d) \quad \text{for any } 1 \leq c < 4. \quad (3.47a)$$

Since obviously $\mathbf{v}_{\varepsilon k} = (\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})(1/\varrho_{\varepsilon k})$, thanks to (3.43a) and (3.47a), we also have that

$$\mathbf{v}_{\varepsilon k} \rightarrow \mathbf{v}_\varepsilon \quad \text{strongly in } L^c(I \times \Omega; \mathbb{R}^d) \quad \text{with any } 1 \leq c < 4. \quad (3.47b)$$

For the limit passage in the momentum equation, one uses the monotonicity of the dissipative stress, i.e., the monotonicity of the quasilinear operator $\mathbf{v} \mapsto \operatorname{div}(\operatorname{div}(\nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v}) - \nu_1 |\mathbf{e}(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v}))$, as well as of the time-derivative operator. One could use the already obtained weak convergences and the so-called Minty trick but, later, we will need a strong convergence of $\mathbf{e}(\mathbf{v}_{\varepsilon k})$ to pass to the limit in the heat equation. Thus we first prove this strong convergence, which then allows for the limit passage in the momentum equation directly. We will use the weak convergence of the inertial force

$$\int_0^T \int_\Omega \left(\frac{\partial(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})}{\partial t} + \operatorname{div}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k}) \right) \cdot \tilde{\mathbf{v}} \, d\mathbf{x} dt \xrightarrow{k \rightarrow \infty} \int_0^T \int_\Omega \left(\frac{\partial(\varrho_\varepsilon \mathbf{v}_\varepsilon)}{\partial t} + \operatorname{div}(\varrho_\varepsilon \mathbf{v}_\varepsilon \otimes \mathbf{v}_\varepsilon) \right) \cdot \tilde{\mathbf{v}} \, d\mathbf{x} dt; \quad (3.48)$$

cf. [49, 50]. Further, relying on the calculus (2.38), we will use the identity

$$\begin{aligned} \int_\Omega \frac{\varrho_{\varepsilon k}(T)}{2} |\mathbf{v}_{\varepsilon k}(T) - \mathbf{v}_\varepsilon(T)|^2 \, d\mathbf{x} &= \int_0^T \int_\Omega \left(\frac{\partial(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})}{\partial t} + \operatorname{div}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k}) \right) \cdot \mathbf{v}_{\varepsilon k} \, d\mathbf{x} dt \\ &\quad + \int_\Omega \frac{\varrho_0}{2} |\mathbf{v}_0|^2 - \varrho_{\varepsilon k}(T) \mathbf{v}_{\varepsilon k}(T) \cdot \mathbf{v}_\varepsilon(T) + \frac{\varrho_{\varepsilon k}(T)}{2} |\mathbf{v}_\varepsilon(T)|^2 \, d\mathbf{x}. \end{aligned} \quad (3.49)$$

We further used that the $\varrho_{\varepsilon k}(T)$ is also bounded in $W^{1,r}(\Omega)$ and $\mathbf{v}_{\varepsilon k}(T)$ is bounded in $L^2(\Omega; \mathbb{R}^d)$, together with some information about the time derivative $\frac{\partial}{\partial t}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k})$, cf. (3.46), so that we can identify the weak limit of $\varrho_{\varepsilon k}(T) \mathbf{v}_{\varepsilon k}(T)$. Specifically, we have that

$$\varrho_{\varepsilon k}(T) \mathbf{v}_{\varepsilon k}(T) \rightarrow \varrho_\varepsilon(T) \mathbf{v}_\varepsilon(T) \quad \text{weakly in } L^2(\Omega; \mathbb{R}^d). \quad (3.50)$$

For the stress $\mathbf{K}_\lambda(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k})$, we will further need strong convergence of $\nabla \mathbf{m}_{\varepsilon k}$. This can be seen from the uniform monotonicity of the operator $\mathbf{m} \mapsto -\operatorname{div}(\kappa_\lambda(\mathbf{F}) \nabla \mathbf{m} / \det \mathbf{F})$. We use the Galerkin approximation (3.27e) tested by $\tilde{\mathbf{r}} = \mathbf{m}_{\varepsilon k} - \widetilde{\mathbf{m}}_k$ with an approximation $\widetilde{\mathbf{m}}_k : I \rightarrow Z_k$ of \mathbf{m}_ε converging to \mathbf{m}_ε strongly in $L^2(I; H^1(\Omega; \mathbb{R}^d))$. Then we can estimate

$$c_{\kappa, \lambda} \|\nabla \mathbf{m}_{\varepsilon k} - \nabla \mathbf{m}_\varepsilon\|_{L^2(I \times \Omega)}^2 \leq \int_0^T \int_\Omega \frac{\kappa_\lambda(\mathbf{F}_{\varepsilon k})}{\det \mathbf{F}_{\varepsilon k}} |\nabla \mathbf{m}_{\varepsilon k} - \nabla \mathbf{m}_\varepsilon|^2 \, d\mathbf{x} dt$$

$$\begin{aligned}
&= \int_0^T \int_{\Omega} \left(\widehat{\mathbf{t}}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) - \mathbf{h}_{\varepsilon k} + \tau \mathbf{r}_{\varepsilon k} + h_C(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{d}_{\varepsilon k} - \frac{\mathbf{m}_{\varepsilon k} \times \mathbf{r}_{\varepsilon k}}{\gamma(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})} \right) \cdot (\mathbf{m}_{\varepsilon k} - \widetilde{\mathbf{m}}_k) \\
&\quad + \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon k}) \nabla \mathbf{m}_{\varepsilon k}}{\det \mathbf{F}_{\varepsilon k}} : \nabla (\mathbf{m}_{\varepsilon} - \widetilde{\mathbf{m}}_k) - \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon k}) \nabla \mathbf{m}_{\varepsilon}}{\det \mathbf{F}_{\varepsilon k}} : \nabla (\mathbf{m}_{\varepsilon k} - \mathbf{m}_{\varepsilon}) \, d\mathbf{x} dt \xrightarrow{k \rightarrow \infty} 0
\end{aligned}$$

with some $\mathbf{d}_{\varepsilon k} \in \text{Dir}(\mathbf{r}_{\varepsilon k})$, where $c_{\kappa, \lambda} := \inf_{F \in \mathbb{R}^{d \times d}} \kappa_{\lambda}(F) / \det F$ is positive thanks to our definition (3.24b). This convergence to 0 for $k \rightarrow \infty$ is due to (3.42d,e). Therefore,

$$\mathbf{m}_{\varepsilon k} \rightarrow \mathbf{m}_{\varepsilon} \quad \text{strongly in } L^2(I; H^1(\Omega; \mathbb{R}^d)), \quad (3.51a)$$

which also improves the weak*-convergence in (3.45f) and ensures convergence $\mathbf{K}_{\lambda}(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) \rightarrow \mathbf{K}_{\lambda}(\mathbf{F}_{\varepsilon}, \nabla \mathbf{m}_{\varepsilon})$ in $L^1(I \times \Omega; \mathbb{R}^{d \times d})$. Moreover, by interpolation with (3.42d),

$$\nabla \mathbf{m}_{\varepsilon k} \rightarrow \nabla \mathbf{m}_{\varepsilon} \quad \text{strongly in } L^c(I; L^2(\Omega; \mathbb{R}^{d \times d})) \text{ for any } 1 \leq c < \infty. \quad (3.51b)$$

Thus $|\nabla \mathbf{m}_{\varepsilon k}|^2 \rightarrow |\nabla \mathbf{m}_{\varepsilon}|^2$ and $\nabla \mathbf{m}_{\varepsilon k} \otimes \nabla \mathbf{m}_{\varepsilon k} \rightarrow \nabla \mathbf{m}_{\varepsilon} \otimes \nabla \mathbf{m}_{\varepsilon}$ strongly in $L^c(I; L^1(\Omega; \mathbb{R}^{d \times d}))$ for any $1 \leq c < \infty$, which is needed for the convergence in (3.27e) when multiplied by $\mathbf{e}(\mathbf{v}_{\varepsilon k}) \in L^p(I; L^{\infty}(\Omega; \mathbb{R}^{d \times d}))$.

We now use the Galerkin approximation of the regularized momentum equation (3.27d) tested by $\widetilde{\mathbf{v}} = \mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k$ with $\widetilde{\mathbf{v}}_k : I \rightarrow V_k$ an approximation of \mathbf{v}_{ε} in the sense that $\widetilde{\mathbf{v}}_k \rightarrow \mathbf{v}_{\varepsilon}$ strongly in $L^{\infty}(I; L^2(\Omega; \mathbb{R}^d))$ and $\nabla^2 \widetilde{\mathbf{v}}_k \rightarrow \nabla^2 \mathbf{v}_{\varepsilon}$ for $k \rightarrow \infty$ strongly in $L^p(I \times \Omega; \mathbb{R}^{d \times d \times d})$ for $k \rightarrow \infty$. Using also the first inequality in (3.22) and (3.49), we can estimate

$$\begin{aligned}
&\frac{1}{2C_{r, \varepsilon}} \|\mathbf{v}_{\varepsilon k}(T) - \mathbf{v}_{\varepsilon}(T)\|_{L^2(\Omega; \mathbb{R}^d)}^2 + \nu_1 c_p \|\mathbf{e}(\mathbf{v}_{\varepsilon k} - \mathbf{v}_{\varepsilon})\|_{L^p(I \times \Omega; \mathbb{R}^{d \times d})}^p + \nu_2 c_p \|\nabla^2(\mathbf{v}_{\varepsilon k} - \mathbf{v}_{\varepsilon})\|_{L^p(I \times \Omega; \mathbb{R}^{d \times d \times d})}^p \\
&\leq \int_{\Omega} \frac{\varrho_{\varepsilon k}(T)}{2} |\mathbf{v}_{\varepsilon k}(T) - \mathbf{v}_{\varepsilon}(T)|^2 \, d\mathbf{x} + \int_0^T \int_{\Gamma} \nu_b |\mathbf{v}_{\varepsilon k} - \mathbf{v}_{\varepsilon}|^p \, dS \, dt \\
&\quad + \int_0^T \int_{\Omega} \left(\nu_1 (|\mathbf{e}(\mathbf{v}_{\varepsilon k})|^{p-2} \mathbf{e}(\mathbf{v}_{\varepsilon k}) - |\mathbf{e}(\mathbf{v}_{\varepsilon})|^{p-2} \mathbf{e}(\mathbf{v}_{\varepsilon})) : \mathbf{e}(\mathbf{v}_{\varepsilon k} - \mathbf{v}_{\varepsilon}) \right. \\
&\quad \left. + \nu_2 (|\nabla^2 \mathbf{v}_{\varepsilon k}|^{p-2} \nabla^2 \mathbf{v}_{\varepsilon k} - |\nabla^2 \mathbf{v}_{\varepsilon}|^{p-2} \nabla^2 \mathbf{v}_{\varepsilon}) : \nabla^2(\mathbf{v}_{\varepsilon k} - \mathbf{v}_{\varepsilon}) \right) \, d\mathbf{x} \, dt \\
&= \int_0^T \int_{\Omega} \left(\left(\sqrt{\frac{\rho \varrho_{\varepsilon k}}{\det \mathbf{F}_{\varepsilon k}}} \mathbf{g} + \mu_0 (\nabla \mathbf{h}_{\varepsilon k})^{\top} \mathbf{m}_{\varepsilon k} \right) \cdot (\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) + \mu_0 \mathbf{h}_{\varepsilon k} \cdot \mathbf{m}_{\varepsilon k} \operatorname{div}(\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) \right. \\
&\quad - (\mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) + \mathbf{K}_{\lambda}(\mathbf{F}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) + \mathbf{S}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})) : \mathbf{e}(\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) \\
&\quad - \nu_1 |\mathbf{e}(\widetilde{\mathbf{v}}_k)|^{p-2} \mathbf{e}(\widetilde{\mathbf{v}}_k) : \mathbf{e}(\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) - \nu_2 (|\nabla^2 \widetilde{\mathbf{v}}_k|^{p-2} \nabla^2 \widetilde{\mathbf{v}}_k) : \nabla^2(\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) \\
&\quad + \left(\frac{\partial}{\partial t} (\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}) + \operatorname{div}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k} \otimes \mathbf{v}_{\varepsilon k}) \right) \cdot \widetilde{\mathbf{v}}_k - \mathcal{J}_{\lambda}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \nabla \mathbf{m}_{\varepsilon k}) : \nabla^2(\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) \Big) \, d\mathbf{x} \, dt \\
&\quad + \int_0^T \int_{\Gamma} (\mathbf{k} + \nu_b |\widetilde{\mathbf{v}}|^{p-2} \widetilde{\mathbf{v}}_k) \cdot (\mathbf{v}_{\varepsilon k} - \widetilde{\mathbf{v}}_k) \, dS \, dt + \int_{\Omega} \left(\frac{\varrho_0}{2} |\mathbf{v}_0|^2 - \varrho_{\varepsilon k}(T) \mathbf{v}_{\varepsilon k}(T) \cdot \widetilde{\mathbf{v}}_k(T) \right. \\
&\quad \left. + \frac{\varrho_{\varepsilon k}(T)}{2} |\widetilde{\mathbf{v}}_k(T)|^2 \right) \, d\mathbf{x} + \mathcal{O}_k \xrightarrow{k \rightarrow \infty} 0 \quad (3.52)
\end{aligned}$$

with $C_{r, \varepsilon} > 0$ from (3.31d) and with some $c_p > 0$ related to the inequality $c_p |G - \widetilde{G}|^p \leq (|G|^{p-2} G - |\widetilde{G}|^{p-2} \widetilde{G}) : (G - \widetilde{G})$ holding for $p \geq 2$. The remainder term \mathcal{O}_k in (3.52) is

$$\mathcal{O}_k = \int_{\Omega} \frac{\varrho_{\varepsilon k}(T)}{2} \mathbf{v}_{\varepsilon k}(T) \cdot (\widetilde{\mathbf{v}}_k(T) - \mathbf{v}_{\varepsilon}(T)) \, d\mathbf{x} + \int_0^T \int_{\Gamma} \nu_b |\mathbf{v}_{\varepsilon k}|^{p-2} \mathbf{v}_{\varepsilon k} \cdot (\widetilde{\mathbf{v}}_k - \mathbf{v}_{\varepsilon}) \, dS \, dt$$

$$+ \int_0^T \int_{\Omega} \nu_1 |\mathbf{e}(\mathbf{v}_{\varepsilon k})|^{p-2} \mathbf{e}(\mathbf{v}_{\varepsilon k}) : \mathbf{e}(\tilde{\mathbf{v}}_k - \mathbf{v}_{\varepsilon}) + \nu_2 |\nabla^2 \mathbf{v}_{\varepsilon k}|^{p-2} \nabla^2 \mathbf{v}_{\varepsilon k} : \nabla^2 (\tilde{\mathbf{v}}_k - \mathbf{v}_{\varepsilon}) \, d\mathbf{x} dt$$

and it converges to zero due to the strong approximation properties of the approximation $\tilde{\mathbf{v}}_k$ of \mathbf{v}_{ε} . Here we used (3.48)–(3.49) and also the strong convergence (3.43a), (3.45b), and (3.47). Knowing already (3.45c) and that $\mathbf{e}(\mathbf{v}_{\varepsilon k} - \tilde{\mathbf{v}}_k) \rightarrow 0$ weakly in $L^p(I; W^{1,p}(\Omega; \mathbb{R}_{\text{sym}}^{d \times d}))$, we have that $\int_0^T \int_{\Omega} \mathbf{T}_{\lambda, \varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) : \mathbf{e}(\mathbf{v}_{\varepsilon k} - \tilde{\mathbf{v}}_k) \, d\mathbf{x} dt \rightarrow 0$. Thus we obtain the desired strong convergence

$$\mathbf{v}_{\varepsilon k} \rightarrow \mathbf{v}_{\varepsilon} \quad \text{strongly in } L^p(I; W^{2,p}(\Omega; \mathbb{R}^d)) \quad (3.53a)$$

and also of $\mathbf{v}_{\varepsilon k}(T) \rightarrow \mathbf{v}_{\varepsilon}(T)$ in $L^2(\Omega; \mathbb{R}^d)$. In fact, executing this procedure for a current time instants t instead of T , we obtain

$$\mathbf{v}_{\varepsilon k}(t) \rightarrow \mathbf{v}_{\varepsilon}(t) \quad \text{strongly in } L^2(\Omega; \mathbb{R}^d) \text{ for any } t \in I. \quad (3.53b)$$

By (3.42d) and the Aubin-Lions theorem, we also obtain

$$\mathbf{m}_{\varepsilon k} \rightarrow \mathbf{m}_{\varepsilon} \quad \text{strongly in } L^c(I; L^{2^*-1/c}(\Omega; \mathbb{R}^d)) \text{ for any } 1 \leq c < \infty. \quad (3.53c)$$

It also implies, by continuity of the trace operator $L^p(I; W^{2,p}(\Omega)) \rightarrow L^p(I \times \Gamma)$, that

$$\mathbf{v}_{\varepsilon k}|_{I \times \Gamma} \rightarrow \mathbf{v}_{\varepsilon}|_{I \times \Gamma} \quad \text{strongly in } L^p(I \times \Gamma; \mathbb{R}^d). \quad (3.53d)$$

Having (3.53) at disposal, the limit passage in the Galerkin-approximation of (3.27d) to the weak solution of (3.25b) is then easy. The variational inequality (3.27e) can be converged by lower weak-semicontinuity of its right-hand-side integral functionals. Convergence in (3.27a–c) is due to Lemmas 3.3 and 3.4.

For further purposes, let us mention that the energy dissipation balance (3.29) is inherited in the limit, i.e.

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \frac{\varrho_{\varepsilon}}{2} |\mathbf{v}_{\varepsilon}|^2 + \frac{\pi_{\lambda}(\mathbf{F}_{\varepsilon}) \varphi(\mathbf{F}_{\varepsilon}, \mathbf{m}_{\varepsilon})}{\det \mathbf{F}_{\varepsilon}} + \frac{\kappa_{\lambda}(\mathbf{F}_{\varepsilon})}{2 \det \mathbf{F}_{\varepsilon}} |\nabla \mathbf{m}_{\varepsilon}|^2 - \mu_0 \mathbf{h}_{\text{ext}} \cdot \mathbf{m}_{\varepsilon} \, d\mathbf{x} \\ & + \int_{\Omega} \xi_{\varepsilon}(\mathbf{F}_{\varepsilon}, \theta_{\varepsilon}; \mathbf{e}(\mathbf{v}_{\varepsilon}), \nabla^2 \mathbf{v}_{\varepsilon}, \mathbf{r}_{\varepsilon}) \, d\mathbf{x} + \int_{\Gamma} \nu_b |\mathbf{v}_{\varepsilon}|^p \, dS + \frac{d}{dt} \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u_{\varepsilon}|^2 \, d\mathbf{x} \\ & = \int_{\Gamma} \mathbf{k} \cdot \mathbf{v}_{\varepsilon} \, dS + \int_{\Omega} \left(\sqrt{\frac{\varrho_{\varepsilon} \varrho}{\det_{\lambda}(\mathbf{F}_{\varepsilon})}} \mathbf{g} \cdot \mathbf{v}_{\varepsilon} - \frac{\pi_{\lambda}(\mathbf{F}_{\varepsilon}) \zeta'_{\mathbf{F}}(\mathbf{F}_{\varepsilon}, \mathbf{m}_{\varepsilon}, \theta_{\varepsilon}) \mathbf{F}_{\varepsilon}^{\top}}{(1 + \varepsilon \theta_{\varepsilon}) \det \mathbf{F}_{\varepsilon}} : \mathbf{e}(\mathbf{v}_{\varepsilon}) \right. \\ & \quad \left. - \frac{\pi_{\lambda}(\mathbf{F}_{\varepsilon}) \zeta'_{\mathbf{m}}(\mathbf{F}_{\varepsilon}, \mathbf{m}_{\varepsilon}, \theta_{\varepsilon})}{(1 + \varepsilon \theta_{\varepsilon}^{1/2}) \det \mathbf{F}_{\varepsilon}} \cdot (\mathbf{r}_{\varepsilon} - \text{skw}(\nabla \mathbf{v}_{\varepsilon}) \mathbf{m}_{\varepsilon}) - \mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}_{\varepsilon} \right) \, d\mathbf{x} \end{aligned} \quad (3.54)$$

with $u_{\varepsilon} = \Delta^{-1} \text{div}(\mathbf{m}_{\varepsilon})$. This follows by legitimacy of the tests which led formally (in the non-regularized situation) to (2.49). Specifically, here (3.25a–d) in the weak formulation are in duality (and can be tested) by $|\mathbf{v}_{\varepsilon}|^2/2$, \mathbf{v}_{ε} , $[\varphi(\mathbf{F}_{\varepsilon}, \mathbf{m}_{\varepsilon})/\det \mathbf{F}_{\varepsilon}]'_{\mathbf{F}}$, and \mathbf{r}_{ε} , cf. Step 10 below.

Step 5: Strong convergence of $\mathbf{r}_{\varepsilon k}$ and limit passage in the thermal part for $k \rightarrow \infty$. For the heat equation, we will still need the strong convergence of $\mathbf{r}_{\varepsilon k} \rightarrow \mathbf{r}_{\varepsilon}$. We cannot exploit the

strong monotonicity of the operator $\mathbf{r} \mapsto \tau \mathbf{r} + h_c(\mathbf{F}, \theta) \text{dir}_\varepsilon(\mathbf{r})$ because the exchange driving force on the right-hand side of (3.25d) is not a compact lower-order term. Thus, as we already passed to the limit in the magneto-mechanical part, we can use the “limsup-trick” and the strict convexity of the potential of the mentioned operator. Specifically,

$$\begin{aligned}
& \int_0^T \left(\int_\Omega \xi_\varepsilon(\mathbf{F}_\varepsilon, \theta_\varepsilon; \mathbf{e}(\mathbf{v}_\varepsilon), \nabla^2 \mathbf{v}_\varepsilon, \mathbf{r}_\varepsilon) \, d\mathbf{x} + \int_\Gamma \nu_b |\mathbf{v}_\varepsilon|^p \, dS \right) dt \\
& \leq \liminf_{k \rightarrow \infty} \int_0^T \left(\int_\Omega \xi_\varepsilon(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) \, d\mathbf{x} + \int_\Gamma \nu_b |\mathbf{v}_{\varepsilon k}|^p \, dS \right) dt \\
& \leq \limsup_{k \rightarrow \infty} \int_0^T \left(\int_\Omega \xi_\varepsilon(\mathbf{F}_{\varepsilon k}, \theta_{\varepsilon k}; \mathbf{e}(\mathbf{v}_{\varepsilon k}), \nabla^2 \mathbf{v}_{\varepsilon k}, \mathbf{r}_{\varepsilon k}) \, d\mathbf{x} + \int_\Gamma \nu_b |\mathbf{v}_{\varepsilon k}|^p \, dS \right) dt \\
& = \int_\Omega \frac{\varrho_0}{2} |\mathbf{v}_0|^2 + \frac{\varphi(\mathbf{F}_0, \mathbf{m}_0)}{\det \mathbf{F}_0} + \frac{\kappa(\mathbf{F}_0) |\nabla \mathbf{m}_0|^2}{2 \det \mathbf{F}_0} - \mu_0 \mathbf{h}(0) \cdot \mathbf{m}_0 \, d\mathbf{x} + \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u(0)|^2 \, d\mathbf{x} \\
& \quad - \liminf_{k \rightarrow \infty} \left(\int_\Omega \left(\frac{\varrho_{\varepsilon k}(T)}{2} |\mathbf{v}_{\varepsilon k}(T)|^2 + \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}(T)) \varphi(\mathbf{F}_{\varepsilon k}(T), \mathbf{m}_{\varepsilon k}(T))}{\det \mathbf{F}_{\varepsilon k}(T)} \right. \right. \\
& \quad \left. \left. + \frac{\kappa_\lambda(\mathbf{F}_{\varepsilon k}(T)) |\nabla \mathbf{m}_{\varepsilon k}(T)|^2}{2 \det \mathbf{F}_{\varepsilon k}(T)} - \mu_0 \mathbf{h}(T) \cdot \mathbf{m}_{\varepsilon k}(T) \, d\mathbf{x} + \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u_{\varepsilon k}(T)|^2 \, d\mathbf{x} \right) \right) \\
& \quad + \lim_{k \rightarrow \infty} \int_0^T \left(\int_\Omega \varrho_{\varepsilon k} \mathbf{g} \cdot \mathbf{v}_{\varepsilon k} + \mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}_{\varepsilon k} - \pi_\lambda(\mathbf{F}_{\varepsilon k}) \frac{\zeta'_{\mathbf{F}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k}) \mathbf{F}_{\varepsilon k}^\top}{(1 + \varepsilon |\theta_{\varepsilon k}|^\alpha) \det \mathbf{F}_{\varepsilon k}} \cdot \mathbf{e}(\mathbf{v}_{\varepsilon k}) \right. \\
& \quad \left. - \frac{\pi_\lambda(\mathbf{F}_{\varepsilon k}) \zeta'_{\mathbf{m}}(\mathbf{F}_{\varepsilon k}, \mathbf{m}_{\varepsilon k}, \theta_{\varepsilon k})}{(1 + \varepsilon |\theta_{\varepsilon k}|^\alpha)^2 \det \mathbf{F}_{\varepsilon k}} \cdot (\mathbf{r}_{\varepsilon k} - \text{skw}(\nabla \mathbf{v}_{\varepsilon k}) \mathbf{m}_{\varepsilon k}) + \int_\Gamma \mathbf{k} \cdot \mathbf{v}_{\varepsilon k} \, dS \right) dt \\
& \leq \int_\Omega \frac{\varrho_0}{2} |\mathbf{v}_0|^2 + \frac{\varphi(\mathbf{F}_0, \mathbf{m}_0)}{\det \mathbf{F}_0} + \frac{\kappa(\mathbf{F}_0) |\nabla \mathbf{m}_0|^2}{2 \det \mathbf{F}_0} - \mu_0 \mathbf{h}(0) \cdot \mathbf{m}_0 \, d\mathbf{x} + \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u(0)|^2 \, d\mathbf{x} \\
& \quad - \int_\Omega \left(\frac{\varrho_\varepsilon(T)}{2} |\mathbf{v}_\varepsilon(T)|^2 + \frac{\pi_\lambda(\mathbf{F}_\varepsilon(T)) \varphi(\mathbf{F}_\varepsilon(T), \mathbf{m}_\varepsilon(T))}{\det \mathbf{F}_\varepsilon(T)} \right. \\
& \quad \left. + \frac{\kappa_\lambda(\mathbf{F}_\varepsilon(T)) |\nabla \mathbf{m}_\varepsilon(T)|^2}{2 \det \mathbf{F}_\varepsilon(T)} - \mu_0 \mathbf{h}(T) \cdot \mathbf{m}_\varepsilon(T) \right) d\mathbf{x} + \int_{\mathbb{R}^d} \frac{\mu_0}{2} |\nabla u_\varepsilon(T)|^2 \, d\mathbf{x} \\
& \quad + \int_0^T \left(\int_\Omega \varrho_\varepsilon \mathbf{g} \cdot \mathbf{v}_\varepsilon + \mu_0 \frac{\partial \mathbf{h}_{\text{ext}}}{\partial t} \cdot \mathbf{m}_\varepsilon - \pi_\lambda(\mathbf{F}_\varepsilon) \frac{\zeta'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top}{(1 + \varepsilon \theta_\varepsilon) \det \mathbf{F}_\varepsilon} \cdot \mathbf{e}(\mathbf{v}_\varepsilon) \right. \\
& \quad \left. - \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)}{(1 + \varepsilon \theta_\varepsilon^{1/2}) \det \mathbf{F}_\varepsilon} \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon) + \int_\Gamma \mathbf{k} \cdot \mathbf{v}_\varepsilon \, dS \right) dt \\
& \stackrel{(3.54)}{=} \int_0^T \left(\int_\Omega \xi_\varepsilon(\mathbf{F}_\varepsilon, \theta_\varepsilon; \mathbf{e}(\mathbf{v}_\varepsilon), \nabla^2 \mathbf{v}_\varepsilon, \mathbf{r}_\varepsilon) \, d\mathbf{x} + \int_\Gamma \nu_b |\mathbf{v}_\varepsilon|^p \, dS \right) dt. \tag{3.55}
\end{aligned}$$

This still gives

$$\mathbf{r}_{\varepsilon k} \rightarrow \mathbf{r}_\varepsilon \quad \text{strongly in } L^2(I \times \Omega; \mathbb{R}^d). \tag{3.56}$$

Strong convergence (3.53a,b,d) allows for passing to the limit also in the dissipative heat sources and the other terms in the Galerkin-approximation of the heat equation (3.27f) are even easier.

Step 6 – non-negativity of temperature: We can now perform various nonlinear tests of the regularized but non-discretized heat equation. The first test can be by the negative part

of temperature $\theta_\varepsilon^- := \min(0, \theta_\varepsilon)$. Let us recall the extension (3.10), which in particular gives $\omega(\mathbf{F}, \mathbf{m}, \theta^-) = \theta^-$ and $\omega'_\mathbf{F}(\mathbf{F}, \mathbf{m}, \theta^-) = \mathbf{0}$ and also $\omega'_\mathbf{m}(\mathbf{F}, \mathbf{m}, \theta^-) = \mathbf{0}$. Note also that $\theta_\varepsilon^- \in L^2(I; H^1(\Omega))$, so that it is indeed a legal test for (3.25f). Here we rely on the data qualification $\nu_1, \nu_2, \nu_b \geq 0$, $\mathcal{K} = \mathcal{K}(\mathbf{F}, \theta) \geq 0$, $\theta_0 \geq 0$, and $h(\theta) \geq 0$ for $\theta \leq 0$, cf. (3.5f,i,n). Realizing that $\nabla \theta^- = 0$ wherever $\theta > 0$ so that $\nabla \theta \cdot \nabla \theta^- = |\nabla \theta^-|^2$ and that $\zeta'_\mathbf{F}(\mathbf{F}, \mathbf{m}, \theta) \theta^- = \zeta'_\mathbf{F}(\mathbf{F}, \mathbf{m}, \theta^-) \theta^- = \mathbf{0}$ and $\zeta'_\mathbf{m}(\mathbf{F}, \mathbf{m}, \theta) \theta^- = \zeta'_\mathbf{m}(\mathbf{F}, \mathbf{m}, \theta^-) \theta^- = \mathbf{0}$ and also $h(\theta) \theta^- = h(\theta^-) \theta^- = 0$, this test gives

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} \|\theta_\varepsilon^-\|_{L^2(\Omega)}^2 &\leq \int_\Omega \theta_\varepsilon^- \frac{\partial w_\varepsilon}{\partial t} + \mathcal{K}(\mathbf{F}_\varepsilon, \theta_\varepsilon) \nabla \theta_\varepsilon \cdot \nabla \theta_\varepsilon^- \, d\mathbf{x} \\
&= \int_\Omega \left(w_\varepsilon \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon^- + \left(\xi_\varepsilon(\mathbf{F}_\varepsilon, \theta_\varepsilon; \mathbf{e}(\mathbf{v}_\varepsilon), \nabla^2 \mathbf{v}_\varepsilon, \mathbf{r}_\varepsilon) + \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_\mathbf{F}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top}{(1+\varepsilon|\theta_\varepsilon|) \det \mathbf{F}_\varepsilon} : \mathbf{e}(\mathbf{v}_{\varepsilon k}) \right. \right. \\
&\quad \left. \left. + \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_\mathbf{m}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)}{(1+\varepsilon|\theta_\varepsilon|^{1/2}) \det \mathbf{F}_\varepsilon} \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon) \right) \theta_\varepsilon^- \right) d\mathbf{x} + \int_\Gamma \left(h(\theta_\varepsilon) + \frac{\nu_b |\mathbf{v}_\varepsilon|^p}{2+\varepsilon|\mathbf{v}_\varepsilon|^p} \right) \theta_\varepsilon^- \, dS \\
&\leq \int_\Omega w_\varepsilon \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon^- \, d\mathbf{x} = \int_\Omega \theta_\varepsilon^- \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon^- \, d\mathbf{x} = - \int_\Omega |\nabla \theta_\varepsilon^-|^2 \operatorname{div} \mathbf{v}_\varepsilon \, d\mathbf{x} \\
&= - \frac{1}{2} \int_\Omega |\nabla \theta_\varepsilon^-|^2 \operatorname{div} \mathbf{v}_\varepsilon \, d\mathbf{x} \leq \|\theta_\varepsilon^-\|_{L^2(\Omega)}^2 \|\operatorname{div} \mathbf{v}_\varepsilon\|_{L^\infty(\Omega)}. \tag{3.57}
\end{aligned}$$

Recalling the assumption $\theta_0 \geq 0$ so that $\theta_{0,\varepsilon}^- = 0$ and exploiting the information $\mathbf{v}_\varepsilon \in L^p(I; W^{1,p}(\Omega; \mathbb{R}^d))$ with $p > d$ inherited from (3.31a), by the Gronwall inequality we obtain $\|\theta_\varepsilon^-\|_{L^\infty(I; L^2(\Omega))} = 0$, so that $\theta_\varepsilon \geq 0$ a.e. on $I \times \Omega$.

Having proved non-negativity of temperature, we can now execute the strategy based of the L^1 -theory for the heat equation which led to the estimates (3.18)–(3.19), i.e. here

$$\|\mathbf{v}_\varepsilon\|_{L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))} \leq C, \quad \|\mathbf{m}_\varepsilon\|_{L^\infty(I; H^1(\Omega; \mathbb{R}^d))} \leq C, \tag{3.58a}$$

$$\|\mathbf{F}_\varepsilon\|_{L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d}))} \leq C_r, \quad \left\| \frac{1}{\det \mathbf{F}_\varepsilon} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r, \tag{3.58b}$$

$$\|\varrho_\varepsilon\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r, \quad \left\| \frac{1}{\varrho_\varepsilon} \right\|_{L^\infty(I; W^{1,r}(\Omega))} \leq C_r \quad \text{for any } 1 \leq r < +\infty, \tag{3.58c}$$

$$\|w_\varepsilon\|_{L^\infty(I; L^1(\Omega))} \leq C, \quad \text{and} \quad \|\theta_\varepsilon\|_{L^\infty(I; L^1(\Omega))} \leq C. \tag{3.58d}$$

By interpolation exploiting the Gagliardo-Nirenberg inequality between $L^2(\Omega)$ and $W^{2,p}(\Omega)$, we have $\|\cdot\|_{L^\infty(\Omega)} \leq C \|\cdot\|_{L^2(\Omega)}^r \|\cdot\|_{W^{2,p}(\Omega)}^{1-r}$ with $0 < r < pd/(pd+4p-2d)$. Using also Korn's inequality, from (3.58a) we thus obtain the estimate

$$\|\mathbf{v}_\varepsilon\|_{L^s(I; L^\infty(\Omega; \mathbb{R}^d))} \leq C_s \quad \text{with } 1 \leq s < \frac{p(pd+4p-2d)}{4p-2d}. \tag{3.58e}$$

By comparison from $\frac{\partial}{\partial t} \varrho_\varepsilon = (\operatorname{div} \mathbf{v}_\varepsilon) \varrho_\varepsilon - \mathbf{v}_\varepsilon \cdot \nabla \varrho_\varepsilon$, from $\frac{\partial}{\partial t} \mathbf{F}_\varepsilon = (\nabla \mathbf{v}_\varepsilon) \mathbf{F}_\varepsilon - (\mathbf{v}_\varepsilon \cdot \nabla) \mathbf{F}_\varepsilon$, and from $\frac{\partial}{\partial t} \mathbf{m}_\varepsilon = \mathbf{r}_\varepsilon + \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon - (\mathbf{v}_\varepsilon \cdot \nabla) \mathbf{m}_\varepsilon$, we also have

$$\left\| \frac{\partial \varrho_\varepsilon}{\partial t} \right\|_{L^p(I; L^r(\Omega))} \leq C, \quad \left\| \frac{\partial \mathbf{F}_\varepsilon}{\partial t} \right\|_{L^p(I; L^r(\Omega; \mathbb{R}^{d \times d}))} \leq C, \quad \text{and} \quad \left\| \frac{\partial \mathbf{m}_\varepsilon}{\partial t} \right\|_{L^2(I \times \Omega; \mathbb{R}^d)} \leq C. \tag{3.58f}$$

The estimates (3.58d) are naturally weaker than (3.40) but, importantly, are uniform with respect to $\varepsilon > 0$, in contrast to (3.40) which is not uniform in this sense. The total energy balance (2.50) holds for ε -solution only as an inequality because the heat sources do not exactly cancel; more in detail, while the regularized adiabatic heat again cancels, the dissipative heat terms are regularized (and smaller) in (3.25f) and in (3.26c) but the corresponding viscous stress in (3.25b) and force in (3.26a) are not regularized. This inequality still allows to execute the above mentioned estimation.

Let us also note that the extension (3.10) becomes now inactive and we can work with the original data defined for non-negative θ only.

Step 7 – further a-priori estimates: Furthermore, having \mathbf{r}_ε estimated in $L^2(I \times \Omega; \mathbb{R}^d)$ uniformly with respect to ε , as in (3.20) we have

$$\begin{aligned} \Delta \mathbf{m}_\varepsilon = & \frac{\det \mathbf{F}_\varepsilon}{\kappa_\lambda(\mathbf{F}_\varepsilon)} \left(\tau \mathbf{r}_\varepsilon + h_c(\mathbf{F}_\varepsilon, \theta_\varepsilon) \operatorname{dir}_\varepsilon(\mathbf{r}_\varepsilon) - \frac{\mathbf{m}_\varepsilon \times \mathbf{r}_\varepsilon}{\gamma(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)} - \mathbf{h}_{\text{ext}} - \nabla u_\varepsilon \right. \\ & \left. + \frac{\varphi'_\mathbf{m}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon) + \zeta'_\mathbf{m}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)}{\det \mathbf{F}_\varepsilon} - \left(\frac{\kappa'_\lambda(\mathbf{F}_\varepsilon)}{\det \mathbf{F}_\varepsilon} - \frac{\kappa_\lambda(\mathbf{F}_\varepsilon) \operatorname{Cof} \mathbf{F}_\varepsilon}{\det \mathbf{F}_\varepsilon^2} \right) : (\nabla \mathbf{F}_\varepsilon \otimes \nabla \mathbf{m}_\varepsilon) \right), \end{aligned}$$

so that we can estimate $\nabla^2 \mathbf{m}_\varepsilon$ by a H^1 -regularity as (3.21), i.e. now

$$\|\nabla^2 \mathbf{m}_\varepsilon\|_{L^2(I \times \Omega; \mathbb{R}^{d \times d \times d})} \leq C. \quad (3.59)$$

Furthermore, we are to prove an estimate of $\nabla \theta_\varepsilon$ based on the test of the heat equation (3.25f) by $\chi_\zeta(\theta_\varepsilon)$ with an increasing nonlinear function $\chi_\zeta : [0, +\infty) \rightarrow [0, 1]$ defined as

$$\chi_\zeta(\theta) := 1 - \frac{1}{(1+\theta)^\zeta}, \quad \zeta > 0, \quad (3.60)$$

simplifying the original idea of L. Boccardo and T. Gallouët [8,9] in the spirit of [18], expanding the estimation strategy in [26, Sect. 8.2]. Importantly, here we have $\chi_\zeta(\theta_\varepsilon(t, \cdot)) \in H^1(\Omega)$, hence it is a legal test function, because $0 \leq \theta_\varepsilon(t, \cdot) \in H^1(\Omega)$ has already been proved and because χ_ζ is Lipschitz continuous on $[0, +\infty)$.

We consider $1 \leq \mu < 2$ and estimate the L^μ -norm of $\nabla \theta_\varepsilon$ by Hölder's inequality as

$$\begin{aligned} \int_0^T \int_\Omega |\nabla \theta_\varepsilon|^\mu \, d\mathbf{x} \, dt \leq & C_1 \underbrace{\left(\int_0^T \|1 + \theta_\varepsilon(t, \cdot)\|_{L^{(1+\zeta)\mu/(2-\mu)}(\Omega)}^{(1+\zeta)\mu/(2-\mu)} \, dt \right)^{1-\mu/2}}_{=: I_{\mu, \zeta}^{(1)}(\theta_\varepsilon)} \underbrace{\left(\int_0^T \int_\Omega \chi'_\zeta(\theta_\varepsilon) |\nabla \theta_\varepsilon|^2 \right)^{\mu/2}}_{=: I_\zeta^{(2)}(\theta_\varepsilon)}. \end{aligned} \quad (3.61)$$

with χ_ζ from (3.60) so that $\chi'_\zeta(\theta) = \zeta/(1+\theta)^{1+\zeta}$ and with a constant C_1 dependent on ζ , μ , and T . Then we interpolate the Lebesgue space $L^{(1+\zeta)\mu/(2-\mu)}(\Omega)$ between $W^{1,\mu}(\Omega)$ and $L^1(\Omega)$ in order to exploit the already obtained $L^\infty(I; L^1(\Omega))$ -estimate in (3.58d). More specifically, by the Gagliardo-Nirenberg inequality, we obtain

$$\|1 + \theta_\varepsilon(t, \cdot)\|_{L^{\mu/\sigma}(\Omega)}^{\mu/\sigma} \leq C_2 \left(1 + \|\nabla \theta_\varepsilon(t, \cdot)\|_{L^\mu(\Omega; \mathbb{R}^d)} \right)^\mu \quad \text{with } \sigma = \frac{2-\mu}{1+\zeta} \quad (3.62)$$

with C_2 depending on σ , C_1 , and C from (3.58d), so that $I_{\mu,\zeta}^{(1)}(\theta_\varepsilon) \leq C_3(1 + \int_0^T \int_\Omega |\nabla \theta_\varepsilon|^\mu \, d\mathbf{x} dt)$ with C_3 depending on C_2 . Combining it with (3.61), we obtain

$$\|\nabla \theta_\varepsilon\|_{L^\mu(I \times \Omega; \mathbb{R}^d)}^\mu = C_1 C_3 (1 + \|\nabla \theta_\varepsilon\|_{L^\mu(I \times \Omega)}^\mu)^{1-\mu/2} I_{\mu,\zeta}^{(2)}(\theta_\varepsilon)^{\mu/2}. \quad (3.63)$$

Furthermore, we estimate $I_\zeta^{(2)}(\theta_\varepsilon)$ in (3.61). Let us denote by \mathcal{X}_ζ a primitive function to $\theta \mapsto \chi_\zeta(\theta) \omega'_\theta(\mathbf{F}, \mathbf{m}, \theta)$ depending smoothly on \mathbf{F} , specifically

$$\mathcal{X}_\zeta(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 \theta \chi_\zeta(r\theta) \omega'_\theta(\mathbf{F}, \mathbf{m}, r\theta) \, dr. \quad (3.64)$$

Like (3.34) but using partial (not convective) time derivative, we have now the calculus

$$\begin{aligned} \int_\Omega \chi_\zeta(\theta) \frac{\partial w}{\partial t} \, d\mathbf{x} &= \int_\Omega \chi_\zeta(\theta) \omega'_\theta(\mathbf{F}, \mathbf{m}, \theta) \frac{\partial \theta}{\partial t} + \chi_\zeta(\theta) \left(\omega'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) : \frac{\partial \mathbf{F}}{\partial t} + \omega'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) \cdot \frac{\partial \mathbf{m}}{\partial t} \right) \, d\mathbf{x} \\ &= \frac{d}{dt} \int_\Omega \mathcal{X}_\zeta(\mathbf{F}, \mathbf{m}, \theta) \, d\mathbf{x} - \int_\Omega [\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) : \frac{\partial \mathbf{F}}{\partial t} + [\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) \cdot \frac{\partial \mathbf{m}}{\partial t} \, d\mathbf{x} \\ &\quad \text{where } \mathcal{X}_\zeta(\mathbf{F}, \mathbf{m}, \theta) := \mathcal{X}_\zeta(\mathbf{F}, \mathbf{m}, \theta) - \chi_\zeta(\theta) \omega(\mathbf{F}, \mathbf{m}, \theta). \end{aligned} \quad (3.65)$$

In view of (3.64), it holds $[\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 \theta \chi_\zeta(r\theta) \omega''_{\mathbf{F}\theta}(\mathbf{F}, \mathbf{m}, r\theta) \, dr - \chi_\zeta(\theta) \omega'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)$ and $[\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta) = \int_0^1 \theta \chi_\zeta(r\theta) \omega''_{\mathbf{m}\theta}(\mathbf{F}, \mathbf{m}, r\theta) \, dr - \chi_\zeta(\theta) \omega'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)$. Altogether, testing (3.25f) with (3.26c) by $\chi_\zeta(\theta_\varepsilon)$ gives

$$\begin{aligned} &\frac{d}{dt} \int_\Omega \mathcal{X}_\zeta(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \, d\mathbf{x} + \int_\Omega \chi'_\zeta(\theta_\varepsilon) \mathcal{K}(\mathbf{F}_\varepsilon, \theta_\varepsilon) |\nabla \theta_\varepsilon|^2 \, d\mathbf{x} \\ &= \int_\Omega \left(\xi_\varepsilon(\mathbf{F}_\varepsilon, \theta_\varepsilon; \mathbf{e}(\mathbf{v}_\varepsilon), \nabla^2 \mathbf{v}_\varepsilon, \mathbf{r}_\varepsilon) \chi_\zeta(\theta_\varepsilon) + \omega(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \chi'_\zeta(\theta_\varepsilon) \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon \right. \\ &\quad + [\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) : \frac{\partial \mathbf{F}_\varepsilon}{\partial t} + [\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \cdot \frac{\partial \mathbf{m}_\varepsilon}{\partial t} \\ &\quad + \chi_\zeta(\theta_\varepsilon) \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)}{(1+\varepsilon \theta_\varepsilon^{1/2}) \det \mathbf{F}_\varepsilon} \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon) \\ &\quad \left. + \chi_\zeta(\theta_\varepsilon) \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top : \mathbf{e}(\mathbf{v}_\varepsilon)}{(1+\varepsilon \theta_\varepsilon) \det \mathbf{F}_\varepsilon} \right) \, d\mathbf{x} + \int_\Gamma \left(h_\varepsilon(\theta_\varepsilon) + \frac{\nu_b |\mathbf{v}_\varepsilon|^p}{2+\varepsilon |\mathbf{v}_\varepsilon|^p} \right) \chi_\zeta(\theta_\varepsilon) \, dS. \end{aligned} \quad (3.66)$$

We realize that $\chi'_\zeta(\theta) = \zeta/(1+\theta)^{1+\zeta}$ as used already in (3.61) and that $\mathcal{X}_\zeta(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \geq c_K \theta_\varepsilon$ with some c_K for $\theta_\varepsilon \geq 0$ due to (3.5c); again K is a compact subset of $\text{GL}^+(d)$ related here with the already proved estimates (3.58b). The convective term in (3.66) is a bit delicate. For any $\delta > 0$, it can be estimated by Hölder inequality as

$$\begin{aligned} \int_\Omega w_\varepsilon \chi'_\zeta(\theta_\varepsilon) \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon \, d\mathbf{x} &\leq \frac{1}{\delta} \int_\Omega \chi'_\zeta(\theta_\varepsilon) |\mathbf{v}_\varepsilon|^2 w_\varepsilon^2 \, d\mathbf{x} + \delta \int_\Omega \chi'_\zeta(\theta_\varepsilon) |\nabla \theta_\varepsilon|^2 \, d\mathbf{x} \\ &= \frac{1}{\delta} \int_\Omega \chi'_\zeta(\theta_\varepsilon) |\mathbf{v}_\varepsilon|^2 w_\varepsilon^2 \, d\mathbf{x} + \delta I_\zeta^{(2)}(\theta_\varepsilon). \end{aligned} \quad (3.67)$$

Denoting by $0 < \mathcal{K}_0 = \inf_{\mathbf{F}, \theta} \mathcal{K}(\mathbf{F}, \theta)$, and using (3.66) integrated over $I = [0, T]$, we further estimate:

$$I_\zeta^{(2)}(\theta_\varepsilon) = \frac{1}{\zeta} \int_0^T \int_\Omega \chi'_\zeta(\theta_\varepsilon) |\nabla \theta_\varepsilon|^2 \, d\mathbf{x} dt \leq \frac{1}{\mathcal{K}_0 \zeta} \int_0^T \int_\Omega \mathcal{K}(\mathbf{F}_\varepsilon, \theta_\varepsilon) \nabla \theta_\varepsilon \cdot \nabla \chi_\zeta(\theta_\varepsilon) \, d\mathbf{x} dt$$

$$\begin{aligned}
&\leq \frac{1}{\mathcal{K}_0\zeta} \left(\int_0^T \int_{\Omega} \mathcal{K}(\mathbf{F}_\varepsilon, \theta_\varepsilon) \nabla \theta_\varepsilon \cdot \nabla \chi_\zeta(\theta_\varepsilon) \, d\mathbf{x} dt + \int_{\Omega} \mathcal{X}_\zeta(\mathbf{F}_\varepsilon(T), \theta_\varepsilon(T)) \, d\mathbf{x} \right) \\
&= \frac{1}{\mathcal{K}_0\zeta} \left(\int_{\Omega} \mathcal{X}_\zeta(\mathbf{F}_0, \theta_{0,\varepsilon}) \, d\mathbf{x} + \int_0^T \int_{\Omega} \left(\xi_\varepsilon(\mathbf{F}_\varepsilon, \theta_\varepsilon; \mathbf{e}(\mathbf{v}_\varepsilon), \nabla^2 \mathbf{v}_\varepsilon, \mathbf{r}_\varepsilon) \chi_\zeta(\theta_\varepsilon) \right. \right. \\
&\quad + [\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \cdot \frac{\partial \mathbf{F}_\varepsilon}{\partial t} + [\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \cdot \frac{\partial \mathbf{m}_\varepsilon}{\partial t} + \omega(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \chi'_\zeta(\theta_\varepsilon) \mathbf{v}_\varepsilon \cdot \nabla \theta_\varepsilon \\
&\quad + \chi_\zeta(\theta_\varepsilon) \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)}{(1+\varepsilon\theta_\varepsilon^{1/2}) \det \mathbf{F}_\varepsilon} \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon) \\
&\quad \left. + \frac{\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top : \mathbf{e}(\mathbf{v}_\varepsilon)}{(1+\varepsilon\theta_\varepsilon) \det \mathbf{F}_\varepsilon} \chi_\zeta(\theta_\varepsilon) \right) d\mathbf{x} + \int_0^T \int_{\Gamma} \left(h_\varepsilon(\theta_\varepsilon) + \frac{\nu_b |\mathbf{v}_\varepsilon|^p}{2+\varepsilon |\mathbf{v}_\varepsilon|^p} \right) \chi_\zeta(\theta_\varepsilon) \, dS dt \Big) \\
&\stackrel{(3.67)}{\leq} \frac{1}{\mathcal{K}_0\zeta} \left(\|\mathcal{X}_\zeta(\mathbf{F}_0, \theta_{0,\varepsilon})\|_{L^1(\Omega)} + \|\nu_1 |\mathbf{e}(\mathbf{v}_\varepsilon)|^p + \tau |\mathbf{r}_\varepsilon|^2 + h_c(\mathbf{F}_\varepsilon, \theta_\varepsilon) |\mathbf{r}_\varepsilon| + \nu_2 |\nabla^2 \mathbf{v}_\varepsilon|^p\|_{L^1(I \times \Omega)} \right. \\
&\quad + \int_0^T \left\| [\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \right\|_{L^{r'}(\Omega; \mathbb{R}^{d \times d})}^{r'} + \left\| \frac{\partial \mathbf{F}_\varepsilon}{\partial t} \right\|_{L^r(\Omega; \mathbb{R}^{d \times d})}^r dt + \|h_{\max}\|_{L^1(I \times \Gamma)} \\
&\quad + \left\| [\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \right\|_{L^2(I \times \Omega; \mathbb{R}^d)}^2 + \left\| \frac{\partial \mathbf{m}_\varepsilon}{\partial t} \right\|_{L^2(I \times \Omega; \mathbb{R}^d)}^2 + \frac{1}{2} \|\sqrt{\nu_b} \mathbf{v}_\varepsilon\|_{L^p(I \times \Gamma; \mathbb{R}^d)}^p \\
&\quad + \left\| \pi_\lambda(\mathbf{F}_\varepsilon) \frac{\zeta'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top : \mathbf{e}(\mathbf{v}_\varepsilon) + \zeta'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon)}{\det \mathbf{F}_\varepsilon} \right\|_{L^1(I \times \Omega)} \\
&\quad \left. + \frac{1}{\delta} \|\mathbf{v}_\varepsilon\|_{L^2(I; L^\infty(\Omega; \mathbb{R}^d))}^2 \|\chi'_\zeta(\theta_\varepsilon) \omega^2(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)\|_{L^\infty(I; L^1(\Omega))} \right) + \frac{\delta}{\mathcal{K}_0} I_\zeta^{(2)}(\theta_\varepsilon); \tag{3.68}
\end{aligned}$$

noteworthy, we choose $\delta < \mathcal{K}_0$, we can absorb the last term in the left-hand side. Due to the assumption (3.5c), we can estimate the adiabatic rates $\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \mathbf{F}_\varepsilon^\top : \mathbf{e}(\mathbf{v}_\varepsilon) / \det \mathbf{F}_\varepsilon$ and $\pi_\lambda(\mathbf{F}_\varepsilon) \zeta'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon) \cdot (\mathbf{r}_\varepsilon - \text{skw}(\nabla \mathbf{v}_\varepsilon) \mathbf{m}_\varepsilon) / \det \mathbf{F}_\varepsilon$ in (3.68), cf. (3.16). We also use the estimates (3.58) and (3.32) and the assumption (3.5i) relying on the already proved non-negativity of temperature. By the qualification (3.5d), we have $|\mathcal{X}_\zeta'_{\mathbf{F}}(\mathbf{F}, \mathbf{m}, \theta)| \leq C(1+\theta)$. This allows for estimation

$$\begin{aligned}
\|[\mathcal{X}_\zeta]'_{\mathbf{F}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)\|_{L^{r'}(\Omega; \mathbb{R}^{d \times d})}^{r'} &\leq C^{r'} \|1 + \theta_\varepsilon\|_{L^{r'}(\Omega)}^{r'} \\
&\leq C_4 + C_4 \|\theta_\varepsilon\|_{L^1(\Omega)}^{r'(1-\mu^{*}/r)} (\|\theta_\varepsilon\|_{L^1(\Omega)} + \|\nabla \theta_\varepsilon\|_{L^\mu(\Omega; \mathbb{R}^d)})^{\mu^{*}/(r-1)}, \tag{3.69}
\end{aligned}$$

where we use the Gagliardo-Nirenberg inequality to interpolate $L^{r'}(\Omega)$ between $L^1(\Omega)$ and $W^{1,\mu}(\Omega)$. Similarly, since $|\mathcal{X}_\zeta'_{\mathbf{m}}(\mathbf{F}, \mathbf{m}, \theta)| \leq C(1+\theta)$ which is again ensured by the qualification (3.5d), we can estimate $\|[\mathcal{X}_\zeta]'_{\mathbf{m}}(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)\|_{L^2(I \times \Omega; \mathbb{R}^d)}^2 \leq C \|1 + \theta_\varepsilon\|_{L^1(I \times \Omega)}$, which is bounded due to (3.58d). The penultimate term in (3.68) is a-priori bounded independently of ε for $\zeta > 0$ fixed because, as $\omega(\mathbf{F}, \mathbf{m}, \theta) = \mathcal{O}(\theta)$ due to (3.5d) and due to $\chi'_\zeta(\theta) = \mathcal{O}(1/\theta)$ uniformly for $\zeta > 0$, so that we have $[\chi'_\zeta(\cdot) \omega^2(\mathbf{F}, \mathbf{m}, \cdot)](\theta) = \mathcal{O}(\theta)$. Thus the estimate (3.58d) guarantees $\chi'_\zeta(\theta_\varepsilon) \omega^2(\mathbf{F}_\varepsilon, \mathbf{m}_\varepsilon, \theta_\varepsilon)$ bounded in $L^\infty(I; L^1(\Omega))$ while $|\mathbf{v}_\varepsilon|^2$ is surely bounded in $L^1(I; L^\infty(\Omega))$, cf. (3.58e).

In view of (3.69), one can summarize (3.68) as $I_\zeta^{(2)}(\theta_\varepsilon) \leq C(1 + \|\nabla \theta_\varepsilon\|_{L^\mu(\Omega; \mathbb{R}^d)})^{\mu^{*}/(r-1)}$. Combining it with (3.63), one obtain the inequality as

$$\|\nabla \theta_\varepsilon\|_{L^\mu(I \times \Omega; \mathbb{R}^d)} \leq C(1 + \|\nabla \theta_\varepsilon\|_{L^\mu(I \times \Omega; \mathbb{R}^d)})^{1-\mu/2+\mu^{*}/(2r-2)}. \tag{3.70}$$

Reminding $\sigma := (2-\mu)/(1+\zeta)$ from (3.62) with $\zeta > 0$ arbitrarily small, one gets after some algebra, the condition $\mu < (d+2)/(d+1)$. Obviously, for r big enough (in particular if $r > d$ as assumed), the exponent in the left-hand side of (3.70) is higher than the exponent in the right-hand side, which gives a bound for $\nabla\theta_\varepsilon$ in $L^\mu(I \times \Omega; \mathbb{R}^d)$. Altogether, we proved

$$\|\theta_\varepsilon\|_{L^\infty(I; L^1(\Omega)) \cap L^\mu(I; W^{1,\mu}(\Omega))} \leq C_\mu \quad \text{with } 1 \leq \mu < \frac{d+2}{d+1}. \quad (3.71a)$$

Next, we again exploit the calculus (3.41) now omitting the index k , with $\nabla \mathbf{F}_\varepsilon$ bounded in $L^\infty(I; L^r(\Omega; \mathbb{R}^{d \times d \times d}))$ and $\nabla \mathbf{m}_\varepsilon$ bounded in $L^\infty(I; L^2(\Omega; \mathbb{R}^{d \times d}))$ and relying on the assumption (3.5d), we have also the bound on ∇w_ε in $L^\mu(I; L^{\mu^*d/(\mu^*+d)}(\Omega; \mathbb{R}^d))$, so that

$$\|w_\varepsilon\|_{L^\infty(I; L^1(\Omega)) \cap L^\mu(I; W^{1,\mu^*d/(\mu^*+d)}(\Omega))} \leq C_\mu. \quad (3.71b)$$

Step 8: Limit passage for $\varepsilon \rightarrow 0$. We use the Banach selection principle as in Step 4, now also taking (3.58) and (3.71) into account instead of the estimates (3.31) and (3.40). For some subsequence and some $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, \theta)$, we now have

$$\begin{aligned} \varrho_\varepsilon \rightarrow \varrho & \quad \text{weakly}^* \text{ in } L^\infty(I; W^{1,r}(\Omega)) \cap W^{1,p}(I; L^r(\Omega)) \\ & \quad \text{and strongly in } C(I \times \overline{\Omega}), \end{aligned} \quad (3.72a)$$

$$\mathbf{v}_\varepsilon \rightarrow \mathbf{v} \quad \text{weakly}^* \text{ in } L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^2(I; W^{2,p}(\Omega; \mathbb{R}^d)), \quad (3.72b)$$

$$\begin{aligned} \mathbf{F}_\varepsilon \rightarrow \mathbf{F} & \quad \text{weakly}^* \text{ in } L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d})) \cap W^{1,p}(I; L^r(\Omega; \mathbb{R}^{d \times d})) \\ & \quad \text{and strongly in } C(I \times \overline{\Omega}; \mathbb{R}^{d \times d}), \end{aligned} \quad (3.72c)$$

$$\mathbf{m}_\varepsilon \rightarrow \mathbf{m} \quad \text{weakly}^* \text{ in } L^\infty(I; H^1(\Omega; \mathbb{R}^d)) \cap H^1(I; L^2(\Omega; \mathbb{R}^d)), \quad (3.72d)$$

$$\theta_\varepsilon \rightarrow \theta \quad \text{weakly}^* \text{ in } L^\mu(I; W^{1,\mu}(\Omega)), \quad 1 \leq \mu < (d+2)/(d+1). \quad (3.72e)$$

Like (3.45a), by the Aubin-Lions theorem, we now have

$$w_\varepsilon \rightarrow w \quad \text{strongly in } L^c(I \times \Omega), \quad 1 \leq c < 1+2/d, \quad (3.72f)$$

and then, using again continuity of $(\mathbf{F}, \mathbf{m}, w) \mapsto [\omega(\mathbf{F}, \mathbf{m}, \cdot)]^{-1}(w)$ as in (3.45b), we have

$$\theta_\varepsilon \rightarrow \theta = [\omega(\mathbf{F}, \mathbf{m}, \cdot)]^{-1}(w) \quad \text{strongly in } L^c(I \times \Omega), \quad 1 \leq c < 1+2/d. \quad (3.72g)$$

By the continuity of $\varphi'_\mathbf{F}$, $\zeta'_\mathbf{F}$, \det , and \mathcal{K} , we have also

$$\mathcal{K}(\mathbf{F}_\varepsilon, \theta_\varepsilon) \rightarrow \mathcal{K}(\mathbf{F}, \theta) \quad \text{strongly in } L^c(I \times \Omega) \text{ for any } 1 \leq c < \infty, \text{ and} \quad (3.72h)$$

$$\mathbf{T}_{\lambda,\varepsilon} \rightarrow \mathbf{T}_\lambda = \frac{[\pi_\lambda \varphi]'_\mathbf{F}(\mathbf{F}, \mathbf{m}) + \pi_\lambda(\mathbf{F}) \zeta'_\mathbf{F}(\mathbf{F}, \mathbf{m}, \theta)}{\det \mathbf{F}} \mathbf{F}^\top \quad \text{strongly in } L^1(I \times \Omega; \mathbb{R}_{\text{sym}}^{d \times d}). \quad (3.72i)$$

The momentum equation (3.25b) (still regularized by ε) is to be treated like in Step 4. Here we exploit the information about $\frac{\partial}{\partial t}(\varrho_\varepsilon \mathbf{v}_\varepsilon)$ in $L^{q'}(I; W^{1,q}(\Omega; \mathbb{R}^d)^*) + L^{p'}(I; W^{2,p}(\Omega; \mathbb{R}^d)^*)$

obtained like in (3.46); here we used also (3.58e). By the Aubin-Lions compact-embedding theorem, we then obtain

$$\varrho_\varepsilon \mathbf{v}_\varepsilon \rightarrow \varrho \mathbf{v} \quad \text{strongly in } L^s(I \times \Omega; \mathbb{R}^d) \quad \text{with } s \text{ from (3.58e)}. \quad (3.73)$$

In fact, the argumentation (3.52) now with C_r instead of $C_{r,\varepsilon}$ is to be slightly modified by using $(\varrho_\varepsilon, \mathbf{v}_\varepsilon, \mathbf{T}_{\lambda,\varepsilon}(\mathbf{F}_\varepsilon, \mathbf{v}_\varepsilon), \theta_\varepsilon)$ in place of $(\varrho_{\varepsilon k}, \mathbf{v}_{\varepsilon k}, \mathbf{T}_{\lambda,\varepsilon}(\mathbf{F}_{\varepsilon k}, \mathbf{v}_{\varepsilon k}), \theta_{\varepsilon k})$ and with $\tilde{\mathbf{v}}_k$ replaced by \mathbf{v} . Also, $\int_0^T \int_\Omega \frac{\partial}{\partial t}(\varrho_{\varepsilon k} \mathbf{v}_{\varepsilon k}) \cdot \tilde{\mathbf{v}}_k \, d\mathbf{x} \, dt$ is to be replaced by the duality $\langle \frac{\partial}{\partial t}(\varrho_\varepsilon \mathbf{v}_\varepsilon), \mathbf{v} \rangle$ with $\langle \cdot, \cdot \rangle$ denoting here the duality between $L^{q'}(I; W^{1,q}(\Omega; \mathbb{R}^d)^*) + L^{p'}(I; W^{2,p}(\Omega; \mathbb{R}^d)^*)$ and $L^q(I; W^{1,q}(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))$.

Limit passage in the heat equation (3.25f) is then simple. Altogether, we proved that $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, \theta)$ solves in the weak sense the problem (3.25)–(3.26) with $\varepsilon = 0$ and with \mathbf{T}_λ from (3.72i) in place of $\mathbf{T}_{\lambda,\varepsilon}$.

Step 9: the original problem. Let us note that the limit \mathbf{F} lives in $L^\infty(I; W^{1,r}(\Omega; \mathbb{R}^{d \times d})) \cap W^{1,p}(I; L^\infty(\Omega; \mathbb{R}^{d \times d}))$, cf. (3.58b,f), and this space is embedded into $C(I \times \bar{\Omega}; \mathbb{R}^{d \times d})$ if $r > d$. Therefore \mathbf{F} and its determinant evolve continuously in time, being valued respectively in $C(\bar{\Omega}; \mathbb{R}^{d \times d})$ and $C(\bar{\Omega})$. Let us recall that the initial condition \mathbf{F}_0 complies with the bounds (3.22) and we used this \mathbf{F}_0 also for the λ -regularized system. Therefore \mathbf{F} satisfies these bounds not only at $t = 0$ but also at least for small times. Yet, in view of the choice (3.22) of λ , this means that the λ -regularization is nonactive and $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, \theta)$ solves, at least for a small time, the original nonregularized problem (3.25)–(3.26) for which the a priori L^∞ -bounds (3.19) hold. By the continuation argument, we may see that the λ -regularization remains therefore inactive within the whole evolution of $(\varrho, \mathbf{v}, \mathbf{F}, \mathbf{m}, \theta)$ on the whole time interval I .

Step 10: energy balances. It is now important that the tests and then all the subsequent calculations leading to the energy balances (2.49) and (2.50) integrated over a current time interval $[0, t]$ are really legitimate.

In the calculus (2.34), we rely on that $[\varphi(\mathbf{F}, \mathbf{m})/\det \mathbf{F}]'_\mathbf{F} \in L^\infty(I \times \Omega; \mathbb{R}^{d \times d})$ is surely in duality with $\frac{\partial}{\partial t} \mathbf{F} \in L^p(I; L^r(\Omega; \mathbb{R}^{d \times d}))$ and $(\mathbf{v} \cdot \nabla) \mathbf{F} \in L^s(I; L^r(\Omega; \mathbb{R}^{d \times d}))$ with s from (3.58e). Moreover, $\frac{\partial}{\partial t}(\varrho \mathbf{v}) \in L^1(I; L^2(\Omega; \mathbb{R}^d)^*) + L^{p'}(I; W^{2,p}(\Omega; \mathbb{R}^d)^*)$ is in duality with $\mathbf{v} \in L^\infty(I; L^2(\Omega; \mathbb{R}^d)) \cap L^p(I; W^{2,p}(\Omega; \mathbb{R}^d))$, as used in (2.38). Further, the calculus (2.38) relies on that $\frac{\partial}{\partial t} \varrho$ and $\operatorname{div}(\varrho \mathbf{v}) = \mathbf{v} \cdot \nabla \varrho + \varrho \operatorname{div} \mathbf{v}$ live in $L^s(I; L^{rs/(r+s)}(\Omega))$ and thus are surely in duality with $|\mathbf{v}|^2 \in L^{s/2}(I; L^\infty(\Omega))$ with $3 \leq s < p(pd+4p-2d)/(4p-2d)$, cf. (3.58e). Eventually, since $\nabla^2 \mathbf{v} \in L^p(I \times \Omega; \mathbb{R}^{d \times d \times d})$, we have $\operatorname{div}^2(\nu_2 |\nabla^2 \mathbf{v}|^{p-2} \nabla^2 \mathbf{v}) \in L^{p'}(I; W^{2,p}(\Omega; \mathbb{R}^d)^*)$ in duality with \mathbf{v} . Also $\operatorname{div}(\nu_1 |e(\mathbf{v})|^{p-2} \mathbf{e}(\mathbf{v})) \in L^{p'}(I; W^{1,p}(\Omega; \mathbb{R}^d)^*)$ is in duality with \mathbf{v} due to the growth condition (3.5e). Altogether, the calculations (2.34)–(2.38) are legitimate.

Recalling in particular $\operatorname{div}(\kappa(\mathbf{F}) \nabla \mathbf{m} / \det \mathbf{F}) \in L^2(I \times \Omega; \mathbb{R}^d)$, we can see that also the calculations (2.39)–(2.48) are legitimate. This ends the proof of Theorem 3.2.

Remark 3.5 (*The classical solutions*). In fact, having $\nabla \mathbf{m}$ estimated from the exchange energy, (3.25e) holds even in the sense of $L^2(I \times \Omega; \mathbb{R}^d)$, not only in the weak sense, similarly as (3.27a,b). Actually, since $\operatorname{div}(\kappa(\mathbf{F}) \nabla \mathbf{m} / \det \mathbf{F}) \in L^2(I \times \Omega; \mathbb{R}^d)$, the inequality (3.4b) can be formulated as a classical inclusion (2.30d) and (3.25d) a.e. in the sense of $L^2(I \times \Omega; \mathbb{R}^d)$.

Remark 3.6 (*Importance of the exchange energy*). At large magnets, in contrast to micromagnetism, the influence of the exchange energy is small and, for very large magnetic continua, eventually negligible, cf. [11]. Yet, this energy controls $\nabla \mathbf{m}$, which ensures for “compactness” and strong convergence in \mathbf{m} and its complete deletion would be analytically problematic in particular because nonconvexity of $\psi(\mathbf{F}, \cdot, \theta)$ in ferromagnetic phase .

Acknowledgments. The author is deeply thankful to Giuseppe Tomassetti for many inspiring discussions and comments to the manuscript. A support from the Ministry of Education of the Czech Republic project CZ.02.1.01/0.0/0.0/15-003/0000493 and the CSF/DFG project GA22-00863K, and from the institutional support RVO:61388998 (ČR) is acknowledged.

References

- [1] J.M. Ball. Singular minimizers and their significance in elasticity. In M.G. Crandall, P.H. Rabinowitz, and R.E.L. Turner, editors, *Directions in Partial Differential Equations*, pages 1–15. Acad. Press, 1987.
- [2] J.M. Ball. Some open problems in elasticity. In P. Newton, P. Holmes, and A. Weinstein, editors, *Geometry, Mechanics, and Dynamics*, pages 3–59. Springer, New York, 2002.
- [3] J.M. Ball. Progress and puzzles in nonlinear elasticity. In J. Schröder and P. Neff, editors, *Poly-, Quasi- and Rank-One Convexity in Applied Mechanics*, CISM Intl. Centre for Mech. Sci. 516, pages 1–15. Springer, Wien, 2010.
- [4] W. Baltensperger and J.S. Helman. Dry friction in micromagnetics. *IEEE Trans. Magnet.*, 27:4772–4774, 1991.
- [5] B. Benešová, J. Forster, C. Liu, and A. Schlömerkemper. Existence of weak solutions to an evolutionary model for magnetoelasticity. *SIAM J. Math. Anal.*, 50:1200–1236, 2018.
- [6] G. Bertotti. *Hysteresis in Magnetism*. Academic Press, San Diego, 1998.
- [7] M. Bertsch, P. Podio Guidugli, and V. Valente. On the dynamics of deformable ferromagnets. *Annali di Mat. Pura ed Applicata*, 179:331–360, 2001.
- [8] L. Boccardo, A. Dall’aglio, T. Gallouët, and L. Orsina. Nonlinear parabolic equations with measure data. *J. Funct. Anal.*, 147:237–258, 1997.
- [9] L. Boccardo and T. Gallouët. Non-linear elliptic and parabolic equations involving measure data. *J. Funct. Anal.*, 87:149–169, 1989.
- [10] W.F. Brown Jr. *Magnetoelastic Interactions*. Springer, Berlin, 1966.
- [11] A. DeSimone. Energy minimizers for large ferromagnetic bodies. *Arch. Rational Mech. Anal.*, 125:99–143, 1993.
- [12] A. DeSimone and P. Podio Guidugli. Inertial and self interactions in structured continua: Liquid crystals and magnetostrictive solids. *Meccanica*, 300:629–640, 1995.
- [13] A. DeSimone and P. Podio Guidugli. On the continuum theory of deformable ferromagnetic solids. *Arch. Rational Mech. Anal.*, 136:201–233, 1996.
- [14] L. Dorfmann and R.W. Ogden. *Nonlinear Theory of Electroelastic and Magnetoelastic Interactions*. Springer, New York, 2014.
- [15] E. Du Trémolet de Lacheisserie, D. Gignoux, and M. Schlenker. *Magnetism: Fundamentals*. Springer, Berlin, 2005.
- [16] B. El Houari et al. Para-ferrimagnetic transition in strong coupling paramagnetic systems: Landau theory approach. *J. Magnetism & Magnetic Mater.*, 166:97–110, 1997.

- [17] J.L. Ericksen. Electromagnetic effects in thermoelastic materials. *Math. Mech. Solids*, 7:165–189, 2002.
- [18] E. Feireisl and J. Málek. On the navier-stokes equations with temperature-dependent transport coefficients. *Diff. Equations Nonlin. Mech.*, 14pp.(electronic), Art.ID 90616, 2006.
- [19] M. Foss, W. J. Hrusa, and V. J. Mizel. The Lavrentiev gap phenomenon in nonlinear elasticity. *Archive Rat. Mech. Anal.*, 167:337–365, 2003.
- [20] E. Fried and M.E. Gurtin. Traction, balances, and boundary conditions for nonsimple materials with application to liquid flow at small-length scales. *Arch. Ration. Mech. Anal.*, 182:513–554, 2006.
- [21] T.L. Gilbert. A Lagrangian formulation of the gyromagnetic equation of the magnetization field. *Phys. Rev.*, 100:1243, 1955.
- [22] G. Grün and P. Weiß. On the field-induced transport of magnetic nanoparticles in incompressible flow: Existence of global solutions. *J. Math. Fluid Mech.*, 23:Art.no.10, 2021.
- [23] M.E. Gurtin, E. Fried, and L. Anand. *The Mechanics and Thermodynamics of Continua*. Cambridge Univ. Press, New York, 2010.
- [24] X. Hu and N. Masmoudi. Global solutions to repulsive Hookean elastodynamics. *Arch. Ration. Mech. Anal.*, 223, 2016.
- [25] M. Kalousek, J. Kortum, and A. Schlömerkemper. Mathematical analysis of weak and strong solutions to an evolutionary model for magnetoviscoelasticity. *Disc. Cont. Dynam. Systems - S*, 14:17–39, 2021.
- [26] M. Kružík and T. Roubíček. *Mathematical Methods in Continuum Mechanics of Solids*. Springer, Cham/Switzerland, 2019.
- [27] L.D. Landau. On the theory of phase transitions. *Zh. Eksp. Teor. Fiz.*, 7:19–32, 1937.
- [28] L.D. Landau and E.M. Lifshitz. On the theory of dispersion of magnetic permeability in ferromagnetic bodies. *Phys. Z. Sowjet.*, 8 (153):153–169, 1935.
- [29] L.D. Landau and E.M. Lifshitz. *Electrodynamics of Continuous Media*. Pergamon Press, Oxford, 1984.
- [30] Z. Lei, C. Liu, and P. Zhou. Global solutions for incompressible viscoelastic fluids. *Archive Rational Mech. Anal.*, 188:371–398, 2008.
- [31] C. Liu and N.J. Walkington. An Eulerian description of fluids containing visco-elastic particles. *Archive Rational Mech. Anal.*, 159:229–252, 2001.
- [32] Z. Martinec. *Principles of Continuum Mechanics*. Birkhäuser/Springer, Switzerland, 2019.
- [33] G.A. Maugin. A continuum theory of deformable ferrimagnetic bodies. I. Field equations. *J. Math. Phys.*, 17:1727–1738, 1976.
- [34] G.A. Maugin. A note on micromagnetics at high temperature. *Appl. Phys.*, 11:185–186, 1976.
- [35] G.A. Maugin. *Continuum Mechanics of Electromagnetic Solids*. North-Holland, Amsterdam, 1988.
- [36] R.D. Mindlin. Micro-structure in linear elasticity. *Archive Ration. Mech. Anal.*, 16:51–78, 1964.
- [37] J. Nečas. Theory of multipolar fluids. In L. Jentsch and F. Tröltzsch, editors, *Problems and Methods in Mathematical Physics*, pages 111–119, Wiesbaden, 1994. Vieweg+Teubner.
- [38] J. Nečas, A. Novotný, and M. Šilhavý. Global solution to the compressible isothermal multipolar fluid. *J. Math. Anal. Appl.*, 162:223–241, 1991.
- [39] J. Nečas and M. Růžička. Global solution to the incompressible viscous-multipolar material problem. *J. Elasticity*, 29:175–202, 1992.

- [40] R.H. Nochetto, A.J. Salgado, and I. Tomas. The equations of ferrohydrodynamics: Modeling and numerical methods. *Math. Models Meth. Appl. Sci.*, 26:2393–2449, 2016.
- [41] P. Podio-Guidugli. On dissipation mechanisms in micromagnetics. *Eur. Phys J. B.*, 19:417–424, 2001.
- [42] P. Podio-Guidugli, T. Roubíček, and G. Tomassetti. A thermodynamically-consistent theory of the ferro/paramagnetic transition. *Archive Rat. Mech. Anal.*, 198:1057–1094, 2010.
- [43] J. Qian and Z. Zhang. Global well-posedness for compressible viscoelastic fluids near equilibrium. *Archive Rational Mech. Anal.*, 198:835–868, 2010.
- [44] M. Růžička. Mathematical and physical theory of multipolar viscoelasticity. *Bonner Mathematische Schriften* 233, Bonn, 1992.
- [45] R.E. Rosensweig. Magnetic fluids. *Ann. Rev. Fluid Mech.*, 19:437–463, 1987.
- [46] T. Roubíček. *Nonlinear Partial Differential Equations with Applications*. Birkhäuser, Basel, 2nd edition, 2013.
- [47] T. Roubíček. Visco-elastodynamics at large strains Eulerian. *Zeitschrift f. angew. Math. Phys.* (to appear), 2021. (Preprint arXiv no. 2102.00261.).
- [48] T. Roubíček. A thermodynamical model for paleomagnetism in Earth’s crust. *Math. Mech. Solids*, pages 1–28, 2022. printed online: DOI: 10.1177/10812865221093281.
- [49] T. Roubíček. Thermodynamics of viscoelastic solids, its Eulerian formulation, and existence of weak solutions. (Preprint arXiv no.2203.06080), 2022.
- [50] T. Roubíček and U. Stefanelli. Visco-elastodynamics of solids undergoing swelling at large strains by an Eulerian approach. (Preprint 2022 arXiv no.2201.10959). *SIAM J. Math. Anal.* (to appear).
- [51] T. Roubíček and G. Tomassetti. A thermodynamically consistent model of magneto-elastic materials under diffusion at large strains and its analysis. *Zeit. angew. Math. Phys.*, 69:Art.no.55, 2018.
- [52] T. Roubíček, G. Tomassetti, and C. Zanini. The Gilbert equation with dry-friction-type damping. *J. Math. Anal. Appl.*, 355:453–468, 2009.
- [53] A Schlömerkemper and J Žabenský. Uniqueness of solutions for a mathematical model for magneto-viscoelastic flows. *Nonlinearity*, 31:2989–3012, 2018.
- [54] S. Scrobogna. On the global well-posedness of a class of 2D solutions for the Rosensweig system of ferrofluids. *J. Diff. Equations*, 266:2718–2761, 2019.
- [55] M. Šilhavý. Multipolar viscoelastic materials and the symmetry of the coefficient of viscosity. *Appl. Math.*, 37:383–400, 1992.
- [56] R.A. Toupin. Elastic materials with couple stresses. *Arch. Ration. Mech. Anal.*, 11:385–414, 1962.
- [57] R.A. Toupin. Theories of elasticity with couple stresses. *Arch. Ration. Mech. Anal.*, 17:85–112, 1964.
- [58] P. Vágner, M. Pavelka, and O. Esen. Multiscale thermodynamics of charged mixtures. *Continuum Mech. Thermodyn.*, 33:237–268, 2021.
- [59] A. Visintin. Modified Landau-Lifshitz equation for ferromagnetism. *Phys. B*, 233:365–369, 1997.
- [60] W. Zhao. Local well-posedness and blow-up criteria of magneto-viscoelastic flow. *Disc. Cont. Dynam. Syst.*, 38:4637–4655, 2018.