

The Finslerian Wormhole model with $f(R, T)$ gravity

Manjunath Malligawad,¹ S. K. Narasimhamurthy,¹ Z. Nekouee,^{1,2} and Mallikarjun Y. Kumbar³

¹*Department of PG Studies and Research in Mathematics, Kuvempu University
Jnana Sahyadri, Shankaraghatta - 577 451, Shivamogga, Karnataka, India*

²*Department of Mathematics, Faculty of Mathematical Sciences,
University of Mazandaran, P. O. Box 47416-95447, Babolsar, Iran*

³*Department of Mathematics, Mahantswamy Arts, Science and Commerce College,
Haunsbhavi - 581 109, Haveri, Karnataka, India*

(*Electronic mail: nmurthysk@gmail.com)

(Dated: 7 February 2023)

In this article, based on Finsler geometry, we study the wormhole model in $f(R, T) = R + 2\lambda T$ gravity theory with an exponential shape function. On the basis of this wormhole solution, we derive the gravitational field equations. Using the exponential shape function, we discuss the character of parameters such as density, radial pressure, transverse pressure, equation-of-state parameters, and energy conditions in $f(R, T)$ gravity. We study the significant role of parameter λ in the violation of energy conditions and also in the equilibrium state of the Finslerian wormhole solution, which is caused by the anisotropic force and hydrostatic force in $f(R, T)$ gravity. Further, we observe in the framework of Finslerian space-time that the $f(R, T)$ gravity model successfully captures the features of wormholes. Further using these features we plotted and visualized the 3-D wormhole structure Fig. 2.

I. INTRODUCTION

The theme of the wormhole was first discussed by German mathematician H. Weyl¹. He explained the structure of a wormhole is an asymptotically flat tube-like structure, and it is a tunnel with two entrances or two distinct points of different space-times or bridge joining two ends of the same space-time. Therefore, the issue of wormholes attracted the attention of many geometers and physicists. Schwarzschild wormhole is the first type of wormhole solution developed by Schwarzschild.

The wormhole is the most well-known and extensively researched idea in General Relativity (GR) and modified theories of gravity. The GR solutions produce wormholes. These hypothetical shortcuts, generally known as Einstein-Rosen bridges, were mathematically proven to exist for the first time by². But³ showed in a paper later published in 1962 this type of wormhole is unstable. If it joins two regions of the same universe and any object moving slower than the speed of light travels from one exterior area to the other if it falls from it. There are two kinds of wormholes static and dynamic wormholes.

The ideology of the traversable wormhole was proposed by⁴ with the intention of using a wormhole for time travel or interstellar travel. It is worth noting that the idea of the traversable wormhole is distinct from the Einstein-Rosen bridge theory. To keep the wormhole geometry, and key constituent is the existence of exotic matter in its throat, which pushes wormhole walls apart and prevents them from collapsing. The throat from contracting, allowing it to be traversed, and the rigidity of the traversable wormhole has been examined in order to reduce violation of the null energy condition. The most frequently researched issue in the wormhole concept is the violation of energy conditions.⁵ proved

that all traversable wormholes, both static and dynamic, violated the null energy conditions.

The energy conditions connected are as $DEC \subset WEC \subset NEC$ and $SEC \subset NEC$. Violating the NEC involves violating all of the energy conditions. In reality, even under conditions weaker than the NEC⁶, traversable wormholes cannot be realized in GR, according to the following topological censorship theorem in $f(R)$ gravity⁷, traversable wormholes were examined by⁸. They supported the wormhole constructions and identified the causes of the null energy condition's dissatisfaction. $f(R)$ and $f(R, T)$ theories have received a lot of attention recently as a way to explain several elements of the universe. By adding the stress-energy tensor term,⁹ generalized the $f(R)$ gravitational theory and developed the $f(R, T)$ MGT. Gravity models depend on the source term since matter and gravity are connected. Using $f(R, T)$ gravity,¹⁰ investigated the cosmic evolution of deceleration and the equation of state (EoS) parameters. In the context of the FLRW model, in the study of static wormholes in the $f(R, T)$ theory of gravity,¹¹ provided some wormhole models with various matter content assumptions.¹² investigated traversable wormholes with one extra space-like compact dimension. In this respect, they considered how it affected energy density, scale factor, and shape function.

In $f(R, T)$ modified gravity theory (MGT)¹³, describe the geometry of the wormhole by using an exponential shape function. The exponential $f(R, T)$ MGT solution is provided by¹⁴, and the concept of exponential $f(R, T)$ MGT is considered original in literature.

Finsler geometry has caught the attention of many physicists in currently because it explain a variety of issues that Einstein's gravity is unable to explain. Riemannian geometry is one of the special cases of Finsler geometry. Modern standard high-energy theories and Finsler-like gravity theories are

consistent with experimental results. Without considering the dark matter hypothesis, Finsler geometry provides a better method for addressing problems with experimental results of flat rotation curves and spiral galaxies¹⁵.

In Finsler geometry traversable wormhole was created by¹⁶. They discovered an exact solution for various types of shape functions, red-shift functions, and EoS. They also talked about the features of wormhole models. By studying whether or if wormhole stable solutions exist without the need for exotic matter, wormhole geometry has also been investigated thoroughly using modified theories, where MGTs allow us to discuss the issue of exotic matter¹⁷.

Within the context of Finsler geometry,¹⁸ investigate the wormhole model in $f(R)$ gravity with an exponential shape function. They also examined the gravitational field equation for finding wormhole solutions in the Finslerian framework by taking into account the anisotropic energy-momentum tensor. Our inspiration for it comes from the exponential $f(R)$ gravity model. By resolving the wormhole field equations using the Finslerian $f(R, T)$ MGT formalism, then we apply this form for the $f(R, T)$ MGT. Specifically we focus on the family of $f(R, T)$ MGT, i.e., $f(R, T) = R + f(T)$, where $f(T) = 2\lambda T$, R represent the Ricci scalar, T represent energy momentum tensor, and λ is a parameter. In this article, we explore the Finslerian wormhole model under $f(R, T)$ MGT with exponential shape function and examine the wormhole solutions based on parameter values showed that the violation of energy conditions.

The following planning guides the structure of the paper. In section (II), we quickly review the ideology of Finsler geometry. In section (III), we described the Finsler geometry formalization in $f(R, T)$ MGT and discussed the energy conditions. In section (IV), we discussed and analyzed the obtained outcomes. The study concludes with section (V), and we provide some concluding observations.

II. PRELIMINARIES AND NOTATIONS OF FINSLER GEOMETRY

Finsler structure on manifold M is defined as function $F : TM \rightarrow [0, \infty)$ which satisfies the below properties:

1. Regularity: F is smooth function on the $TM \setminus \{0\}$.
2. Positive Homogeneity: $F(x, cy) = cF(x, y)$ for all $c > 0$.
3. Strong Convexity: The $n \times n$ Hessian matrix

$$g_{\mu\nu} = \frac{\partial^2(\frac{1}{2}F^2)}{\partial y^\mu \partial y^\nu} = \frac{1}{2} \dot{\partial}_\mu \dot{\partial}_\nu F^2, \quad (1)$$

Finsler structure F is positive definite on $TM \setminus \{0\}$, and it is the function of (x^i, y^i) . The pair (M, F) is called Finsler space.

A Finslerian metric is referred to as Riemannian if F^2 is quadratic in y . For the Finsler manifold, the geodesic equation

is as follows:

$$\frac{d^2 x^\mu}{d\tau^2} + 2G^\mu = 0, \quad (2)$$

where

$$G^\mu = \frac{1}{4} g^{\mu\nu} \left(\frac{\partial^2 F^2}{\partial x^\nu \partial y^\nu} y^\nu - \frac{\partial F^2}{\partial x^\nu} \right), \quad (3)$$

are called geodesic spray coefficients.

We observed that along the geodesic the Finslerian structure $F(x, y)$ is constant. The Finslerian modified Ricci tensor equation suggested by¹⁹ as follows.

$$Ric_{\mu\nu} = \frac{\partial^2(\frac{1}{2}F^2 Ric)}{\partial y^\mu \partial y^\nu}, \quad (4)$$

here Ric stands for Ricci scalar. Ric is a geometric invariant with the following expression

$$Ric = g^{\mu\nu} R_{\mu\nu}. \quad (5)$$

on any basis, Eq. (5) is true. Moreover, in Eq. (5), $R_{\mu\nu}$ is a representation of the flag's original curvature. It might be stated as,

$$R_\nu^\mu = \frac{1}{F^2} \left(2 \frac{\partial G^\mu}{\partial x^\nu} - y^\nu \frac{\partial^2 G^\mu}{\partial x^\nu \partial y^\nu} + 2G^\nu \frac{\partial^2 G^\mu}{\partial y^\nu \partial y^\nu} - \frac{\partial G^\mu}{\partial y^\nu} \frac{\partial G^\nu}{\partial y^\nu} \right). \quad (6)$$

Thus, Ric is as follows,

$$Ric \equiv R_\mu^\mu, \quad R_\mu^\mu = \frac{1}{F^2} \left(2 \frac{\partial G^\mu}{\partial x^\mu} - y^\nu \frac{\partial^2 G^\mu}{\partial x^\nu \partial y^\mu} + 2G^\nu \frac{\partial^2 G^\mu}{\partial y^\nu \partial y^\mu} - \frac{\partial G^\mu}{\partial y^\nu} \frac{\partial G^\nu}{\partial y^\mu} \right). \quad (7)$$

Finslerian modified formula for scalar curvature is

$$S = g^{\mu\nu} Ric_{\mu\nu}, \quad (8)$$

and Einstein tensor formula is

$$G_{\mu\nu} = Ric_{\mu\nu} - \frac{1}{2} g_{\mu\nu} S. \quad (9)$$

Since it is derived from Ric , connections do not affect the Finslerian modified Einstein tensor. It is solely reliant on the Finslerian structure. As a result, the Finslerian gravitational field equations are also insensitive to the connections.

We choose the⁴ wormhole metric to search for wormhole structure and consider the Finsler structure to be of the following form,

$$F^2 = e^{2a(r)} y^t y^t - \left(1 - \frac{b(r)}{r} \right)^{-1} y^r y^r - r^2 \bar{F}^2(\theta, \phi, y^\theta, y^\phi), \quad (10)$$

where \bar{F} is a Finsler structure in two dimensions, and we assume that \bar{F} has the form,

$$\bar{F}^2 = y^\theta y^\theta + A(\theta, \phi) y^\phi y^\phi. \quad (11)$$

Within the Finsler framework in Eq. (10) $a(r)$ stands for the redshift function, while $b(r)$ is the shape function²⁰. $a(r)$ explains the redshift effect and tidal force in the wormhole space-time. The shape function $b(r)$ specifies how a wormhole is shaped. The wormhole throat is positioned at a minimal surface radius $b(r_0) = r_0$, and the nonmonotonic radial coordinate r falls from infinity to r_0 before increasing from r_0 back to infinity. To prevent the appearance of an event horizon, everywhere $a(r)$ must be a finite. At the throat singularity does not exist for the traversable wormhole metric. $b(r)$ needs to meet the requirements⁴ below to produce the wormhole solutions:

- Radial co-ordinate r has a range of values from $r_0 \leq r \leq \infty$, where r_0 is the throat radius.
- Within the throat $b(r)$ fulfill the requirement $b(r_0) = r_0$, as well as for out of the throat, i.e., for $r > r_0, 0 < 1 - \frac{b(r)}{r}$.
- Shape function $b(r)$ at the throat must satisfy the flaring out requirement, i.e., $b'(r_0) < 1$, where the derivative is with respect to r .
- The minimum requirement for the space-time geometry to be asymptotically flat $\frac{b(r)}{r} \rightarrow 0$ as $|r| \rightarrow \infty$.

The geometry of the wormhole space-time is described by the requirements listed above.

For two-dimensional Finsler structure \bar{F} Eq. (11) one can obtain the Finslerian metric as,

$$\bar{g}_{ij} = \text{diag}(1, A(\theta, \phi)), \quad (12)$$

$$\bar{g}^{ij} = \text{diag}\left(1, \frac{1}{A(\theta, \phi)}\right), \quad (13)$$

where $(i, j = \theta, \phi)$. The \bar{Ric} of the Finslerian structure \bar{F} can be found by applying the formula from the reference¹⁶.

$$\bar{Ric} = \frac{1}{2A} \left[-\frac{\partial^2 A}{\partial \theta^2} + \frac{1}{2A} \left(\frac{\partial A}{\partial \theta} \right)^2 \right]. \quad (14)$$

For the Finsler structure \bar{F} , we can better understand wormholes in the Finslerian theory by using the $\bar{Ric} = \eta$. Here it is assumed to be constant. In three separate circumstances, we find the solution to the differential equation (14) by solving it ($\eta > 0, \eta = 0, \eta < 0$). Consequently, the Finsler structure \bar{F} Eq. (11) changes to,

$$\bar{F}^2 = y^\theta y^\theta + C \sin^2(\sqrt{\eta} \theta) y^\phi y^\phi \quad \text{for } (\eta > 0), \quad (15)$$

$$\bar{F}^2 = y^\theta y^\theta + C \theta^2 y^\phi y^\phi \quad \text{for } (\eta = 0), \quad (16)$$

$$\bar{F}^2 = y^\theta y^\theta + C \sinh^2(\sqrt{-\eta} \theta) y^\phi y^\phi \quad \text{for } (\eta < 0). \quad (17)$$

Now let's use $C = 1$. For the solution $\eta > 0$ the Finslerian wormhole structure Eq. (10) can be written as

$$F^2 = e^{2a(r)} y^t y^t - \left(1 - \frac{b(r)}{r}\right)^{-1} y^r y^r - r^2 \left(y^\theta y^\theta + \sin^2(\sqrt{\eta} \theta) y^\phi y^\phi \right). \quad (18)$$

The Riemannian metric is denoted by α , and the appropriate Riemannian wormhole structure is provided by

$$\alpha^2 = e^{2a(r)} y^t y^t - \left(1 - \frac{b(r)}{r}\right)^{-1} y^r y^r - r^2 \left(y^\theta y^\theta + \sin^2(\theta) y^\phi y^\phi \right). \quad (19)$$

Using Eq. (19), we can written Eq. (18) as,

$$F^2 = \alpha^2 + r^2 (\sin^2 \theta - \sin^2 \sqrt{\eta} \theta) y^\phi y^\phi. \quad (20)$$

If we take $r^2 (\sin^2 \theta - \sin^2 \sqrt{\eta} \theta) y^\phi y^\phi = \beta^2$ into consideration, then Eq. (20) yields,

$$F^2 = \alpha^2 (1 + s^2), \quad (21)$$

where

$$s = \frac{\beta}{\alpha} = \frac{b_\phi y^\phi}{\alpha}.$$

We consider $b_\phi = r \sqrt{\sin^2 \theta - \sin^2 \sqrt{\eta} \theta}$. Additionally, the $\beta = b_\mu y^\mu$ denotes the differential 1-form where $b_\mu = (0, 0, 0, b_\phi)$. Thus, Eq. (21) yields,

$$F = \alpha \varphi(s), \quad (22)$$

where $\varphi(s) = \sqrt{1 + s^2}$. The Finslerian wormhole structure F is indicated by Eq. (22), which represents the Finsler space with an (α, β) -metric.

Concerning the Riemannian metric α , the given symbol " $|$ " signifies the covariant derivative. The equations for $K_V(\alpha)$ and $K_V(\beta)$ are then provided by,

$$K_V(\alpha) = \frac{1}{2\alpha} (V_\mu|_\nu + V_\nu|_\mu) y^\mu y^\nu, \\ K_V(\beta) = \left(V^\mu \frac{\partial b_\nu}{\partial x^\mu} + b_\mu \frac{\partial V^\mu}{\partial x^\nu} \right) y^\nu.$$

The Finsler space's Killing equation, $K_V(F) = 0$, is derived from isometric transformations²¹.

$$\left(\varphi(s) - s \frac{\partial \varphi(s)}{\partial s} \right) K_V(\alpha) + \frac{\partial \varphi(s)}{\partial s} K_V(\beta) = 0. \quad (23)$$

From Eq. (22), the Killing equation (23) becomes,

$$\alpha K_V(\alpha) + \beta K_V(\beta) = 0,$$

and it pays off

$$K_V(\alpha) = 0 \quad \text{and} \quad K_V(\beta) = 0 \quad (24)$$

as a result we have

$$V_{\mu|v} + V_{v|\mu} = 0, \quad (25)$$

$$V^\mu \frac{\partial b_v}{\partial x^\mu} + b_\mu \frac{\partial V^\mu}{\partial x^\nu} = 0. \quad (26)$$

The Riemannian space-time and the Killing equation Eq. (25) are equivalents. Eq. (26) restricts Eq. (25). The Lie derivatives $L_V(\alpha) = 0$ and $L_V(\beta) = 0$, are comparable to the Killing equations Eq. (25), and Eq. (26) respectively. In addition, we have $\alpha^2 = h_{\mu\nu}$. We can get the Riemannian wormhole metric as,

$$h_{\mu\nu} = \text{diag} \left(e^{2a(r)}, - \left(1 - \frac{b(r)}{r} \right)^{-1}, -r^2, -r^2 \sin^2 \theta \right).$$

The Riemannian space-time isometric symmetry is broken as a result of the Killing equation (26).

The redshift function should be positive for every $r > r_0$ if there are no horizons or singularities. In our current study, the redshift function $a(r)$ is taken to be constant as a result, $a'(r) = 0$. One of the crucial properties of traversable wormholes is that the tidal gravitational forces that traveler experiences must be small or $a'(r) = 0$. The zero-tidal-force solutions have been examined by⁴, who also took into account the vanishing redshift function. As a result, the Finslerian wormhole structure Eq. (10) becomes,

$$F^2 = y^t y^t - \left(1 - \frac{b(r)}{r} \right)^{-1} y^r y^r - r^2 \left(y^\theta y^\theta + \sin^2(\sqrt{\eta}\theta) y^\phi y^\phi \right). \quad (27)$$

One can derive the Finsler metric as,

$$g_{\mu\nu} = \text{diag} \left(1, - \left(1 - \frac{b(r)}{r} \right)^{-1}, -r^2, -r^2 \sin^2(\sqrt{\eta}\theta) \right), \quad (28)$$

$$g^{\mu\nu} = \text{diag} \left(1, - \left(1 - \frac{b(r)}{r} \right), \frac{-1}{r^2}, \frac{-1}{r^2 \sin^2(\sqrt{\eta}\theta)} \right). \quad (29)$$

When looking at the Finslerian wormhole crucial role is played by the Ricci scalar $\bar{Ric} = \eta$ of the two-dimensional Finsler structure \bar{F} . We conclude that the structure F has a constant flag curvature because $\eta = \text{constant}$. The geodesic spray coefficients can be found by putting the Finslerian wormhole structure Eq. (27) in Eq. (2).

$$G^t = 0, \quad (30)$$

$$G^r = \frac{rb' - b}{4r(r-b)} y^r y^r - \frac{r-b}{2} y^\theta y^\theta - \frac{r-b}{2} \sin^2(\sqrt{\eta}\theta) y^\phi y^\phi, \quad (31)$$

$$G^\theta = \frac{1}{r} y^r y^\theta - \frac{\sqrt{\eta}}{2} \sin(\sqrt{\eta}\theta) \cos(\sqrt{\eta}\theta) y^\phi y^\phi, \quad (32)$$

$$G^\phi = \frac{1}{r} y^r y^\phi + \sqrt{\eta} \cot(\sqrt{\eta}\theta) y^\theta y^\phi. \quad (33)$$

By substituting the expressions from Eqs. (30-33) in Eq. (7), we obtain,

$$F^2 Ric = \frac{rb' - b}{r^2(r-b)} y^r y^r + \left(\eta - 1 + \frac{b}{2r} + \frac{b'}{2} \right) y^\theta y^\theta + \left(\eta - 1 + \frac{b}{2r} + \frac{b'}{2} \right) \sin^2(\sqrt{\eta}\theta) y^\phi y^\phi. \quad (34)$$

From Eq. (4), we deduced Finslerian modified Ricci tensor components as,

$$Ric_{tt} = 0, \quad (35)$$

$$Ric_{rr} = \frac{rb' - b}{r^2(r-b)}, \quad (36)$$

$$Ric_{\theta\theta} = \left(\eta - 1 + \frac{b}{2r} + \frac{b'}{2} \right), \quad (37)$$

$$Ric_{\phi\phi} = \left(\eta - 1 + \frac{b}{2r} + \frac{b'}{2} \right) \sin^2(\sqrt{\eta}\theta). \quad (38)$$

By substituting expressions Eq. (35-38) and Eq. (29) in Eq. (8), we can determine the scalar curvature for the Finslerian wormhole space-time as,

$$S = -\frac{2}{r^2} (b' + \eta - 1). \quad (39)$$

Using the components of the Finslerian-modified Ricci tensor and the scalar curvature, from Eq. (9) one can quickly determine the components of the Finslerian-modified Einstein tensor as,

$$G_t^t = \frac{1}{r^2} (b' + \eta - 1), \quad (40)$$

$$G_r^r = \frac{b}{r^3} + \frac{1}{r^2} (\eta - 1), \quad (41)$$

$$G_\theta^\theta = G_\phi^\phi = \frac{rb' - b}{r^3}. \quad (42)$$

From components of the Finslerian modified Einstein tensor Eq. (40-42) depend only on r and are direction-independent. With the help of the formula, the Chern connection is determined by,

$$\Gamma_{\nu\nu}^\alpha = \frac{\partial^2 G^\alpha}{\partial y^\nu \partial y^\nu} - A_{\nu\nu|\beta}^\alpha \frac{y^\beta}{F}. \quad (43)$$

The term $A_{\mu\nu\nu} = g_{\mu\alpha} A_{\nu\nu}^\alpha$ in the above equation, which represents the Cartan connection, is defined as,

$$A_{\mu\nu\nu} = \frac{F}{4} \frac{\partial^3 F^2}{\partial y^\mu \partial y^\nu \partial y^\nu}, \quad (44)$$

for all indices μ, ν and ν , we obtain $A_{\mu\nu\nu} = 0$, to the Finslerian wormhole structure Eq. (27). From this, we obtain that the Cartan connections for the Finslerian wormhole space-time are vanish. As a result, we can calculate the Chern connection coefficients using Eq. (43). So we have,

$$\begin{aligned} \Gamma_{\mu\nu}^\nu &= 0 \quad (\text{for all indices } \mu, \nu), \quad \Gamma_{r\theta}^\theta = \Gamma_{r\phi}^\phi = \frac{1}{r}, \\ \Gamma_{rr}^r &= \frac{rb' - b}{2r(r-b)}, \quad \Gamma_{rt}^r = \Gamma_{r\theta}^r = \Gamma_{r\phi}^r = 0, \\ \Gamma_{\theta\theta}^r &= -(r-b), \quad \Gamma_{\phi\phi}^r = -(r-b)\sin^2(\sqrt{\eta}\theta), \\ \Gamma_{\theta\theta}^\theta &= \Gamma_{\theta t}^\theta = \Gamma_{\theta\phi}^\theta = 0, \\ \Gamma_{\phi t}^\phi &= \Gamma_{\phi\phi}^\phi = 0, \quad \Gamma_{\phi\theta}^\phi = \sqrt{\eta}\cot(\sqrt{\eta}\theta). \end{aligned} \quad (45)$$

Now we define the derivative $\frac{\delta}{\delta x^\mu}$ as,

$$\frac{\delta}{\delta x^\mu} = \frac{\partial}{\partial x^\mu} - \frac{\partial G^\alpha}{\partial y^\mu} \frac{\partial}{\partial y^\alpha}.$$

We can determine the covariant derivative of G_V^μ by the following equation

$$G_{\nu|\mu}^\mu = \frac{\delta G_V^\mu}{\delta x^\mu} + \Gamma_{\mu\alpha}^\mu G_V^\alpha - \Gamma_{\mu\nu}^\alpha G_{\alpha}^\mu. \quad (46)$$

From Eq. (46) all the covariant derivative components becomes $G_{t|\mu}^\mu = G_{r|\mu}^\mu = G_{\theta|\mu}^\mu = G_{\phi|\mu}^\mu = 0$. This leads us to the conclusion that the covariant derivative of G_V^μ for the Finslerian modified Einstein tensor is conserved.

III. FORMALISM OF FINSLER GEOMETRY IN $f(R, T)$ GRAVITY

$f(R, T)$ MGT has received a lot of attention in recent research and seems to be a good fit for addressing wormhole construction problems (since it doesn't have any misleading characteristics). For understanding the observed universe, $f(R, T)$ MGT, this is the combination of two functions R and T , where R stands for the Ricci scalar and T for the trace of the energy-momentum tensor, is frequently studied in the research⁹. Since the distribution of the matter needed to create wormholes remains a difficult problem for physicists. So, we take the universal anisotropic energy-momentum tensor as²²,

$$T_V^\mu = (\rho + p_t)u^\mu u_\nu + (p_r - p_t)x^\mu x_\nu - p_t\delta_V^\mu. \quad (47)$$

When taken in the radial direction, energy density is represented by ρ , u^μ is the four-velocity such that $u^\mu u_\mu = 1$ and x^μ is the space-like unit vector, such that $x^\mu x_\mu = -1$. p_r is the radial pressure, which is determined by the direction of the space-like unit vector x^μ , where p_t represents the transverse pressure measures perpendicular to x^μ .

Now let's have a look at the parameter anisotropic factor $\Delta = p_t - p_r$, which measures the anisotropy where $p_t \neq p_r$. We have $T = \rho - p_r - 2p_t$, which displays the trace of the anisotropic energy-momentum tensor. We can determine the geometry of wormholes based on the anisotropic factor. If $\Delta < 0$ (negative) geometry of the wormhole is attractive, so anisotropic force is directed inward. If $\Delta > 0$ (positive) geometry of the wormhole is repulsive, then anisotropic force is directed outward. The matter distribution of the wormholes exhibits isotropic pressure if $\Delta = 0$.

The achievement of the $f(R)$ theory in the cosmology inspired us to investigate wormhole solutions because $f(R, T)$ MGT reduces to $f(R)$ gravity in the limit $\lambda \rightarrow 0$. In this section, we'll look at the functional form $f(R, T) = R + f(T)$, where $f(T) = 2\lambda T$ is a function of the trace of the stress-energy tensor of matter. As a result, it is crucial to research how λ affects wormhole solutions with shape functions when there is matter present. The equation for the Finslerian gravitational field is provided by,

$$G_V^\mu = (8\pi_F + 2\lambda)T_V^\mu + \lambda(2\rho + T)g_V^\mu, \quad (48)$$

where $4\pi_F$ stands for the volume of the Finsler structure \bar{F} in two dimensions. If we take \bar{F} as the Finslerian sphere², then π_F becomes equal to π . By combining components of the Finslerian modified Einstein tensor Eq. (40-42) and energy-momentum tensor Eq. (47), by taking ($G = C = 1$) as relativistic units we can arrive at the gravitational field equations from Eq. (48) for Finslerian wormhole space-time Eq. (27) as follows,

$$G_t^t = (8\pi_F + 2\lambda)T_t^t + \lambda(2\rho + T)g_t^t,$$

$$\frac{1}{r^2}(b' + \eta - 1) = (8\pi_F + 2\lambda)\rho + \lambda(p_r + 2p_t). \quad (49)$$

$$G_r^r = (8\pi_F + 2\lambda)T_r^r + \lambda(2\rho + T)g_r^r,$$

$$-\frac{b}{r^3} - \frac{1}{r^2}(\eta - 1) = (8\pi_F + 3\lambda)p_r + \lambda(\rho + 2p_t). \quad (50)$$

$$G_\mu^\mu = (8\pi_F + 2\lambda)T_\mu^\mu + \lambda(2\rho + T)g_\mu^\mu \quad (\mu = \theta, \phi),$$

$$\frac{b - b'r}{2r^3} = (8\pi_F + 4\lambda)p_t + \lambda(\rho + 2p_r). \quad (51)$$

If we know the distribution of the matter, gravitational field equations enable us to identify the space-time geometry and vice-versa. Additionally, it has the interesting property that, using the field equations, one can analyze both the distribution of matter and the whole structure of space-time if they are aware of some of the energy stress tensor's components and some of the space-time geometry.

In the context of $f(R, T) = R + 2\lambda T$ gravity, an exponential shape function¹³

$$b(r) = r_0 e^{1 - \frac{r}{r_0}}, \quad (52)$$

is taken into consideration in the Finslerian wormhole model to investigate the energy conditions. The shape function Eq. (52) meets every requirement for supporting wormholes mentioned above (Sec. II), i.e., fulfills the requirements $r > r_0$,

$b(r_0) = r_0, 1 - \frac{b(r)}{r} > 0$ where r_0 is the wormhole's throat radius, the flaring-out condition, and the asymptotic flatness condition. Fig. 1(a) depicts the properties of the exponential shape function Eq. (52) with $r_0 = 0.5$ and demonstrates that the shape function meets all important requirements. We constructed an embedded 2-D diagram in Fig. 1(b) and a 3-D diagram in Fig. 2 for the shape function Eq. (52) for better visualization of the wormhole see Appendix.




FIG. 1. (a) Behavior of the exponential shape function Eq. (52) at $r_0 = 0.5$. (b) Embedded 2-D plot of the wormhole defined by Eq. (52).




FIG. 2. Embedded 3-D plot of the wormhole defined by Eq. (52).

By inserting Eq. (52) into Eqs. (49-51), energy density ρ , radial pressure p_r , and transverse pressure p_t are obtained respectively as follows,

$$\rho = \frac{-e^{1-\frac{r}{r_0}} + \eta - 1}{r^2(8\pi_F + 2\lambda)}, \quad (53)$$

$$p_r = -\frac{r_0 e^{1-\frac{r}{r_0}} + r(\eta - 1)}{r^3(8\pi_F + 2\lambda)}, \quad (54)$$

$$p_t = \frac{e^{1-\frac{r}{r_0}}(r + r_0)}{2r^3(8\pi_F + 2\lambda)}. \quad (55)$$

Fig. 3, Fig. 4, and Fig. 5 depict the plot of p_r , p_t , ρ respectively as function of r and η for the Finslerian wormhole solution (27) at different value of λ , i.e., $8\pi_F \neq -2\lambda$ ($\lambda \neq -12.568$) so we are taking $\lambda \geq -12.5$ (considering $\lambda = -12.5$) and $\lambda \leq -12.6$ (considering $\lambda = -12.6$) with wormhole throat placed at $r_0 = 0.5$.




FIG. 3. (a) Radial pressure $p_r = p_r(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$. (b) Radial pressure $p_r = p_r(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$.




FIG. 4. (a) Transverse pressure $p_t = p_t(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$. (b) Transverse pressure $p_t = p_t(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$.




FIG. 5. (a) Energy density $\rho = \rho(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (WEC term). (b) Energy density $\rho = \rho(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (WEC term).

In addition, by combining the Eqs. (53-55), we obtain the

following stress-energy components,

$$\rho + p_r = -\frac{e^{1-\frac{r}{r_0}}(r+r_0)}{r^3(8\pi_F + 2\lambda)}, \quad (56)$$

$$\rho + p_t = \frac{e^{1-\frac{r}{r_0}}(r_0-r) + 2r(\eta-1)}{2r^3(8\pi_F + 2\lambda)}, \quad (57)$$

$$\rho + p_r + 2p_t = 0, \quad (58)$$

$$\rho - |p_r| = \frac{-e^{1-\frac{r}{r_0}} + \eta - 1 - r^2 \left| \frac{r_0+r}{r^3} \right| r^2 e^{1-\frac{r}{r_0}}}{r^2(8\pi_F + 2\lambda)}, \quad (59)$$

$$\rho - |p_t| = \frac{-2e^{1-\frac{r}{r_0}} + 2\eta - 2 - \left| \frac{r_0+r}{r^3} \right| r^2 e^{1-\frac{r}{r_0}}}{2r^3(8\pi_F + 2\lambda)}, \quad (60)$$

$$p_t - p_r = \frac{e^{1-\frac{r}{r_0}}(3r_0+r) + 2r(\eta-1)}{2r^3(8\pi_F + 2\lambda)}, \quad (61)$$

and

$$\rho - p_r - 2p_t = \frac{-e^{1-\frac{r}{r_0}} + \eta - 1}{r^2(4\pi_F + \lambda)}. \quad (62)$$

The energy conditions are particularly effective tools for predicting the behavior of strong gravitational fields and the wormhole's geometry. In GR there are seven distinct kinds of energy conditions²⁴, but we are just concentrating on four of them. Null energy condition (NEC), weak energy condition (WEC), strong energy condition (SEC), and dominant energy condition (DEC). These energy conditions are characterized in terms of ρ , p_r , and p_t .

- According to the WEC, any time-like observer's measurement of energy density must be always positive, i.e., $\rho \geq 0$, and for all l , $\rho + p_l \geq 0$.
- According to the SEC, gravity should always be attractive. Regarding the components of the energy-momentum tensor, i.e., for all l , $\rho + p_l \geq 0$ and $\rho + \sum_l p_l \geq 0$.
- The DEC states that any observer's measurement of the energy flux is null or time-like and should be non-negative, i.e., $\rho \geq 0$, which results in $\rho \geq |p_l|$.
- For SEC and WEC basic requirement is NEC, i.e., for all l , $\rho + p_l \geq 0$.
All of the above-mentioned energy conditions are implied to be invalid by the NEC violation.

The NEC follows from the SEC, but the reverse need not be true. Moreover, the SEC need not illustrate the WEC. The DEC implies both the NEC and the WEC, while the reverse doesn't need to be true in each case. Furthermore,

the DEC need not imply the SEC. The SEC²⁵ is a result of gravity's attractive characteristics, and its shape originates directly from an analysis of a spherically symmetrical metric in the general relativity system. To describe the geometry of wormholes, the SEC violation is essential. Although the violation of energy requirements is quite acceptable in some quantum fields, the violation of NEC (commonly referred to as exotic matter) is a basic aspect of static traversable wormholes in the context of GR. Due to this NEC violation, all of the standard energy conditions were also violated.

From Figs. (5-12), we plotted the four energy conditions, and anisotropic factor for different values of λ as a function of r and η with wormhole throat at $r_0 = 0.5$ for the current Finslerian wormhole model in $f(R, T)$ MGT.




FIG. 6. (a) $\rho + p_r = (\rho + p_r)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (NEC term). (b) $\rho + p_r = (\rho + p_r)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (NEC term).




FIG. 7. (a) $\rho + p_t = (\rho + p_t)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (NEC term). (b) $\rho + p_t = (\rho + p_t)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (NEC term).

We will need barotropic EoS for the radial and transverse pressures. we express it by

$$p_r = \omega_r \rho, \quad p_t = \omega_t \rho, \quad (63)$$

where ω_t and ω_r stand for the transverse and radial EoS parameters, respectively. SEC is specified by Eq. (58), which states that the inhomogeneous and anisotropic component fulfills. In this specific instance, $-1 \leq \omega_r \leq 1$ followed by $-1 \leq \omega_t \leq 0$, along with $\omega_r - 1 \geq 0$ and $\omega_r + 1 \leq 0$, and we




FIG. 8. (a) $\rho - |p_r| = (\rho - |p_r|)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (DEC term). (b) $\rho - |p_r| = (\rho - |p_r|)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (DEC term).




FIG. 9. (a) $\rho - |p_t| = (\rho - |p_t|)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (DEC term). (b) $\rho - |p_t| = (\rho - |p_t|)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (DEC term).

observed that $\omega_t \leq -1$ and $\omega_t \geq 0$ respectively. From Eq. 63 ω_r and ω_t can be written by using Eqs. (53-55) as follows,

$$\omega_r = \frac{r_0 e^{1-\frac{r}{r_0}} + (\eta - 1)r}{(e^{1-\frac{r}{r_0}} - \eta + 1)r}, \quad (64)$$

$$\omega_t = \frac{e^{1-\frac{r}{r_0}}(r_0 + r)}{(-e^{1-\frac{r}{r_0}} + \eta - 1)2r}. \quad (65)$$

The following expression is satisfied by the barotropic EoS parameters ω_r and ω_t

$$\omega_r + 2\omega_t + 1 = 0. \quad (66)$$

With reference to the radial EoS parameter ω_r , the transverse pressure p_t can be expressed as follows,

$$p_t = -\frac{1}{2}(1 + \omega_r)\rho. \quad (67)$$

It is evident that $1 + \omega_r \geq 0$ from 13(a). As a result, Eq. (67) suggests that $p_t \geq 0$. Accordingly, Fig. 4 makes this point clear. In accordance with the indicated range of radial coordinate r and the η , with the wormhole throat at $r_0 = 0.5$, the radial EoS parameter ω_r and the transverse EoS parameter ω_t are shown in Fig. 13.




FIG. 10. $\rho + p_r + 2p_t = (\rho + p_r + 2p_t)(r, \eta)$ at $r_0 = 0.5$ (SEC term).




FIG. 11. (a) $T = \rho - p_r - 2p_t = (\rho - p_r - 2p_t)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \geq -12.5$ (SEC term). (b) $T = \rho - p_r - 2p_t = (\rho - p_r - 2p_t)(r, \eta)$ at $r_0 = 0.5$ when $\lambda \leq -12.6$ (SEC term).

Now we are considering F_a stands for the anisotropic wormhole force, F_h for the hydrostatic force, and F_g for the gravity force. The formulas for F_a, F_h and F_g are provided by

$$F_a = \frac{2(p_t - p_r)}{r}, \quad (68)$$

$$F_h = -p'_r, \quad (69)$$

$$F_g = -a'(r)(\rho + p_r). \quad (70)$$

The following is the conservation equation

$$F_a + F_h + F_g = 0. \quad (71)$$

The equations for F_a, F_h , and F_g for the Finslerian wormhole space-time (27) are obtained from Eqs. (53-55), and are used in Eqs. (68-70).

$$F_a = \frac{e^{1-\frac{r}{r_0}}(3r_0 + r) + (\eta - 1)2r}{r^4(8\pi_F + 2\lambda)}, \quad (72)$$




FIG. 12. (a) $\Delta = \Delta(r, \eta)$ the anisotropic factor at $r_0 = 0.5$ when $\lambda \geq -12.5$. (b) $\Delta = \Delta(r, \eta)$ the anisotropic factor at $r_0 = 0.5$ when $\lambda \leq -12.6$.




FIG. 13. (a) $\omega_r = \omega_r(r, \eta)$ radial EoS parameter at $r_0 = 0.5$. (b) $\omega_t = \omega_t(r, \eta)$ transverse EoS parameter at $r_0 = 0.5$.

$$F_h = -\frac{e^{1-\frac{r}{r_0}}(3r_0 + r) + (\eta - 1)2r}{r^4(8\pi_F + 2\lambda)}, \quad (73)$$

$$F_g = a'(r) \left(\frac{e^{1-\frac{r}{r_0}}(r + r_0)}{r^3(8\pi_F + 2\lambda)} \right) = 0. \quad (74)$$

Since we are studying tideless wormholes, we taken $a(r) = \text{constant}$ implies $a'(r) = 0$. For this condition the force due to the contribution of gravity Eq. (74) becomes zero in our study model. The tidal gravitational forces that traveler experiences must be minimal for the wormhole to be traversable. So that the condition $a'(r) = 0$ supports the features of a traversable wormhole. Because of this conservation equation (71), becomes

$$F_a = -F_h. \quad (75)$$

We deduce that the Finslerian wormhole solution is in equilibrium because our observation in Eq. (75) shows that the opposing effects of hydrostatic and anisotropic forces cancel each other.

In Figs. 14 and 15, the variation of the hydrostatic force F_h , and the anisotropic force F_a against the radial coordinate r with the change in the η and at $r_0 = 0.5$ shown for distinct values of λ .




FIG. 14. (a) $F_a = F_a(r, \eta)$ variation of anisotropic force at $r_0 = 0.5$ when $\lambda \geq -12.5$. (b) $F_a = F_a(r, \eta)$ variation of anisotropic force at $r_0 = 0.5$ when $\lambda \leq -12.6$.




FIG. 15. (a) $F_h = F_h(r, \eta)$ variation of hydrostatic force at $r_0 = 0.5$ when $\lambda \geq -12.5$. (b) $F_h = F_h(r, \eta)$ variation of hydrostatic force at $r_0 = 0.5$ when $\lambda \leq -12.6$.

Considering the following formulas are used to express the energy density gradient $\frac{d\rho}{dr}$, radial pressure gradient $\frac{dp_r}{dr}$, and transverse pressure gradient $\frac{dp_t}{dr}$, respectively,

$$\frac{d\rho}{dr} = \frac{e^{1-\frac{r}{r_0}}(r + 2r_0) - (\eta - 1)2r_0}{r_0 r^3(8\pi_F + 2\lambda)}, \quad (76)$$

$$\frac{dp_r}{dr} = \frac{e^{1-\frac{r}{r_0}}(r + 3r_0) + (\eta - 1)2r}{r^4(8\pi_F + 2\lambda)}, \quad (77)$$

$$\frac{dp_t}{dr} = -\frac{e^{1-\frac{r}{r_0}}(r^2 + 3rr_0 + 3r_0^2)}{2r_0 r^4(8\pi_F + 2\lambda)}. \quad (78)$$

Using Eq. (74-76) we have following,

$$\frac{dp_r}{d\rho} = \frac{r_0 e^{1-\frac{r}{r_0}}(r + 3r_0) + (\eta - 1)2rr_0}{r e^{1-\frac{r}{r_0}}(r + 2r_0) - (\eta - 1)2rr_0}, \quad (79)$$

$$\frac{dp_t}{d\rho} = -\frac{e^{1-\frac{r}{r_0}}(r^2 + 3rr_0 + 3r_0^2)}{2[r e^{1-\frac{r}{r_0}}(r + 2r_0) - (\eta - 1)2rr_0]}. \quad (80)$$

Showing the gradient $\frac{d\rho}{dr}$, $\frac{dp_r}{dr}$, and $\frac{dp_t}{dr}$, how they are varying in Figs. (16-18), respectively, for different value of λ that change with r for given value of η for the wormhole throat at $r_0 = 0.5$.




FIG. 16. (a) Variation of $\frac{dp_r}{dr} = \frac{dp_r}{dr}(r, \eta)$ at $r_0 = 0.5$ when $\lambda \ge -12.5$. (b) Variation of $\frac{dp_r}{dr} = \frac{dp_r}{dr}(r, \eta)$ at $r_0 = 0.5$ when $\lambda \le -12.6$.




FIG. 17. (a) Variation of $\frac{dp_t}{dr} = \frac{dp_t}{dr}(r, \eta)$ at $r_0 = 0.5$ when $\lambda \ge -12.5$. (b) Variation of $\frac{dp_t}{dr} = \frac{dp_t}{dr}(r, \eta)$ at $r_0 = 0.5$ when $\lambda \le -12.6$.

IV. DISCUSSION AND RESULTS

Modified theories have the flexibility to experiment with the effective stress-energy tensor to find an alternative approach for solving the exotic matter problem while still adhering to all energy conditions. $f(R, T)$ gravity is one of the modified theories of gravity that has received a lot of interest. With an exponential shape function that corresponds




FIG. 18. Variation of $\frac{dp_t}{dr} = \frac{dp_t}{dr}(r)$ at wormhole throat $r_0 = 0.5$.

to the family $f(R, T)$ of extended theories of gravity, we have developed a Finslerian wormhole model in our present study. We focused our studies to the case $f(R, T) = R + f(T)$, where $f(T) = 2\lambda T$ and λ is constant parameter. The sensitivity of $f(R, T)$ gravity wormholes depends on a range of parameter choices. Due to this, several scholars have been working on $f(R, T)$ MGT among those,²⁶ discover that the parameter λ has a significant impact in determining the composition of matter in wormholes. In Ref.¹⁸, authors investigate the wormhole model using an exponential shape function within the context of Finsler geometry by taking $\lambda = 0$. The equilibrium configurations of white dwarfs are investigated by²⁷ using an MGT that depends on the parameter λ . As a result, numerous physicists and geometrists have this generalized geometry as interesting. Consequently, the Finslerian space-time manifold is of significance to us when we study the wormhole configuration in $f(R, T)$ MGT. In general, the "exotic matter" violates the weak/null energy conditions and has the unique capability to keep the wormhole tunnel open. We can reveal that NEC is an artifact of the general relativity that supports a wormhole. A key requirement of static traversable wormholes, according to GR, is the violation of different energy conditions, often known as exotic matter.

In the current work, Fig. 1 shows that the exponential shape function Eq. (52) meets all essential geometric conditions, and as a result, we obtained the structure of the wormhole. We plotted WEC, NEC, DEC, and SEC energy conditions in Figs. (5-12). We plotted WEC in Fig. 5 and observed that in Fig. 5(a) energy density condition is negative and also tends to zero with the change of radial coordinate r with η at $r_0 = 0.5$, for the value $\lambda \ge -12.5$ and in Fig. 5(b) energy density condition is positive and also tends to zero with the change of radial coordinate r with η at $r_0 = 0.5$, for the value $\lambda \le -12.6$. Therefore, we conclude that WEC is violated in the Finslerian wormhole $f(R, T)$ gravity model for the value $\lambda \ge -12.5$, within the specified range of r and η .

In the Figs. (6-9) we plotted the energy condition terms, using the specified range of r and η at $r_0 = 0.5$, and found $\rho + p_r$ in Fig. 6, $\rho + p_t$ in Fig. 7, $\rho - |p_r|$ in Fig. 8, and $\rho - |p_t|$ in Fig. 9, negative for the value of $\lambda \ge -12.5$ and positive for the value of $\lambda \le -12.6$. With Eq. (56), we determined that the η is not a factor in the NEC term $\rho + p_r$, which is the function of r only. But we have demonstrated the term $\rho + p_r$ graphically in a 3-D plot in Fig. 6 and discovered that the result is negative for the value $\lambda \ge -12.5$ in Fig. 6(a), and positive for the value $\lambda \ge -12.6$ in Fig. 6(b). From Fig. 7(a) we observed $\rho + p_t$ is negative for $\lambda \ge -12.5$, and positive for $\lambda \le -12.6$. As a result, we conclude NEC is violated in Finslerian $f(R, T)$ gravity at $\lambda \ge -12.5$, hence Finslerian wormhole to be traversable. Also, we observed DEC in Fig. 9(a) violates for the value $\lambda \ge -12.5$. From these findings, we draw the following conclusions in the framework of Finslerian $f(R, T)$ gravity the wormholes must be filled with exotic matter.

From Eq. (58) in Fig. 10 $\rho + p_r + p_t$ is equal to zero for

the given range of r and η , and we observed the violation of SEC. The η determines whether the Finslerian wormhole is attractive or repulsive. According to Fig. 12(a), the Finslerian wormhole is repulsive for $1 \geq \eta \geq 0.8$, it is initially repulsive and then becomes attractive after crossing the certain value of r for $0.2 \leq \eta < 0.8$ for $\lambda \geq -12.5$, and it is opposite when $\lambda \leq -12.6$ in Fig. 12(b). We observed from Fig. 13(a) that the radial EoS parameter ω_r takes the values between -1 and 1 for the specified value of r and η . Further, if we replace \bar{F} with the Finslerian sphere, then ω_r is obtained to be non-negative with values less than 1. The transverse EoS parameter ω_t lies between the value of -1 and 0 in Fig. 13(b)²⁸ because of Eq. (58) and $-1 \leq \omega_r \leq 1$. The wormhole matter distribution is affected by three forces because of anisotropic pressure. The Finslerian wormhole solution in equilibrium as a result of the combined effect of the anisotropic force F_a and hydrostatic force F_h in Fig. 14(a) and Fig. 15(a) respectively, for $\lambda \leq -12.5$. In the beginning, the anisotropic force and the hydrostatic force have positive values for the range $1 \geq \eta \geq 0.6$ with a variation of r and at the value of $\lambda \geq -12.5$, and negative values for the value of $0.2 \leq \eta < 0.6$ respectively. As a result, they are repulsive and attractive. Their nature changes as r reach a certain value.

The energy density gradient $\frac{dp}{dr}$ is positive in Fig. 16(a) for $\lambda \geq -12.5$ and negative in Fig. 16(b) for $\lambda \leq -12.6$, with change of r for each value of the η at $r_0 = 0.5$. We can observe from Fig. 17(a) that the radial pressure gradient $\frac{dp_r}{dr}$ is found to be dependent on the value of η . $\frac{dp_r}{dr}$ is positive For $\eta = 0.6, 0.8$ and 1, for distinct values of r . But $\frac{dp_r}{dr}$ is positive for $\eta = 0.2$ until $r = 0.8$ and for $\eta = 0.4$ is positive until $r = 0.9$, after it turns negative due to the change in r at $\lambda \geq -12.5$, but it is different when $\lambda \leq -12.6$ in Fig. 17(b). From Fig. 17, we observe the transverse pressure gradient $\frac{dp_t}{dr}$, which is independent of η , is negative for $\lambda \geq -12.5$ and positive for $\lambda \leq -12.6$ at $r_0 = 0.5$ for the specified values of r .

Energy conditions	Terms	$\lambda = -12.5$	$\lambda = -12.6$
	p_r	≤ 0	≥ 0
	p_t	≥ 0	≤ 0
WEC	ρ	≤ 0	≥ 0
NEC	$\rho + p_r$	≤ 0	≥ 0
	$\rho + p_t$	≤ 0	≥ 0
DEC	$\rho - p_r $	≤ 0	≥ 0
	$\rho - p_t $	≤ 0	≥ 0
SEC	$\rho + p_r + 2p_t$	$= 0$	$= 0$
	$\rho - p_r - 2p_t$	≤ 0	≥ 0

TABLE I. Summary of a result for the energy conditions at $r_0 = 0.5$, $r \geq 0$ ($0 \leq r \leq 25$), and $\eta \geq 0$ ($0 \leq \eta \leq 1$) with different value of λ

V. CONCLUSION

In the present paper, we explore an extended gravity model $f(R, T)$, it has arbitrary connectivity between geometry and

matter in the Finslerian framework, which is represented by the trace of the stress-energy tensor. Therefore, the $f(R, T)$ gravity model's assumptions could result in some significant deviations from those of standard general relativity or other generalized gravity models. By this model, the gravitational field equations have been derived. We constructed an exponential Finslerian wormhole that relates to the $f(R, T) = R + f(T)$ where $f(T) = 2\lambda T$ of extended theories of gravity. Here, the parameter λ plays a very important role in the construction of an exponential Finslerian wormhole. Many physicians and geometers work on $f(R, T)$ gravity and give the importance of parameter λ in the construction of a wormhole.

The energy conditions WEC, NEC, and DEC are violated in our $f(R, T)$ gravity model when $\lambda \leq -12.5$, but SEC is equal to zero (see table (I)). This violation provides strong evidence for the presence of wormholes and exotic matter in the Finslerian world. The wormholes are discovered to be phantom when $\omega_r < -1$ ²⁹, but in our $f(R, T)$ gravity wormhole model is not phantom because we obtain the radial EoS parameter $\omega_r > -1$. Therefore, background theory is crucial for understanding geometry, determining the type of matter distribution, and determining the type of filled fluid. Observing all our exponential Finslerian wormhole models with $f(R, T)$ MGT is true physically, which is the overall conclusion. Finally, the $f(R, T)$ gravity model accurately captures the characteristics of wormholes in the context of Finslerian space-time, which can be visualized 3-D state wormhole model.

ACKNOWLEDGMENT

The author Manjunath Malligawad acknowledges the financial support for this study was provided by Govt. of Karnataka, India, Backward Classes Welfare Department(BCWD) (Application No:2021PHD9521296). We also want to express our gratitude to the editor and the honorable referee for their insightful remarks and suggestions, which allowed us to significantly improve the article.

APPENDIX

We constructed embedded 2-D and 3-D diagrams for the shape function Eq. (52) for better visualisation of the wormhole. We used an equatorial plane $\theta = \frac{\pi}{2}$ at a fixed time or $t = \text{constant}$, and $\eta = 1$, from these conditions Eq. (27) reduce into the form

$$F^2 = - \left(1 - \frac{b(r)}{r} \right)^{-1} dr^2 - r^2 d\phi^2. \quad (81)$$

The above equation is represent in cylindrical coordinates as

$$F^2 = -dz^2 - dr^2 - r^2 d\phi^2. \quad (82)$$

In Euclidean three dimensional space $z = z(r)$ represents the embedded surface, we can rewrite Eq. (82) as

$$F^2 = - \left(1 + \left(\frac{dz}{dr} \right)^2 \right) dr^2 - r^2 d\phi^2, \quad (83)$$

comparing Eq. (81) and Eq. (83)

$$\frac{dz}{dr} = \pm \sqrt{\left(1 - \frac{b(r)}{r}\right)^{-1} - 1}, \quad (84)$$

using Eq. (84) we plotted the embedded surface of the wormhole.

- ¹H. Weyl, “Feld und Materie”, *Ann. Phys. Berl.* **370**(14), 541–563 (1921).
- ²A. Einstein, N. Rosen, “The particle problem in the general theory of relativity”, *Phys. Rev.* **48**(1), 73–77 (1935).
- ³Robert W. Fuller, John A. Wheeler, “Causality and Multiply Connected Space Time”, *Phys. Rev.* **128**(2), 919–929 (1962).
- ⁴M. S. Morris, K. S. Thorne, “Wormholes in space-time and their use for interstellar travel: a tool for teaching general relativity”, *Am. J. Phys.* **56**, 395–412 (1988).
- ⁵D. Hochberg, M. Visser, “General Dynamic Wormholes and Violation of the Null Energy Condition”, *arXiv:gr-qc/9901020*, (1999).
- ⁶Gregory J. Galloway, “On the topology of the domain of outer communication”, *Class. Quantum Grav.* **12**, L99 (1995).
- ⁷J. L. Friedman, K. Schleich, D. M. Witt, “Topological Censorship”, *Phys. Rev. Lett.* **71**, 1486–1489 (1993).
- ⁸Francisco S. N. Lobo, Miguel A. Oliveira, “Wormhole geometries in $f(R)$ modified theories of gravity”, *Phys. Rev. D* **80**(10), 104012 (2009).
- ⁹T. Harko, “ $f(R, T)$ gravity”, *Phys. Rev. D* **84**, 024020 (2011).
- ¹⁰E. H. Baffou, M. J. S. Houndjo, M. E. Rodrigues, A. V. Kpadonou, J. Tossa, “Cosmological Evolution in $f(R, T)$ theory with Collisional Matter”, *Phys. Rev. D* **92**, 084043 (2015).
- ¹¹P. H. R. S. Moraes, P. K. Sahoo, “Modeling wormholes in $f(R, T)$ gravity”, *Phys. Rev. D* **96**, 044038 (2017).
- ¹²S. Najafi, T. Rostami, S. Jalalzadeh, “Five dimensional cosmological traversable wormhole”, *Ann. Phys.* **354**, 288–297 (2015).
- ¹³P. H. R. S. Moraes, “An exponential shape function for wormholes in modified gravity”, *Chin. Phys. Lett.* **36**, 120401 (2019).
- ¹⁴P. H. R. S. Moraes, P. K. Sahoo, “Wormholes in exponential $f(R, T)$ gravity”, *Eur. Phys. J. C* **79**, 677 (2019).
- ¹⁵Z. Chang, X. Li, “Modified Newton’s gravity in Finsler space as a possible alternative to dark matter hypothesis”, *Phys. Rev. B* **668**, 453–456 (2008).
- ¹⁶F. Rahman, N. Paul, A. Banerjee, S. S. De, S. Ray, A. A. Usmani, “The Finslerian wormhole models”, *Eur. Phys. J. C* **76**, 246 (2016).
- ¹⁷T. Clifton, Pedro G. Ferreira, A. Padilla, C. Skordis, “Modified gravity and cosmology”, *Phys. Rep.* **513**, 1–189 (2012).
- ¹⁸H. M. Manjunatha, S. K. Narasimhamurthy, “The wormhole model with an exponential shape function in the Finslerian framework”, *Chin. J. Phys.* **77**, 1561–1578 (2022).
- ¹⁹H. Akbar-Zadeh, “Sur les espaces de Finsler a courbures sectionnelles constantes”, *Acad. Roy. Belg. Bull. Cl. Sci.* **74**, 281–322 (1988).
- ²⁰R. A. Konoplya, “How to tell the shape of a wormhole by its quasinormal modes”, *Phys. Lett. B* **784**, 43–49 (2018).
- ²¹X. Li, Z. Chang, “Exact solution of vacuum field equation in Finsler space-time”, *Phys. Rev. D* **90**, 064049 (2014).
- ²²M. K. Mak, T. Harko, “Anisotropic stars in general relativity”, *Proc. Roy. Soc. Lond. A* **459**, 393–408 (2003).
- ²³X. Li, “Special Finslerian generalization of the Reissner-Nordström space-time”, *Phys. Rev. D* **98**, 084030 (2018).
- ²⁴M. Visser, “Lorentzian Wormholes: From Einstein to Hawking”, (AIP Press, Springer, New York) (1996).
- ²⁵A. K. Mishra, U. K. Sharma, V. C. Dubey, A. Pradhan, “Traversable wormholes in $f(R, T)$ gravity”, *Astrophys. Space Sci.* **365**, 34 (2020).
- ²⁶A. Chanda, S. Dey, B. C. Paul, “Morris-Thorne Wormholes in $f(R, T)$ modified theory of gravity”, *Gen. Rel. Gra.* **53**, 78 (2021).
- ²⁷G. A. Carvalho, R. V. Lobato, P. H. R. S. Moraes, José D. V. Arbañil, R. M. Marinho Jr, E. Otoniel, M. Malheiro, “Stellar equilibrium configurations of white dwarfs in the $f(R, T)$ gravity”, *Eur. Phys. J. C* **77**, 871 (2017).
- ²⁸M. Cataldo, S. Campo, “Two-fluid evolving Lorentzian wormholes”, *Phys. Rev. D* **85**, 104010 (2012).
- ²⁹P. H. R. S. Moraes, P. K. Sahoo, Barkha Taori, Parbati Sahoo, “Phantom energy-dominated universe as a transient stage in $f(R)$ cosmology”, *Int. J. Mod. Phys. D* **28**, 1950124 (2019).