

# Examination of Nonlinear Longitudinal Processes with Latent Variables, Latent Processes, Latent Changes, and Latent Classes in the Structural Equation Modeling Framework: The R package **nlpsem**

Jin Liu

Virginia Commonwealth University

---

## Abstract

We present the R package **nlpsem**, which provides a comprehensive set of functions to assess longitudinal processes with individual measurement occasions within the structural equation modeling (SEM) framework. This package focuses on providing computational tools for nonlinear longitudinal models, particularly intrinsically nonlinear models, across four distinct scenarios: (1) univariate longitudinal processes captured by latent variables, with or without covariates, including time-invariant covariates (TICs) and time-varying covariates (TVCs); (2) multivariate longitudinal processes for evaluating correlations or causations between longitudinal variables; (3) multiple-group frameworks for models in scenarios (1) and (2), enabling the examination of differences between manifested classes; and (4) mixture models for scenarios (1) and (2), assuming that trajectories originate from heterogeneous latent classes. By interfacing with the R package **OpenMx**, **nlpsem** enables flexible specification of structural equation models and generates maximum likelihood estimators using the full information maximum likelihood technique. The package includes an algorithm to obtain initial values from raw data, thereby facilitating computation and enhancing the likelihood of model convergence. Additionally, **nlpsem** provides functions for goodness-of-fit analyses, clustering analyses, plots, and predicted trajectories. This paper constitutes a companion to the package, detailing each model scenario, the estimation technique, implementation details, output interpretation, and showcasing examples through a dataset on intelligence development.

*Keywords:* structural equation modeling, latent growth curve models, latent change score models, multivariate growth models, multiple-group growth modeling, growth mixture modeling, time-invariant covariates, time-varying covariates, individual measurement occasions.

---

## 1. Introduction

Longitudinal data are prevalent across various fields, including biomedical, behavioral, and social sciences. Their repeated measurements on same subjects over time provide unique insights into patterns of change, distinguishing them from cross-sectional studies. However, such richness in data also brings forth challenges in data analysis, requiring specialized statistical methods and software tools tailored for these complexities.

As statistical software continues to evolve, it becomes an indispensable ally for researchers,

furnishing them with sophisticated tools to unravel intricate patterns in data. With the ever-growing complexity and volume of longitudinal data, there is an escalating demand for robust computational solutions adept at navigating both the foundational and the more nuanced challenges inherent to this type of data. Analyzing longitudinal data transcends merely tracking changes over time. It delves deep into uncovering individual differences in change trajectories, discerning how initial characteristics influence future paths, scrutinizing associations between multiple longitudinal variables, and extracting profound insights from the patterns that materialize as time progresses.

Two primary modeling families are commonly employed for analyzing longitudinal data: the mixed-effects modeling framework and the structural equation modeling (SEM) framework. These frameworks utilize latent variables to capture the underlying patterns of change in the data. In the context of linear trajectories, these latent variables are often referred to differently based on the modeling approach:

- In the SEM framework, they are termed *growth factors*. The mean and variance of these growth factors provide insights into the average trajectory and individual variability around this average, respectively.
- In the mixed-effects modeling framework, the equivalent terms are *fixed effects* (for the mean trajectory) and *random effects* (for individual deviations from the average).

For instance, in a simple linear growth model, an intercept (representing the starting point) and a slope (indicating the rate of change) are the primary growth factors. Estimating the means (or fixed effects) and variances (or random effects) of these components enables the assessment of between-individual differences in within-individual change processes. While the linear longitudinal model is frequently employed and can often provide a reasonable approximation of change within limited time spans, it may not always be sufficient. Especially when the process under investigation spans a long duration or when measurements are taken frequently, trajectories can exhibit nonlinearity with varying rates of change. In such scenarios, relying solely on linear models can lead to oversimplifications, failing to capture the intricacies of individual trajectories. This necessitates the exploration of models with nonlinear functions to better understand and assess between-individual differences in within-individual change.

### 1.1. Foundational Challenges in Nonlinear Longitudinal Analysis and Existing Solutions

Longitudinal data analysis, with its intricate nonlinear change patterns, poses numerous challenges. As outlined in Chapter 9 of [Grimm, Ram, and Estabrook \(2016\)](#), nonlinear longitudinal models can be categorized into three distinct types based on their nonlinearity with respect to: (1) time, (2) parameters, and (3) growth factors. The third category, termed as intrinsically nonlinear<sup>1</sup>, is particularly complex. This type is defined by a growth function

---

<sup>1</sup>The term ‘intrinsic nonlinearity’ was initially used to describe a nonlinear regression function where at least one of the first derivatives of the function with respect to the parameters depends on one or more of the parameters ([Bates and Watts 1988](#), Chapter 2). Subsequently, researchers extended this concept to encompass nonlinear longitudinal models ([Harring 2014](#)). According to the original definition, both Type II and Type III nonlinear longitudinal models can be considered intrinsically nonlinear. However, ‘intrinsic nonlinearity’ in this project exclusively refers to Type III nonlinear longitudinal models, which necessitate multidimensional integrations of joint likelihood over the growth factors and do not possess a closed-form likelihood function following [Rohloff, Kohli, and Chung \(2022\)](#).

derivative that depends on other growth factors with respect to a growth factor. Both the mixed-effects modeling and latent growth curve modeling (LGCM) frameworks, members of the SEM family, are proficient in handling these three types. They excel particularly in fitting the first two types, yielding results that are both theoretically and empirically equivalent (Bauer 2003; Curran 2003).

There are a plethora of computational tools available for analyzing longitudinal data. For the first two types of nonlinear longitudinal models, there exists a rich array of specialized tools. Within the mixed-effects modeling paradigm, packages such as **lme4** (Bates, Mächler, Bolker, and Walker 2014, 2015), **nlme** (Pinheiro and Bates 2000; Pinheiro, Bates, and R Core Team 2023), and **lcm** (Proust-Lima, Philipps, and Liqueur 2017) have garnered significant recognition. While **lme4** and **nlme** are esteemed for their robust capabilities, **lcm** distinguishes itself by adeptly fitting joint and mixture models. On the other hand, the SEM framework, represented by tools like **OpenMx** (Neale, Hunter, Pritikin, Zahery, Brick, Kirkpatrick, Estabrook, Bates, Maes, and Boker 2016; Pritikin, Hunter, and Boker 2015; Hunter 2018; Boker, Neale, Maes, Wilde, Spiegel, Brick, Estabrook, Bates, Mehta, von Oertzen, Gore, Hunter, Hackett, Karch, Brandmaier, Pritikin, Zahery, and Kirkpatrick 2020) and **Mplus** (Muthén and Muthén 2017), provides a more generalized approach. However, this enhanced flexibility necessitates a meticulous effort in model specification and rigorous validation.

Fitting the third type of nonlinear longitudinal model, specifically the intrinsically nonlinear model, often requires specialized approximation techniques. The marginal maximum likelihood estimation (MLE) procedure (Harring, Cudeck, and du Toit 2006; Du Toit and Cudeck 2009; Cudeck and Harring 2010) is the go-to method within the mixed-effects modeling domain, while the SEM framework typically employs the Taylor series expansion (Browne and du Toit 1991; Preacher and Hancock 2015). Notably, Zopluoglu, Harring, and Kohli (2014) introduced the R routine **fitPMM**, which implements the marginal MLE procedure for longitudinal models characterized by a linear-linear functional form and a random knot, a hallmark of Type III nonlinear longitudinal models. However, this tool is limited to a specific functional form and does not cater to time-invariant covariates. This limitation hinders researchers from fully exploring how individual attributes impact trajectory heterogeneity.

The **nlpssem** package fills this gap, providing a comprehensive suite of tools tailored for intrinsically nonlinear LGCMs within the SEM framework. The current version of **nlpssem** focuses on the most prevalent and practically significant functional forms, such as the negative exponential function, which underscores individual growth rate ratios (Sterba 2014), the Jenss-Bayley function that centers on individual growth acceleration ratios (Grimm *et al.* 2016, Chapter 12), and the bilinear spline function (or linear-linear piecewise function) that highlights individual knots (Liu, Perera, Kang, Kirkpatrick, and Sabo 2021a). Recognizing the complexities associated with Type III nonlinear models, **nlpssem** also provides parsimonious versions aligned with Type II models and supports estimations for models with quadratic functional forms (Type I). While the primary focus is on nonlinear models, **nlpssem** does accommodate linear longitudinal models to cater to a broader research audience.

Unstructured measurement schedules pose a significant challenge in longitudinal data analysis, a phenomenon prevalent across diverse research domains, including biomedical, behavioral, and social sciences. For example, in clinical trials, individual patient site visits might be slated for varying study days<sup>2</sup>. Such irregularities also arise when time measurements are exact or

---

<sup>2</sup>The term ‘study day’ in clinical trials typically commences at either randomization or dosing, starting

when data collection is self-initiated. A common analytical strategy partitions the assessment duration into several time intervals, allowing for a single response within each interval for each participant.

However, this approach has its drawbacks. Research by Blozis and Cho (2008) and Coulombe, Selig, and Delaney (2015) underscores that overlooking individual time variations can skew results. The mixed-effects modeling paradigm offers a solution by treating measurement time as a continuous variable, a feature seamlessly incorporated in renowned packages like **lme4** (Bates *et al.* 2014, 2015), **nlme** (Pinheiro and Bates 2000; Pinheiro *et al.* 2023), and **lcm** (Proust-Lima *et al.* 2017). The SEM framework, however, presents unique challenges. Predominantly, its computational tools cater to wide-format longitudinal data, making the integration of individual measurement times intricate. To address this, Mehta and West (2000) and Mehta and Neale (2005) introduced the ‘definition variables’ technique. In this method, observed variables, termed ‘definition variables’, modify model parameters to align with individual-specific values. While platforms like **OpenMx** (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020) and **Mplus** (Muthén and Muthén 2017) are robust, the process of specifying ‘definition variables’ within them can be intricate (Sterba 2014; Grimm *et al.* 2016). This underscores the pressing need for tools adept at effortlessly managing such specifications, especially when navigating the complexities of unstructured measurement schedules in longitudinal research. Addressing this gap, **nlpsem** is specifically designed to incorporate individual measurement occasions, streamlining the process for researchers. For a detailed comparison of the strengths and limitations of **nlpsem** in relation to other tools in longitudinal data analysis, please refer to Table 1.

=====  
 Insert Table 1 about here  
 =====

## 1.2. Advanced Challenges in Longitudinal Data Analysis: The **nlpsem** Solution

In the domain of longitudinal data analysis, researchers are confronted with evolving research objectives and intricate data structures. Particularly for nonlinear trajectories, the challenge intensifies: it is not merely about understanding the time-dependent status, but also about the time-dependent change. Such estimations are paramount, as they provide a deeper insight into the underlying dynamics of the data. Furthermore, when multiple longitudinal variables come into play, the intricacy intensifies. Beyond merely charting the progression of each variable, it is imperative to decode the interrelationships they share. Whether these associations represent simple correlations or extend to causative interactions, akin to those in cross-sectional data, they mandate rigorous scrutiny.

To traverse these multifaceted challenges, researchers require tools that epitomize sophistication and versatility. While packages like **lcm** offer valuable insights, especially in discerning correlations between multiple longitudinal outcomes, they sometimes fall short of the expansive flexibility inherent to the SEM framework. This is where **nlpsem** emerges as a beacon. It embodies the versatility of SEM, adeptly navigating nonlinear trajectories and the intricate dynamics between multiple longitudinal variables. Beyond the purview of mixed-effects

models, SEM offers a comprehensive perspective on the multifarious aspects of longitudinal data. Within this ambit, **nlpsem** encompasses:

1. **Estimation of Rate of Change Over Time:** Delving into both intrinsic and extrinsic nonlinear processes, as elucidated in [Liu and Perera \(2023a\)](#); [Grimm, Zhang, Hamagami, and Mazzocco \(2013b\)](#).
2. **Baseline Characteristics' Influence on Trajectories:** Unearthing how foundational attributes shape longitudinal progressions, as detailed in [Liu \*et al.\* \(2021a\)](#).
3. **Holistic Exploration of Joint Processes:** This includes:
  - **Impact of Time-Varying Covariates (TVC) on Outcomes:** A scenario where one process serves as a longitudinal outcome, while another acts as a TVC, as explored in [Liu \(2022a\)](#) and [Liu and Perera \(2023b\)](#).
  - **Interplay Among Multiple Longitudinal Processes:** A situation where all processes are viewed as longitudinal outcomes, with a focus on their interrelations, as discussed in [Liu and Perera \(2021\)](#).
  - **Dissecting Direct and Mediated Effects of Predictors:** A methodology that posits a predictor influencing a longitudinal outcome both directly and via a mediator, as detailed in [Liu and Perera \(2022a\)](#).
4. **Temporal Variations Across Groups:** A study of the evolutionary patterns of different groups over time.
5. **Unearthing Latent Classes:** A deep dive into concealed data structures and the variability they exhibit, as explored in [Liu and Perera \(2022c\)](#).

The package **nlpsem** advances from the rudimentary univariate longitudinal processes, such as estimating growth status or rate of change, to the more complex multivariate processes. Specifically, it encompasses: (1) longitudinal outcomes impacted by time-varying covariates, (2) synchronized and correlated growth models, and (3) longitudinal mediation models. In addition, **nlpsem** offers tools for multiple group and mixture models, which are in harmony with the models described above. Computationally, all estimations within the package are bolstered by the robustness of **OpenMx** ([Neale \*et al.\* 2016](#); [Pritikin \*et al.\* 2015](#); [Hunter 2018](#); [Boker \*et al.\* 2020](#)), a trusted name within the SEM framework lauded for its accuracy. A distinguishing feature of **nlpsem** is its integration of the 'definition variables' methodology, which assures a smooth incorporation of unstructured temporal designs.

The optimization process in any statistical package is fundamental for ensuring accuracy and efficiency. In this regard, **nlpsem** is particularly robust. It seamlessly integrates with multiple renowned optimization engines provided by **OpenMx**. These include:

1. NPSOL, originating from the Nonlinear Programming at Systems Optimization Laboratory ([Gill, Murray, Saunders, and Wright 1986](#)).
2. SLSQP, a method grounded in Sequential Least-Squares Quadratic Programming ([Johnson 2014](#); [Kraft 1994](#)).
3. CSOLNP, a C++-based optimizer developed for Solving Nonlinear Programs ([Zahery, Maes, and Neale 2017](#)).

These engines are recognized for their precision and efficiency across various applications, and their compatibility with **nlpsem** ensures optimal results for users.

To summarize, in the vast landscape of statistical software dedicated to longitudinal analysis, **nlpsem** provides a distinctive edge. Its ability to handle inherently nonlinear longitudinal models addresses a fundamental need in analyzing real-world processes that often exhibit non-linear patterns. Beyond this, based on our extensive literature review from the past three years, **nlpsem** stands alone in its capability to provide multiple advanced analyses. These analyses cater to complex scenarios that researchers frequently grapple with in modern research. Combined with the trusted optimization engines it incorporates, users of **nlpsem** can be confident not only in the versatility but also in the accuracy of their results.

As for the structure of this article, it unfolds as follows: Section 2 delineates the specifications of each model within the package. Section 3 delves into the intricacies of the full information maximum likelihood (FIML) technique. Section 4 elaborates on the implementation of the seven estimation functions, shedding light on the initial value algorithm. Section 5 discusses computational nuances and post-fit analyses. Section 6 showcases examples rooted in the package’s dataset, and finally, Section 7 provides both practical insights and methodological considerations, alongside potential future directions.

## 2. Model Specification

In this section, we outline the specifications for each modeling family incorporated in the package. The initial two subsections detail the modeling frameworks for univariate longitudinal processes, specifically latent growth curve models (LGCMs) and latent change score models (LCSMs), both with and without time-invariant covariates. Subsections 2.3-2.5 focus on frameworks designed for multivariate longitudinal processes, encompassing LGCMs or LCSMs with a time-varying covariate (TVC), multivariate LGCMs or LCSMs, and longitudinal mediation models. Concluding the section, Subsections 2.6 and 2.7 introduce multiple group and mixture models. These models challenge the single population assumption present in the aforementioned frameworks, offering extensions for heterogeneous populations characterized by both observed and latent classes.

### 2.1. Latent Growth Curve Models

The LGCM, a modeling family in the SEM framework, is often employed to analyze growth status over time. As introduced in Section 1, the linear and first two types of nonlinear LGCMs are mathematically and empirically equivalent to the corresponding model in the mixed-effects modeling framework. This subsection briefly describes the LGCM, providing an overview of three intrinsically nonlinear functional forms, their corresponding reduced non-intrinsically nonlinear functions, and the other two commonly used functional forms, linear and quadratic curves, with unstructured time frames. A general form of the LGCM with individual measurement occasions is

$$\mathbf{y}_i = \mathbf{\Lambda}_i \times \boldsymbol{\eta}_i + \boldsymbol{\epsilon}_i, \quad (1)$$

where  $\mathbf{y}_i$  is a  $J \times 1$  vector of the repeated measurements of individual  $i$  (where  $J$  is the number of measures). The elements in the vector  $\boldsymbol{\eta}_i$  ( $K \times 1$ ) are often called growth factors, which are



latent variables that constitute growth status in LGCMs for the  $i^{th}$  individual (where  $K$  is the number of growth factors). Furthermore,  $\mathbf{\Lambda}_i$  is a  $J \times K$  matrix of the corresponding factor loadings. Note that the subscript  $i$  in  $\mathbf{\Lambda}_i$  indicates that the factor loadings, which are functions of measurement times, are at the individual level. The vector  $\boldsymbol{\epsilon}_i$  ( $J \times 1$ ) represents residuals of individual  $i$  (usually assumed to follow an identical and independent normal distribution). That is,  $\boldsymbol{\epsilon}_i \sim \text{MVN}(\mathbf{0}, \theta_\epsilon \mathbf{I})$ , where  $\theta_\epsilon$  is the residual variance and  $\mathbf{I}$  is  $J \times J$  identity matrix). The vector of growth factors can be further written as deviations from the corresponding mean values

$$\boldsymbol{\eta}_i = \boldsymbol{\mu}_\eta + \boldsymbol{\zeta}_i, \quad (2)$$

where  $\boldsymbol{\mu}_\eta$  (a  $K \times 1$  vector) represents growth factor means, and  $\boldsymbol{\zeta}_i$  (a  $K \times 1$  vector) denotes deviations of the  $i^{th}$  individual from factor means. Suppose it is of interest to evaluate the effects of TICs on the heterogeneity of growth factors and, consequently, on growth curves. In that case, one may further regress growth factors on the covariates,

$$\boldsymbol{\eta}_i = \boldsymbol{\alpha} + \mathbf{B}_{\text{TIC}} \times \mathbf{X}_i + \boldsymbol{\zeta}_i, \quad (3)$$

where  $\boldsymbol{\alpha}$  is a  $K \times 1$  vector of growth factor intercepts (equivalent to the mean vector of growth factors if the TICs are centered), and  $\mathbf{B}_{\text{TIC}}$  is a  $K \times m$  matrix of regression coefficients (with  $m$  representing the number of TICs) from TICs to growth factors. The vector  $\mathbf{X}_i$  ( $m \times 1$ ) is for TICs, which could be continuous or categorical, of individual  $i$ . In the current version of **nlpsem**, growth factors in Equation 1 are assumed to follow a (conditional) multivariate normal distribution, a common assumption in practice. That is,  $\boldsymbol{\zeta}_i \sim \text{MVN}(\mathbf{0}, \boldsymbol{\Psi}_\eta)$ , where  $\boldsymbol{\Psi}_\eta$  is a  $K \times K$  matrix and indicates growth factor variance and unexplained growth factor variance in the scenarios without and with TICs, respectively.

Table 2 provides the model specification of LGCM with multiple commonly used functional forms and corresponding interpretation of growth coefficients. For instance, consider the linear functional form, the most straightforward function. There are two free coefficients in an individual linear trajectory: intercept ( $\eta_{0i}$ ) and linear slope ( $\eta_{1i}$ ). The intercept and slope are allowed to vary from individual to individual. Examining individual differences in the intercept and slope, and thus, the between-individual differences in within-individual changes is one primary interest when exploring longitudinal records (Biesanz, Deeb-Sossa, Papadakis, Bollen, and Curran 2004; Zhang, McArdle, and Nesselroade 2012; Grimm *et al.* 2013b).

=====  
 Insert Table 2 about here  
 =====

Although the linear function's coefficients are straightforward to interpret, they do not help effectively describe more complex change patterns. When the trajectory of a longitudinal process exhibits a certain degree of nonlinearity to time, particularly if the process is observed over a long period, more complex functional forms with additional growth coefficients are needed to depict change patterns. For instance, an individual quadratic functional form, which includes an acceleration growth factor ( $\eta_{2i}$ , representing the quadratic component of change) in addition to the intercept ( $\eta_{0i}$ ) and the linear component of change ( $\eta_{1i}$ ), serves as a candidate for describing nonlinear trajectories, particularly when estimating growth acceleration is of interest. A LGCM with a quadratic functional form belongs to Type I of nonlinear longitudinal models, as its factor loadings are functions that depend solely on measurement

times. In practice, other nonlinear functions listed in Table 2 can also provide valuable insights regarding growth status. For example,  $\eta_{1i}$  in an individual negative exponential function indicates the individual's growth capacity.

As illustrated in Table 2, the second term of the individual growth curve of the negative exponential function contains two growth factors,  $\eta_{1i}$  and  $b_i$ . This implies that the derivatives of the growth curve with respect to  $\eta_{1i}$  or  $b_i$  would be functions of  $b_i$  and  $\eta_{1i}$ , respectively (i.e., intrinsically nonlinear model). Consequently, the negative exponential function with individual coefficient  $b$  cannot be modeled directly in the SEM framework, as the factor loadings matrix cannot depend on a growth factor. In practice, two ways to address this issue are: (1) performing Taylor series expansion to linearize the function (Preacher and Hancock 2015), and (2) using a reduced model assuming that the growth rate ratio is consistent across the entire population. These two approaches can also be employed to handle  $c_i$  in the Jenss-Bayley function and  $\gamma_i$  in the bilinear spline function with an unknown knot. After linearization, the factor-loading matrices of these intrinsically nonlinear models depend on the growth factor means and measurement occasions and can thus be fit in the SEM framework. Technical details of linearization via Taylor series expansion for the negative exponential function, Jenss-Bayley function, and bilinear spline function with an unknown knot are documented in Sterba (2014), Grimm *et al.* (2016, Chapter 12), and Liu *et al.* (2021a), respectively. Note that the reduced versions of the three models belong to Type II of nonlinear longitudinal models, as their factor loading matrices depend on population-level coefficients  $b$ ,  $c$ , or  $\gamma$  in addition to measurement times. The coefficient  $b$ ,  $c$ , or  $\gamma$  is estimated as an additional parameter in the corresponding parsimonious model. The package **nlpsem** provides a function `getLGCM()` that enables the estimation of the three intrinsically nonlinear LGCMs, the corresponding reduced models, and LGCMs with linear or quadratic functional forms.

It is important to note that the bilinear spline function with an unknown knot is a piecewise function that must be unified when implemented in SEM software. Multiple reparameterization techniques are available to unify the expression pre- and post-knot (i.e.,  $\gamma_i$ , or  $\gamma$  in the reduced model). More technical details can be found in Harring *et al.* (2006), Grimm *et al.* (2016, Chapter 11), and Liu *et al.* (2021a). The function `getLGCM()` implements the reparameterization approach developed in Liu *et al.* (2021a), wherein the initial status, first, and second slopes are reparameterized as the measurement at the knot ( $\eta'_0$ ), the mean of the two slopes ( $\eta'_1$ ), and the half-difference of the two slopes ( $\eta'_2$ ). The function `getLGCM()` also allows for the transformation of the reparameterized growth factors  $\eta'_0$ ,  $\eta'_1$ , and  $\eta'_2$  to their original forms to obtain coefficients that are interpretable and related to the developmental theory by implementing the (inverse) transformation functions and matrices developed by Liu *et al.* (2021a).

## 2.2. Latent Change Score Models

The primary focus of employing LGCMs is to characterize time-dependent status. However, when exploring nonlinear longitudinal processes, one might also be interested in assessing the growth rate (i.e., rate-of-change) (Grimm *et al.* 2013b; Grimm, Castro-Schilo, and Davoudzadeh 2013a; Zhang *et al.* 2012) and the cumulative value of the growth rate over time (Liu and Perera 2023a). In such cases, one needs to turn to LCSMs, which emphasize time-dependent changes. LCSMs, also known as latent difference score models (McArdle 2001; McArdle and Hamagami 2001; McArdle 2009), were developed to incorporate difference



equations with discrete measurement occasions into the SEM framework. [Liu \(2022b, 2021\)](#); [Liu and Perera \(2023a\)](#) proposed a novel specification for LCSMs that allows for unequal study waves and individual measurement occasions around each wave through the ‘definition variables’ approach. These three works form the theoretical foundation for the computational tool of LCSMs in **nlpsem**. The model specification of LCSMs begins with the concept of classical test theory

$$y_{ij} = y_{ij}^* + \epsilon_{ij}, \quad (4)$$

where  $y_{ij}$ ,  $y_{ij}^*$ , and  $\epsilon_{ij}$  represent the observed score, latent true score, and residual for individual  $i$  at time  $j$ , respectively. This specification suggests that the observed score for an individual at a specific occasion can be decomposed into a latent true score and a residual. At baseline (i.e.,  $j = 1$ ), the true score corresponds to the growth factor indicating the initial status. For each post-baseline (i.e.,  $j \geq 2$ ), the true score at time  $j$  is a linear combination of the score at the previous time point  $j - 1$  and the amount of true change that occurs between time  $j - 1$  and time  $j$

$$y_{ij}^* = \begin{cases} \eta_{0i}, & \text{if } j = 1 \\ y_{i(j-1)}^* + \delta y_{ij}, & \text{if } j = 2, \dots, J \end{cases}, \quad (5)$$

where  $\delta y_{ij}$  is the amount of change that occurs during the  $(j - 1)^{th}$  time interval (i.e., from  $j - 1$  to  $j$ ) for the  $i^{th}$  individual. Such interval-specific changes can be further expressed as the product of the corresponding interval-specific slopes and the time interval. [Liu and Perera \(2023a\)](#) pointed out two ways to express the interval-specific slopes, depending on research interests when exploring a longitudinal process. If the interest is only in assessing the growth rate over time, one may view a process with  $J$  measurements as a linear piecewise function with  $J - 1$  segments (i.e., a nonparametric functional form)

$$\delta y_{ij} = dy_{ij} \times (t_{ij} - t_{i(j-1)}) \quad (j = 2, \dots, J), \quad (6)$$

$$dy_{ij} = \eta_{1i} \times \gamma_{j-1} \quad (j = 2, \dots, J), \quad (7)$$

where  $dy_{ij}$  is the slope during the  $(j - 1)^{th}$  time interval, which can be further expressed as the product of the shape factor indicating the slope in the first time interval and the corresponding relative rate. Suppose one also wants to capture other features of change patterns, such as growth acceleration or growth capacity, and resorts to a parametric nonlinear functional form, such as quadratic, negative exponential, and Jenss-Bayley functions. In this scenario, the slope within an interval is not constant. [Liu and Perera \(2023a\)](#) and [Liu \(2022b\)](#) proposed approximating the interval-specific changes as the product of the instantaneous slope midway through the corresponding interval and the interval length

$$\delta y_{ij} \approx dy_{ij\_mid} \times (t_{ij} - t_{i(j-1)}), \quad (8)$$

in which  $dy_{ij\_mid}$  is the instantaneous slope at the midpoint of the  $(j - 1)^{th}$  time interval (i.e., from time  $(j - 1)$  to time  $J$ ). Table 3 provides the expression of  $dy_{ij}$  for the LCSM with the linear piecewise function and  $dy_{ij\_mid}$  for the LCSMs with quadratic, negative exponential, and Jenss-Bayley functional forms. Note that the LCSMs also allow for intrinsically nonlinear functional forms. In particular, [Grimm et al. \(2013b\)](#) and [Liu \(2022b\)](#) first proposed the LCSMs for the negative exponential function with a random ratio of growth rate and the Jenss-

Bayley function with a random ratio of growth acceleration, respectively. The package **nlpsem** provides the function `getLCSM()` to allow for the implementation of these two intrinsically nonlinear LCSMs, the corresponding reduced model with a fixed ratio of growth rate or growth acceleration, and the LCSMs with linear piecewise and quadratic functional forms.

=====  
 Insert Table 3 about here  
 =====

Latent Change Score Models (LCSMs) can also be expressed in matrix form, with a general expression analogous to Equation 1. Depending on the scenario, they can be further represented as Equations 2 and 3 for cases without and with TICs, respectively. Commonly, the distribution assumptions for LCSMs parallel those made for LGCMs, as introduced in Subsection 2.1. Table 3 outlines the growth factors and corresponding factor loadings for each model, as provided by the `getLCSM()` function. The interpretation of growth factors for each parametric LCSM remains consistent with those of the corresponding LGCM. Notably, the first element in vector  $\boldsymbol{\eta}_i$  symbolizes the latent variable representing the initial status, while the remaining elements compose the growth rate. Unlike the corresponding LGCM, the factor loadings of an LCSM are expressed differently since an LCSM is constructed from the perspective of accumulated change. Specifically, the factor-loading matrices for LCSMs are contingent upon factors related to interval-specific growth rates and time intervals. These time intervals function as ‘definition variables’ in LCSMs with unstructured time frames.

One advantage of employing LCSMs is that this modeling framework facilitates the evaluation of interval-specific slopes, interval-specific changes, and the amounts of change from baseline. Two approaches exist for performing such evaluations. The first method involves deriving the expressions for the means and variances of interval-specific slopes, interval-specific changes, and the amounts of change from baseline, treating them as non-estimable parameters. Table 3 provides these expressions, and the function `getLCSM()` is capable of generating estimates for these parameters. As an alternative, when applicable, interval-specific slopes, interval-specific changes, and the amounts of change from baseline can be defined as additional latent variables, allowing for the calculation of factor scores in a post-fit manner.

### 2.3. Longitudinal Models with Time-varying Covariates

A fundamental interest in evaluating longitudinal processes involves assessing the impact of a covariate on between-individual differences in within-individual change. As introduced in subsections 2.1 and 2.2, time-invariant covariates (TICs) can be incorporated when analyzing longitudinal processes with LGCMs or LCSMs, allowing them to account for variability in growth factors, and then growth curves. However, covariates in longitudinal studies are not necessarily TICs. Grimm (2007) proposed using LGCMs with a time-varying covariate (TVC) to simultaneously analyze bivariate longitudinal variables, wherein the primary process is treated as the longitudinal outcome, and the other variable is considered as the TVC. Table 4 presents the model specification details. The advantages and drawbacks of this specification have been documented in recent research, such as Grimm *et al.* (2016, Chapter 8), Liu (2022a), and Liu and Perera (2023b). To address some limitations of the model proposed by Grimm (2007), Liu (2022a) developed three methods to decompose a TVC into an initial trait and a set of temporal states, enabling separate evaluation of the baseline and temporal effects of

the TVC on outcome trajectories. All three decomposition methods are based on the LCSM specification with a piecewise linear function introduced in Subsection 2.2. Table 4 provides technical details for the three TVC decomposition methods and their corresponding model specifications.

=====  
 Insert Table 4 about here  
 =====

It is important to note that the initial trait remains consistent across all three decomposition methods, represented by the true score of the baseline value, which signifies the initial status of the TVC (i.e.,  $\eta_{0i}^{[x]}$  in Table 4). In contrast, the temporal states differ across the methods. In particular, interval-specific slopes (i.e.,  $\mathbf{dx}_i$  in Table 4, a  $J \times 1$  vector), interval-specific changes (i.e.,  $\mathbf{\delta x}_i$  in Table 4, a  $J \times 1$  vector), and change-from-baseline amounts (i.e.,  $\mathbf{\Delta x}_i$  in Table 4, a  $J \times 1$  vector) characterize each approach, respectively. Regressing the growth factors of the longitudinal outcome on the initial trait allows for the evaluation of the baseline effect (i.e.,  $\beta_{\text{TVC}}$  in Table 4). Regressing each post-baseline observed measurement of the longitudinal outcome on the corresponding temporal state enables the assessment of the temporal effect (i.e.,  $\kappa$  in Table 4). Liu (2022a) provides the technical details for the three decomposition approaches, along with Grimm (2007), serves as the theoretical foundation for the `getTVCmodel()` function, which enables the implementation of LGCMs in Subsection 2.1 and parametric LCSMs in Subsection 2.2 using the four possible ways to include a TVC.

## 2.4. Parallel Processes and Correlated Growth Models

Alternatively, all processes under investigation can be viewed as longitudinal outcomes and analyzed using multivariate growth models (MGMs) (Grimm *et al.* 2016, Chapter 8), also known as parallel process or correlated growth models (McArdle 1988; Grimm 2007). MGMs allow for the examination of covariances among cross-process growth factors, thereby capturing the relationships among multiple longitudinal processes. A general model specification for a bivariate growth model can be expressed as

$$\begin{pmatrix} \mathbf{y}_i \\ \mathbf{z}_i \end{pmatrix} = \begin{pmatrix} \mathbf{\Lambda}_i^{[y]} & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_i^{[z]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[y]} \\ \boldsymbol{\eta}_i^{[z]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[y]} \\ \boldsymbol{\epsilon}_i^{[z]} \end{pmatrix}, \quad (9)$$

where  $\mathbf{z}_i$  is a vector of repeated measurements for the second longitudinal outcome of individual  $i$ . In applications,  $\mathbf{y}_i$  and  $\mathbf{z}_i$  often assume the same functional forms when using MGMs; however, the time structures of  $\mathbf{y}_i$  and  $\mathbf{z}_i$  may differ. The growth factors  $\begin{pmatrix} \boldsymbol{\eta}_i^{[y]} & \boldsymbol{\eta}_i^{[z]} \end{pmatrix}^T$  can be further expressed as deviations from the corresponding mean values

$$\begin{pmatrix} \boldsymbol{\eta}_i^{[y]} \\ \boldsymbol{\eta}_i^{[z]} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\mu}_{\boldsymbol{\eta}}^{[y]} \\ \boldsymbol{\mu}_{\boldsymbol{\eta}}^{[z]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\zeta}_i^{[y]} \\ \boldsymbol{\zeta}_i^{[z]} \end{pmatrix}, \quad (10)$$

where  $\boldsymbol{\mu}_{\boldsymbol{\eta}}^{[y]}$  and  $\boldsymbol{\mu}_{\boldsymbol{\eta}}^{[z]}$  are  $K \times 1$  vectors of growth factor mean values, while  $\boldsymbol{\zeta}_i^{[y]}$  and  $\boldsymbol{\zeta}_i^{[z]}$  are  $K \times 1$  vectors of individual  $i$  deviations from factor means.

In practice,  $\begin{pmatrix} \zeta_i^{[y]} & \zeta_i^{[z]} \end{pmatrix}^T$  is often assumed to follow a multivariate normal distribution

$$\begin{pmatrix} \zeta_i^{[y]} \\ \zeta_i^{[z]} \end{pmatrix} \sim \text{MVN}\left(\mathbf{0}, \begin{pmatrix} \Psi_{\eta}^{[y]} & \Psi_{\eta}^{[yz]} \\ \Psi_{\eta}^{[yz]} & \Psi_{\eta}^{[z]} \end{pmatrix}\right), \quad (11)$$

where  $\Psi_{\eta}^{[y]}$  and  $\Psi_{\eta}^{[z]}$  are  $K \times K$  variance-covariance matrices of outcome-specific growth factors, while  $\Psi_{\eta}^{[yz]}$ , also a  $K \times K$  matrix, indicates the covariances between cross-outcome growth factors. Assuming both  $\mathbf{y}_i$  and  $\mathbf{z}_i$  take linear functions,  $\Psi_{\eta}^{[yz]}$  allows for the estimation of intercept-intercept and slope-slope covariance (correlation) and the covariance between intercept and slope across outcomes. To simplify the model specification, the individual outcome-specific residuals,  $\zeta_i^{[y]}$  and  $\zeta_i^{[z]}$ , are assumed to follow identical and independent normal distributions over time, and the residual covariances are assumed to be homogeneous over time. That is,

$$\begin{pmatrix} \epsilon_i^{[y]} \\ \epsilon_i^{[z]} \end{pmatrix} \sim \text{MVN}\left(\mathbf{0}, \begin{pmatrix} \theta_{\epsilon}^{[y]} \mathbf{I} & \theta_{\epsilon}^{[yz]} \mathbf{I} \\ \theta_{\epsilon}^{[yz]} \mathbf{I} & \theta_{\epsilon}^{[z]} \mathbf{I} \end{pmatrix}\right), \quad (12)$$

where  $\epsilon_i^{[y]}$  and  $\epsilon_i^{[z]}$  are residual variances of the longitudinal outcomes  $\mathbf{y}_i$  and  $\mathbf{z}_i$ , respectively,  $\epsilon_i^{[yz]}$  is the residual covariance between the two processes, and  $\mathbf{I}$  is a  $J \times J$  identity matrix if both  $\mathbf{y}_i$  and  $\mathbf{z}_i$  have  $J$  repeated measurements.

Earlier studies have proposed and developed MGMs using LGCMs or LCSMs to describe univariate longitudinal processes. For instance, a multivariate version of an intrinsically nonlinear LGCM, the bilinear spline growth curve model with an unknown knot, and its reduced model were developed and examined in recent work (Liu and Perera 2021). Moreover, Blozis (2004) created MGMs with quadratic and negative exponential functions (the reduced version). Ferrer and McArdle (2003) suggested employing multivariate LCSMs to analyze multivariate nonlinear developmental processes, while Liu (2021) developed a multivariate version of the LCSM with a piecewise linear functional form. However, some earlier works were limited to MGMs with structured time frames. The current work extends these previous models. Specifically, the package `nlpsem` provides the `getMGM()` function, which implements a multivariate version of each univariate longitudinal model defined in Tables 2 and 3, within the framework of individual measurement occasions. Additionally, `getMGM()` accommodates multiple longitudinal outcomes with varying time structures, tackling a prevalent challenge in analyzing multivariate longitudinal processes by accounting for diverse assessment schedules across outcomes. In clinical trials, for instance, multiple longitudinal outcomes might be collected during specific site visits, with each outcome assessed at different frequencies.

## 2.5. Longitudinal Mediation Models

The third type of statistical model employed to investigate multivariate longitudinal processes is longitudinal mediation models. These models were developed to assess two potential pathways through which the predictor variable affects the outcome variable: paths that lead from the predictor to the outcome, both directly and indirectly through a mediator, enabling the simultaneous evaluation of direct effects and indirect impacts on the outcome variable of the predictor (Baron and Kenny 1986). Since causal relationships often require time to unfold and

causal effects may change over time, longitudinal data are preferred when testing mediation hypotheses. Previous studies, such as Hayes (2009); MacKinnon, Krull, and Lockwood (2000); MacKinnon, Lockwood, Hoffman, West, and Sheets (2002); Cheung and Lau (2008); Shrout and Bolger (2002); Selig and Preacher (2009); Gollob and Reichardt (1987); Cole and Maxwell (2003); Maxwell and Cole (2007); MacKinnon (2008); Cheong, MacKinnon, and Khoo (2003); Soest and Hagtvet (2011), have detailed the drawbacks of using cross-sectional data and the advantages of employing longitudinal data for mediational analyses. Similar to MGMs, longitudinal mediation models can assess each univariate process. However, unlike MGMs that evaluate the covariance (correlation) between cross-outcome growth factors, longitudinal mediation models estimate regression coefficients of the unidirectional paths between these cross-outcome growth factors.

Cheong *et al.* (2003) proposed analyzing longitudinal mediation processes within the LGCM framework and developed a parallel linear growth model to examine how the baseline predictor affects the change in the outcome through the change in the mediator. Soest and Hagtvet (2011) extended this model by incorporating additional regression paths. Moreover, MacKinnon (2008, Chapter 8) noted that the baseline predictor could also be longitudinal. Recently, Liu and Perera (2022a) developed two parallel bilinear growth models to investigate mediational relationships among multiple nonlinear processes. In the proposed models, the linear-linear piecewise function with an unknown knot is used to capture the underlying change patterns of each longitudinal process, where the slopes of the two linear segments represent the short- and long-term growth rate. Table 5 presents the specifications of two parallel linear growth models and two parallel bilinear growth models with all possible paths within the framework of individual measurement occasions. Since the measurement at the knot conveys time-dependent effects and is more meaningful than the baseline value in a longitudinal mediation model, longitudinal mediation models with linear-linear functional forms employ an alternative reparameterization technique, proposed by Grimm *et al.* (2016, Chapter 11), to unify the pre- and post-knot expressions. Specifically, the initial status and two segment-specific slopes are reparameterized as the minimum value between 0 and  $t_{ij} - \gamma$ , the measurement at the knot, and the maximum value between 0 and  $t_{ij} - \gamma$ . The function `getMediation()` facilitates the implementation of these four longitudinal mediation models with all possible paths within the framework of individual measurement occasions. In addition to estimating the direct effects of the predictor on the outcome variable, the function `getMediation()` enables the evaluation of indirect effects on the outcome through the mediator, as well as the total effect. This comprehensive assessment captures the relationships between predictor, mediator, and outcome variables over time, providing a more nuanced understanding of the underlying processes.

=====  
 Insert Table 5 about here  
 =====

## 2.6. Multiple-group Models for Longitudinal Processes

Subsections 2.1 and 2.2 introduce LGCMs and LCSMs with time-invariant covariates (TICs). When TICs are categorical variables, these models assist in evaluating differences in average growth trajectories between groups but have limitations in examining differences in other

aspects of longitudinal processes, such as variability in growth curves or differences in multivariate longitudinal processes between such manifest categorical TICs. This section introduces the multiple-group modeling framework as an alternative method to assess group differences in any aspect of processes, providing additional insights into how and why individuals differ in their development. With a specification of the corresponding model for each manifest group, it is straightforward to extend a one-group model to the multiple-group modeling framework. The function `getMGroup()` in **nlpsem** allows for the implementation of a multiple-group model with each of the models introduced in Subsections 2.1-2.5 as the group-specific model (or sub-model), under the assumption that the change pattern and model type are invariant across groups. A general expression for a multiple-group model with  $G$  manifest groups (where  $G$  is a finite number) can be expressed as

$$p(\text{sub-model}|mc_i = g) = \sum_{g=1}^G p(mc_i = g) \times p(\text{sub-model}|mc_i = g), \quad (13)$$

where  $mc_i$  represents the manifest class label for the  $i^{th}$  individual, and  $p()$  denotes the proportion of each manifest group, summing up to 1 across groups.

## 2.7. Mixture Models for Longitudinal Processes

Mixture models for analyzing longitudinal processes, often referred to as growth mixture models (GMMs) within the SEM framework, bear similarities to multiple-group models. Similar to multiple-group models, GMMs relax the single-population assumption that all individuals are homogeneous and can be captured by a single profile of all models introduced in Subsections 2.1-2.5, mixing these groups through linear combinations. However, unlike multiple-group models where group information is a manifested indicator, GMMs introduced in this section relax the assumption and combine individuals from  $G$  heterogeneous latent subpopulations (where  $G$  is a finite number). In statistical terms, GMMs can also be viewed as a cluster analysis technique for longitudinal processes. Contrasting traditional methods for cluster analysis, such as K-means, GMMs employ a probability-based clustering approach, allowing each individual to belong to multiple latent classes with varying probabilities; an individual is then classified into the class with the highest probability. The vector of these probabilities follows a multinomial distribution that may be further regressed on the TICs suggesting group components. For scenarios without such TICs, a general specification of a GMM with  $G$  latent classes is given by

$$p(\text{sub-model}|lc_i = g) = \sum_{g=1}^G \pi(lc_i = g) \times p(\text{sub-model}|lc_i = g), \quad (14)$$

where  $lc_i$  is the latent class label for the  $i^{th}$  individual, and  $\pi()$  is the latent mixing component variable dividing the sample into  $K$  latent classes. Note that  $\pi()$  needs to satisfy two constraints: (1)  $0 \leq \pi() \leq 1$ , and (2)  $\sum_{g=1}^G \pi() = 1$ .

For scenarios with TICs informing class formation, a general specification of a GMM with  $G$



latent classes is

$$p(\text{sub-model}|lc_i = g, \mathbf{x}_{gi}) = \sum_{g=1}^G \pi(lc_i = g|\mathbf{x}_{gi}) \times p(\text{sub-model}|lc_i = g), \quad (15)$$

$$\pi(lc_i = g|\mathbf{x}_{gi}) = \begin{cases} \frac{1}{1 + \sum_{g=2}^G \exp(\beta_{g0}^{(g)} + \beta_g^{(g)T} \mathbf{x}_{gi})} & \text{reference Group } (g = 1) \\ \frac{\exp(\beta_{g0}^{(g)} + \beta_g^{(g)T} \mathbf{x}_{gi})}{1 + \sum_{g=2}^G \exp(\beta_{g0}^{(g)} + \beta_g^{(g)T} \mathbf{x}_{gi})} & \text{other Groups } (g = 2, \dots, G) \end{cases}, \quad (16)$$

where  $\mathbf{x}_{gi}$  is the TIC vector of the  $i^{th}$  individual,  $\beta_0^{(g)}$ , and  $\beta^{(g)}$  are the intercept and coefficients of the multinomial logistic functions in the  $g^{th}$  latent class. Numerous earlier works, such as Muthén and Shedden (1999); Bouveyron, Celeux, Murphy, and Raftery (2019); Bauer and Curran (2003); Muthén (2004); Grimm and Ram (2009); Grimm, Ram, and Estabrook (2010); Liu, Perera, Kang, Sabo, and Kirkpatrick (2021b); Liu and Perera (2022c,b, 2023b), have developed GMMs with various sub-models (i.e., within-class models) and examined their pros and cons. The majority of developed GMMs are for assessing the heterogeneity in a univariate longitudinal process. However, some recent studies have also developed GMMs to explore the heterogeneity in a multivariate longitudinal process (Liu and Perera 2022b, 2023b). The package **nlpsem** enables the examination of heterogeneity for univariate and multivariate longitudinal processes. In particular, it provides the function `getGMM()` to allow for the implementation of a GMM with each of the models introduced in Subsections 2.1-2.5 as the submodel.

### 3. Model Estimation

The **nlpsem** package is developed by interfacing with the **OpenMx** package (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). This section briefly introduces the iterative algorithm implemented in **OpenMx**. Iterative estimation algorithms require initial values for the model parameters. Given a specified model and initialized parameters, the package calculates the expected model-implied mean vector  $\boldsymbol{\mu}_i$  and variance-covariance structure  $\boldsymbol{\Sigma}_i$ . The expectation is then compared to the raw data to determine the goodness of fit for the model. This comparison is based on the difference between the model-implied mean vector and variance-covariance structure and the observed data distribution, also known as the function value. **OpenMx** (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020) supports multiple fit functions; however, the models implemented in **nlpsem** exclusively use the maximum likelihood (also known as full information maximum likelihood, or FIML, among SEM researchers). The FIML technique enables the specification of a separate likelihood calculation for each individual, accounting for individual heterogeneity in the likelihood contributions. This feature also allows for the analysis of dropout data under the missing at random assumption by leveraging available information.

The function value of the FIML technique is  $-2 \ln L$ , where  $L$  is the likelihood function across all  $N$  individuals. This article denotes the parameter space as  $\Theta$ . For a single-group analysis (i.e., the models introduced in Subsections 2.1-2.5), assuming there are  $p$  variables measured

for the  $i^{th}$  individual in a vector  $\boldsymbol{\omega}_i$ ,  $L$  is defined as follows

$$L(\Theta) = \prod_{i=1}^N \frac{\exp\left(-\frac{1}{2}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i)^T \boldsymbol{\Sigma}_i^{-1}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i)\right)}{\sqrt{(2\pi)^p |\boldsymbol{\Sigma}_i|}}, \quad (17)$$

where  $\boldsymbol{\mu}_i$  and  $\boldsymbol{\Sigma}_i$  are the model-implied mean vector and variance-covariance structure of the  $i^{th}$  individual. For a multiple-group model with  $G$  manifested groups (i.e., the model introduced in Subsection 2.6), the likelihood function across all individuals is defined as

$$L(\Theta) = \prod_{i=1}^N \left( \sum_{g=1}^G p(mc_i = g) \times \frac{\exp\left(-\frac{1}{2}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})^T \boldsymbol{\Sigma}_i^{(g)-1}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})\right)}{\sqrt{(2\pi)^p |\boldsymbol{\Sigma}_i^{(g)}|}} \right), \quad (18)$$

where  $\boldsymbol{\mu}_i^{(g)}$  and  $\boldsymbol{\Sigma}_i^{(g)}$  are the submodel-implied mean vector and variance-covariance structure of the  $i^{th}$  individual in the  $g^{th}$  manifested group. For a growth mixture model with  $G$  latent classes (i.e., the models introduced in Subsection 2.7), the likelihood function across all  $N$  individuals is defined as

$$L(\Theta) = \prod_{i=1}^N \left( \sum_{g=1}^G \pi(lc_i = g) \times \frac{\exp\left(-\frac{1}{2}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})^T \boldsymbol{\Sigma}_i^{(g)-1}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})\right)}{\sqrt{(2\pi)^p |\boldsymbol{\Sigma}_i^{(g)}|}} \right) \quad (19)$$

and

$$L(\Theta) = \prod_{i=1}^N \left( \sum_{g=1}^G \pi(lc_i = g | \mathbf{x}_{gi}) \times \frac{\exp\left(-\frac{1}{2}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})^T \boldsymbol{\Sigma}_i^{(g)-1}(\boldsymbol{\omega}_i - \boldsymbol{\mu}_i^{(g)})\right)}{\sqrt{(2\pi)^p |\boldsymbol{\Sigma}_i^{(g)}|}} \right) \quad (20)$$

for scenarios without and with the TICs informing cluster formation (i.e.,  $\mathbf{x}_{gi}$ ), in which  $\boldsymbol{\mu}_i^{(g)}$  and  $\boldsymbol{\Sigma}_i^{(g)}$  are the submodel-implied mean vector and variance-covariance structure of the  $i^{th}$  individual in the  $g^{th}$  latent class. Optimizers work iteratively to minimize the FIML function value, continuing until convergence is achieved, which is indicated by the status code 0 in **OpenMx** (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). **OpenMx** provides three optimization engines: NPSOL (Gill *et al.* 1986), SLSQP (Johnson 2014; Kraft 1994), and CSOLNP (Zahery *et al.* 2017). All these engines are compatible with the **nlpsem** package. However, we particularly advocate for CSOLNP as the primary optimizer for estimation functions within **nlpsem**. This recommendation stems from CSOLNP's proven efficiency in managing non-linearly constrained optimization challenges. Its ability to deliver quicker outcomes, coupled with its efficient memory usage, positions CSOLNP as a reliable and effective choice for **nlpsem**'s computational demands. Nonetheless, an exception is noted by Pritikin, Rappaport, and Neale (2017): they suggest employing SLSQP when generating likelihood-based confidence intervals (see Section 5.1 for details). This is because both NPSOL and CSOLNP encounter challenges with constrained optimization, and likelihood-based confidence intervals appear optimally formulated when using SLSQP optimizer for constrained optimization.

## 4. Implementation

This package currently includes seven estimation functions:

1. `getLGCM()` for latent growth curve models (LGCMs), without or with time-invariant covariates (TICs),
2. `getLCSM()` for latent change score models (LCSMs), without or with TICs,
3. `getTVModel()` for LGCMs or LCSMs with a time-varying covariate (TVC),
4. `getMGM()` for multivariate version of LGCMs or LCSMs,
5. `getMediation()` for longitudinal mediation analysis,
6. `getMGroup()` for multiple-group version of the models (1)-(5), and
7. `getMIX()` for mixture model version of the models (1)-(5).

The optimization for the seven estimation functions relies on the built-in optimizers of the package **OpenMx**. This section describes the calls of these estimation functions, presents the process of having additional but non-estimable parameters, and details the initialization of the initiative optimizers.

#### 4.1. Function `getLGCM()`

The call of `getLGCM()` is

```
getLGCM(dat, t_var, y_var, curveFun, intrinsic = TRUE, records, growth_TIC =
NULL, starts = NULL, res_scale = NULL, tries = NULL, OKStatus = 0, jitterD =
"runif", loc = 1, scale = 0.25, paramOut = FALSE, names = NULL).
```

The `dat` argument represents the input dataset, which must be in a wide format (i.e., one row per individual) to satisfy the requirements of the **OpenMx** package (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). This dataset typically comprises repeated measurements, corresponding measurement occasions, and time-invariant covariates (TICs) that account for the variability of growth factors, if present. The `y_var` and `t_var` arguments denote the prefixes of the column names for the measurement value and time at each point, respectively. Together with `records`, these arguments delineate the column name of the longitudinal outcome and the measurement occasions over time. For instance, if the dataset contains four repeated measurements ( $Y_1, Y_2, Y_3$ , and  $Y_4$ ) recorded at  $T_1, T_2, T_3$ , and  $T_4$ , the user needs to specify `t_var = "T"`, `y_var = "Y"`, `records = 1:4`. The `curveFun` argument designates the functional form, chosen from any function listed in Table 2, to define the change patterns of the repeated measurements. The `intrinsic` argument is a boolean flag that indicates whether an intrinsically nonlinear longitudinal model (i.e., the third type of nonlinear longitudinal model in the current package) is constructed. The choice of `curveFun` and `intrinsic` often depends on the trajectory shapes of raw data and the research questions of interest.

The `growth_TIC` argument specifies the TICs added to the model, if any, to account for the variability of growth factors. By default, it is set to `NULL`, indicating that no growth TICs are incorporated into the fitted LGCM. The `starts` argument establishes the set of initial values for the iterative algorithms. If set to the default value of `NULL`, the estimation function will derive a set of initial values based on the raw data. Alternatively, the `getLGCM()` function allows for user-specified initial values. When `starts = NULL`, the `res_scale` argument is

required, as it defines the proportion of variance unexplained by the growth factors and aids in determining the initial values of the residual variance  $\theta_\epsilon$ .

The `tries` argument establishes the number of additional attempts to fit a specified model in **OpenMx** (often referred to as a `mxModel` object). With its default value of `NULL` (indicating zero additional trials unless the initial attempt), the `getLGCM()` function will call the `mxRun()` function to send a `mxModel` object to an optimizer and return the free parameters with their optimized values. Assigning a numerical value to `tries` permits the corresponding additional attempts to fit a model until the optimizer produces an acceptable solution defined by the `OKStatus` argument or reaches the maximum number of attempts. When assigning a numeric value to `tries`, the user must also define the `jitterD`, `loc`, `scale`, and `OKStatus` arguments. The `jitterD` argument, in conjunction with `loc` and `scale`, defines the distribution and its location and scale parameters, from which a set of random values are drawn to perturb initial values between attempts. The `jitterD` argument is a character string, with supported values including "runif" (uniform distribution), "rnorm" (normal distribution), or "rcauchy" (Cauchy distribution). The `loc` and `scale` arguments are numeric inputs. By default, the setup employs a uniform distribution with a location parameter of 1 and a scale parameter of 0.25, adhering to the default settings in the **OpenMx** package (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). The `OKStatus` argument is an integer vector that defines the acceptable solution of the optimizer status code, with its default value being 0, indicating a successful optimization without error returns. The `paramOut` argument specifies whether the point estimates and standard errors of the free and additional non-estimable parameters will be extracted from the optimized `mxModel`. Parameter names provided by the `names` argument must be supplied if `paramOut = TRUE`.

## 4.2. Function `getLCSM()`

The call of `getLCSM()` is

```
getLCSM(dat, t_var, y_var, curveFun, intrinsic = TRUE, records, growth_TIC =
NULL, starts = NULL, res_scale = NULL, tries = NULL, OKStatus = 0, jitterD =
"runif", loc = 1, scale = 0.25, paramOut = FALSE, names = NULL).
```

The arguments in `getLCSM()` mirror those in `getLGCM()`, with a notable distinction in the `curveFun` argument, which supports different functional forms for `getLCSM()`: nonparametric, quadratic, negative exponential, and Jenss-Bayley curves.

## 4.3. Function `getTVCmodel()`

The call of `getTVCmodel()` is

```
getTVCmodel(dat, t_var, y_var, curveFun, intrinsic = TRUE, records, y_model,
TVC, decompose, growth_TIC = NULL, starts = NULL, res_scale = NULL, res_cor
= NULL, tries = NULL, OKStatus = 0, jitterD = "runif", loc = 1, scale = 0.25,
paramOut = FALSE, names = NULL).
```

In comparison to the `getLGCM()` and `getLCSM()` functions, the `getTVCmodel()` function incorporates four additional arguments: `y_model`, `TVC`, `decompose`, and `res_cor`. Specifically, the `y_model` argument establishes the model describing longitudinal outcome change patterns, providing options 'LGCM' and 'LCSM'. For example, if `y_model = "LGCM"` is specified, a LGCM will be fit to capture these change patterns. The `TVC` and `records` arguments together

define the column names for the TVC over time. For example, to specify four repeated TVC measurements,  $x_1$ ,  $x_2$ ,  $x_3$ , and  $x_4$ , one should set `TVC = "x"`. The `decompose` argument, an integer, indicates the decomposing option, with supported values of 0 (adding a TVC directly), 1 (adding a decomposed TVC with interval-specified slopes), 2 (adding a decomposed TVC with interval-specified changes), and 3 (adding a decomposed TVC with change from baseline values). If `starts = NULL`, `res_scale` must be specified, and its configuration is contingent upon the modeling context. For models where the longitudinal outcome directly regresses on the TVC, `res_scale` should be a single numeric value, indicating the initial proportion of variance in the longitudinal outcome that the growth factors do not capture. Conversely, when the longitudinal outcome is regressed on a decomposed TVC, `res_scale` requires a two-element numeric vector, detailing the unexplained variance proportions for longitudinal outcome and the TVC. In this latter scenario, `res_cor` must also be specified to denote the initial residual correlation between the two longitudinal variables.

#### 4.4. Function `getMGM()`

The call of `getMGM()` is

```
getMGM(dat, t_var, y_var, curveFun, intrinsic = TRUE, records, y_model, starts = NULL, res_scale = NULL, res_cor = NULL, tries = NULL, OKStatus = 0, jitterD = "runif", loc = 1, scale = 0.25, paramOut = FALSE, names = NULL).
```

In the `getMGM()` function, parameters are akin to those found in the three preceding functions but have been adapted to accommodate multiple longitudinal outcomes. In contrast to the scenario of univariate longitudinal processes where `y_var` (`t_var`) and `records` are each represented by a single character string or a numeric vector, the analysis of multivariate longitudinal processes necessitates their specification as a vector of character strings and a list of numeric vectors, respectively. The length of `y_var`, `t_var`, and `records` must be identical, as it reflects the number of longitudinal processes being analyzed simultaneously within the multivariate model. Each element in `y_var`, in conjunction with the relevant index vector from the `records` argument, defines the repeated measurements for the associated longitudinal outcome. Similarly, each `t_var` element, when paired with the corresponding index vector from the `records` argument, establishes the measurement occasions for the respective longitudinal outcome. This configuration enables the `getMGM()` function to effectively analyze multiple longitudinal processes simultaneously while retaining flexibility in model specification. Suppose the input data contains two longitudinal processes: (1)  $Y_1$ ,  $Y_2$ ,  $Y_4$ , and  $Y_5$  recorded at  $T_1$ ,  $T_2$ ,  $T_4$ , and  $T_5$ , and (2)  $Z_1$ ,  $Z_3$ ,  $Z_4$ , and  $Z_5$  recorded at  $T_1$ ,  $T_3$ ,  $T_4$ , and  $T_5$ . To analyze this bivariate longitudinal process using all available data, one needs to set `y_var = c("Y", "Z")`, `t_var = c("T", "T")`, and `records = list(c(1:2, 4:5), c(1, 3:5))`. The `res_scale` argument in the `getMGM()` function should maintain a length identical to that of `y_var`, `t_var`, and `records`, as it represents a numeric vector where each value delineates the proportion of unexplained variance in the corresponding longitudinal process attributable to the growth factors to derive initial values of the corresponding residual variance. In addition, `res_cor` provides the initial value(s) of the residual correlation(s) of any two longitudinal processes under investigation.

#### 4.5. Function `getMediation()`

The call of `getMediation()` is

```
getMediation(dat, t_var, y_var, m_var, x_type, x_var, curveFun, records, starts
= NULL, res_scale = NULL, res_cor = NULL, tries = NULL, OKStatus = 0, jitterD =
"runif", loc = 1, scale = 0.25, paramOut = FALSE, names = NULL).
```

In the `getMediation()` function, the `y_var` and `m_var` arguments, in conjunction with `records`, determine the column names for the longitudinal outcome and the longitudinal mediator, respectively. This function accommodates both baseline and longitudinal predictors, specified by `x_type = "baseline"` and `x_type = "longitudinal"`. For a longitudinal mediation model with a baseline predictor, a two-element character vector is required for the `t_var` argument, while a list containing two numeric vectors is needed for the `records` argument, specifying the measurement times for the outcome and mediator. In a model with a longitudinal predictor, an additional numeric vector for the `records` argument is necessary, allowing the `x_var` argument and the third numeric vector to define the column names of the predictors. The `t_var` argument then requires an extra element, along with the third numeric vector of the `records` argument, to delineate the column names of measurement occasions of the longitudinal predictor. Similarly, the `res_scale` argument should contain two elements for a model with a baseline predictor and three elements for a model with a longitudinal predictor.

#### 4.6. Function `getMGroup()`

The call of `getMGroup()` is

```
getMGroup(dat, grp_var, sub_Model, t_var, records, y_var, curveFun, intrinsic
= NULL, y_model = NULL, m_var = NULL, x_type = NULL, x_var = NULL, TVC = NULL,
decompose = NULL, growth_TIC = NULL, starts = NULL, res_scale = NULL, res_cor
= NULL, tries = NULL, OKStatus = 0, jitterD = "runif", loc = 1, scale = 0.25,
paramOut = FALSE, names = NULL).
```

The `getMGroup()` function necessitates the inclusion of two supplementary arguments, `grp_var` and `sub_Model`. The `grp_var` argument specifies the column name representing the manifest group. This enables the determination of the number of manifest groups present in the model. The `sub_Model` argument allows for the selection of a single-group model (i.e., LGCMs, LCSMs, TVC models, MGMs, or longitudinal mediation models) to be incorporated as a submodel. The remaining arguments are comparable to those detailed in Subsections 4.1-4.5. To address group-specific specifications, the `getMGroup()` function requires that arguments associated with initial value derivation (i.e., `starts`, `res_scale`, and `res_cor`) be structured as a list of inputs corresponding to the relevant single-group model.

#### 4.7. Function `getMIX()`

The call of `getMIX()` is

```
getMIX(dat, prop_starts, sub_Model, cluster_TIC = NULL, t_var, y_var, curveFun,
intrinsic = NULL, records, y_model = NULL, m_var = NULL, x_var = NULL, x_type =
NULL, TVC = NULL, decompose = NULL, growth_TIC = NULL, starts = NULL, res_scale
= NULL, res_cor = NULL, tries = NULL, OKStatus = 0, jitterD = "runif", loc = 1,
scale = 0.25, paramOut = FALSE, names = NULL).
```

The implementation of `getMIX()` calls for three additional arguments: `prop_starts`, `sub_Model`, and `cluster_TIC`. The `prop_starts` argument specifies the initial proportions for each latent



class in the mixture model. The length of this argument indicates the number of latent classes and the sum of these elements totaling 1. The `sub_Model` argument allows users to select a single-group model (i.e., LGCMs, LCSMs, TVC models, MGMs, or longitudinal mediation models) for integration as a submodel within the mixture model. The `cluster_TIC` argument designates the TICs, if any, to be included in the model to inform cluster formation. By default, it is set to `NULL`, indicating that no cluster TICs are incorporated in the fitted mixture model. Analogous to the `getMGroup()` function, the remaining arguments are consistent with those outlined in Subsections 4.1-4.5. The `getMIX()` function necessitates that arguments about initial value derivation (i.e., `starts`, `res_scale`, and `res_cor`) be formatted as a list of inputs corresponding to the respective single-group model to accommodate class-specific specifications.

#### 4.8. Initial Values

Iterative estimation algorithms necessitate initialization using a set of initial values for the parameter space  $\Theta$ . Selecting suitable initial values improves the likelihood of model convergence and reduces computational burdens. In each estimation function, the `starts` argument dictates the algorithm's initialization process. The estimation functions provided in the **nlpsem** package allow for user-specified initial values or derive them from raw data. As the longitudinal models in **nlpsem** are centered around the concept of 'growth factors' that are latent variables, the most crucial step in deriving initial values involves calculating these growth factors for each individual from raw data. The derivation algorithm may differ based on the target being a single-group, multiple-group, or mixture model. For single-group models, 'raw' growth factors of each longitudinal process are derived by fitting a linear regression using `lm()` for linear functions or fitting a nonlinear regression with `nls()` for nonlinear functions for each individual.

Subsequent steps depend on the model specification. For models without covariates or mediators, estimation functions compute the mean values and variance-covariance matrix of these 'raw' growth factors. When evaluating the impacts of TICs, estimation functions further initialize path coefficients that quantify such impacts by regressing 'raw' growth factors on the TICs. In cases assessing the effect of a TVC, the longitudinal outcome is further regressed on the corresponding TVC measurement. When examining the effect of a decomposed TVC, baseline and temporal effects are separately initialized by regressing the 'raw' growth factors on the initial trait and regressing the longitudinal outcome on the corresponding temporal state. In longitudinal mediation models, path coefficients' initialization occurs stepwise. First, the mediator's 'raw' growth factors are regressed on predictor-related parameters, followed by regressing the outcome's 'raw' growth factors on both predictor and mediator-related parameters. Direct initialization of residual variances and covariances from raw data is unfeasible, so estimation functions necessitate user-provided information via the `res_scale` and `res_cor` arguments.

For multiple-group models, the procedure above is executed for each manifested group. In mixture models, the K-means algorithm is initially performed with a pre-specified number of latent classes, assigning individuals to clusters. The procedure is then conducted for each latent class using the assigned labels.

#### 4.9. Additional Non-estimable Parameters

The **nlpsem** package streamlines the estimation of non-estimable parameters by utilizing the `mxAlgebra()` and `mxSE()` functions within the **OpenMx** package. Specifically, `mxAlgebra()` outlines the derivation of additional parameters from free parameters, computing point estimates, while `mxSE()` generates the corresponding standard errors based on the (multivariate) delta method (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). The seven estimation functions introduced earlier accommodate four categories of non-estimable parameters. Firstly, as discussed in Subsection 2.1, when analyzing a bilinear spline growth curve model with an unknown knot, a reparameterization process is essential to unify the pre- and post-knot expressions. However, such reparameterized parameters are not directly tied to developmental theory and consequently lack meaningful interpretations. The **nlpsem** package integrates the inverse transformation functions and matrices developed in Liu *et al.* (2021a) and Liu and Perera (2021) to derive parameters of the LGCM and its multivariate version with the bilinear spline functional form in their original form.

Second, as introduced in Subsection 2.2, one advantage of employing the LCSM is that it enables the estimation of means and variances of interval-specific slopes, interval-specific changes, and change-from-baseline values for a longitudinal outcome. These means and variances can also be considered additional parameters. Thirdly, the package incorporates parameters of conditional distributions for models involving covariates (and a mediator). For example, in an LGCM with a TIC, the estimated parameters related to the longitudinal outcome are the vector of growth factor intercepts and the unexplained variance-covariance structure of the growth factors, as indicated in Equation 3. In practice, the mean vector of growth factors conditional on the TIC holds greater interest since these coefficients directly pertain to developmental theory. Therefore, **nlpsem** facilitates the derivation of parameters for such conditional distributions. Finally, the fourth category of additional parameters includes the indirect and total effects of the predictor on the outcome generated by longitudinal mediation models.

## 5. Post-fit Computations

The **nlpsem** package provides a comprehensive set of post-fit computations, analyses, and evaluations applicable to all seven estimation functions. However, certain analyses, such as posterior classification, the enumeration process, and the computation of latent kappa statistics (Dumenci 2011; Dumenci, Perera, Keefe, Ang, Slover, Jensen, and Riddle 2019), are specifically tailored to mixture models. This section presents an overview of these post-fit analyses and their applications.

### 5.1. Statistical Significance: Wald p-values and Confidence Intervals

It is essential to obtain p-values and confidence intervals to quantify the uncertainty of estimates. Similar to the **OpenMx** package, the **nlpsem** package produces point estimates and standard errors for each estimable or derived parameter by default. Additionally, **nlpsem** provides the `getEstimateStats()` function, allowing for the evaluation of Wald p-values (Wald 1943) and confidence intervals. While generating p-values is not the primary focus of **OpenMx**, the `getEstimateStats()` function provides an algorithm to derive the z-score and associated p-value for each parameter based on the corresponding point estimate and standard error, with the assumption that the sampling distribution of the estimate is approximately normal.

Furthermore, `getEstimateStats()` allows for three types of confidence intervals: Wald confidence intervals (Casella and Berger 2002), likelihood-based confidence intervals (also known as ‘likelihood profile confidence intervals’) (Madansky 1965; Matthews 1988), and bootstrap confidence intervals (Efron and Tibshirani 1994, Chapter 12). Wald confidence intervals require the assumption of asymptotic normal distribution, while likelihood-based and bootstrap confidence intervals relax this assumption. Specifically, likelihood-based confidence intervals rely on the asymptotic chi-square distribution of the log-likelihood ratio test statistic, and bootstrap confidence intervals depend on resampling distributions. The `getEstimateStats()` function produces likelihood-based and bootstrap confidence intervals using **OpenMx** built-in functions, `mxCI()` and `mxBootstrap()`, respectively (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020). Both functions require an optimized `mxModel` object as input, seamlessly integrating with the **nlpsem** package to provide comprehensive statistical information for users.

The call of `getEstimateStats()` is

```
getEstimateStats(model = NULL, est_in, p_values = TRUE, CI = TRUE, CI_type =
"Wald", rep = NA, conf.level = 0.95).
```

The `getEstimateStats()` function takes several arguments, including `model`, an optimized `mxModel` object; `est_in`, a data frame comprising point estimates and standard errors of parameters for which users aim to derive p-values and/or confidence intervals; `p_values`, a logical value denoting whether to generate p-values (default is `TRUE`); `CI`, a logical value denoting whether to generate confidence intervals (default is `TRUE`); `CI_type`, a character string indicating the type of confidence interval (default is `"Wald"`); `conf.level`, a numeric value representing the desired probability for the confidence interval (default is `0.95`); and `rep`, a numeric value specifying the number of resampling instances when constructing bootstrap CIs (default is `NA`).

By default, the function generates p-values and Wald confidence intervals for the parameters enumerated in the `est_in` argument without requiring an input of the `model` argument. The `CI_type` argument enables users to designate alternative confidence intervals, such as likelihood-based intervals (`CI_type = "likelihood"`) or bootstrap intervals (`CI_type = "bootstrap"`). When specifying these two types of confidence intervals, the `model` argument is mandatory, and the `getEstimateStats()` function generates the corresponding confidence intervals for all free parameters. Additionally, `rep` must be specified when constructing bootstrap CIs.

## 5.2. Model Selection between Intrinsically Nonlinear Longitudinal Models and Their Parsimonious Alternatives

As detailed in Section 2, fitting intrinsically nonlinear longitudinal models within the SEM framework necessitates linearizing functional forms through Taylor series expansion. These linearizations provide valuable insights, especially when estimating individual growth rate parameters, individual growth acceleration parameters, or individual knots in the context of developmental theory. However, it is essential to balance the benefits of these intricate models against the interpretability and feasibility of their simpler counterparts. Specifically, before delving into the parameters of such complex models, researchers must ascertain if a more straightforward version—with fixed growth rates, growth accelerations, or knots—serves as an adequate alternative. Often, this decision is influenced by both the research’s focus and

the available statistical evidence. To aid in this decision-making, the **nlpsem** package provides tools grounded in statistical principles. One can compare an intrinsically nonlinear longitudinal model with its parsimonious version using the likelihood ratio test (LRT), given that the simpler model is nested within the more complex one. Our `getLRT()` function, leveraging the capabilities of the `mxCompare()` function in **OpenMx**, facilitates these comparisons. What sets `mxCompare()` apart from other R functions like `lrtest` from **lmtest** is its ability to conduct bootstrap likelihood ratio tests. This feature is particularly valuable as it enhances the p-value's accuracy, especially when the standard assumptions of the LRT might be challenged (Feng and McCulloch 1996).

The call of `getLRT()` is

```
getLRT(full, reduced, boot = FALSE, rep = NA).
```

The `getLRT()` function is tailored to compare a full model (e.g., an intrinsically nonlinear longitudinal model) with its parsimonious counterpart and accepts the following arguments: `full`, the `mxModel` object of the full model; `reduced`, the `mxModel` object of the reduced model; `boot`, a logical value indicating whether to use the bootstrap resampling distribution to compute the p-value; and `rep`, an integer specifying the number of replications employed to approximate the bootstrap distribution. By default, the function does not utilize the bootstrap distribution to compute the p-value, but users can enable this by setting `boot = TRUE`. When employing the bootstrap distribution, the `rep` argument governs the number of replications for approximating the distribution, with a default value of `NA`.

### 5.3. Derivation of Individual Factor Scores

As previously introduced, one of the primary goals in analyzing longitudinal data is to explore between-individual differences in within-individual changes. In the context of longitudinal processes within the SEM framework, these between-individual differences are represented by the variance of growth factors. For instance, using the point estimate and associated standard error of a growth factor variable, a researcher may perform a Wald test to determine whether the variability is significantly greater than 0. If it is found to be significant, a natural follow-up question arises as to whether a value for each individual can be derived. Such individual values in the SEM framework are referred to as factor scores, indicating an individual's relative position or standing on a latent variable. The `getIndFS()` function, built upon the **OpenMx** function `mxFactorScores()` (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020), facilitates the estimation of factor scores for all specified latent variables. Note that this function is capable of calculating factor scores for both free latent variables (e.g., growth factors) and derived latent variables (e.g., interval-specific slopes, interval-specific changes, and change from baseline). The function `getIndFS()` generates factor scores and their standard errors for each latent variable.

The call of `getIndFS()` is

```
getIndFS(model, FS_type = "Regression").
```

The `getIndFS()` function is designed to calculate or estimate factor scores, along with their associated standard errors, for an optimized model obtained from the **nlpsem** package. This function provides three options for the type of factor scores to compute by interfacing with the `mxFactorScores()` function in the **OpenMx** package, including `ML` (maximum likelihood), `WeightedML` (weighted maximum likelihood), and `Regression` (Neale *et al.* 2016; Pritikin *et al.* 2015; Hunter 2018; Boker *et al.* 2020).

The **Regression** method, which is the default option of the `getIndFS()` function, generates factor scores using a general linear prediction formula that predicts the factor scores from the non-missing manifest variables for each row of the raw data. In contrast, both **ML** and **WeightedML** methods create a new model for each data row and estimate factor scores as free parameters in a model with a single data row. The **ML** method optimizes the conditional likelihood of the data given the factor scores, while the **WeightedML** method optimizes the joint likelihood of the data and factor scores, taking the latent variable distribution into account during their estimation. More technical details regarding the pros and cons of each method of obtaining factor scores are well-documented in earlier studies, such as [Priestley and Subba Rao \(1975\)](#); [Estabrook and Neale \(2013\)](#).

#### 5.4. Posterior Classification

An essential post-fit evaluation of a mixture model involving latent classes is a posterior classification to compute a vector of probabilities, which helps determine the probability of each individual belonging to each latent class. This posterior classification is grounded in the Bayes rule. Assuming a mixture model does not contain a time-invariant covariate (TIC) that indicates class components, the posterior probabilities can be calculated as follows

$$p(lc_i = g) = \frac{\pi(lc_i = g) \times p(\text{sub-model}|lc_i = g)}{\sum_{g=1}^G \pi(lc_i = g) \times p(\text{sub-model}|lc_i = g)}. \quad (21)$$

On the other hand, if a mixture model contains TICs that indicate class components, the posterior probabilities can be computed as

$$p(lc_i = g) = \frac{\pi(lc_i = g|\mathbf{x}_{gi}) \times p(\text{sub-model}|lc_i = g)}{\sum_{g=1}^G \pi(lc_i = g|\mathbf{x}_{gi}) \times p(\text{sub-model}|lc_i = g)}. \quad (22)$$

The call of `getPosterior()` is

```
getPosterior(model, nClass, label = FALSE, cluster_TIC = NULL).
```

The `getPosterior()` function facilitates the calculation of posterior probabilities for each individual's membership in latent classes of an optimized mixture model with **OpenMx**. This function accepts four arguments: `model`, an optimized mixture model; `nClass`, representing the number of latent classes specified for the model; `label`, a boolean flag indicating whether the dataset contains true labels (default is `FALSE`); and `cluster_TIC`, specifying the column index of time-invariant covariates that inform class formation in the observed data, when applicable (default is `NULL`, signifying that no TICs indicating class components are included in the mixture model).

Upon execution, the function returns an object comprising the following slots: (1) a matrix of posterior probabilities for each individual, (2) a vector of cluster memberships, (3) the entropy of the input model, and (4) the accuracy value of the input model if the true labels are available.

#### 5.5. Model Selection and Enumeration Process with Summary Tables

The **nlpsem** package provides the `getSummary()` function to facilitate model comparison and selection based on estimated likelihood and information criteria such as Akaike Information

Criterion (AIC) and Bayesian Information Criterion (BIC). This function is an alternative to the likelihood ratio test provided by the `getLRT()` function, particularly when the models under comparison are not nested. The `getSummary()` function streamlines the model selection and comparison process across various scenarios.

The `getSummary()` function is capable of generating a summary table for any optimized models fitted by the estimation functions provided by `nlpsem`. However, it is essential to ensure that comparisons are fair and meaningful by comparing models with only one distinct aspect. For instance, comparing a series of LGCMs without any covariates but with different functional forms is appropriate, as it helps in selecting the optimal functional form<sup>3</sup>. In this context, the summary table generated by `getSummary()` presents a comprehensive overview of each model's performance, including the number of free parameters, estimated likelihood, AIC, BIC, and residual variances. Such information allows researchers to make informed decisions, for example, the most suitable functional form for their data from the statistical perspective.

Another practical application of `getSummary()` is in the enumeration process. Within the SEM literature, the enumeration process for mixture models is typically performed in an exploratory fashion. This process involves fitting a series of candidate models, including a single-group model and the corresponding mixture models with different numbers of latent classes. The 'best' model and the optimal number of latent classes are then determined using statistical criteria such as the BIC (Nylund, Asparouhov, and Muthén 2007). Following the SEM literature convention, it is recommended to conduct the enumeration process without including any covariates to prevent potential confounding effects (Diallo, Morin, and Lu 2017; Nylund-Gibson and Masyn 2016). For multivariate longitudinal outcomes, the enumeration process can be conducted either collectively across all outcomes or individually for each outcome. When considering multivariate longitudinal outcomes collectively in the enumeration process, models with fewer latent classes tend to be selected, as the BIC penalizes model complexity. On the other hand, executing the enumeration process independently for each outcome allows for the identification of the optimal number of latent classes specific to each longitudinal trajectory. Ultimately, the decision regarding the number of latent classes for a multivariate longitudinal model should be empirically based (Liu and Perera 2022b).

The call of `getSummary()` is

```
getSummary(model_list, HetModels = FALSE).
```

The `getSummary()` function is developed for facilitating the process of model comparison and selection, taking in two arguments: `model_list` and `HetModels`. The `model_list` argument is expected to be a list of optimized models from `nlpsem`. These models should be comparable for meaningful comparisons, exhibiting differences in only one aspect. The `HetModels` argument, a logical value, denotes whether a model within the `model_list` incorporates heterogeneous classes. When `HetModels` is set to `TRUE`, the function outputs a summary table that accentuates the enumeration process, which includes a comparison of single-group models and respective mixture models with varying quantities of latent classes.

## 5.6. Visualization of Estimated Growth Curves and Growth Changes Over Time

The `nlpsem` package provides `getFigure()` function to facilitate the visualization of (class-

---

<sup>3</sup>Note that comparing an LGCM with a linear functional form and TICs to an LGCM with a quadratic functional form but no TICs is not meaningful, as the models differ in more than one aspect.



specific) estimated growth status and changes over time.

The call of `getFigure()` is

```
getFigure(model, nClass = NULL, cluster_TIC = NULL, grp_var = NULL, sub_Model,
y_var, curveFun, y_model = NULL, t_var, records, m_var = NULL, x_type = NULL,
x_var = NULL, xstarts, xlab = "Time", outcome = "Process").
```

The `getFigure()` function incorporates multiple arguments the same as those utilized in the estimation functions. For instance, the `sub_Model` argument allows users to select a (class-specific) model. Note that `getFigure()` generates estimated growth curves when `sub_model = "LGCM"` or `y_model = "LGCM"`. Conversely, it provides estimated growth rates and changes from baseline when `sub_model = "LCSM"` or `y_model = "LCSM"`.

In addition to the arguments shared with the estimation functions, `getFigure()` also includes three supplementary arguments: `xstarts`, `xlab`, and `outcome`. The `xstarts` argument signifies the onset of the longitudinal study, enabling the provision of interpretable estimates. On the other hand, `xlab` and `outcome` are used for labeling the generated figures. To elaborate, `xlab` represents the unit of the time structure, while `outcome` denotes the name(s) of the longitudinal processes. For example, should the data record the time unit in months, then `xlab` should be assigned as `xlab = "Month"`. If the model scrutinizes two longitudinal processes, such as the advancement of reading and math abilities, the `outcome` argument should be configured as `outcome = c("Reading Ability", "Math Ability")`. This ensures that the labels on the generated figures are both pertinent, thereby aiding their interpretation.

## 5.7. Latent Kappa Statistic

The latent kappa statistic (Dumenci 2011) is a measure designed to quantify the agreement between latent groups. While the kappa statistic is commonly used to assess inter-rater reliability for manifested categorical variables, the latent kappa statistic focuses on evaluating agreement for latent categorical variables. Recent research has extended this metric to assess the agreement between latent class assignments across various clustering algorithms. Such applications include evaluating the agreement between cluster assignments based on different longitudinal outcomes (Dumenci *et al.* 2019; Liu and Perera 2022b), assessing the impact of TICs on latent class assignments (Liu and Perera 2022c), and examining the effects of considering multiple longitudinal outcomes in the clustering algorithm on clustering results (Liu and Perera 2022b). The `nlpsem` package provides the `getLatentKappa()` function to facilitate comparisons between clustering algorithms. This function enables researchers to assess the consistency of latent class assignments, providing valuable insights into the robustness of clustering results and facilitating a more accurate evaluation and interpretation of heterogeneity among individuals.

The call of `getLatentKappa()` is

```
getLatentKappa(label1, label2, conf.level = 0.95).
```

The `getLatentKappa()` function accepts three arguments: `label1`, `label2`, and `conf.level`. The `label1` and `label2` arguments represent the latent class assignments determined by the first and second clustering algorithms, respectively. Meanwhile, the `conf.level` argument indicates the probability for confidence intervals associated with the kappa statistic.

## 6. Tutorial: Synthetic Examples

### 6.1. Example Data

The **nlpsem** package incorporates a sample dataset, **RMS\_dat**, obtained from the publicly accessible segment of the Early Childhood Longitudinal Study, Kindergarten Class of 2010-11 (ECLS-K:2011). This study, conducted by the National Center for Education Statistics (NCES), spanned nine rounds of data collection from the fall of 2010 to the spring of 2016. The ECLS-K:2011 dataset provides comprehensive information about children's early life experiences, encompassing aspects such as health, developmental milestones, educational progress, and experiences leading up to kindergarten<sup>4</sup>.

The **RMS\_dat** dataset consists of 500 observations spanning 49 variables, thereby capturing a broad spectrum of factors pertinent to early childhood education research. These include variables related to academic performance, such as reading, mathematics, and science scores across all study waves, as well as demographic and environmental variables like age-in-month at each study wave, sex, race, family income, and others. Additionally, the dataset incorporates teacher evaluations of children's behavioral and learning traits, such as approach to learning, self-control, interpersonal skills, and attention focus. The dataset underwent careful preprocessing for computation and interpretation purposes. We conducted three main preprocessing procedures. First, we set the initial time point to zero across all time-varying measurements. This manipulation enables the initial status estimates to be interpreted as the values corresponding to the onset of the study. We implemented this in R, modifying each measurement occasion (T1 through T9) by subtracting the initial time point (T1). Second, we standardized the time-invariant covariates (TICs) to expedite computational processes. For this, we employed the `scale()` function in R, which centers these variables around a mean of zero and scales them to have a standard deviation of one. In addition, we standardized the time-varying covariate (TVC), in this case, the reading scores at each time point (R1 through R9). Following the approach outlined by [Liu \(2022a\)](#); [Liu and Perera \(2023b\)](#), we calculated the mean and variance of the baseline reading scores (R1) and used these baseline parameters to standardize the reading scores across all time points.

```
R> library(nlpsem)
R> # Load ECLS-K (2011) data
R> data("RMS_dat")
R> RMS_dat0 <- RMS_dat
R> # Re-baseline the data so that the estimated initial status is for the
R> # starting point of the study
R> baseT <- RMS_dat0$T1; xstarts <- mean(baseT)
R> RMS_dat0$T1 <- RMS_dat0$T1 - baseT
R> RMS_dat0$T2 <- RMS_dat0$T2 - baseT
R> RMS_dat0$T3 <- RMS_dat0$T3 - baseT
R> RMS_dat0$T4 <- RMS_dat0$T4 - baseT
R> RMS_dat0$T5 <- RMS_dat0$T5 - baseT
```

---

<sup>4</sup>The dataset supplied in this package has been formatted only for demonstration purposes and does not include the original survey weights. Thus, researchers intending to perform in-depth analyses should resort to the entire dataset on the NCES data products page and ensure the appropriate use of complex survey weights.

```

R> RMS_dat0$T6 <- RMS_dat0$T6 - baseT
R> RMS_dat0$T7 <- RMS_dat0$T7 - baseT
R> RMS_dat0$T8 <- RMS_dat0$T8 - baseT
R> RMS_dat0$T9 <- RMS_dat0$T9 - baseT
R> # Standardize time-invariant covariates (TICs)
R> ## ex1 and ex2 are standardized growth TICs in models
R> RMS_dat0$ex1 <- scale(RMS_dat0$Approach_to_Learning)
R> RMS_dat0$ex2 <- scale(RMS_dat0$Attention_focus)
R> ## gx1 and gx2 are standardized cluster TICs in models
R> RMS_dat0$gx1 <- scale(RMS_dat0$INCOME)
R> RMS_dat0$gx2 <- scale(RMS_dat0$EDU)
R> # Standardize time-varying covariate (TVC)
R> BL_mean <- mean(RMS_dat0[, "R1"])
R> BL_var <- var(RMS_dat0[, "R1"])
R> RMS_dat0$Rs1 <- (RMS_dat0$R1 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs2 <- (RMS_dat0$R2 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs3 <- (RMS_dat0$R3 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs4 <- (RMS_dat0$R4 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs5 <- (RMS_dat0$R5 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs6 <- (RMS_dat0$R6 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs7 <- (RMS_dat0$R7 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs8 <- (RMS_dat0$R8 - BL_mean)/sqrt(BL_var)
R> RMS_dat0$Rs9 <- (RMS_dat0$R9 - BL_mean)/sqrt(BL_var)
R> head(RMS_dat0)

```

##	ID	R1	R2	R3	R4	R5	R6	R7						
## 14	10000014	61.0533	71.0552	78.5028	109.9882	112.6338	119.0805	131.0204						
## 29	10000029	58.7342	63.3107	87.4390	108.3098	117.8031	124.4843	123.0577						
## 47	10000047	51.5844	66.6677	67.5425	81.5954	92.1793	104.7755	112.0417						
## 66	10000066	53.6920	61.6393	74.8572	91.1721	105.4468	122.3648	126.2234						
## 95	10000095	54.8889	73.2573	77.7501	107.0961	113.7871	108.4502	127.0300						
## 129	10000129	91.2245	107.1388	120.3719	136.5753	137.1338	139.9153	140.1828						
##		R8	R9	M1	M2	M3	M4	M5	M6					
## 14	143.4052	137.0652	51.2953	50.6399	57.3925	80.1414	85.2682	94.1731						
## 29	126.5784	137.9527	44.1411	54.3557	64.7107	82.9958	83.2294	106.2275						
## 47	121.5341	140.2891	30.9925	44.7628	47.0755	57.7788	66.8194	78.5502						
## 66	141.0226	148.8337	24.6459	47.0937	56.6954	69.5438	76.1157	86.6673						
## 95	132.5852	136.9246	29.6065	50.2221	61.0513	81.7005	77.4089	95.1059						
## 129	152.3430	155.9560	60.4633	74.5698	82.6556	83.0306	110.9810	119.8762						
##		M7	M8	M9	S2	S3	S4	S5	S6	S7				
## 14	106.1280	115.0474	125.8766	24.6497	30.4830	32.3496	40.9454	46.5182	65.5331					
## 29	123.4263	132.7925	127.5096	35.8322	43.7028	40.6455	47.7815	53.1821	60.4607					
## 47	94.3809	99.7010	115.3470	41.2768	46.7668	42.0820	43.3976	55.4341	73.3641					
## 66	122.5060	118.6628	131.0051	43.2319	53.8406	59.3788	56.0984	61.3869	77.8254					
## 95	109.7579	118.7727	120.9763	43.4005	46.8256	50.2495	47.4000	55.3191	68.1581					
## 129	119.4172	122.1821	136.5360	47.8725	63.9814	66.4430	68.9873	73.0444	78.6502					
##		S8	S9	T1	T2	T3	T4	T5	T6	T7	T8	T9	SEX	RACE

```

## 14 69.0916 75.9290 0 4.08 10.36 18.12 23.91 29.69 39.03 54.05 65.07 1 1
## 29 77.6913 68.3381 0 5.76 11.54 17.46 23.28 29.92 41.00 53.92 65.89 2 1
## 47 80.3503 82.6842 0 6.02 11.77 17.73 23.77 29.73 42.51 53.53 64.57 1 1
## 66 79.7523 86.9595 0 6.68 11.94 18.12 23.94 30.51 43.53 54.61 67.50 1 3
## 95 58.9861 82.4376 0 5.29 11.74 17.95 23.90 30.31 42.41 53.43 67.43 2 1
## 129 82.9082 89.5211 0 5.69 11.94 17.62 24.07 29.62 41.66 53.56 65.56 2 8
##      LOCALE INCOME SCHOOL_TYPE Approach_to_Learning Self_control Interpersonal
## 14      2      17              1              3.1667      3.6667      3.2
## 29      1       8              1              2.8571      3.0000      3.2
## 47      1      17              1              2.5714      3.5000      4.0
## 66      1      18              2              2.2857      3.0000      3.2
## 95      2      16              1              3.5714      3.5000      3.4
## 129     4      18              1              3.7143      3.3333      3.2
##      External_prob_Behavior Internal_prob_Behavior Attention_focus
## 14              1.0              1.00      5.5000
## 29              2.0              1.00      6.0000
## 47              1.6              1.00      4.5000
## 66              1.6              1.25      4.8333
## 95              1.0              1.00      5.5000
## 129             1.0              1.00      7.0000
##      Inhibitory_Ctrl EDU      ex1      ex2      gx1      gx2
## 14      6.8000 5 0.1827911 0.51616548 0.9808159 -0.1505668
## 29      4.3333 5 -0.3144371 0.92312846 -0.7372345 -0.1505668
## 47      2.6667 5 -0.7732811 -0.29776049 0.9808159 -0.1505668
## 66      3.5000 6 -1.2321251 -0.02647897 1.1717104 0.3513226
## 95      6.0000 6 0.8327532 0.51616548 0.7899214 0.3513226
## 129     6.0000 8 1.0622555 1.73705443 1.1717104 1.3551014
##      Rs1      Rs2      Rs3      Rs4      Rs5      Rs6      Rs7      Rs8
## 14 0.48312698 1.3656884 2.022860 4.801112 5.034558 5.603411 6.656981 7.749808
## 29 0.27849104 0.6823186 2.811385 4.653011 5.490694 6.080239 5.954357 6.265021
## 47 -0.35240288 0.9785382 1.055730 2.295749 3.229666 4.341147 4.982312 5.819915
## 66 -0.16642957 0.5348353 1.701175 3.140791 4.400382 5.893216 6.233696 7.539568
## 95 -0.06081585 1.5600004 1.956442 4.545915 5.136325 4.665400 6.304870 6.795058
## 129 3.14541497 4.5496829 5.717363 7.147141 7.196423 7.441861 7.465465 8.538474
##      Rs9
## 14 7.190370
## 29 7.268682
## 47 7.474845
## 66 8.228815
## 95 7.177963
## 129 8.857282

```

## 6.2. getLGCM() Examples


The function `getLGCM()` applies latent growth curve models (LGCMs) to illustrate mathematics development through linear-linear trajectories without the inclusion of any covariates.

In this section, we fit two distinct models: intrinsically nonlinear (i.e., a linear-linear function with a random knot) and its parsimonious counterpart (i.e., a linear-linear function with a fixed knot). We performed a likelihood ratio test as part of this demonstration. This statistical comparison allows us to ascertain whether the parsimonious model can serve as an effective substitute for the more complex model without a significant compromise in fit or information conveyance. In addition, we visualized the estimated developmental statuses of mathematics ability for both models (refer to Figure 1a). As indicated in Figure 1b, both LGCMs successfully capture the underlying patterns of mathematics development. However, when utilizing AIC, BIC, or the likelihood ratio test as comparative metrics, the intrinsically nonlinear model appears as a superior fit for the development of mathematics ability.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # LGCM_TICs, Bilinear Spline Functional Form, Random knot
R> Math_LGCM_BLS_f <- getLGCM(
R> + dat = RMS_dat0, t_var = "T", y_var = "M", curveFun = "bilinear spline",
R> + intrinsic = TRUE, records = 1:9, growth_TIC = NULL, res_scale = 0.1
R> + )@mxOutput
R> # LGCM_TICs, Bilinear Spline Functional Form, Fixed knot
R> Math_LGCM_BLS_r <- getLGCM(
R> + dat = RMS_dat0, t_var = "T", y_var = "M", curveFun = "bilinear spline",
R> + intrinsic = FALSE, records = 1:9, growth_TIC = NULL, res_scale = 0.1
R> + )@mxOutput
R> # Perform LRT to compare two models
R> getLRT(full = Math_LGCM_BLS_f, reduced = Math_LGCM_BLS_r, boot = FALSE,
R> + rep = NA)
R> # Figure of Growth Status for LGCM_TICs, Bilinear Spline Functional Form,
R> # Random knot
R> Figure1 <- getFigure(
R> + model = Math_LGCM_BLS_f, nClass = NULL, cluster_TIC = NULL, sub_Model = "LGCM",
R> + y_var = "M", curveFun = "BLS", y_model = "LGCM", t_var = "T", records = 1:9,
R> + m_var = NULL, x_var = NULL, x_type = NULL, xstarts = xstarts, xlab = "Month",
R> + outcome = "Mathematics"
R> + )
R> show(Figure1) ## See Figure 1(a) in the manuscript
R> # Figure of Growth Status for LGCM_TICs, Bilinear Spline Functional Form,
R> # Fixed knot
R> Figure2 <- getFigure(
R> + model = Math_LGCM_BLS_r, nClass = NULL, cluster_TIC = NULL, sub_Model = "LGCM",
R> + y_var = "M", curveFun = "BLS", y_model = "LGCM", t_var = "T", records = 1:9,
R> + m_var = NULL, x_var = NULL, x_type = NULL, xstarts = xstarts, xlab = "Month",
R> + outcome = "Mathematics"
R> + )
R> show(Figure2) ## See Figure 1(b) in the manuscript
```

##	# of Free Param	-2loglik	Degree of Freedom	Diff in loglik
## Full Model	15	31261.60	4485	NA
## Reduced Model	11	31347.39	4489	85.78891

##	Diff in DoF	p.values	AIC	BIC
## Full Model	NA	<NA>	31291.60	31354.82
## Reduced Model	4	<0.0001	31369.39	31415.75



(a) Bilinear Growth Curve (Random Knot)

(b) Bilinear Growth Curve (Fix Knot)

Figure 1: Estimated Development Status of Mathematics Ability from Latent Growth Curve Models

### 6.3. getLCSM() Examples

The function `getLCSM()` demonstrates reading development employing latent change score models (LCSMs) with a nonparametric functional form. In this section, we implement two models: one without any covariates and the other incorporating two standardized growth time-invariant covariates (TICs), specifically the baseline teacher-reported approach to learning and attentional focus. In addition to these model fits, we provide a summary table for both models and compute p-values and confidence intervals for the model incorporating growth TICs. Furthermore, we present visualizations of the estimated change-from-baseline and growth rate for the model without TICs in Figure 2.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # LCSM, Nonparametric Functional Form
R> paraNonP_LCSM <- c(
R> + "mueta0", "mueta1", paste0("psi", c("00", "01", "11")), paste0("rel_rate", 2:8),
R> + "residuals", paste0("slp_val_est", 1:8), paste0("slp_var_est", 1:8),
R> + paste0("chg_inv_val_est", 1:8), paste0("chg_inv_var_est", 1:8),
R> + paste0("chg_bl_val_est", 1:8), paste0("chg_bl_var_est", 1:8)
R> + )
R> Read_LCSM_NonP <- getLCSM(
R> + dat = RMS_dat0, t_var = "T", y_var = "R", curveFun = "nonparametric",
R> + intrinsic = FALSE, records = 1:9, growth_TIC = NULL, res_scale = 0.1,
R> + paramOut = TRUE, names = paraNonP_LCSM
R> + )@mxOutput
```



```

R> # LCSM_TIC, Nonparametric Functional Form
R> paraNonP_LCSM_TIC <- c(
R> + "alpha0", "alpha1", paste0("psi", c("00", "01", "11")), paste0("rel_rate", 2:8),
R> + "residuals", paste0("beta1", c(0:1)), paste0("beta2", c(0:1)),
R> + paste0("mux", 1:2), paste0("phi", c("11", "12", "22")), "mueta0", "mueta1",
R> + paste0("slp_val_est", 1:8), paste0("slp_var_est", 1:8),
R> + paste0("chg_inv_val_est", 1:8), paste0("chg_inv_var_est", 1:8),
R> + paste0("chg_bl_val_est", 1:8), paste0("chg_bl_var_est", 1:8)
R> + )
R> Read_LCSM_NonP_TIC <- getLCSM(
R> + dat = RMS_dat0, t_var = "T", y_var = "R", curveFun = "nonparametric",
R> + intrinsic = FALSE, records = 1:9, growth_TIC = c("ex1", "ex2"), res_scale = 0.1,
R> + paramOut = TRUE, names = paraNonP_LCSM_TIC
R> + )
R> Read_LCSM_NonP_TIC_o <- Read_LCSM_NonP_TIC@mxOutput
R> Read_LCSM_NonP_TIC_e <- Read_LCSM_NonP_TIC@Estimates
R> # Get summary of two LCSMs with Nonparametric Functional Form
R> getSummary(model_list = list(Read_LCSM_NonP, Read_LCSM_NonP_TIC_o))
R> # Figure of LCSM, Nonparametric Functional Form
R> Figure3 <- getFigure(
R> + model = Read_LCSM_NonP, sub_Model = "LCSM", y_var = "R", curveFun = "NonP",
R> + y_model = "LCSM", t_var = "T", records = 1:9, xstarts = xstarts, xlab = "Month",
R> + outcome = "Reading"
R> + )
R> show(Figure3) ## See Figures 2(a) and 2(b) in the manuscript
R> # Three types of confidence intervals of LCSM_TIC, Nonparametric Functional Form
R> mxOption(model = NULL, key = "Default optimizer", "SLSQP", reset = FALSE)
R> set.seed(20181022)
R> getEstimateStats(
R> + model = Read_LCSM_NonP_TIC_o, est_in = Read_LCSM_NonP_TIC_e,
R> + CI_type = "all", rep = 1000
R> + )

```

```

##      Model No_Params      -2ll      AIC      BIC Y_residuals
## 1 Model1          13 32288.39 32314.39 32369.18    45.1379
## 2 Model2          22 34575.48 34619.48 34712.20    45.1411

```

```
## An object of class "StatsOutput"
```

```
## Slot "wald":
```

```

##      Estimate      SE p.value wald_lbound wald_ubound
## alpha0      55.5596 0.6005 <0.0001    54.3826    56.7366
## alpha1       2.3772 0.0724 <0.0001     2.2353     2.5191
## psi00      136.1660 9.8748 <0.0001   116.8117   155.5203
## psi01      -1.1696 0.2349 <0.0001    -1.6300    -0.7092
## psi11       0.1076 0.0113 <0.0001     0.0855     0.1297
## rel_rate2     0.7137 0.0435 <0.0001     0.6284     0.7990
## rel_rate3     1.2059 0.0459 <0.0001     1.1159     1.2959

```


## rel_rate4	0.5228	0.0339	<0.0001	0.4564	0.5892
## rel_rate5	0.6870	0.0339	<0.0001	0.6206	0.7534
## rel_rate6	0.2888	0.0174	<0.0001	0.2547	0.3229
## rel_rate7	0.2923	0.0170	<0.0001	0.2590	0.3256
## rel_rate8	0.2570	0.0166	<0.0001	0.2245	0.2895
## residuals	45.1411	1.0797	<0.0001	43.0249	47.2573
## beta10	2.8588	0.8811	0.0012	1.1319	4.5857
## beta11	-0.0307	0.0273	0.2608	-0.0842	0.0228
## beta20	2.1026	0.8775	0.0166	0.3827	3.8225
## beta21	0.0440	0.0273	0.1070	-0.0095	0.0975
## mux1	0.0000	0.0447	>0.9999	-0.0876	0.0876
## mux2	0.0000	0.0447	>0.9999	-0.0876	0.0876
## phi11	0.9980	0.0631	<0.0001	0.8743	1.1217
## phi12	0.7759	0.0565	<0.0001	0.6652	0.8866
## phi22	0.9980	0.0631	<0.0001	0.8743	1.1217
## mueta0	55.5595	0.6359	<0.0001	54.3132	56.8058
## mueta1	2.3772	0.0724	<0.0001	2.2353	2.5191
## slp_val_est1	2.3772	0.0724	<0.0001	2.2353	2.5191
## slp_val_est2	1.6966	0.0700	<0.0001	1.5594	1.8338
## slp_val_est3	2.8666	0.0684	<0.0001	2.7325	3.0007
## slp_val_est4	1.2427	0.0721	<0.0001	1.1014	1.3840
## slp_val_est5	1.6331	0.0651	<0.0001	1.5055	1.7607
## slp_val_est6	0.6866	0.0362	<0.0001	0.6156	0.7576
## slp_val_est7	0.6950	0.0349	<0.0001	0.6266	0.7634
## slp_val_est8	0.6111	0.0352	<0.0001	0.5421	0.6801
## slp_var_est1	0.1076	0.0113	<0.0001	0.0855	0.1297
## slp_var_est2	0.0548	0.0065	<0.0001	0.0421	0.0675
## slp_var_est3	0.1565	0.0156	<0.0001	0.1259	0.1871
## slp_var_est4	0.0294	0.0042	<0.0001	0.0212	0.0376
## slp_var_est5	0.0508	0.0060	<0.0001	0.0390	0.0626
## slp_var_est6	0.0090	0.0012	<0.0001	0.0066	0.0114
## slp_var_est7	0.0092	0.0012	<0.0001	0.0068	0.0116
## slp_var_est8	0.0071	0.0010	<0.0001	0.0051	0.0091
## chg_inv_val_est1	13.8240	0.4211	<0.0001	12.9987	14.6493
## chg_inv_val_est2	9.9338	0.4097	<0.0001	9.1308	10.7368
## chg_inv_val_est3	17.7667	0.4237	<0.0001	16.9363	18.5971
## chg_inv_val_est4	7.0613	0.4099	<0.0001	6.2579	7.8647
## chg_inv_val_est5	10.4670	0.4172	<0.0001	9.6493	11.2847
## chg_inv_val_est6	7.9308	0.4186	<0.0001	7.1104	8.7512
## chg_inv_val_est7	8.3602	0.4203	<0.0001	7.5364	9.1840
## chg_inv_val_est8	7.3259	0.4215	<0.0001	6.4998	8.1520
## chg_inv_var_est1	3.6401	0.3829	<0.0001	2.8896	4.3906
## chg_inv_var_est2	1.8796	0.2218	<0.0001	1.4449	2.3143
## chg_inv_var_est3	6.0125	0.5980	<0.0001	4.8404	7.1846
## chg_inv_var_est4	0.9498	0.1368	<0.0001	0.6817	1.2179
## chg_inv_var_est5	2.0868	0.2456	<0.0001	1.6054	2.5682
## chg_inv_var_est6	1.1981	0.1623	<0.0001	0.8800	1.5162

```


## chg_inv_var_est7  1.3313  0.1751 <0.0001    0.9881    1.6745
## chg_inv_var_est8  1.0223  0.1471 <0.0001    0.7340    1.3106
## chg_bl_val_est1   13.8240  0.4211 <0.0001   12.9987   14.6493
## chg_bl_val_est2   23.7578  0.4419 <0.0001   22.8917   24.6239
## chg_bl_val_est3   41.5246  0.4908 <0.0001   40.5626   42.4866
## chg_bl_val_est4   48.5858  0.5144 <0.0001   47.5776   49.5940
## chg_bl_val_est5   59.0528  0.5565 <0.0001   57.9621   60.1435
## chg_bl_val_est6   66.9836  0.5901 <0.0001   65.8270   68.1402
## chg_bl_val_est7   75.3438  0.6287 <0.0001   74.1116   76.5760
## chg_bl_val_est8   82.6698  0.6637 <0.0001   81.3690   83.9706
## chg_bl_var_est1    3.6401  0.3829 <0.0001    2.8896    4.3906
## chg_bl_var_est2   10.7512  0.9991 <0.0001    8.7930   12.7094
## chg_bl_var_est3   32.8437  2.9315 <0.0001   27.0981   38.5893
## chg_bl_var_est4   44.9637  3.9800 <0.0001   37.1630   52.7644
## chg_bl_var_est5   66.4238  5.8517 <0.0001   54.9547   77.8929
## chg_bl_var_est6   85.4633  7.4952 <0.0001   70.7730  100.1536
## chg_bl_var_est7  108.1279  9.4500 <0.0001   89.6062  126.6496
## chg_bl_var_est8  130.1774 11.3692 <0.0001  107.8942  152.4606
##
## Slot "likelihood":
##           Estimate lik_lbound lik_ubound
## beta01         2.8588      1.1069      4.6111
## beta11        -0.0307     -0.0847      0.0205
## beta02         2.1026      0.3505      3.8544
## beta12         0.0440     -0.0073      0.0980
## Y_rel_rate2     0.7137      0.6318      0.8034
## Y_rel_rate3     1.2059      1.1198      1.3007
## Y_rel_rate4     0.5228      0.4581      0.5913
## Y_rel_rate5     0.6870      0.6227      0.7562
## Y_rel_rate6     0.2888      0.2555      0.3241
## Y_rel_rate7     0.2923      0.2599      0.3268
## Y_rel_rate8     0.2571      0.2253      0.2905
## Y_residuals    45.1411    43.0795    47.3331
## phi1           0.9980      0.8833      1.1334
## phi2           0.7759      0.6730      0.8963
## phi3           0.9980      0.8833      1.1334
## Y_psi00        136.1660   118.2743   157.2625
## Y_psi01        -1.1696    -1.5725    -0.7773
## Y_psi11         0.1076      0.0876      0.1316
## mux1           0.0000     -0.0877      0.0877
## mux2           0.0000     -0.0877      0.0877
## Y_mueta0        55.5596    54.3749    56.7388
## Y_mueta1         2.3772      2.2348      2.5198
##
## Slot "bootstrap":
##           Estimate boot_lbound boot_ubound
## beta01         2.8588      1.1640      4.4200

```

## beta11	-0.0307	-0.0845	0.0170
## beta02	2.1026	0.4246	3.7234
## beta12	0.0440	-0.0150	0.1022
## Y_rel_rate2	0.7137	0.6476	0.7835
## Y_rel_rate3	1.2059	1.1188	1.2912
## Y_rel_rate4	0.5228	0.4618	0.5840
## Y_rel_rate5	0.6870	0.6259	0.7470
## Y_rel_rate6	0.2888	0.2566	0.3236
## Y_rel_rate7	0.2923	0.2642	0.3214
## Y_rel_rate8	0.2570	0.2304	0.2871
## Y_residuals	45.1411	42.3520	48.0030
## phi1	0.9980	0.8918	1.1049
## phi2	0.7759	0.6696	0.8801
## phi3	0.9980	0.8759	1.1089
## Y_psi00	136.1660	106.7621	166.3167
## Y_psi01	-1.1696	-1.9469	-0.5296
## Y_psi11	0.1076	0.0857	0.1321
## mux1	0.0000	-0.0878	0.0811
## mux2	0.0000	-0.0881	0.0864
## Y_mueta0	55.5596	54.6802	56.5723
## Y_mueta1	2.3772	2.2658	2.4974



(a) Estimated Change from Baseline



(b) Estimated Growth Rate

Figure 2: Latent Change Score Models with Nonparametric Function for Reading Ability

Figures 2a and 2b depict the estimated values for both change-from-baseline and interval-specific growth rates, both of which are derived from the estimated growth factors. It is important to note that the estimated growth factors from the model integrating growth TICs are conditional on these TICs, thus differing from the model excluding growth TICs. Examining the summary table output reveals that Model 1 (the nonparametric LCSM excluding growth TICs) exhibits a smaller estimated likelihood, AIC, and BIC compared to Model 2 (the nonparametric LCSM including two growth TICs). However, this difference does not inherently signify that Model 1 outperforms Model 2, given that the data employed to fit

both models vary. When setting `CI_type = "all"` in the `getEstimateStats()` function, it generates three tables corresponding to Wald, likelihood-based, and bootstrap confidence intervals. Note that the function generates likelihood-based and bootstrap confidence intervals solely for free parameters, while it provides Wald confidence intervals for both free parameters and those that cannot be estimated. These non-estimable parameters include the conditional mean values of the growth factors, the means and variances of interval-specific slopes, interval-specific changes, and the amounts of change from baseline.

#### 6.4. `getTVCmodel()` Examples

The function `getTVCmodel()` is implemented to construct univariate longitudinal outcome models with a time-varying covariate (TVC). It exemplifies the latent growth curve model with an intrinsically linear-linear functional form for mathematics development while simultaneously examining the impact of reading ability on mathematics development over time. This section contains the fitting of two distinct models. Both models view reading ability as the TVC and the baseline teacher-reported approach to learning as the time-invariant covariate (TIC). However, the first model directly incorporates the TVC, while the second introduces a decomposed TVC, partitioned into the baseline value and a set of interval-specific slopes. Alongside these model fittings, we computed p-values and Wald confidence intervals for the model incorporating the decomposed TVC. We provide plots of the estimated growth status of mathematics development for both models, demonstrated in Figures 3a and 3b.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # LGCM with TVC and TICs, Bilinear Spline Functional Form (random knot)
R> set.seed(20191029)
R> Math_TVC_BLS_f <- getTVCmodel(
R> + dat = RMS_dat0, t_var = "T", y_var = "M", curveFun = "BLS", intrinsic = TRUE,
R> + records = 1:9, y_model = "LGCM", TVC = "Rs", decompose = 0, growth_TIC = "ex1",
R> + res_scale = 0.1, tries = 10
R> + )@mxOutput
R> # LGCM with decomposed TVC and TICs, Bilinear Spline Functional Form (random knot)
R> paraBLS_TVC.f <- c(
R> + "Y_alpha0", "Y_alpha1", "Y_alpha2", "Y_alphag",
R> + paste0("Y_psi", c("00", "01", "02", "0g", "11", "12", "1g", "22", "2g", "gg")),
R> + "Y_residuals", "X_mueta0", "X_mueta1", paste0("X_psi", c("00", "01", "11")),
R> + paste0("X_rel_rate", 2:8), paste0("X_abs_rate", 1:8), "X_residuals",
R> + paste0("betaTIC", c(0:2, "g")), paste0("betaTVC", c(0:2, "g")), "muTIC", "phiTIC",
R> + "Y_mueta0", "Y_mueta1", "Y_mueta2", "Y_mu_knot", "covBL", "kappa", "Cov_XYres"
R> + )
R> set.seed(20191029)
R> Math_TVCslp_BLS_f <- getTVCmodel(
R> + dat = RMS_dat0, t_var = "T", y_var = "M", curveFun = "BLS", intrinsic = TRUE,
R> + records = 1:9, y_model = "LGCM", TVC = "Rs", decompose = 1, growth_TIC = "ex1",
R> + res_scale = c(0.1, 0.1), res_cor = 0.3, tries = 10, paramOut = TRUE,
R> + names = paraBLS_TVC.f
R> + )
R> Math_TVCslp_BLS_f_o <- Math_TVCslp_BLS_f@mxOutput
```

```

R> Math_TVCslp_BLS_f_e <- Math_TVCslp_BLS_f@Estimates
R> # Get Wald confidence intervals for LGCM with decomposed TVC and TICs, Bilinear
R> # Spline Functional Form (random knot)
R> getEstimateStats(est_in = Math_TVCslp_BLS_f_e, CI_type = "Wald")
R> # Figure of LGCM with TVC and TICs, Bilinear Spline Functional Form (random knot)
R> Figure4 <- getFigure(
R> + model = Math_TVC_BLS_f, sub_Model = "TVC", y_var = "M", curveFun = "BLS",
R> + y_model = "LGCM", t_var = "T", records = 1:9, xstarts = xstarts, xlab = "Month",
R> + outcome = "Mathematics"
R> + )
R> # Figure of LGCM with decomposed TVC and TICs, Bilinear Spline Functional Form
R> # (random knot)
R> Figure5 <- getFigure(
R> + model = Math_TVCslp_BLS_f_o, sub_Model = "TVC", y_var = "M", curveFun = "BLS",
R> + y_model = "LGCM", t_var = "T", records = 1:9, xstarts = xstarts, xlab = "Month",
R> + outcome = "Mathematics"
R> + )
R> show(Figure4) ## See Figure 3(a) in the manuscript
R> show(Figure5) ## See Figure 3(b) in the manuscript

```

```
## An object of class "StatsOutput"
```

```
## Slot "wald":
```

##	Estimate	SE	p.value	wald_lbound	wald_ubound
## Y_alpha0	37.1157	0.4057	<0.0001	36.3205	37.9109
## Y_alpha1	1.7073	0.0171	<0.0001	1.6738	1.7408
## Y_alpha2	0.6839	0.0157	<0.0001	0.6531	0.7147
## Y_alphag	37.9199	0.5011	<0.0001	36.9378	38.9020
## Y_psi00	47.2134	4.1851	<0.0001	39.0108	55.4160
## Y_psi01	-0.1004	0.1358	0.4597	-0.3666	0.1658
## Y_psi02	-0.1912	0.1246	0.1249	-0.4354	0.0530
## Y_psi0g	-4.4622	3.6339	0.2195	-11.5845	2.6601
## Y_psi11	0.0845	0.0083	<0.0001	0.0682	0.1008
## Y_psi12	-0.0015	0.0055	0.7851	-0.0123	0.0093
## Y_psi1g	-0.7959	0.1853	<0.0001	-1.1591	-0.4327
## Y_psi22	0.0231	0.0079	0.0035	0.0076	0.0386
## Y_psi2g	-0.3249	0.1777	0.0675	-0.6732	0.0234
## Y_psigg	42.3164	6.7900	<0.0001	29.0082	55.6246
## Y_residuals	28.3922	0.8032	<0.0001	26.8180	29.9664
## X_mueta0	0.0093	0.0556	0.8672	-0.0997	0.1183
## X_mueta1	0.2086	0.0058	<0.0001	0.1972	0.2200
## X_psi00	1.2124	0.0857	<0.0001	1.0444	1.3804
## X_psi01	-0.0065	0.0017	0.0001	-0.0098	-0.0032
## X_psi11	0.0007	0.0001	<0.0001	0.0005	0.0009
## X_rel_rate2	0.6947	0.0384	<0.0001	0.6194	0.7700
## X_rel_rate3	1.2479	0.0431	<0.0001	1.1634	1.3324
## X_rel_rate4	0.4903	0.0303	<0.0001	0.4309	0.5497
## X_rel_rate5	0.7110	0.0330	<0.0001	0.6463	0.7757



```

## X_rel_rate6    0.2823 0.0169 <0.0001    0.2492    0.3154
## X_rel_rate7    0.3019 0.0168 <0.0001    0.2690    0.3348
## X_rel_rate8    0.2568 0.0165 <0.0001    0.2245    0.2891
## X_abs_rate1    0.2086 0.0058 <0.0001    0.1972    0.2200
## X_abs_rate2    0.1449 0.0055 <0.0001    0.1341    0.1557
## X_abs_rate3    0.2603 0.0055 <0.0001    0.2495    0.2711
## X_abs_rate4    0.1023 0.0057 <0.0001    0.0911    0.1135
## X_abs_rate5    0.1483 0.0055 <0.0001    0.1375    0.1591
## X_abs_rate6    0.0589 0.0032 <0.0001    0.0526    0.0652
## X_abs_rate7    0.0630 0.0031 <0.0001    0.0569    0.0691
## X_abs_rate8    0.0536 0.0031 <0.0001    0.0475    0.0597
## X_residuals    0.3587 0.0088 <0.0001    0.3415    0.3759
## betaTIC0       0.9665 0.3904 0.0133    0.2013    1.7317
## betaTIC1       0.0099 0.0169 0.5580   -0.0232    0.0430
## betaTIC2      -0.0175 0.0165 0.2889   -0.0498    0.0148
## betaTICg       0.0593 0.4690 0.8994   -0.8599    0.9785
## betaTVC0       7.6231 0.3796 <0.0001    6.8791    8.3671
## betaTVC1       0.0708 0.0168 <0.0001    0.0379    0.1037
## betaTVC2      -0.0289 0.0158 0.0674   -0.0599    0.0021
## betaTVCg      -1.5263 0.4644 0.0010   -2.4365   -0.6161
## muTIC          0.0000 0.0447 >0.9999   -0.0876    0.0876
## phiTIC         0.9980 0.0631 <0.0001    0.8743    1.1217
## Y_mueta0       37.1868 0.5447 <0.0001   36.1192   38.2544
## Y_mueta1        1.7079 0.0171 <0.0001    1.6744    1.7414
## Y_mueta2        0.6836 0.0157 <0.0001    0.6528    0.7144
## Y_mu_knot      37.9057 0.5006 <0.0001   36.9245   38.8869
## covBL          0.4000 0.0523 <0.0001    0.2975    0.5025
## kappa         21.3467 1.2692 <0.0001   18.8591   23.8343
## Cov_XYres      0.5510 0.0619 <0.0001    0.4297    0.6723
##
## Slot "likelihood":
## data frame with 0 columns and 0 rows
##
## Slot "bootstrap":
## data frame with 0 columns and 0 rows

```

The model outputs highlight the merits of incorporating a decomposed time-varying covariate (TVC) into the analysis. First, the decomposed TVC enables the examination of the baseline and temporal effects of reading ability on mathematics development. According to the model results, the baseline effect of reading ability on the early-stage development of mathematics ability stands at 0.0708. This implies that for every standardized unit increase in baseline reading ability, the growth rate of mathematics ability before Grade 3 increases 0.0708. The temporal effect of reading ability on mathematics ability is reported to be 21.3467. This suggests, for instance, that the final mathematics score in the spring semester of Grade 1 increases by 21.3467 for each standardized unit growth in the reading ability growth rate within that semester. Secondly, the decomposed model facilitates the exploration of the relationship between the TVC baseline value and the TIC. For example, the covariance between read-




Figure 3: Latent Growth Curve Models with Bilinear Spline Function (Random Knots) for Mathematics Ability (TVC: Standardized Reading Ability over Time; Growth TIC: Teacher-reported Approach)

ing ability and the teacher-reported approach to learning was 0.4000 at the inception of the ECLS-K: 2011 study. More importantly, as displayed in Figure 3, incorporating a TVC into a longitudinal model tends to result in underestimated growth factors and growth trajectories since the longitudinal outcome is regressed on the TVC (or its temporal states). However, the extent of underestimation is noticeably less in the model with a decomposed TVC (Figure 3b) compared to the model where the TVC is directly integrated (Figure 3a).

### 6.5. `getMGM()` Examples

The `getMGM()` function is designed to construct a multivariate growth model (MGM) to analyze the development of reading and mathematics abilities and the correlations between these two developmental processes over time. In addition to the model fitting, we compute p-values and Wald confidence intervals for this bivariate longitudinal model. We present visualizations of the estimated growth trajectories for both reading and mathematics abilities in Figures 4a and 4b. Apart from the growth factors associated with each univariate developmental process, this constructed model enables us to estimate the covariances between the growth factors of different outcomes and the residual covariance. As demonstrated in the output, the relationships between the developmental processes of the two abilities are positive, evidenced by positive intercept-intercept ( $YZ\_psi00$ , p-value < 0.0001) and pre-knot slope-slope ( $YZ\_psi11$ , p-value < 0.0001) covariances. This suggests that students who demonstrated a higher reading ability at the inception of the ECLS-K:2011 study also tended to exhibit a higher level of mathematics ability and vice versa. Similarly, students with more rapid growth in reading ability during the early stage were generally associated with more rapid development in mathematics ability, and vice versa.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # Multivariate Latent Growth Curve Model, Bilinear Spline (random knot)
R> paraBLS_PLGCM_f <- c(
```


```

R> + "Y_mueta0", "Y_mueta1", "Y_mueta2", "Y_knot",
R> + paste0("Y_psi", c("00", "01", "02", "0g", "11", "12", "1g", "22", "2g", "gg")),
R> + "Y_res",
R> + "Z_mueta0", "Z_mueta1", "Z_mueta2", "Z_knot",
R> + paste0("Z_psi", c("00", "01", "02", "0g", "11", "12", "1g", "22", "2g", "gg")),
R> + "Z_res",
R> + paste0("YZ_psi", c(c("00", "10", "20", "g0", "01", "11", "21", "g1",
R> + "02", "12", "22", "g2", "0g", "1g", "2g", "gg"))),
R> + "YZ_res"
R> + )
R> RM_PLGCM.f <- getMGM(
R> + dat = RMS_dat0, t_var = c("T", "T"), y_var = c("R", "M"), curveFun = "BLS",
R> + intrinsic = TRUE, records = list(1:9, 1:9), y_model = "LGCM", res_scale =
R> + c(0.1, 0.1), res_cor = 0.3, paramOut = TRUE, names = paraBLS_PLGCM_f
R> + )
R> RM_PLGCM.f_o <- RM_PLGCM.f@mxOutput
R> RM_PLGCM.f_e <- RM_PLGCM.f@Estimates
R> # Get Wald confidence intervals of the constructed model
R> getEstimateStats(est_in = RM_PLGCM.f_e, CI_type = "Wald")
R> # Figure of the constructed model
R> Figure6 <- getFigure(
R> + model = RM_PLGCM.f_o, sub_Model = "MGM", y_var = c("R", "M"), curveFun = "BLS",
R> + y_model = "LGCM", t_var = c("T", "T"), records = list(1:9, 1:9), xstarts =
R> + xstarts, xlab = "Month", outcome = c("Reading", "Mathematics")
R> + )
R> show(Figure6) ## See Figures 4(a) and 4(b) in the manuscript

## An object of class "StatsOutput"
## Slot "wald":
##           Estimate      SE p.value wald_lbound wald_ubound
## ...
## YZ_psi00  88.5558 7.5042 <0.0001    73.8478    103.2638
## YZ_psi10   1.5594 0.3149 <0.0001     0.9422     2.1766
## YZ_psi20  -0.6429 0.1454 <0.0001    -0.9279    -0.3579
## YZ_psig0 -15.3384 3.4818 <0.0001   -22.1626   -8.5142
## YZ_psi01   0.6831 0.2057  0.0009     0.2799     1.0863
## YZ_psi11   0.0687 0.0108 <0.0001     0.0475     0.0899
## YZ_psi21   0.0018 0.0047  0.7017    -0.0074     0.0110
## YZ_psig1  -0.2231 0.1141  0.0505    -0.4467     0.0005
## YZ_psi02  -0.4524 0.1882  0.0162    -0.8213    -0.0835
## YZ_psi12  -0.0188 0.0094  0.0455    -0.0372    -0.0004
## YZ_psi22   0.0018 0.0046  0.6956    -0.0072     0.0108
## YZ_psig2   0.2468 0.1194  0.0387     0.0128     0.4808
## YZ_psi0g -18.8062 5.1876  0.0003   -28.9737   -8.6387
## YZ_psi1g   0.1887 0.2711  0.4864    -0.3426     0.7200
## YZ_psi2g   0.1961 0.1280  0.1255    -0.0548     0.4470
## YZ_psigg  -1.6773 3.8826  0.6657    -9.2871     5.9325

```

```
## YZ_res      7.6825 0.7061 <0.0001      6.2986      9.0664
##
## Slot "likelihood":
## data frame with 0 columns and 0 rows
##
## Slot "bootstrap":
## data frame with 0 columns and 0 rows
```



(a) Development Status of Reading Ability

(b) Development Status of Mathematics Ability

Figure 4: Multivariate Latent Growth Curve Models with Bilinear Spline Function (Random Knots) for Reading Ability and Mathematics Ability

## 6.6. getMediation() Examples

The `getMediation()` function is deployed to build a longitudinal mediation model. In this section, we set up two such models. The first model, employing a linear-linear functional form and a baseline predictor, was designed to analyze how the baseline approach to learning influences mathematics development via the mediation of reading ability development. Conversely, the second model, also adopting a linear-linear functional form but with a longitudinal predictor, examined how the developmental trajectory of reading ability influences the developmental process of science ability via the mediation of mathematics ability development. Both models are substantiated with p-values and Wald confidence intervals.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # Longitudinal Mediation Model with Baseline Predictor, Bilinear Spline
R> paraMed2_BLS <- c(
R> + "muX", "phi11", "alphaM1", "alphaMr", "alphaM2", "mugM",
R> + paste0("psi", c("M1M1", "M1Mr", "M1M2", "MrMr", "MrM2", "M2M2")), "_r"),
R> + "alphaY1", "alphaYr", "alphaY2", "mugY",
R> + paste0("psi", c("Y1Y1", "Y1Yr", "Y1Y2", "YrYr", "YrY2", "Y2Y2")), "_r"),
R> + paste0("beta", rep(c("M", "Y"), each = 3), rep(c(1, "r", 2), 2)),
R> + paste0("beta", c("M1Y1", "M1Yr", "M1Y2", "MrYr", "MrY2", "M2Y2"))),
```

```

R> + "muetaM1", "muetaMr", "muetaM2", "muetaY1", "muetaYr", "muetaY2",
R> + paste0("Mediator", c("11", "1r", "12", "rr", "r2", "22")),
R> + paste0("total", c("1", "r", "2")),
R> + "residualsM", "residualsY", "residualsYM"
R> + )
R> RM_BLS_LGCM <- getMediation(
R> + dat = RMS_dat0, t_var = rep("T", 2), y_var = "M", m_var = "R", x_type =
R> + "baseline", x_var = "ex1", curveFun = "BLS", records = list(1:9, 1:9),
R> + res_scale = c(0.1, 0.1),
R> + res_cor = 0.3, paramOut = TRUE, names = paraMed2_BLS
R> + )
R> # Get Wald confidence intervals for the constructed model
R> getEstimateStats(est_in = RM_BLS_LGCM@Estimates, CI_type = "Wald")
R> # Longitudinal Mediation Model with Longitudinal Predictor, Bilinear Spline
R> paraMed3_BLS <- c(
R> + "muetaX1", "muetaXr", "muetaX2", "mugX",
R> + paste0("psi", c("X1X1", "X1Xr", "X1X2", "XrXr", "XrX2", "X2X2")),
R> + "alphaM1", "alphaMr", "alphaM2", "mugM",
R> + paste0("psi", c("M1M1", "M1Mr", "M1M2", "MrMr", "MrM2", "M2M2")), "_r"),
R> + "alphaY1", "alphaYr", "alphaY2", "mugY",
R> + paste0("psi", c("Y1Y1", "Y1Yr", "Y1Y2", "YrYr", "YrY2", "Y2Y2")), "_r"),
R> + paste0("beta", c("X1Y1", "X1Yr", "X1Y2", "XrYr", "XrY2", "X2Y2",
R> + "X1M1", "X1Mr", "X1M2", "XrMr", "XrM2", "X2M2",
R> + "M1Y1", "M1Yr", "M1Y2", "MrYr", "MrY2", "M2Y2")),
R> + "muetaM1", "muetaMr", "muetaM2", "muetaY1", "muetaYr", "muetaY2",
R> + paste0("mediator", c("111", "11r", "112", "1rr", "1r2", "122", "rr2",
R> + "r22", "rrr", "222")),
R> + paste0("total", c("11", "1r", "12", "rr", "r2", "22")),
R> + "residualsX", "residualsM", "residualsY",
R> + "residualsMX", "residualsYX", "residualsYM"
R> + )
R> set.seed(20191029)
R> RMS_BLS_LGCM <- getMediation(
R> + dat = RMS_dat0, t_var = rep("T", 3), y_var = "S", m_var = "M", x_type =
R> + "longitudinal",
R> + x_var = "R", curveFun = "bilinear spline", records = list(2:9, 1:9, 1:9),
R> + res_scale = c(0.1, 0.1, 0.1), res_cor = c(0.3, 0.3), tries = 10, paramOut =
R> + TRUE, names = paraMed3_BLS)
R> # Get Wald confidence intervals for the constructed model
R> getEstimateStats(est_in = RMS_BLS_LGCM@Estimates, CI_type = "Wald")

## An object of class "StatsOutput"
## Slot "wald":
##           Estimate      SE p.value wald_lbound wald_ubound
## ...
## betaM1      0.0623  0.0231  0.0070      0.0170      0.1076
## betaMr      5.5471  0.6945 <0.0001      4.1859      6.9083

```

```

## betaM2      -0.0468  0.0118  0.0001      -0.0699      -0.0237
## betaY1       0.0149  0.0139  0.2837      -0.0123      0.0421
## betaYr       1.2907  0.5133  0.0119       0.2847      2.2967
## betaY2      -0.0212  0.0135  0.1163      -0.0477      0.0053
## betaM1Y1     0.3807  0.0317 <0.0001      0.3186      0.4428
## betaM1Yr     0.1206  0.9362  0.8975     -1.7143      1.9555
## betaM1Y2     0.0548  0.0505  0.2779     -0.0442      0.1538
## betaMrYr     0.7277  0.0309 <0.0001      0.6671      0.7883
## betaMrY2    -0.0012  0.0020  0.5485     -0.0051      0.0027
## betaM2Y2     0.4813  0.1434  0.0008       0.2002      0.7624
## ...
## Mediator11   0.0237  0.0090  0.0085       0.0061      0.0413
## Mediator1r   0.0075  0.0585  0.8980     -0.1072      0.1222
## Mediator12   0.0034  0.0034  0.3173     -0.0033      0.0101
## Mediatorrrr  4.0368  0.5298 <0.0001      2.9984      5.0752
## Mediatorrr2 -0.0066  0.0111  0.5521     -0.0284      0.0152
## Mediator22  -0.0225  0.0088  0.0106     -0.0397     -0.0053
## total1      0.0386  0.0156  0.0133       0.0080      0.0692
## totalr      5.3350  0.6953 <0.0001      3.9722      6.6978
## total2     -0.0469  0.0133  0.0004     -0.0730     -0.0208
## ...
##
## Slot "likelihood":
## data frame with 0 columns and 0 rows
##
## Slot "bootstrap":
## data frame with 0 columns and 0 rows

## An object of class "StatsOutput"
## Slot "wald":
##           Estimate      SE p.value wald_lbound wald_ubound
## ...
## betaX1Y1     0.3987  0.0313 <0.0001      0.3374      0.4600
## betaX1Yr     0.6677  1.2151  0.5827     -1.7139      3.0493
## betaX1Y2     0.0653  0.0571  0.2528     -0.0466      0.1772
## betaXrYr     0.7445  0.0347 <0.0001      0.6765      0.8125
## betaXrY2    -0.0020  0.0023  0.3845     -0.0065      0.0025
## betaX2Y2     0.4755  0.1700  0.0052      0.1423      0.8087
## betaX1M1     0.1540  0.0371 <0.0001      0.0813      0.2267
## betaX1Mr     4.4495  1.2668  0.0004      1.9666      6.9324
## betaX1M2    -0.1999  0.0700  0.0043     -0.3371     -0.0627
## betaXrMr     0.1960  0.0348 <0.0001      0.1278      0.2642
## betaXrM2     0.0090  0.0028  0.0013      0.0035      0.0145
## betaX2M2     0.8529  0.1818 <0.0001      0.4966      1.2092
## betaM1Y1     0.2718  0.0541 <0.0001      0.1658      0.3778
## betaM1Yr    -2.6244  1.5247  0.0852     -5.6128      0.3640
## betaM1Y2     0.0092  0.0928  0.9210     -0.1727      0.1911

```



```

## betaMrYr      0.2927  0.0367 <0.0001      0.2208      0.3646
## betaMrY2      0.0028  0.0024  0.2433      -0.0019      0.0075
## betaM2Y2      0.3738  0.1587  0.0185      0.0628      0.6848
...
## mediator111   0.1084  0.0229 <0.0001      0.0635      0.1533
## mediator11r  -1.0462  0.6148  0.0888      -2.2512      0.1588
## mediator112   0.0037  0.0370  0.9203      -0.0688      0.0762
## mediator1rr   0.1954  0.3497  0.5763      -0.4900      0.8808
## mediator1r2   0.0019  0.0039  0.6261      -0.0057      0.0095
## mediator122   0.0244  0.0269  0.3644      -0.0283      0.0771
## mediatorrrr2  0.2179  0.0308 <0.0001      0.1575      0.2783
## mediatorrr22  0.0021  0.0018  0.2433      -0.0014      0.0056
## mediatorrrrr  -0.0008  0.0010  0.4237      -0.0028      0.0012
## mediator222   0.1777  0.0713  0.0127      0.0380      0.3174
## total11       0.2624  0.0273 <0.0001      0.2089      0.3159
## total1r       3.5987  1.2804  0.0049      1.0892      6.1082
## total12      -0.1700  0.0633  0.0072      -0.2941     -0.0459
## totalrr       0.4140  0.0339 <0.0001      0.3476      0.4804
## totalr2       0.0103  0.0024 <0.0001      0.0056      0.0150
## total22       1.0306  0.1773 <0.0001      0.6831      1.3781
## ...
##
## Slot "likelihood":
## data frame with 0 columns and 0 rows
##
## Slot "bootstrap":
## data frame with 0 columns and 0 rows

```


Similar to the MGMs, these longitudinal models can estimate the growth trajectories for each univariate development and the relationship among these processes over time. However, the relationships between these processes are captured by the coefficients of unidirectional paths. For example, the output from the first model indicates that the baseline approach to learning exerts a positive influence on the growth rate of reading (betaM1, p-value = 0.0070) during the early developmental stage, as well as reading (betaMr, p-value < 0.0001) and mathematics ability at the knot (betaYr, p-value = 0.0119). Additionally, the impact of reading ability on mathematics ability during the early stage is also positive (betaM1Y1, p-value < 0.0001).

By using these path coefficients, we are able to calculate both the indirect effect (mediation effect) of the baseline approach to learning on mathematics development through reading development and the total effect of the approach to learning on mathematics development. For example, through the growth rate of reading ability during the early period, the indirect effect of the baseline approach to learning on the growth rate of mathematics ability was 0.0237. Therefore, the total effect of the baseline approach to learning on the early growth rate of mathematics ability is 0.0386 ( $0.0386 = 0.0149 + 0.0237$ ). The path coefficients, indirect effects, and total effects for the second longitudinal model can be interpreted in a similar manner.

### 6.7. getMGroup() Examples

The `getMGroup()` function is utilized to build a multiple-group latent growth curve model. This model employs a linear-linear functional form and random knot, allowing the examination of differences in mathematics development from Grade K through Grade 5 between girls and boys. As evidenced in Figure 5a, the development of mathematical ability in boys slightly outpaces that of girls. However, this discrepancy is not statistically significant, as evident by the overlapping confidence intervals between the two manifest groups.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # Multiple group Growth Models
R> MGroup_Math_BLS_LGCM_f <- getMGroup(
R> + dat = RMS_dat0, grp_var = "SEX", sub_Model = "LGCM", y_var = "M", t_var = "T",
R> + records = 1:9, curveFun = "BLS", intrinsic = TRUE, res_scale = list(0.1, 0.1)
R> + )@mxOutput
R> # Figure of the constructed model
R> Figure7 <- getFigure(
R> + model = MGroup_Math_BLS_LGCM_f, nClass = 2, cluster_TIC = NULL, grp_var = "SEX",
R> + sub_Model = "LGCM", y_var = "M", curveFun = "BLS", y_model = "LGCM", t_var = "T",
R> + records = 1:9, m_var = NULL, x_var = NULL, x_type = NULL, xstarts = xstarts,
R> + xlab = "Month", outcome = "Mathematics"
R> + )
R> show(Figure7) ## See Figure 5(a) in the manuscript
```



(a) Mathematics Development by Sex

(b) Mathematics Development by 3 Latent Classes

Figure 5: Multiple Group and Mixture Latent Growth Curve Models with Bilinear Spline Function for Mathematics Ability

### 6.8. getMIX() Examples

The `getMIX()` function is employed to construct mixture latent growth curve models, which incorporate a linear-linear functional form and a fixed knot. We first performed an enumeration process, where we built a set of potential models consisting of one to three latent classes.

Utilizing the `getSummary()` function with `HetModels = TRUE`, we were able to obtain estimates of the likelihood, AIC, BIC, class-specific residuals, and class-specific proportions for each of the three models. Both the estimated likelihood and the information criteria consistently pointed towards the three latent classes model as being optimal. We then plotted the estimated growth status for each of these three latent classes in Figure 5b. The diagram reveals that students in the third latent class outperformed the other two classes in mathematics. Although the other two classes showed overlapping patterns in mathematics development during the early stage, the growth pace of the first group slowed down earlier than the second group.

```
R> mxOption(model = NULL, key = "Default optimizer", "CSOLNP", reset = FALSE)
R> # LGCM, Bilinear Spline Functional Form (single group)
R> Math_BLS_LGCM1 <- getLGCM(
R> + dat = RMS_dat0, t_var = "T", y_var = "M", curveFun = "BLS", intrinsic = FALSE,
R> + records = 1:9, res_scale = 0.1
R> + )@mxOutput
R> # Mixture Models without Cluster TICs (2 latent classes)
R> Math_BLS_LGCM2 <- getMIX(
R> + dat = RMS_dat0, prop_starts = c(0.45, 0.55), sub_Model = "LGCM", y_var = "M",
R> + t_var = "T", records = 1:9, curveFun = "BLS", intrinsic = FALSE,
R> + res_scale = list(0.3, 0.3)
R> + )@mxOutput
R> # Mixture Models without Cluster TICs (3 latent classes)
R> set.seed(20191029)
R> Math_BLS_LGCM3 <- getMIX(
R> + dat = RMS_dat0, prop_starts = c(0.33, 0.34, 0.33), sub_Model = "LGCM",
R> + y_var = "M", t_var = "T", records = 1:9, curveFun = "BLS", intrinsic =
R> + FALSE, res_scale = list(0.3, 0.3, 0.3), tries = 10
R> + )@mxOutput
R> # Figure of Mixture Models with 3 latent classes
R> Figure8 <- getFigure(
R> + model = Math_BLS_LGCM3, nClass = 3, cluster_TIC = NULL, sub_Model = "LGCM",
R> + y_var = "M", curveFun = "BLS", y_model = "LGCM", t_var = "T", records = 1:9,
R> + m_var = NULL, x_var = NULL, x_type = NULL, xstarts = xstarts, xlab = "Month",
R> + outcome = "Mathematics"
R> + )
R> show(Figure8) ## See Figure 5(b) in the manuscript
```

##	Model	No_Params	-2ll	AIC	BIC	Y_res_c1	Y_res_c2	Y_res_c3
##	Model1	11	31347.39	31369.39	31415.75	34.0030	NA	NA
##	Model2	23	31134.61	31180.61	31277.55	34.2367	31.5802	NA
##	Model3	35	31008.30	31078.30	31225.81	29.5689	31.0093	31.4662
##	%Class1	%Class2	%Class3					
##	100%	NA	NA					
##	67.4%	32.6%	NA					
##	22.4%	41.6%	36.0%					

## 7. Concluding Remarks

The developed R package, **nlpsem**, aims to facilitate comprehensive evaluations of nonlinear longitudinal processes, including intrinsically nonlinear functional forms within the SEM framework. It currently supports three commonly used intrinsically nonlinear functional forms, namely, the individual ratio of growth rate under the negative exponential function, the individual ratio of growth acceleration within the Jenss-Bayley function, and the individual knot in the bilinear spline function, also known as the linear-linear piecewise function. In addition, this package is versatile enough to handle parsimonious models and models with a quadratic functional form, both under Type I and Type II of nonlinear longitudinal models. Despite not primarily focusing on models for linear longitudinal processes, **nlpsem** incorporates functionalities for them, making it a comprehensive tool for researchers in the field. The package provides computational resources for univariate longitudinal processes, with the option to include or exclude time-invariant covariates. Further, it facilitates estimations for multivariate longitudinal processes, including a longitudinal outcome with time-varying covariates, correlated growth models for multiple outcomes, and longitudinal mediation models. Multiple group and mixture models are accommodated within **nlpsem**, where the sub-model can be any of the types above. Built on the **OpenMx** package, it enables flexible SEM specification and data-driven parameter estimation through built-in optimizers. Note that the package allows for unstructured time frame compatibility by employing the definition variables approach.

Despite its capabilities, **nlpsem** has limitations, which also pave the way for future developments. First, other nonlinear functional forms, such as logistic and Gompertz functions, are not currently supported. The inclusion of such additional forms could enhance the flexibility and applicability of the package. Second, formal statistical hypothesis testing needs to be developed for complex longitudinal models to evaluate the impact of removing or adding specific paths on the overall model. Third, although several nonlinear longitudinal models with certain functional forms have been well-documented and validated by simulation studies, others still need to be explored. Conducting further simulation studies to investigate the performance of these lesser-known models under different scenarios will provide essential insights and improve the robustness and validity of the **nlpsem** package.

## References

- Baron RM, Kenny DA (1986). “The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations.” *Journal of personality and social psychology*, **52**(6), 1173–1182. URL <https://doi.org/10.1037//0022-3514.51.6.1173>.
- Bates DM, Mächler M, Bolker B, Walker S (2014). *lme4: Linear mixed-effects models using Eigen and S4*. R package version 1.1-7, URL <http://CRAN.R-project.org/package=lme4>.
- Bates DM, Mächler M, Bolker B, Walker S (2015). “Fitting Linear Mixed-Effects Models Using lme4.” *Journal of Statistical Software*, **67**(1), 1–48. URL <https://doi.org/10.18637/jss.v067.i01>.

- Bates DM, Watts DG (1988). *Nonlinear Regression Analysis and Its Applications*. John Wiley & Sons, Ltd. URL <https://doi.org/10.1002/9780470316757.ch2>.
- Bauer DJ (2003). “Estimating Multilevel Linear Models as Structural Equation Models.” *Journal of Educational and Behavioral Statistics*, **28**(2), 135–167.
- Bauer DJ, Curran PJ (2003). “Distributional Assumptions of Growth Mixture Models: Implications for Overextraction of Latent Trajectory Classes.” *Psychological Methods*, **8**(3), 338–363. URL <https://doi.org/10.1037/1082-989X.8.3.338>.
- Biesanz JC, Deeb-Sossa N, Papadakis AA, Bollen KA, Curran PJ (2004). “The role of coding time in estimating and interpreting growth curve models.” *Psychological methods*, **9**(1), 30–52. URL <https://doi.org/10.1037/1082-989X.9.1.30>.
- Blozis SA (2004). “Structured Latent Curve Models for the Study of Change in Multivariate Repeated Measures.” *Psychological Methods*, **9**(3), 334–353. URL <https://doi.org/10.1037/1082-989X.9.3.334>.
- Blozis SA, Cho Y (2008). “Coding and Centering of Time in Latent Curve Models in the Presence of Interindividual Time Heterogeneity.” *Structural Equation Modeling: A Multidisciplinary Journal*, **15**(3), 413–433.
- Boker SM, Neale MC, Maes HH, Wilde MJ, Spiegel M, Brick TR, Estabrook R, Bates TC, Mehta P, von Oertzen T, Gore RJ, Hunter MD, Hackett DC, Karch J, Brandmaier AM, Pritikin JN, Zahery M, Kirkpatrick RM (2020). *OpenMx 2.17.2 User Guide*. URL <https://vipbg.vcu.edu/vipbg/OpenMx2/docs/OpenMx/2.17.2/OpenMxUserGuide.pdf>.
- Bouveyron C, Celeux G, Murphy T, Raftery A (2019). *Model-Based Clustering and Classification for Data Science: With Applications in R (Cambridge Series in Statistical and Probabilistic Mathematics)*. Cambridge: Cambridge University Press. URL <https://doi.org/10.1017/9781108644181>.
- Browne MW, du Toit SHC (1991). “Models for learning data.” In LM Collins, JL Horn (eds.), *Best methods for the analysis of change: Recent advances, unanswered questions, future directions*, chapter 4, pp. 47–68. American Psychological Association., Washington, DC, US.
- Casella G, Berger R (2002). *Statistical Inference*. 2nd edition. Duxbury Press.
- Cheong J, Mackinnon DP, Khoo ST (2003). “Investigation of Mediational Processes Using Parallel Process Latent Growth Curve Modeling.” *Structural equation modeling: a multidisciplinary journal*, **10**(2), 238–262. URL [https://doi.org/10.1207/S15328007SEM1002\\_5](https://doi.org/10.1207/S15328007SEM1002_5).
- Cheung GW, Lau RS (2008). “Testing Mediation and Suppression Effects of Latent Variables: Bootstrapping With Structural Equation Models.” *Organizational Research Methods*, **11**(2), 296–325. URL <https://doi.org/10.1177/1094428107300343>.
- Cole DA, Maxwell SE (2003). “Testing mediational models with longitudinal data: questions and tips in the use of structural equation modeling.” *Journal of abnormal psychology*, **112**(4), 558–577. URL <https://doi.org/10.1037/0021-843X.112.4.558>.

- Coulombe P, Selig JP, Delaney HD (2015). “Ignoring individual differences in times of assessment in growth curve modeling.” *International Journal of Behavioral Development*, **40**(1), 76–86.
- Cudeck R, Harring JR (2010). “Developing a random coefficient model for nonlinear repeated measures data.” In SM Chow, E Ferrer, F Hsieh (eds.), *The Notre Dame series on quantitative methodology. Statistical methods for modeling human dynamics: An interdisciplinary dialogue*, chapter 2, pp. 289–318. Routledge/Taylor & Francis Group., New York, NY: Routledge.
- Curran PJ (2003). “Have Multilevel Models Been Structural Equation Models All Along?” *Multivariate Behavioral Research*, **38**(4), 529–569.
- Diallo TMO, Morin AJS, Lu H (2017). “The impact of total and partial inclusion or exclusion of active and inactive time invariant covariates in growth mixture models.” *Psychological Methods*, **22**(1), 166–190. URL <https://doi.org/10.1037/met0000084>.
- Du Toit SHC, Cudeck R (2009). “Estimation of the nonlinear random coefficient model when some random effects are separable.” *Psychometrika*, **74**, 65–82. URL <https://doi.org/10.1007/s11336-008-9107-7>.
- Dumenci L (2011). “The psychometric latent agreement model (PLAM) for discrete latent variables measured by multiple items.” *Organizational Research Methods*, **14**(1), 91–115. URL <https://doi.org/10.1177/1094428110374649>.
- Dumenci L, Perera RA, Keefe FJ, Ang DC, Slover J, Jensen MP, Riddle DL (2019). “Model-based pain and function outcome trajectory types for patients undergoing knee arthroplasty: a secondary analysis from a randomized clinical trial.” *Osteoarthritis and Cartilage*, **27**(6), 878–884. URL <https://doi.org/10.1016/j.joca.2019.01.004>.
- Efron B, Tibshirani RJ (1994). *An introduction to the bootstrap*. CRC press.
- Estabrook R, Neale MC (2013). “A Comparison of Factor Score Estimation Methods in the Presence of Missing Data: Reliability and an Application to Nicotine Dependence.” *Multivariate Behavioral Research*, **48**, 1–27. URL <https://doi.org/10.1080/00273171.2012.730072>.
- Feng Z, McCulloch CE (1996). “Using bootstrap likelihood ratios in finite mixture models.” *Journal of the Royal Statistical Society, Series B*, **58**, 609–617. [Google Scholar].
- Ferrer E, McArdle JJ (2003). “Alternative Structural Models for Multivariate Longitudinal Data Analysis.” *Structural Equation Modeling: A Multidisciplinary Journal*, **10**(4), 493–524. URL [https://doi.org/10.1207/S15328007SEM1004\\_1](https://doi.org/10.1207/S15328007SEM1004_1).
- Gill PE, Murray W, Saunders MA, Wright MH (1986). “User’s Guide for NPSOL (Version 4.0): A Fortran Package for Nonlinear Programming.” *Technical report*, Stanford University, Systems Optimization Lab. URL <https://apps.dtic.mil/sti/citations/ADA169115>.
- Gollob H, Reichardt C (1987). “Taking Account of Time Lags in Causal Models.” *Child Development*, **58**(1), 80–92. URL <https://doi.org/10.2307/1130293>.

- Grimm KJ (2007). “Multivariate longitudinal methods for studying developmental relationships between depression and academic achievement.” *International Journal of Behavioral Development*, **31**(4), 328–339. URL <https://doi.org/10.1177/0165025407077754>.
- Grimm KJ, Castro-Schilo L, Davoudzadeh P (2013a). “Modeling intraindividual change in nonlinear growth models with latent change scores.” *GeroPsych: The Journal of Gerontopsychology and Geriatric Psychiatry*, **26**(3), 153–162. URL <https://doi.org/10.1024/1662-9647/a000093>.
- Grimm KJ, Ram N (2009). “Nonlinear Growth Models in Mplus and SAS.” *Structural Equation Modeling: A Multidisciplinary Journal*, **16**(4), 676–701. URL <https://doi.org/10.1080/10705510903206055>.
- Grimm KJ, Ram N, Estabrook R (2010). “Nonlinear Structured Growth Mixture Models in Mplus and OpenMx.” *Multivariate Behavioral Research*, **45**(6), 887–909. URL <https://doi.org/10.1080/00273171.2010.531230>.
- Grimm KJ, Ram N, Estabrook R (2016). *Growth Modeling: Structural Equation and Multi-level Modeling Approaches*. Guilford Press.
- Grimm KJ, Zhang Z, Hamagami F, Mazzocco M (2013b). “Modeling Nonlinear Change via Latent Change and Latent Acceleration Frameworks: Examining Velocity and Acceleration of Growth Trajectories.” *Multivariate Behavioral Research*, **48**(1), 117–143. URL <https://doi.org/10.1080/00273171.2012.755111>.
- Harring JR (2014). “A Spline Regression Model for Latent Variables.” *Educational and Psychological Measurement*, **74**(2), 197–213. URL <https://doi.org/10.1177/0013164413504295>.
- Harring JR, Cudeck R, du Toit SHC (2006). “Fitting Partially Nonlinear Random Coefficient Models as SEMs.” *Multivariate Behavioral Research*, **41**(4), 579–596.
- Hayes AF (2009). “Beyond Baron and Kenny: Statistical mediation analysis in the new millennium.” *Communication Monographs*, **76**(4), 408–420. URL <https://doi.org/10.1080/03637750903310360>.
- Hunter MD (2018). “State Space Modeling in an Open Source, Modular, Structural Equation Modeling Environment.” *Structural Equation Modeling*, **25**(2), 307–324. URL <https://doi.org/10.1080/10705511.2017.1369354>.
- Johnson SG (2014). “The NLOpt nonlinear-optimization package.” URL [Availableonline: https://nlopt.readthedocs.io/en/latest/NLOpt\\_Introduction/](https://nlopt.readthedocs.io/en/latest/NLOpt_Introduction/).
- Kraft D (1994). “Algorithm 733: TOMP–Fortran modules for optimal control calculations.” *ACM Transactions on Mathematical Software (TOMS)*, **20**(3), 262–281. URL <https://doi.org/10.1145/192115.192124>.
- Liu J (2021). “Extending Latent Basis Growth Model to Explore Joint Development in the Framework of Individual Measurement Occasions.” URL <https://arxiv.org/abs/2107.01773>.



- Liu J (2022a). “Decomposing Impact on Longitudinal Outcome of Time-varying Covariate into Baseline Effect and Temporal Effect.” URL <https://arxiv.org/abs/2210.16916>.
- Liu J (2022b). “Jenss–Bayley Latent Change Score Model With Individual Ratio of the Growth Acceleration in the Framework of Individual Measurement Occasions.” *Journal of Educational and Behavioral Statistics*, **47**(5), 507–543. URL <https://doi.org/10.3102/10769986221099919>.
- Liu J, Perera R (2022a). “Assessing mediational processes using piecewise linear growth curve models with individual measurement occasions.” *Behavior Research Methods (Advance online publication)*. URL <https://doi.org/10.3758/s13428-022-01940-2>.
- Liu J, Perera RA (2021). “Estimating Knots and Their Association in Parallel Bilinear Spline Growth Curve Models in the Framework of Individual Measurement Occasions.” *Psychological Methods (Advance online publication)*. URL <https://doi.org/10.1037/met0000309>.
- Liu J, Perera RA (2022b). “Extending Growth Mixture Model to Assess Heterogeneity in Joint Development With Piecewise Linear Trajectories in the Framework of Individual Measurement Occasions.” *Psychological Methods (Advance online publication)*. URL <http://dx.doi.org/10.1037/met0000500>.
- Liu J, Perera RA (2022c). “Extending Mixture of Experts Model to Investigate Heterogeneity of Trajectories: When, Where and How to Add Which Covariates.” *Psychological Methods (Advance online publication)*. URL <https://doi.org/10.1037/met0000436>.
- Liu J, Perera RA (2023a). “Estimating Rate of Change for Nonlinear Trajectories in the Framework of Individual Measurement Occasions: A New Perspective on Growth Curves.” *Behavior Research Methods (In press)*.
- Liu J, Perera RA (2023b). “Further Exploration of the Effects of Time-varying Covariate in Growth Mixture Models with Nonlinear Trajectories.” *Behavior Research Methods (Advance online publication)*. URL <https://doi.org/10.3758/s13428-023-02183-5>.
- Liu J, Perera RA, Kang L, Kirkpatrick RM, Sabo RT (2021a). “Obtaining interpretable parameters from reparameterizing longitudinal models: transformation matrices between growth factors in two parameter spaces.” *Journal of Educational and Behavioral Statistics*. URL <https://doi.org/10.3102/10769986211052009>.
- Liu J, Perera RA, Kang L, Sabo RT, Kirkpatrick RM (2021b). “Two-step growth mixture model to examine heterogeneity in nonlinear trajectories.” *Journal of Behavioral Data Science*, **Advance online publication**. URL <https://doi.org/10.35566/jbds/v1n2/p4>.
- MacKinnon DP (2008). *Introduction to statistical mediation analysis*. Fourth edition. Taylor & Francis Group/Lawrence Erlbaum Associates., New York.
- MacKinnon DP, Krull JL, Lockwood CM (2000). “Equivalence of the Mediation, Confounding and Suppression Effect.” *Prevention Science*, **1**, 173–181. URL <https://doi.org/10.1023/A:1026595011371>.
- MacKinnon DP, Lockwood CM, Hoffman JM, West SG, Sheets V (2002). “A comparison of methods to test mediation and other intervening variable effects.” *Psychological Methods*, **7**(1), 83–104. URL <https://doi.org/10.1037/1082-989x.7.1.83>.

- Madansky A (1965). “Approximate Confidence Limits for the Reliability of Series and Parallel Systems.” *Technometrics*, **7**(4), 495–503. URL <https://www.jstor.org/stable/1266390>.
- Matthews DE (1988). “Likelihood-Based Confidence Intervals for Functions of Many Parameters.” *Biometrika*, **75**(1), 139–144. URL <https://www.jstor.org/stable/2336444>.
- Maxwell SE, Cole DA (2007). “Bias in cross-sectional analyses of longitudinal mediation.” *Psychological methods*, **12**(1), 23–44. URL <https://doi.org/10.1037/1082-989X.12.1.23>.
- McArdle JJ (1988). “Dynamic but Structural Equation Modeling of Repeated Measures Data.” In J Nesselroade, R Cattell (eds.), *Handbook of Multivariate Experimental Psychology*, chapter 17, pp. 561–614. Springer, Boston, MA.
- McArdle JJ (2001). “A latent difference score approach to longitudinal dynamic structural analysis.” In R Cudeck, SHC du Toit, D Sorbom (eds.), *Structural equation modeling: Present and future*, pp. 342–380. Lincolnwood, IL: Scientific Software International.
- McArdle JJ (2009). “Latent variable modeling of differences and changes with longitudinal data.” *Annual review of psychology*, **60**, 577–605. URL <https://doi.org/10.1146/annurev.psych.60.110707.163612>.
- McArdle JJ, Hamagami F (2001). “Latent difference score structural models for linear dynamic analyses with incomplete longitudinal data.” In LM Collins, AG Sayer (eds.), *Decade of behavior. New methods for the analysis of change*, p. 139–175. American Psychological Association. URL <https://doi.org/10.1037/10409-005>.
- Mehta PD, Neale MC (2005). “People are variables too: Multilevel structural equations modeling.” *Psychological Methods*, **10**(3), 259–284.
- Mehta PD, West SG (2000). “Putting the individual back into individual growth curves.” *Psychological Methods*, **5**(1), 23–43.
- Muthén B (2004). “Latent Variable Analysis:.” In D Kaplan (ed.), *The SAGE Handbook of Quantitative Methodology for the Social Sciences*, chapter 19, pp. 345–368. SAGE Publications, Washington, DC, US.
- Muthén B, Shedden K (1999). “Finite Mixture Modeling with Mixture Outcomes Using the EM Algorithm.” *Biometrics*, **55**(2), 463–469. URL <https://doi.org/10.1111/j.0006-341x.1999.00463.x>.
- Muthén LK, Muthén BO (2017). *Mplus: Statistical Analysis with Latent Variables: User’s Guide (Version 8)*.
- Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kirkpatrick RM, Estabrook R, Bates TC, Maes HH, Boker SM (2016). “OpenMx 2.0: Extended structural equation and statistical modeling.” *Psychometrika*, **81**(2), 535–549. URL <https://doi.org/10.1007/s11336-014-9435-8>.
- Nylund KL, Asparouhov T, Muthén BO (2007). “Deciding on the Number of Classes in Latent Class Analysis and Growth Mixture Modeling: A Monte Carlo Simulation Study.”

- Structural Equation Modeling: A Multidisciplinary Journal*, **14**(4), 535–569. URL <https://doi.org/10.1080/10705510701575396>.
- Nylund-Gibson K, Masyn KE (2016). “Covariates and mixture modeling: Results of a simulation study exploring the impact of misspecified effects on class enumeration.” *Structural Equation Modeling: A Multidisciplinary Journal*, **23**(6), 782–797. URL <https://doi.org/10.1080/10705511.2016.1221313>.
- Pinheiro JC, Bates DM (2000). *Mixed-Effects Models in S and S-PLUS*. Springer, New York. doi:10.1007/b98882.
- Pinheiro JC, Bates DM, R Core Team (2023). *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1-162, URL <https://CRAN.R-project.org/package=nlme>.
- Preacher KJ, Hancock GR (2015). “Meaningful aspects of change as novel random coefficients: A general method for reparameterizing longitudinal models.” *Psychological Methods*, **20**(1), 84–101.
- Priestley MB, Subba Rao T (1975). “The estimation of factor scores and Kalman filtering for discrete parameter stationary processes.” *International Journal of Control*, **21**, 971–975. URL <https://doi.org/10.1080/00207177508922050>.
- Pritikin JN, Hunter MD, Boker SM (2015). “Modular open-source software for Item Factor Analysis.” *Educational and Psychological Measurement*, **75**(3), 458–474. URL <https://doi.org/10.1177/0013164414554615>.
- Pritikin JN, Rappaport LM, Neale MC (2017). “Likelihood-Based Confidence Intervals for a Parameter With an Upper or Lower Bound.” *Structural Equation Modeling: A Multidisciplinary Journal*, **24**(3), 395–401. URL <https://doi.org/10.1080/10705511.2016.1275969>.
- Proust-Lima C, Philipps V, Lique B (2017). “Estimation of extended mixed models using latent classes and latent processes: The R package lcmm.” *Journal of Statistical Software*, **78**(2). URL <http://dx.doi.org/10.18637/jss.v078.i02>.
- Rohloff CT, Kohli N, Chung S (2022). “The Impact of Functional Form Complexity on Model Overfitting for Nonlinear Mixed-Effects Models.” *Multivariate Behavioral Research*. URL <https://doi.org/10.1080/00273171.2022.2119360>.
- Selig JP, Preacher KJ (2009). “Mediation Models for Longitudinal Data in Developmental Research.” *Research in Human Development*, **6**(2-3), 144–164. URL <https://doi.org/10.1080/15427600902911247>.
- Shrout PE, Bolger N (2002). “Mediation in experimental and nonexperimental studies: New procedures and recommendations.” *Psychological Methods*, **7**(4), 422–445. URL <https://doi.org/10.1037/1082-989X.7.4.422>.
- Soest T, Hagtvet KA (2011). “Mediation Analysis in a Latent Growth Curve Modeling Framework.” *Structural equation modeling: a multidisciplinary journal*, **18**(2), 289–314. URL <https://doi.org/10.1080/10705511.2011.557344>.

- Sterba SK (2014). “Fitting Nonlinear Latent Growth Curve Models With Individually Varying Time Points.” *Structural Equation Modeling: A Multidisciplinary Journal*, **21**(4), 630–647. URL <https://doi.org/10.1080/10705511.2014.919828>.
- Wald A (1943). “Tests of statistical hypotheses concerning several parameters when the number of observations is large.” *Transactions of the American Mathematical Society*, **54**(3), 426–482. URL <https://doi.org/10.2307/1990256>.
- Zahery M, Maes H, Neale M (2017). “CSOLNP: Numerical Optimization Engine for Solving Non-linearly Constrained Problems.” *Twin Research and Human Genetics*, **20**(4), 290–297. URL <https://doi.org/10.1017/thg.2017.28>.
- Zhang Z, McArdle JJ, Nesselroade JR (2012). “Growth rate models: emphasizing growth rate analysis through growth curve modeling.” *Journal of Applied Statistics*, **39**(6), 1241–1262. URL <https://doi.org/10.1080/02664763.2011.644528>.
- Zopluoglu C, Harring JR, Kohli N (2014). “FitPMM: An R routine to fit finite mixtures of piecewise mixed-effects models with unknown random knots.” *Applied Psychological Measurement*, **38**, 583–584. URL <https://doi.org/10.1177/0146621614540482>.

Table 1: Comparison of Tools Addressing Basic Challenges in Longitudinal Data Analysis

	nlme	lme4	lcmm	fitPMM	OpenMx	MPlus 8	nlpssem
SEM Framework					✓	✓	✓
First and Second Type of Nonlinear Models	✓	✓	✓	✓	✓	✓	✓
Third Type of Nonlinear Models				✓	✓	✓	✓
Time-invariant Covariates	✓	✓	✓		✓	✓	✓
Allow for Individual Measurement Occasions	✓	✓	✓		✓	✓	✓
Allow for Latent Classes			✓		✓	✓	✓
User-friendly <sup>a</sup>	✓	✓	✓	✓			✓

<sup>a</sup> User-friendly implies that the tool provides built-in functionalities or templates for common analyses, reducing the need for users to manually specify or script the model. This facilitates a more intuitive and efficient user experience, especially for those who may not be well-versed in the intricacies of model specification.

### Affiliation:

Jin Liu

E-mail: [Veronica.Liu0206@gmail.com](mailto:Veronica.Liu0206@gmail.com)

Table 2: Model Specification for Commonly Used Latent Growth Curve Models with Individual Measurement Occasions

Linear Function		
Individual Growth Curve <sup>a</sup>	$y_{ij} = \eta_{0i} + \eta_{1i} \times t_{ij} + \epsilon_{ij}$	
Growth Factors <sup>b</sup>	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i})$	
Factor Loadings <sup>b</sup>	$\boldsymbol{\Lambda}_i = (1 \quad t_{ij})$	
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual linear component of change	
Quadratic Function		
Individual Growth Curve <sup>a</sup>	$y_{ij} = \eta_{0i} + \eta_{1i} \times t_{ij} + \eta_{2i} \times t_{ij}^2 + \epsilon_{ij}$	
Growth Factors <sup>b</sup>	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i})$	
Factor Loadings <sup>b</sup>	$\boldsymbol{\Lambda}_i = (1 \quad t_{ij} \quad t_{ij}^2)$	
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual linear component of change $\eta_{2i}$ : the individual quadratic component of change (i.e., half of the individual growth acceleration)	
Negative Exponential Function		
	Intrinsically Nonlinear Model	Reduced Non-intrinsically Nonlinear Model
Individual Growth Curve <sup>a</sup>	$y_{ij} = \eta_{0i} + \eta_{1i} \times (1 - \exp(-b_i \times t_{ij})) + \epsilon_{ij}$	$y_{ij} = \eta_{0i} + \eta_{1i} \times (1 - \exp(-b \times t_{ij})) + \epsilon_{ij}$
Growth Factors <sup>b</sup>	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i} \quad b_i - \mu_b)$	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i})$
Factor Loadings <sup>b</sup>	$\boldsymbol{\Lambda}_i \approx (1 \quad 1 - \exp(-\mu_b \times t_{ij}) \quad \mu_{\eta_1} \times \exp(-\mu_b t_{ij}) \times t_{ij})$	$\boldsymbol{\Lambda}_i = (1 \quad 1 - \exp(-b \times t_{ij}))$
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual change from initial status to asymptotic level (i.e., the individual growth capacity) $b$ ( $b_i$ ) <sup>c</sup> : a growth rate parameter that controls the curvature of the growth trajectory (for individual $i$ )	
Jenss-Bayley Function		
	Intrinsically Nonlinear Model	Reduced Non-intrinsically Nonlinear Model
Individual Growth Curve <sup>a</sup>	$y_{ij} = \eta_{0i} + \eta_{1i} \times t_{ij} + \eta_{2i} \times (\exp(c_i \times t_{ij}) - 1) + \epsilon_{ij}$	$y_{ij} = \eta_{0i} + \eta_{1i} \times t_{ij} + \eta_{2i} \times (\exp(c \times t_{ij}) - 1) + \epsilon_{ij}$
Growth Factors <sup>b</sup>	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i} \quad c_i - \mu_c)$	$\boldsymbol{\eta}_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i})$
Factor Loadings <sup>b</sup>	$\boldsymbol{\Lambda}_i \approx (1 \quad t_{ij} \quad \exp(\mu_c \times t_{ij}) - 1 \quad \mu_{\eta_2} \times \exp(\mu_c t_{ij}) \times t_{ij})$	$\boldsymbol{\Lambda}_i = (1 \quad t_{ij} \quad \exp(c \times t_{ij}) - 1)$
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual slope of linear asymptote with the assumption $c_i < 0$ ( $c < 0$ ) <sup>d</sup> $\eta_{2i}$ : the individual change from initial status to the linear asymptote intercept $c$ ( $c_i$ ) <sup>e</sup> : a growth acceleration parameter that controls the rate of change of the growth trajectory's curvature (for individual $i$ )	
Bilinear Spline Function with an Unknown Knot		
	Intrinsically Nonlinear Model	Reduced Non-intrinsically Nonlinear Model
Individual Growth Curve <sup>a</sup>	$y_{ij} = \begin{cases} \eta_{0i} + \eta_{1i} \times t_{ij} + \epsilon_{ij}, & t_{ij} < \gamma_i \\ \eta_{0i} + \eta_{1i} \times \gamma_i + \eta_{2i} \times (t_{ij} - \gamma_i) + \epsilon_{ij}, & t_{ij} \geq \gamma_i \end{cases}$	$y_{ij} = \begin{cases} \eta_{0i} + \eta_{1i} \times t_{ij} + \epsilon_{ij}, & t_{ij} < \gamma \\ \eta_{0i} + \eta_{1i} \times \gamma + \eta_{2i} \times (t_{ij} - \gamma) + \epsilon_{ij}, & t_{ij} \geq \gamma \end{cases}$
Growth Factors <sup>b</sup>	$\boldsymbol{\eta}'_i = (\eta_{0i} + \gamma_i \eta_{1i} \quad \frac{\eta_{1i} + \eta_{2i}}{2} \quad \frac{\eta_{2i} - \eta_{1i}}{2} \quad \gamma_i - \mu_\gamma)$	$\boldsymbol{\eta}'_i = (\eta_{0i} + \gamma \eta_{1i} \quad \frac{\eta_{1i} + \eta_{2i}}{2} \quad \frac{\eta_{2i} - \eta_{1i}}{2})$
Factor Loadings <sup>b</sup>	$\boldsymbol{\Lambda}'_i \approx \begin{pmatrix} 1 & t_{ij} - \mu_\gamma &  t_{ij} - \mu_\gamma  & -\mu'_{\eta_2} - \frac{\mu'_{\eta_2}(t_{ij} - \mu_\gamma)}{ t_{ij} - \mu_\gamma } \end{pmatrix}$	$\boldsymbol{\Lambda}'_i = (1 \quad t_{ij} - \gamma \quad  t_{ij} - \gamma )$
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual slope of the first linear piece $\eta_{2i}$ : the individual slope of the second linear piece $\gamma$ ( $\gamma_i$ ): the (individual) transition time from 1 <sup>st</sup> linear piece to 2 <sup>nd</sup> linear piece (i.e., knot)	

<sup>a</sup> In the function of the individual growth curve,  $y_{ij}$ ,  $t_{ij}$ , and  $\epsilon_{ij}$  are the observed measurement, recorded time, and residual of the  $i^{th}$  individual at the  $j^{th}$  time point.

<sup>b</sup> In the vector of growth factors and the corresponding factor loadings,  $\mu_b$ ,  $\mu_c$ , and  $\mu_\gamma$  are the mean values of individual growth rate parameters, of individual growth acceleration parameters, and of individual knots for the negative exponential function, Jenss-Bayley function, and bilinear spline function with an unknown knot, respectively.

<sup>c</sup> There are multiple interpretations for  $b(b_i)$ . For example,  $\exp(-b_i \times (t_{i(j+1)} - t_{ij}))$  represents the ratio of the instantaneous growth rates at time points  $t_{i(j+1)}$  and  $t_{ij}$ . This value reflects how much the growth rate has changed between  $t_{ij}$  and  $t_{i(j+1)}$ , depending on the growth rate parameter  $b_i$ . With the assumption that measurements are taken at equally-spaced waves with scaled intervals,  $\exp(b_i)$  represents the ratio of the instantaneous rates at any adjacent time points.

<sup>d</sup> If  $c_i > 0$  ( $c > 0$ ), the Jenss-Bayley function does not have a linear asymptote as the nonlinear component continues to grow with time.

<sup>e</sup> There are multiple interpretations for  $c(c_i)$ . For example,  $\exp(c_i \times (t_{i(j+1)} - t_{ij}))$  represents the ratio of the instantaneous growth accelerations at time points  $t_{i(j+1)}$  and  $t_{ij}$ . This value reflects how much the growth acceleration has changed between  $t_{ij}$  and  $t_{i(j+1)}$ , depending on the growth acceleration parameter  $c_i$ . With the assumption that measurements are taken at equally-spaced waves with scaled intervals,  $\exp(c_i)$  represents the ratio of the instantaneous accelerations at any adjacent time points.

Table 3: Model Specification for Commonly Used Latent Change Score Models with Individual Measurement Occasions

Quadratic Function		
Individual Growth Rate <sup>b</sup>	$dy_{ij,\text{mid}} = \eta_{1i} + 2 \times \eta_{2i} \times t_{ij,\text{mid}}$	
Growth Factors of Growth Rate <sup>c</sup>	$\eta_{di} = (\eta_{1i} \quad \eta_{2i})$	
Factor Loadings of Growth Rate <sup>c</sup>	$\Lambda_{di} = (1 \quad 2 \times t_{ij,\text{mid}})$	
Growth Factors of Growth Status <sup>d</sup>	$\eta_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i})$	
Factor Loadings of Growth Status <sup>d</sup>	$\Lambda_i = (1 \quad \Omega_i \times \Lambda_{di})$	
Negative Exponential Function		
Intrinsically Nonlinear Model		Reduced Non-intrinsically Nonlinear Model
Individual Growth Rate <sup>b</sup>	$dy_{ij,\text{mid}} = b_i \times \eta_{1i} \times \exp(-b_i \times t_{ij,\text{mid}})$	$dy_{ij,\text{mid}} = b \times \eta_{1i} \times \exp(-b \times t_{ij,\text{mid}})$
Growth Factors of Growth Rate <sup>c,e</sup>	$\eta_{di} = (\eta_{1i} \quad b_i - \mu_b)$	$\eta_{di} = \eta_{1i}$
Factor Loadings of Growth Rate <sup>c,e</sup>	$\Lambda_{di} \approx (\mu_b \times \exp(-\mu_b \times t_{ij,\text{mid}}) \quad \mu_{\eta_1} \times \exp(-\mu_b t_{ij,\text{mid}}) \times (1 - \mu_b t_{ij,\text{mid}}))$	$\Lambda_{di} = b \times \exp(-b \times t_{ij,\text{mid}})$
Growth Factors of Growth Status <sup>d,e</sup>	$\eta_i = (\eta_{0i} \quad \eta_{1i} \quad b_i - \mu_b)$	$\eta_i = (\eta_{0i} \quad \eta_{1i})$
Factor Loadings of Growth Status <sup>d,e</sup>	$\Lambda_i \approx (1 \quad \Omega_i \times \Lambda_{di})$	$\Lambda_i = (1 \quad \Omega_i \times \Lambda_{di})$
Jenss-Bayley Function		
Intrinsically Nonlinear Model		Reduced Non-intrinsically Nonlinear Model
Individual Growth Rate <sup>b</sup>	$dy_{ij,\text{mid}} = \eta_{1i} + c_i \times \eta_{2i} \times \exp(c_i \times t_{ij,\text{mid}})$	$dy_{ij,\text{mid}} = \eta_{1i} + c \times \eta_{2i} \times \exp(c \times t_{ij,\text{mid}})$
Growth Factors of Growth Rate <sup>c,e</sup>	$\eta_{di} = (\eta_{1i} \quad \eta_{2i} \quad c_i - \mu_c)$	$\eta_{di} = (\eta_{1i} \quad \eta_{2i})$
Factor Loadings of Growth Rate <sup>c,e</sup>	$\Lambda_{di} \approx (1 \quad \mu_c \times \exp(\mu_c \times t_{ij,\text{mid}}) \quad \mu_{\eta_2} \times \exp(\mu_c t_{ij,\text{mid}}) \times (1 + \mu_c t_{ij,\text{mid}}))$	$\Lambda_{di} \approx (1 \quad c \times \exp(c \times t_{ij,\text{mid}}))$
Growth Factors of Growth Status <sup>d,e</sup>	$\eta_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i} \quad c_i - \mu_c)$	$\eta_i = (\eta_{0i} \quad \eta_{1i} \quad \eta_{2i})$
Factor Loadings of Growth Status <sup>d,e</sup>	$\Lambda_i \approx (1 \quad \Omega_i \times \Lambda_{di})$	$\Lambda_i = (1 \quad \Omega_i \times \Lambda_{di})$
Nonparametric Function		
Individual Growth Rate <sup>b</sup>	$dy_{ij} = \eta_{1i} \times \gamma_{j-1}$	
Growth Factors of Growth Rate <sup>c</sup>	$\eta_{di} = \eta_{1i}$	
Factor Loadings of Growth Rate <sup>c</sup>	$\Lambda_{di} = \gamma_{j-1}$	
Growth Factors of Growth Status <sup>d</sup>	$\eta_i = (\eta_{0i} \quad \eta_{1i})$	
Factor Loadings of Growth Status <sup>d</sup>	$\Lambda_i = (1 \quad \Omega_i \times \Lambda_{di})$	
Interpretation of Growth Coef.	$\eta_{0i}$ : the individual initial status $\eta_{1i}$ : the individual slope during the first time interval $\gamma_j$ : the relative growth rate of the $j^{\text{th}}$ interval	

<sup>a</sup> This table does not include the specifications for LCSMs with linear and bilinear spline functions, as LGCMS with these two functional forms can estimate interval-specific slopes, eliminating the need for LCSMs to estimate growth rates. Additionally, this table presents the model specifications for the LCSM with a piecewise linear function. Note that the specification of this model serves as the foundation for the models with a decomposed TVC, which is introduced in Subsection 2.3.

<sup>b</sup> In the individual growth rate function,  $dy_{ij,\text{mid}}$  and  $t_{ij,\text{mid}}$  are the instantaneous slope midway through the  $(j-1)^{\text{th}}$  time interval and the corresponding time.

<sup>c</sup> The growth factors of the growth rate  $\eta_{di}$  consists of those associated with the growth rates, which are present in the respective growth rate function. The corresponding factor loadings are provided in the matrix  $\Lambda_{di}$  representing the factor loadings of the growth rate (where  $j = 2, 3, \dots, J$ ). The mean vector and variance-covariance matrix of growth factors of growth rate are  $\mu_{\eta_d}$  and  $\Psi_{\eta_d}$ , respectively. With  $\eta_{di}$ ,  $\Lambda_{di}$ ,  $\mu_{\eta_d}$ , and  $\Psi_{\eta_d}$ , we are able to derive (1) mean and variance of interval-specific slopes:  $\mu_{dy,\text{mid}} = \Lambda_{\eta_d} \times \mu_{\eta_d}$  and  $\sigma_{dy,\text{mid}}^2 = \Lambda_{\eta_d} \times \Psi_{\eta_d} \times \Lambda_{\eta_d}^T$ , and (2) mean and variance of interval-specific changes:  $\mu_{\Delta y_{ij}} = \Lambda_{\eta_d} \times \mu_{\eta_d} \times (t_{ij} - t_{i(j-1)})$  and  $\sigma_{\Delta y_{ij}}^2 = \Lambda_{\eta_d} \times \Psi_{\eta_d} \times \Lambda_{\eta_d}^T \times (t_{ij} - t_{i(j-1)})^2$ . In the equations of means and variances of interval-specific slopes and interval-specific changes,  $j = 2, 3, \dots, J$ .

<sup>d</sup> The vector of growth factor of the growth status consists of growth factors associated with both the growth rates and the initial status, which together determine the growth status. The corresponding factor loadings are provided in the matrix representing the factor loadings of the growth status (where

$j = 1, 2, \dots, J$ ), in which  $\Omega_i = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots & 0 \\ t_{i2} - t_{i1} & 0 & \dots & \dots & \dots & 0 \\ t_{i2} - t_{i1} & t_{i3} - t_{i2} & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ t_{i2} - t_{i1} & t_{i3} - t_{i2} & t_{i4} - t_{i3} & \dots & \dots & t_{ij} - t_{i(j-1)} \end{pmatrix}$  so that  $\Omega_i \times \Lambda_{di}$  represents the accumulative value since the

initial status of the corresponding factor loading of the growth rate. With  $\Omega_i$ , we are able to derive the mean and variance of change from baseline:  $\mu_{\Delta y_{ij}} = \Omega_i \times \Lambda_{di} \times \mu_{\eta_d}$  and  $\sigma_{\Delta y_{ij}}^2 = \Omega_i \times \Lambda_{\eta_d} \times \Psi_{\eta_d} \times \Lambda_{\eta_d}^T \times \Omega_i^T$ . In the equations of means and variances of change from baseline,  $j = 2, 3, \dots, J$ .

<sup>e</sup> In the vector of growth factors and the corresponding factor loadings,  $\mu_b$  and  $\mu_c$  are the mean values of  $b_i$  and of  $c_i$  for the negative exponential function and Jenss-Bayley function, respectively.

Table 4: Model Specification for Four Possible Ways of Adding Time-varying Covariate with Individual Measurement Occasions

LGCM with a TVC and a TIC	
Model Specification	$\mathbf{y}_i = \mathbf{\Lambda}_i^{[y]} \times \boldsymbol{\eta}_i^{[y]} + \kappa \times \mathbf{x}_i + \boldsymbol{\epsilon}_i^{[y]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + \mathbf{B}_{\text{TIC}} \times \mathbf{X}_i + \boldsymbol{\zeta}_i^{[y]}$
LGCM with a Decomposed TVC into Baseline and Interval-specific Slopes and a TIC	
Individual Function of TVC	$x_{ij} = x_{ij}^* + \epsilon_{ij}^{[x]}$ $x_{ij}^* = \begin{cases} \eta_{0i}^{[x]}, & \text{if } j = 1 \\ x_{i(j-1)}^* + dx_{ij} \times (t_{ij} - t_{i(j-1)}), & \text{if } j = 2, \dots, J \end{cases}$ $dx_{ij} = \eta_{1i}^{[x]} \times \gamma_{j-1} \quad (j = 2, \dots, J)$
Model Specification	$\begin{pmatrix} \mathbf{x}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \mathbf{\Lambda}_i^{[x]} & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[x]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \kappa \times \begin{pmatrix} \mathbf{0} \\ d\mathbf{x}_i \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[x]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\mathbf{x}_i = \mathbf{\Lambda}_i^{[x]} \times \boldsymbol{\eta}_i^{[x]} + \boldsymbol{\epsilon}_i^{[x]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + (\mathbf{B}_{\text{TIC}} \quad \boldsymbol{\beta}_{\text{TVC}}) \times \begin{pmatrix} \mathbf{X}_i \\ \eta_{0i}^{[x]} \end{pmatrix} + \boldsymbol{\zeta}_i^{[y]}$ $d\mathbf{x}_i = (0 \quad dx_{i2} \quad dx_{i3} \quad \dots \quad dx_{iJ})^T$
LGCM with a Decomposed TVC into Baseline and Interval-specific Changes and a TIC	
Individual Function of TVC	$x_{ij} = x_{ij}^* + \epsilon_{ij}^{[x]}$ $x_{ij}^* = \begin{cases} \eta_{0i}^{[x]}, & \text{if } j = 1 \\ x_{i(j-1)}^* + \delta x_{ij}, & \text{if } j = 2, \dots, J \end{cases}$ $\delta x_{ij} = dx_{ij} \times (t_{ij} - t_{i(j-1)}) \quad (j = 2, \dots, J)$ $dx_{ij} = \eta_{1i}^{[x]} \times \gamma_{j-1} \quad (j = 2, \dots, J)$
Model Specification	$\begin{pmatrix} \mathbf{x}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \mathbf{\Lambda}_i^{[x]} & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[x]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \kappa \times \begin{pmatrix} \mathbf{0} \\ \delta \mathbf{x}_i \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[x]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\mathbf{x}_i = \mathbf{\Lambda}_i^{[x]} \times \boldsymbol{\eta}_i^{[x]} + \boldsymbol{\epsilon}_i^{[x]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + (\mathbf{B}_{\text{TIC}} \quad \boldsymbol{\beta}_{\text{TVC}}) \times \begin{pmatrix} \mathbf{X}_i \\ \eta_{0i}^{[x]} \end{pmatrix} + \boldsymbol{\zeta}_i^{[y]}$ $\delta \mathbf{x}_i = (0 \quad \delta x_{i2} \quad \delta x_{i3} \quad \dots \quad \delta x_{iJ})^T$
LGCM with a Decomposed TVC into Baseline and Change-from-baseline and a TIC	
Individual Function of TVC	$x_{ij} = x_{ij}^* + \epsilon_{ij}^{[x]}$ $x_{ij}^* = \begin{cases} \eta_{0i}^{[x]}, & \text{if } j = 1 \\ \eta_{0i}^{[x]} + \Delta x_{ij}, & \text{if } j = 2, \dots, J \end{cases}$ $\Delta x_{ij} = \Delta x_{i(j-1)} + dx_{ij} \times (t_{ij} - t_{i(j-1)})$ $dx_{ij} = \eta_{1i}^{[x]} \times \gamma_{j-1} \quad (j = 2, \dots, J)$
Model Specification	$\begin{pmatrix} \mathbf{x}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \mathbf{\Lambda}_i^{[x]} & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[x]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \kappa \times \begin{pmatrix} \mathbf{0} \\ \Delta \mathbf{x}_i \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[x]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\mathbf{x}_i = \mathbf{\Lambda}_i^{[x]} \times \boldsymbol{\eta}_i^{[x]} + \boldsymbol{\epsilon}_i^{[x]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + (\mathbf{B}_{\text{TIC}} \quad \boldsymbol{\beta}_{\text{TVC}}) \times \begin{pmatrix} \mathbf{X}_i \\ \eta_{0i}^{[x]} \end{pmatrix} + \boldsymbol{\zeta}_i^{[y]}$ $\Delta \mathbf{x}_i = (0 \quad \Delta x_{i2} \quad \Delta x_{i3} \quad \dots \quad \Delta x_{iJ})^T$



Table 5: Model Specification for Longitudinal Mediation Models with Individual Measurement Occasions

Baseline Covariate, Longitudinal Mediator, and Longitudinal Outcome		
Linear Function	Model Specification	$\begin{pmatrix} \mathbf{m}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \Lambda_i^{[m]} & \mathbf{0} \\ \mathbf{0} & \Lambda_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[m]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[m]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\boldsymbol{\eta}_i^{[u]} = (\eta_{0i}^{[u]} \quad \eta_{1i}^{[u]})^T \quad (u = m, y)$ $\Lambda_i^{[u]} = (0 \quad t_{ij}) \quad (u = m, y; j = 1, \dots, J)$ $\boldsymbol{\eta}_i^{[m]} = \boldsymbol{\alpha}^{[m]} + \mathbf{B}^{[x \rightarrow m]} \times x_i + \boldsymbol{\zeta}_i^{[m]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + \mathbf{B}^{[x \rightarrow y]} \times x_i + \mathbf{B}^{[m \rightarrow y]} \times \boldsymbol{\eta}_i^{[m]} + \boldsymbol{\zeta}_i^{[y]}$
	Growth Factor Intercept	$\boldsymbol{\alpha}_i^{[u]} = (\alpha_{0i}^{[u]} \quad \alpha_{1i}^{[u]})^T \quad (u = m, y)$
	Path Coef.	$\mathbf{B}^{[x \rightarrow m]} = \begin{pmatrix} \beta_0^{[x \rightarrow m]} & \beta_1^{[x \rightarrow m]} \end{pmatrix}^T; \mathbf{B}^{[x \rightarrow y]} = \begin{pmatrix} \beta_0^{[x \rightarrow y]} & \beta_1^{[x \rightarrow y]} \end{pmatrix}^T$ $\mathbf{B}^{[m \rightarrow y]} = \begin{pmatrix} \beta_{01}^{[m \rightarrow y]} & 0 \\ \beta_{01}^{[m \rightarrow y]} & \beta_{11}^{[m \rightarrow y]} \end{pmatrix}$
Bilinear Function	Model Specification	$\begin{pmatrix} \mathbf{m}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \Lambda_i^{[m]} & \mathbf{0} \\ \mathbf{0} & \Lambda_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[m]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[m]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\boldsymbol{\eta}_i^{[u]} = (\eta_{1i}^{[u]} \quad \eta_{\gamma i}^{[u]} \quad \eta_{2i}^{[u]})^T \quad (u = m, y)$ $\Lambda_i^{[u]} = (\min(0, t_{ij} - \gamma^{[u]}) \quad 1 \quad \max(0, t_{ij} - \gamma^{[u]})) \quad (u = m, y; j = 1, \dots, J)$ $\boldsymbol{\eta}_i^{[m]} = \boldsymbol{\alpha}^{[m]} + \mathbf{B}^{[x \rightarrow m]} \times x_i + \boldsymbol{\zeta}_i^{[m]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + \mathbf{B}^{[x \rightarrow y]} \times x_i + \mathbf{B}^{[m \rightarrow y]} \times \boldsymbol{\eta}_i^{[m]} + \boldsymbol{\zeta}_i^{[y]}$
	Growth Factor Intercept	$\boldsymbol{\alpha}_i^{[u]} = (\alpha_{1i}^{[u]} \quad \alpha_{\gamma i}^{[u]} \quad \alpha_{2i}^{[u]})^T \quad (u = m, y)$
	Path Coef.	$\mathbf{B}^{[x \rightarrow m]} = \begin{pmatrix} \beta_1^{[x \rightarrow m]} & \beta_\gamma^{[x \rightarrow m]} & \beta_2^{[x \rightarrow m]} \end{pmatrix}^T; \mathbf{B}^{[x \rightarrow y]} = \begin{pmatrix} \beta_1^{[x \rightarrow y]} & \beta_\gamma^{[x \rightarrow y]} & \beta_2^{[x \rightarrow y]} \end{pmatrix}^T$ $\mathbf{B}^{[m \rightarrow y]} = \begin{pmatrix} \beta_{11}^{[m \rightarrow y]} & 0 & 0 \\ \beta_{1\gamma}^{[m \rightarrow y]} & \beta_\gamma^{[m \rightarrow y]} & 0 \\ \beta_{12}^{[m \rightarrow y]} & \beta_\gamma^{[m \rightarrow y]} & \beta_{22}^{[m \rightarrow y]} \end{pmatrix}$
Longitudinal Covariate, Longitudinal Mediator, and Longitudinal Outcome		
Linear Function	Model Specification	$\begin{pmatrix} \mathbf{x}_i \\ \mathbf{m}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \Lambda_i^{[x]} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \Lambda_i^{[m]} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \Lambda_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[x]} \\ \boldsymbol{\eta}_i^{[m]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[x]} \\ \boldsymbol{\epsilon}_i^{[m]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\boldsymbol{\eta}_i^{[u]} = (\eta_{0i}^{[u]} \quad \eta_{1i}^{[u]})^T \quad (u = x, m, y)$ $\Lambda_i^{[u]} = (1 \quad t_{ij}) \quad (u = x, m, y; j = 1, \dots, J)$ $\boldsymbol{\eta}_i^{[x]} = \boldsymbol{\mu}_\eta + \boldsymbol{\zeta}_i^{[x]}$ $\boldsymbol{\eta}_i^{[m]} = \boldsymbol{\alpha}^{[m]} + \mathbf{B}^{[x \rightarrow m]} \times \boldsymbol{\eta}_i^{[x]} + \boldsymbol{\zeta}_i^{[m]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + \mathbf{B}^{[x \rightarrow y]} \times \boldsymbol{\eta}_i^{[x]} + \mathbf{B}^{[m \rightarrow y]} \times \boldsymbol{\eta}_i^{[m]} + \boldsymbol{\zeta}_i^{[y]}$
	Growth Factor Intercept	$\boldsymbol{\alpha}_i^{[u]} = (\alpha_{0i}^{[u]} \quad \alpha_{1i}^{[u]})^T \quad (u = m, y)$
	Path Coef.	$\mathbf{B}^{[x \rightarrow m]} = \begin{pmatrix} \beta_{00}^{[x \rightarrow m]} & 0 \\ \beta_{01}^{[x \rightarrow m]} & \beta_{11}^{[x \rightarrow m]} \end{pmatrix}; \mathbf{B}^{[x \rightarrow y]} = \begin{pmatrix} \beta_{00}^{[x \rightarrow y]} & 0 \\ \beta_{01}^{[x \rightarrow y]} & \beta_{11}^{[x \rightarrow y]} \end{pmatrix}$ $\mathbf{B}^{[m \rightarrow y]} = \begin{pmatrix} \beta_{00}^{[m \rightarrow y]} & 0 \\ \beta_{01}^{[m \rightarrow y]} & \beta_{11}^{[m \rightarrow y]} \end{pmatrix}$
Longitudinal Covariate, Longitudinal Mediator, and Longitudinal Outcome		
Bilinear Function	Model Specification	$\begin{pmatrix} \mathbf{x}_i \\ \mathbf{m}_i \\ \mathbf{y}_i \end{pmatrix} = \begin{pmatrix} \Lambda_i^{[x]} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \Lambda_i^{[m]} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \Lambda_i^{[y]} \end{pmatrix} \times \begin{pmatrix} \boldsymbol{\eta}_i^{[x]} \\ \boldsymbol{\eta}_i^{[m]} \\ \boldsymbol{\eta}_i^{[y]} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}_i^{[x]} \\ \boldsymbol{\epsilon}_i^{[m]} \\ \boldsymbol{\epsilon}_i^{[y]} \end{pmatrix}$ $\boldsymbol{\eta}_i^{[u]} = (\eta_{1i}^{[u]} \quad \eta_{\gamma i}^{[u]} \quad \eta_{2i}^{[u]})^T \quad (u = x, m, y)$ $\Lambda_i^{[u]} = (\min(0, t_{ij} - \gamma^{[u]}) \quad 1 \quad \max(0, t_{ij} - \gamma^{[u]})) \quad (u = x, m, y; j = 1, \dots, J)$ $\boldsymbol{\eta}_i^{[x]} = \boldsymbol{\mu}_\eta + \boldsymbol{\zeta}_i^{[x]}$ $\boldsymbol{\eta}_i^{[m]} = \boldsymbol{\alpha}^{[m]} + \mathbf{B}^{[x \rightarrow m]} \times \boldsymbol{\eta}_i^{[x]} + \boldsymbol{\zeta}_i^{[m]}$ $\boldsymbol{\eta}_i^{[y]} = \boldsymbol{\alpha}^{[y]} + \mathbf{B}^{[x \rightarrow y]} \times \boldsymbol{\eta}_i^{[x]} + \mathbf{B}^{[m \rightarrow y]} \times \boldsymbol{\eta}_i^{[m]} + \boldsymbol{\zeta}_i^{[y]}$
	Growth Factor Intercept	$\boldsymbol{\alpha}_i^{[u]} = (\alpha_{1i}^{[u]} \quad \alpha_{\gamma i}^{[u]} \quad \alpha_{2i}^{[u]})^T \quad (u = m, y)$
	Path Coef.	$\mathbf{B}^{[x \rightarrow m]} = \begin{pmatrix} \beta_{11}^{[x \rightarrow m]} & 0 & 0 \\ \beta_{1\gamma}^{[x \rightarrow m]} & \beta_\gamma^{[x \rightarrow m]} & 0 \\ \beta_{12}^{[x \rightarrow m]} & \beta_\gamma^{[x \rightarrow m]} & \beta_{22}^{[x \rightarrow m]} \end{pmatrix}; \mathbf{B}^{[x \rightarrow y]} = \begin{pmatrix} \beta_{11}^{[x \rightarrow y]} & 0 & 0 \\ \beta_{1\gamma}^{[x \rightarrow y]} & \beta_\gamma^{[x \rightarrow y]} & 0 \\ \beta_{12}^{[x \rightarrow y]} & \beta_\gamma^{[x \rightarrow y]} & \beta_{22}^{[x \rightarrow y]} \end{pmatrix}$ $\mathbf{B}^{[m \rightarrow y]} = \begin{pmatrix} \beta_{11}^{[m \rightarrow y]} & 0 & 0 \\ \beta_{1\gamma}^{[m \rightarrow y]} & \beta_\gamma^{[m \rightarrow y]} & 0 \\ \beta_{12}^{[m \rightarrow y]} & \beta_\gamma^{[m \rightarrow y]} & \beta_{22}^{[m \rightarrow y]} \end{pmatrix}$