Revisiting the electronic properties of disclinated graphene sheets

Nicolás Fernández^{1*}, Pierre Pujol², Mario Solís^{3,4} and Teófilo Vargas⁵

¹Facultad de Ciencias Físicas, Universidad Nacional Mayor de San Marcos, Avenida Venezuela s/n Cercado de Lima, Lima, 15081, Lima, Perú.

²Laboratoire de Physique Théorique, CNRS and Université de Toulouse, UPS, Toulouse, F-31062, France.

³Centro Atómico Bariloche, CNEA and CONICET, S. C. de Bariloche, R8402AGP, Rio Negro, Argentina.

⁴Instituto Balseiro, UNCuyo, S. C. de Bariloche, R8402AGP, Rio Negro, Argentina. ⁵Grupo de Física Teórica GFT and Grupo de Astronomía SPACE, Universidad Nacional Mayor de San Marcos, Avenida Venezuela s/n Cercado de Lima, Lima, 15081, Lima, Perú.

*Corresponding author(s). E-mail(s): nicolas.fernandez@unmsm.edu.pe; Contributing authors: pierre.pujol@irsamc.ups-tlse.frmar; mario.solis@ib.edu.ar; tvargasa@unmsm.edu.pe;

Abstract

The interplay between topological defects, such as dislocations or disclinations, and the electronic degrees of freedom in graphene has been extensively studied. In the literature, for the study of this kind of problems, it is in general used either a gauge theory or a curved spatial Riemannian geometry approach, where, in the geometric case, the information about the defects is contained in the metric and the spin-connection. However, these topological defects can also be associated to a Riemann-Cartan geometry where curvature and torsion plays an important role. In this article we study the interplay between a wedge disclination in a planar graphene sheet and the properties of its electronic degrees of freedom. Our approach relies in its relation with elasticity theory through the so called elastic-gauge, where their typical coefficients, as for example the Poisson's ratio, appear directly in the metric, and consequently also in the electronic spectrum.

Keywords: Graphene, Topological defects, Dislocations, Disclinations.

1 Introduction

Graphene has received much attention in recent years due to its elastic and electronic properties, which offer key industrial impact in nanoelectronic devices [1]. Since its electronic band structure was studied by Wallace and Semenoff [2, 3], and then produced by Novoselov et. al. [4], a very wide variety of theoretical and experimental work has been conducted. Among them, one can mention the study of the electronic properties and how the electrons interact with, for example, an electromagnetic field, topological defects, disorder or impurities. The most important feature of graphene is the linear energy spectrum at two Dirac points in the Brillouin zone, which provide a low energy description with massless Dirac fermions. Another interesting result comes from the study of material irregularities, which have dramatic effects on its electronic properties. The topological defects give rise to interesting modifications of the electronic properties. They are classified into two general types: disclinations and dislocations [5, 6]. From the atomistic point of view, a disclination adds or removes one atom in the lattice, and a dislocation can be viewed as a pair of disclinations. If one considers the system as a continuum medium, it can be studied using elasticity theory [7]. In this approach, one can add or remove a piece of material and then glue the new boundaries. This procedure is known as Volterra's process [8]. In both cases, the key quantities that specify these irregularities are the Burgers and Franck vectors: they express the deviation with respect to a defect-free configuration. The presence of these defects introduces an extra field which interacts and modifies the free electronic dynamics. This new field enter in the electron physics either through a gauge field or through a geometric approach. In the first case, the gauge field describes the modifications of the electronic degrees of freedom due to the presence of disclinations or dislocations, and, in the continuum limit, it acts as an point flux producing an Aharonov-Bohm effect [9]. This approach was developed extensively in the last three decades in the context of fullerenes and graphite. The insightful studies of Gonzalez et. al. [10, 11] demostrated that the folded carbon lattice mimics a gauge field. An interesting application to graphite system was developed by Lammert and Crespi [12], where they studied graphite in the presence of a disclination. They viewed this lattice as a graphite cone where the disclination is at its apex. Within this approach they calculated the LDOS (local density of states); similar results were previously reported [13] within a tight-binding formulation (see [14] and references therein for a summary of these techniques and applications), and also in the presence of a magnetic field. In the second case, the topological defects were considered sources of geometric distortions, and these distortions produce a non-trivial geometry background in which the fermionic dynamics develop: the euclidean flat metric is promoted to a curved one.

The presence of defects (dislocations or disclinations) produces a singularity point in the manifold which is phenomenologically characterized by a conical metric (see for i.e. [15]). This metric can also be related to a cosmic string [16] in highenergy physics. The conical metric provides the fermions with a spin connection, which plays the role of the gauge field in the previously mentioned approach. In the case of graphene, the application of this technique was reported by Osipov and Kochetov [17], where they obtained the solution of the massless Dirac equation in the conical geometry and its LDOS. They reported that there are no extended states with non-zero density of states, and their results showed a different power energy dependence with respect to [12]. Similar studies were done by Vozmediano et. al. [18] where the authors computed the LDOS, or Cortijo et. al. [19] who solved the Dirac equation for ripples in graphene. However, much of this studies were focused in the interplay between the fermionic dynamics and the geometry instead of the interplay between electronic-elastic properties, or how the previous results in elasticity theory [7] are connected in this approach. An important progress in geometric-elastic properties was achieved by Katanaev and Volovich [20] (see [21] for a review). They relates the Burgers and Franck vectors with torsion and curvature, respectively. The inclusion of torsion was previously mentioned long time ago [22–24]. Disclinations carry curvature but not torsion, while dislocations carry torsion but not curvature. In their approach, the Riemann-Cartan geometry is used, where the curvature and torsion are non-trivial. The final ingredient related to the elastic properties was finally realized in [25], where the authors implemented an elastic-gauge to derive the elastic wedge disclinated metric from a conical metric [7]. The importance of this gauge lies in its explicit dependence on the elastic constants. With this ideas, the interplay between fermions (in graphene) was studied by Furtado et. al. [26, 27]. They also included the interaction with an external magnetic field and determined the electronic spectrum. However, the consequences of the elastic-gauge in graphene were not previously reported. As we show below, the elastic-gauge is responsible of an important change in the electronic behavior. In this work, we study the electronic dynamics within the Katanaev-Volovich presence of a wedge disclination. We will focus on how its electronic dynamics is changed by the presence of a disclination. Our major goal is the interesting interplay between the electronic and elastic properties that appear in the spectrum. We also compute the LDOS and we find a different power dependence than the one obtained in previous works which focused on a conical shape [12, 17].

2 Wedge disclination and its associated metric

In the geometric approach of defects [21], the flat Minkowskian space-time metric is replaced by a curved metric in a Riemann-Cartan manifold; the line element is

$$ds^{2} = g_{\mu\nu}dx^{\mu}dx^{\nu} = dt^{2} - g_{ij}dx^{i}dx^{j}, \quad (1)$$

where μ, ν , are the space-time indices and g_{ij} is the spatial curved metric in euclidean signature (+,+). The temporal component of the metric is flat indicating that the topological defects are not time-dependent. In this perspective, the system is characterized by the curvature and torsion tensors

$$R_{\mu\nu}{}^{ab} = \partial_{\mu}\omega_{\nu}{}^{ab} - \partial_{\nu}\omega_{\mu}{}^{ab} + \omega_{\mu}{}^{a}_{c}\omega_{\nu}{}^{cb} - \omega_{\nu}{}^{a}_{c}\omega_{\mu}{}^{cb},$$

$$T_{\mu\nu}{}^{a} = \partial_{\mu}h_{\nu}{}^{a} - \partial_{\nu}h_{\mu}{}^{a} + \omega_{\mu}{}^{b}_{b}h_{\nu}{}^{b} - \omega_{\nu}{}^{b}_{b}h_{\mu}{}^{b},$$
(2)

where a, b are tangent-space indices, $\omega_{\mu}{}^{ab}$ is the spin-connection and $h_{\mu}{}^{b}$ is an associated tetrad. According to the geometric approach [20, 21, 25], the Burgers and Franck vectors are associated to

$$b^{a} = \int_{S} dx^{\mu} \wedge dx^{\nu} T_{\mu\nu}{}^{a}, \qquad (3)$$

$$\Omega^{ab} = \int_S dx^{\mu} \wedge dx^{\nu} R_{\mu\nu}{}^{ab}, \qquad (4)$$

in this context, the presence of a non-trivial torsion indicates the presence of dislocation while the curvature is associated to disclinations. The Action for topological defects was obtained in [20, 21], where in the absence of disclinations, the curvature vanish. In this case, the Riemann-Cartan geometry is reduced to a space of absolute parallelism giving an analogue of teleparallel gravity. In this theory, the Riemann curvature is determined by the torsion tensor which depends on the Weitzenböck connection (see [28, 29] and references therein for more details). In the rest of the paper we use the notation of [28], where \circ , • denote the quantities given by the Levi-Civita and Weitzenböck connection, respectively. The equation of motion for geometry in the presence of a disclination is

$$\overset{\circ}{R}_{\mu\nu} - \frac{1}{2}g_{\mu\nu}\overset{\circ}{R} = \kappa\Theta_{\mu\nu}\,,\qquad(5)$$

where κ is a constant which makes the equation well defined from a dimensional point of view 1 and $\Theta_{\mu\nu}$ is the energy-momentum tensor. The only non-vanishing component of the energymomentum tensor for a disclination at the origin is

$$\Theta_{zz} = \frac{2\pi}{\sqrt{g}} \theta \delta^2(x) \,, \tag{6}$$

Which depend explicitly on the deficit angle θ . The Dirac delta in eq. (6) produce a singularity in the geometry. Following the same steps of [20, 21], the metric, written in cylindrical coordinates, is given by

$$g_{ij}dx^i dx^j = d\tilde{r}^2 + \alpha^2 \tilde{r}^2 d\varphi^2 \,, \tag{7}$$

where $0 \leq \tilde{r} < \infty$ and $0 \leq \varphi < 2\pi$. The parameter α is related with the deficit angle θ of disclination as $\alpha = 1 + \theta$. One way to see how the metric in eq. (7) arise because of the wedge disclination is to consider the Volterra's process [8]. First, consider the system as a disk, then remove a wedge piece of the material with angle θ . After that, glue the two sides while maintaining the system flat, and the disclination core is located at the center of the disk. An important point that deserves a clarification is that, in previous works, the metric used for a system with conical geometry do not contain information about the elasticity parameters [7]. This gave rise to the proposal of Katanaev [25] to use what he called an a elastic-gauge, in which the elasticity parameters appear explicitly in the metric. For the planar geometry we consider here we follow this route and use the metric that contains the elasticity parameters. With this choice, we find

$$\tilde{r} = \frac{r^{\gamma}}{\tilde{\gamma}R^{\gamma-1}} \equiv \rho \,, \tag{8}$$

¹In gravity context is $\kappa = 8\pi G_N$.

where R is the radii of the disk (or cylinder base) and

$$\gamma = \frac{-\theta\sigma + \sqrt{\theta^2 \sigma^2 + 4(1+\theta)(1-\sigma)}}{2(1-\sigma)}, \qquad (9)$$

here the Poisson's ratio is denoted by σ . As a result, the elastic-gauge relates the general conical metric (7) with a specific conical metric containing information of the elastic properties of the sample and reproduce a well known result [7] in elasticity theory. Hence, using eq. (8) in (7), we find

$$g_{ij}dx^i dx^j = d\rho^2 + \alpha^2 \rho^2 d\varphi^2 \,, \tag{10}$$

with $0 \le r < R$, $0 \le \varphi < 2\pi$, $\alpha = 1 + \theta$, where we used the variable ρ defined in (8). Therefore, the complete metric (1) is

$$ds^2 = dt^2 - d\rho^2 - \alpha^2 \rho^2 d\varphi^2 , \qquad (11)$$

where $0 < \rho \leq R/\gamma$, $0 < \varphi < 2\pi$, $\alpha = 1 + \theta$ and ρ , γ are given by eqs. (8) and (9) respectively. This result is extremely important for the next sections, as it modifies the electronic spectrum and LDOS as we show below.

3 Wedge disclinations and Dirac equation

The metric in eq. (11) provides a natural description of the considered defect. We now can turn to the study of the electronic degrees of freedom in an elastic medium with a wedge disclination from the perspective of the Weitzenböck geometry. In this perspective we introduce the Dirac equation in the farmework of the teleparallel Fock-Ivanenko covariant derivative [28, 29]. The equation we want to study is:

$$i\gamma^{\mu} \mathcal{D}_{\mu} \psi - m\psi = 0, \qquad (12)$$

where we set $\hbar = v_F = 1$. Here,

$${}^{\bullet}\mathcal{D}_{\mu} = \partial_{\mu} + \frac{i}{2} {}^{\bullet}K_{\mu}{}^{ab}S_{ab} , \qquad (13)$$

where $\overset{\bullet}{K}_{\mu}{}^{ab}$ is the contorsion tensor, $S_{ab} = \frac{i}{4} [\gamma_a, \gamma_b]$ is the Lorentz spin-1/2 generator, and $\gamma^{\mu} = h^{\mu}{}_a \gamma^a$ are the curved Dirac matrices in

terms of a nontrivial tetrad field $h^{\mu}{}_{a}$. The spacetime and the tangent-space metrics are related by $g_{\mu\nu} = \eta_{ab} h^{a}{}_{\mu} h^{b}{}_{\nu}$ and the contorsion tensor can be determined by the following relation

$${}^{\bullet}_{K_{\mu}}{}^{ab} = h^{a}{}_{\rho}{}^{\bullet}_{K^{\rho}}{}_{\mu\nu}h^{\ \nu}_{c}\eta^{cb}\,, \qquad (14)$$

with $\overset{\bullet}{K}{}^{\rho}{}_{\mu\nu} = \overset{\bullet}{\Gamma}{}^{\rho}{}_{\mu\nu} - \overset{\circ}{\Gamma}{}^{\rho}{}_{\mu\nu}$, where $\overset{\bullet}{\Gamma}{}^{\rho}{}_{\mu\nu} = h_a{}^{\rho}\partial_{\mu}h^a{}_{\nu}$ is the Weitzenböck connection and $\overset{\circ}{\Gamma}{}^{\rho}{}_{\mu\nu} = \frac{1}{2}g^{\sigma\rho} \left(\partial_{\mu}g_{\rho\nu} + \partial_{\nu}g_{\rho\mu} - \partial_{\rho}g_{\mu\nu}\right)$ is the Levi-Civita connection of the Riemannian geometry. The Dirac equation (12) can be written in the Schrödinger's form as

$$i\partial_t \psi = \left(\gamma^t m - i\gamma^t \gamma^i \partial_i - i\gamma^t \gamma^i \tilde{K}^{ab}_{\ i} \ S_{ab}\right) \psi \,. \tag{15}$$

The Dirac matrices obey the Clifford's algebra

$$\{\gamma^{\mu}, \gamma^{\nu}\} = 2g^{\mu\nu}, \qquad (16)$$

and can be chosen to be the Pauli matrices $\gamma^t = \gamma^0 = \sigma^3$, $\gamma^1 = i\sigma^2$ and $\gamma^2 = -i\sigma^1$, or explicitly

$$\gamma^{t} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \gamma^{1} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \gamma^{2} = \begin{pmatrix} 0 & -i \\ -i & 0 \\ (17) \end{pmatrix}$$

with this choice we find: $\overset{\bullet}{K}{}^{12}{}_{\varphi} = 1 - \alpha$ and $S_{12} = \frac{1}{2}\sigma^3$. Therefore, the Dirac Hamiltonian is given by

$$H = \gamma^{t} m - i\gamma^{t} \Gamma(\varphi) \partial_{\rho} - i\gamma^{t} \Gamma'(\varphi) \frac{\partial_{\varphi}}{\alpha \rho} + \frac{1 - \alpha}{2\alpha \rho} \gamma^{t} \Gamma'(\varphi) \sigma^{3},$$
(18)

where $\Gamma(\varphi) = \gamma^1 \cos(\varphi) + \gamma^2 \sin(\varphi)$ and $\Gamma'(\varphi) = -\gamma^1 \sin(\varphi) + \gamma^2 \cos(\varphi)$, as shorthand notation. Inspired in the result of [30], in the process of a parallel transport of a spinor along a closed trajectory C around the defect, the holonomy matrix $U(C) = \exp\left(-\oint_C \Gamma_\mu(x)dx^\mu\right)$, being Γ_μ is the spinorial connection, provide the quantum phase acquired by the wave function, i.e., $\psi = U(C)\psi_0$, and the Hamiltonian becomes

$$H = \gamma^t m - i\gamma^t \Gamma(\varphi) \partial_\rho - i\gamma^t \Gamma'(\varphi) \frac{\partial_\varphi}{\alpha \rho} \,. \tag{19}$$

Noticing that the total angular momentum $J = -i\partial_{\varphi} + \frac{1}{2}\sigma^3$ commutes with the Hamiltonian (19), the eigenfunctions are classified by the eigenvalues $j = l \pm 1/2, \ l = 0, \pm 1, \pm 2, \ldots$ We can therefore

analyze the stationary state of the Dirac equation using the following anzatz

$$\psi_0 = e^{-iEt} \begin{pmatrix} u(\rho)e^{il\varphi} \\ v(\rho)e^{i(l+1)\varphi} \end{pmatrix}, \qquad (20)$$

which give the following system of differential equations

$$(E - m)u(\rho) = -i\partial_{\rho}v(\rho) - i\frac{l+1}{\alpha\rho}v(\rho), \quad (21)$$
$$(E + m)v(\rho) = -i\partial_{\rho}u(\rho) + i\frac{l}{\alpha\rho}u(\rho). \quad (22)$$

In the following, we focus on the case of massless chiral fermion in order to describes the low energy electronic degrees of freedom of graphene.

4 Massless fermions in disclinated planar Graphene

We now specify our previous system of differential equations on massless fermions, which is the relevant case for the description of graphene. Solving for $u(\rho)$ we get

$$u(\rho) = C_1 \rho^{\xi} J_{\eta}(E\rho) , \qquad (23)$$

where we consider only the Bessel function of the first kind as it is the only one finite at the origin. In this section we use $\xi = \frac{\alpha - 1}{2\alpha}$ and $\eta = \pm \frac{\alpha - 2l - 1}{2\alpha}$. Then $v(\rho)$ is given by

$$\mathbf{v}(\rho) = \pm i C_1 \rho^{\xi} J_{\tilde{\eta}}(E\rho) \,, \tag{24}$$

where C_1 is a constant to be determined below, $\tilde{\eta} = \pm (\eta + 1)$ and the signs \pm corresponds to positive and negative energy solutions, respectively. Hence, there are two independent solutions for each energy state, one for $\eta, \tilde{\eta} > 0$ and another for $\eta, \tilde{\eta} < 0$. Now we focus in case of positive energy and we want to obtain the constant C_1 . The normalization condition is written as

$$\int \left(|u(\rho)|^2 + |\mathbf{v}(\rho)|^2 \right) \sqrt{-g} \, d\rho \, d\varphi = 1 \,, \qquad (25)$$

where g is the determinant of the metric. Using the equations (23) and (24) find

$$2\pi\alpha C_1^2 \int_0^{R/\gamma} \rho^{2\xi+1} \left(J_\eta^2(E\rho) + J_{\eta+1}^2(E\rho) \right) d\rho = 1,$$
(26)

and we obtain

$$C_1 = \frac{1}{E^{\eta}} \sqrt{f(\eta, \xi, R, \gamma)} \,_2 F_3(\zeta) \,, \qquad (27)$$

where $f(\eta, \xi, R, \gamma)$ is a constant whose dependence on the parameters do not need to be made explicit here. The $_2F_3(\zeta)$ is an Hyper-geometric function of arguments $\zeta = \{\eta + 1 + \xi, \eta + 1/2; 2\eta + 1, \eta + 1, 2 + \eta + \xi; -(R/\gamma)^2 E^2\}$. For fixed values of Eand taking $R \to \infty$ we get

$$C_1 \propto E^{1/2} \left(\frac{\gamma}{R}\right)^{\xi + \frac{1}{2}} \tag{28}$$

Reanalyzing the normalization condition (26) for $E\rho \ll 1$, we find

$$\int_{0}^{R/\gamma} \rho^{2\xi+1} \left(\left(\frac{E\rho}{2}\right)^{2\eta} + \left(\frac{E\rho}{2}\right)^{2\tilde{\eta}} \right) d\rho = 1,$$
(29)

The integrability of the previous expression induce restrictions in the values of the angular momentum quantum number as: $l < \min(2\theta+1, 3\theta+2)$ for $\eta, \tilde{\eta} > 0$ and $l > \max(-1 - \theta, 0)$ for $\eta, \tilde{\eta} < 0$. Furthermore, the parameter α is bounded by $0 < \alpha < 1$, because of the removal of a part of the material. To study the wave function near the center of the disk, it is useful to consider small arguments in (23), (24), i.e., with fixed values of E and taking $\rho \to 0$. For $\eta, \tilde{\eta} > 0$, the dependence on E and ρ is

$$\begin{pmatrix} u(\rho) \\ v(\rho) \end{pmatrix} \sim \begin{pmatrix} E^{1/2+\eta} \rho^{\xi+\eta} \\ E^{1/2+\tilde{\eta}} \rho^{\xi+\tilde{\eta}} \end{pmatrix}, \qquad (30)$$

and for $\eta, \tilde{\eta} < 0$,

$$\begin{pmatrix} u(\rho) \\ v(\rho) \end{pmatrix} \sim \begin{pmatrix} E^{1/2-\eta} \rho^{\xi-\eta} \\ E^{1/2-\tilde{\eta}} \rho^{\xi-\tilde{\eta}} \end{pmatrix}.$$
 (31)

We can express the above results in terms of the deficit angle θ , we obtain

$$\begin{pmatrix} u(\rho) \\ v(\rho) \end{pmatrix} \sim \begin{pmatrix} E^{\frac{1+2\theta-2l}{2+2\theta}} \rho^{\frac{\theta-l}{1+\theta}} \\ E^{\frac{3+4\theta-2l}{2+2\theta}} \rho^{\frac{2\theta-l+1}{1+\theta}} \end{pmatrix}, \quad \eta, \tilde{\eta} > 0,$$
(32)

and

$$\begin{pmatrix} u(\rho) \\ v(\rho) \end{pmatrix} \sim \begin{pmatrix} E^{\frac{1+2l}{2+2\theta}} \rho^{\frac{l}{1+\theta}} \\ E^{\frac{-2\theta+2l-1}{2+2\theta}} \rho^{\frac{-\theta+l-1}{1+\theta}} \end{pmatrix}, \quad \eta, \tilde{\eta} < 0.$$
(33)

In particular, with the meaningful values for the angle $\theta = 0, -1/6, -1/3$, we get l = 0 and, therefore, $\psi \sim E^{1/2}, \psi \sim E^{2/5}\rho^{-1/5}$, and $\psi \sim E^{1/4}\rho^{-1/2}$, respectively. In the same way, for $\theta = 1/6, 1/3, 1/2$, we get l = 1 and $\psi \sim E^{2/7}\rho^{-1/7}$, $\psi \sim E^{1/8}\rho^{-1/4}$ and $\psi \sim \rho^{-1/3}$, respectively.

In order to compute the local density of states, we notice that it diverges for $\rho \to 0$. This divergence can be regularized by considering the total density of states on a small disk $0 < \rho \leq \delta$, as was done in [17] and [12]. The local density of states is defined as

$$\rho(\mathbf{r}, E) = \sum_{i} |\psi_i(\mathbf{r})|^2 \delta(E - E_i), \qquad (34)$$

thus, we calculate the total density of states as

$$DoS = \left(\int_0^{\frac{\delta^{\gamma}}{\gamma R^{\gamma-1}}} u(\rho) d\rho\right)^2 + \left(\int_0^{\frac{\delta^{\gamma}}{\gamma R^{\gamma-1}}} v(\rho) d\rho\right)^2,$$
(35)

and we find the dependence for E and δ as

$$DoS \propto \begin{cases} E^{\frac{4\theta - 2l + 3}{1 + \theta}} \delta^{\gamma \frac{6\theta - 2l + 4}{1 + \theta}} & \eta, \tilde{\eta} > 0, \\ E^{\frac{-2\theta + 2l - 1}{1 + \theta}} \delta^{\gamma \frac{2l}{1 + \theta}} & \eta, \tilde{\eta} < 0. \end{cases}$$
(36)

In the leading order for equation (36), we obtain

$$DoS \propto \begin{cases} E^{1/2} \delta^{\gamma} , & \theta = -1/3, \quad l = 0 , \\ E^{4/5} \delta^{8\gamma/5} , & \theta = -1/6, \quad l = 0 , \\ E \delta^{2\gamma} , & \theta = 0, \quad l = 0 , \\ E^{4/7} \delta^{12\gamma/7} , & \theta = 1/6, \quad l = 1 , \\ E^{1/4} \delta^{3\gamma/2} , & \theta = 1/3, \quad l = 1 , \\ \delta^{4\gamma/3} , & \theta = 1/2, \quad l = 1 , \end{cases}$$
(37)

for values $\theta = 0, -1/6, -1/3$ of the deficit angle. In the inextensional limit ($\sigma = 1/2$, i.e., rigid membrane), our results are in agreement with previous works, [17] and [12]. At last, we study the discretization of spectrum because of the finite size system. We consider the billiard boundary condition [31] with an infinite potential at the boundary $\rho = R/\gamma$ assuming a circular contour, i.e., with $\varphi = \alpha$ at the boundary. Such a condition for the spinor components can be written as

$$\frac{\psi_2}{\psi_1} = ie^{i\varphi}, \qquad (38)$$

where $\psi_1 = C_1 \rho^{\xi} J_{\eta}(ER/\gamma) e^{il\varphi}$ and $\psi_2 = iC_1 \rho^{\xi} J_{\eta+1}(ER/\gamma) e^{i(l+1)\varphi}$, so (38) becomes

$$J_{\eta}(ER/\gamma) = J_{\eta+1}(ER/\gamma), \qquad (39)$$

Considering the asymptotic behavior of the Bessel functions for $ER/\gamma \gg 1$,

$$J_{\eta}(ER/\gamma) \approx \sqrt{\frac{2}{\pi ER/\gamma}} \cos\left(E\rho - \frac{\eta\pi}{2} - \frac{\pi}{4}\right),$$
(40)

The boundary condition impose

$$\cos\left(ER/\gamma - \frac{\eta\pi}{2} - \frac{\pi}{4}\right) = \sin\left(ER/\gamma - \frac{\eta\pi}{2} - \frac{\pi}{4}\right) + \frac{\pi}{4}$$

hence the argument must be equal to $\pi/4 + n$. This give us the spectrum in the presence of wedge disclination as

$$E = \frac{\pi\gamma}{R} \left(n + \frac{1}{2} + \left| \frac{\alpha - 2l - 1}{4\alpha} \right| \right) \,. \tag{42}$$

It is important to point out that the result depends on the elasticity properties through the definition of γ , given in equation (9), with an explicit dependence on the Poisson's ratio. This last result is to put in perspective with the remark that when a wedge of material is removed, the electrons spread over a smaller area than the one available in a material without defects, whereas, when a wedge of material is added, the electrons spread over a larger area. Geometrically, these correspond to positive and negative curvature surfaces respectively.

5 Conclusions

In this work we showed how the fermionic dynamics in graphene is affected by a wedge disclination. We used the geometric approach developed in [21] and implemented the line element of a wedge disclination found in elasticity theory [7, 21]within the elastic-gauge [25]. This choice of metric corresponds to the case where the graphene sheet is kept planar despite the fact of a presence of a disclination. Our main results are the LDOS in equation (37), where we found a different power dependence in radius δ than in previous works made on a conical geometry [12, 17] and the finite size spectrum in equation (42) was obtained using the billiard condition. A similar result was reported in [26] where the authors studied the Landau's levels. However, it is important to stress that in this work, our results contain the explicit presence of the Poisson's ratio, which appear for example in eq. (42), a result directly arising from the planar geometry which manifest itself through the so called elastic-gauge. As a consequence, the result for the spectrum, eq. (42) show the explicitly interplay between the electronic and elastic properties in a disclinated graphene sheet. Our results may be verified in an experimental set-up and provide, in principle, a nondestructive manner to compute the Poisson's ratio. We are confident that this analysis can be extended to a graphene sheet with several disclinations with equal or different deficit angles, and the study of the Landau levels of graphene with disclinations.

Acknowledgments

The authors would like the thank the LA CoNGA Physics program which was at the origin of this collaboration. M. S. is supported by CNEA, Conicet and UNCuyo-Instituto Balseiro. T. V. is supported by the Vice-rectorate for Research and Postgraduate B20131101 and CONCYTEC through the Research Academic grant. M. S. would like to thank the Facultad de Ciencias Físicas, Universidad Nacional Mayor de San Marcos for its hospitality during the first stage of this project.

Declarations

Author contribution

All authors contributed to the study equally.

Data Availability Statement

No Data associated in the manuscript.

References

- A.C. Neto, F. Guinea, N.M. Peres, K.S. Novoselov, A.K. Geim, The electronic properties of graphene. Reviews of modern physics 81(1), 109 (2009)
- [2] P.R. Wallace, The band theory of graphite. Physical review 71(9), 622 (1947)
- [3] G.W. Semenoff, Condensed-matter simulation of a three-dimensional anomaly. Physical Review Letters 53(26), 2449 (1984)
- [4] K.S. Novoselov, A.K. Geim, S.V. Morozov, D.e. Jiang, Y. Zhang, S.V. Dubonos, I.V. Grigorieva, A.A. Firsov, Electric field effect in atomically thin carbon films. science **306**(5696), 666–669 (2004)
- [5] W.T. Read, W. Shockley, Dislocation models of crystal grain boundaries. Physical review 78(3), 275 (1950)
- [6] W.F. Harris, Disclinations. Scientific American 237(6), 130–145 (1977)
- [7] L.D. Landau, E.M. Lifshitz, A.M. Kosevich, L.P. Pitaevskii, *Theory of elasticity: volume* 7, vol. 7 (Elsevier, 1986)
- [8] R.A. Puntigam, H.H. Soleng, Volterra distortions, spinning strings, and cosmic defects.

Classical and Quantum Gravity 14(5), 1129 (1997)

- [9] K. Kawamura, Scattering of electrons due to screw dislocations. Zeitschrift f
 ür Physik B Condensed Matter 30(1), 1–11 (1978)
- [10] J. González, F. Guinea, M.A.H. Vozmediano, Continuum approximation to fullerene molecules. Physical review letters 69(1), 172 (1992)
- [11] J. Gonzalez, F. Guinea, M.A. Vozmediano, The electronic spectrum of fullerenes from the dirac equation. Nuclear Physics B 406(3), 771–794 (1993)
- [12] P.E. Lammert, V.H. Crespi, Topological phases in graphitic cones. Physical review letters 85(24), 5190 (2000)
- [13] R. Tamura, M. Tsukada, Disclinations of monolayer graphite and their electronic states. Physical Review B 49(11), 7697 (1994)
- M.A. Vozmediano, M. Katsnelson, F. Guinea, Gauge fields in graphene. Physics Reports 496(4-5), 109–148 (2010)
- [15] K. Tod, Conical singularities and torsion. Classical and Quantum Gravity 11(5), 1331 (1994)
- [16] A. Vilenkin, Gravitational field of vacuum domain walls and strings. Physical Review D 23(4), 852 (1981)
- [17] V.A. Osipov, E.A. Kochetov, Dirac fermions on graphite cones. Journal of Experimental and Theoretical Physics Letters 73(10), 562– 565 (2001)
- [18] M.A. Vozmediano, F. de Juan, A. Cortijo, in *Journal of Physics: Conference Series*, vol. 129 (IOP Publishing, 2008), p. 012001
- [19] A. Cortijo, M.A. Vozmediano, Effects of topological defects and local curvature on the electronic properties of planar graphene. Nuclear Physics B 763(3), 293–308 (2007)

- [20] M. Katanaev, I. Volovich, Theory of defects in solids and three-dimensional gravity. Annals of Physics 216(1), 1–28 (1992)
- [21] M.O. Katanaev, Geometric theory of defects. Physics-Uspekhi 48(7), 675 (2005)
- [22] K. Kondo, in Proc. 2nd Japan Nat. Congr. Applied Mechanics, vol. 2 (1952), pp. 41–47
- [23] B.A. Bilby, R. Bullough, E. Smith, Continuous distributions of dislocations: a new application of the methods of non-riemannian geometry. Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 231(1185), 263–273 (1955)
- [24] E. Kröner, Continuum theory of defects, in" physics of defects" (les houches, session 35), ed. r. balian et al. ed: North-Holland, Amsterdam (1980)
- [25] M.O. Katanaev, Wedge dislocation in the geometric theory of defects. Theoretical and mathematical physics 135(2), 733–744 (2003)
- [26] C. Furtado, B.G. da Cunha, F. Moraes, E.B. de Mello, V. Bezzerra, Landau levels in the presence of disclinations. Physics Letters A 195(1), 90–94 (1994)
- [27] C. Furtado, F. Moraes, On the binding of electrons and holes to disclinations. Physics Letters A 188(4-6), 394–396 (1994)
- [28] H.I. Arcos, J.G. Pereira, Torsion gravity: a reappraisal. International Journal of Modern Physics D 13(10), 2193–2240 (2004)
- [29] S. Bahamonde, K. Dialektopoulos, C. Escamilla-Rivera, G. Farrugia, V. Gakis, M. Hendry, M. Hohmann, J. Said, J. Mifsud, E. Di Valentino, Teleparallel gravity: from theory to cosmology. Reports on Progress in Physics (2022)
- [30] C. Furtado, F. Moraes, A.d.M. Carvalho, Geometric phases in graphitic cones. Physics Letters A 372(32), 5368–5371 (2008)
- [31] M.V. Berry, R. Mondragon, Neutrino billiards: time-reversal symmetry-breaking