

H_I filaments as potential compass needles? Comparing the magnetic field structure of the Small Magellanic Cloud to the orientation of GASKAP-H_I filaments

Y. K. Ma,^{1,2}★ N. M. McClure-Griffiths,¹ S. E. Clark,^{3,4} S. J. Gibson,⁵ J. Th. van Loon,⁶
J. D. Soler,⁷ M. E. Putman,⁸ J. M. Dickey,⁹ M. -Y. Lee,¹⁰ K. E. Jameson,¹¹ L. Uscanga,¹²
J. Dempsey,^{1,13} H. Dénes,^{14,15} C. Lynn,¹ and N. M. Pingel¹⁶

¹Research School of Astronomy & Astrophysics, Australian National University, Canberra, ACT 2611, Australia

²Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

³Department of Physics, Stanford University, Stanford, CA 94305, USA

⁴Kavli Institute for Particle Astrophysics & Cosmology, P.O. Box 2450, Stanford University, Stanford, CA 94305, USA

⁵Department of Physics and Astronomy, Western Kentucky University, 1906 College Heights Blvd., Bowling Green, KY 42101, USA

⁶Lennard-Jones Laboratories, Keele University, ST5 5BG, UK

⁷Istituto di Astrofisica e Planetologia Spaziali (IAPS), INAF, Via Fosso del Cavaliere 100, 00133 Roma, Italy

⁸Department of Astronomy, Columbia University, New York, NY 10027, USA

⁹School of Natural Sciences, Private Bag 37, University of Tasmania, Hobart, TAS 7001, Australia

¹⁰Korea Astronomy and Space Science Institute, 776 Daedeok-daero, Yuseong-gu, Daejeon 34055, Republic of Korea

¹¹Owens Valley Radio Observatory, California Institute of Technology, Big Pine, CA 93513, USA

¹²Departamento de Astronomía, Universidad de Guanajuato, A.P. 144, Guanajuato 36000, Guanajuato, Mexico

¹³CSIRO Information Management and Technology, GPO Box 1700, Canberra, ACT 2601, Australia

¹⁴School of Physical Sciences and Nanotechnology, Yachay Tech University, Hacienda San José S/N, 100119, Urcuquí, Ecuador

¹⁵ASTRON - The Netherlands Institute for Radio Astronomy, 7991 PD Dwingeloo, The Netherlands

¹⁶Department of Astronomy, University of Wisconsin-Madison, 475 North Charter Street, Madison, WI 53706-15821, USA

Accepted 2023 February 07. Received 2023 February 05; in original form 2022 December 01

ABSTRACT

High-spatial-resolution H_I observations have led to the realisation that the nearby (within few hundreds of parsecs) Galactic atomic filamentary structures are aligned with the ambient magnetic field. Enabled by the high quality data from the Australian Square Kilometre Array Pathfinder (ASKAP) radio telescope for the Galactic ASKAP H_I (GASKAP-H_I) survey, we investigate the potential magnetic alignment of the $\gtrsim 10$ pc-scale H_I filaments in the Small Magellanic Cloud (SMC). Using the Rolling Hough Transform (RHT) technique that automatically identifies filamentary structures, combined with our newly devised ray-tracing algorithm that compares the H_I and starlight polarisation data, we find that the H_I filaments in the northeastern end of the SMC main body (“Bar” region) and the transition area between the main body and the tidal feature (“Wing” region) appear preferentially aligned with the magnetic field traced by starlight polarisation. Meanwhile, the remaining SMC volume lacks starlight polarisation data of sufficient quality to draw any conclusions. This suggests for the first time that filamentary H_I structures can be magnetically aligned across a large spatial volume ($\gtrsim$ kpc) outside of the Milky Way. In addition, we generate maps of the preferred orientation of H_I filaments throughout the entire SMC, revealing the highly complex gaseous structures of the galaxy likely shaped by a combination of the intrinsic internal gas dynamics, tidal interactions, and star formation feedback processes. These maps can further be compared with future measurements of the magnetic structures in other regions of the SMC.

Key words: ISM: magnetic fields – ISM: structure – galaxies: ISM – Magellanic Clouds – galaxies: magnetic fields – radio lines: ISM

1 INTRODUCTION

The μ G-strength magnetic fields in galaxies affect nearly all aspects of galactic astrophysics (e.g., Beck & Wielebinski 2013; Beck 2016), including the propagation of cosmic rays (Aab et al. 2015; Seta et al.

2018), the rate at which stars form (Price & Bate 2008; Federrath & Klessen 2012; Birnboim et al. 2015; Krumholz & Federrath 2019), the stellar initial mass function (Krumholz & Federrath 2019; Sharda et al. 2020; Mathew & Federrath 2021), the large-scale gas dynamics (Beck et al. 2005; Kim & Stone 2012), and possibly even the rotation curves of galaxies (Chan & Del Popolo 2022; Khademi et al. 2022, however see also Elstner et al. 2014). Detailed mapping of

★ E-mail: yikki.ma@anu.edu.au

the magnetic field strengths and structures in galaxies is challenging, but important for a full understanding of the astrophysical processes above. In addition, it has wide applicabilities such as tracing gas flows (e.g., Beck et al. 1999; Heald 2012), disentangling the 3D structures of galaxies (e.g., Panopoulou et al. 2021), and furthering our fundamental understanding in the origin and evolution of the magnetic fields in galaxies (e.g., Beck 2016; Federrath 2016a).

The linear polarisation of starlight is amongst the first phenomena utilised to measure the magnetic fields in galaxies (Hiltner 1951). While starlight is generally intrinsically unpolarised, the intervening dust in the interstellar medium (ISM) can induce linear polarisation in the observed starlight (Hall 1949; Hiltner 1949). The magnetic moment vector of an asymmetric dust grain is aligned to the ambient magnetic field via the radiative torque alignment effect (Hoang & Lazarian 2014), forcing the long axes of the dust particles to be perpendicular to the magnetic field direction. From this, the preferential extinction along the long axis of the dust grains leads to the linear polarisation signal parallel to the plane-of-sky magnetic field orientation (e.g., Andersson et al. 2015; Hoang & Lazarian 2016). Meanwhile, the same dust grains can re-emit in the infrared and sub-millimetre wavelengths, with the emission also linearly polarised but with the polarisation plane being perpendicular to the magnetic field instead (e.g., Hildebrand 1988; Planck Collaboration XIX 2015; Lopez-Rodriguez et al. 2022). These two methods can be exploited to probe the plane-of-sky magnetic fields in the colder phases of the ISM. For the line-of-sight component of the magnetic field, one can utilise the rotation measure (RM) of background polarised radio continuum sources (e.g., Ma et al. 2020; Tahani et al. 2022), or the polarised Zeeman-splitting measurements (e.g., with H I absorption, Heiles & Tröland 2005; or with OH masers, Ogbodo et al. 2020).

The linear polarisation state is commonly described by the Stokes Q and U parameters defined as

$$Q = PI \cdot \cos(2\theta), \text{ and} \quad (1)$$

$$U = PI \cdot \sin(2\theta), \quad (2)$$

where PI and θ are the polarised intensity and the polarisation position angle, respectively. We follow the convention of the International Astronomical Union (IAU) on the θ , which measures the polarisation E -vector from north through east (Contopoulos & Jappell 1974). We further define, in line with the literature, fractional Stokes q and u parameters as

$$q = Q/I, \text{ and} \quad (3)$$

$$u = U/I, \quad (4)$$

where I is the total intensity (or, Stokes I) of the emission.

High spatial resolution observations have revealed that the H I gas in the Milky Way is organised into highly filamentary structures (e.g., McClure-Griffiths et al. 2006; Clark et al. 2014; Martin et al. 2015; Kalberla et al. 2016; Blagrove et al. 2017; Soler et al. 2020; Skolidis et al. 2022; Campbell et al. 2022; Soler et al. 2022; Syed et al. 2022). Upon comparisons with starlight and dust polarisation data, it has been found that the elongation of these slender (with presumed widths of $\lesssim 0.1$ pc) H I filaments is often aligned with their ambient magnetic field orientations (McClure-Griffiths et al. 2006; Clark et al. 2014, 2015; Martin et al. 2015; Kalberla et al. 2016; Clark & Hensley 2019, see Skolidis et al. 2022 for a counter-example). However, it remains unclear whether such magnetic alignment is common within the entirety of the Milky Way as well as amongst galaxies with different astrophysical conditions, as the studies above focused on the neighbourhood around the Sun only (within a few hundreds of parsecs). The limitation is imposed by a combination

of the paucity of starlight polarisation data throughout the Galactic volume, the angular resolution of the H I as well as dust polarisation data, and the complexity of studying the Milky Way from within.

From simulations, it has been suggested that filamentary H I structures can be formed by turbulence, shocks, or thermal instabilities, with the role of the magnetic field still under debate (e.g., Hennebelle 2013; Federrath 2016b; Inoue & Inutsuka 2016; Villagran & Gazol 2018; Gazol & Villagran 2021). In fact, various numerical studies have led to results ranging from no preferred orientation of the H I filaments with respect to the magnetic field (Federrath 2016b), to the filaments preferentially oriented parallel (Inoue & Inutsuka 2016; Villagran & Gazol 2018) or perpendicular (Gazol & Villagran 2021) to the magnetic field. Extending the observational study of the relative orientation between magnetic fields and H I filamentary structures to nearby galaxies is therefore crucial, as the simpler external perspective will allow us to verify, despite the very different spatial scales probed, if the magnetically aligned H I filaments are a general trend across a vast galactic volume. The main hurdle to achieving this is obtaining H I data of sufficient quality, specifically the spatial resolution, velocity resolution, and sensitivity.

Apart from improving our understanding of the physical nature of H I filaments as discussed above, the alignment of the filaments with the ambient magnetic fields, if established, will open up the possibility of using the H I data as a tomographic probe of the magnetic field. This is because the plane-of-sky magnetic field orientation can then be dissected across pseudo-distance separated by the radial velocity (e.g., Clark & Hensley 2019). It also allows the study of magnetic field tangling along the line of sight (Clark 2018).

At a distance of about 62 kpc (e.g., Scowcroft et al. 2016; Graczyk et al. 2020), the Small Magellanic Cloud (SMC) is one of the closest galaxies from us. Its proximity makes it among the best targets for the investigation of the relative orientation between magnetic fields and H I filaments. The SMC is a low-mass ($M_\star = 3 \times 10^8 M_\odot$; Skibba et al. 2012), gas-rich ($M_{\text{H I}} = 4 \times 10^8 M_\odot$; Brüns et al. 2005), low-metallicity ($Z \approx 0.004 \approx 0.3 Z_\odot$; Choudhury et al. 2018) irregular galaxy undergoing an episode of enhanced star formation ($\approx 0.26 M_\odot \text{ yr}^{-1}$; see Massana et al. 2022). The galaxy consists of two major components (see, e.g., Gordon et al. 2011): the main body called “the Bar” which is unrelated to an actual galactic bar, and a peripheral feature called “the Wing” which is believed to have formed by tidal interactions with the Large Magellanic Cloud (LMC; Besla et al. 2012). The tidal forces are believed to have also created the gaseous bridge connecting the two Magellanic Clouds (Besla et al. 2012), aptly named the Magellanic Bridge. The overall 3D structures of both the gaseous and stellar components of the SMC are highly complex, and remain poorly understood (see, e.g., Di Teodoro et al. 2019; Murray et al. 2019; Tatton et al. 2021, and references therein).

The SMC has previously been observed and studied using the Australia Telescope Compact Array (ATCA) in H I emission (Staveley-Smith et al. 1997). The angular resolution of these data ($1.6' \approx 30$ pc) is a drastic improvement over those of single dish observations, leading to the distinct identification of numerous shell structures throughout the galaxy (Staveley-Smith et al. 1997; Stanimirovic et al. 1999). With the Australian Square Kilometre Array Pathfinder (ASKAP) radio telescope (Hotan et al. 2021), the SMC was observed in H I during the commissioning phase with 16 antennas (McClure-Griffiths et al. 2018), and recently with the full 36 antennas array as part of the pilot observations for the GASKAP-H I survey (Pingel et al. 2022, see Dickey et al. 2013 for a description of the GASKAP survey). The latter pilot survey data have clearly revealed the highly filamentary structures of the SMC (see Section 2.1), enabling our study here

regarding the links between these H_I structures and the associated ambient magnetic field.

Apart from the early studies observing the polarised synchrotron emission from within the SMC (Haynes et al. 1986; Loiseau et al. 1987), the magnetic field of the SMC was first explored in great detail by Mao et al. (2008), using both RM of polarised background extragalactic radio sources (EGSs) and polarised stars within the SMC. An extensive starlight polarisation catalogue of the SMC was made available by Lobo Gomes et al. (2015), leading to their study of the plane-of-sky magnetic field in the northeastern end of the SMC Bar, the SMC Wing, and the start of the Magellanic Bridge (see Section 2.2). Recently, the line-of-sight magnetic field has been revisited with RM values from new ATCA observations of EGSs (Livingston et al. 2022). The current picture of the galactic-scale magnetic field in the SMC consists of:

- A coherent magnetic field along the line-of-sight ($\approx 0.2\text{--}0.3\ \mu\text{G}$) directed away from the observer across the entire galaxy;
- Two trends in the plane-of-sky magnetic field orientation ($\approx 0.9\text{--}1.6\ \mu\text{G}$), one aligned with the elongation of the SMC Bar, and the other along the direction towards the SMC Wing and the Magellanic Bridge; and
- A turbulent magnetic field component that dominates in strength ($\approx 1.5\text{--}5.0\ \mu\text{G}$) over the ordered / coherent counterparts, by a factor of ≈ 1.5 in the plane-of-sky and ≈ 10 along the line of sight.

However, the current spatial coverage of the data (both EGS RM and starlight polarisation) remain too coarse to construct a detailed map of the magnetic structure of the SMC.

Are the H_I filaments in the SMC preferentially aligned to the ambient magnetic field similar to the case in the solar neighbourhood, despite the vastly different astrophysical characteristics (e.g., metallicity, mass, star formation rate, and tidal influences) and spatial scales probed (≈ 0.1 pc in the Milky Way; ≈ 9 pc in the SMC)? How are the 3D H_I structures linked to the different astrophysical processes occurring in the SMC, including its overall magnetic field structure? Motivated by these questions, we investigate in this work the relative orientation between H_I structures in the SMC as traced by the new GASKAP-H_I data and the magnetic fields traced by starlight polarisation reported by Lobo Gomes et al. (2015).

This paper is organised as follows. We describe the data and the associated processing required for our study in Section 2, and devise a new ray-tracing algorithm that enables our careful comparison between the H_I and starlight polarisation data as outlined in Section 3. In Section 4, we (1) evaluate whether the SMC H_I filaments are magnetically aligned, (2) test whether the GASKAP-H_I data can trace the small-scale turbulent magnetic field, and (3) present the plane-of-sky magnetic field structure of the SMC as traced by H_I filaments. We discuss the implications of our work in Section 5, and conclude our study in Section 6.

2 DATA AND DATA PROCESSING

2.1 H_I filaments from GASKAP

We use new GASKAP-H_I data of the SMC for this study (Pingel et al. 2022). The 20.9-hour ASKAP data were taken in December 2019 during Phase I of the Pilot Survey, and were combined with single-dish data from the Parkes Galactic All-Sky Survey (GASS; McClure-Griffiths et al. 2009). The resulting data cube presents an unprecedented view of the H_I emission of the SMC (see Figure 1), with the highest combination of angular resolution (synthesised beam

of $30''$), velocity resolution ($0.98\ \text{km s}^{-1}$), and sensitivity ($1.1\ \text{K per channel}$).

It is immediately apparent that the SMC exhibits a vast network of filamentary structures throughout the entire galaxy. We proceed to apply the Rolling Hough Transform¹ (RHT; Clark et al. 2014) algorithm to the GASKAP-H_I cube to automatically locate these filaments. Other algorithms that have been used in the literature for the study of elongated structures include the Hessian analysis (e.g., Polychroni et al. 2013; Kalberla et al. 2016) and the anisotropic wavelet analysis (e.g., Patrikeev et al. 2006; Frick et al. 2016). While the former has been shown to lead to comparable results as the RHT (Soler et al. 2020), the differences of the latter with the RHT have not been explored in details, and is beyond the scope of this work.

In particular, we apply the convolutional RHT algorithm (see BI-CEP/Keck Collaboration et al. 2022, for details) which is a significant improvement in the computational efficiency. For each 2D image, the RHT first performs an unsharp mask procedure, subtracting from the image a smoothed version of itself. The smoothing is done by convolving the image with a circular top-hat function with radius R_{sm} . Next, a bitmask is created by checking the value of the resulting difference map – `True` if the pixel value is greater than zero, and `False` otherwise. This bitmask can be regarded as a map of small-scale structures, including potential filaments, edges of structures, etc. Finally, the algorithm “rolls” through each pixel in the bitmask image and quantifies the distribution of surrounding linear structure. This is done by extracting a circular window with diameter D_W around each pixel, and applying a Hough transform (Hough 1962) to the bitmasked data in the window, with the sampling done through the centre of the circular window only (i.e., $\rho = 0$ in the formulation of Duda & Hart 1972). A simplified explanation of the operation here is that we sample straight lines passing through the centre pixel of the circular window, each with different θ ranging from 0° to 180° . For each of the straight lines, the fraction of `True`-valued pixels has been evaluated and compared with the `threshold` parameter. If the computed fraction exceeds `threshold`, the fraction value subtracted by the `threshold` is written to the final output 4D-hypercube (with the axes being the two spatial coordinates, velocity, and θ). Otherwise, zero is written to the hypercube instead. In other words, a non-zero pixel in the RHT 4D-hypercube means a filament with orientation θ passes through the 3D location (position-position-velocity) of the corresponding pixel.

The original RHT algorithm outlined above performs well for Galactic sky regions and velocity ranges where the emission is ubiquitous (e.g., Clark et al. 2014; Jelić et al. 2018; Campbell et al. 2022). However, this is far from the case for the SMC in H_I, for which the presence of emission is highly dependent on the location and the radial velocity (Stanimirovic et al. 1999; McClure-Griffiths et al. 2018; Di Teodoro et al. 2019; Pingel et al. 2022). Upon application of this original RHT to the new GASKAP-H_I cube of the SMC, we find that it can sometimes erroneously identify filamentary structures in very low signal-to-noise sky areas. This prompts us to implement an intensity cutoff procedure in the RHT algorithm – the bitmask formation step above would additionally compare the H_I intensity of the input image with a determined cutoff value, and will set the bitmask pixel value as `False` if the intensity is lower than the cutoff. For our application here, we adopt a cutoff value of ($5.7\ \text{K}/P_{\text{PB}}$), where the $5.7\ \text{K}$ corresponds to five times the rms noise near the centre of the images, and P_{PB} is the primary beam attenuation level.

¹ Available on <https://github.com/seclark/RHT>.




Figure 1. The H I peak intensity image of the SMC from the GASKAP-H I Pilot Survey I observations (Pingel et al. 2022), highlighting the vast network of filamentary structures in this galaxy. The locations of the 20 Lobo Gomes et al. (2015) starlight polarisation fields considered in this study are each shown as a square with the size reflecting the true field of view of 8×8 sq. arcmin. The approximate spatial division between the Bar and the Wing regions of the SMC is outlined by the grey dotted line, and the Magellanic Bridge is situated outside of the covered sky area in the direction indicated by the arrow to the lower left.

We apply the modified RHT algorithm² independently to each of the 223 velocity channels³ from 40.91 to 257.85 km s⁻¹, with the three RHT parameters set as $R_{\text{sm}} = 12 \text{ px} = 25 \text{ pc}$, $D_W = 83 \text{ px} = 175 \text{ pc}$, and $\text{threshold} = 0.7$. The conversions to physical scales above assume a distance of 62 kpc to the SMC (e.g., Scowcroft et al. 2016; Graczyk et al. 2020) with a pixel scale of $7''$ for the GASKAP-H I data (Pingel et al. 2022). A sample of the RHT output is illustrated in Figure 2. Our choice of R_{sm} , in units of the synthesised beam, is similar to that of Clark et al. (2014) with Galactic Arecibo L-Band Feed Array H I (GALFA-H I; Peek et al. 2011) data (2.8 for us

here compared to their 2.5). Meanwhile, our chosen D_W in units of R_{sm} , which determines the aspect ratios of the identified filamentary structures, is about 7, again similar to the choice of Clark et al. (2014) of 10. Finally, our choice of threshold is identical to Clark et al. (2014). To ensure that our results are not critically dependent on the RHT parameter choice, we repeat our analysis using different sets of parameters, reported in Appendix A.

In this study, we do not count the number of H I filaments identified, since the RHT algorithm only reports whether a pixel is part of a filamentary structure, but not group the many pixels together as a filament. The quantification of the number of filaments in the SMC will require additional algorithms that take into account the spatial and radial-velocity coherence of the RHT output, which is beyond the scope of this work.

Finally, we note that the GASKAP-H I SMC maps are in orthographic projection, and given the large angular extent of the maps,

² This can be toggled on by using the `cutoff_mask` parameter in the convolutional RHT algorithm.

³ The radial velocities presented throughout this work are with respect to the local standard of rest (LSR).




Figure 2. Illustration of the automatically identified filaments using RHT. Left panel shows a zoomed-in image of the central area of the GASKAP-H_I SMC map (Pingel et al. 2022) at $v_{\text{LSR}} = 133.74 \text{ km s}^{-1}$, and the right panel shows in cubehelix colour scheme (Green 2011) the corresponding RHT back-projection map, where any non-zero pixels are regarded as a filament in our study.

sky curvature is apparent (see Figure 1). This means that the vertical axis of the map is in general not parallel to the sky north-south axis. As RHT operates on the maps' cartesian grid, there can be angle offsets between the output hypercube's θ -axis and the sky θ . This has been corrected for in our analysis throughout this paper.

2.2 Starlight polarisation data

To trace the plane-of-sky magnetic field orientation in the SMC, we use the Lobo Gomes et al. (2015) starlight polarisation catalogue derived from a V-band optical survey towards the SMC using the Cerro Tololo Inter-American Observatory (CTIO). The survey has covered a total of 28 fields in the northeastern Bar and the Wing of the SMC, as well as part of the Magellanic Bridge, with a field of view of 8×8 sq. arcmin each. The polarisation properties of 7,207 stars have been reported, with the foreground polarisation contribution of the Milky Way determined and subtracted in Stokes qu space by making use of the polarised starlight from Galactic stars in the same sky area. To compare with our GASKAP-H_I data of the SMC, we focus on the 20 starlight fields in the SMC only (Figure 1), encompassing a total of 5,999 stars with detected linear polarisation.

In Lobo Gomes et al. (2015), the preferred orientation(s) of the starlight polarisation angle ($\theta_\star$) of each of their fields was obtained by fitting a single- or double-component Gaussian function to the histogram of $\theta_\star$. In other words, they have only used the angle information of the starlight polarisation vector (in Stokes qu plane), without taking the polarisation fraction ($p_\star$) into account. Here, we re-analyse the starlight polarisation data with a full vector approach as outlined below.

Consider that the SMC is permeated by a magnetic field composed of two components in superposition – a large-scale magnetic field with a coherence length $\gg 100$ pc, and a small-scale isotropic magnetic field with a coherence length $\lesssim 100$ pc (e.g., Beck 2016, see also Livingston et al. 2022). As each of the Lobo Gomes et al. (2015) fields spans ≈ 150 pc across in the plane of sky, the two magnetic field components will leave different imprints on the observed starlight polarisation when we consider each starlight field as a whole. On the Stokes qu plane, all stars start at the origin ($q = u = 0$) since they are intrinsically unpolarised. The large-scale magnetic field in




Figure 3. Results from our re-analysis of the Lobo Gomes et al. (2015) starlight polarisation data. Green line segments are directed along the $\bar{\theta}_\star$ that trace the plane-of-sky magnetic field orientation for fields that we find a coherent starlight polarisation angle, while red crosses mark fields that do not exhibit a coherent starlight polarisation angle. The background image shows the same but zoomed in GASKAP-H_I peak intensity map as in Figure 1.

the intervening volume shifts all stars coherently in a single direction in the Stokes qu plane as determined by its magnetic field orientation, while the small-scale magnetic field scatters the stars isotropically in the Stokes qu plane.

In light of the expected effects of the two magnetic field components on the observed starlight polarisation, we re-analyse the Lobo Gomes et al. (2015) data accordingly. The large-scale magnetic field

contribution is evaluated by the vector mean in Stokes qu space:

$$\bar{q}_\star = \frac{1}{N_\star} \sum_i^{N_\star} q_i, \text{ and} \quad (5)$$

$$\bar{u}_\star = \frac{1}{N_\star} \sum_i^{N_\star} u_i, \quad (6)$$

where i is the index for the $N_\star$ stars within each of the starlight fields. This can be further converted to $\bar{p}_\star$ and $\bar{\theta}_\star$ by

$$\bar{p}_\star = \sqrt{\bar{q}_\star^2 + \bar{u}_\star^2}, \text{ and} \quad (7)$$

$$\bar{\theta}_\star = 0.5 \tan^{-1}(\bar{u}_\star / \bar{q}_\star). \quad (8)$$

Meanwhile, the effect of the small-scale magnetic field is captured through the 2D standard deviation ($\sigma_{p\star} = \sqrt{\sigma_{q\star} \cdot \sigma_{u\star}}$) of the star sample in Stokes qu plane, with $\sigma_{q\star}$ and $\sigma_{u\star}$ being the 1D standard deviation of Stokes q and u , respectively. The uncertainties in $\bar{p}_\star$, $\bar{\theta}_\star$, and $\sigma_{p\star}$ are estimated by bootstrapping – for each starlight field we correspondingly draw with replacement $N_\star$ stars, and obtain the values of the three parameters as above. This process is repeated 10^6 times, and the standard deviations out of the 10^6 values are taken as the uncertainty values of the three parameters⁴. The values of $\bar{\theta}_\star$, $\bar{p}_\star$, and $\sigma_{p\star}$ of each field, as well as the number of SMC stars per field ($N_\star$) are all listed in Table 1, with the corresponding 2D histograms shown in Figure C1 under Appendix C. Finally, the sky distribution of $\bar{\theta}_\star$ is shown in Figure 3.

We deem the resulting $\bar{\theta}_\star$ of seven out of the total of 20 fields as uncertain, since their signal-to-noise ratios of $\bar{p}_\star$ are low (< 3). All these uncertain values are placed in parentheses in Table 1.

We compare our newly obtained $\bar{\theta}_\star$ of each field with the corresponding results from Lobo Gomes et al. (2015). Since their approach can yield up to two polarisation angles for each starlight field, we identify the primary polarisation component for such case, defined as the listed Gaussian component with the highest peak in their $\theta_\star$ histogram. The resulting angles are labelled as θ_{LG15} and listed in Table 1. In almost all fields, the values of our $\bar{\theta}_\star$ show good agreement with θ_{LG15} (to within 10°), with the only exceptions being field 7 (angle difference of $26^\circ \pm 6^\circ$) and field 12 (angle difference of $11^\circ \pm 7^\circ$).

2.3 3D dust extinction data

To model the extinction effect experienced by starlight through the SMC (Section 3), we require 3D information of SMC dust extinction. Yanchulova Merica-Jones et al. (2021) derived a relation between the dust extinction (A_V) and the hydrogen column density (N_{H}) for the southwestern end of the SMC Bar region as

$$\frac{A_V}{N_{\text{H}}} = \frac{A_V}{N_{\text{H}} + 2N_{\text{H}_2}} = 3.2\text{--}4.2 \times 10^{-23} \text{ mag cm}^2 \text{ H}^{-1}, \quad (9)$$

where N_{H} and N_{H_2} are the column densities for atomic and molecular hydrogen, respectively. While we have the full 3D (position-position-velocity) information for H I from our new GASKAP-H I observations covering the entire SMC, the same for H₂ is not available.

We therefore attempt to convert the 2D N_{H_2} map from Jameson et al. (2016), obtained through *Herschel* observations of dust emission, to an approximate 3D distribution of H₂ throughout the SMC.

To achieve this, we first obtain a 2D $N_{\text{H I}}$ map from the GASKAP-H I data. From this, we compute a molecular-to-atomic hydrogen column density ratio map ($N_{\text{H}_2}/N_{\text{H I}}$), and subsequently apply it to each velocity slice of the GASKAP-H I cube to obtain the 3D H₂ cube. In other words, we assume that the H I and H₂ number densities are correlated, which is generally not the case (e.g., Wannier et al. 1983; Lee et al. 2012). However, we point out that the exact details of the implementation of the H₂ data likely will not significantly affect our results here, as we find that the SMC is dominated by H I, with the median H₂-to-H I column density ratio being a mere 0.06.

Finally, we apply Equation 9 to the H I and H₂ cubes to obtain the 3D dust extinction cube of the SMC, with the middle of the quoted range (i.e., $3.7 \times 10^{-23} \text{ mag cm}^2 \text{ H}^{-1}$) adopted as the applied value. Each velocity slice of this cube is a map of extinction (in units of mag) that V-band starlight is subjected to while traversing through the corresponding volume.

3 RAY TRACING OF STARLIGHT POLARISATION

We proceed to perform a careful comparison between the orientation of H I filaments (Section 2.1) and the magnetic field traced by starlight polarisation (Section 2.2). For this, we devise a ray-tracing analysis of starlight polarisation, with the effect of diminishing starlight intensity due to dust extinction (Section 2.3) taken into account. Our goal here is to obtain the expected linear polarisation signature of each of the Lobo Gomes et al. (2015) stars, assuming that the H I filaments in the SMC are indeed aligned with the ambient magnetic fields that are also experienced by the dust grains imprinting linear polarisation signals in the observed starlight. This assumption will be confirmed if we find a match between the expected (from ray tracing) and the observed starlight polarisation. In essence, we use the locations of polarised SMC stars reported in Lobo Gomes et al. (2015), and send the starlight through the GASKAP-H I cube. When the starlight is intercepted by H I filaments, linear polarisation signal along the filament orientation is added to it accordingly⁵. Note that the results from the ray-tracing analysis here are a representation of the H I data, and the ray tracing is done (instead of averaging all spatial pixels in the H I data) to completely remove the possibility of sampling bias imposed by the positions where polarised stars were found in Lobo Gomes et al. (2015). Furthermore, we adopt this ray-tracing approach instead of directly comparing the orientation angles of the filaments and starlight polarisation (e.g., McClure-Griffiths et al. 2006; Clark et al. 2014) since, for our case here studying the SMC, the observed starlight often traverses through multiple H I filaments along the sightline. The contributions by these filaments are correctly combined by our ray-tracing analysis. The details of the ray tracing are described below.

First, we need to determine the 3D positions where we place the stars within the GASKAP-H I cube. While the plane-of-sky locations (i.e., in right ascension and declination) of the stars can be directly adopted from the Lobo Gomes et al. (2015) catalogue, the choice along the velocity axis is less straightforward. Putting the stars on the far side of the cube may not be a good choice, since this would be assuming that all of the polarised SMC stars are physically behind all the gas in the SMC. Instead, we calculate, for each of the 20 starlight polarisation fields, the H I intensity weighted mean velocity (v_{mean})

⁴ We repeat this bootstrapping for ten times, each time resampling 10^6 times as stated, and find that the resulting uncertainties are always almost identical, meaning that these obtained uncertainty values have certainly converged.

⁵ The preferential extinction of starlight along the polarisation plane perpendicular to the magnetic field will lead to a net polarisation signal added along the magnetic field orientation.

Table 1. Observables of each of the Lobo Gomes et al. (2015) starlight polarisation fields covered by the new GASKAP-H α SMC field

Field No.	θ_{LG15} (deg)	$\bar{\theta}_\star$ (deg)	$\bar{p}_\star$ (%)	$\sigma_{p\star}$ (%)	$N_\star$	v_{mean} (km s $^{-1}$)
1	120.9 $\pm$ 0.3	120.5 $\pm$ 2.5	0.45 $\pm$ 0.04	0.92 $\pm$ 0.03	491	154.81
2	58.4 $\pm$ 0.7	52.6 $\pm$ 4.1	0.20 $\pm$ 0.03	0.61 $\pm$ 0.02	492	153.39
3	102.2 $\pm$ 0.6	106.0 $\pm$ 4.5	0.34 $\pm$ 0.05	1.11 $\pm$ 0.03	471	145.00
4	169.2 $\pm$ 1.6	178.0 $\pm$ 7.6	0.30 $\pm$ 0.07	0.95 $\pm$ 0.05	170	144.30
5	–	117.0 $\pm$ 10.8	0.32 $\pm$ 0.10	0.76 $\pm$ 0.06	47	166.12
6	100.6 $\pm$ 2.1	(96.7 $\pm$ 15.8)	(0.18 $\pm$ 0.07)	1.29 $\pm$ 0.06	247	165.88
7	178.8 $\pm$ 0.7	25.0 $\pm$ 5.6	0.22 $\pm$ 0.04	1.06 $\pm$ 0.03	640	146.40
8	73.7 $\pm$ 0.4	74.8 $\pm$ 4.7	0.21 $\pm$ 0.03	0.91 $\pm$ 0.02	735	164.11
9	67.5 $\pm$ 0.2	69.0 $\pm$ 2.3	0.49 $\pm$ 0.04	0.96 $\pm$ 0.03	605	152.18
10	142.5 $\pm$ 0.3	142.3 $\pm$ 4.7	0.26 $\pm$ 0.05	1.09 $\pm$ 0.03	560	153.17
11	158.3 $\pm$ 0.2	156.0 $\pm$ 1.7	0.86 $\pm$ 0.05	1.08 $\pm$ 0.03	453	154.75
12	147.9 $\pm$ 0.7	159.0 $\pm$ 7.0	0.36 $\pm$ 0.08	1.21 $\pm$ 0.06	205	151.92
13	126.2 $\pm$ 2.5	(139.0 $\pm$ 23.4)	(0.14 $\pm$ 0.08)	1.06 $\pm$ 0.06	122	158.27
14	–	(92.4 $\pm$ 44.1)	(0.03 $\pm$ 0.06)	1.07 $\pm$ 0.06	165	157.20
15	144.5 $\pm$ 1.0	142.4 $\pm$ 5.1	0.54 $\pm$ 0.09	1.11 $\pm$ 0.06	139	157.43
16	134.5 $\pm$ 4.8	(121.7 $\pm$ 24.5)	(0.19 $\pm$ 0.11)	1.15 $\pm$ 0.09	87	169.38
17	54.0 $\pm$ 1.6	(52.4 $\pm$ 37.1)	(0.07 $\pm$ 0.07)	1.06 $\pm$ 0.06	136	149.78
18	88.5 $\pm$ 2.9	(94.0 $\pm$ 16.2)	(0.21 $\pm$ 0.09)	1.16 $\pm$ 0.08	131	150.06
19	–	(102.4 $\pm$ 40.7)	(0.10 $\pm$ 0.12)	1.10 $\pm$ 0.11	40	153.40
20	75.0 $\pm$ 3.2	72.7 $\pm$ 10.4	0.38 $\pm$ 0.12	1.01 $\pm$ 0.10	63	159.22

NOTE – Parameters that are deemed uncertain, as described in the text, are placed in parentheses.

and place the corresponding polarised SMC stars there. The values of v_{mean} are listed in Table 1.

The above choice of 3D stellar positions involves two key assumptions. The first one being that for any given line of sight, the H α velocity has a monotonic trend with the macroscopic physical distance. While effects such as the gas dynamics of small H α clouds and turbulence can break this monotonic trend within a velocity range of $\sim 1\text{--}10\text{ km s}^{-1}$, we require the H α velocity to follow the macroscopic physical distance monotonically for the ray-tracing experiment to be a good analogy with attenuation along the line of sight. The second assumption is that the H α velocity profile traces stellar density across the physical distances corresponding to the associated H α velocities. This ensures that using v_{mean} as the ray-tracing starting point for a large ($\gtrsim 100$) number of stars will give statistically meaningful results. We further attempt using the radial velocity measurements from the *Gaia* DR3 (Gaia Collaboration et al. 2022) instead of v_{mean} as the stars' positions along the line of sight for the 57 cross-matched stars, as reported in Appendix B.

Our next step is to direct starlight from v_{mean} through the higher ($v \geq v_{\text{mean}}$) and lower ($v < v_{\text{mean}}$) velocity portions of the H α cube independently. The former (latter) case would imply that the higher (lower) velocity H α gas is physically closer to us, since the gas as well as the associated dust causing the stellar extinction needs to intercept the traversing starlight to cause the observed linear polarisation. While many optical and ultraviolet absorption line studies have suggested that the lower velocity gas component of the SMC is physically closer to us (e.g., Mathewson et al. 1986; Danforth et al. 2002; Welty et al. 2012), we do not make this assumption a-priori. In addition, despite being astrophysically unrealistic (see above), we also perform the ray-tracing analysis through the entire SMC H α cube (40.91–256.85 km s $^{-1}$) for both cases of starting from the lower and higher velocity ends for completeness⁶.

For each step through the radial velocity axis, we check individually for each of the stars if the corresponding starlight is being intercepted by any H α filaments. If so, we add starlight polarisation signal accordingly as follows, taking into account the possibility of overlapping filaments with different orientations (θ_i) at a single velocity step. The added linear polarisation at velocity v is expressed in Stokes QU space as

$$Q(v) = \frac{F(v) \cdot I(v)}{n} \cdot \sum_i^n \cos 2\theta_i, \text{ and} \quad (10)$$

$$U(v) = \frac{F(v) \cdot I(v)}{n} \cdot \sum_i^n \sin 2\theta_i, \quad (11)$$

where $F(v)$ is the (unitless) attenuated fractional starlight flux density due to dust extinction (see next paragraph), $I(v)$ is the H α intensity, and the summation index i goes through the list of the n intercepting filaments, all evaluated at the sky position of the star at velocity v . This operation does not only give the correct orientation of the polarisation signal to be added, but also accounts for the depolarisation effect among multiple filaments. For example, consider the extreme case of two orthogonal intervening filaments, which are expected to cancel out one another and add no linear polarisation signal to the traversing starlight. Our scheme above would correctly yield $Q(v) = U(v) = 0$. Finally, the added polarisation signal (barring the depolarisation effect above) is proportional to the H α intensity, since we expect the amount of extinction leading to the observed polarisation to be proportional to the dust and gas column densities.

The attenuated fractional starlight flux density $F(v)$ introduced in the above paragraph incorporates the amount of dust extinction sustained by the starlight over its journey up till v . This term is

during the traversal along the line of sight (see below), the results do not only depend on the velocity range considered, but also the direction along the velocity axis that the starlight propagates through.

⁶ Since our ray-tracing algorithm takes into account the extinction of starlight

Table 2. Results from our new starlight polarisation ray-tracing analysis through the GASKAP-HI cube

Field No.	$\bar{\theta}_{\text{HI}}$ (deg)	Low Velocity Range ($v < v_{\text{mean}}$)				$\bar{\theta}_{\text{HI}}$ (deg)	High Velocity Range ($v \geq v_{\text{mean}}$)			
		$\bar{p}_{\text{HI}}$ (%)	$\sigma_{p_{\text{HI}}}$ (%)	$\Delta\bar{\theta}$ (deg)	C (10^{-3})		$\bar{p}_{\text{HI}}$ (%)	$\sigma_{p_{\text{HI}}}$ (%)	$\Delta\bar{\theta}$ (deg)	C (10^{-3})
1	69.0 ± 1.9	0.45 ± 0.03	0.63 ± 0.01	51.5 ± 3.1	2.41	71.7 ± 1.4	0.45 ± 0.02	0.49 ± 0.01	48.8 ± 2.9	2.49
2	52.0 ± 7.1	0.20 ± 0.04	1.05 ± 0.02	0.6 ± 8.2	9.49	37.8 ± 3.2	0.20 ± 0.03	0.52 ± 0.01	14.8 ± 5.2	2.19
3	95.9 ± 1.1	0.34 ± 0.01	0.28 ± 0.01	10.0 ± 4.6	1.46	(146.8 ± 13.5)	(0.34 ± 0.12)	3.01 ± 0.07	(40.9 ± 14.2)	11.97
4	(147.0 ± 24.7)	(0.30 ± 0.18)	2.69 ± 0.12	(31.0 ± 25.9)	9.97	110.2 ± 3.7	0.30 ± 0.04	0.50 ± 0.02	67.9 ± 8.4	2.45
5	54.4 ± 6.1	0.32 ± 0.07	0.46 ± 0.03	62.6 ± 12.4	1.54	54.1 ± 4.6	0.32 ± 0.05	0.35 ± 0.02	62.9 ± 11.7	0.96
6	(118.0 ± 16.8)	(0.18 ± 0.08)	1.36 ± 0.04	(21.3 ± 23.1)	5.84	22.4 ± 3.5	0.18 ± 0.02	0.30 ± 0.01	(74.3 ± 16.2)	0.69
7	83.0 ± 1.6	0.22 ± 0.01	0.30 ± 0.01	58.0 ± 5.8	1.15	63.6 ± 1.5	0.22 ± 0.01	0.30 ± 0.01	38.6 ± 5.8	2.10
8	85.1 ± 2.0	0.21 ± 0.01	0.39 ± 0.01	10.3 ± 5.1	3.63	64.0 ± 2.1	0.21 ± 0.01	0.41 ± 0.01	10.8 ± 5.1	1.98
9	84.6 ± 1.8	0.49 ± 0.02	0.62 ± 0.01	15.6 ± 2.9	3.06	76.1 ± 1.9	0.49 ± 0.03	0.74 ± 0.01	7.1 ± 3.0	4.15
10	136.8 ± 3.3	0.26 ± 0.03	0.72 ± 0.01	5.5 ± 5.7	3.50	66.9 ± 1.2	0.26 ± 0.01	0.27 ± 0.01	75.4 ± 4.9	1.23
11	169.0 ± 3.5	0.86 ± 0.10	2.17 ± 0.05	12.9 ± 3.9	14.10	37.1 ± 2.7	0.86 ± 0.07	1.61 ± 0.04	61.1 ± 3.1	5.76
12	108.4 ± 8.2	0.36 ± 0.09	1.36 ± 0.05	50.6 ± 10.8	12.68	(70.7 ± 14.3)	(0.36 ± 0.14)	2.20 ± 0.08	(88.3 ± 16.0)	18.93
13	44.5 ± 2.6	0.14 ± 0.02	0.16 ± 0.01	(85.6 ± 23.5)	1.23	106.7 ± 4.5	0.14 ± 0.02	0.24 ± 0.01	(32.2 ± 23.8)	1.01
14	49.3 ± 3.6	0.03 ± 0.00	0.04 ± 0.00	(43.1 ± 44.2)	0.33	(16.9 ± 17.0)	(0.03 ± 0.01)	0.18 ± 0.01	(75.4 ± 47.2)	0.61
15	48.0 ± 2.6	0.54 ± 0.05	0.57 ± 0.02	85.6 ± 5.7	3.77	(99.0 ± 30.4)	(0.54 ± 0.40)	6.00 ± 0.41	(43.4 ± 30.8)	24.43
16	61.6 ± 5.8	0.19 ± 0.04	0.35 ± 0.02	(60.0 ± 25.1)	8.33	(61.3 ± 18.2)	(0.19 ± 0.09)	0.91 ± 0.06	(60.3 ± 30.5)	12.81
17	67.2 ± 3.4	0.07 ± 0.01	0.09 ± 0.00	(14.8 ± 37.2)	0.47	88.2 ± 8.1	0.07 ± 0.01	0.20 ± 0.01	(35.7 ± 38.0)	2.31
18	92.2 ± 2.5	0.21 ± 0.02	0.23 ± 0.01	(1.8 ± 16.4)	1.74	14.3 ± 5.2	0.21 ± 0.04	0.45 ± 0.02	(79.7 ± 17.1)	5.05
19	99.2 ± 4.7	0.10 ± 0.02	0.11 ± 0.01	(3.2 ± 41.0)	1.65	(127.4 ± 10.3)	(0.10 ± 0.04)	0.22 ± 0.02	(25.0 ± 42.0)	3.56
20	97.0 ± 7.2	0.38 ± 0.06	0.61 ± 0.05	24.4 ± 12.7	6.51	(155.5 ± 24.6)	(0.38 ± 0.24)	2.06 ± 0.16	(82.9 ± 26.7)	102.14

Field No.	$\bar{\theta}_{\text{HI}}$ (deg)	Full Velocity Range ($v_0 = 40.91 \text{ km s}^{-1}$)				$\bar{\theta}_{\text{HI}}$ (deg)	Full Velocity Range ($v_0 = 256.85 \text{ km s}^{-1}$)			
		$\bar{p}_{\text{HI}}$ (%)	$\sigma_{p_{\text{HI}}}$ (%)	$\Delta\bar{\theta}$ (deg)	C (10^{-3})		$\bar{p}_{\text{HI}}$ (%)	$\sigma_{p_{\text{HI}}}$ (%)	$\Delta\bar{\theta}$ (deg)	C (10^{-3})
1	70.2 ± 1.3	0.45 ± 0.02	0.43 ± 0.01	50.3 ± 2.9	1.30	70.1 ± 1.3	0.45 ± 0.02	0.42 ± 0.01	50.4 ± 2.9	1.28
2	40.9 ± 3.2	0.20 ± 0.02	0.51 ± 0.01	11.7 ± 5.2	1.94	40.5 ± 3.1	0.20 ± 0.02	0.49 ± 0.01	12.1 ± 5.1	1.84
3	98.8 ± 1.7	0.34 ± 0.02	0.45 ± 0.01	7.2 ± 4.8	1.46	99.0 ± 1.9	0.34 ± 0.03	0.51 ± 0.01	6.9 ± 4.8	1.67
4	116.7 ± 5.7	0.30 ± 0.06	0.78 ± 0.03	61.4 ± 9.5	2.43	115.8 ± 5.5	0.30 ± 0.06	0.75 ± 0.03	62.3 ± 9.3	2.41
5	53.9 ± 4.2	0.32 ± 0.03	0.26 ± 0.02	63.1 ± 11.6	0.65	54.1 ± 4.1	0.32 ± 0.03	0.26 ± 0.02	62.9 ± 11.5	0.62
6	21.7 ± 4.9	0.18 ± 0.03	0.46 ± 0.01	(75.0 ± 16.6)	0.88	21.3 ± 4.6	0.18 ± 0.03	0.43 ± 0.01	(75.4 ± 16.4)	0.79
7	77.0 ± 1.3	0.22 ± 0.01	0.24 ± 0.01	52.0 ± 5.7	0.81	75.7 ± 1.2	0.22 ± 0.01	0.24 ± 0.01	50.7 ± 5.7	0.83
8	71.5 ± 1.8	0.21 ± 0.01	0.35 ± 0.01	3.3 ± 5.0	1.42	70.7 ± 1.8	0.21 ± 0.01	0.36 ± 0.01	4.1 ± 5.0	1.43
9	81.0 ± 1.3	0.49 ± 0.02	0.50 ± 0.01	11.9 ± 2.6	1.83	80.9 ± 1.3	0.49 ± 0.02	0.50 ± 0.01	11.8 ± 2.6	1.86
10	76.4 ± 2.2	0.26 ± 0.02	0.49 ± 0.01	65.9 ± 5.2	1.78	74.5 ± 2.0	0.26 ± 0.02	0.43 ± 0.01	67.8 ± 5.1	1.57
11	24.7 ± 3.1	0.86 ± 0.09	1.97 ± 0.04	48.7 ± 3.5	5.80	27.4 ± 3.1	0.86 ± 0.09	1.96 ± 0.04	51.4 ± 3.6	5.76
12	93.8 ± 9.1	0.36 ± 0.10	1.52 ± 0.06	65.2 ± 11.5	9.87	92.4 ± 9.4	0.36 ± 0.10	1.56 ± 0.06	66.7 ± 11.7	10.12
13	79.4 ± 6.5	0.14 ± 0.03	0.34 ± 0.02	(59.5 ± 24.2)	1.23	83.8 ± 6.7	0.14 ± 0.03	0.35 ± 0.02	(55.1 ± 24.3)	1.24
14	38.9 ± 5.3	0.03 ± 0.01	0.08 ± 0.00	(53.5 ± 44.4)	0.25	38.3 ± 6.0	0.03 ± 0.01	0.08 ± 0.00	(54.1 ± 44.5)	0.27
15	52.2 ± 4.7	0.54 ± 0.09	1.06 ± 0.06	89.8 ± 6.9	3.76	53.2 ± 5.1	0.54 ± 0.10	1.16 ± 0.06	89.2 ± 7.2	4.08
16	61.5 ± 7.4	0.19 ± 0.04	0.43 ± 0.03	(60.1 ± 25.6)	5.14	61.9 ± 7.5	0.19 ± 0.05	0.44 ± 0.03	(59.7 ± 25.6)	5.17
17	70.2 ± 3.4	0.07 ± 0.01	0.09 ± 0.00	(17.8 ± 37.2)	0.41	70.8 ± 3.4	0.07 ± 0.01	0.09 ± 0.00	(18.4 ± 37.2)	0.42
18	86.5 ± 4.9	0.21 ± 0.04	0.42 ± 0.02	(7.5 ± 16.9)	2.47	86.1 ± 5.2	0.21 ± 0.04	0.45 ± 0.02	(7.9 ± 17.0)	2.60
19	107.6 ± 4.6	0.10 ± 0.02	0.10 ± 0.01	(5.2 ± 41.0)	1.27	108.0 ± 4.7	0.10 ± 0.02	0.10 ± 0.01	(5.6 ± 41.0)	1.29
20	98.4 ± 7.4	0.38 ± 0.06	0.62 ± 0.05	25.8 ± 12.8	6.71	98.7 ± 7.4	0.38 ± 0.06	0.62 ± 0.05	26.0 ± 12.8	6.81

NOTE – Parameters that are deemed uncertain are placed in parentheses. Units for the conversion factor C is $\% \text{ K}^{-1} \text{ km}^{-1} \text{ s}$.

necessary since the polarised intensity added at each velocity step should be proportional to the starlight flux density as it traverses through the same velocity step. The value of $F(v)$ can be obtained by first summing the A_V velocity cube (Section 2.3) from the starting velocity of the starlight (v_{mean} , 40.91, or 256.85 km s^{-1} , depending on where the stars are placed along the velocity axis) to the velocity channel right before v . The summed A_V is then converted from magnitude to flux density, with the intrinsic starlight flux density defined to be unity (since only the proportionality matters here). These all are captured by the following equation:

$$\log_{10} F(v_i) = -\frac{2}{5} \sum_{j=0}^{i-1} A_V(v_j), \quad (12)$$

where the summation index j goes through each of the relevant velocity channels, with v_0 corresponding to the starting velocity of the starlight, and v_i being the velocity step where the starlight flux density is being evaluated.

From the above four runs of our ray-tracing experiment with different starting velocities and velocity ranges considered, we corre-

spondingly obtain four sets of the expected linear polarisation signal from the 5,999 SMC stars. We note that all stars in all cases are intercepted by at least one HI filament, and in most cases by multiple. We extract the per-field polarisation behaviour from these four cases of ray tracing by following the identical procedures as we did to the Lobo Gomes et al. (2015) data in Section 2.2. At this stage, the Stokes Q and U values are in units of K km s^{-1} since they are brightness temperature summed across velocity channels. We convert them to Stokes q and u in units of $\%$ by applying a conversion factor C (in units of $\% \text{ K}^{-1} \text{ km}^{-1} \text{ s}$), such that the obtained ray-traced $\bar{p}_{\text{HI}}$ values here exactly match the observed $\bar{p}_{\star}$ values on a per-field basis (see next paragraph for a more detailed discussion). The resulting $\bar{\theta}_{\text{HI}}$, $\bar{p}_{\text{HI}}$, $\sigma_{p_{\text{HI}}}$, and C values are listed in Table 2, with the corresponding 2D histograms shown in Figures C2–C5 under Appendix C. The sub-script “HI” is chosen here to stress again that the ray-traced starlight polarisation results are a representation of the HI data.

Our application of the conversion factor C to each of the combinations of the four ray-tracing cases and 20 starlight fields forces the ray-traced $\bar{p}_{\text{HI}}$ values to match the observed $\bar{p}_{\star}$ obtained from a re-analysis of the Lobo Gomes et al. (2015) data (Section 2.2). The

values of C encapsulate information such as the gas-to-dust ratio in number density and the intrinsic properties of the dust (specifically, the efficacy in producing the observed starlight polarisation). While we obviously cannot then draw meaningful conclusions from comparing between the ray-traced $\bar{\rho}_{\text{H I}}$ and the observed $\bar{\rho}_*$, we can still compare the $\sigma_p/\bar{p}$ values to assess the ability of a ray-tracing experiment through the GASKAP-H_I cube to uncover the small-scale magnetic field in the SMC (Section 4.3). Furthermore, the scaling does not affect the study of the large-scale magnetic field orientation with $\bar{\theta}_{\text{H I}}$ (Section 4.1).

Finally, we remark that the differences between our formulation and that of Clark & Hensley (2019) are our implementation of extinction along the line of sight, as well as their incorporation of the RHT amplitude. For the former, the inclusion of the extinction term is appropriate for our comparison with starlight polarisation data, while their approach of excluding the extinction term is suitable for their comparison of the H_I4PI (HI4PI Collaboration et al. 2016) and GALFA-H_I (Peek et al. 2018) cubes with the polarised dust emission from *Planck* at 353 GHz (Planck Collaboration XIX 2015). Meanwhile for the latter, our exclusion of the RHT amplitude represents a different view in the RHT outputs compared to that of Clark & Hensley (2019), with the RHT 4D-hypercube seen as a deterministic depiction of the filament locations (i.e., any non-zero values delineate filamentary structures) rather than a probabilistic one (i.e., the RHT amplitude describes the probability of being part of an H_I filament). We repeat our analysis with the RHT amplitude incorporated into Equations 10 and 11 similar to Clark & Hensley (2019), and find that the results are almost identical, with the resulting $\bar{\theta}_{\text{H I}}$ differing by 5° in the worst case and by less than 1° on average.

4 RESULTS

4.1 Magnetic alignment of H_I filaments

To test whether magnetic alignment of H_I filaments exists in the SMC, we compute the polarisation angle difference ($\Delta\bar{\theta} = |\bar{\theta}_* - \bar{\theta}_{\text{H I}}|$) between the Lobo Gomes et al. (2015) observations (see Section 2.2) and each of our four cases of ray-tracing experiment (see Section 3). The results are listed in Table 2 and plotted in Figure 4.

We recognise a notable trend in $\Delta\bar{\theta}$ for the case of ray tracing through the low velocity range of the H_I cube (top panel of Figure 4) – the values of $\Delta\bar{\theta}$ are close to 0° for most of the fields in the SMC Bar and the start of the SMC Wing (approximately fields 1 to 11). Meanwhile, no obvious trends can be seen for the other velocity ranges. Below, we will first statistically quantify this apparent alignment in the low velocity range, followed by exhaustively investigating the potential trends of $\Delta\bar{\theta}$ with diagnostics from H_I, H α , and starlight polarisation data.

4.1.1 Statistical significance of the preferential magnetic alignment

We first compute the average $\Delta\bar{\theta}$ from ray tracing through the low velocity portion of the H_I. Considering all fields (1–20, less the uncertain fields), the mean, median, and inverse-variance weighted mean of $\Delta\bar{\theta}$ are $32^\circ \pm 2^\circ$, $20^\circ \pm 3^\circ$, and $29^\circ \pm 1^\circ$, respectively. The listed uncertainties are the corresponding standard errors. Meanwhile, considering fields 1–11 only (again excluding the uncertain fields) these three average values decrease to $25^\circ \pm 2^\circ$, $13^\circ \pm 3^\circ$, and $24^\circ \pm 2^\circ$, respectively. These are all lower than the 45° expected if the H_I filament orientation is independent of the magnetic field orientation. To




Figure 4. Starlight polarisation angle difference ($\Delta\bar{\theta}$) between the Lobo Gomes et al. (2015) observations and our ray-tracing experiment through the GASKAP-H_I cube, with the four panels showing different velocity ranges adopted for the ray tracing. The translucent data points represent fields without a coherent starlight polarisation angle from either ray tracing or the actual starlight observations, while the two horizontal dash lines at 20° and 70° represent the cutoff value adopted for alignment and anti-alignment, respectively (Sections 4.1.1).




Figure 5. Starlight polarisation angle difference ($\Delta\bar{\theta}$) between the [Lobo Gomes et al. \(2015\)](#) observations and our ray-tracing experiment through the GASKAP-HI cube, plotted against the magnetic field orientation as traced by the observed starlight polarisation. Only starlight fields 1–11 with reliably determined $\bar{p}_\star$ and $\bar{p}_{\text{HI}}$ (see Tables 1 and 2) are shown here. The two horizontal dash lines at 20° and 70° represent the cutoff value adopted for alignment and anti-alignment, respectively (Sections 4.1.1).

evaluate the statistical significance, we perform two statistical tests as described below.

First, we apply the one-sample Kolmogorov-Smirnov (KS) test to our results, comparing the $\Delta\bar{\theta}$ distributions against a uniform distribution within $[0^\circ, 90^\circ)$. Our null hypothesis is that the cumulative distribution function (CDF) of the data is less than or equal to the CDF of a uniform distribution for all $\Delta\bar{\theta}$ values, while the alternative hypothesis is that the data CDF is greater than that of a uniform distribution for at least some $\Delta\bar{\theta}$. The resulting p -values considering all fields and fields 1–11 (both excluding the uncertain fields) are 0.061 and 0.008, respectively. These indicate a preference of the alternative hypothesis above and, combined with the low average $\Delta\bar{\theta}$ determined above, suggest a preferred alignment of the HI filaments with magnetic fields in the concerned SMC volume (namely, the northeastern end of the SMC Bar and the Bar-Wing transition region).

For the second statistical test, we first draw a cutoff level of $\Delta\bar{\theta}$ at 20° , below which the HI filaments and magnetic fields are defined as aligned. This adopted value is in line with the typical degree of alignment of Galactic HI filaments with magnetic fields ([Clark et al. 2014](#)) and structures such as the Galactic plane ([Soler et al. 2020](#)). Similarly, a cutoff level at 70° can be defined, above which the two are classed as perpendicular to each other. This gives six out of 12 fields that exhibit apparent magnetic alignment of HI filaments out of all starlight fields, again excluding the uncertain fields. We then evaluate the likelihood that this alignment fraction is purely by chance drawn from a uniform distribution within $[0^\circ, 90^\circ)$. This is done by drawing 10^8 sets of 12 $\Delta\bar{\theta}$ values from such uniform distribution, and counting how many sets have at least six $\Delta\bar{\theta}$ values of less than 20° . We find that the likelihood of such chance alignment occurring is only 3.2% (i.e., p -value of 0.032). The case of fields 1–11, which corresponds to the northeastern Bar and the Bar-Wing transition region, is similarly investigated to evaluate the likelihood for six out of nine fields to show an apparent magnetic alignment. We find that this arises in only 0.5% of the cases (i.e., p -value of 0.005). These all again suggest that the agreement in orientation between HI filaments and magnetic fields is astrophysical instead of randomly by chance.




Figure 6. The preferred orientation of HI filaments, obtained by applying the ray-tracing algorithm through the full velocity range of the HI cube (Section 3; but with the extinction term turned off), shown as the flow-line pattern generated by the LIC algorithm ([Cabral & Leedom 1993](#)). The colour map shows the corresponding “polarised intensity” from ray tracing. This map can be compared with future polarised dust emission observations of the SMC, with the flow-line pattern here representing the predicted polarisation B -vector ($= E$ -vector $+90^\circ$) if HI filaments are indeed tracing the plane-of-sky magnetic field orientation.

4.1.2 Coherence of $\Delta\bar{\theta}$ between starlight fields

Next, we maintain our focus on the starlight fields where we find magnetic alignment of HI filaments, looking into their spatial distribution and relationship with the magnetic field orientation. Such information on the spatial coherence can reflect the underlying astrophysics shaping both the magnetic fields and HI structures, as well as affecting the statistical significance above (since, the analyses in Section 4.1.1 have implicitly assumed that there are no spatial correlations between different fields).

We plot the $\Delta\bar{\theta}$ against the observed starlight $\bar{\theta}_\star$ in Figure 5, with the six fields demonstrating magnetic alignment of HI filaments ($\Delta\bar{\theta} < 20^\circ$) being fields 2, 3, 8, 9, 10, and 11. These fields cover a broad range in the observed $\bar{\theta}_\star$ that traces the magnetic field orientation, from 52.6° to 156.0° . In particular, we note rapid angle changes between nearby fields – from $52.6^\circ \pm 4.1^\circ$ to $106.0^\circ \pm 4.5^\circ$ from field 2 to 3, and from $69.0^\circ \pm 2.3^\circ$ to $142.3^\circ \pm 4.7^\circ$ from field 9 to 10, both across $15' = 270$ pc. Overall, despite the strongly fluctuating magnetic field orientation amongst these fields, the HI filament orientations seem to remain preferentially aligned with their respective local magnetic field.

4.1.3 Correlation with other diagnostics

Finally, we wrap up our investigation in $\Delta\bar{\theta}$ by looking into the potential physical properties of the fields that may have led to the alignment or misalignment of the HI filaments with the magnetic field. Diagnostics are derived from the new GASKAP-HI data ([Pingel et al. 2022](#)), the starlight polarisation data ([Lobo Gomes et al. 2015](#)), and the data from the Wisconsin H α Mapper (WHAM) survey ([Smart et al. 2019](#)). For HI and H α , we compare the $\Delta\bar{\theta}$ values against the moment 0 (velocity-integrated intensity), moment 1 (mean velocity), and moment 2 (velocity dispersion), as well as visually inspecting the




Figure 7. The preferred orientation of H I filaments in the low-velocity (presumably near-side; upper panel) and the high-velocity (presumably far-side; lower panel) portions of the SMC, outlined by the green tick marks. Only sightlines with SMC H I column densities of higher than 10^{21} cm^{-2} are considered here. The background greyscale maps are the peak H I intensity maps from GASKAP-H I (Pingel et al. 2022).

velocity profiles for each field. Meanwhile for starlight polarisation, we compare $\Delta\theta$ against $\bar{p}_\star$ and $\sigma_{p\star}$ (Section 2.2). We do not find any notable trends with any of these parameters.

4.2 The preferred orientation of H I filaments in the SMC

We use the results from applying the RHT algorithm to the GASKAP-H I data to further obtain the preferred orientation of H I filaments across the SMC. This can allow us to identify the astrophysical processes shaping the H I structures of this galaxy, and can also be used to compare against future observations of the SMC magnetic fields to further test the magnetic alignment of H I filaments, as elaborated in Sections 5.3 and 5.5.

First, we produce the preferred orientation map of H I filaments by




Figure 8. Map of the angle difference of the preferred H I filament orientations between the low- and high-velocity portions of the SMC (see Figure 7), shown as the colour dots. The background greyscale map shows the peak H I intensity map.




Figure 9. Histogram of the angle difference of the preferred H I filament orientations between the low- and high-velocity portions of the SMC (see Figures 7 and 8).

combining the full radial velocity range of the SMC. Specifically, we follow the ray-tracing analysis procedures as described in Section 3, but performed for each pixel of the GASKAP-H I map instead of on a per-star basis. Furthermore, we have turned off the dust extinction effect (i.e., $A_V(v_j) = 0$ for all v_j in Equation 12). The resulting Stokes Q and U maps are smoothed to $8'$ to extract the underlying $\approx 150 \text{ pc}$ -scale pattern. The resulting “polarised intensity” map is shown in Figure 6, with the corresponding position angle tracing the preferred orientation of H I filamentary structures shown as the flow-line pattern produced by the Line Integral Convolution (LIC) algorithm (Cabrál & Leedom 1993). The operations here are similar to that of Clark & Hensley (2019), which studied the case of the Milky Way by comparing the HI4PI cube (HI4PI Collaboration et al. 2016) with *Planck* polarised dust emission (Planck Collaboration XIX 2015) (see also Section 3).

Next, we produce preferred orientation maps separately for the low-velocity ($v < v_{\text{mean}}$; presumably near-side) and the high-velocity ($v \geq v_{\text{mean}}$; presumably far-side) portions of the H I cube. This is again done by a per-pixel ray tracing through the H I cube without dust extinction ($A_V(v_j) = 0$), but restricted in velocity space accordingly separated by v_{mean} . The 2D spatial domain is then divided into independent boxes of $70 \times 70 \text{ px} \approx 8' \times 8'$, within which the Stokes Q and U values are summed and subsequently converted to θ . This operation is only performed for boxes with a hydrogen column density above 10^{21} cm^{-2} in order to focus on the main gaseous body of the SMC. The θ maps for the two portions of the SMC are plotted in Figure 7, the angle difference map is plotted in Figure 8, and the histogram of angle difference is shown in Figure 9.

Finally, we generate a preferred orientation map of the H I filaments for the low-velocity portion (presumably near-side) of the SMC only, with the attenuation of starlight flux density due to dust extinction taken into account (Equations 10–12). The resulting Stokes Q and U maps are again smoothed to $\approx 8'$, and the corresponding position angle map is shown in Figure 10 as the flow-line pattern from LIC over the Digitized Sky Surveys 2 (DSS2; Lasker et al. 1996) optical image of the SMC. This map can be compared to future starlight polarisation observations.

4.3 Relationship between ray-traced and observed $\sigma_p/\bar{p}$

As pointed out in Section 2.2, the ratio between $\sigma_{p\star}$ and $\bar{p}_\star$ of the observed starlight can be indicative of the relative strength between the small- ($\ll 100 \text{ pc}$) and large-scale ($\gg 100 \text{ pc}$) magnetic field. It is therefore of interest to explore whether the H I data have sufficient angular resolution to enable similar measurements. This can be evaluated by seeing whether the $\sigma_{p\text{H I}}/\bar{p}_{\text{H I}}$ parameter from the ray-traced starlight polarisation corresponds well with the observed $\sigma_{p\star}$ and $\bar{p}_\star$. We plot the ray-traced against observed $\sigma_p/\bar{p}$ of fields 1–11 through the low velocity portion of the GASKAP-H I cube in Figure 11, and we do not find good agreement between the two sets of $\sigma_p/\bar{p}$ values. We further note that for all of the fields except 2 and 11, the ray-traced $\sigma_{p\text{H I}}/\bar{p}_{\text{H I}}$ values are lower than the observed counterparts. The mismatch between the two is likely due to the limited spatial resolution of the GASKAP-H I data (see Section 5.2.2).

5 DISCUSSION

5.1 The physical nature of the H I filaments

5.1.1 Physical scales of the H I filaments

We first consider the physical scales of the SMC H I filaments that we study here, and attempt to identify analogues in the Milky Way. Given the spatial resolution of $30'' = 9 \text{ pc}$ of the GASKAP-H I observations (Pingel et al. 2022) and our chosen RHT parameters (specifically, $D_W = 83 \text{ px} = 175 \text{ pc}$ and $\text{threshold} = 0.7$), the H I filamentary structures that we uncover have widths of $\sim 9 \text{ pc}$ and lengths of $\gtrsim 120 \text{ pc}$. This is clearly in a vastly different physical scale range compared to the individual H I filaments on the local bubble wall with widths $\lesssim 0.1 \text{ pc}$ and lengths $\sim \text{few pc}$ (Clark et al. 2014). However, as pointed out in their work, the Clark et al. (2014) filaments are highly spatially correlated in orientation, together forming coherent bundles of filaments across 10s of pc. Meanwhile, high spatial resolution studies of discrete elongated H I clouds in the Milky Way have found that they can be further composed of fine strands of H I filaments. For example, the Riegel-Crutcher cloud exhibits as an elongated H I cloud with a width of $\approx 5 \text{ pc}$ and a length of $\approx 20 \text{ pc}$, and is constituted by

countless H I filaments with widths of $\lesssim 0.1 \text{ pc}$ (McClure-Griffiths et al. 2006). Similarly, the local velocity cloud towards the Ursa Major cirrus (Skalidis et al. 2022) has a width of $\approx 5 \text{ pc}$ and a length of $\approx 15 \text{ pc}$, and is formed by groups of $\lesssim 1 \text{ pc}$ -wide H I filaments. Combining all the examples above, we hypothesise that the H I filamentary structures in the SMC uncovered by our GASKAP-H I observations may actually be bundles of fine H I filaments with coherent orientations across $\gtrsim 120 \text{ pc}$. Alternatively, we can be tracing the general anisotropic features of the H I gas as seen in the Galactic plane (e.g., Soler et al. 2022).

Finally, we note the existence of several gigantic filamentary H I structures in the Milky Way. These can be seen in, for example, the Canadian Galactic Plane Survey (CGPS) data ($\approx \text{few degrees}$ in length; see Gibson 2010) as well as the GALFA-H I data ($\approx 10 \text{ deg}$ in length; Peek et al. 2018). In addition, there is a recent discovery of an enormous Galactic H I filament with width $\approx 50 \text{ pc}$ and length $\approx 1,200 \text{ pc}$ (the ‘‘Maggie’’ filament; Soler et al. 2020; Syed et al. 2022). Similar H I filaments can account for a small fraction of our SMC H I filaments, but given the apparent rarity of such class of H I filaments in the Milky Way we deem it improbable as the primary explanation for most of the SMC filaments.

5.1.2 The alignment with magnetic fields

We report and statistically assess the alignment of H I filaments in the SMC with magnetic fields in Section 4.1, with the trend seen in the northeastern Bar and the beginning of the Wing region (approximately fields 1–11). The rest of the SMC volume lacks starlight polarisation data of sufficient quality to draw conclusions. This is the first time that such a relation of magnetically aligned H I filaments has been identified beyond the Milky Way, enabled by the unprecedented combination of the angular resolution, velocity resolution, and surface brightness sensitivity of the new GASKAP-H I data. The results suggest that magnetically aligned H I filaments may also be seen beyond the Milky Way, which is a key piece of information for future numerical studies of the astrophysics governing the formation of these filamentary H I structures.

The ability of the GASKAP-H I data to see magnetic alignment of filaments at all is, in fact, somewhat surprising. Clark et al. (2014) has explored the effects of the spatial resolution of the H I data on the alignment of the subsequently identified filamentary structures with magnetic fields. Upon comparing the results using GALFA-H I data at a resolution of $4' = 0.1 \text{ pc}$ (Peek et al. 2011) with those from GASS data at a resolution of $16' = 0.5 \text{ pc}$ (McClure-Griffiths et al. 2009), they have found that the degree of alignment can worsen from within $\approx 16^\circ$ from the former to within $\approx 36^\circ$ from the latter. Meanwhile, our spatial resolution of the SMC in H I is $30'' = 9 \text{ pc}$, significantly worse than even the GASS data. We suspect that the key here is to have matching spatial scales traced by both the H I and the starlight polarisation data. In particular, we opt for an analysis on a per-field basis, with both data sets tracing $\approx 150 \text{ pc}$ scales. Meanwhile, the Clark et al. (2014) analyses studied much smaller scales ($\lesssim 0.1 \text{ pc}$) in both data sets.

Finally, as pointed out in Section 4.1.2, the starlight fields in the SMC Bar region exhibit a range of plane-of-sky magnetic field orientation across $\sim 100 \text{ pc}$ scale (see Section 5.2 for more discussions). Despite such a rapidly varying magnetic structure, the H I filament orientation remains following the magnetic fields. This strongly suggests that we are not looking at a chance alignment of the two across multiple spatially correlated starlight fields, and the statistical evaluation of the magnetic alignment in Section 4.1.1 is valid.




Figure 10. Result of ray tracing through the low velocity portion (presumably the near-side of the SMC) of the GASKAP-H I data cube, with the effect of starlight attenuation due to dust extinction taken into account. The position angle is shown as the flow-line pattern generated by the LIC algorithm (Cabral & Leedom 1993). This can be compared with future starlight polarisation observations to test whether H I filaments are aligned with the magnetic field throughout the entirety of the SMC. The colour optical map is from the DSS2 (Lasker et al. 1996).




Figure 11. Ray-traced against observed values of $\sigma_p / \bar{p}$. Only starlight fields 1–11 with reliably determined $\bar{p}$ (see Tables 1 and 2) are shown here. The diagonal grey line marks where the ray-traced and observed values agree.

5.2 The magnetic field structure of the SMC

5.2.1 Information from starlight polarisation

In the Bar region of the SMC (defined here as fields 1–9), we do not find any consistent trends in the observed starlight polarisation amongst the starlight fields. This shows that the large-scale magnetic fields in the Bar are not ordered on scales much larger than that corresponding to the field-of-view of the starlight observations ($8' \approx 150$ pc). Meanwhile, within each of the starlight fields we find consistent coherent starlight polarisation signals ($\bar{p}_\star$ and $\bar{\theta}_\star$) indicative of an ordered magnetic field on ≈ 150 pc scale. Finally, the large scatter of the per-field starlight polarisation ($\sigma_{p\star}$) is a manifestation of the strong turbulent magnetic field on $\ll 150$ pc scale. All these can be explained by the perturbation of the magnetic field by some $\gtrsim 150$ pc structures (e.g., supershells that are known to be ubiquitous in the SMC; Staveley-Smith et al. 1997, and tidal forces), as well as the injection of turbulent energy at $\lesssim 10$ pc scale by stellar feedback processes (e.g., MacLow 2004).

The above interpretation is in an *apparent* conflict with the conclusion from studies of RM of extragalactic sources behind the SMC that found a coherent magnetic field pointed away from the observer consistently across the entire SMC Bar (Mao et al. 2008; Livingston et al. 2022), which would not have been observed if there is no coherent magnetic field on $\gg 150$ pc scale. However, we point out that the lack of an ordered⁷ plane-of-sky magnetic field does not necessarily imply a lack of a coherent line-of-sight magnetic field. For example, imagine an initial perfectly coherent magnetic field along the line-of-sight only, with the magnetic field strength component in the plane-of-sky being zero. If this frozen-in magnetic field is then perturbed by turbulence, the resulting magnetic field configuration can have a significant unordered plane-of-sky component, while still preserving some degree of coherence along the line-of-sight.

We next move onto the SMC Wing region (defined here as fields 10–20), which is believed to have formed due to tidal interactions with the LMC. Four (namely, 10, 11, 12, and 15) of the five fields that individually show coherent starlight polarisation signals exhibit a consistent magnetic field orientation with $\theta \approx 150^\circ$ across ≈ 1 kpc, with the remaining field (20) situated far from the SMC Bar (about $3^\circ \approx 3.2$ kpc in projected distance) having a distinct $\theta \approx 70^\circ$. The trend of $\theta \approx 150^\circ$ has also been pointed out by Lobo Gomes et al. (2015) using their analysis methods (trend III in their section 5). The magnetic fields are oriented along the general elongation of the SMC Wing itself but with a slight offset of $\approx 20^\circ$. Combined with the recent results of a general negative RM of extragalactic sources behind the SMC Wing (Livingston et al. 2022), we obtain a picture of a coherent magnetic field on scales $\gtrsim 1$ kpc along the Wing’s elongation. This resembles the ~ 20 kpc tidal tail of the Antennae galaxies, which was found through studying its synchrotron emission to host a regular magnetic field along its entirety, believed to have come from tidal stretching of the original disk magnetic fields (Basu et al. 2017). Apart from the concerned physical scales, one key difference between the two cases is the strength ratio between the large-scale regular and the small-scale turbulent magnetic fields: the former is believed to be stronger for the case of the Antennae galaxies tidal tail (Basu et al. 2017), while the latter likely dominates in the SMC Wing on $\ll 150$ pc scale as reflected by the consistently high $\sigma_{p\star}$ compared to $\bar{p}_\star$ for all of the starlight fields.

Finally, we point out again and further discuss a common characteristic in both the Bar and Wing regions of the SMC – the $\ll 150$ pc turbulent magnetic field strength is much higher than the ordered magnetic field strength, as inferred from the consistently high $\sigma_{p\star}/\bar{p}_\star$ ratio in all observed starlight fields. This is in agreement with the conclusions of numerous previous studies of the SMC magnetic fields (e.g., Mao et al. 2008; Lobo Gomes et al. 2015; Livingston et al. 2022), in line with our knowledge of the highly complex neutral and ionised gas dynamics in the SMC (e.g., Le Coarer et al. 1993; Staveley-Smith et al. 1997; Smart et al. 2019), and in contrast with most spiral galaxies that have comparable strengths between the turbulent and ordered components (see e.g., Beck & Wielebinski 2013; Beck 2016). Given that the $\sigma_{p\star}$ values are larger than the corresponding $\bar{p}_\star$ values, we argue that the traditional Davis-Chandrasekhar-Fermi method (Davis 1951; Chandrasekhar & Fermi 1953), which uses the spread of starlight polarisation angle as a measure of the turbulent-to-ordered magnetic field strength, cannot be

directly applied to the SMC. This is because the starlight polarisation angles span the full 180° , meaning that the angle spread loses physical meaning.

5.2.2 H I filaments as a tracer of the small-scale magnetic field

We explore using our ray-tracing analysis method on the GASKAP-H I cube to obtain information on the small-scale magnetic field in Section 4.3, and do not find a good match between the ray-traced and observed $\sigma_p/\bar{p}$ values, with the former underestimating the latter in most of the starlight fields. We discuss the possible reasons behind this mismatch below.

The most probable reason behind the low ray-traced $\sigma_{p\text{H I}}/\bar{p}_{\text{H I}}$ values is the limited spatial resolution of the GASKAP-H I data. The $30'' = 9$ pc resolution of the data sets the absolute minimum scale that our study is sensitive to, while our RHT parameter choice of $R_{\text{sm}} = 12 \text{ px} = 25 \text{ pc}$ may further coarsen the effective resolution. If the spatial resolution of our data is comparable to or poorer than the outer scale of turbulence in the SMC, the map of filaments identified by the RHT algorithm and thus our ray-tracing analysis may not be able to capture the corresponding intricate features that actual starlight polarisation data can. Recent spatial power spectrum and structure function analyses of the SMC in H I have concluded that the turbulence is being driven on a very large (galactic) scale (Szkotkowski et al. 2019), suggesting that our H I data at ≈ 9 pc resolution should well resolve the turbulent structures in the SMC spatially. Meanwhile, the RM structure function using extragalactic sources behind the SMC suggested an upper limit to the outer scale of turbulence in the SMC of 250 pc (Livingston et al. 2022), and similar studies through the Milky Way disk have indicated outer scales of turbulence of ~ 10 pc in the spiral arms and ~ 100 pc in the interarm regions (Haverkorn et al. 2008). All these could be reconciled if the *magnetic* outer scale of turbulence that can be resolved by the starlight polarisation data but not our ray-tracing analysis are much smaller than that of the gas density. This would mean that the physical conditions portrayed by the H I filaments may be less turbulent and more coherent than the actual reality traced by observed starlight polarisation, leading to the lower $\sigma_{p\text{H I}}/\bar{p}_{\text{H I}}$ values from our ray-tracing analysis.

We further consider whether the polarised dust extinction in the Milky Way can be a reasonable explanation to the mismatch in $\sigma_p/\bar{p}$. The procedure of Galactic foreground removal by the Lobo Gomes et al. (2015) starlight polarisation catalogue concerns the large-scale coherent component only, while the contributions by the turbulent magnetic fields in the Milky Way (if present) cannot be removed on a per-star basis due to the stochastic nature. The Galactic foreground can therefore introduce extra scatter in the observed starlight polarisation (i.e., higher $\sigma_{p\star}$ and therefore $\sigma_{p\star}/\bar{p}_\star$), but not to the ray-traced starlight since we did not take the Galactic contributions into account. Along the line of sight towards the SMC (Galactic latitude: $b = -44.3^\circ$), the approximate path lengths through the Galactic H I thin and thick disks (with half-widths of ~ 100 and ~ 400 pc, respectively; Dickey 2013) are 140 and 570 pc, respectively. At these distances, the $8'$ field-of-view of the starlight polarisation observations convert to about 0.3 and 1.3 pc, respectively. These values are much smaller than the ~ 10 – 100 pc outer scale of turbulence in the Milky Way (Haverkorn et al. 2008). As the large-scale and small-scale magnetic fields are of comparable strengths in the Milky Way (e.g., Beck 2016), the turbulent magnetic field must be significantly weaker than the large-scale counterpart at scales much smaller than the outer scale of turbulence. Therefore we deem this unlikely as the primary explanation of the mismatch in $\sigma_p/\bar{p}$.

⁷ The distinction between a coherent and an ordered magnetic field being that, the former has a constant magnetic field *direction* (without any flips), while the latter only need to have a constant *orientation* (flips in direction are permitted; e.g., Jaffe et al. 2010; Beck 2016).

5.3 The preferred orientation of H I filaments

Moving along the Bar from the northeastern end ($RA \approx 1^h 10^m$, $Decl \approx -71.5^\circ$) to the southwestern end ($RA \approx 0^h 45^m$, $Decl \approx -73.5^\circ$), we identify three distinct regions. First, the H I filaments are preferentially oriented along the elongation of the Bar, seen in both the low- and high-velocity ranges (Figure 7). Noting that the H I velocity gradient is also oriented northeast-southwest along the elongation of the SMC Bar (Di Teodoro et al. 2019), the orientation of the H I filaments here can be controlled by the gas dynamics in the galaxy, similar to the case of the disk-parallel H I filaments in the Milky Way (Soler et al. 2022). However, we point out that the internal gas dynamics of the SMC may be much more complex than that revealed by H I data alone (see Murray et al. 2019). Second, at $RA \approx 0^h 55^m$, $Decl \approx -72.3^\circ$, the H I filaments switch in the preferred orientation abruptly to be along northwest-southeast. This is seen in the low-velocity portion only, and is lined up with the SMC Wing to the southeast. These H I structures here are likely shaped by the tidal stretching from interactions with the LMC that have also formed the SMC Wing and the Magellanic Bridge (Besla et al. 2012; Wang et al. 2022). We further note that in this same sky area within the SMC, the stellar proper motion (Niederhofer et al. 2021) that is believed to be tracing the effect of tidal stretching exhibits a consistent direction as our H I filament orientation. Third and finally, starting from $RA \approx 0^h 55^m$, $Decl \approx -72.5^\circ$ to the southwest, the filaments are preferentially oriented east-west, with a significant perpendicular component to the Bar elongation, seen most clearly in the low-velocity and also in the high-velocity. This can be shaped by feedback processes from star formation that transport gas away from the galaxy into the circumgalactic medium. To summarise, the preferred orientation of H I filaments across the SMC Bar shows highly complex geometries, possibly shaped by multiple astrophysical processes that the SMC is subjected to.

Meanwhile, we note that the preferred H I filament orientation in the SMC Wing also exhibits highly complex structures. Overall, it appears as though the H I filaments are wrapping around the elongated structure of the SMC Wing.

5.4 The 3D structure of the SMC

In Section 4.1, we find moderate evidence for alignment between the orientation of starlight polarisation and that of the H I filaments in the low velocity portion of the northeastern Bar and the start of the Wing regions. In comparison, the match with other H I velocity ranges (high velocity portion, as well as the full velocity range in both ray-tracing directions) are considerably poorer. This information can be used to help decipher the complex 3D structure of the SMC (see, e.g., Panopoulou et al. 2021, and references therein for similar cases in the Milky Way). In particular, since starlight polarisation is induced by the foreground dusty ISM, our results suggest that the aforementioned SMC regions are physically closer to us than the higher velocity portion. This is in agreement with the result of Mathewson et al. (1986), which found that the radial velocities of the sample of 26 SMC stars are consistently higher than that of the associated Ca II absorption from the SMC ISM. Assuming that their stellar radial velocities correspond to the ambient gas radial velocities, this would mean that the lower velocity gas component of the SMC is physically closer to us. The same conclusion has also been reached from the many newer optical and/or ultraviolet absorption line studies (e.g., Danforth et al. 2002; Welty et al. 2012).

We note that the remaining areas of the SMC, namely the southwestern end of the Bar and the majority of the Wing, remain relatively

unexplored. Future, deep starlight polarisation surveys covering the entirety of the SMC will be key to unravelling the overall 3D structure of the gaseous component of this galaxy.

Finally, we identify from Figures 8 and 9 that filamentary H I orientations in the low- and high-velocity portions of the SMC are similar across large areas in both the Bar and the Wing regions. The mean and median θ differences are about 35° and 30° , respectively, with 35 % of the evaluated areas having θ differences of less than 20° . We further perform a one-sample KS test against a uniform distribution (similar to Section 4.1.1) and obtain a p -value of 2×10^{-23} . This suggests that the two velocity components of the SMC are physically linked.

5.5 Future prospects

5.5.1 Ray-tracing analysis

To enable a detailed comparison between the new GASKAP-H I data of the SMC (Pingel et al. 2022) and starlight polarisation data (Lobo Gomes et al. 2015), we develop the new ray-tracing analysis method (Section 3), with which we establish the alignment of H I filaments with the ≈ 150 pc-scale magnetic field in the SMC Bar region (Section 4.1). The same analysis method can be applied to similar future Galactic and Magellanic studies using recent and future data such as:

- Starlight polarisation: SOUTH-POL (Magalhães et al. 2012), Polar-Areas Stellar-Imaging in Polarization High-Accuracy Experiment (PASIPHAIE; Tassis et al. 2018), and the Galactic Plane Infrared Polarization Survey (GPIPS; Clemens et al. 2020);
- H I emission: GALFA-H I (Peek et al. 2011, 2018), GASKAP-H I (Dickey et al. 2013), The H I/OH/Recombination line survey of the Milky Way (THOR; Beuther et al. 2016), and the Dominion Radio Astrophysical Observatory (DRAO) H I Intermediate Galactic Latitude Survey (DHIGLS; Blagrove et al. 2017);
- Diffuse synchrotron emission: the Global Magneto-Ionic Medium Survey (GMIMS; Wolleben et al. 2009), the Polarisation Sky Survey of the Universe's Magnetism survey (POSSUM; Gaensler et al. 2010), the LOFAR Two-metre Sky Survey (LoTSS; Shimwell et al. 2017), the C-Band All Sky Survey (C-BASS; Jones et al. 2018), and the S-band Polarization All Sky Survey (S-PASS; Carretti et al. 2019).

In particular, our work here concludes that the $\sigma_{pH I} / \bar{p}_{H I}$ parameter from GASKAP-H I observations cannot be used to trace the small-scale magnetic field of the SMC, because of the lack of spatial resolution. We plan to apply the same analysis to future GASKAP-H I data of the Milky Way. The much higher ($\lesssim 1$ pc) spatial resolution will allow us to test whether the H I data can be used as a good tracer of the turbulent magnetic field in the ISM.

5.5.2 Polarised starlight and dust emission of the SMC

We identify the preferential alignment of H I filaments with the magnetic fields traced by starlight polarisation in the northeastern end of the Bar region and the Bar-Wing transition region of the SMC. Subsequently, we use the GASKAP-H I data to produce maps of the preferred orientation of H I filaments across the SMC (Figures 6–10). These maps can be compared with future starlight polarisation and polarised dust emission data for further direct confirmation of the alignment of these H I structures with the magnetic field.

In particular, the starlight data can be compared with the H I emission on the near side of the SMC (Figure 10). This is especially intriguing in the SMC Wing, as this can shed light on both its 3D

(Section 5.4) and magnetic (Section 5.3) structures that are still not fully explored.

New, high spatial resolution observations of the polarised dust emission that probes the entire line of sight through the SMC, using forthcoming instruments such as the Prime-cam (CCAT-Prime Collaboration et al. 2023) and the Simons Observatory (Hensley et al. 2022), similar to the few other nearby galaxies observed with the Stratospheric Observatory for Infrared Astronomy (SOFIA) High-resolution Airborne Wideband Camera Plus (HAWC+ Jones et al. 2020; Lopez-Rodriguez et al. 2022), can be compared with our H I filament orientation map (Figure 6). This will test whether the magnetic alignment of H I filaments persists through the full SMC volume. If this will be confirmed, we will have higher confidence in using the GASKAP-H I data for a tomographic view of the SMC’s plane-of-sky magnetic field structure. Our Stokes $Q(v)$ and $U(v)$ cubes prior to the ray-tracing steps above retain the information of magnetic fields along the line of sight decomposed by radial velocity (see, e.g., Clark 2018; Clark & Hensley 2019). These can be compared with the POSSUM data (Gaensler et al. 2010) also from ASKAP that measures the polarised synchrotron emission. Applications of continuum polarimetric techniques such as RM-Synthesis (Brentjens & de Bruyn 2005) to broadband spectro-polarimetric data can similarly decompose the polarised synchrotron emission by Faraday depth (see, e.g., Van Eck et al. 2019). These two ASKAP datasets can enable 3D-3D comparisons that can lead to crucial knowledge in how magnetic fields link the diffuse ISM probed by the polarised continuum emission and the neutral ISM probed by the H I filaments. This will be a major step forward compared to the recent work on M 51 in the 2D-2D domain (Fletcher et al. 2011; Kierdorf et al. 2020; Borlaff et al. 2021).

6 CONCLUSIONS

We investigate whether the H I filaments in the SMC are aligned with the magnetic fields, as is the case in the solar neighbourhood in the Milky Way (e.g., McClure-Griffiths et al. 2006; Clark et al. 2014; Clark & Hensley 2019). Our work has been enabled by the new, sensitive, high resolution H I observations using the ASKAP telescope (Hotan et al. 2021) by the GASKAP-H I survey (Dickey et al. 2013; Pingel et al. 2022), in addition to the recently released starlight polarisation catalogue of the SMC (Lobo Gomes et al. 2015). The RHT algorithm (Clark et al. 2014; BICEP/Keck Collaboration et al. 2022) is applied to the GASKAP-H I cube to automatically identify filamentary structures, and the Lobo Gomes et al. (2015) data are re-analysed with a vector approach to extract the large- and small-scale magnetic field information.

We devise a new ray-tracing analysis to perform a careful comparison between the H I filament orientation and the starlight polarisation data, and find a preferential alignment of the low radial velocity H I filaments with the large-scale magnetic fields traced by starlight polarisation in two regions of the SMC: the northeastern end of the Bar region, and the Bar-Wing transition region. The remainder of the Bar region, as well as the Wing region, do not yet have sufficient coverage by starlight polarisation observations for such detailed comparisons with H I data. This is the first time that the alignment of H I filaments with the ambient magnetic field is seen across large spatial volume ($\gtrsim 1$ kpc) and outside of the Milky Way. The results further suggest that the lower velocity H I component in the SMC Bar and Bar-Wing transition area is physically closer to us than the higher velocity component, consistent with previous findings (Mathewson et al. 1986; Danforth et al. 2002; Welty et al. 2012).

We produce maps tracing the preferred orientation of H I filaments across the SMC, revealing the highly complex structures likely shaped by a combination of the intrinsic internal gas motion of the SMC, tidal forces from the LMC, and stellar feedback mechanisms. These maps can further be compared with future measurements of the magnetic field structure of the SMC from starlight and dust polarisation, as well as with the diffuse polarised synchrotron emission from POSSUM (Gaensler et al. 2010). We also find that the orientation of the H I structures between the low- and high-velocity portions of the SMC are similar, suggesting that the two velocity components are physically linked.

ACKNOWLEDGEMENTS

We thank the anonymous referee for the comments, especially on the discussions on the statistical robustness of the bootstrapping procedures. We thank Christoph Federrath, Isabella Gerrard, Gilles Joncas, Marc-Antoine Miville-Deschênes, Snežana Stanimirović, and Josh Peek for the fruitful discussions on this work. We thank Rainer Beck for the careful reading of the manuscript and the thoughtful suggestions that have improved the presentation of this paper. YKM thanks Michael Kramer and Sui Ann Mao for their gracious extended host at the Max-Planck-Institut für Radioastronomie in Bonn, Germany. This research was partially funded by the Australian Government through the Australian Research Council. LU acknowledges support from the University of Guanajuato (Mexico) grant ID CIIC 164/2022. This scientific work uses data obtained from Inyarrimanha Ilgari Bundara / the Murchison Radio-astronomy Observatory. We acknowledge the Wajarri Yamaji People as the Traditional Owners and native title holders of the Observatory site. The Australian SKA Pathfinder is part of the Australia Telescope National Facility which is managed by CSIRO. Operation of ASKAP is funded by the Australian Government with support from the National Collaborative Research Infrastructure Strategy. ASKAP uses the resources of the Pawsey Supercomputing Centre. Establishment of ASKAP, the Murchison Radio-astronomy Observatory and the Pawsey Supercomputing Centre are initiatives of the Australian Government, with support from the Government of Western Australia and the Science and Industry Endowment Fund. The Parkes radio telescope is part of the Australia Telescope National Facility which is funded by the Australian Government for operation as a National Facility managed by CSIRO. We acknowledge the Wiradjuri people as the traditional owners of the Observatory site.

DATA AVAILABILITY

The GASKAP-H I Pilot Survey data are available on the CSIRO ASKAP Science Data Archive⁸ (CASDA). The auxiliary data products from this article will be shared on reasonable request to the corresponding author.

REFERENCES

- Aab A., et al., 2015, *ApJ*, **804**, 15
- Andersson B.-G., Lazarian A., Vaillancourt J. E., 2015, *ARA&A*, **53**, 501
- BICEP/Keck Collaboration et al., 2022, arXiv e-prints, p. arXiv:2210.05684

⁸ <https://research.csiro.au/casda/>.

- Basu A., Mao S. A., Kepley A. A., Robishaw T., Zweibel E. G., Gallagher John. S. I., 2017, *MNRAS*, **464**, 1003
- Beck R., 2016, *A&ARv*, **24**, 4
- Beck R., Wielebinski R., 2013, in Oswalt T. D., Gilmore G., eds, *Planets, Stars and Stellar Systems. Vol. 5: Galactic Structure and Stellar Populations*. Springer, Berlin, p. 641, doi:10.1007/978-94-007-5612-0_13
- Beck R., Ehle M., Shoutenkov V., Shukurov A., Sokoloff D., 1999, *Nature*, **397**, 324
- Beck R., Fletcher A., Shukurov A., Snodin A., Sokoloff D. D., Ehle M., Moss D., Shoutenkov V., 2005, *A&A*, **444**, 739
- Besla G., Kallivayalil N., Hernquist L., van der Marel R. P., Cox T. J., Kereš D., 2012, *MNRAS*, **421**, 2109
- Beuther H., et al., 2016, *A&A*, **595**, A32
- Birnboim Y., Balberg S., Teyssier R., 2015, *MNRAS*, **447**, 3678
- Blagrove K., Martin P. G., Joncas G., Kothés R., Stil J. M., Miville-Deschênes M. A., Lockman F. J., Taylor A. R., 2017, *ApJ*, **834**, 126
- Borlaff A. S., et al., 2021, *ApJ*, **921**, 128
- Brentjens M. A., de Bruyn A. G., 2005, *A&A*, **441**, 1217
- Brüns C., et al., 2005, *A&A*, **432**, 45
- CCAT-Prime Collaboration et al., 2023, *ApJS*, **264**, 7
- Cabral B., Leedom L. C., 1993, in , *Proc. 20th Annual Conference on Computer Graphics and Interactive Techniques*. (ACM, New York, NY), p. 263, doi:10.1145/166117.166151
- Campbell J. L., et al., 2022, *ApJ*, **927**, 49
- Carretti E., et al., 2019, *MNRAS*, **489**, 2330
- Chan M. H., Del Popolo A., 2022, *MNRAS*, **516**, L72
- Chandrasekhar S., Fermi E., 1953, *ApJ*, **118**, 113
- Choudhury S., Subramaniam A., Cole A. A., Sohn Y. J., 2018, *MNRAS*, **475**, 4279
- Clark S. E., 2018, *ApJ*, **857**, L10
- Clark S. E., Hensley B. S., 2019, *ApJ*, **887**, 136
- Clark S. E., Peek J. E. G., Putman M. E., 2014, *ApJ*, **789**, 82
- Clark S. E., Hill J. C., Peek J. E. G., Putman M. E., Babler B. L., 2015, *Phys. Rev. Lett.*, **115**, 241302
- Clemens D. P., et al., 2020, *ApJS*, **249**, 23
- Contopoulos G., Jappel A., 1974, *Transactions of the IAU XV B: Proceedings of the Fifteenth General Assembly, Sydney 1973 and Extraordinary Assembly, Poland 1973*. (Dordrecht: Reidel)
- Danforth C. W., Howk J. C., Fullerton A. W., Blair W. P., Sembach K. R., 2002, *ApJS*, **139**, 81
- Davis L., 1951, *Physical Review*, **81**, 890
- Di Teodoro E. M., et al., 2019, *MNRAS*, **483**, 392
- Dickey J. M., 2013, in Oswalt T. D., Gilmore G., eds, *Planets, Stars and Stellar Systems, Vol. 5, Galactic Structure and Stellar Populations*. (Springer, Dordrecht), p. 549, doi:10.1007/978-94-007-5612-0_11
- Dickey J. M., et al., 2013, *Publ. Astron. Soc. Australia*, **30**, e003
- Duda R. O., Hart P. E., 1972, *Commun. Assoc. Comput. Mach.*, **15**, 11
- Elstner D., Beck R., Gressel O., 2014, *A&A*, **568**, A104
- Federrath C., 2016a, *Journal of Plasma Physics*, **82**, 535820601
- Federrath C., 2016b, *MNRAS*, **457**, 375
- Federrath C., Klessen R. S., 2012, *ApJ*, **761**, 156
- Fletcher A., Beck R., Shukurov A., Berkhuijsen E. M., Horellou C., 2011, *MNRAS*, **412**, 2396
- Frick P., Stepanov R., Beck R., Sokoloff D., Shukurov A., Ehle M., Lundgren A., 2016, *A&A*, **585**, A21
- Gaensler B. M., Landecker T. L., Taylor A. R., POSSUM Collaboration 2010, in *American Astronomical Society Meeting Abstracts #215*. p. 470.13
- Gaia Collaboration et al., 2022, arXiv e-prints, p. arXiv:2208.00211
- Gazol A., Villagran M. A., 2021, *MNRAS*, **501**, 3099
- Gibson S. J., 2010, in Kothés R., Landecker T. L., Willis A. G., eds, *Astronomical Society of the Pacific Conference Series Vol. 438, The Dynamic Interstellar Medium: A Celebration of the Canadian Galactic Plane Survey*. p. 111 (arXiv:1009.0966)
- Gordon K. D., et al., 2011, *AJ*, **142**, 102
- Graczyk D., et al., 2020, *ApJ*, **904**, 13
- Green D. A., 2011, *Bulletin of the Astronomical Society of India*, **39**, 289
- HI4PI Collaboration et al., 2016, *A&A*, **594**, A116
- Hall J. S., 1949, *Science*, **109**, 166
- Haverkorn M., Brown J. C., Gaensler B. M., McClure-Griffiths N. M., 2008, *ApJ*, **680**, 362
- Haynes R. F., Klein U., Wielebinski R., Murray J. D., 1986, *A&A*, **159**, 22
- Heald G. H., 2012, *ApJ*, **754**, L35
- Heiles C., Troland T. H., 2005, *ApJ*, **624**, 773
- Hennebelle P., 2013, *A&A*, **556**, A153
- Hensley B. S., et al., 2022, *ApJ*, **929**, 166
- Hildebrand R. H., 1988, *QJRAS*, **29**, 327
- Hiltner W. A., 1949, *Science*, **109**, 165
- Hiltner W. A., 1951, *ApJ*, **114**, 241
- Hoang T., Lazarian A., 2014, *MNRAS*, **438**, 680
- Hoang T., Lazarian A., 2016, *ApJ*, **831**, 159
- Hotan A. W., et al., 2021, *Publ. Astron. Soc. Australia*, **38**, e009
- Hough P. V., 1962, *Methods and Means for Recognizing Complex Patterns*, <https://www.osti.gov/biblio/4746348>
- Inoue T., Inutsuka S.-i., 2016, *ApJ*, **833**, 10
- Jaffe T. R., Leahy J. P., Banday A. J., Leach S. M., Lowe S. R., Wilkinson A., 2010, *MNRAS*, **401**, 1013
- Jameson K. E., et al., 2016, *ApJ*, **825**, 12
- Jelić V., Prelogović D., Haverkorn M., Remeijn J., Klindžić D., 2018, *A&A*, **615**, L3
- Jones M. E., et al., 2018, *MNRAS*, **480**, 3224
- Jones T. J., et al., 2020, *AJ*, **160**, 167
- Kalberla P. M. W., Kerp J., Haud U., Winkel B., Ben Bekhti N., Flöer L., Lenz D., 2016, *ApJ*, **821**, 117
- Khademi M., Nasiri S., Tabatabaei F. S., 2022, arXiv e-prints, p. arXiv:2205.09069
- Kierdorf M., et al., 2020, *A&A*, **642**, A118
- Kim W.-T., Stone J. M., 2012, *ApJ*, **751**, 124
- Krumholz M. R., Federrath C., 2019, *Frontiers in Astronomy and Space Sciences*, **6**, 7
- Lasker B. M., Doggett J., McLean B., Sturch C., Djorgovski S., de Carvalho R. R., Reid I. N., 1996, in Jacoby G. H., Barnes J., eds, *Astronomical Society of the Pacific Conference Series Vol. 101, Astronomical Data Analysis Software and Systems V*. p. 88
- Le Coarer E., Rosado M., Georgelin Y., Viale A., Goldes G., 1993, *A&A*, **280**, 365
- Lee M.-Y., et al., 2012, *ApJ*, **748**, 75
- Livingston J. D., McClure-Griffiths N. M., Mao S. A., Ma Y. K., Gaensler B. M., Heald G., Seta A., 2022, *MNRAS*, **510**, 260
- Lobo Gomes A., Magalhães A. M., Pereyra A., Rodrigues C. V., 2015, *ApJ*, **806**, 94
- Loiseau N., Klein U., Greybe A., Wielebinski R., Haynes R. F., 1987, *A&A*, **178**, 62
- Lopez-Rodriguez E., et al., 2022, *ApJ*, **936**, 92
- Ma Y. K., Mao S. A., Ordog A., Brown J. C., 2020, *MNRAS*, **497**, 3097
- MacLow M.-M., 2004, *Ap&SS*, **289**, 323
- Magalhães A. M., et al., 2012, in Hoffman J. L., Bjorkman J., Whitney B., eds, *American Institute of Physics Conference Series Vol. 1429, Stellar Polarimetry: from Birth to Death*. pp 244–247, doi:10.1063/1.3701933
- Mao S. A., Gaensler B. M., Stanimirović S., Haverkorn M., McClure-Griffiths N. M., Staveley-Smith L., Dickey J. M., 2008, *ApJ*, **688**, 1029
- Martin P. G., Blagrove K. P. M., Lockman F. J., Pinheiro Gonçalves D., Boothroyd A. I., Joncas G., Miville-Deschênes M. A., Stephan G., 2015, *ApJ*, **809**, 153
- Massana P., et al., 2022, *MNRAS*, **513**, L40
- Mathew S. S., Federrath C., 2021, *MNRAS*, **507**, 2448
- Mathewson D. S., Ford V. L., Visvanathan N., 1986, *ApJ*, **301**, 664
- McClure-Griffiths N. M., Dickey J. M., Gaensler B. M., Green A. J., Haverkorn M., 2006, *ApJ*, **652**, 1339
- McClure-Griffiths N. M., et al., 2009, *ApJS*, **181**, 398
- McClure-Griffiths N. M., et al., 2018, *Nature Astronomy*, **2**, 901
- Murray C. E., Peek J. E. G., Di Teodoro E. M., McClure-Griffiths N. M., Dickey J. M., Dénes H., 2019, *ApJ*, **887**, 267
- Niederhofer F., et al., 2021, *MNRAS*, **502**, 2859
- Ogodo C. S., Green J. A., Dawson J. R., Breen S. L., Mao S. A., McClure-Griffiths N. M., Robishaw T., Harvey-Smith L., 2020, *MNRAS*, **493**, 199

- Panopoulou G. V., Dickinson C., Readhead A. C. S., Pearson T. J., Peel M. W., 2021, *ApJ*, **922**, 210
- Patrikeev I., Fletcher A., Stepanov R., Beck R., Berkhuijsen E. M., Frick P., Horellou C., 2006, *A&A*, **458**, 441
- Peek J. E. G., et al., 2011, *ApJS*, **194**, 20
- Peek J. E. G., et al., 2018, *ApJS*, **234**, 2
- Pingel N. M., et al., 2022, *Publ. Astron. Soc. Australia*, **39**, e005
- Planck Collaboration XIX 2015, *A&A*, **576**, A104
- Polychroni D., et al., 2013, *ApJ*, **777**, L33
- Price D. J., Bate M. R., 2008, *MNRAS*, **385**, 1820
- Scowcroft V., Freedman W. L., Madore B. F., Monson A., Persson S. E., Rich J., Seibert M., Rigby J. R., 2016, *ApJ*, **816**, 49
- Seta A., Shukurov A., Wood T. S., Bushby P. J., Snodin A. P., 2018, *MNRAS*, **473**, 4544
- Sharda P., Federrath C., Krumholz M. R., 2020, *MNRAS*, **497**, 336
- Shimwell T. W., et al., 2017, *A&A*, **598**, A104
- Skalidis R., et al., 2022, *A&A*, **665**, A77
- Skibba R. A., et al., 2012, *ApJ*, **761**, 42
- Smart B. M., Haffner L. M., Barger K. A., Hill A., Madsen G., 2019, *ApJ*, **887**, 16
- Soler J. D., et al., 2020, *A&A*, **642**, A163
- Soler J. D., et al., 2022, *A&A*, **662**, A96
- Stanimirovic S., Staveley-Smith L., Dickey J. M., Sault R. J., Snowden S. L., 1999, *MNRAS*, **302**, 417
- Staveley-Smith L., Sault R. J., Hatzidimitriou D., Kesteven M. J., McConnell D., 1997, *MNRAS*, **289**, 225
- Syed J., et al., 2022, *A&A*, **657**, A1
- Szotkowski S., et al., 2019, *ApJ*, **887**, 111
- Tahani M., et al., 2022, *A&A*, **660**, A97
- Tassis K., et al., 2018, arXiv e-prints, [p. arXiv:1810.05652](https://arxiv.org/abs/1810.05652)
- Tatton B. L., et al., 2021, *MNRAS*, **504**, 2983
- Van Eck C. L., et al., 2019, *A&A*, **623**, A71
- Villagran M. A., Gazol A., 2018, *MNRAS*, **476**, 4932
- Wang J., Hammer F., Yang Y., 2022, *MNRAS*, **515**, 940
- Wannier P. G., Lichten S. M., Morris M., 1983, *ApJ*, **268**, 727
- Welty D. E., Xue R., Wong T., 2012, *ApJ*, **745**, 173
- Wolleben M., et al., 2009, in Strassmeier K. G., Kosovichev A. G., Beckman J. E., eds, IAU Symp. Vol. 259, Cosmic Magnetic Fields: From Planets, to Stars and Galaxies. pp 89–90, doi:10.1017/S1743921309030117
- Yanchulova Merica-Jones P., et al., 2021, *ApJ*, **907**, 50

APPENDIX A: TESTS OF RHT PARAMETER CHOICE

As mentioned in Section 2.1, we repeat our analysis by adopting an array of RHT parameter choice to test its effects on our results of magnetically aligned H_i filaments in the SMC. The parameters chosen for our analysis in the main text were $R_{\text{sm}} = 12$ px, $D_W = 83$ px, and $\text{threshold} = 0.7$. We adjust each of these parameters slightly in turn: $R_{\text{sm}} = [8, 12, 16]$ px, $D_W = [71, 83, 95]$ px, and $\text{threshold} = [0.6, 0.7, 0.8]$. This amounts to 27 runs of our analysis, including the initial run described in the main text.

For each of the analysis runs, we repeat our two statistical tests (ray-tracing through the lower velocity portion of the GASKAP-H_i cube, for fields 1–11 only; see Section 4.1.1) and list the p -values in Table A1. In all cases, a lower p -value favours the alternative hypothesis (i.e., H_i filaments are preferentially aligned with the magnetic field traced by starlight polarisation), and we set a cutoff of 0.05 below which we deem the results as statistically significant. We further list the number of stars (out of the total of 4,494 in the concerned starlight fields) that are rejected from our ray-tracing analysis. These stars are not intercepted by any identified H_i filaments in our ray-tracing analysis, and therefore have Stokes $q = u = 0$.

Out of the total of 27 analysis runs, we find that a significant fraction of them reach the same conclusion of magnetic alignment of H_i




Figure B1. *Gaia* DR3 (*Gaia Collaboration et al. 2022*) *G*-band magnitude against *Lobo Gomes et al. (2015)* *V*-band magnitude for the 106 stars found by a positional cross-match within an angular separation of $5''$ between the two lists of stars. The 57 stars that further show good photometric matches (in the magenta box) are used for our per-star ray-tracing analysis using the *Gaia* radial velocity measurements.

filaments with the SMC (with a p -value of less than 0.05). Specifically, 18 / eight of the parameter combinations (66.7% / 33.3%) give p -values of less than 0.05 in at least one / both of the two statistical tests, respectively. In particular, we find that setting the threshold parameter to 0.7 or 0.8, which imposes a tighter restraint to the RHT filament finder by requiring the identified filaments to be more spatially coherent than when set to 0.6, do report more consistently p -values of less than 0.05. This means that when the RHT is configured to identify more coherent and therefore most probably more genuine ISM structures, the effect of magnetic alignment of H_i filaments appear to be more pronounced.

APPENDIX B: RAY TRACING USING *GAIA* DR3 RADIAL VELOCITIES

For our ray-tracing analysis (Section 3), we place the 5,999 *Lobo Gomes et al. (2015)* stars within the GASKAP-H_i cube (*Pingel et al. 2022*) at the H_i-intensity-weighted mean velocities (v_{mean}). This is because most of these polarised SMC stars do not have radial velocity measurements in the literature.

In the recently released *Gaia* DR3 (*Gaia Collaboration et al. 2022*), we find that 57 of the *Lobo Gomes et al. (2015)* stars have reported radial velocities. This prompts us to attempt a dedicated ray-tracing experiment for these 57 stars using their radial velocity values as the locations to place them at within the H_i cube, described as follows.


We first perform a cross-match between the *Lobo Gomes et al. (2015)* and the *Gaia* DR3 catalogues by finding closest positional matches within $5''$ amongst the two lists of stars, followed by plotting the *Gaia G*-band magnitude against the *Lobo Gomes et al. (2015)* *V*-band magnitude (Figure B1). This allows us to identify the 57 stars that further show good photometric matches between the two catalogues that we use for our per-star ray-tracing analysis.

Next, we perform the ray-tracing analysis as described in Section 3, but for the 57 stars above only. Furthermore, the stars are placed along

Table A1. Results from re-running our analysis (fields 1–11; $v < v_{\text{mean}}$) using various choices of RHT parameters

	threshold = 0.6			threshold = 0.7			threshold = 0.8		
	$R_{\text{sm}} = 8 \text{ px}$	$R_{\text{sm}} = 12 \text{ px}$	$R_{\text{sm}} = 16 \text{ px}$	$R_{\text{sm}} = 8 \text{ px}$	$R_{\text{sm}} = 12 \text{ px}$	$R_{\text{sm}} = 16 \text{ px}$	$R_{\text{sm}} = 8 \text{ px}$	$R_{\text{sm}} = 12 \text{ px}$	$R_{\text{sm}} = 16 \text{ px}$
<i>p</i> -values (Test 1 / Test 2)									
$D_W = 71 \text{ px}$	0.124 / 0.628	0.338 / 0.118	0.525 / 0.628	0.060 / 0.118	0.010 / 0.005	0.034 / 0.118	0.015 / 0.005	0.085 / 0.030	0.129 / 0.118
$D_W = 83 \text{ px}$	0.209 / 0.322	0.171 / 0.118	0.085 / 0.030	0.146 / 0.118	0.008 / 0.005	0.034 / 0.030	0.077 / 0.030	0.194 / 0.118	0.067 / 0.030
$D_W = 95 \text{ px}$	0.005 / 0.030	0.076 / 0.030	0.005 / 0.005	0.108 / 0.030	0.007 / 0.005	0.065 / 0.030	0.025 / 0.118	0.033 / 0.030	0.040 / 0.030
Rejected Star Count									
$D_W = 71 \text{ px}$	0	0	0	0	0	0	1330	684	415
$D_W = 83 \text{ px}$	0	0	0	0	0	1	2166	1148	738
$D_W = 95 \text{ px}$	0	0	0	12	4	4	2809	1744	1128

NOTE – Test 1: One-sample KS test; Test 2: Matching within 20° test (see Section 4.1.1).
 Parameter choice of our main analysis also listed here ($R_{\text{sm}} = 12 \text{ px}$, $D_W = 83 \text{ px}$, and threshold = 0.7).
p-values of lower than 0.05 is deemed as statistically significant here, and are marked in boldface.

**Figure B2.** Angle difference between the ray-traced and the observed starlight polarisation for the 57 stars cross-matched between the *Gaia* DR3 (Gaia Collaboration et al. 2022) and the Lobo Gomes et al. (2015) catalogues.

the velocity axis at the reported *Gaia* DR3 radial velocities⁹ instead of at v_{mean} , the starlight is sent through the lower velocity portion of the H α cube only as per our findings in Section 4.1, and each of the stars is studied individually instead of on a per-field basis. The resulting angle difference of these 57 stars between the ray-traced and the observed starlight polarisation is shown in Figure B2.

Evidently, we do not find a good match between the observed starlight polarisation angle and the H α filament orientation here. This is likely because, as pointed out in Section 5.2.2, that the ray-tracing analysis through the GASKAP-H α cube may not capture the contributions of the turbulent magnetic field, which dominates in the SMC (e.g., Mao et al. 2008; Lobo Gomes et al. 2015; Livingston et al. 2022). Therefore, the per-star analysis here does not result in a good match in orientations between the magnetic field and H α filaments. In the future, when radial velocity measurements become available for most, if not all, of the Lobo Gomes et al. (2015) polarised SMC stars, one can return to our ray-tracing analysis as presented in the main text to re-assess the use of the stellar radial velocities.

⁹ We have converted the radial velocities from the barycentric frame of *Gaia* to our local standard of rest frame.

APPENDIX C: 2D HISTOGRAMS OF OBSERVED AND RAY-TRACED STARLIGHT POLARISATION

We include the 2D histograms of the polarised stars in the SMC on the Stokes *qu* plane, separated by the Lobo Gomes et al. (2015) fields of observations (Figure 1). Figure C1 shows the actual observed polarised SMC stars reported by Lobo Gomes et al. (2015), while the results from ray-tracing through the GASKAP-H α cube are shown in Figures C2, C3, C4, and C5 for the case of the lower velocity portion, higher velocity portion, full velocity range in ascending velocity, and full velocity range in descending velocity, respectively. As described in Section 2.2, the parameters $\bar{p}$ and $\bar{\theta}$ represents the contribution of the large-scale ($\approx 150 \text{ pc}$) ordered magnetic field, while σ_p reflects the small-scale ($\ll 150 \text{ pc}$) turbulent magnetic field.

This paper has been typeset from a T_EX/L^AT_EX file prepared by the author.




Figure C1. 2D histograms of the linear polarisation properties of the Lobo Gomes et al. (2015) observed stars in the SMC. Each panel corresponds to a starlight field. The resulting $\bar{p}$ and $\bar{\theta}$ from taking a vector mean out of all stars for each field are represented by the blue crosses, while the 2D standard deviations are represented by the radii of the blue circles.




Figure C2. Similar to Figure C1, but from ray tracing through the GASKAP-H_I cube for the lower velocity portion only.




Figure C3. Similar to Figure C1, but from ray tracing through the GASKAP-H1 cube for the higher velocity portion only.




Figure C4. Similar to Figure C1, but from ray tracing through the GASKAP-H_I cube for the full velocity range. The starlight traverses through the H_I cube in ascending velocity, starting at $v = 40.91 \text{ km s}^{-1}$.




Figure C5. Similar to Figure C1, but from ray tracing through the GASKAP-Hi cube for the full velocity range. The starlight traverses through the Hi cube in descending velocity, starting at $v = 256.85 \text{ km s}^{-1}$.