Benchmark Studies on the Isomerization Enthalpies for Interstellar Molecular Species

Emmanuel E. Etim

Department of Chemical Sciences, Federal University Wukari, Katsina-Ala Road, P.M.B. 1020 Wukari, Taraba State, Nigeria Department of Chemistry, Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India

*email: emmaetim@gmail.com

Abstract: With the well-established correlation between the relative stabilities of isomers and their interstellar abundances coupled with the prevalence of isomeric species among the interstellar molecular species, isomerization remains a plausible formation route for isomers in the interstellar medium. The present work reports an extensive investigation of the isomerization energies of 246 molecular species from 65 isomeric groups using the Gaussian-4 theory composite method with atoms ranging from 3 to 12. From the results, the high abundances of the most stable isomers coupled with the energy sources in interstellar medium drive the isomerization process even for barriers as high as 67.4 kcal/mol. Specifically, the

cyanides and their corresponding isocyanides pairs appear to be effectively synthesized via this process. The following



potential interstellar molecules; CNC, NCCN, c-C5H, methylene ketene, methyl Ketene, CH3SCH3, C5O, 1,1-ethanediol, propanoic acid, propan-2-ol, and propanol are identified and discussed. The study further reaffirms the importance of thermodynamics in interstellar formation processes on a larger scale and accounts for the known isomeric species. In all the isomeric groups, isomerization appears to be an effective route for the formation of the less stable isomers (which are probably less abundant) from the most stable ones that are perhaps more abundant.

Keywords: Isomers, Energy barrier, Interstellar chemistry, Astrochemistry, Hydrogen Bonding,

Introduction: That the interstellar medium (ISM) is chemically rich is not an argument with the discoveries of over 200 different molecular species in this thin space between the stars which was earlier regarded as a vacuum dotted with stars, black holes, and other celestial bodies [1-6]. These molecules are important in various fields such as atmospheric chemistry, astrochemistry, prebiotic chemistry, astrophysics, astronomy, astrobiology, etc, and in our understanding of the solar system, "the world around us" with each successfully detected interstellar molecule telling the story of the chemistry and physics of the environment from where it was detected. They serve as the most important tools for probing deep into the interior of the molecular clouds and the molecular clouds are significant because it is from them that stars and consequently new planets are formed. The symmetric rotors serve as interstellar thermometers while the metal-bearing species provide useful information regarding how the simple molecules that were present on the early earth may have given rise to the complex systems and processes of contemporary biology is one question the

biologically related interstellar molecules can be used to address. Molecules also provide the cooling mechanism for the clouds through their emission [3,7-8].

A careful look at the list of known interstellar and circumstellar molecules reveals some interesting chemistries among these molecular species. Tables 1 and 2 contain the list of the currently (as of June 2021) known interstellar and circumstellar species arranged according to the number of atoms making up the molecules.

2 atoms	3 atoms	4 atoms	5 atoms	6 atoms	7 atoms
H ₂ , CO, CSi,	$H_2O, H_2S,$	NH_3, H_2CO	CH ₄ , SiH ₄	CH ₃ OH, CH ₃ SH	CH ₂ CHOH
СР	HCN, $TiO_{2,}$	H ₂ CS, C ₂ H ₂	CH ₂ NH.	C ₂ H ₄ , HC ₄ H	c-C ₂ H ₄ O
CS, NO, NS,	\mathbf{SO}_2	2) - 2 2	NH ₂ CN	- 2 47 - 4	
SO		HNCS	CH CO	CH ₃ CN, CH ₃ NC	HC(O)CH ₃
HCI NaCl	MgCN,	H ₃ O ⁺ , SiC ₃	$H_2CO,$	HCONH ₂ ,	Н ₃ СССН
KCl	MgNC, NaCN		neoon		
	FeCN,	C_3S, H_2CN	HCCCN,	$HC_2C(0)H$	CH ₃ NH ₂
AICI, AIF, PN	KCN,	c-C ₃ H, l-C ₃ H	HUUNU	HC_3NH^+	CH ₂ CHCN
SiN, SiO, SiS	AIOH.	HCCN, CH ₂	c-C ₃ H ₂ , l-	HCAN	HC ₄ CN
NH. OH. C2	$H_2Cl, H_2O^+,$	110 01 (j 011)j	C_3H_2		1104011
1,11,011,02	$H_2Cl^*,\ N_2O,$	C_2CN, C_3O	CH ₂ CN,	C_5N, C_5H	C ₆ H
CN, HF, FeO	$\rm NH_2, OCS$	HCNH⁺,	H_2COH^+	H ₂ C ₄ , H ₂ CCNH	CH ₃ NCO
LiH, CH, CH^+	CH ₂ , HCO,	\mathbf{HOCO}^+	C ₄ Si	C N ²	
CO^+ SO^+ SU	C ₃	C₂N, HNCO		C ₅ N	пс ₅ 0,
NO^+	С.Н. С.О.	0311, 221100	C ₅	c-H ₂ C ₃ O	HOCH ₂ CN
110	C_2S , AINC,	HSCN, C_3N ,	HNCCC	HNCHCN C-S	нисносн
O_2 , N_2 , CF^+	HNO	PH ₃	G 11	SiH ₃ CN,	micenceni,
PO, HD	SiCN. N ₂ H ⁺	HMgNC	C ₄ H		HC ₄ NC,
	SiNC, c-	нссо	C4H	z-HNCHCN,	c-C₃HCCH,
SiH, AlO,	SiC ₂	NCCP,	HC(O)CN	MgC ₄ H	
\mathbf{ArH}^{+} , \mathbf{OH}^{+} ,	C ₂ N HCO+	MgCCH,	СН ₃ О,	CH-CO+	
CN^{-} , SH^{+} ,	HOC^+	HOCN,	HNCNH,	cii;co ;	
HCI, 110 , CrO , NS^+ .	HCS^+ , H_3^+	CNCN	H_2NCO^+ ,	CH ₂ CCH,	
010, 115 ,	OCN, HCP,	що	NH ₃ D, CH ₂ CL	H ₂ CCCS,	
VO, HeH ⁺	S ₂ H, HCS.	$\mathbf{H}_{2}\mathbf{O}_{2},$	<i></i> ,		
	HSC, NCO,	t-HONO	MgC ₃ N,		
	CaNC		MH_2OH ,		
	Carve,		HC ₃ O ⁺ ,		
	NCS, HO ₂		HC_3S^+ ,		
			- <u>-</u> -		
			C_4S ,		

Table 1: Interstellar molecules between 2 and 7 atoms

t-HC(O)SH,
H ₂ CCS,
NCCNH ⁺

Table 2: In			veen 8 and	>12 atoms
-------------	--	--	------------	-----------

8 atoms	9 atoms	10 atoms	11 atoms	12 atoms	>12 atoms
CH ₃ COOH	(CH ₃) ₂ O	(CH ₃)CO	HC ₉ N	C ₆ H ₆	HC ₁₁ N
HCOOCH ₃	CH ₃ CH ₂ CN	HOC ₂ H ₄ OH	CH ₃ C ₆ H	C ₃ H ₇ CN	C ₆₀
HOCH ₂ CHO	CH ₃ CH ₂ OH	H ₃ CCH ₂ COH	HCOOC ₂ H ₅	C ₂ H ₅ OCH ₃	C ₆₀ ⁺
H ₃ C ₃ CN,	CH ₃ CH ₂ SH	CH ₃ C ₄ CN,	CH ₃ OCOCH ₃	branched-	C ₇₀
C ₇ H, (NH ₂) ₂ CO	CH ₃ C ₄ H	СН ₃ СНСНО,	CH ₃ COCH ₂ OH	$C_3H_7CN_1$	c-C ₆ H ₅ CN
H_2C_6	HC ₇ N	CH ₃ OCH ₂ OH	c-C ₅ H ₆	c-C₅H₅CN	C ₁₀ H ₇ CN
H(CC) ₃ H	C ₈ H		NH ₂ CH ₂ CH ₂ OH		c-C ₉ H ₈
H ₂ CCHCHO	CH ₃ CONH ₂				
CH ₂ CCHCN	C ₈ H ⁻				
H ₂ NCH ₂ CN,	CH ₃ CHCH ₂ ,				
CH ₃ CHNH	CH ₃ NHCHO,				
CH ₃ SiH ₃ ,	HC ₇ O				
(NH ₂) ₂ CO	HCCCHCHCN				
HCCCH ₂ CN,	H ₂ CCHC ₃ N				
$\mathrm{HC}_{5}\mathrm{NH}^{+},$	H ₂ CCCHCCH				
CH ₂ CHCCH,					

Some of the chemistries existing among the interstellar molecular species include the dominance of the linear carbon chain species of the form C_n , H_2C_n , HC_nN , $CH_3(CC)_nH$, $CH_3(C\equiv C)_nCN$ and C_nX (X=N, O, Si, S, H, P, N⁻, H⁻) which account for more than 20% of all the known interstellar and circumstellar molecular species; the presence of about 30 alkyl group containing interstellar molecules mostly observed from the same or similar molecular clouds; periodic trends in which elements from the same group appear to form similar

compounds with similar properties as in the case of oxygen and sulphur (group 6/16 elements) in which of the over 25 S-containing molecules observed in ISM, about 20 have the corresponding O-containing molecules uniquely detected in ISM and the abundance of S-compound relative to its O-analogue is approximately equal to the cosmic S/O ratio, 1/42 as seen in methyl mercaptan, thioisocyanic acid, etc, except where the effect of interstellar hydrogen bonding dominates [9-10]; successive hydrogen addition where larger species are believed to be formed from the smaller unsaturated species via successive hydrogen addition in which both species could be shown to be chemically and spatially related [11]. Isomerism is yet another prevailing chemistry among the interstellar molecular species. Apart from the diatomics and a few species which cannot form isomers, about 40% of all the known interstellar molecules have isomeric analogues ranging from the isomeric pairs to the isomeric triads which are believed to have a common precursor for their formation process [12]. Table 3 lists some of the known isomeric species and their isomeric groups.

Isomeric Group	Astronomically observed isomers
CHN	HCN, HNC
CNMg	MgCN, MgNC
CNSi	SiCN, SiNC
CHO^+	HOC^+ , HCO^+
CHSN	HSCN, HNCS
HC ₃	$c-C_3H$, $l-C_3H$
H_2C_3	$c-C_3H_2$, $1-C_3H_2$
CN_2H_2	NH ₂ CN, HNCNH
C_3H_2O	HCCCHO, $c-H_2C_3O$
C_4H_2	$H(CC)_2H$, H_2C_4
C_6H_2	H_2C_{6} $H(CC)_3H$
C_4H_3N	CH ₂ CCHCN, CH ₃ CCCN
C_2H_6O	$(CH_3)_2O, CH_3CH_2OH$
C_3H_6O	$(CH_3)_2CO, CH_3CH_2C(O)H$
$C_3H_6O_2$	$HC(O)OCH_2CH_3$, $CH_3OC(O)CH_3$
C_4H_7N	$n-CH_3CH_2CH_2CN$, $b-CH_3CH_2CH_2CN$
CHON	HNCO, HCNO, HOCN
$C_2H_4O_2$	CH ₃ COOH, HC(O)OCH ₃ , HOCH ₂ C(O)H
C_2H_4O	$CH_2CH(OH)$ c- C_2H_4O , $HC(O)CH_3$
C_2H_3N	CH ₃ CN, CH ₃ NC, H ₂ CCNH
C ₃ HN	HC ₂ CN, HC ₂ NC HNCCC

Table 3: Known interstellar isomeric pairs, triads and their isomeric groups.

Because of the conditions (low temperature and pressure) in the interstellar medium, there is hardly a consensus as to how these molecules are formed but some of the chemistries listed above and the effect of thermodynamics serve as clues as to how these molecules could be formed in ISM. Focusing here on isomerization, the prevalence of isomeric species among the known astromolecules coupled with the energy sources in ISM such as the shock waves (which could arise from the interaction of the Earth's magnetic field with the solar wind, molecular outflows during star formation, supernova blasts and galaxies colliding with each other) which provide energy for both the formation and distribution of large interstellar species, place isomerization as one of the most plausible routes for the formation of interstellar molecules [13]. As observed in some studies, the most stable isomers are found to be more abundant than their less stable analogues except where other factors dominate; thus, the isomerization of the most stable isomer (which is probably the most abundant) to the less stable isomers can be a very effective and efficient formation mechanism in ISM. Also, apart from the energy sources in ISM, the high abundance of the most stable isomer can drive the isomerization process irrespective of the energy barrier.

According to the minimum energy principle [14], isomerization is the most important process in determining the relative abundances of the isomers in ISM. Isomers of the same generic formula are said to have a common intermediate in their formation and destruction routes. After reaching the generic formula, the equilibration process is said to occur. This implies an internal isomerization with a low activation barrier, assisted isomerization, or catalytic isomerization at the grain/ice surface [14].

The present work aims at estimating accurate isomerization enthalpies for 246 different molecular species from 65 isomeric groups using the Gaussian 4 theory composite method [15]. The molecules range from the 3 atomic species to those with 12 atoms with at least one known interstellar molecule from each isomeric group. The results account for the extent and effectiveness of isomerization as a plausible formation route in ISM; and the rationale behind the successful observation of the known species. Among other things, potential candidates for astronomical searches are highlighted and discussed.

Computational Details: A concerted effort between theory and experiment is found in the heart of some of the most successful scientific studies. Considering the large range of molecules examined in this study, only very few of them have experimentally measured enthalpies of formation while others are so unstable that they cannot be probed experimentally in the terrestrial laboratory but all of these can be comfortably handled computationally. The Gaussian 4 (G4) theory composite method is employed in estimating accurate standard enthalpies of formation ($\Delta_{\rm f} {\rm H}^{\rm O}$) for all the molecular systems investigated in this work. The G4 composite method is very accurate for several systems and benchmark studies; in its release note, it has an average absolute deviation of 0.8 kcal/mol from experimental values for the enthalpies of formation of 270 molecular species. The G4 method is a modification of the G3 method which has an average absolute deviation of 1.19 kcal/mol for the same number of molecular systems [15-16]. In the G4 method, geometry optimization and zero point energy (ZPE) are carried out at the B3LYP level of theory using the 6-31G(2df,p) basis set. The ZPE is scaled by 0.9854. Single-point calculations and energy are done using the MP4/6-31G(d) method modified by corrections from additional calculations (with MP4 and other methods) while core correlation is obtained via higher-level correction terms. The Gaussian 09 suite of programs is used for all the computational studies reported here. Only stable geometries with no imaginary frequencies are considered. The standard

enthalpies of formation are obtained from the atomization energies using the approach described in previous studies [4-6, 17-23]

Results and Discussion

The standard enthalpies of formation ($\Delta_f H^O$) for all the 246 molecular species from 65 isomeric groups examined in this study are contained in the supporting information. The relative enthalpies for each isomeric group are presented and discussed in this section. The different isomeric groups investigated are grouped according to the number of atoms beginning from 3 to 12.

Isomers with 3 atoms

In Table 4, the relative enthalpies or isomerization energies of the 26 molecular species from 13 isomeric groups with 3 atoms are shown alongside the current astronomical status of these species. At least one isomer from each of these isomeric groups is a known interstellar molecule [24-43]. The isomerization enthalpies range from 0.2 to 41.5 kcal/mol for groups where both isomers have been detected. This range illustrates the effectiveness of the isomerization mechanism as a plausible route for the formation of these molecular species in ISM; it also shows how far or the extent to which the energy sources within the ISM can drive some chemical processes. For the isomeric groups where only one isomer has been detected, the observed range of the relative enthalpy of formation suggests that species like NaNC, AICN, ONC⁻ and even HOC can be formed from their most stable isomers that have been detected already. The high abundances of the stable isomers coupled with the energy sources in ISM imply the possibility of these less stable isomers being formed from their most stable isomers via the isomerization mechanism.

With the exception of C_2N group where only the less stable isomer is been detected, in all other cases where only one isomer is detected, it is the most stable isomer which supports the fact that the most stable isomer is probably the most abundant and the most abundant species is easily detected compared to the less stable isomer. Where both isomers have been detected, the most stable isomer is found to be the most abundant except where other processes dominate. CNC is more stable than CCN but the less stable isomer has been detected while the most stable isomer is yet to be astronomically observed. From literature perusal, there is no information regarding the spectroscopic parameters of CNC that would have warranted it astronomical searches. Thus, the detection of CNC awaits the availability of accurate spectroscopic parameters.

Ions (both cations and anions) play important role in the formation processes of interstellar molecules. Under the conditions of the ISM, neutral atoms and molecules tend to be unreactive toward molecular hydrogen but in the presence of ions, most of the reactions become very efficient since the ions can easily react without having to overcome the reaction barrier [7,44]. This can be seen in the case of the CHO⁺ group where HOC⁺ with a relative enthalpy of formation of 37.3 kcal/mol has been detected. Theoretical calculations and laboratory experiments have shown that the isomerization process in the CHO⁺ isomers has essentially no barrier [14]. Thus, with the availability of accurate spectroscopic parameters,

the HSC^+ and ONC^- ions (where their most stable isomers are already detected) could be successfully detected.

Isomerization appears to be a favourable route for the cyanide/isocyanide pair. Table 4 contains 7 cyanide/isocyanide pairs of which both cyanide and isocyanide have been detected in three groups with isomerization energy ranging from 0.2 to 13.5 kcal/mol. Except for the ONC⁻ ion with a relative enthalpy of formation of 17.1kcal/mol; the remaining members of the cyanide/isocyanide pairs have relative enthalpy of formation in the range of 0.0 to 7.3 kcal/mol which is well within the range of those already detected. Thus, NaNC, AICN and CNC are most likely to be formed from their corresponding analogues and could be successfully detected.

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$	Astronomical	Ref.
Isomeric group	Isomers	(kcal/mol)	status	
CNH	HCN	0.0	Observed	[24]
	HNC	13.5	Observed	[25]
CNNa	NaCN	0.0	Observed	[26]
	NaNC	2.5	Not observed	
		0.0		
CNMg	MgNC	0.0	Observed	[27]
	MgCN	0.7	Observed	[28-30]
CNAl	AINC	0.0	Observed	[31]
	AlCN	7.3	Not observed	
CNSi	SiNC	0.0	Observed	[32]
	SiCN	0.2	Observed	[33]
CUO	UCO	0.0	Observed	[2.4]
СНО	HCO	0.0	Observed	[34]
	HOC	38.2	Not observed	
СНР	НСР	0.0	Observed	[35]
	HPC	75.7	Not observed	[]
C_2N	CNC	0.0	Not observed	
	CCN	2.1	Observed	[36]
C ₂ P	ССР	0.0	Observed	[37]
- 2-	CPC	85.3	Not observed	['']
CHO^+	HCO^+	0.0	Observed	[38]

Table 4:	Isomerization	enthalpies	for isomers	with 3 atoms
----------	---------------	------------	-------------	--------------

	HOC ⁺	37.3	Observed	[39-40]
CHS^+	HCS^+	0.0	Observed	[41]
	HSC^+	94.1	Not observed	
CNO ⁻	OCN ⁻	0.0	Observed	[42]
	ONC ⁻	17.1	Not observed	
CHS	HCS	0.0	Observed	[43]
	HSC	41.5	Observed	[43]

Isomers with 4 atoms

For the isomers with 4 atoms, 25 molecular species from 10 isomeric groups are examined. Table 5 contains the isomerization energies of these molecular species and their current astronomical status. Fourteen of these molecular species have been detected from different astronomical sources [45-63]. In the CHON, CHSN, and C_3H groups where more than one isomer has been detected, the relative enthalpy of formation ranges from 3.1 to 67.4 kcal/mol. For the remaining 7 isomeric groups with only one known molecular species from each, the relative enthalpy of formation ranges from 21.6 to 49.8 kcal/mol. This range falls within that of the known species, thus, pointing to the possibility of detecting the less stable isomers of these groups that could be formed via the isomerization process.

Among the isomers with 4 atoms examined here, there are eight cyanide/isocyanide pairs. In two of these pairs, both cyanide and isocyanide have been detected with a relative enthalpy of formation in the range of 2.8 to 29.0 kcal/mol. For the other cyanide/isocyanide pairs where only the cyanides are observed, the relative enthalpy of formation ranges from 18.7 to 24.7 kcal/mol which is below the 29.0 kcal/mol relative enthalpy of formation calculated for the known cyanide/isocyanide pairs. Thus, isomerization remains a plausible mechanism for the formation of the less stable isocyanides from their corresponding cyanides which are probably more abundant.

The C_2N_2 isomeric group is the isoelectronic analogue of the C_2NP isomeric group. Just as there is interesting chemistry between the O and S-containing interstellar molecular species; such also exists for the N and P-containing species, though it is not well explored as that of O and S. Over 80% of all the known interstellar and circumstellar molecules have been detected via their rotational spectral features because, at the low temperature of the ISM, rotational excited states are easily populated. NC₂N (the most stable isomer of the C₂N₂ group) is microwave inactive meaning that it cannot be astronomically detected via radio astronomy. There are reasons to believe the presence and detectability of NC₂N; its protonated analogue, NC₂NH⁺ has been detected [64]. NCCP, the isoelectronic analogue of NC₂N is also known [62]. Both the protonated analogues of NC₂N and its isoelectronic analogue that have been observed are likely to be less abundant than NC₂N because of the reactive nature of the ion and the less cosmic abundance of P as compared to N. Thus, infrared astronomy of NCCN or radio astronomy of its isotopologues (which are microwave active) is likely to be successful.

The observed isomers in all the groups shown in Table 5 are also the most stable isomers.

Isomeric	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$	Astronomical status	Reference
group	Terrenia esid	(KCal/mol)	Observed	[45]
CHON	Isocyanic acid	0.00	Observed	[45]
	Cvanic acid	29.0	Observed	[46-47]
	5			
	Fulminic acid	67.4 (0.0)	Observed	[48]
	Isofulminic	86.1 (18.7)	Not observed	
	acid			
CHSN	HNCS	0.0	Observed	[49]
	HSCN	11.2	Observed	[50]
	HCNS	40.6 (0.0)	Not observed	
	HSNC	43.3 (2.8)	Not observed	
C_3H	c-C ₃ H	0.0	Observed	[51]
	l-C ₃ H	3.1	Observed	[52]
C ₃ N	I-C ₃ N	0.0	Observed	[53-54]
	C ₂ NC	22.7	Not observed	
	c-C ₃ N	28.8	Not observed	
C ₂ NH	HC ₂ N	0.0	Observed	[55]
	HCNC	28.0	Not observed	
~ ~ ~				
C ₃ O	I-C ₃ O	0.0	Observed	[56-57]
	c-C ₃ O	21.6	Not observed	
~ ~ ~	1 7 7			[#0, c0]
C_3S	1-C ₃ S	0.0	Observed	[58-60]
	c-C ₃ S	26.3	Not observed	
SiC	a C Si	0.0	Observed	[61]
SIC ₃	1 C Si	0.0	Not observed	[01]
	1-0351	47.0	not observed	
C ₂ NP	NCCP	0.0	Observed	[62]
	CNCP	24.1	Not observed	[]
C ₂ N ₂	NC ₂ N	0.0	Not observed	
- 2 2	CNCN	24.7	Observed	[63]
L				

Table 5: Isomerization enthalpies for isomers with 4 atoms

Isomers with 5 atoms

Isomerization energies for 20 molecular species from 6 isomeric groups are presented in Table 6 alongside their current astronomical status. Nine of these species are known astromolecules [65-73]. In the C₃HN and C₃H₂ groups where more than one isomer has been detected, the isomerization energy ranges from 13.7 to 45.4 kcal/mol, and the isomerization energies for some of the isomers with five atoms whose most stable isomers have been detected fall within this range. These include; ethynol, CH₂NN, NH₂NC, CH₂NC, HCONC and c-C₂NHO. These species can as well be formed from their most stable isomers via the isomerization mechanism.

Again, isomerization appears to be a favourable route for the formation of isocyanides from their corresponding cyanides. As shown in Table 6, an isomerization enthalpy as high as 45.4 kcal/mol is noted for an isocyanide formation. This strongly supports the formation of other isocyanides from their corresponding cyanides. The barrier for other isocyanides whose corresponding cyanides have been detected ranges from 12.9 to 43.4 kcal/mol which is within the limit of the one that has been observed.

The fact that the most stable isomers are probably the most abundant and are easily detected in ISM is well demonstrated among these isomers. From table 6, all the observed isomers are the most stable ones in their respective isomeric groups as compared to the ones that have not been detected.

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$ (kcal/mol)	Astronomical status	Reference
C ₃ HN	HCCCN	0.0	Observed	[65]
	HCCNC	24.1	Observed	[66]
	HNCCC	45.4	Observed	[67]
	CC(H)CN	50.5	Not observed	
	HCNCC	72.6	Not observed	
C_2H_2O	Ketene	0.0	Observed	[68]
	Ethynol	38.8	Not observed	
	Oxirene	81.9	Not observed	
C ₃ H ₂	c-C ₃ H ₂	0.0	Observed	[69]

Table 6: Isomerization enthalpies for isomers with 5 atoms

	l-C ₃ H ₂	13.7	Observed	[70]
N_2H_2C	NH ₂ CN	0.0	Observed	[71]
	CH ₂ NN	26.3	Not observed	
	NH ₂ NC	43.4	Not observed	
C_2H_2N	CH ₂ CN	0.0	Observed	[72]
	CH ₂ NC	22.2	Not observed	
C ₂ HNO	CNCHO	0.0	Observed	[73]
	HCONC	12.9	Not observed	
	c-C ₂ NHO	30.2	Not observed	
	HNCCO	69.2	Not observed	
	HC ₂ NO	83.3	Not observed	

Isomers with 6 atoms

The isomerization energies and the current astronomical status of the different isomeric species with 6 atoms investigated in this study are presented in Table 7. Figures 1 to 5 display some of the cyclic isomers highlighted in Table 7. One-third of all the molecular species presented in Table 7 have all been detected from several astronomical sources [74-85]. In the C_2H_3N and H_2C_3O groups where more than one isomer has been detected, the isomerization enthalpies range from 7.3 to 23.1 kcal/mol. Except for some of the cyclic isomers which are highly unstable, most of the unknown isomers (c- C_5N^* , C_4NC , c- C_5N^{**} , c- C_5H^a , SiH₃NC, HC₃NC, methylene ketene, hydroxymethylimine) have isomerization enthalpies in the range of the known isomers, suggesting their possible astronomical observation if they could be formed via isomerization as the ones that have been observed. Ketenimine for instance is reported to be formed from methyl cyanide via tautomerization (i.e., an isomerization pathway in which the migration of hydrogen atom from the methyl group to the nitrogen atom is accompanied by a rearrangement of bonding electrons) and this process is said to be driven by shock waves that provide the energy for both the formation and distribution of large interstellar species [80].

The C₅H isomers; $1-C_5H$ and $c-C_5H^a$ are almost identical in energy. If the spectroscopic parameters of $c-C_5H^a$ (Figure 2a) are accurately probed, either experimentally or theoretically; this species ($c-C_5H$) could be detected in ISM. The four cyanide/isocyanide pairs among the isomers with 6 atoms presented in Table 7 have isomerization enthalpies in the range of 4.5 to 23.0 kcal/mol. Interestingly, the C₂H₃N group where methyl cyanide and methyl isocyanide have been detected has the highest relative enthalpy of formation of 23.0 kcal/mol which implies the possible detection of the other isocyanides with lower barriers (4.5 to 20.7kcal/mol) assuming isomerization as their plausible formation route.

Methylene ketene and methyl ketene have been proposed as potential interstellar molecules for many reasons. The ketenes are found to be more stable than their corresponding isomers (Tables 6, 7, and 9); they are less affected by interstellar hydrogen bonding assuming surface formation processes; ketene and ketenyl radical (H_2C_2O and HC_2O respectively) are known interstellar molecules [64; 68].

Just as every known O-containing interstellar molecule points to the presence and the detectability of the S-analogue. For every known S-species, the O-analogue is not only present in detectable abundance, but it can also be said to have even been overdue for astronomical detection. C_5O is the O-analogue of C_5S that has been detected. Thus, it is an important potential interstellar molecule.

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$	Astronomical status	Reference
		(kcal/mol)		
C ₅ N	$1-C_5N$	0.0	Observed	[74]
	c-C ₅ N*	17.4	Not observed	
	C ₄ NC	20.7	Not observed	
	c-C ₅ N**	21.0	Not observed	
	c-C ₅ N***	100.1	Not observed	
C ₅ H	l-C₅H	0.0	Observed	[75-77]
	$c-C_5H^a$	1.0	Not observed	
	c-C ₅ H ^b	34.7	Not observed	
	c-C ₅ H ^c	60.8	Not observed	
		0.0		
C_2H_3N	Methyl cyanide	0.0	Observed	[78]
	Methyl isocyanide	23.0	Observed	[79]
	Ketenimine	23.1	Observed	[80]
	Ethynamine	42.3	Not observed	
	2H-azirine	46.5	Not observed	
	1H-azirine	80.8	Not observed	
SiCH ₃ N	SiH ₃ CN	0.0	Observed	[62]
	SiH ₃ NC	4.5	Not observed	
	c-SiCH ₃ N ^d	33.4	Not observed	
	H ₂ SiCNH	46	Not observed	
	c-SiCH ₃ N ^e	50.2	Not observed	
	H ₂ NCSiH	57.0	Not observed	
C ₄ HN	HC ₄ N	0.0	Observed	[81]
	HC ₃ NC	17.4	Not observed	
H ₂ C ₃ O	Methylene ketene	0.0	Not observed	
	Propynal	7.3	Observed	[82]
	Cyclopropenone	10.6	observed	[83]

Table 7: Isomerization enthalpies for isomers with 6 atoms

H ₃ CON	Formamide	0.0	Observed	[84]
	Hydroxymethylim ine	12.6	Not observed	
	Nitrosomethane	60.8	Not observed	
C_5S	$1-C_5S$	0.0	Observed	[62; 85]
	$c-C_5S^{\#}$	26.9	Not observed	
	c-C ₅ S ^{##}	41.8	Not observed	
	c-C ₅ S ^{###}	84.1	Not observed	
	C ₄ SC	114.2	Not observed	
C ₅ O	1-C ₅ O	0.0	Not observed	
	c-C ₅ O ^o	28.6	Not observed	
	c-C ₅ O ^{σσ}	46.3	Not observed	
	$c-C_5O^{\sigma\sigma\sigma}$	84.5	Not observed	
	C ₄ OC	114.3	Not observed	

*Figure 1a, **Figure 1b, ***Figure 1c; ^aFigure 2a, ^bFigure 2b, ^cFigure 2c; [#]Figure 3a, ^{##}Figure 3b, ^{###}Figure 3c; ^dFigure 4a, ^eFigure 4b; ^oFigure 5a, ^{oo}Figure 5b, ^{ooo}Figure 5c.



Figure 1: Optimized structures of cyclic C_5N isomers.



Figure 2: Optimized structures of cyclic C₅H isomers.



Figure 3: Optimized structures of cyclic C₅S isomers.



Figure 4: Optimized structures of cyclic SiCH₃N isomers.



Figure 5: Optimized structures of cyclic C₅O isomers.

Isomers with 7 atoms

Table 8 shows the isomerization energies and the current astronomical status of different isomeric species with 7 atoms investigated in this study. Of the 22 molecular species in the Table, 7 have been astronomically observed [86-93]. In the H₄C₂O isomeric group where all the isomers have been detected, the isomerization energies range from 12.2 to 27.8 kcal/mol. All the non-observed isomers in the H₃C₃N and C₃H₄ isomeric groups including, cyanomethanol, iminoacetaldehyde, and methyl cyanate (from C₂H₃NO isomeric group) have isomerization enthalpies in the range calculated for the H₄C₂O isomers which have all been detected. This points to the possibility of detecting these molecules in ISM with relative enthalpy of formation within those of the known interstellar molecules. There is no cyanide/isocyanide pair observed among these systems, however, the relative enthalpy of formation in the range of 19.9 to 27.3 kcal/mol is within the range of those noted for other observed cyanide/isocyanide pairs.

As in the previous examples, all the observed isomers are the most stable ones in their respective groups. In the H_4C_2O isomeric group where all the stable isomers have been observed, the most stable isomer; acetaldehyde was first observed before the other isomers. As would be expected, the most stable isomer (acetaldehyde) is present in high abundance in

the different astronomical sources where it has been detected as compared to the abundances of the less stable isomers in the same sources [94-96].

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$	Astronomical status	Reference
	Acetaldebyde		Observed	[06 07]
H_4C_2O	Acctaidenyde	0.0	Observed	[00-07]
	Vinyl alcohol (syn)	12.2	Observed	[88]
	Vinyl alcohol (anti)	13.9	Observed	[88]
	Ethylene oxide	27.8	Observed	[89]
H_3C_3N	Acrylonitrile	0.0	Observed	[90]
	Isocyanoethene	19.9	Not observed	
CaHa	CH2C2H	0.0	Observed	[91]
	H ₂ CCCH ₂	7.8	Not observed	[71]
	c-C ₃ H ₄	23.6	Not observed	
C ₂ H ₃ NO	Methyl isocyanate	0.0	Observed	[92-93]
	Cyanomethanol	13.6	Not observed	
	Iminoacetaldehyde	20.1	Not observed	
	Methyl cyanate	27.3	Not observed	
	2-Aziridinone	36.0	Not observed	
	2-Oxiranimine	40.8	Not observed	
	Methyl fulminic acid	56.5	Not observed	
	2-Iminoethenol	60.9	Not observed	
	Nitrosoethene	69.3	Not observed	
	2H-1,2-Oxazete	79.8	Not observed	
	Methyl isofulminic acid	84.8	Not observed	
	N- Hydroxyacetylenamin	93.6	Not observed	
	(Aminooxy)acetylene	94.7	Not observed	

Table 8: Isomerization enthalpies for isomers with 7 atoms

Isomers with 8 atoms

Table 9 contains the isomerization energies for 33 isomeric species from 7 groups containing eight atoms each. The astronomical statuses of these species are also shown in the table. Figure 6 highlights the cyclic isomers from the C_2H_5N and CH_4N_2O groups. As shown in Table 9, 11 of these species have been astronomically detected [97-109]. The isomerization energies of these species range from 3.1 to 50.0 kcal/mol. Except for 1,2-dioxetane, 1,3-dioxetane, and epoxyproprene, the isomerization energies for all the unknown isomers with 8 atoms fall within the range (of 13.0 to 46.9 kcal/mol) of the known isomers (3.1 to 50.0 kcal/mol). Thus, some of the unknown isomers in Table 9 could be formed from their most stable isomers (which are probably more abundant) via the isomerization process.

The three cyanide/isocyanide pairs among these isomers have isomerization enthalpies ranging from 19.0 to 25.6 kcal/mol. Though no cyanide/isocyanide pair has been detected among these molecular species, the relative enthalpy of formation is within the range of those where the pairs have been detected. CH₃CCNC and H₂NCH₂NC could be formed from their corresponding cyanides that have been detected, thus, they could also be detected provided their spectroscopic parameters are accurately known. Methyl ketene, the most stable isomer of the H₄C₃O group is yet to be astronomically detected. The reasons for its presence and detectability in ISM are the same as discussed for methylene ketene (among the isomers with 6 atoms).

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$ (kcal/mol)	Astronomical status	References
C ₄ H ₃ N	CH ₃ CCCN	0.0	Observed	[97]
	CH ₂ CCHCN	3.1(0.0)	Observed	[98-99]
	HCCCH ₂ CN	13.0	Not observed	
	CH ₃ CCNC	25.6	Not observed	
	CH ₂ CCHNC	25.7 (22.6)	Not observed	
$C_2H_4N_2$	H ₂ NCH ₂ CN	0.0	Observed	[100]
	H ₂ NCH ₂ NC	19.0	Not observed	
		0.0		
$C_2H_4O_2$	Acetic acid	0.0	Observed	[101]
	Methylformate	17.7	Observed	[102-103]
	Glycolaldehyde	33.2	Observed	[104]
	1,3-dioxetane	52.9	Not observed	
	1,2-dioxetane	103.0	Not observed	
H_4C_3O	Methyl ketene	0.0	Not observed	
	Propenal	2.3	Observed	[105]
	Cyclopropanone	17.3	Not observed	
	Propynol	30.8	Not observed	
	Propargyl alcohol	35.2	Not observed	
	Methoxy ethyne	42.0	Not observed	
	1-cyclopropenol	44.0	Not observed	
	2-cyclopropenol	45.5	Not observed	
	Epoxypropene	65.8	Not observed	
H ₂ C ₆	HC ₆ H	0.0	Observed	[106]
	H_2C_6	50.0	Observed	[107]
				51003
C_2H_5N	CH ₃ CHNH	0.0	Observed	[108]
	H ₂ CCHNH ₂	2.8	Not observed	

Table 9: Isomerization enthalpies for isomers with 8 atoms

	CH3NCH2	8.1	Not observed	
	$c-C_2H_5N^m$	19.8	Not observed	
CH ₄ N ₂ O	H ₂ NCONH ₂	0.0	Observed	[109]
	H ₂ NNHCHO	11.0	Not observed	
	HN ₂ CH ₂ OH	35.0	Not observed	
	c- CH ₄ N ₂ O ⁿ	42.4	Not observed	
	CH ₃ NHNO	43.4	Not observed	
	H ₂ NCHNHO	46.9	Not observed	

^mFigure 5a; ⁿFigure 5b.



Figure 6: Optimized structures of cyclic C₂H₅N (A) and CH₄N₂O (B) isomers.

Isomers with 9 atoms

In Table 10, 28 isomeric species from 7 groups are presented with their isomerization enthalpies and current astronomical status. Eight of these species are known interstellar molecular species [110-120]. Figure 7 shows some of the cyclic molecules with 9 atoms. Among the species shown in Table 10, only ethanol and dimethyl ether from the C_2H_6O group have been observed in more than one isomeric form with a relative enthalpy of formation of 12.6 kcal/mol. CH₃SCH₃, the S-analogue of dimethyl ether remains a potential candidate for astronomical detection for many reasons; the well-established chemistry of S and O-containing interstellar molecules, the high abundance of CH₃CH₂SH (the most stable isomer of the group which can isomerize to CH₃SCH₃) and the low relative enthalpy of formation (1.4 kcal/mol) as compared to that of dimethyl ether (12.6 kcal/mol). Assuming accurate spectroscopic parameters are available, interstellar CH₃SCH₃ will soon become a reality.

Isomerization remains one of the plausible formation routes for the formation of the less stable isomers of the different groups here whose most stable isomers have been detected. The isomerization enthalpies for the two cyanide/isocyanide pairs range from 20.8 to 27.1 kcal/mol. Though none of these pairs has been detected, the possibility of their formation from their corresponding cyanides (via isomerization) which could lead to their successful detection cannot be ruled out. HC_6NC is the highest member of the $HC_{2n}NC$ linear chains with experimentally measured rotational transitions that can be used for its astronomical search [121]. However, a recent study using a combined experimental and theoretical approach has provided accurate rotational constants for higher members of the $HC_{2n}NC$ linear chains [6]. HNC remains the only member of this series that has been detected in ISM [122].

Isomoria group	Isomars	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$	Astronomical	References
isomeric group	Isomers	(kcal/mol)	status	
C ₂ H ₆ O	Ethanol	0.0	Observed	[110-111]
	Dimethyl ether	12.6	Observed	[112]
C_2H_6S	CH ₃ CH ₂ SH	0.0	Observed	[113]
	CH ₃ SCH ₃	1.4	Not observed	
C ₂ H ₅ ON	Acetamide	0.0	Observed	[114]
	N-methylformamide	9.7	Not observed	
	Nitrosoethane	64.7	Not observed	
	1-aziridnol	77.9	Not observed	
	Cyanoethoxyamide	134.4	Not observed	
C_3H_5N	Cyanoethane	0.0	Observed	[115]
	Isocyanoethane	20.8	Not observed	
	Propylenimine	22.0	Not observed	
	2-propen-1-imine	25.3	Not observed	
	N-methylene ethenamine	26.0	Not observed	
	Azastene	32.2	Not observed	

Table 10: Isomerization enthalpies for isomers with 9 atoms

	Cyclopropanimine	37.2	Not observed	
	Methylene azaridine	43.8	Not observed	
	Propargylamine	45.1	Not observed	
	1- azabicyclo(1.1.0)buta ne	53.5	Not observed	
C_5H_4	CH ₃ C ₄ H	0.0	Observed	[116]
	$H_2C_3HC_2H$	5.2	Not observed	
	H ₂ C ₅ H ₂	8.1	Not observed	
	$c-C_5H_4^x$	26.1	Not observed	
	$c-C_5H_4^y$	31.6	Not observed	
C_3H_6	CH ₃ CHCH ₂	0.0	Observed	[117]
	$c-C_3H_6^z$	8.2	Not observed	
C ₇ HN	HC ₇ N	0.0	Observed	[118-120]
	HC ₆ NC	27.1	Not observed	

^xFigure 6a; ^yFigure 6b; ^zFigure 6c.



Figure 7: Optimized structures of cyclic C₅H₄ (A and B) and C₃H₆ (C) isomers.

Isomers with 10 atoms

Isomerization energies and astronomical statuses for 16 molecular species from three isomeric groups comprising 10 atoms each are presented in Table 11. Of these three groups, only the C_3H_6O group contains isomers (propanone and propanal) that have been detected in more than one isomeric form with a relative enthalpy of formation of 8.1 kcal/mol [123-127]. $CH_3(CC)_2NC$ is the isocyanide analogue of $CH_3(CC)_2CN$ that has been detected. It has a relative enthalpy of formation of 25.3 kcal/mol which is within the range of the known cyanide/isocyanide pairs. However, there is little or no information regarding the rotational spectrum of $CH_3(CC)_2NC$ that could warrant its astronomical search. Thus, the availability of

accurate spectroscopic parameters for this species remains the starting point in the steps toward its astronomical detection.

1,1-ethanediol; the most stable isomer of the $C_2H_6O_2$ group has not been detected whereas ethylene glycol, the next stable isomer of the group with an isomerization enthalpy of 6.1kcal/mol has been detected in good abundance. The delayed astronomical detection of 1,1-ethanediol is directly linked to a lack of spectroscopic parameters for this molecule. The rotational spectrum of 1,1-ethanediol is yet to be probed either experimentally or theoretically. Once this is done, this molecule could be successfully observed.

Isomeric group	Isomers	Relative $\Delta_{\rm f} {\rm H}^{\rm O}$ (kcal/mol)	Astronomical status	References
$C_2H_6O_2$	1,1-Ethanediol	0.0	Not observed	
	Ethylene glycol	6.1	Observed	[123]
	Methoxy methanol	15.9	Not observed	
	Ethyl hydroperoxide	51.9	Not observed	
	Dimethane peroxide	57.7	Not observed	
C ₃ H ₆ O	Propanone	0.0	Observed	[124-125]
	Propanal	8.1	Observed	[126]
	Propen-2-ol	13.8	Not observed	
	1-propen-1-ol	18.6	Not observed	
	Methoxy ethene	25.6	Not observed	
	2-propene-1-ol	27.1	Not observed	
	1,2-epoxypropane	30.0	Not observed	
	Cyclopropanol	31.2	Not observed	
	Oxetane	33.2	Not observed	
C ₆ H ₃ N	CH ₃ (CC) ₂ CN	0.0	Observed	[127]
	CH ₃ (CC) ₂ NC	25.3	Not observed	

Table 11: Isomerization enthal	pies for isomers with 10 atoms
--------------------------------	--------------------------------

Isomers with 11 atoms

There are currently seven known interstellar molecules (HC_9N , CH_3C_6H , ethyl formate, methyl acetate, hydroxyacetone, cyclopentadiene and ethanolamine) containing 11 atoms [128-134]. In Table 12, the 9 isomeric species from two groups containing 11 atoms are

shown with their isomerization enthalpies and current astronomical status. For the known isomers in the $C_3H_6O_2$ group, the isomerization energy ranges from 11.8 to 14.3 kcal/mol. The non-detection or delayed detection of propanoic acid (the most stable isomer of the group) has been traced to the effect of interstellar hydrogen bonding on the surface of the dust grains which reduces the overall abundance of this species in the gas phase, thus, influencing its successful astronomical observation. HC₉N is the second largest member of the cyanopolyyne chain that has been detected in ISM. As seen in other cyanide/isocyanide pairs, the isomerization of HC₉N to HC₈NC with the relative enthalpy of formation of 27.2 is achievable. An accurate rotational constant for HC₈NC is now available from a recent study [6; 17]. Thus, HC₈NC could be astronomically searched.

Isomeric group	Isomers	Relative ∆ _f H ^O (kcal/mol)	Astronomical status	References
C ₃ H ₆ O ₂	Propanoic acid	0.0	Not observed	
	Ethylformate	11.8	Observed	[128]
	Methyl acetate	14.3	Observed	[129]
	Lactaldehyde	28.1	Not observed	
	Dioxolane	36.1	Not observed	
	Glycidol	52.1	Not observed	
	Dimethyldioxirane	81.7	Not observed	
C ₉ HN	HC ₉ N	0.0	Observed	[130]
	HC ₈ NC	27.2	Not observed	

Table 12: Isomerization enthalpies for isomers with 11 atoms

Isomers with 12 atoms

Only very few cyclic molecules are known among the interstellar molecular species [17; 20-21. From the different isomeric groups considered in this study, it is clear that the cyclic isomers are found to be among the less stable isomers in their respective groups as compared to their corresponding straight-chain analogues. This low stability could affect their interstellar abundance and influences their astronomical observation. The C_6H_6 isomeric group is one of the few groups where the cyclic molecules are found to be the most stable species. Apart from cyanocyclopentadiene and 2-cyanocyclopentadiene that have been recently observed in the interstellar medium [135-136], the other four known interstellar molecules with 12 atoms, [137-140] their corresponding isomers, and isomerization enthalpies are presented in Table 13.

The only known branched-chain interstellar molecule; isopropyl cyanide is almost equivalent in energy to its linear chain analogue; propyl cyanide with a relative enthalpy of formation of 0.4kcal/mol. Propan-2-ol and propanol (the two most stable isomers of the C_3H_6O group) are yet to be astronomically observed. These species have also been shown to be strongly affected by interstellar hydrogen bonding as compared to ethyl methyl ether that has been detected [6; 18-21]. The stability of ethyl methyl ether is also enhanced by the stabilizing effect of the two alkyl substituents while propanol and propan-2-ol have only one alkyl substituent each.

Isomeric group	Isomers	Relative Δ _f H ^O (kcal/mol)	Astronomical status	References
C ₆ H ₆	Benzene	0.0	Detected	[137]
	Fulvene	34.2	Not detected	
	3,4-dimethylenecyclopropene	62.0	Not detected	
	1,5-hexadiene-3-yne	62.3	Not detected	
	2,4-hexadiyne	65.2	Not detected	
	1,2,3,4-hexateraene	71.7	Not detected	
	1,3-hexadiyne	73.2	Not detected	
	Trimethylenecyclopropane	79.8	Not detected	
	1,4-hexadiyne	82.3	Not detected	
	1,5-hexadiyne	85.8	Not detected	
C ₃ H ₈ O	Propan-2-ol	0.0	Not observed	
	Propanol	3.7	Not observed	
	Ethyl methyl ether	8.2	Observed	[139]
C_4H_7N	Isopropyl cyanide	0.0	Observed	[138]
	Propyl cyanide	0.4	Observed	[140]
	2-isocyanopropane	18.2	Not observed	
	2-aminobutadiene	18.2	Not observed	
	3-pyrroline	20.2	Not observed	
	2,2-dimethylethylenimine	20.5	Not observed	
	But-1-en-1-imine	22.6	Not observed	
	2,3-butadiene-1-amine	38.1	Not observed	
	N-vinylazaridine	40.1	Not observed	
	N-methyl-1-propyn-1-amine	41.4	Not observed	
	3-butyn-1-amine	43.2	Not observed	
	N-methyl propargylamine	49.3	Not observed	

Table 13: Isomerization enthalpies for isomers with 12 atoms

2-azabicyclo(2.1.0)pentane	51.9	Not observed	
----------------------------	------	--------------	--

Potential Interstellar Molecules

Knowing the right candidates for astronomical searches is vital in reducing the number of unsuccessful astronomical searches considering the time, energy and resources involved in these projects. From the present study, few molecular species have been identified as potential interstellar molecules. These are briefly summarized below:

Isocyanomethylidyne, CNC: This is the isocyanide analogue of C₂N that has been detected [141]. CNC is found to be more stable than its cyanide analogue. Table 14 contains the relative energies of CNC and CCN from different quantum chemical calculation methods while Figure 8 shows the optimized geometries of these isomers at the MP2(full)/6-311++G** level with the bond angle in degrees and bond distance in angstroms. As discussed under the isomers with 3 atoms, all the observed species are the most stable ones and these stable species are also found to be more abundant. Where both species have been detected, the most stable is reported to present in high abundance than the less stable. For instance, HCN is found to be more abundant than HNC in different molecular clouds and this is also the case for the MgNC/MgCN abundance ratio measured in the asymptotic giant branch (AGM) stars [142-145]. CNC is more stable than CCN which implies that CNC should be present in high abundance in ISM than CCN that has been detected. Thus, CNC remains a potential candidate for astronomical detection. CNC is microwave inactive with a zero-dipole moment; thus, infrared astronomy remains the best approach for its astronomical observation.

Mathad	Relative energy (kcal/mol)		
Method	CNC	CCN	
CCSD/6-311++G**	0.0	1.5	
MP2(full)/6-311++G**	0.0	6.2	
B3LYP/6-311++G**	0.0	2.0	
G4	0.0	3.7	

Table 14: Relative energies of CNC and CCN



Figure 8: Optimized geometries of CNC and CCN at MP2(full)/6-311++G** level.

NCCN: The rationale for the choice of NCCN as a potential interstellar molecule is well discussed under the isomers with 4 atoms.

c-C₅H: With the limited number of known cyclic interstellar molecules, it is always exciting finding rings that are as stable as their corresponding chains. $c-C_5H$ (Figure 2a) is the cyclic analogue of C₅H that has been detected. From Table 7, the relative enthalpy of formation between the linear chain and this cyclic analogue is just 1 kcal/mol. Table 15 shows the relative energies of these species at different quantum chemical calculation methods. While the MP2 method predicts the cyclic isomer to be more stable than the linear, others methods predict the reverse but the magnitude of the difference in energy is much at the MP2 level as compared to other methods. Figure 9 displays the optimized structures of these species. As shown in Table 15, $c-C_5H$ is microwave active with a very good dipole moment making its astronomical searches in the radio frequency possible. If the spectroscopic parameters of c-C5H can be accurately probed, either experimentally or theoretically, the possibility of its astronomical observation is very high.

Method	Relative energy (kcal/mol)		Dipole moment (Debye)	
	c-C ₅ H	1-C ₅ H	c-C ₅ H	1-C_5H
MP2(full)/6-311++G**	0.0	18.7	3.4	4.2
MP2/aug-cc-pVTZ	0.0	17.5	3.4	4.2
G4	10.9	0.0	3.7	4.6
CCSD/6-311++G**	3.0	0.0	3.4	4.6
B3LYP/6-311++G**	10.4	0.0	3.7	5.2
B3LYP/aug-cc-pVTZ	10.3	0.0	3.7	5.2

Table 15: Relative energies of linear and cyclic stable isomers of C₅H



Figure 9: Optimized structures of C₅H most stable isomers.

Methylene and Methyl Ketenes: As highlighted in the text, the ketenes are found to be more stable than their corresponding isomers in all the isomeric groups examined in this study (Tables 6, 7 and 9) and as such be present in detectable amounts in ISM. Ketenes have also been shown to be less affected by hydrogen bonding on the surface of the interstellar dust grains as compared to their corresponding isomers. Also, the fact that ketene (H₂C₂O), ketenyl radical (HC₂O) and isomers of ketenes (like propynal, cyclopropenone, and propenal) are known interstellar molecular species further supports the presence and detectability of

methyl ketene and methylene ketene in ISM. Thus, methyl ketene and methylene kenetne remain potential interstellar molecules pending their astronomical searches. No unsuccessful astronomical search has been reported for any of these ketenes.

C₅**O**: This is the oxygen analogue of C₅S which is a known interstellar molecule. Without any exception, an interstellar O-containing molecular species is more abundant than its S-analogue. This could simply be traced to the cosmic abundance of O and S. Thus, for every known S-species, the O-analogue is not only present in detectable abundance, but it can also be said to have even been overdue for astronomical detection because for sure the O-species are more abundant than their S-analogue and as such could be detected with less difficulty as compared to its S-analogue. The column density of 2.14x 10¹² cm⁻² reported for C₅S suggests thehigh abundance of C₅O in ISM [62; 85]. The microwave spectrum of C₅O that could guide its successful astronomical observation is available [146]. Interstellar C₅O is just a matter of time.

CH₃SCH₃: This is the S-analogue of dimethyl ether; a known interstellar molecule. The interstellar chemistry of S- and O-containing species is well established. Every known O-containing interstellar molecule points to the presence and detectability of the S-analogue. As earlier mentioned, except where the effect of interstellar hydrogen bonding dominates, the ratio of an interstellar sulphur molecule to its oxygen analogue is close to the cosmic S/O ratio. Assuming dimethyl ether is formed from ethanol via isomerization with a barrier of 12.6 kcal/mol, then the isomerization of C_2H_5SH to CH_3SCH_3 is a much more feasible process with a relative enthalpy of formation of just 1.4 kcal/mol (nine times lower than that of CH_3OCH_3) compared to that of the O-analogue. Thus, because of the unique chemistry of S- and O-containing interstellar molecules and the low relative enthalpy of formation, CH_3SCH_3 is considered a potential candidate for astronomical observation assuming its spectroscopic parameters are accurately known.

1,1-Ethanediol, Propanoic Acid, Propan-2-ol, and Propanol: These molecules are found to be the most stable isomers in their respective groups (Tables 11, 12, and 13). Whereas their less stable isomers have been astronomically detected; successful detection of these most stable isomers is highly feasible. The spectroscopic parameters for propanoic acid, propan-2-ol, and propanol are known but those of 1,1-ethanediol are yet to come by. Though the delayed astronomical observations of some of these species have been linked to the effect of interstellar hydrogen bonding; there are high chances for their successful astronomical detection.

Conclusion: An extensive investigation of the isomerization energies of 246 molecular species from 65 isomeric groups is reported in this study. From the results, isomerization is found to be one of the plausible mechanisms for the formation of molecules in ISM. The high abundances of the most stable isomers coupled with the energy sources in ISM drive the isomerization process even for barriers as high as 67.4 kcal/mol. However, the chemical reaction pathways can also influence the final abundance of any species in the ISM. Specifically, the isomerization process is found to be very effective in converting cyanides to their corresponding isocyanides and vice versa. Thus, for every cyanide or isocyanide, the corresponding isocyanide or cyanide could be synthesized via isomerization. Also, from the results, the following potential interstellar molecules; CNC, NCCN, c-C₅H,

methylene ketene, methyl Ketene, CH_3SCH_3 , C_5O , 1,1-ethanediol, propanoic acid, propan-2ol, and propanol are highlighted and discussed

Acknowledgement: EEE acknowledges a research fellowship from the Indian Institute of Science, Bangalore, and fruitful discussions with Prof. E. Arunan of the same Institute. The visiting position from the Indian Institute of Technology Bombay where the final part of this work was done is also acknowledged.

References

[1] M. Agúndez, C. Cabezas, B. Tercero, N. Marcelino, J. D. Gallego, P. de Vicente, J. Cernicharo, "Discovery of the propargyl radical (CH2CCH) in TMC-1: One of the most abundant radicals ever found and a key species for cyclization to benzene in cold dark clouds", A&A **647** (2021) L10.

[2] B. A. McGuire, "Census of Interstellar, Circumstellar, Extragalactic, Protoplanetary Disk, and Exoplanetary Molecules". Astrophysical Journal Supplement Series, **239** (2018) 17.

[3] E. E. Etim, E. Arunan, "Rotational Spectroscopy and Interstellar Molecules", Planex Newletter, **5**(2015) 16.

[4] E. E. Etim, E. Arunan, "Interstellar Isomeric Species: Energy,Stability and AbundanceRelationship",EuropeanPhysicalJournalPlus,131(2016)448.https://doi.org/10.1140/epjp/i2016-16448-0

[5] E. E. Etim, E. Arunan, "Partition Function and Astronomical Observation of Interstellar Isomers: Is there a link?", Advances in Space Research, **59**(2017)1161. http://dx.doi.org/10.1016/j.asr.2016.11.021

[6] E. E. Etim, E. Arunan, "Accurate Rotational Constants for linear Interstellar Carbon Chains: Achieving Experimental Accuracy", Astrophysics and Space Science, 362 (2017)
4.DOI 10.1007/s10509-016-2979-6

[7] H. J. Fraser, M. R. S. McCoustra., D. A. Williams, "The Molecular Universe", A &G; **43** (2002) 10. https://doi.org/10.1046/j.1468-4004.2002.43210.x

[8]A. G. G. M. Tielens, "The Molecular Universe", Rev. Mod. Phys. 85 (2013) 1021.

[9] M. A. Frerking, R. A. Linke, P. Thaddeus, "Interstellar Isothiocyanic Acid", ApJ, 234 (1979) 143.

[10] R. A. Linke, M. A. Frerking, P. Thaddeus, "Interstellar Methyl Mercaptan", ApJ, **234**(1979) L139.

[11] J. M. Hollis, P. R. Jewell, F. J. Lovas, A. Remijan, H. Møllendal, "Green Bank Telescope Detection of New Interstellar Aldehydes: Propenal and Propanal" ApJ **610** (2004)L21.

[12] E. Oladimeji, E. E. Etim, M. Ojo, "The thermodynamic properties of interstellar isomers with 3 atoms", Advances in Space Research **70** (2022) 3745.

[13] F. J. Lovas, J. M. Hollis, A. J. Remijan, P. R. Jewell, "Detection of Ketenimine (CH₂CNH) in Sagittarius B2(N) Hot Cores", ApJ, **645** (2006) 137.

[14] M. Lattelais, F. Pauzat, Y. Ellinger, C. Ceccarelli, "Interstellar complex organic molecules and the minimum energy principle", Astrophys J, **696** (2009) L133.

[15] L. A. Curtiss, P. C. Redfern, K. Raghavachari, "Gaussian-4 theory" Jour. Chem. Phys, **126** (2007)084108

[16] L. A. Curtiss, K. Raghavachari, P. C. Redfern, V. Rassolov, J. A. Pople, "Gaussian-3 (G3) theory for molecules containing first and second-row atoms," J. Chem. Phys., **109**(1998)7764.

[17] E. E. Etim, P. Gorai, A.Das, S. Charabati, E. Arunan, "Systematic Theoretical Study on the Interstellar Carbon Chain Molecules", ApJ, **832**(2016) 144. Doi:10.3847/0004-637X/832/2/144

[18] E. E. Etim, P. Gorai, A. Das, E. Arunan, "Interstellar Protonated Molecular Species. Advances in Space Science", **60**(2017)709. https://doi.org/10.1016/j.asr.2017.04.003, 2017

[19] E. E. Etim, P. Gorai, A. Das, S. K. Chakrabarti, E. Arunan, "Interstellar HydrogenBonding", Advances in Space Research, 61(2018)https://doi.org/10.1016/j.asr.2018.03.003

[20] E. E. Etim, P. Gorai, R. Ghosh, A. Das, "Detectable Interstellar Anions: Examining the Key Factors", Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, **230**(2020)118011. https://doi.org/10.1016/j.saa.2019.118011

[21] E. E. Etim, R. O. A. Adelagun, C. Andrew, E. Oladimeji, "Optimizing the Searches for Interstellar Heterocycles", Advances in Space Research Journal, **68** (2022) 3508. https://doi.org/10.1016/j.asr.2021.06.003

[22] E. E. Etim, M. E. Khan, O. E. Godwin, G. O. Ogofotha, "Quantum Chemical Studies on $C_4H_4N_2$ Isomeric Molecular Species", J. Nig. Soc. Phys. Sci. **3** (2022) 429

[23] M. E. Khan, E. E. Etim, V. J. Anyam, A. Abel ,I. G. Osigbemhe, C. T. Agber, "Computational studies on Emodin ($C_{15}H_{10}O_5$) from Methanol extract of Pteridium acquilinum leaves", J. Nig. Soc. Phys. Sci. **3**(2021)360.

[24] B. Zuckerman, M. Morris, P. Palmer, B. E. Turner, "Observations of CS, HCN, U89.2, and U90.7 in NGC 2264", ApJ, **173**(1972)125.

[25] G. L. Blackman, R. D. Brown, P. D. Godfrey, H. I. Gunn, "The Microwave Spectrum of HNC: Identification of U90.7", Nature, **261**(1976)395.

[26] B. E. Turner, T. C. Steimle, L. Meerts, "Detection of Sodium Cyanide (NaCN) in IRC 10216", ApJ. **426** (1994)97.

[27] K. Kawaguchi, E. Kagi, T. Hirano, S. Takano, S. Saito, "Laboratory Spectroscopy of MgNC - The First Radioastronomical Identification of Mg-Bearing Molecule", ApJ, **406** (1993)39.

[28] L. M. Ziurys, A. J. Apponi, M. Guélin, J. Cernicharo, "Detection of MgCN in IRC +10216: A New Metal-Bearing Free Radical", ApJ, **445** (1995) 47.

[29] M. Guélin, J. Cernicharo, C. Kahane, J. Gomez-Gonzales, "A New Free Radical in IRC +10216", A&A **157** (1986) L17.

[30] M. Guelin, R. Lucas, J. Cernicharo, "MgNC and the Carbon-Chain Radicals in IRC +10216", A&A, **280** (1993)19.

[31] L. M. Ziurys, C. Savage, J. L. Highberger, A. J. Apponi, M. Guélin, J.; Cernicharo, "More Metal Cyanide Species: Detection of AlNC (X1Sigma+) toward IRC +10216", ApJ, **564** (2002) 45.

[32] M. Guélin, S. Muller, J. Cernicharo, M. McCarthy, P. Thaddeus, "Detection of the SiNC Radical in IRC+10216", A&A, **426** (2004) L49.

[33] M. Guélin, S. Muller, J. Cernicharo, A. J. Apponi, M. McCarthy, C. A. Gottlieb, P. Thaddeus, "Astronomical Detection of the Free Radical SiCN", A&A, **369** (2000) L9.

[34] L. E. Snyder, J. M. Hollis, B. L. Ulich, "Radio Detection of the Interstellar Formyl Radical", ApJ **208** (1976) L91.

[35] M. Agúndez, J. Cernicharo, M. Guélin, "Discovery of Phosphaethyne (HCP) in Space: Phosphorus Chemistry in Circumstellar Envelopes", ApJ **662** (2007) L91.

[36] J. K. Anderson, L. M. Ziurys, "Detection of CCN (X2Πr) in IRC+10216: Constraining Carbon-chain Chemistry", ApJ **795** (2014) L1.

[37] D. T. Halfen, D. T. Clouthier, L. M. Ziurys, "Detection of the CCP Radical (X $2\Pi r$) in IRC +10216: A New Interstellar Phosphorus-containing Species", ApJ **677** (2008) L101.

[38] D. Buhl, L. E. Snyder, The Detection of a MM-Wave Transition of Methlacetylene, in Molecules in the Galactic Environment (eds. M. A. Gordon and L. E. Snyder), Wiley-Interscience, New York, 1973, 187-195.

[39] R.C. Woods, C. S. Gudeman, R. L. Dickman, P. F. Goldsmith, G. R. Huguenin, W. M. Irvine, A. Hjalmarson, L. A. Nyman, H. Olofsson, "The [HCO+]/[HOC+] Abundance Ratio in Molecular Clouds", ApJ **270** (1983) 583.

[40] L. M. Ziurys, A. J. Apponi, M. Guélin, J. Cernicharo, "Detection of MgCN in IRC +10216: A New Metal-Bearing Free Radical", ApJ, **445** (1995) 47.

[41] P. Thaddeus, M. Guélin, R. A. Linke, "Three New "Nonterrestrial" Molecules", ApJ 246 (1971) L41.

[42] B. T. Soifer, R. C. Puetter, R. W. Russell, S. P. Willner, P. M. Harvey, F. C. Gillett, "The 4-8-micron spectrum of the infrared source W33A", ApJ, **232** (1979) 53.

[43] M. Agúndez, N. Marcelino, J. Cernicharo, M. Tafafa, "Detection of interstellar HCS and its metastable isomer HSC: new pieces in the puzzle of sulfur chemistry", A&A, **611** (2018) L1.

[44] A. M. Shaw, A. M. Astrochemistry: From Astronomy to Astrobiology. Wiley, 2006.

[45] L. E. Snyder, D. Buhl, "Interstellar Isocyanic Acid", ApJ., 177 (1972) 619.

[46] S. Brunken, C. A. Gottlieb, M. C. McCarthy, P. Thaddeus, "Laboratory Detection of HOCN and Tentative Identification in Sgr B2", ApJ, **697** (2009) 880.

[47] S. Brunken, A. Belloche, S. Martin, L. Verheyen, K. M. Menten, "Interstellar HOCN in the Galactic centre region", A&A, **516** (2010) A109.

[48] N. Marcelino, J. Cernicharo, B. Tercero, E. Roueff, "Discovery of Fulminic Acid, HCNO in Dark Clouds" ApJ., **690** (2009) 27.

[49] M. A. Frerking, R.A. Linke, P. Thaddeus, "Interstellar Isothiocyanic Acid", ApJ 234 (1979) L143.

[50] D. T. Halfen, L. M. Ziurys, S. Brünken, C. A. Gottlieb, M. C. McCarthy, P. Thaddeus, "Detection of a New Interstellar Molecule: Thiocyanic Acid HSCN. ApJ, **702** (2009) 124. [51] P. Thaddeus, J. M. Vrtilek, C. A. Gottlieb, "Laboratory and Astronomical Identification of Cyclopropenylidene, C₃H₂". ApJ **299** (1985) L63.

[52] S. Yamamoto, S. Saito, M. Ohishi, S. Suzuki, S. I. Ishikawa, N. Kaifu, A. Murakami, "Laboratory and Astronomical Detection of the Cyclic C₃H Radical", ApJ **322** (1987) L55.

[53] M. Guélin, P. Thaddeus, "Tentative Detection of the C_3N Radical", ApJ **212** (1977) L81.

[54] P. Friberg, A. Hjalmarson, W. M. Irvine, M. Guélin, Interstellar C_3N - Detection in Taurus Dark Clouds. ApJ **241** (1980) L99.

[55] M. Guélin, J. Cernicharo, "Astronomical Detection of HCCN Radical- Toward a New Family of Carbon-Chain Molecules?" A&A, **244** (1991) L21.

[56] H. E. Matthews, W. M. Irvine, P. Friberg, R. D. Brown, P. D. Godfrey, "A New Interstellar Molecule- Tricarbon Monoxide", Nature **310** (1984) 125

[57] R. D. Brown, P. D. Godfrey, D. M. Cragg, D. H. N. Rice, W. M. Irvine, P. Friberg, H. Suzuki, M. Ohishi, N. Kaifu, M. Morimoto, "Tricarbon Monoxide in TMC-1", ApJ **297** (1985) 302.

[58] N. Kaifu, H. Suzuki, M. Ohishi, T. Miyaji, S. Ishikawa, T. Kasuga, M. Morimoto, "Detection of intense unidentified lines in TMC-1", ApJ **317** (1987) L111.

[59] S. Yamamoto, S. Saito, K. Kawaguchi, N. Kaifu, H. Suzuki, "Laboratory Detection of a New Carbon-Chain Molecule C3S and Its Astronomical Identification", ApJ 317 (1987): L119.

[60] J. Cernicharo, C. Kahane, M. Guélin, H. Hein,"Sulfur in IRC +10216", A&A **181** (1987) L9.

[61] A. J. Apponi, M. C. McCarthy, C. A. Gottlieb, P. Thaddeus, "Astronomical Detection of Rhomboidal SIC3", ApJ **516** (1999) L103.

[62] M. Agúndez, J. Cernicharo, M. Guélin, "New molecules in IRC +10216: confirmation of C_5S and tentative identification of MgCCH, NCCP, and SiH₃CN" A&A **570** (2014) A45.

[63] M. Agúndez, N. Marcelino, J. Cernicharo, "Discovery of Interstellar Isocyanogen (CNCN): Further Evidence that Dicyanopolyynes Are Abundant in Space", ApJL **861** (2018) K22.

[64] M. Agúndez, J. Cernicharo, M. Guélin, "Discovery of interstellar ketenyl (HCCO), a surprisingly abundant radical", A&A **577** (2015) L5.

[65] B. E. Turner, Detection of Interstellar Cyanoacetylene. ApJ, 163 (1971) 35.

[66] K. Kawaguchi, M. Ohishi, S. I. Ishikawa, N. Kaifu, "Detection of Isocyanoacetylene HCCNC in TMC-1", ApJ, 386 (1992) 51.

[67] K. Kawaguchi, S. Takano, M. Ohishi, S. I. Ishikawa, K. Miyazawa, N. Kaifu, K. Yamashita, S. Yamamoto, S. Saito, Y. Ohshima, Y. Endo, "Detection of HNCCC in TMC-1", ApJ, 396 (1992) 49.

[68] B. E. Turner, "Microwave Detection of Interstellar Ketene", ApJ 213 (1977) L75.

[69] P. Thaddeus, C. A. Gottlieb, A. Hjalmarson, L. E. B. Johansson, W. M. Irvine, P. Friberg, R. A. Linke, "Astronomical Identification of the C3H Radical", ApJ 294 (1985) L49.

[70] J. Cernicharo, C. A. Gottlieb, M. Guélin, T. C. Killian, P. Thaddeus, J. M. Vrtilek, "Astronomical Detection of H2CCCC", ApJ 368 (1991) L43.

[71] B. E. Turner, H. S. Liszt, N. Kaifu, A. G. Kisliakov, "Microwave Detection of Interstellar Cyanamide", ApJ, 201 (1975) 149.

[72] W. M. Irvine, P. Friberg, A. Hjalmarson, S. Ishikawa, N. Kaifu, K. Kawaguchi, S. C. Madden, H. E. Matthews, M. Ohishi, S. Saito, H. Suzuki, P. Thaddeus, B. E. Turner, S. Yamamoto, L. M. Ziurys, "Identification of the Interstellar Cyanomethyl Radical (CH₂CN) in the Molecular Clouds TMC-1 and Sagittarius B2. ApJ, **334** (1988) 107.

[73] A. J. Remijan, J. M. Hollis, F. J. Lovas, W. D. Stork, P. R. Jewell, D. S. Meier, "Detection of Interstellar Cyanoformaldehyde (CNCHO)", ApJ, **675** (2008) L85.

[74] M. Guélin, N. Neininger, J. Cernicharo, Astronomical Detection of the Cyanobutadiynyl Radical C_5N . A&A **355** (1998) L1.

[75] J. Cernicharo, C. Kahane, J. Gómez-González, M. Guélin, "Tentative Detection of the C_5H Radical", A&A **164** (1986) L1.

[76] J. Cernicharo, C. Kahane, J. Gómez-González, M. Guélin, "Detection of the 2Pi3/2 state of C₅H", A&A, 167 (1986) L5.

[77] J. Cernicharo, M. Guelin, C. M. Walmsley, "Detection of the Hyperfine Structure of the C_5H Radical", A&A, **172** (1987) L5.

[78] P. M. Solomon, K. B. Jefferts, A. A. Penzias, R. W. Wilson, "Detection of Millimeter Emission Lines from Interstellar Methyl Cyanide", ApJ. **168** (1971) 107-110.

[79] J. Cernicharo, C. Kahane, M. Guélin, Gomez-Gonzalez, J. Tentative Detection of CH_3NC towards Sgr B2. A&A, **189** (1988) 1.

[80] F. J. Lovas, A. J. Remijan, J. M. Hollis, P. R. Jewell, L. E. Snyder, Hyperfine Structure Identification of Interstellar Cyanoallene toward TMC-1. ApJ **637** (2006) L37.

[81] J. Cernicharo, M. Guélin, J. R. Pardo, "Detection of the Linear Radical HC4N in IRC +10216", ApJ, **615** (2004) 145.

[82] W. M. Irvine, R. D. Brown, D. M. Craig, P. Friberg, P. D. Godfrey, N. Kaifu, H. E. Matthews, M. Ohishi, H. Suzuki, H. Takeo, "A New Interstellar Polyatomic Molecule: Detection of Propynal in the Cold Cloud TMC-1", ApJ, **335** (1988) 89.

[83] J. M. Hollis, F. J. Lovas, J. R. Anthony, A. J. Remijan, P. R. Jewell, V. V. Ilyushin, I. Kleiner, "Detection of Acetamide (CH₃CONH₂): The Largest Interstellar Molecule with a Peptide Bond", ApJ.,643 (2006) 25.

[84] R. H. Rubin, G. W. Swenson, R. C. Solomonm, Jr, H. L. Flygare, "Microwave Detection of Interstellar Formamide", ApJ, **169** (1971) 39.

[85] M. B. Bell, L. W. Avery, P. A. Feldman, "C3S and C_5S in IRC +10216", ApJ **417** (1993) L37.

[86] N. Fourikis, M. W. Sinclair, B. J. Robinson, P. D. Godfrey, R. D. Brown, "Rotational Transition in Interstellar Acetaldehyde", AuJPh, **27** (1974) 425.

[87] W. Gilmore, M. Morris, P. Palmer, D. R. Johnson, F. J. Lovas, B. E. Turner, B. Zuckerman, "Observation of the 6(16)-5(15) Transitions of acetaldehyde in Sagittarius B2", ApJ, **204** (1976) 43.

[88] B. E. Turner, A. J. Apponi, "Microwave Detection of Interstellar Vinyl Alcohol, CH₂CHOH", ApJ, **561** (2001) 207.

[89] J. E. Dickens, W. M. Irvine, M. Ohishi, M. Ikeda, S. Ishikawa, A. Nummelin, A. Hjalmarson, "Detection of Interstellar Ethylene Oxide (c-C2H4O)". ApJ, **489** (1997)753.

[90] F. F. Gardner, G. Winnewisser, "The Detection of Interstellar Vinyl Cyanide (Acrylonitrile)", ApJ,195 (1975) L127.

[91] D. Buhl, L. E. Snyder, "Unidentified Interstellar Microwave Line", Nature 228 (1970) 267.

[92] D. T. Halfen, V. V. Ilyushin, L. M. Ziurys, "Interstellar Detection of Methyl Isocyanate CH_3NCO in Sgr B2(N): A Link from Molecular Clouds to Comets". ApJ **812** (2015) L5

[93] J.Cernicharo, Z. Kisiel, B. Tercero, L. Kolesniková, I. R. Medvedev, A. López, S. Fortman, M. Winnewisser, F. de Lucia, J. L. Alonso, J. C. Guillemin, "A rigorous detection of interstellar CH3NCO: An important missing species in astrochemical networks", A&A 587 (2016) L4.

[94] A. Nummelin, P. Bergman, A. Hjalmarson, P. Friberg, W. M. Irvine, T. J. Millar, M. Ohishi, S. Saito, "Abundances of ethylene oxide and acetaldehyde in hot molecular cloud cores", A&A, 337 (1998) 275.

[95] A. Nummelin, P. Bergman, A. Hjalmarson, P. Friberg, W. M. Irvine, T. J. Millar, M. Ohishi, S. Saito, "Abundances of ethylene oxide and acetaldehyde in hot molecular cloud cores", A&A, 337 (1998) 275.

[96] M. Ikeda, M. Ohishi, J. E. Dickens, P. Bergman, A. Hjalmarson, W. M. Irvine, "Survey Observations of c-C2H4O and CH3CHO toward Massive Star-forming Regions", ApJ, 560 (2001) 792.

[97] N. W. Broten, J. W. MacLeod, L. W. Avery, W. M. Irvine, B. Höglund, P. Friberg, P. Hjalmarson, "The Detection of Interstellar Methylcyanoacetylene", ApJ, 276 (1984) 25.

[98] F. J. Lovas, J. M. Hollis, A. J. Remijan, P. R. Jewell, "Detection of Ketenimine (CH₂CNH) in Sagittarius B2(N) Hot Cores", ApJ, **645** (2006) 137.

[99] J. N. Chin, R. I. Kaiser, C. Lemme, C. Henkel, "Detection of Interstellar Cyanoallene and its Implications for Astrochemistry", AIP Conf. Proc. 855 (2006) 149.

[100] A. Belloche, K. M. C. Menten, C. Comito, H. S. P. Müller, P. Schilke, J. Ott, S. Thorwirth, C.Hieret, "Detection of amino acetonitrile in Sgr B2(N)", A&A., **482** (2008) 179.

[101] M. D. Mehringer, L. E. Snyder, Y. Miao, F. Lovas, "Detection and Confirmation of Interstellar Acetic Acid" ApJ 480 (1997) L71.

[102] E. Churchwell, G. Winnewisser, "Observations of Methyl Formate in the Galactic Center", A&A., 45 (1975) 229.

[103] R. D. Brown, J. G. Crofts, P. D. Godfrey, F. F. Gardner, B. J. Robinson, J. B. Whiteoak, "Discovery of Interstellar Methyl Formate", ApJ, **197** (1975) L29.

[104] J. M. Hollis, F. J. Lovas, P. R. Jewell, "Interstellar Glycolaldehyde: The First Sugar", ApJ, **540** (2000) 107.

[105] J. M. Hollis, P. R. Jewell, F. J. Lovas, A. Remijan, A. Møllendal, "Green Bank Telescope Detection of New Interstellar Aldehydes: Propenal and Propanal", ApJ, 610 (2004) 21.

[106] W. D. Langer, T. Velusamy, T. B. H. Kuiper, R. Peng, M. C. McCarthy, M. J. Travers, A. Kovács, C. A. Gottlieb, P. Thaddeus, "First Astronomical Detection of the Cumulene Carbon Chain molecule H2C6 in TMC-1", ApJ, **480** (1997) 63.

[107] J. Cernicharo, A. M. Heras, A. G. G. M. Tielens, J. R. Pardo, F. Herpin, M. Guélin, L. B. F. M. Waters, "Infrared Space Observatory's Discovery of C_4H_2 , C_6H_2 , and Benzene in CRL 618", ApJ, **546** (2001) L123.

[108] Loomis, et al., "The Detection of Interstellar Ethanimine (CH₃CHNH) from Observations taken during the GBT PRIMOS Survey", ApJL, **765** (2013) L9.

[109] A. J. Remijan, L. E. Snyder, B. A. McGuire, H. L. Kuo, L. W. Looney, D. N. Friedel, G. Y. Golubiatnikov, F. J. Lovas, V. V. Ilyushin, E. A. Alekseev, S. F. Dyubko, B. J. McCall, J. M. Hollis, "Observational Results of a Multi-telescope Campaign in Search of Interstellar Urea [(NH₂)₂CO]", ApJ., **783** (2014) 77.

[110] B. Zuckerman, B. E. Turner, D. R. Johnson, F. O. Clark, F. J. Lovas, N. Fourikis, P. Palmer, M. Morris, A. E. Lilley, J. A. Ball, C. A. Gottlieb, H. Penfield, "Detection of Interstellar trans -Ethyl Alcohol", ApJ., **196** (1975) 99.

[111] J. C. Pearson, K. V. L. N. Sastry, E. Herbst, F. C. De Lucia, "Gauche Ethyl Alcohol: Laboratory Assignments and Interstellar Identification", ApJ, **480** (1997) 420.

[112] L. E. Snyder, D. Buhl, P. R. Schwartz, F. O. Clark, D. R. Johnson, F. J. Lovas, P. T. Giguere, "Radio Detection of Interstellar Dimethyl Ether", ApJ., 191 (1974) 79.

[113] L. Kolesniková, B. Tercero, J. Cernicharo, J. L. Alonso, A. M. Daly, B. P. Gordon, S. T. Shipman, "Spectroscopic Characterization and Detection of Ethyl Mercaptan in Orion", ApJ 784 (2014) L7.

[114] J. M. Hollis, A. J. Remijan, P. R. Jewell, F. J. Lovas, "Cyclopropenone (c-H2C3O): A New Interstellar Ring Molecule", ApJ **642** (2006) 933

[115] D. R. Johnson, F. J. Lovas, C. A. Gottlieb, E. W. Gottlieb, M. M. Litvak, M. Guélin, P. Thaddeus, "Detection of Interstellar Ethyl Cyanide", ApJ., **218** (1977) 370.

[116] C. M. Walmsley, P.R. Jewell, L. E. Snyder, G. Winnewisser, "Detection of Interstellar Methyldiacetylene (CH3C4H) in the Dark Dust Cloud TMC 1", A&A 134 (1984) L11.

[117] N. Marcelino, J. Cernicharo, M. Agúndez, E. Roueff, M. Gerin, J. Martín-Pintado, R. Mauersberger, C. Thum, "Discovery of Interstellar Propylene (CH₂CHCH₃): Missing Links in Interstellar Gas-Phase Chemistry", ApJ 665 (2007) L127.

[118] H. W. Kroto, C. Kirby, D. R. M. Walton, L. W. Avery, N. W. Broten, J. M. MacLeod, T. Oka, "The Detection of Cyanohexatriyne in Heiles's Cloud 2", ApJ 219(1978) L133.

[119] L. T. Little, G. H. MacDonald, P. W. Riley, D. N. Matheson, "Observations of Interstellar HC_5N and HC7N in Dark Dust Clouds", MNRAS 183(1978) 45P.

[120] G. Winnewisser, C. M. Walmsley, "The Detection of HC5N and HC7N in IRC +10216", A&A 70(1978) L37.

[121] P. Botschwina, A. Heyl, "The isocyanopolyynes HC4NC and HC6NC: Microwave spectra and ab initio calculations", J. Chem. Phys., **109** (1998) 3108.

[122] G. L. Blackman, R. D. Brown, P. D. Godfrey, H. I. Gunn, "The Microwave Spectrum of HNC: Identification of U90.7.", Nature, **261** (1976) 395.

[123] J. M. Hollis, P. R. Jewell, F. J. Lovas, A. Remijan, H. Møllendal, "Green Bank Telescope Detection of New Interstellar Aldehydes: Propenal and Propanal" ApJ **610** (2004)L21.

[124] J. M. Hollis, F. J. Lovas, P. R. Jewell, L. H. Coudert, "Interstellar Antifreeze: Ethylene Glycol" ApJ, **571** (2002) 59.

[125] F. Combes, M. Gerin, A. Wooten, G. Wlodarczak, F. Clausset, P. I. Encrenaz, "Acetone in Interstellar Space". A&A.,180 (1987) L13.

[126] L. E. Snyder, F. J. Lovas, D. M. Mehnringer, N. Y. Miao, Y. J. Kuan. J. M. Hollis, P. R. Jewell, "Confirmation of Interstellar Acetone", ApJ, **578** (2002) 245.

[127] L. E. Snyder, J. M. Hollis, P. R. Jewell, F. J. Lovas, A. Remijan, "Confirmation of Interstellar Methylcyanodiacetylene (CH3C5N)", ApJ, **647** (2006) 412.

[128] A. Belloche, R. T. Garrod, H. S. P. Müller, K. M. Menten, C. Comito, P. Schilke, "Increased Complexity in Interstellar Chemistry: Detection and Chemical Modelling of Ethyl Formate and n-Propyl Cyanide in Sagittarius B2(N)", A&A., **499** (2009) 215

[129] B. Tercero, I. Kleiner, J. Cernicharo, H. V. L. Nguyen, A. López, G. M. Muñoz-Caro, "Discovery of Methyl Acetate and Gauche EthylFormate in Orion", ApJL., **770** (2013) 13.

[130] N. W. Broten, T. Oka, L. W. Avery, J. M. MacLeod, H. W. Kroto, "The Detection of HC₉N in Interstellar Space", ApJ **223** (1978) L105.

[131] A. J. Remijan, J. M. Hollis, L. E. Snyder, P. R. Jewell, F. J. Lovas, "Methyltriacetylene (CH₃C₆H) toward TMC-1: The Largest Detected Symmetric Top", ApJ **643** (2006) L37.

[132] Y. Zhou, D. H. Quan, X. Zhang, S. L. Qin, "Detection of hydroxyacetone in protostar IRAS 16293-2422 B", Res Astron Astrophys **20** (2020) 12.

[133] J. Cernicharo, C. Agúndez, C. Cabezas, B. Tercero, N. Marcelino, J. R. Pardo, P. de Vicente, "Pure hydrocarbon cycles in TMC-1: Discovery of ethynyl cyclopropenylidene, cyclopentadiene, and indene", A&A 649 (2021):L15.

[134] V. M. Rivilla et al., "Discovery in space of ethanolamine, the simplest phospholipid head group", Proc Nat Acad Sci **118** (2021) :e2101314118. https://doi.org/10.1073/pnas.2101314118

[135] M. C. McCarthy, K. L. K. Lee, R. A. Loomis, A. M. Burkhardt, C. N. Shingledecker, S. B. Charnley, M. A. Cordiner, E. Herbst, S. Kalenskii, E. R. Willis, C. Xue, A. J. Remijan, B. A. McGuire, "Interstellar detection of the highly polar five-membered ring cyanocyclopentadiene", Nature Astron 5 (2021)176.

[136] K. L. K. Lee, P. B. Changala, R. A. Loomis, A. B. Burkhardt, C. Xue, M. A. Cordiner, S. B. Charnley, M. C. McCarthy, B. A. McGuire, "Interstellar Detection of 2-cyanocyclopentadiene, C_5H_5CN , a Second Five-membered Ring toward", ApJL 910 (2021) L2.

[137] J. Cernicharo, A. M. Heras, A. G. G. M. Tielens, J. R. Pardo, F. Herpin, M. Guélin, L. B. F. M. Waters, "Infrared Space Observatory's Discovery of C_4H_2 , C_6H_2 , and Benzene in CRL 618", ApJ, **546** (2001) L123.

[138] A. Belloche, R. T. Garrod, H. S. P. Müller, K. M. Menten, C. Comito, P. Schilke, "Increased Complexity in Interstellar Chemistry: Detection and Chemical Modelling of Ethyl Formate and n-Propyl Cyanide in Sagittarius B2(N)", A&A., 499 (2009) 215

[139] G. W. Fuchs, U. Fuchs, T. F. Giesen, F. Wyrowski, "Trans-ethyl methyl ether in space: A new look at a complex molecule in selected hot core regions", A&A, 444 (2005) 521.

[140] A. Belloche, R. T. Garrod, H. S. P. Müller, K. M. Menten, "Detection of a branched alkyl molecule in the interstellar medium: iso-propyl cyanide", Science, 345 (2014) 1584.

[141] J. K. Anderson, L. M. Ziurys, "Detection of CCN (X2Πr) in IRC+10216: Constraining Carbon-chain Chemistry", ApJ **795** (2014) L1.

[142] M. Guélin, J. Cernicharo, C. Kahane, J. Gomez-Gonzales, "A New Free Radical in IRC +10216", A&A **157** (1986) L17.

[143] M. Guelin, R. Lucas, J. Cernicharo, "MgNC and the Carbon-Chain Radicals in IRC +10216", A&A, **280** (1993) 19.

[144] P. P. Tennekes, J. Harju, M. Juvela, L. V. Tóth, "HCN and HNC mapping of the protostellar core Chamaeleon-MMS1", A&A, **456** (2006) 1037

[145] W. M. Irvine, F. P. Schloerb, "Cyanide and Isocyanide Abundances in the Cold, Dark Cloud TMC-1", ApJ, **282** (1984) 516.

[146] T. Ogata, Y. Ohshima, Y., Endo, "Rotational Spectra and Structures of Carbon Monoxides C₅O, C₇O and C₉O", J. Am. Chem. Soc, **117**(1995) 3593.