The formation and evolution of Andromeda IX

Hedieh Abdollahi¹, Atefeh Javadi², Mohammad Taghi Mirtorabi¹, Elham Saremi^{2,3,4}, Habib Khosroshahi², Jacco Th. van Loon⁵, Iain McDonald^{6,7}, Elahe Khalouei², Sima T. Aghdam², and Maryam Saberi⁸

Physics Department, Faculty of Physics and Chemistry, Alzahra University, Vanak, 1993891176, Tehran, Iran

email: Abdollahi.hedieh2013@gmail.com

²School of Astronomy, Institute for Research in Fundamental Sciences (IPM), Tehran, 19568-36613, Iran

³Instituto de Astrofisica de Canarias, C/ Via Làctea s/n, 38205 La Laguna, Tenerife, Spain
⁴Departamento de Astrofisica, Universidad de La Laguna, 38205 La Laguna, Tenerife, Spain
⁵Lennard-Jones Laboratories, Keele University, ST5 5BG, UK

⁶Jodrell Bank Centre for Astrophysics, Alan Turing Building, University of Manchester, M13 9PL, UK

⁷Department of Physical Sciences, The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK

⁸Rosseland Centre for Solar Physics, University of Oslo, P.O. Box 1029, Blindern, NO-0315, Oslo, Norway

Abstract. Local Group (LG), the nearest and most complete galactic environment, provides valuable information on the formation and evolution of the Universe. Studying galaxies of different sizes, morphologies, and ages can provide this information. For this purpose, we chose the And IX dSph galaxy, which is one of the observational targets of the Isaac Newton Telescope (INT) survey. A total of 50 long-period variables (LPVs) were found in And IX in two filters, Sloan *i'* and Harris V at a half-light radius of 2.5 arcmin. The And IX's star formation history (SFH) was constructed with a maximum star formation rate (SFR) of about 0.00082 \pm 0.00031 M_{\odot} yr⁻¹, using LPVs as a tracer. The total mass return rate of LPVs was calculated based on the spectral energy distribution (SED) of about $2.4 \times 10^{-4} M_{\odot}$ yr⁻¹. The distance modulus of $24.56^{+0.05}_{-0.15}$ mag was estimated based on the tip of the red giant branch (TRGB).

Keywords. stars: AGB and LPV – stars: formation – stars: mass-loss– galaxies: evolution – galaxies: star formation – galaxies: individual: And IX

1. Introduction

Answering questions about the formation and evolution of the universe requires studying dwarf galaxies. While dwarf galaxies are the simplest system known, they are the most diverse in terms of star formation and chemical enrichment (Mateo 1998).

A comprehensive survey (up to now) in the optical bands has been launched to explore the evolution of dwarf galaxies using the Isaac Newton Telescope (INT) (Saremi et al. 2017, 2020). Probing long-period variable (LPV) stars, deriving the star formation history (SFH), and estimating the mass-loss of stars in galaxies are among the reasons for initiating the INT project. As a result of this survey, we can compare the SFHs of different galaxy types and study the evolution and quenching of dwarf galaxies (Hamedani Golshan et al. 2017; Saremi et al. 2019; Navabi et al. 2021). The method to reconstruct SFH was first applied by Javadi et al. (2011b) to rebuild the SFH of M 33.

Hedieh Abdollahi et al

The asymptotic giant branch (AGB) stars were selected because they are at their most luminous stage (~ $10^4 L_{\odot}$ (van Loon et al. 2005; Yuan et al. 2018)) and are easier to detect. AGBs are evolved stars and their luminosity is related to their birth mass (Rezaei et al. 2014; Hashemi et al. 2019). Moreover, their high mass-loss ($10^{-7} < \dot{M} \leq 10^{-3}$ $M_{\odot} \text{ yr}^{-1}$ (van Loon et al. 1999; Javadi et al. 2013, 2019; Boyer et al. 2017)) contributes to enhancing the ISM.

The spheroidal dwarf galaxy And IX, located ~ 39^{+5}_{-2} kpc (Weisz et al. 2019) along the major axis of M 31 galaxy, was selected as our candidate for study.

2. Photometry and detection of LPVs

2.1. Observations and photometry

Nine observations were made from 2015 - 2017 using the 2.5-m wide field camera (WFC) at INT in the Sloan i', Harris V, and RGO I filters. Data reduction is done using THELI (Transforming Heavenly Light into Image), an image processing pipeline adapted to multi-chip cameras. The DAOPHOT package performs the photometry procedures, and ADDSTAR task can evaluate the completeness of the photometry (Stetson 1987). Fully described details of the photometric procedure can be found in Saremi et al. (2020).

2.2. Detection of LPVs as SFH tracers

We detected LPVs using the method introduced by Stetson (1996). In this method, each star is assigned a variability index, which is a function of its magnitude and magnitude error. In each magnitude interval, a threshold for variability index is determined to separate variable stars from non-variables. More than 90% of the stellar population in this magnitude range should have variability indexes higher than the threshold.

To construct the SFH of And IX based on the LPVs, Milky Way foreground contamination should be excluded from candidates with an acceptable threshold. Omitting contamination was done by cross-correlating the INT catalog with the *Gaia* DR3 (Gaia Collaboration et al. 2021) and TRILEGAL simulation of Milky Way populations (Girardi et al. 2005).

We estimated the AGBs' birth mass through the birth mass-luminosity relation using PADOVA evolutionary tracks (Marigo et al. 2017). AGBs are at the end of their evolutionary path, and their luminosity is a function of their birth mass (Javadi et al. 2017). Age and pulsation duration are also calculated by mass-age and mass-pulsation relations (Javadi et al. 2011b; Saremi et al. 2021). The star formation rate (SFR) based on the mass, age, and pulsation duration over time represents the SFH. Fig. 1 presents SFH in two half-light radii (5 arcmin) of And IX in three metallicities. The highest peak of SFR reached at a level of $0.00082 \pm 0.00031 \text{ M}_{\odot} \text{ yr}^{-1}$ in 6 Gyr ago at Z = 0.0001. The metallicity of galaxies varies during their evolution, so in this study we consider metallicity Z = 0.0002 as well as a more metal-rich estimate Z = 0.0003 in addition to the main metallicity Z = 0.0001 (Collins et al. 2010; McConnachie et al. 2012; Kirby et al. 2013; Wojno et al. 2020).

3. Probing of dust in And IX

We modeled the spectral energy distribution (SED) by DUSTY code to estimate the mass-loss ratio of LPVs (Ivezi & Elitzur 1997). DUSTY assumes radiatively driven wind to solve hydrodynamic equations for AGB stars. Output parameters should be scaled for And IX as DUSTY solved equation by default parameters (gas-to-dust mass ratio $\psi_{\odot} = 200$, L = 10⁴ L_{\odot}, and $\rho_{dust} = 3$ g cm⁻³).



Figure 1. SFHs of And IX for metallicities of Z = 0.0001 (pink), Z = 0.0002 (orange), and Z = 0.0003 (blue) within two half-light radii (~ 0.022 deg²).

4. Results and conclusions

According to the different age gradients of the population in the inner and outer parts of the galaxy, the outside-in star formation scenario could be a galaxy evolution scenario.

The total mass return rate of our detected LPVs at Z = 0.0003, is estimated to be ~ $10^{-4} M_{\odot} \text{ yr}^{-1}$. By decreasing metallicity to Z = 0.0002, the total mass return increased by 50%. The total mass return rate is about $2.4 \times 10^{-4} M_{\odot} \text{ yr}^{-1}$ in And IX at Z = 0.0001. The average mass-loss rate by C-rich stars is more than 80% of the total mass return rate in three metallicities.

Following are papers that attempt to estimate the total mass return rate of the variables and the SFH observed by the INT survey (Saremi et al. 2017, 2020) for other dwarf galaxies.

References

Boyer et al. 2017, ApJ, 851, 152 Collins M. L. M., et al. 2010, MNRAS, 407, 2411 Gaia Collaboration et al., 2021, A&A...649A...1G Girardi, L.,Groenewegen, M. A. T.,Hatziminaoglou, E., da Costa, L., 2005, A&A, 436, 895G Hamedani Golshan R., Javadi A., van Loon J. Th., et al. 2017, MNRAS, 466, 1764 Hashemi S. A., Javadi A., van Loon J. Th., 2019, MNRAS, 483, 4751 Ivezi e., Elitzur M., 1997, Monthly Notices of the Royal Astronomical Society, 287, 799–811

- Javadi A., van Loon J. Th., Mirtorabi M. T., 2011b, MNRAS, 414, 3394
- Javadi A., van Loon J. Th., Khosroshahi H., Mirtorabi M. T., 2013, MNRAS, 432, 2824
- Javadi A., van Loon J. Th., Khosroshahi H., et al., 2017, MNRAS, 464, 2103
- Javadi A., van Loon J. Th., 2019, IAUS, 343, 283
- Kirby, E. N., Cohen, J. G., Guhathakurta, P., Cheng L., Bullock, J. S., Gallazzi, A., 2013, ApJ, 779, 102
- McConnachie, A. W., 2012, AJ, 144, 4M
- Marigo P., Girardi L., Bressan A., et al. 2017, ApJ, 835, 77
- Mateo, Mario L. 1998, ARA&A...36...435M
- Navabi M., Saremi E., Javadi A., et al. 2021, ApJ, 910, 127
- Rezaei Kh S., Javadi A., Khosroshahi H., van Loon J. Th., 2014, MNRAS, 445, 2214
- Saremi E., et al., 2017, Journal of Physics: Conference Series, 869, article id. 012068
- Saremi E., Javadi A., van Loon J. Th., Khosroshahi H. G., Rezaeikh S., Hamedani Golshan R., Hashemi S. A., 2019, Proceedings of IAU Symposium, 344, 125
- Saremi E., et al. 2020, ApJ, 894, 135S
- Saremi E., Javadi A., Navabi M., et al. 2021, ApJ, 923, 164
- Stetson P. B., 1987, PASP, 99, 191
- Stetson P. B., 1996, PASP, 108, 851
- van Loon J. Th., Groenewegen M. A. T., de Koter A., 1999, A&A, 351, 559
- van Loon J. Th., Cioni M.-R. L., Zijlstra A. A., Loup C., 2005, A&A, 438, 273
- Weisz D. R., et al. 2019, MNRAS, 489, 763W
- Wojno J., Gilbert K. M., Kirby E. N., Escala I., Beaton R. L., Tollerud E. J., Majewski S. R., Guhathakurta P., 2020, ApJ, 895, 78W
- Yuan W., et al. 2018, AJ, 156, 11