Do we need delocalised wavefunctions for the excited state dynamics of 1,1-difluoroethylene?

Sandra Gómez,*,^{†,‡,¶} Nadja K. Singer,[‡] Leticia González,[‡] and

Graham A. Worth[†]

[†]Department of Chemistry, University College London, 20 Gordon Street, London WC1H 0AJ, United Kingdom

‡Institute of Theoretical Chemistry, Faculty of Chemistry, University of Vienna, Währingerstrasse 17, 1090 Vienna, Austria

¶Current address: Departamento de Química Física, University of Salamanca, 37008,

Spain

E-mail: sandra.gomez@usal.es

Abstract

In this work we set up a model Hamiltonian to study the excited state quantum dynamics of 1,1-diffuoroethylene, a molecule that has equivalent atoms exchanged by a torsional symmetry operation leading to equivalent minima on the potential energy surface. In systems with many degrees of freedom where the minimum energy geometry is not unique, the ground state wavefunction will be delocalised among multiple minima. In this small test system, we probe the excited state dynamics considering localised (in a single minimum) and delocalised (spread over among multiple minima) wavefunctions and check whether this choice would influence the final outcome of the quantum dynamics calculations. Our molecular Hamiltonian comprises seven electronic states, including valence and Rydberg states, computed with the MS-CASPT2 method and projected onto the vibrational coordinates of the twelve normal modes of 1,1-difluoroethylene in its vibrational ground state. This Hamiltonian has been symmetrised along the torsional degree of freedom to make both minima completely equivalent and the model is supported by the excellent agreement with the experimental absorption spectrum. Quantum dynamics results show that the different initial conditions studied do not appreciably affect the excited state populations or the absorption spectrum when the dynamics is simulated assuming a delta pulse excitation.

keywords: quantum dynamics, excited states, 1,1-difluoroethylene, superpositions

1 Introduction

Symmetry is a powerful tool used to simplify problems and predict properties in a wide range of physical systems. In the context of quantum mechanics applied to molecules, symmetry operations must leave the molecular Hamiltonian invariant, i.e., it always belongs to the totally symmetric irreducible representation. For small molecules belonging to non-Abelian symmetry groups, the probability of finding the system in a particular conformation must be exactly the same as finding it in a conformation equivalent by symmetry. The nuclear wavefunction and potential energy surfaces must then be constructed taking the permutations of the symmetry group into account. The global delocalised wavefunction can be expressed as a superposition of localised wavefunctions at the single minima transformed via symmetry operations. These superpositions have been detected in the laboratory since the Young double slit experiment¹ and much later in buckyballs,² and in systems up to 800 atoms.³ As soon as the system moves away from this relaxed state (i.e., stops being an eigenstate), the expectation values of observables and the wavefunction itself change with time. A question would be whether parts of the superposition might interact differently, maybe due to their relative phase sign, when compared to wavefunctions that are localised in a single minima.

One interesting application where nuclear superpositions are invoked is the theoretical description of nuclear spin isomers (NSIs). The 1920s brought NSIs into the limelight, when the theory about the existence of two different species of the hydrogen molecule (ortho and para) was developed.^{4,5} However, even now the separation, identification, and conversion mechanisms between NSIs is a challenging problem and only a few polyatomic molecules (CH₃F,^{6–9} 13 CH₃F,^{10–12} H₂CO,¹³ 13 CCH₄,¹⁴ CH₃OH,¹⁵ H₂O,^{16–20} and C₂H₄^{21,22}) have been separated in their isomeric forms. In the last ten years theories connecting NSIs with superpositions of wavepackets have been developed.^{23–31} Using reduced dimensionality models, an association between the nuclear spin and an intermolecular torsion has been made^{23,24,27,32–35} and interference of torsional wavepackets within these reduced models leads to different excited state dynamics that can be associated to one or the other NSI, allowing their differentiation.^{25,26,36}

Even though the connection between nuclear spin isomers and rotational levels has been extensively reported, the relation to torsional superpositions (as part of the vibrational wavefunction) has been mostly studied from the theoretical point of view. In this work we will focus on torsional superpositions as initial conditions for the dynamics, without making further connections to NSIs. To the best of our knowledge, every study that has been done using torsional superpositions in the excited state has used reduced dimensionality models. This is the first time that a full dimensional system is considered, using the case of the 1,1-diffuoroethylene (1,1-DFE) molecule in the gas phase, which possesses twelve internal degrees of freedom.

In excited state dynamics, the choice of the initial conditions can critically affect the outcome of a simulation.^{37–39} For mixed quantum-classical trajectory methods, such as Tully surface hopping⁴⁰ or ab initio multiple spawning,⁴¹ in gas phase simulations geometries and velocities are usually selected from a Wigner distribution that mimics the lowest eigenstate of the ground state quantum harmonic oscillator. In direct dynamics variational multiconfigurational Gaussian (DD-vMCG)⁴² methods, a single multi-dimensional Gaussian function that is the solution for the ground state quantum harmonic oscillator carries the initial population. Grid methods like multiconfigurational time dependent Hartree (MCTDH)⁴³ are very flexible and almost any type of initial wavefunction can be formed and used in the propagation.

For this reason, here dynamics simulations were carried out using the MCTDH method,^{44,45} treating explicitly all degrees of freedom at a quantum level. Three different initial conditions were considered: two superpositions, one with a positive relative phase and one negative, and a wavefunction localised in a single minimum, i.e., without applying any symmetry operations on it. The deactivation of the population from the valence $\pi\pi$ * state towards the Rydberg state is found to be not particularly affected by the choice of the initial conditions, i.e., superpositions give the same outcome as starting the dynamics in a localised minima. If this behaviour can be extrapolated to larger systems with multiple minima, this result implies an enormous saving in computer time, since only a small part of the configuration space needs to be initially covered.

To describe the full-dimensional potential energy surfaces (PES) of 1,1-DFE, we use a vibronic coupling model⁴⁶ up to sixth order and fitted to high level ab-initio quantum chemistry calculations at different geometries. Since torsional angles of 90 degrees are needed to arrive at the conical intersections present in this system, the corresponding normal mode needs to be transformed to the actual dihedral angle, and high-order coupling terms (up to the sixth order) need to be included to describe the processes involving this motion with accuracy. In an earlier study on 1,1-DFE using multistate complete active space perturbation theory up to second order (MS-CASPT2) in MOLCAS, we found that a number of Rydberg states must be included to correctly describe the non-adiabatic dynamics starting from the bright valence state.⁴⁷

To prove that our Hamiltonian is indeed a good model, we calculate the absorption spectrum after excitation to the excited-state manifold. The excellent agreement between theoretical and experimental spectrum supports our choice of a Hamiltonian and allows the assignment of the spectral bands. Using a localised initial condition and starting the excited state dynamics from the zwitterionic π^{*2} and Rydberg π -3s, π -3p_x, π -3p_y, and π -3p_z states, a reduced dimensionality model then allows us to theoretically unravel the origin of the spectral progression of bands detected in the experimental absorption spectrum of 1,1-DFE.⁴⁸

2 Methods

2.1 Construction of the nuclear Hamiltonian

To analytically describe the PES, we used a vibronic coupling Hamiltonian.^{46,49} This method, diabatic by ansatz, expresses the PES as a Taylor expansion around a given point, normally the optimised geometry (Frank-Condon point). The Hamiltonian \boldsymbol{H} can be thus defined as

$$\boldsymbol{H} = H^{(0)} \boldsymbol{1} + \boldsymbol{W}^{(0)} + \boldsymbol{W}^{(1)} + \boldsymbol{W}^{(2)} + \boldsymbol{W}^{(3)} + \dots,$$
(1)

where each matrix has a dimension of 7, the number of electronic states involved. As a coordinate basis to express the Hamiltonian, we choose the mass-frequency scaled ground-state normal vibrations, which in the case of 1,1-DFE means twelve normal modes. The zero-order Hamiltonian $H^{(0)}$ corresponds to the kinetic energy operator. For the normal

modes q_1 - q_3 and q_5 - q_{12} , the kinetic energy operator \hat{T}_q reads

$$\hat{T}_q = -\sum_{\alpha=1}^{12'} \frac{\omega_\alpha}{2} \frac{\partial^2}{\partial Q_\alpha^2},\tag{2}$$

where ω_{α} is the ground-state normal mode frequency and the prime on the summation indicates $\alpha = 4$ is omitted.

The normal mode q_4 is replaced by the torsion angle θ between the CH₂ group and the molecular plane, considering the fluoride atoms to be fixed due to the large difference in atomic masses. Since a torsional degree of freedom implies a complete rotation along the symmetry axis whereas the normal mode is a harmonic back and forth motion, we transformed the kinetic energy operator following reference 50. The kinetic energy operator \hat{T} for this degree of freedom now reads

$$\hat{T}_{\theta} = -\frac{1}{2\mu r^2} \frac{\partial^2}{\partial \theta^2},\tag{3}$$

where μr^2 corresponds to the moment of inertia with a mass $\mu = 2$ amu, and r is the groundstate equilibrium distance from the hydrogen atoms to the inertial axis (along the C=C bond), which is 0.974 Å. The total zero-order Hamiltonian, $H^{(0)}$, is then described as

$$H^{(0)}(Q) = \hat{T}_{\theta} + \hat{T}_{q},$$
(4)

the sum of both terms.

The matrices **W** form the diabatic potentials and couplings. Following the standard vibronic coupling model scheme, for modes that are close to harmonic (q₁, q₃, q₅, q₇, q₈, q₉ and q₁₂), the zeroth order potential energy matrix $W_{ij}^{(0)}$ is diagonal and formed by the ground-state harmonic oscillator potential shifted by the vertical energies:

$$W_{ij}^{(0)}(Q) = (E_i + V)\,\delta_{ij} = E_i + \sum_{\alpha} \frac{1}{2}\omega_{\alpha}Q_{\alpha}^2 \quad ; \alpha = 1, 3, 5, 7, 8, 9, 12.$$
(5)

Higher order terms then enter the appropriate higher order diabatic potential matrices.

Going back to equation 1, the order of the expansion gives the name for the model: linear vibronic coupling model for expansions truncated at $W^{(1)}$, quadratic vibronic coupling model for truncations at $W^{(2)}$, quartic vibronic coupling model for truncations at $W^{(4)}$, etc. These terms are non-diagonal matrices and are responsible for the coupling, mediated by vibrations, between pairs of electronic states, hence the name of vibronic coupling model. In the 1,1-DFE model, we included expansion terms up to sixth order. The terms up to second order are

$$W_{ii}^{(1)}(Q) = \sum_{\alpha} \kappa_{\alpha}^{(i)} Q_{\alpha}$$

$$W_{ij}^{(1)}(Q) = \sum_{\alpha} \lambda_{\alpha}^{(i,j)} Q_{\alpha}, \quad i \neq j$$

$$W_{ii}^{(2)}(Q) = \sum_{\alpha,\beta} \frac{1}{2} \gamma_{\alpha,\beta}^{(i)} Q_{\alpha} Q_{\beta}$$

$$W_{ij}^{(2)}(Q) = \sum_{\alpha,\beta} \frac{1}{2} \mu_{\alpha,\beta}^{(i,j)} Q_{\alpha} Q_{\beta}, \quad i \neq j$$
(6)

showing how the dependence with the degrees of freedom is linear for the first term, quadratic for the second, etc. Higher-order terms were also required for the diabatic potentials of some modes, in particular for the on-diagonal (intra-state) mode coupling, and are written

$$W_{ij}^{(m)}(Q_{\alpha}) = \frac{1}{m!} C_{\alpha}^{m,(i,j)} Q_{\alpha}^{m}$$
$$W_{ij}^{(m+n)}(Q) = \frac{2}{(m+n)!} C_{\alpha,\beta}^{m,n,(i,j)} Q_{\alpha}^{m} Q_{\beta}^{n}.$$
(7)

Since 1,1-DFE belongs to the C2v symmetry point group we can apply symmetry to neglect the terms that would make the molecular Hamiltonian antisymmetric. Since the Hamiltonian must be invariant under the symmetry operations of the C_{2v} symmetry point group, to which 1,1-DFE belongs, some of the Hamiltonian terms are zero by definition. The terms up to second order which are nonzero obey the following product rules for the irreps of the vibrations, $\Gamma_{\alpha}, \Gamma_{\beta}$, and the irreps of the electronic wavefunctions, Γ_i, Γ_j

$$\kappa_{\alpha}^{(i)} \neq 0: \ \Gamma_{\alpha} \supset \Gamma_{A}$$

$$\lambda_{\alpha}^{(i,j)} \neq 0: \ \Gamma_{\alpha} \otimes \Gamma_{i} \otimes \Gamma_{j} \supset \Gamma_{A}$$

$$\gamma_{\alpha,\beta}^{(i)} \neq 0: \ \Gamma_{\alpha} \otimes \Gamma_{\beta} \supset \Gamma_{A}$$

$$\mu_{\alpha,\beta}^{(i,j)} \neq 0: \ \Gamma_{\alpha} \otimes \Gamma_{\beta} \otimes \Gamma_{i} \otimes \Gamma_{j} \supset \Gamma_{A}.$$
(8)

This means that the excited state gradients, κ , are only non-zero for totally symmetric modes, while the linear vibronic coupling between states, λ , is due to modes with the correct symmetry that when multiplied by the symmetry of the pair of states leads to a totally symmetric irreducible representation. The diagonal γ terms will be non-zero if the direct product of both modes involved is totally symmetric, whereas for the off-diagonal quadratic coupling terms, μ , the product must also include the symmetry of the electronic states. Symmetry rules for the higher order terms follow as straighforward extensions of these.

Some modes with strong anharmonicity require diabatic potential functions in place of the Taylor expansions described above in order to go to higher orders without drastically increasing the number of parameters to fit. The off-diagonal coupling, however, always follows the scheme above. These anharmonic functions were chosen purely on how well they fitted the ab initio data. Modes q_2 and q_{11} are described by Morse potentials:

$$W_{Mor,ii}(Q_{\alpha}) = D_0^{(i)} \left\{ 1 - \exp\left[-a_{\alpha}^{(i)} \left(Q_{\alpha} - Q_{0,\alpha}^{(i)} \right) \right] \right\}^2 + \epsilon_{\alpha}^{(i)}$$
(9)

and modes \mathbf{q}_6 and \mathbf{q}_{10} used a Lennard-Jones form

$$W_{LJ,ii}(Q_{\alpha}) = -D_0^{(i)} \left[\left(\frac{(\alpha_{\alpha}^{(i)})^{12}}{Q_{\alpha} - Q_{0,\alpha}^{(i)}} \right)^{12} - \left(\frac{2(\alpha_{\alpha}^{(i)})^6}{Q_{\alpha} - Q_{0,\alpha}^{(i)}} \right)^6 \right] + \epsilon_{\alpha}^{(i)}.$$
(10)

The torsional q_4 mode, as explained above, is replaced by an angle, θ . Potentials for this

mode are periodic and were fitted as a function of the cosine of the torsion angle:

$$W_{ii}(\theta) = \kappa_4^{(i)} \cos(\theta) + \frac{1}{2} \gamma_4^{(i)} \cos^2(\theta), \qquad (11)$$

where $\kappa_4^{(i)}$ and $\gamma_4^{(i)}$ are fitting constants. Any diabatic coupling and higher order terms coupling q_4 to the other modes used expressions of the form in Eqs. (6) and (7), using $\cos(\theta)$ in place of Q, and taking into account that this function is symmetric at $\theta = 0$ and anti-symmetric at $\theta = \frac{\pi}{2}$.

The parameters for the functions were obtained using the VCHam tool in the Quantics suite⁵¹ to calculated ab-initio data. The 1,1-DFE normal modes were obtained at the planar D_{2h} optimised ground state geometry at the MP2/aug-cc-pVDZ level of theory using the Gaussian 09 program.⁵² The normal modes and the corresponding symmetry labels using the C_{2v} point group are depicted in Figure 1. We use C_{2v} labels as this point group is retained throughout rotation around the torsional axis. A comparison between the optimised MP2 geometrical parameters and the experimental values is shown in Table 1. A table with frequencies and description of the motions can be found in the Supporting Information in SI-Table 1.

Table 1: Bond lengths (in Ångstrom) and angles (in degrees) for the geometries obtained bu optimisation of the ground state of 1,1-DFE employing MP2/aug-cc-pVDZ in comparison with experimental data from Ref. 53.

	Exp^{53}	MP2
r(CC)	1.315	1.335
r(CF)	1.323	1.337
r(CH)	1.078	1.087
$\not\preccurlyeq$ (FCF)	109.1	109.6
\neq (HCH)	121.8	121.5
$\not\prec$ (CCF)	125.5	125.2
∢(HCC)	119.1	119.3

The electronic state energies of the lowest seven singlet states at torsion angles of 0° and 90° calculated at the SA(11)-MS-CAS(2,6)PT2/aug-cc-pvDZ level of theory are listed



Figure 1: Normal mode coordinate basis with symmetry labels belonging to the C_{2v} point group. The torsional normal mode q_4 is replaced by the full torsion θ in the model Hamiltonian.

in Table 2. The active space includes the Rydberg orbitals 3s, $3p_x$, $3p_y$, and $3p_z$ and the π and π^* orbitals. The state-average procedure is performed over eleven roots to account for every single and double excitation related to the chosen active space (as explained in Ref. 47), even if here we were only interested in the dynamics in the lowest seven states. A picture of the active space and further details about the electronic structure can be found in Ref. 47.

Table 2: Vertical excitations (ΔE , in eV) of 1,1-DFE at the SA(11)-MS-CAS(2,6)PT2/augcc-pvDZ level of theory. Three electronic states are labelled following the Mulliken notation for ethylene,⁵⁴ N for the closed shell π^2 , V for the $\pi\pi^*$ and Z for the π^{*2} state. The energies are calculated at the optimised geometry (FC, for Frank Condon) and at 90° of torsion. The experimental (exp) values are taken from Ref. 48.

State	ΔE at FC	\exp^{48}	symmetry	ΔE at 90°
Ν	0.00		A_1	3.30
V	7.45	7.49	A_1	3.47
π -3s	7.07	7.06	B_2	7.91
π -3p _x	8.10		A_2	9.03
π -3p _y	8.20		B_1	9.26
π -3p _z	8.93	8.97	A_1	10.03
Ζ	13.77		A_1	9.27

The full dimensional potential energy surface consists of 12 nuclear degrees of freedom and 7 electronic states: N, V, Z, π -3s, π -3p_x, π -3p_y, and π -3p_z. N stands for neutral, V for valence and Z for zwitterionic using the Mulliken notation for ethylene.⁵⁴ These labels represent the character of the wavefunction at the Franck-Condon point and are thus used to label the diabatic states of our model. Since we need two minima equivalent by torsion to construct the initial wavefunction as a torsional superposition, we transformed our potentials using an orbital diabatic basis. In this basis, the N and V potentials cross and interchange character (see Figure 2b). In this new basis, we rename the states as π_1 and π_2 , since they are now completely equivalent. The rest of the states mentioned above keep their original character and thus their labels do not change. The 7 diabatic potential energy curves along the torsion were fitted to ab-initio calculations using the SA(11)-MS-CAS(2,6)PT2/aug-ccpVDZ level of theory performed at different geometries. Adiabatic curves and single ab-initio points are plotted in Figure 2a. The corresponding diabats are plotted in Figure 2b. The potentials along the original normal mode q_4 can be seen in Figure 2c showing the need of transforming the torsional coordinate in order to describe the crossing to the ground state. The two lowest states cross at 90 degree of torsion.

To provide data for the fitting, calculations at the FC geometry and at 90 degree of torsion were performed along every normal mode providing 3234 data points, corresponding to 462 single point electronic structure calculations. The model has 553 parameters to fit which are not independent and so the fitting was done in a series of steps to allow control of the procedure. The diabatic potentials for the torsional degree of freedom were fitted first, using the planar geometry at 0 degree as the initial point for the Taylor expansion. Next, for every normal mode a 1D fit to the ab-initio data was performed, starting the expansion at 90 degrees of torsion. Explicit couplings to the torsional degree of freedom were then included to force the two minima at planar geometries (0 and 180 degree) to be equivalent. For every 1D fit, the configuration interaction vector of the electronic structure calculation was monitored, to ensure that the diabats correctly follow their electronic character. For the states π_1 and π_2 , however, the same diabatic functions must be used to impose symmetry, which caused some deviations between the ab-initio data and the fitted curves.



Figure 2: Potential energy curves as a function of the torsional angle θ of 1,1-DFE with all other coordinates at the equilibrium geometry. Fit made using the VCHam tool in the Quantics⁵¹ suite to ab-initio points calculated with SA(11)-MS-CAS(2,6)PT2/aug-cc-pvDZ. a) Adiabatic curves and ab-initio points. b) Diabatic curves. The orbital-diabatic potentials π_1 and π_2 (purple and blue respectively) cross at $\pi/2$ radians and have two equivalent minima at 0 and π radians, each one from a different diabat. c) Ab-initio calculated energy points as a function of the torsional normal mode q_4 .

After the initial guesses for every diabatic 1D potential were found, a global fit was performed including couplings up to sixth order for specific coordinates (especially couplings that involved the torsional motion). The total standard deviation with respect to the abinitio points was 0.257 eV, with the largest errors being associated with high energy points that are not well fitted by the model potentials and do not play a role in the dynamics. The 1D adiabatic potential energy curves and the corresponding ab-initio points for three key modes (q_3 , q_6 and q_{10}) are plotted in Figure 3. Cuts along the other degrees of freedom can be found in the Supporting Information. The middle column shows the potential curves at 90 degree of torsion, where the fit was performed. Note that to its right and its left, at both 0 and 180 degrees of torsion, the potentials are equivalent, conserving the symmetry. However, these two minima correspond to different diabatic potentials (π_1 and π_2) which switch energetic order at 90 degree. To preserve the symmetry, these diabatic potentials thus need to be fitted with the same analytical functions and parameters for each normal mode.

All parameters obtained for the model Hamiltonian are listed in the Supporting Information, along with cuts like that in Fig. 3 for all modes showing the quality of the fit.

2.2 Initial conditions

Calculations were performed on different size systems (2D, 5D and 12D). The initial wavefunction in each calculation was constructed as the Hartree product of the harmonic oscillator ground state for every normal mode and the torsional ground state wavefunction in the π_1 minimum. This was followed by propagation in imaginary time (energy relaxation)⁵⁵ to get the ground-state vibrational eigenfunction. The barrier between the minima is high enough that on the timescale of the relaxation (100 fs) there is no transfer of population to the π_2 minimum. Three different initial conditions for the torsional wavefunction were then constructed from this localised wavefunction obtained by relaxation; taken directly as the eigenstate of the π_1 and π_2 diabatic potential energy curves. A scheme can be seen in Figure 4.

This multidimensional initial wavefunctions created in this way for one or two torsional minima corresponding to states π_1 and π_2 was consequently stored, and read in to be further electronically excited to the π_2 and π_1 states, respectively. This way, either a localised torsional wavefunction or a superposition of them was created in the S₂ adiabatic state (blue state in Figure 2a). In the case of the localised initial condition, the wavefunction carried the whole population, while for the superposition each half was given an amplitude of $1/\sqrt{2}$.



Figure 3: Adiabatic potential energy curves along key normal mode coordinates of 1,1-DFE calculated with SA(11)-MS-CAS(2,6)PT2/aug-cc-pvDZ. The right and left columns correspond to torsional angles of π (180 degree) and 0 degrees, respectively. The middle column corresponds to a torsional angle of $\pi/2$ (90 degree), where the fit was performed. All other coordinates are at equilibrium values. The labels q_i correspond to the normal mode coordinate along which the ab-initio energies were calculated and fitted. The corresponding normal modes can be seen in Figure 1.



Figure 4: Scheme depicting the initial conditions used in this work. Either the symmetric (blue), the antisymmetric (orange) or the localised (just the eigenstate of the left side minima with positive amplitude) were vertically excited to initiate the dynamics.

3 Results

3.1 Dynamics started on the $\pi\pi^*$ valence state

Quantum dynamics simulations of 1,1,-DFE were performed for symmetric and antisymmetric superpositions as well as for a system localised in one single minima.

For each case, a propagation using the MCTDH method was performed for 150 fs, using the multiset formalism, i.e., different single particle functions (SPFs) are optimised for each electronic state. The mean fields were calculated at each integration step and integrated with the Adams-Bashforth-Moulton⁵⁶ predictor-corrector method. The torsional coordinate was described using a Fast Fourier Transform (FFT) collocation grid, while the other degrees of freedom used harmonic oscillator discrete variable representations (DVRs) as primitive basis functions.

Since convergence of the MCTDH method depends on how the modes are combined within a SPF, on the number of them and on the primitive grid size, we first started converging calculations on a reduced system, to steadily increase the number of degrees of freedom. We performed calculations with two (torsion θ and CC stretching q_{10}), five (torsion θ , CC stretching q_{10} , CF₂ pyramidalisation q_3 , CF symmetric stretch q_6 and a combination of asymetric CF₂ stretch and CH₂ bend, q_7) and twelve degrees of freedom, i.e., full dimensionality (the motions are depicted in Figure 1). Calculations for the three different initial conditions in all sized systems (2, 5, or 12 degrees of freedom) used the same primitive grids, but were converged individually with respect to the SPF basis. The number of primitive basis functions, along with mode combinations and the number of SPFs per state for the largest calculations performed is found in Table 3.

The dynamics including only two degrees of freedom, q_{10} and θ , shows no interchange of population between the π_1 - π_2 states and the π - $3p_x$ state during 150 fs (Figure 5a), indicating that the CC double bond vibration is not enough to drive the system away from the Frank-Condon region or excite the torsional motion to reach the conical intersections. This agrees with previous work on ethene which showed that for population transfer to occur the pyramidalisation motion also needs to be included ⁵⁹ The populations of the π_1 and π_2 states have been summed up in the superposition calculations to allow comparison with the localised initial condition that started the dynamics with 100% of its population in π_2 . The differences of using or not using a superposition as initial condition are unnoticeable. As we can see in Figure 5b, at least five modes are needed to drive the dynamics, which are qualitatively similar to the full dimensionality result shown in Figure 5c. The population, initially excited to the V state, is very quickly transferred to the π -3s and π -3p_x Rydberg states. This transfer is initially mediated by q_7 , a combination of asymmetric CF_2 stretch and CH_2 bend, and the torsional degree of freedom q_4 with the same symmetry as the electronic states and it is driven by the gradient created by the CF symmetric (q_6) and the CC double bond (q₁₀) stretches. In the 5D and 12D cases, the π -3p_x state traps the population during the first 30 fs, after which it slowly relaxes to the π_1 - π_2 states and the π -3s. In the 12D case (Figure 5c), the population transferred to the π -3s state is larger than in the reduced dimensionality system, indicating that other degrees of freedom play a role in the excited state dynamics. When comparing the localised and delocalised initial condition 12D

Table 3: Basis used for the largest MCTDH calculations starting with localised and delocalised (symmetric and antisymmetric) wavefunctions on the $\pi\pi^*$ state. The primitive basis was the harmonic oscillator DVR⁵⁷ for every mode except the torsion, which used a periodic FFT grid.⁵⁸ N_i is the number of primitive basis functions for each degree of freedom in the particles, while n_i is the number of SPFs used for each of the seven electronic states. Config is the total number of configurations in the MCTDH wavefunction.

Type of WF	particle	N_i	n_i	Config.
2mode				
local	el	7		7
	$ heta, q_{10}$	$100,\!55$	$(1,\!1,\!1,\!1,\!1,\!1,\!1)$	
sym / anti	el	7		7
	$ heta, q_{10}$	$100,\!55$	$(1,\!1,\!1,\!1,\!1,\!1,\!1)$	
5mode				
local	el	7		35278
	$ heta, ext{q}_7$	100,27	$(10,\!29,\!26,\!8,\!3,\!3,\!4)$	
	q ₁₀	55	(16, 20, 20, 7, 3, 3, 4)	
	q_3,q_6	$31,\!27$	$(10,\!30,\!20,\!10,\!3,\!3,\!4)$	
sym / anti	el	7		54269
	$ heta, ext{q}_7$	$100,\!27$	(23, 32, 32, 10, 4, 3, 4)	
	q_{10}	55	$(19,\!23,\!28,\!8,\!3,\!3,\!4)$	
	q_3,q_6	$31,\!27$	$(10,\!30,\!30,\!10,\!4,\!3,\!4)$	
12mode				
local	el	7		229112
	$ heta, q_{10}$	100,55	(17, 27, 37, 11, 4, 3, 3)	
	q_1, q_2, q_5, q_8	11	$(5,\!9,\!10,\!6,\!3,\!3,\!3)$	
	$\mathbf{q}_3, \mathbf{q}_6, \mathbf{q}_7$	$31,\!27,\!27$	$(17,\!29,\!30,\!9,\!4,\!3,\!4)$	
	q_9, q_{11}, q_{12}	11	(10, 11, 12, 6, 4, 3, 3)	
sym / anti	el	7		619117
	$ heta, q_{10}$	$100,\!55$	$(30,\!40,\!40,\!20,\!4,\!3,\!4)$	
	$\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_5, \mathbf{q}_8$	11	(11, 10, 12, 9, 3, 3, 3)	
	$\mathbf{q}_3, \mathbf{q}_6, \mathbf{q}_7$	$31,\!27,\!27$	(30, 40, 40, 20, 4, 3, 4)	
	q_9, q_{11}, q_{12}	11	(13, 13, 13, 9, 4, 3, 3)	

runs, we can observe small differences in the populations, probably due to the difficulty of converging the SPF basis in the larger systems.



Figure 5: Diabatic time-dependent electronic state populations for calculations including (a) two nuclear degrees of freedom: torsion and CC stretch, (b) five nuclear degrees of freedom: torsion, CC stretch and scissoring, and pyramidalisation of CH_2 and CF_2 fragments, (c) twelve degrees of freedom, i.e., full dimensionality. In solid lines the population after exciting a localised initial wavefunction (local) is depicted. In dashed and in dotted lines the populations after exciting the symmetric superposition (sym) and the antisymmetric (anti), respectively, are shown.

During the dynamics, the autocorrelation function (overlap of wavefunctions at time = t and time = 0) was stored every 0.5 fs and later Fourier transformed to obtain the absorption spectrum. In Figure 6, the spectrum for the localised initial condition (local) and superpositions with positive (sym) and negative (anti) relative phase are shown for the full system with twelve degrees of freedom. The minimal differences that can be observed are again probably due to lack of basis set convergence. The main spacing is due to the CC stretching mode (q_{10}) with a harmonic frequency of 0.2192 eV and the shoulder peaks next to the main



Figure 6: Absorption spectra calculated as the Fourier transform of the autocorrelation function from full dimensionality dynamics with a damping time (phenomenological broadening) of $\tau = 100$ fs comparing an initial condition localised in one minimum (local) a symmetric superposition (sym) and an antisymmetric superposition (anti) started on the V state.

progression are due to the CF_2 symmetric stretch (q₆, harmonic frequency 0.1132 eV). By comparing to the spectra from reduced dimensionality calculations, the overall width and shape comes from the torsional degree of freedom and the CF symmetric stretches which couple the electronic states, driving the initial wavepacket away from the Frank-Condon region.

In contrast to a linear vibronic coupling model, our high-order Hamiltonian makes the assignment of which modes couple which states and drive the dynamics impossible after a few femtoseconds. To aid in this task, we performed excited state dynamics based on the 5D propagation shown in Figure 5b, but using only the localised initial condition and removing one degree of freedom each time. The populations and the absorption spectra are depicted in Figure 7. These simulations show clearly that the torsion and the CF symmetric and CC stretches are crucial to obtain population transfer to any state on the ultrafast timescale, since removing any of these DoFs inhibits the deactivation of the bright state. The CF₂ pyramidalization is needed to allow a transfer to the π -3s state. Although q₇, a combination of asymmetric CF₂ stretch and CH₂ bend, was initially thought to couple the valence and the π -3s state for symmetry reasons, its presence does not affect the outcome

of the dynamics. The spectra shown in Figure 7e supports the assumption that the torsion dominates the spectrum, since without it we obtain a vibrational progression where the maximum is the 0-0 band. The CF symmetric stretch is responsible for the shoulder peaks and the pyramidalization narrows down the general shape of the spectra when allowing the transfer to the π -3s state.



Figure 7: Simulations corresponding to the five mode dynamics in figure 5 with a localised initial condition, after removing one degree of freedom. a) Diabatic time-dependent populations after removing the torsional θ mode (solid line) and after removing the CF symmetric stretch q₆ (dashed), b) without including the CF₂ pyramidalisation q₃, c) ignoring the CC stretch mode q₁₀ and d) ignoring q₇, a combination of asymmetric CF₂ stretch and CH₂ bend. e) Absorption spectra computed as the Fourier transform of the autocorrelation function for each of these cases.

In our previous work on 1,1-DFE using on-the-fly dynamics,⁴⁷ the π -3s state acted as a doorway state, keeping population from the V state trapped for a while before transferring it back. This process was essential for the deactivation of the molecule to its electronic ground state. Using our parametrised potentials and quantum dynamics, we do not see any trapping of the population on this Rydberg state. Instead, its population keeps growing and after 150 fs it is the most populated state. To distinguish between the π_1 - π_2 state populations we calculated the adiabatic populations for the 5D localised system integrating the transformed wavefunction over the primitive grids. The electronic ground state population was found to never exceed the value of 0.6%. Therefore, the outcome of the dynamics is substantially different in the two cases and deserves a closer look why this could be.

A quantum dynamics method (like MCTDH) is fully quantum and thus expected to provide a better description of the nuclear motion than that obtained by propagating independent classical trajectories, as done in surface hopping. An important difference in the two calculations is that in surface hopping, the trajectories were selected from a Wigner distribution, which is a quasiprobability function that provides an approximation for the lowest eigenstate of the quantum harmonic oscillator; instead, the MCTDH simulations use the actual lowest energy "localised" eigenfunction of the Hamiltonian for the initial wavefunction. However, to use the MCTDH method, one must pre-compute the potential energy surfaces in advance and fit them to an analytical potential form on a base of nuclear coordinates. This procedure potentially restricts the system as the choice of coordinates and the fitted function may not allow the system to evolve in the same way as with on-the-fly dynamical simulations. As an example, one could think of the different MCTDH dynamics that would result if we had used the torsional normal mode in the model potential and not substituted it by the full dihedral angle, as depicted in Figure 2. While the quality of the potential used in the MCTDH simulations is supported by the calculated spectrum, the latter only samples the potential energy surfaces around the Frank-Condon point and does not say anything about the surfaces in general. Furthermore, due to its construction the vibronic coupling model does not include any coupling between modes that requires high-order polynomials.

Further work is thus needed to see which of the two sets of dynamical simulations is correct. A possible protocol would be to run on-the-fly quantum dynamics with the DDvMCG method,^{60,61} but this would require calculating frequencies, gradients, couplings and energies at the CASPT2 level for seven states and each pair of states, which would be computationally very expensive.

3.2 Localised dynamics started on the Rydberg states

In order to extend our excited state dynamics study beyond the the valence $\pi\pi^*$ state, we investigate the dynamics of the system including the manifold of electronic states that cross, couple and affect the behaviour after light irradiation. In Figure 8a, the transition dipole matrix elements that couple the ground state and excited state electronic wavefunctions are plotted at different torsional angles. The values, calculated at the same level of theory as the potentials, refer to the adiabatic electronic states. However, at the ground state optimised geometry the V and π -3s states switch character (see for example Figure 2) and we accommodated that change switching the values of the matrix elements μ_{12} and μ_{13} at the planar geometry. Since at the Frank-Condon region the transition dipole moment only has negligible values for the coupling with the π -3p_x state, we started 5D dynamics using the localised initial condition on every other electronic state and calculated the absorption spectra from the autocorrelation function (Figure 8b). Every spectrum has been weighted by the corresponding matrix element of the transition dipole moment at the FC point. The spectrum starting in the V state has been computed from dynamics in full dimensionality (it is the same as Figure 6) and therefore has been shifted in energy by the difference in the zero point energies due to the extra degrees of freedom. In Ref. 48, the first band is concluded to be a mixture of the Rydberg π -3s and the valence $\pi\pi$ * states. The vibrational progression observed at higher energies starts with peaks from the π -3p_y state, followed by the π -3p_z and probably higher lying Rydberg states. The Z state modulates the intensity of these peaks with a broad gaussian-like shape.

4 Conclusions

We set out to investigate the importance of superpositions due to symmetrically equivalent minima to describe the initial wavepacket in photo-excited quantum dynamics simulations. The answer is found to be in the negative and no significant difference was seen in the



Figure 8: a) Elements of the adiabatic state transition dipole moment matrix along the torsional degree of freedom calculated with MS-CASPT2. The sub-indices reflect the pair of electronic states involved in the transition. b) Absorption spectra calculated as the Fourier transform of the autocorrelation function from dynamics including five degrees of freedom (except for the V case, where the full dimensionality result was taken), as specified in Table 4 with a damping of $\tau = 50$ fs. Separate calculations were made used a localised initial wavefunction started on the valence, zwitterionic and Rydberg electronic states. The amplitude has been weighted by the transition dipole moments. The experimental spectrum shown in grey was taken from Ref. 48.

Table 4: Basis used for the MCTDH simulations starting with a localised wavepacket on the Rydberg and zwitterionic states. The primitive basis was the harmonic oscillator DVR for every mode except the torsion, which uses a periodic FFT grid. N_i is the number of primitive basis per dimension in every particle, while n_i is the number of SPFs used per electronic state. Config is the total number of configurations in the MCTDH wavefunction.

Initial state	particle	N_i	n_i	Config.
π -3s	el		7	7130
	$ heta, q_7$	$100,\!27$	$(5,\!12,\!15,\!3,\!3,\!3,\!3)$	
	q ₁₀	55	(10, 14, 12, 4, 4, 3, 3)	
	$\mathrm{q}_3,\mathrm{q}_6$	$10,\!27$	$(7,\!20,\!20,\!4,\!4,\!3,\!3)$	
π -3p _x	el		7	46519
	$ heta, q_7$	$100,\!27$	$(12,\!32,\!32,\!10,\!4,\!3,\!3)$	
	q ₁₀	55	$(19,\!22,\!23,\!9,\!4,\!3,\!4)$	
	q_3, q_6	$10,\!27$	$(10,\!30,\!30,\!10,\!4,\!3,\!4)$	
π -3p _y	el		7	45105
	$ heta, q_7$	$100,\!27$	$(9,\!32,\!31,\!10,\!4,\!3,\!4)$	
	q ₁₀	55	(16, 22, 23, 10, 4, 3, 4)	
	q_3, q_6	$10,\!27$	$(10,\!30,\!30,\!10,\!4,\!3,\!4)$	
π -3p _z	el		7	26490
	$ heta, q_7$	$100,\!27$	$(7,\!13,\!28,\!10,\!4,\!4,\!3)$	
	q ₁₀	55	(11, 19, 25, 10, 4, 3, 4)	
	q_3, q_6	$10,\!27$	$(10,\!23,\!27,\!10,\!4,\!3,\!4)$	
Z	el		7	68355
	$ heta, q_7$	$100,\!27$	(24, 32, 32, 10, 4, 3, 4)	
	q ₁₀	55	(24, 32, 32, 10, 4, 3, 4)	
	q_3, q_6	$10,\!27$	$(10,\!30,\!30,\!10,\!4,\!3,\!4)$	

excited state quantum dynamics of the 1,1-diffuoroethylene molecular system using either localised (single minimum) or delocalised (symmetry-adapted) initial conditions. We observe that a qualitative correct behaviour of the deactivation of the V state is achieved with five degrees of freedom from the total of twelve. The torsion, CC and CF symmetric stretches, combination of asymmetric CF_2 stretch and CH_2 bend, and the CF_2 pyramidalisation drive the excited state dynamics on this system and explain the peaks observed in the absorption spectra after Fourier transforming the autocorrelation function.

In contrast to previous SH dynamics,⁴⁷ the current simulations find that there is barely population transfer to the electronic ground state, flowing into the Rydberg π -3s instead. This outcome is consistent for all three initial conditions investigated, illustrating that using a localised nuclear wavefunction is perfectly reasonable and the relative phases on the parts of the symmetric and antisymmetric superpositions do not affect the dynamics, leading to numerically identical results. There are minor differences for the full-dimensional system when starting with a localised wavefunction which maybe is due to convergence. As a way of validating our parameterised Hamiltonian, we performed dynamics using a localised initial condition on the valence, zwitterionic, and Rydberg electronic states and calculated absorption spectra weighted by the oscillator strengths that couple those states with the electronic ground state. An excellent agreement with the experimental spectrum measured in Reference 48 supports our model.

5 Competing interests

There are no competing interests to declare.

6 Data availability

Extra information such as ground state normal mode frequencies, potential energy curves along the twelve degrees of freedom and ab initio points against which the curves are fitted, parameters for the one-dimensional diabatic potentials and intra-state, linear and bilinear coupling parameters can be found in the Supporting Information. The quantics inputs and operator file are deposited at the UCL Repository https://rdr.ucl.ac.uk DOI: 10.5522/04/21318339.

Acknowledgement

We thank the MOLecules In Motion COST Action (CM1405) for funding short term scientific missions from the University of Vienna to University College London that made this collaboration possible. SG acknowledges NextGenerationEU funds (María Zambrano Grant for the attraction of international talent). GW thanks the EPSRC for funding through grant EP/S028781/1.

References

- Young, T. II. The Bakerian Lecture. On the theory of light and colours. <u>Philos. Trans.</u> <u>R. Soc. London</u> 1802, <u>92</u>, 12–48.
- (2) Arndt, M.; Nairz, O.; Vos-Andreae, J.; Keller, C.; van der Zouw, G.; Zeilinger, A.
 Wave-particle duality of C60 molecules. Nature 1999, 401, 680.
- (3) Eibenberger, S.; Gerlich, S.; Arndt, M.; Mayor, M.; Tüxen, J. Matter-wave interference with particles selected from a molecular library with masses exceeding 10000 amu. <u>Phys.</u> Chem. Chem. Phys **2013**, 15, 14696.
- (4) Hund, F. Zur Deutung der Molekelspektren 2. Z. Phys **1927**, <u>42</u>, 93.
- (5) Heisenberg, W. Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik. Z. Phys 1927, 41, 239.

- (6) Panfilov, V.; Strunin, V.; Chapovsky, P. Light-induced drift of CH₃F molecules. <u>Sov.</u> Phys. J. Exp. Theor. Phys. **1983**, 58, 510.
- (7) Chapovsky, P.; Krasnoperov, L.; Panfilov, V.; Strunin, V. Studies on separation and conversation of spin modification of CH₃F molecules. Chem. Phys. **1985**, 97, 449.
- (8) Bakarev, A.; ., P. C. Anomalously large isotopic effect observed in the conversion of different nuclear-spin modifications of CH₃F molecules. <u>J. Exp. Theor. Phys. Lett.</u> 1986, <u>44</u>, 4.
- (9) Chapovsky, P. L. Conversion of Nuclear Spin Modifications of CH₃F Molecules in the Gaseous Phase. Sov. Phys. J. Exp. Theor. Phys. **1990**, 70, 895.
- (10) Nagels, B.; Schuurman, M.; Chapovsky, P. L.; Hermans, L. J. F. Nuclear Spin Conversion in Molecules: Experiments on CH₃¹³F Support a Mixing-of-States Model. <u>Phys.</u> Rev. A **1996**, 54, 2050.
- (11) Nagels, B.; Hermans, L. J. F.; Chapovsky, P. L. Quantum Zeno Effect Induced by Collisions. Phys. Rev. Lett. **1997**, 79, 3097.
- (12) Cacciani, P.; Coslóu, J.; Herlemont, F.; Khelkhal, M.; Lecointre, J. Nuclear spin conversion in the gaseous phase in the presence of a static electric field: Intramolecular magnetic interactions and the role of collisions. Phys. Rev. A 2004, 69, 1.
- (13) Peters, G.; Schramm, B. Nuclear spin state relaxation in formaldehyde: dependence of the rate constant on pressure. Chem. Phys. Lett. **1999**, 302, 181.
- (14) Chapovsky, P. L.; Cosléou, J.; Herlemont, F.; Khelkhal, M.; Legrand, J. Separation and conversion of nuclear spin isomers of ethylene. <u>Chem. Phys. Lett.</u> 2000, <u>322</u>, 424, 2000.
- (15) Zhen-Dong Sun, M. G.; Zheng, Y. Separation and conversion dynamics of nuclear-spin isomers of gaseous methanol. Nat. Commun. 2015, 6, 6877.

- (16) Konyukhov, V.; Prokhorov, A.; Tikhonov, V.; Faizulaev, V. Rotationally selective condensation of heavy water in a supersonic carbon dioxide jet. J. Exp. Theor. Phys. 1986, 43, 65.
- (17) Tikhonov, V. I.; Volkov, A. A. Separation of Water into Its Ortho and Para Isomers. Science 2002, 296, 2363.
- (18) Kravchuk, T.; Reznikov, M.; Tichonov, P.; Avidor, N.; Meir, Y.; Bekkerman, A.; Alexandrowicz, G. A magnetically focused molecular beam of ortho-water. <u>Science</u> 2011, <u>331</u>, 319.
- (19) Horke, D. A.; Chang, Y.-P.; Dlugolecki, K.; Küpper, J. Separating para and ortho water. Angew. Chem. Int. Ed. 2014, 53, 11965.
- (20) Kilaj, A.; Gao, H.; Rösch, D.; Rivero, U.; Küpper, J.; Willitsch, S. Observation of different reactivities of para and ortho-water towards trapped diazenylium ions. <u>Nat.</u> Commun. **2018**, 9, 2096.
- (21) Sun, Z.-D.; Takagi, K.; Matsushima, F. Separation and conversion dynamics of four nuclear spin isomers of ethylene. Science **1938**, 310, 2005.
- (22) Zhivonitko, V. V.; Kovtunov, K. V.; Chapovsky, P. L.; Koptyug, I. V. Nuclear spin isomers of ethylene: enrichment by chemical synthesis and application for NMR signal enhancement. Angew. Chem. Int. Ed. **2013**, 52, 13251.
- (23) Grohmann, T.; Leibscher, M. Nuclear spin selective alignment of ethylene and analogues. J. Chem. Phys. 2011, 134, 204316.
- (24) Floß, J.; Grohmann, T.; Leibscher, M.; Seideman, T. Nuclear spin selective laser control of rotational and torsional dynamics. J. Chem. Phys. 2012, 136, 084309.
- (25) Obaid, R.; Kinzel, D.; Oppel, M.; González, L. Discrimination of nuclear spin isomers

exploiting the excited state dynamics of a quinodimethane derivative. <u>J. Chem. Phys.</u> **2014**, 141, 164323.

- (26) Obaid, R.; Kinzel, D.; Oppel, M.; González, L. Separating nuclear spin isomers using a pump-dump laser scheme. <u>Theor. Chem. Acc.</u> 2015, 134, 46.
- (27) Belz, S.; Deeb, O.; González, L.; Grohman, T.; Kinzel, D.; Leibscher, M.; Manz, J.; Obaid, R.; Oppel, M.; Xavier, G. D.; Zilberg, S. Nuclear spin selective torsional states: implications of molecular symmetry. Z. Phys. Chem. **2013**, 227, 1021.
- (28) Grohmann, T.; Haase, D.; Jia, D.; Manz, J.; Yang, Y. Nuclear spin blockade of laser ignition of intramolecular rotation in the model boron rotor B13+11. <u>J. Chem. Phys.</u> 2018, 149, 184302.
- (29) Jia, D.; Manz, J.; Yang, Y. From coherent quasi-irreversible quantum dynamics towards the second law of thermodynamics: The model boron rotor B13+. <u>AIP Advances</u> 2018, 8, 045222.
- (30) Lu, X.-Q.; Man, Y.; Ruß, V.; Xu, Y.; Yang, Y.; Li, S.-D. The unified quantum mechanical structure of tubular molecular rotors with multiple equivalent global minimum structures: the 18*C2h → D9d case of La-[B2@B18]-La. <u>Phys. Chem. Chem. Phys.</u> 2021, 23, 19146–19149.
- (31) Yang, H.; Man, Y.; Wang, H.; Yang, Y. Spreading probability distribution and increase of entropy due to coherent quantum dynamics: Proof-of-principle for a model molecular boron rotor. Chem. Phys. **2022**, 562, 111659.
- (32) Gómez, S.; Oppel, M.; González, L. Discrimination of 1,1-difluoroethylene nuclear spin isomers in the presence of non-adiabatic coupling terms. <u>Chem. Phys. Lett.</u> 2017, <u>683</u>, 205.

- (33) Deeb, O.; Leibscher, M.; Manz, J.; von Muellern, W.; Seideman, T. Toward separation of nuclear spin isomers with coherent light. ChemPhysChem 2007, 8, 322.
- (34) Grohmann, T.; Deeb, O.; Leibscher, M. Quantum separation of para- and ortho-fulvene with coherent light: the influence of the conical intersection. <u>Chem. Phys.</u> 2007, <u>338</u>, 252.
- (35) Belz, S.; Grohmann, T.; Leibscher, M. Quantum dynamical simulations for nuclear spin selective laser control of ortho- and para-fulvene. J. Chem. Phys. 2009, 131, 034305.
- (36) Waldl, M.; Oppel, M.; González, L. Controlling the excited-state dynamics of nuclear spin isomers using the dynamic Stark effect. J. Phys. Chem. A 2016, 120, 4907.
- (37) Barbatti, M.; Sen, K. Effects of different initial condition samplings on photodynamics and spectrum of pyrrole. Int. J. Quantum Chem. 2016, 116, 762–771.
- (38) Suchan, J.; Hollas, D.; Curchod, B. F. E.; Slavíček, P. On the importance of initial conditions for excited-state dynamics. Faraday Discuss. 2018, 212, 307–330.
- (39) Avagliano, D.; Lorini, E.; González, L. Sampling effects in quantum mechanical/molecular mechanics trajectory surface hopping non-adiabatic dynamics. <u>Philos.</u> Trans. R. Soc., A **2022**, 380, 20200381.
- (40) Tully, J. C.; Preston, R. K. Trajectory Surface Hopping Approach to Nonadiabatic Molecular Collisions: The Reaction of H⁺ with D₂. J. Chem. Phys, **1971**, 55, 562.
- (41) Ben-Nun, M.; Martínez, T. J. Adv. Chem. Phys.; John Wiley and Sons, 2002; p 439.
- (42) Richings, G. W.; Polyak, I.; Spinlove, K. E.; Worth, G. A.; Burghardt, I.; Lasorne, B.
 Quantum dynamics simulations using Gaussian wavepackets: the vMCG method. <u>Int.</u> Rev. Phys. Chem. **2015**, 34, 269.

- (43) Beck, M. H.; Meyer, H.-D. An efficient and robust integration scheme for the equations of motion of the multiconfiguration time-dependent Hartree (MCTDH) method. <u>Z.</u> Phys. D. Atom. Mol. Cl. **1997**, 42, 113.
- (44) Beck, M. H.; Jäckle, A.; Worth, G. A.; Meyer, H.-D. The multiconfiguration timedependent Hartree (MCTDH) method: a highly efficient algorithm for propagating wavepackets. Phys. Rep. 2000, 324, 1.
- (45) Worth, G. A.; Meyer, H.-D.; Köppel, H.; Cederbaum, L. S.; Burghardt, I. Using the MCTDH wavepacket propagation method to describe multimode nonadiabatic dynamics. Int. Rev. Phys. Chem. 2008, 27, 569–606.
- (46) Köppel, H.; Domcke, W.; Cederbaum, L. S. Multi-mode molecular dynamics beyond the Born-Oppenheimer approximation. Adv. Chem. Phys. **1984**, 57, 59.
- (47) Gómez, S.; Ibele, L. M.; González, L. The 3s Rydberg state as a doorway state in the ultrafast dynamics of 1,1-diffuoroethylene. Phys. Chem. Chem. Phys. 2019, 21, 4871.
- (48) Limão Vieira, P.; Vasekova, E.; Sekhar, B.; Mason, N.; Hoffman, S. VUV electronic state spectroscopy of 1,1-difluoroethene and difluorochloromethane by high resolution synchrotron radiation. Phys. Chem. Chem. Phys. 2006, 8, 4766.
- (49) Worth, G. A.; Cederbaum, L. S. Beyond Born-Oppenheimer: Conical intersections and their impact on molecular dynamics. Ann. Rev. Phys. Chem. 2004, 55, 127–158.
- (50) Cattarius, C.; Worth, G. A.; Meyer, H.-D.; Cederbaum, L. S. All mode dynamics at the conical intersection of an octa-atomic molecules: Multi-configuration time-dependent Hartree (MCTDH) investigation on the butatriene cation. <u>J. Chem. Phys.</u> 2001, <u>115</u>, 2088–2100.
- (51) Worth, G. A.; Giri, K.; Richings, G. W.; Beck, M. H.; Jäckle, A.; Meyer, H.-D. QUAN-

TICS, a suite of programs for molecular QUANTum dynamICS simulations, Version 1.1. 2015.

- (52) Frisch, M. J. et al. Gaussian 09 Revision A.1. Gaussian Inc. Wallingford CT 2009.
- (53) L.M. Sverdlov, M. K.; Krainov, E. Vibrational spectra of polyatomic molecules. <u>J. Mol.</u> Struct. **1975**, 26, 136.
- (54) Mulliken, R. S. Electronic structures of polyatomic molecules and valence. IV. Electronic states, quantum theory of the double bond. Phys. Rev. 1933, 43, 279.
- (55) Auer, J.; Krotscheck, E.; Chin, S. A. A fourth-order real-space algorithm for solving local Schrödinger equations. J. Chem. Phys. 2001, 115, 6841.
- (56) Burden, R. L.; Faires, J. D. <u>Numerical analysis</u>, 5th ed.; Boston PWS-Kent Pub. Co, 1993.
- (57) Light, J. C.; Hamilton, I. P.; Lill, J. V. Generalized discrete variable approximation in quantum mechanics. J. Chem. Phys. **1985**, 82, 1400.
- (58) Kosloff, D.; Kosloff, R. A Fourier-method solution for the time-dependent Schrödinger equation as a tool in molecular dynamics. J. Comput. Phys. **1983**, 52, 35.
- (59) Ben-Nun, M.; Martínez, T. J. Photodynamics of ethylene: ab initio studies of conical intersections. Chem. Phys. 2000, 259, 237.
- (60) Christopoulou, G.; Freibert, A.; Worth, G. A. Improved algorithm for the direct dynamics variational multi-configurational gaussian method. <u>J. Chem. Phys.</u> 2021, <u>154</u>, 124127–13.
- (61) Richings, G. W.; Worth, G. A. Multi-state non-adiabatic direct-dynamics on propagated diabatic potential energy surfaces. Chem. Phys. Lett. 2017, 683, 606–612.