arXiv:2302.08461v1 [math.GR] 16 Feb 2023

REGULAR SEMIGROUPS WEAKLY GENERATED BY ONE ELEMENT

LUÍS OLIVEIRA

ABSTRACT. In this paper we study the regular semigroups weakly generated by a single element x, that is, with no proper regular subsemigroup containing x. We show there exists a regular semigroup F_1 weakly generated by x such that all other regular semigroups weakly generated by xare homomorphic images of F_1 . We define F_1 using a presentation where both sets of generators and relations are infinite. Nevertheless, the word problem for this presentation is decidable. We describe a canonical form for the congruence classes given by this presentation, and explain how to obtain it. We end the paper studying the structure of F_1 . In particular, we show that the 'free regular semigroup FI_2 weakly generated by two idempotents' is isomorphic to a regular subsemigroup of F_1 weakly generated by $\{xx', x'x\}$.

1. INTRODUCTION

An element s of a semigroup S is called *regular* if sts = s for some $t \in S$. Note that s' = tst is a (von Neumann) *inverse* of s, that is, ss's = s and s'ss' = s'. We denote by V(s) the set of all inverses of s in S. A *regular* semigroup is a semigroup with all elements regular. As usual, we denote by E(S) the set of idempotents of S.

A subset of a regular semigroup S may not generate a regular subsemigroup. Hence, the following notion of "weakly generated" seems natural: a regular subsemigroup T of S is weakly generated by a subset X if T has no proper regular subsemigroup containing X. Of course, this does not mean that T is effectively generated by X. Very often, the subsemigroup generated by X is a proper non-regular subsemigroup of T. In particular, we say that S is weakly generated by X if S has no proper regular subsemigroup containing X. We should also point out that the same set X may weakly generate several distinct regular subsemigroups of S.

The interest in studying the structure of regular semigroups weakly generated by a set X came first from the theory of e-varieties of regular semigroups (see [2, 3] for the definition of e-variety). The existence of bifree objects on an e-variety V depends on the validity of the following property in V: for

²⁰²⁰ Mathematics Subject Classification. (Primary) 20M17, (Secondary) 20M05, 20M10.

Key words and phrases. Regular semigroup, Weakly generated semigroup, Word problem.

all $S \in \mathbf{V}$ and all matched subset A of S, there exists a unique regular subsemigroup of S weakly generated by A (see [7]).

In [6] we studied the structure of the regular semigroups weakly generated by idempotents. We proved the existence of a *free regular semigroup* FI(X)*weakly generated by a set X of idempotents* in the sense that FI(X) is weakly generated by X and all other regular semigroups weakly generated by X are homomorphic images of FI(X). We got FI(X) by introducing a presentation with both sets of generators and relations infinite. Despite that fact, the word problem for that presentation is decidable since a procedure to obtain a canonical representative for each congruence class was described.

If FI_k denotes FI(X) for |X| = k, we proved that FI_2 contains copies of all FI_k as subsemigroups. Thus, any regular semigroup generated by a finite set of idempotents must strongly divide FI_2 , that is, must be a homomorphic image of some regular subsemigroup of FI_2 . The case FI_1 is of no interest since any regular semigroup weakly generated by a single idempotent is just the trivial semigroup.

We intend to look now to the general non-idempotent case. In other words, we plan to see if there exists a regular semigroup F(X) weakly generated by a set X (not necessarily of idempotents) such that all other regular semigroups weakly generated by X are homomorphic images of F(X). The case where X is a singleton set is now distinct from the idempotent case: there are nontrivial regular semigroups weakly generated by a single element. So, in this paper, we focus only on the case where X is a singleton set and prove that F(X) exists for |X| = 1.

This paper will follow the structure of [6]. In fact, we will use the same terminology, but the concepts will be more complex here. For example, terms such as landscape and mountain will be used here again but to refer to words with a more intricate structure. The results we will present are also similar to the ones of [6], and some of the proofs use even the same arguments (or with just small adaptations), although they refer to more complex objects. For the sake of completeness, we decided to include those proofs again in this paper.

We will denote F(X) by F_1 for |X| = 1, and we will represent by xthe unique element of X. In fact, we will usually not mention X but its unique element x instead. As stated above, the main goal of this paper is to prove the existence of F_1 . We begin by introducing F_1 using a presentation similarly to [6]. It is an interesting exercise to compare the presentation for F_1 with the presentation for FI(X). We will see their resemblance, but also notice a new 'ingredient' appearing in F_1 , namely conjugation. We then prove that the presentation introduced here has decidable word problem by obtaining a canonical form for each congruence class. To attest the richness of the structure of F_1 , we will prove that F_1 has a regular subsemigroup weakly generated by the idempotents $\{xx', x'x\}$ isomorphic to FI₂, where x'is the only inverse of x in F_1 . Since all regular semigroups weakly generated by a finite set of idempotents strongly divide FI_2 , they also strongly divide F_1 .

This paper is organized as follows. In the next section, we recall some basic concepts needed for this paper. We include also a brief description of the construction of FI_2 since we will need it later. We change also the terminology used in [6] so that it does not overlap with the terminology used here. In Section 3 we introduce a presentation and define F_1 . We present also a solution for the word problem for this presentation. We prove that F_1 is a regular semigroup weakly generated by x in Section 4. To achieve this result we need to characterize first the Green's relations on F_1 . Since we characterize the Green's relations in this section, we end it by describing also the idempotents, the inverses and the natural partial order. However, as it will become evident, this description will be just theoretical and not very useful for practical purposes. In Section 5, we show that all regular semigroups weakly generated by x are homomorphic images of F_1 . Finally, in the last section, we prove that F_1 has a regular subsemigroup weakly generated by $\{xx', x'x\}$ isomorphic to FI₂; thus concluding that all regular semigroups generated by a finite set of idempotents strongly divide F_1 .

2. Preliminaries

Given a semigroup S, S^1 denotes the monoid obtained by adding an identity element to S if necessary. The quasi-orders $\leq_{\mathscr{L}}, \leq_{\mathscr{R}}$ and $\leq_{\mathscr{J}}$ on S are defined as follows:

 $s\leq_{\mathscr{L}}t\Leftrightarrow S^1s\subseteq S^1t,\quad s\leq_{\mathscr{R}}t\Leftrightarrow sS^1\subseteq tS^1,\quad s\leq_{\mathscr{J}}t\Leftrightarrow S^1sS^1\subseteq S^1tS^1.$

Thus, the Green's equivalence relations \mathscr{L}, \mathscr{R} and \mathscr{J} are just the relations

$$\mathscr{L} = \leq_{\mathscr{L}} \cap \geq_{\mathscr{L}}, \quad \mathscr{R} = \leq_{\mathscr{R}} \cap \geq_{\mathscr{R}} \quad \text{ and } \quad \mathscr{J} = \leq_{\mathscr{J}} \cap \geq_{\mathscr{J}},$$

where $\geq_{\mathscr{L}}, \geq_{\mathscr{R}}$ and $\geq_{\mathscr{J}}$ are respectively the dual relations of $\leq_{\mathscr{L}}, \leq_{\mathscr{R}}$ and $\leq_{\mathscr{J}}$. Let $\mathscr{H} = \mathscr{L} \cap \mathscr{R}$ and $\mathscr{D} = \mathscr{L} \vee \mathscr{R}$ be the other two Green's relations. For a regular semigroup S, we can use S instead of S^1 in the previous definitions. We will denote by K_a the \mathscr{K} -class of the element $a \in S$, for $\mathscr{K} \in \{\mathscr{H}, \mathscr{L}, \mathscr{R}, \mathscr{D}, \mathscr{J}\}$.

A crucial concept for this paper is also the notion of sandwich set S(e, f) of two idempotents e and f of S. This concept has a few equivalent definitions. We list some of them next:

$$\begin{split} S(e,f) &= V(ef) \cap E(fSe) = fV(ef)e \\ &= \left\{g \in E(S) \mid fg = g = ge \text{ and } egf = ef\right\}. \end{split}$$

The sandwich set S(e, f) is nonempty precisely when ef is a regular element of S. Thus, S(e, f) is always nonempty if S is regular. In fact, S(e, f) is always a rectangular band and a subsemigroup of S, whenever it is nonempty. There is another interesting property of these sets: if $e, e_1, f, f_1 \in E(S)$ are such that $e \mathcal{L} e_1$ and $f \mathcal{R} f_1$, then $S(e, f) = S(e_1, f_1)$. Hence, for regular semigroups, one can extend the definition of sandwich set to all elements of

S as follows: for $a, b \in S$, let S(a, b) = S(a'a, bb') for some (any) $a' \in V(a)$ and $b' \in V(b)$.

If S is a regular semigroup, there is also another important relation on S, the *natural partial order* \leq :

$$s \leq t \quad \Leftrightarrow \quad s = et = tf \text{ for some } e, f \in E(S).$$

A useful fact about the natural partial order is that we can choose the idempotents e and f such that $e \mathscr{R} s \mathscr{L} f$.

Let X be a nonempty set. As usual, we denote by X^+ the free semigroup on X. The elements of X are called *letters* in this context, while the elements of X^+ are called *words*. The *content* of a word u is the set of letters from X that occur in u, and the *length* of u is the number of letters (counting repetitions) that occur in u. For $u \in X^+$, let $\sigma(u)$ and $\tau(u)$ denote the first and the last letter of u, respectively.

Next, we recall the construction of FI₂ introduced in [6] for any set X. As explained before, we need to modify the notation and terminology used in [6] so that it does not overlap with the one used in this paper. Hence h, H and ρ will replace the symbols g, G and ρ used in [6]. Given a triple h = (a, b, c), we denote a, b and c by h^r , h^c and h^l , respectively.

Let $X = \{e, f\}$ be a two element set, and consider

$$H_0 = \{1\}$$
 and $H_1 = \{(1, e, 1), (1, f, 1)\},\$

where 1 is a new symbol not in X. We identify each $x \in X$ with the triple $(1, x, 1) \in H_1$. Thus $x^r = x^l = 1$. Recursively for i > 1, assume that H_{i-1} is a set of triples and let $\overline{H}_i = H_{i-1} \times H_{i-2} \times H_{i-1}$ and

$$\mathbf{H}_{i} = \left\{ h \in \overline{\mathbf{H}}_{i} \mid h^{l} \neq h^{r}, \ h^{c} \in \{(h^{l})^{l}, (h^{l})^{r}\} \cap \{(h^{r})^{l}, (h^{r})^{r}\} \right\}.$$

Note that

$$\mathbf{H}_2 = \{h_{2,1}, \, h_{2,2}\}$$

for $h_{2,1} = (e, 1, f)$ and $h_{2,2} = (f, 1, e)$, and

$$\mathbf{H}_3 = \{h_{3,1}, h_{3,2}, h_{3,3}, h_{3,4}\}$$

for $h_{3,1} = (h_{2,1}, e, h_{2,2}), h_{3,2} = (h_{2,1}, f, h_{2,2}), h_{3,3} = (h_{2,2}, e, h_{2,1})$ and $h_{3,4} = (h_{2,2}, f, h_{2,1})$. Let $\mathbf{H} = \bigcup_{i \in \mathbb{N}_0} \mathbf{H}_i$.

Consider the following two relations on the free semigroup H⁺:

$$\varrho_e = \{(1h, h), (h1, h), (h^2, h) \mid h \in \mathbf{H}\}\$$

and

$$\varrho_s = \left\{ (h^c h^l h, h), (h h^r h^c, h), (h^r h^c h h^c h^l, h^r h^c h^l) \mid h \in \mathcal{H}_i \text{ with } i \ge 2 \right\}.$$

Let FI_2^1 be the semigroup given by the presentation $\langle \mathrm{H}, \varrho_e \cup \varrho_s \rangle$, that is, $\operatorname{FI}_2^1 = \mathrm{H}^+ / \varrho$ where ϱ is the smallest congruence on H^+ containing $\varrho_e \cup \varrho_s$. Note that ϱ_e just tells us that FI_2^1 is an idempotent generated monoid with identity element 1ϱ . Using both ϱ_e and ϱ_s , one can show easily that $(h^r h^c)\varrho$ and $(h^c h^l)\varrho$ are idempotents of FI_2^1 . Then ϱ_s just turns $h\varrho$ into an element

of the sandwich set $S((h^r h^c) \varrho, (h^c h^l) \varrho)$. The semigroup FI₂ is defined as FI₂¹ \{1 ϱ } ({1 ϱ }} is the group of units of FI₂¹). We can also easily see that 1 ϱ is constituted by all words of H⁺ with content {1}.

A nontrivial *i-mountain* (called mountain in [6]) is a word $h_0h_1 \cdots h_{2n} \in$ H⁺, with $n \ge 1$ and $h_i \in$ H for all $0 \le i \le 2n$, such that

- (i) $h_0 = 1 = h_{2n};$
- (*ii*) $h_{i-1} \in \{h_i^r, h_i^l\}$ for all $1 \le i \le n$; and
- (*iii*) $h_i \in \{h_{i-1}^r, h_{i-1}^l\}$ for all $n < i \le 2n$.

In [6] we proved that each ρ -class of FI(X) contains a unique nontrivial i-mountain and solved the word problem for FI(X) by given a process to construct it. In this paper we will use the term mountain to refer to more complex words.

Let i-M (denoted by M(X) in [6] for |X| = 2) denote the set of all nontrivial i-mountains of H⁺. In [6] we defined an operation \odot in i-M that turns i-M into a semigroup isomorphic to FI₂. The concept of *i*-river (called river in [6]) and the process of *uplifting of i*-rivers (called uplifting of rivers in [6]) were central. In this paper we will use the terms river and uplifting of rivers again. Their definitions will be very similar to the ones introduced in [6]. They differ only on the 'context' where they are defined. It will be immediate to see what i-river and uplifting of i-rivers mean from the notions of river and uplifting of rivers introduced here, but the readers may consult [6] for further details.

The semigroup FI_2 is regular and weakly generated by the two idempotents e and f (identifying each letter with the corresponding ρ -class). It has the following universal property: all regular semigroups weakly generated by two idempotents are homomorphic images of FI_2 (under a homomorphism that sends $e\rho$ and $f\rho$ into those two idempotents). In Section 6 we show that FI_2 is embedded into the semigroup F_1 that we will construct in the next section.

3. The presentation $\langle G, R \rangle$

This section is devoted to the construction of F_1 and, after some initial terms are introduced, we will divide it into subsections for a better organization. As we will see, the construction of F_1 resembles that of FI_2 , and it will be easier to define and work with the monoid F_1^1 obtained from F_1 by adding a new identity. In contrast to the FI_2 case, we will work now with 5-tuples since more information is needed to be included in those tuples. Conjugation will have now an important role too and the two extra entries will contain information about when and where to apply conjugation.

We begin by defining a presentation $\langle G, R \rangle$ for F_1^1 . From now on, and if nothing is said in contrary, X will be the singleton set $\{x\}$. Let A be the set $\{1, x, x'\}$. We endow A with a natural involution ' by setting (x')' = x and 1' = 1. We call *anchors* to the elements of A. As should be expected, x'will represent the inverse of x in F_1^1 , while 1 will correspond to its identity

element. For that reason, we will make some abuse of terminology from now on by calling a' the inverse (anchor) of a, for any $a \in A$.

The generator set G will contain the set A. All other elements from G will be special 5-tuples. We will denote the set of all those 5-tuples by G^5 . Thus $G = A \cup G^5$. The set G^5 will be defined recursively as the union of sets G_i for $i \ge 1$. In fact, each G_i will be the union of two other sets $G_{i,e}$ and $G_{i,d}$. We will describe these sets in more detail below.

We will use the notation

$$g = (g^l, g^{la}, g^c, g^{ra}, g^r)$$

to refer to the entries of a 5-tuple g of G^5 . The entries g^{la} and g^{ra} will be always anchors. For that reason, they will be called the *left anchor* and the *right anchor* of g, respectively. The entries g^l , g^c and g^r will be called the *left, middle* and *right* entries of g. The entries g^l and g^r will be 5-tuples of G_{i-1} if $g \in G_i$ for i > 1; and the entry g^c will be a 5-tuple of G_{i-2} if $g \in G_i$ for i > 2.

As we can see, the left, middle and right entries of a 5-tuple $g \in G^5$ are usually 5-tuples again. If we need to refer to an entry of g^l for example, we will use the abbreviation g^{ls} to refer to the entry $(g^l)^s$ of g^l , for $s \in$ $\{l, la, c, ra, r\}$. But, some entries of g^l may be again 5-tuples. In general, we convene that

$$g^{s_1\cdots s_n} = (((g^{s_1})^{s_2})\cdots)^{s_n},$$

for $s_1, \dots, s_{n-1} \in \{l, c, r\}$ and $s_n \in \{l, la, c, ra, r\}$, whenever the right side of this equality makes sense. We consider also $g^1 = g$.

To help following the remaining of this section, we will divide it into subsections. In the first two subsections we define the sets $G_{i,e}$ and $G_{i,d}$, respectively. It is important that the reader pays some attention to the subtleties involved in these definitions as they may be important to understand some of the arguments presented during the paper. In the following subsection we introduce the set of relations R. This set will give us an interpretation for the 5-tuples of G^5 . In the forth subsection some special words are described. Finally, the operation of uplifting of rivers is introduced in the fifth subsection and used in the last subsection to present a solution to the word problem for this presentation.

3.1. The sets $G_{i,e}$. We begin by setting $g_{xx'}$ as the 5-tuple

$$g_{xx'} = (1, 1, x, x', 1)$$

The 5-tuple $g_{xx'}$ will represent the element xx' of F_1 . For technical reasons that will become clear later, it is convenient to consider the 5-tuple $g_{xx'}$ instead of xx'. Set $G_{1,e} = \{g_{xx'}\}$ and define

$$\mathbf{G}_{i,e} = \left\{ \left(g, (g^{la})', g^l, (g^{ra})', g \right), \left(g, (g^{ra})', g^l, (g^{la})', g \right) : g \in \mathbf{G}_{i-1,e} \right\}$$

 $\mathbf{6}$

for $i \geq 2$. To help clarifying which tuples belong to these sets, note that $G_{2,e}$ has two elements:

 $q_{2,e,1} = (q_{xx'}, 1, 1, x, q_{xx'})$ and $q_{2,e,2} = (q_{xx'}, x, 1, 1, q_{xx'});$

while $G_{3,e}$ has four elements:

$$\begin{array}{ll} (g_{2,e,1},1,g_{xx'},x',g_{2,e,1})\,, & (g_{2,e,1},x',g_{xx'},1,g_{2,e,1})\,, \\ (g_{2,e,2},1,g_{xx'},x',g_{2,e,2}) & \text{and} & (g_{2,e,2},x',g_{xx'},1,g_{2,e,2})\,. \end{array}$$

Analyzing the structure of the 5-tuples of $G_{i,e}$ in more detail for i > 1, we see that $g \in G_{i,e}$ if and only if

- (i) $g^l = g^r \in \mathcal{G}_{i-1,e}$,

(*ii*) $g^c = g^{l^2} = g^{lr}$, and (*iii*) either $\{g^{la}, g^{ra}\} = \{1, x\}$ if *i* even or $\{g^{la}, g^{ra}\} = \{1, x'\}$ if *i* odd.

Hence, the set $G_{i,e}$ has twice the number of elements of $G_{i-1,e}$, and so $G_{i,e}$ has 2^{i-1} elements. Further,

$$\mathbf{G}_{i,e} \subseteq \mathbf{G}_{i-1,e} \times A \times \mathbf{G}_{i-2,e} \times A \times \mathbf{G}_{i-1,e},$$

for i > 1, if one considers $G_{0,e} = \{1\}$. Let $G_e = \bigcup_{i \in \mathbb{N}} G_{i,e}$. The 5-tuples of G_e will correspond to the 5-tuples of G with the same left and right entry.

By construction, both g^{l^2} and g^{lr} are equal to g^c for each $g \in G_{i,e}$ with $i \ge 2$. Of course, they have different meanings: g^c represents the middle entry of g, while g^{l^2} and g^{lr} represent the left and right entries of g^l , respectively. In the special words that we will need to consider later, the 5-tuples of G will have to be 'anchored' to each other. For that, we need to attribute a side to g^c inside g^l in order to identify 'the anchor of g^c in g^l with respect to g'. Note that only one of the elements q^{l^2a} and q^{lra} is equal to $(q^{la})'$. For each q, we define

$$l_a = \begin{cases} l^2 & \text{if } g^{l^2 a} = (g^{la})' \\ lr & \text{if } g^{lra} = (g^{la})' \end{cases}$$

Therefore, the anchors $g^{l_a a}$ and $(g^{l_a})'$ are equal.

We reinforce that the purpose of the notation l_a is to identify which anchor g^{l^2a} or g^{lra} is equal to $(g^{la})'$, and that l_a differs according to g. Hence, the use of l_a only makes sense when we refer to a specific 5-tuple $g \in G_i$ with $i \geq 2$. Note also that g^{l_a} and g^c have the same 'value'. However, when we write q^{l_a} , we want to emphasize not only its 'value', but also a specific entry of the 5-tuple q^l , either its left or its right entry.

In a similar manner we will use also the notation r_a . For each $g \in G_{i,e}$ with $i \geq 2$,

$$r_a = \begin{cases} r^2 & \text{if } g^{r^2 a} = (g^{ra})' \\ rl & \text{if } g^{rla} = (g^{ra})' \end{cases}$$

We alert again that the use of r_a only makes sense when associated with a specific 5-tuple $g \in G_{i,e}$ with $i \ge 2$.

3.2. The sets $G_{i,d}$. We now introduce the 5-tuples of G with distinct left and right entries. They will be described recursively in the sets $G_{i,d}$ for $i \ge 1$. In contrast with the previous case, the left and right entries of a 5-tuple g from $G_{i,d}$ may belong not only to $G_{i-1,d}$, but also to $G_{i-1,e}$. Thus, we set $G_i = G_{i,e} \cup G_{i,d}$. As we will see next, the definition of the sets $G_{i,d}$ is more intricate than the definition of the sets $G_{i,e}$.

In fact, the definition of the sets $G_{i,d}$ is only interesting for i > 2, since we set $G_{1,d}$ and $G_{2,d}$ as empty sets. Basically, what it says is that for i = 1 and i = 2 there are no 5-tuples in G with distinct left and right entries. Now, assume that i > 2 and that $G_{j,d}$ is defined for j < i. Consequently, the sets G_j are also defined. Let $G_{i,d}$ be the set of all 5-tuples

$$g \in G_i = G_{i-1} \times A \times G_{i-2} \times A \times G_{i-1}$$

such that

(i)
$$g^{l} \neq g^{r};$$

(ii) $(g^{c}, g^{la}) = (g^{l^{2}}, (g^{l^{2}a})')$ or $(g^{c}, g^{la}) = (g^{lr}, (g^{lra})');$
(iii) $(g^{c}, g^{ra}) = (g^{rl}, (g^{rla})')$ or $(g^{c}, g^{ra}) = (g^{r^{2}}, (g^{r^{2}a})').$

From (i) we know that the left entry of g is distinct from its right entry. Note that (ii) tells us, in particular, that g^c is the left or the right entry of g^l . In fact, it tells us something stronger: either g^c is the left entry of g^l and the left anchor of g is the inverse of the left anchor of g^l , or g^c is the right entry of g^l and the left anchor of g is the inverse of the inverse of the right anchor of g^l . Condition (iii) is just the dual of (ii) with respect to g^r .

Next, we list the elements of $G_{3,d}$ as an example to help understand these sets. It has already 8 elements:

$g_{3,d,1} = (g_{2,e,1}, 1, g_{xx'}, 1, g_{2,e,2})$	$(g_{2,e,1}, 1, g_{xx'}, x', g_{2,e,2})$,
$g_{3,d,2} = (g_{2,e,1}, x', g_{xx'}, x', g_{2,e,2}) ,$	$(g_{2,e,1}, x', g_{xx'}, 1, g_{2,e,2})$,
$g_{3,d,3} = (g_{2,e,2}, 1, g_{xx'}, 1, g_{2,e,1})$,	$(g_{2,e,2}, 1, g_{xx'}, x', g_{2,e,1})$,
$g_{3,d,4} = (g_{2,e,2}, x', g_{xx'}, x', g_{2,e,1}) ,$	$(g_{2,e,2}, x', g_{xx'}, 1, g_{2,e,1})$.

As the reader may have notice, we have attributed a 'name' only to the elements of the first column. The reason is because only these elements will be important for Section 6. The elements from the second column will not be used there. It is also a combinatorial exercise to compute the cardinality of $G_{i,d}$ and G_i . Since we do not need that information for this paper, we leave that computation for the reader. We just point out the obvious fact that the size of the sets $G_{i,d}$, $G_{i,e}$ and G_i increases exponentially with *i*.

Like in the previous subsection, for each $g \in G_{i,d}$, we need to attribute a side to g^c inside g^l . In this case, there are two distinct situations accordingly to $g^l \in G_{i-1,e}$ or $g^l \in G_{i-1,d}$. If $g^l \in G_{i-1,e}$, then g^{l^2} and g^{lr} are both equal to g^c . In this case, due to condition (*ii*) above, we can define l_a in the same manner as we did in the previous subsection. If $g^l \in G_{i-1,d}$, then the definition of l_a is more natural since only one of left and right entries of g^l

is equal to g^c : let $l_a = l^2$ if $g^{l^2} = g^c$ or let $l_a = lr$ if $g^{lr} = g^c$. The notation r_a is now introduced similarly (see the previous subsection). We alert once more that these two notations only make sense when referring to a specific 5-tuple g.

By definition of both $G_{i,d}$ and $G_{i,e}$, observe that

$$g^{l_a a} = (g^{la})'$$
 and $g^{r_a a} = (g^{ra})'$

in all cases where the notation makes sense (that is, except for $g = g_{xx'}$). This observation will be used frequently in this paper without further notice. A quick inspection to the definition of g^{la} and g^{ra} allow us to conclude also that $g^{la}, g^{ra} \in \{1, x\}$ if v(g) even, while $g^{la}, g^{ra} \in \{1, x'\}$ if v(g) odd. In fact, if $g \in G_e$, we can say more: $\{g^{la}, g^{ra}\} = \{1, x\}$ if v(g) even, while $\{g^{la}, g^{ra}\} = \{1, x'\}$ if v(g) even, while $\{g^{la}, g^{ra}\} = \{1, x'\}$ if v(g) odd. If $g \in G_{i,d}$, we may have $g^{la} = g^{ra}$ as can be observed in some elements of $G_{3,d}$.

Note that the set $G_{i,e}$ is also contained in \overline{G}_i ; whence $G_i \subseteq \overline{G}_i$. Let $G_d = \bigcup_{i \in \mathbb{N}} G_{i,d}$ and $G^5 = \bigcup_{i \in \mathbb{N}} G_i = G_d \cup G_e$. Set $G = G^5 \cup A$ and $G' = G^5 \cup \{1\}$. Thus G^5 is the set of all 5-tuples of G, and it is divided into the set G_e of all 5-tuples of G with the same left and right entries, and the set G_d of all 5-tuples of G with distinct left and right entries. The *height* v(g) of an element $g \in G'$ is defined as follows:

$$v(1) = 0$$
 and $v(g) = i$ if $g \in G_i$.

3.3. The relation R. Let us look now to the relation R. We will denote by F_1^1 the semigroup given by the presentation $\langle G, R \rangle$, that is, $F_1^1 = G^+ / \rho$ where ρ is the smallest congruence containing R. We will write $u \approx v$ to indicate that the words u and v of G^+ are ρ -equivalent. We will use also the notation [u] to refer to the ρ -class $u\rho$. We want:

- (i) [x'] to be an inverse of [x] such that $[xx'] = [g_{xx'}]$;
- (*ii*) [1] to be the identity element of F_1^1 ; and
- (*iii*) [g] to be an idempotent for all $g \in \mathbf{G}^5$.

Thus, we include in R the relation

$$\rho_e = \{(xx'x, x), (x'xx', x'), (g_{xx'}, xx'), (1g, g), (g1, g), (g^2, g) : g \in \mathcal{G}'\}.$$

We immediately see now that we could have omitted $g_{xx'}$ from G. However, its inclusion gives a better feeling about the construction of G^5 : this list derives form the 'idempotent' xx', and not from x or x'. Furthermore, we will see that the congruence classes of the 5-tuples of G^5 constitute a transversal set for the \mathscr{D} -classes of F_1 . We should also point out that we could have replaced xx' with x'x in the theory developed in this paper (we just needed to adapt the definition of G^5 for the x'x case). But the definition of R is not yet finished. We still need to include the conditions that relate g with its entries. For that, we need to explain carefully some notation we will use in order to avoid future ambiguities. Given three letters $g_1, g \in G'$ and $a \in A$, the triplet g_1ag is *left anchored* if

$$(g_1, a) = (g^l, g^{la})$$
 or $(g_1, a) = (g^r, g^{ra})$.

The notion of right anchored is not quite the dual: the triplet gag_1 is right anchored if

$$(g_1, a) = (g^l, (g^{la})')$$
 or $(g_1, a) = (g^r, (g^{ra})')$

Observe that in the definition of left anchored, the anchor a is one of the anchors of g, while in the definition of right anchored, the anchor a is the inverse of one of the anchors of g. In both cases, we say that g_1 is anchored to g. Finally, a triplet g_1ag_2 , with $g_1, g_2 \in G'$ and $a \in A$, is anchored if g_1 is anchored to g_2 or g_2 is anchored to g_1 .

For $s \in \{l, r\}$, we may denote the anchored triplets $g^s g^{sa}g$ and $g(g^{sa})'g^s$ by $g^s \cdot g$ and $g \cdot g^s$, respectively. Note that this notation may become ambiguous if we don't indicate explicitly the value of s. For example, if $g \in G_e$ and $h = g^l$, then the expression $h \cdot g$ is ambiguous. Note that h is also equal to g^r and, therefore, $h \cdot g$ may refer to either $g^l g^{la}g$ or $g^r g^{ra}g$, with $g^{la} \neq g^{ra}$. Hence, we must be careful when using this 'dot' notation.

There is another instance where this 'dot' notation may be useful but ambiguous too if we don't clarify it, namely $g^c \cdot g^s$ and $g^s \cdot g^c$ for $s \in \{l, r\}$. Note that g^c may be equal to both g^{sl} and g^{sr} . We define

$$g^{c} \cdot g^{s} = g^{c}g^{s_{a}a}g^{s} = g^{c}(g^{sa})'g^{s}$$
 and $g^{s} \cdot g^{c} = g^{s}(g^{s_{a}a})'g^{c} = g^{s}g^{sa}g^{c}$.

Clearly, these triplets are anchored by the way they are defined.

Next, we introduce the (non-anchored) triplets g^L and g^R . Let $g^L = (g^{la})'g^lg^{la}$ and $g^R = (g^{ra})'g^rg^{ra}$. Note that

$$g^{L} \in \{1g^{l}1, xg^{l}x', x'g^{l}x\}.$$

In fact, we can say also that $g^L \neq xg^l x'$ if v(g) even, and $g^L \neq x'g^l x$ if v(g) odd. Similar conclusions are also true for g^R . Note now that

$$g \cdot g^l \cdot g = gg^L g$$
, $g^c \cdot g^l \cdot g = g^c g^L g$, $g \cdot g^r \cdot g = gg^R g$ and $g \cdot g^r \cdot g^c = gg^R g^c$.
The part observation is also useful. Since $a^r \cdot g \cdot g^l = a^r a^{ra} g(a^{la})' a^l$ and

The next observation is also useful. Since $g^r \cdot g \cdot g^\iota = g^r g^{ru} g(g^{\iota u})' g^\iota$ and

$$g^r \cdot g^c \cdot g^l = g^r (g^{r_a a})' g^c g^{l_a a} g^l = g^r g^{r a} g^c (g^{l a})' g^l ,$$

the words $g^r \cdot g \cdot g^l$ and $g^r \cdot g^c \cdot g^l$ of length 5 differ only on the middle letter: we have g in $g^r \cdot g \cdot g^l$ and g^c in $g^r \cdot g^c \cdot g^l$.

Before we continue, this is a good place to make a pause and explain the main difference between the theory developed here and the one developed in [6]. This explanation may help also the readers familiar with [6] following this paper. We have already mentioned the close resemblance between the structure of these two papers. The main difference is on what we will consider to be landscapes and mountains. In [6], they were words from H⁺ where any two consecutive letters had a special connection (see Section 2). Here, they will be words constituted by alternating letters from G' and A such that any triplet subword with two letters from G' and one from A is

anchored. The notation introduced above will help us, in some cases, to give a less cumbersome aspect to those words.

Let ρ_s be the following relation on G⁺:

$$\rho_s = \left\{ (g^c g^L g, g), (gg^R g^c, g), (g^r \cdot g \cdot g^l, g^r \cdot g^c \cdot g^l) | g \in \mathcal{G} \text{ with } \upsilon(g) \ge 2 \right\}.$$

We set $R = \rho_e \cup \rho_s$. Observe that [1] is composed by all words from G^+ of content {1}. Thus $F_1^1 \setminus \{[1]\}$ is a subsemigroup of F_1^1 . Let $F_1 = F_1^1 \setminus \{[1]\}$.

The next result gives a list of properties about the product in F_1^1 and terminates with an interpretation for ρ_s . We already know that [g] is an idempotent of F_1^1 due to the definition of ρ_e . We will prove that $[g^c g^R]$ and $[g^L g^c]$ are also idempotents of F_1^1 . Then, the definition of ρ_s is just to assure that $[g] \in S([g^R g^c], [g^c g^L])$.

Lemma 3.1. Let $g \in G$ with v(g) > 1. Then:

$$\begin{array}{ll} (i) & g \approx g^c g \approx gg^c \approx gg^L g \approx gg^R g. \\ (ii) & [gg^L], & [gg^R], & [g^Lg], & [g^Rg] \in E({\rm F}_1^1). \\ (iii) & [g^l \cdot g], & [g^Lg], & [g^r \cdot g], & [g^Rg] \in {\rm L}_{[g]} \ and & [g \cdot g^l], & [gg^L], & [g \cdot g^r], & [gg^R] \in {\rm R}_{[g]}. \\ (iv) & g^L g^c g^L \approx g^L \ and & g^R g^c g^R \approx g^R. \\ (v) & [g^c g^L], & [g^c g^R], & [g^L g^c], & [g^R g^c] \in E({\rm F}_1^1). \\ (vi) & [g] \in S([g^R g^c], & [g^c g^L]). \end{array}$$

Proof. Note that $gg^c \approx gg^R g^c g^c \approx gg^R g^c \approx g$ and $gg^L g \approx gg^c g^L g \approx g^2 \approx g$ by definition of ρ_e and ρ_s . The proof of (i) is now complete by duality. Then (ii) follow obviously from (i). Further, $[g^L g]$ and $[g^R g]$ also belong to $\mathcal{L}_{[g]}$. Then, since $(g^{la})'g^l \cdot g = g^L g$ and $(g^{ra})'g^r \cdot g = g^R g$, we must have both $[g^l \cdot g]$ and $[g^r \cdot g]$ in $\mathcal{L}_{[g]}$ too. Thus (iii) holds by duality.

Note now that

$$g^{L}g^{c}g^{L} = (g^{la})'g^{l}g^{la}g^{c}(g^{la})'g^{l}g^{la} = (g^{la})'g^{l}(g^{l_{a}a})'g^{c}g^{l_{a}a}g^{l}g^{la} = \\ = \begin{cases} (g^{la})'g^{l}(g^{l})^{L}g^{l}g^{la} & \text{if } l_{a} = l^{2} \\ (g^{la})'g^{l}(g^{l})^{R}g^{l}g^{la} & \text{if } l_{a} = lr \end{cases}$$

Applying statement (i) to g^l we get $g^L g^c g^L \approx (g^{la})' g^l g^{la} = g^L$. Similarly, we show that $g^R g^c g^R \approx g^R$ and (iv) is proved. Then (v) is an obvious consequence of (iv).

Finally, to prove (vi), we already know that [g], $[g^R g^c]$ and $[g^c g^L]$ are idempotents of F_1^1 such that $g^c g^L g \approx g$ and $gg^R g^c \approx g$. So, we only need to prove that $g^R g^c gg^c g^L \approx g^R g^c g^L$. But

$$g^R g^c g g^c g^L \approx g^R g g^L = (g^{ra})' g^r \cdot g \cdot g^l g^{la} \approx (g^{ra})' g^r \cdot g^c \cdot g^l g^{la} = g^R g^c g^L.$$

Hence, we proved that $[g] \in S([g^R g^c], [g^c g^L]).$

3.4. Special words. To facilitate the organization of this paper, if nothing is said in contrary, we will use g and h (with possible indices) to refer to elements of G', a and b (with possible indices) to refer to anchors, and u and

v (also with possible indices) to refer to words from G^+ . Next, we list some general terms and notations that we will use throughout this paper. These terms identify special words from G^+ and their characteristics that will be important to us. We use the same terminology of [6] and the notions are very much alike as the reader can verify. The main difference, as mentioned already, is on what we consider now a landscape.

- **Landscape:** alternating word $u = g_0 a_1 g_1 \cdots a_n g_n \in G^+$ $(n \ge 0)$ of letters $g_j \in G'$ and anchors $a_j \in A$ such that all triplet subwords $g_{i-1}a_ig_i$ are anchored. We denote by L the set of all landscapes of G^+ . Note also that single letters $g \in G'$ are particular landscapes; whence $G' \subseteq L$. We make notice that some of the terms below refer to properties only of the letters g_i of a landscape (and not to its anchors a_i).
- **Ridge:** letter g_i of a landscape $u = g_0 a_1 g_1 \cdots a_n g_n$ such that $v(g_{i-1}) = v(g_{i+1}) = v(g_i) 1$. Hence, a ridge is never the first nor the last letter of a landscape.
- **Peak:** highest ridge of a landscape u. Note that u can have several peaks, but all of them have the same height. If u has only one peak, we denote it by $\kappa(u)$. The height v(u) of the landscape u is the maximum between the height of its peaks, the height of $\sigma(u)$, and the height of $\tau(u)$. Note that this definition of height agrees with the notion of height for elements of G introduced earlier.
- **River:** letter g_i of a landscape $u = g_0 a_1 g_1 \cdots a_n g_n$ such that $v(g_{i-1}) = v(g_{i+1}) = v(g_i) + 1$. As with ridges, rivers are never endpoints of landscapes.
- **Hill:** landscape $u = g_0 a_1 g_1 \cdots a_n g_n$ with $n \ge 1$ such that either $g_{i-1} \in \{g_i^l, g_i^r\}$ for all $1 \le i \le n$, or $g_i \in \{g_{i-1}^l, g_{i-1}^r\}$ for all $1 \le i \le n$. In the former case we have an *uphill* since $v(g_i) = v(g_{i-1}) + 1$, while on the latter case we have a *downhill* since $v(g_i) = v(g_{i-1}) 1$. We denote by L_1, L_1^+ and L_1^- the set of all hills, uphills and downhills, respectively. Further, L_2^+ will denote the set of all landscapes composed by an uphill followed by a downhill.
- **Valley:** landscape composed by a downhill followed by an uphill. Thus, a valley has always one (and only one) river at its lower height letter. We denote by L_2^- the set of all valleys of G^+ , and by L_2 the set $L_2^+ \cup L_2^-$.
- **Mountain range:** landscape u with $\sigma(u) = \tau(u) = 1$. Thus, the height of a nontrivial mountain range is the height of its peaks. Note that all mountain ranges have length 4k + 1 for some $k \in \mathbb{N}_0$, and u = 1 is the only mountain range of length 1. If $u \neq 1$, we denote by $\lambda_l(u)$ the maximal prefix of u which is also an uphill, and by $\lambda_r(u)$ the maximal suffix of u which is also a downhill. We call $\lambda_l(u)$ and $\lambda_r(u)$ the *left hill* and the *right hill* of u, respectively. For technical

reasons, we define also $\lambda_l(1) = \lambda_r(1) = 1$. We denote by MR the set of all mountain ranges of G^+ .

Mountain: mountain range u with no rivers. Thus, it is either the trivial mountain range u = 1, or it is composed by the left hill $\lambda_l(u)$ followed by the right hill $\lambda_r(u)$. Nontrivial mountains have only one ridge. We denote by M^1 and M the sets of all mountains and nontrivial mountains, respectively. So $M = MR \cap L_2^+$.

We shall use the previous terminology also to refer to subwords of a given word $u \in G^+$. For example, a hill of $u \in L$ is a subword of u that is also a hill. A hill of $u \in L$ is called maximal if it is not properly contained in another hill of u, while a valley of u is called maximal if it is not properly contained in another valley of u. Thus, each nontrivial mountain range is uniquely decomposable into its left hill, followed by a possible empty sequence of maximal valleys between its consecutive ridges, and ending with its right hill. However, we must be careful since u_1u_2 is not a landscape if u_1 and u_2 are two landscapes such that $g = \tau(u_1) = \sigma(u_2)$: it appears a double gg at the junction of u_1 with u_2 . Therefore, we will use the notation $u_1 * u_2$ to represent the landscape obtained from u_1u_2 by replacing gg with g. Note that $u_1u_2 \approx u_1 * u_2$ since [g] is an idempotent of F_1^1 .

We will represent landscapes as line graphs where the anchors label the edges and the other letters label the vertices. We will include information about the height of the letters from G' in this line graphs as follows: we draw them such that all vertices labeled by letters with the same height belong to the same imaginary horizontal line; and labels of vertices in higher imaginary horizontal lines have higher height. We try to illustrate this idea in Figure 1. For example, looking carefully to that picture, we can conclude that $(g_1, a_2) = (g_2^s, g_2^{sa})$ and $(a_7, g_7) = ((g_6^{ta})', g_6^t)$ for some $s, t \in \{l, r\}$. Using these line graphs, we can see that the terminology introduced above has a natural interpretation. We draw again the line graph of Figure 1 in Figure 2 (omitting some unnecessary information) to illustrate some of those terms.

Next, we associate a mountain to each element of G. We begin by observing that $11g_{xx'}x1$, $1x'g_{xx'}11$ and $11g_{xx'}11$ are mountains. It may seem strange the appearance of double ones (11). This is so because one of them is consider as a letter from G' while the other is an anchor. Of course, since we know that 1 will represent the identity element of F_1^1 , we could have replaced these double ones by a single one or even omit them. However, this would mean that we would have to treat this particular letter different from the other letters from G'. For technical reason it is better not to do so, and consider the letter 1 from G' as any other letter. Therefore, we will see very often these double ones (or even longer sequences of ones) in our landscapes.

We define $\beta_1(x) = 11g_{xx'}x1$, $\beta_1(x') = 1x'g_{xx'}11$ and $\beta_1(g_{xx'}) = 11g_{xx'}11$ since $x \approx 11g_{xx'}x1$, $x' \approx 1x'g_{xx'}11$ and $g_{xx'} \approx 11g_{xx'}11$. We define also $\beta_1(1) = 1$, the trivial mountain. The definition of $\beta_1(g)$ for a 5-tuple g of

LUÍS OLIVEIRA



FIGURE 1. Illustration of a landscape in a line graph.



FIGURE 2. Illustration of some terminology.

height greater than 1 is more complex. These words are long and, therefore, we will define first their left and right hills.

Let
$$g \in \mathbf{G}^5$$
 such that $v(g) = 2n$ or $v(g) = 2n + 1$ for some $n \in \mathbb{N}$. Set

$$\beta_{1,l}(g) = g^{c^n} g^{c^{n-1}l_a a} g^{c^{n-1}l_a} g^{c^{n-1}l_a} g^{c^{n-1}} \cdots g^{c^2} g^{cl_a a} g^{cl} g^{cl_a} g^c g^{l_a a} g^l g^{l_a} g$$

and

$$\beta_{1,r}(g) = g (g^{ra})' g^r (g^{raa})' g^c (g^{cra})' g^{cr} \cdots g^{c^{n-1}} (g^{c^{n-1}ra})' g^{c^{n-1}r} (g^{c^{n-1}raa})' g^{c^n}$$

The 'brackets' under these words highlight the four element blocks used as patterns in their construction. To build $\beta_{1,l}(g)$, we add to the left of each g^{c^i} the block $g^{c^{i+1}}g^{c^il_aa}g^{c^il}g^{c^ila}$; while, to build $\beta_{1,r}(g)$, we add to the right of each g^{c^i} the block $(g^{c^ira})'g^{c^ir}(g^{c^ir_aa})'g^{c^{i+1}}$. Consequently, $\beta_{1,l}(g)$ is an uphill from g^{c^n} to g, while $\beta_{1,r}(g)$ is a downhill from g to g^{c^n} . Furthermore, $g^{c^n} = 1$ if v(g) = 2n, and $g^{c^n} = g_{xx'}$ if v(g) = 2n + 1.

We now define

$$\beta_1(g) = \begin{cases} \beta_{1,l}(g) * \beta_{1,r}(g) & \text{if } \upsilon(g) = 2n \\ 11\beta_{1,l}(g) * \beta_{1,r}(g)11 & \text{if } \upsilon(g) = 2n+1 \end{cases}$$

Clearly, $\beta_1(g)$ is a mountain if v(g) = 2n. If v(g) = 2n+1, then $\beta_1(g)$ is also a mountain since both $11g_{xx'}$ and $g_{xx'}11$ are hills. We depict the mountain $\beta_1(g)$ in Figure 3 to help visualizing it. We denote by $\lambda_l(g)$ and $\lambda_r(g)$ the left hill and the right hill of $\beta_1(g)$, respectively, for each $g \in G^5$. Hence $\beta_1(g) = \lambda_l(g) * \lambda_r(g)$ for all $g \in G^5$. Note also that $\lambda_l(g) = \beta_{1,l}(g)$ and $\lambda_r(g) = \beta_{1,r}(g)$ if v(g) is even; while $\lambda_l(g) = 11\beta_{1,l}(g)$ and $\lambda_r(g) = \beta_{1,r}(g)11$ if v(g) is odd.



FIGURE 3. Illustration of the mountain $\beta_1(g)$.

Let $u = h_0 \cdots h_k$ with $h_i \in G$ for $0 \le i \le k$. Extend β_1 to u by setting $\beta_1(u) = \beta_1(h_0) \ast \cdots \ast \beta_1(h_k)$.

Then $\beta_1(u) \in MR$. In the following result we prove that $u \approx \beta_1(u)$ for all words $u \in G^+$.

Lemma 3.2. $\beta_1(u) \approx u$ for all $u \in G^+$.

Proof. Note that we just need to prove that $\beta_1(h) \approx h$ for all $h \in G$. Indeed, if $\beta_1(h) \approx h$ for all $h \in G$, then

$$\beta_1(u) \approx \beta_1(h_0)\beta_1(h_1)\cdots\beta_1(h_k) \approx h_0h_1\cdots h_k = u$$
.

We have already observed that $\beta_1(a) \approx a$ for all $a \in A$. Clearly

$$g^{c^{i}}(g^{c^{i-1}})^{l_{a}a}g^{c^{i-1}l}g^{c^{i-1}la}g^{c^{i-1}} = (g^{c^{i-1}})^{c}(g^{c^{i-1}})^{L}g^{c^{i-1}} \approx g^{c^{i-1}}$$

by definition of ρ_s . Since we have also $11g_{xx'} \approx g_{xx'}$, we conclude that $\lambda_l(g) \approx \beta_{1,l}(g) \approx g$ for all $g \in \mathbf{G}^5$. Dually, we can conclude also that $\lambda_r(g) \approx \beta_{1,r}(g) \approx g$. Consequently

$$\beta_1(g) = \lambda_l(g) * \lambda_r(g) \approx \lambda_l(g)\lambda_r(g) \approx g^2 \approx g.$$

We have proved that $\beta_1(h) \approx h$ for all $h \in G$ as wanted.

3.5. Uplifting of rivers. Let $u = g_0 a_1 g_1 \cdots a_n g_n$ be a landscape and assume that g_i is a river of u. Then $g_{i-1}a_ig_ia_{i+1}g_{i+1}$ is a subword of u such that $g_{i-1}a_ig_i$ is a right anchored triplet and $g_ia_{i+1}g_{i+1}$ is a left anchored triplet. In other words,

$$(g_i, a_i) = (g_{i-1}^s, (g_{i-1}^{sa})')$$
 and $(g_i, a_{i+1}) = (g_{i+1}^t, g_{i+1}^{ta}),$

for some $s, t \in \{l, r\}$.

Let h_i be the 5-tuple $(g_{i+1}, a'_{i+1}, g_i, a_i, g_{i-1})$. If $g_{i-1} \neq g_{i+1}$, then $h_i \in G_d$; and if $g_{i-1} = g_{i+1}$ and $a_i \neq a'_{i+1}$, then $h_i \in G_e$. However, $h_i \notin G^5$ if $g_{i-1} = g_{i+1}$ and $a_i = a'_{i+1}$. So, set

$$v = \begin{cases} g_0 a_1 \cdots g_{i-1} a_{i+2} g_{i+2} \cdots g_n & \text{if } g_{i-1} = g_{i+1} \text{ and } a_i = a_{i+1}, \\ g_0 \cdots g_{i-1} a_i h_i a_{i+1} g_{i+1} \cdots g_n & \text{otherwise.} \end{cases}$$

If $h_i \in G^5$, then the triplet $g_{i-1}a_ih_i$ is left anchored, while the triplet $h_ia_{i+1}g_{i+1}$ is right anchored; whence v is also landscape. If $h_i \notin G^5$, then $g_{i-1}a_{i+2}g_{i+2} = g_{i+1}a_{i+2}g_{i+2}$ is an anchored triplet and again v is a landscape too. We have shown that v is always a landscape. We say that v is obtained from u by *uplifting a river* and write $u \to v$ (or $u \xrightarrow{g_i} v$ if one needs to identify the river uplifted). Thus \to is a binary relation defined on the set L of all landscapes. We illustrate the two distinct instances of this operation in Figure 4.

The word v either keeps the length of u or decreases it by 4 units. If $h_i \notin \mathbf{G}^5$, the uplifting of g_i eliminates the river g_i , but may create a new river g_{i-1} if neither g_{i-1} nor g_{i+1} are ridges of u. If $h_i \in \mathbf{G}^5$, then the uplifting of g_i replaces the river g_i with a ridge h_i . However, g_{i-1} and g_{i+1} become rivers of v unless they were ridges of u, respectively.

Let us also analyze the impact of this operation onto the 'anchors subsequence' of the landscape u. Continuing to use the notation above, if $h_i \in \mathbf{G}^5$, then this anchors subsequence remains unchanged. However, if $h_i \notin \mathbf{G}^5$, then we obtain a new anchors subsequence by deleting a subword of the form xx', x'x or 11 from the original one.

The next result tells us that the relation \rightarrow is contained in ρ .

Lemma 3.3. With the notation introduced above, $u \approx v$.



FIGURE 4. Illustration of the uplifting of a river.

Proof. If $h_i \notin \mathbf{G}^5$, then

$$g_{i-1}a_ig_ia_{i+1}g_{i+1} = \begin{cases} g_{i-1}g_{i-1}^Lg_{i-1} & \text{if } (g_i, a_{i+1}) = (g_{i+1}^l, g_{i+1}^{la}), \\ g_{i-1}g_{i-1}^Rg_{i-1} & \text{if } (g_i, a_{i+1}) = (g_{i+1}^r, g_{i+1}^{ra}); \end{cases}$$

Since $g_{i-1}g_{i-1}^Lg_{i-1} \approx g_{i-1}$ and $g_{i-1}g_{i-1}^Rg_{i-1} \approx g_{i-1}$ by Lemma 3.1.(*i*), we conclude that $g_{i-1}a_ig_ia_{i+1}g_{i+1} \approx g_{i-1}$ and so $u \approx v$. If $h_i \in \mathbf{G}^5$, then

 $g_{i-1}a_ih_ia_{i+1}g_{i+1} = h_i^r \cdot h_i \cdot h_i^l$ and $g_{i-1}a_ig_ia_{i+1}g_{i+1} = h_i^r \cdot h_i^c \cdot h_i^l$.

By definition of ρ_s , we conclude that $g_{i-1}a_ih_ia_{i+1}g_{i+1} \approx g_{i-1}a_ig_ia_{i+1}g_{i+1}$; whence $u \approx v$ also in this case.

We will denote by $\xrightarrow{*}$ the reflexive and transitive closure of \rightarrow . Due to the previous result, we know that $\xrightarrow{*}$ is contained in ρ . Thus, if u is a landscape, then

$$U(u) = \{ v \in \mathbf{G}^+ \mid u \xrightarrow{*} v \} \subseteq [u].$$

If we assume that u has length 2n + 1 and height i, then the landscapes from U(u) all have length at most 2n + 1 and height at most $j = \lfloor i + n/2 \rfloor$. They all have in common with u its first and last letters. If we look to the number of rivers, we conclude they all must have at most $\lfloor n/2 \rfloor$ rivers each. Hence, for each $v \in U(u)$, we record the number of rivers of v in a j-tuple $r(v) = (r^1(v), \cdots, r^j(v))$, where $r^k(v)$ denotes the number of rivers of v of height k - 1. The sum of all entries of r(v) gives us the number of rivers of v and so is at most $\lfloor n/2 \rfloor$.

Let $R(u) = \{r(v) \mid v \in U(u)\}$. The set R(u) is obviously finite. We order the elements of R(u) using the lexicographic order. If $v \in U(u)$ and $v \to v_1$ by uplifting a river of height k, then $v_1 \in U(u)$,

$$r^{k}(v_{1}) = r^{k}(v) - 1, \ r^{k+1}(v) \le r^{k+1}(v_{1}) \le r^{k+1}(v) + 2 \text{ and } r^{k_{1}}(v_{1}) = r^{k_{1}}(v)$$

for all other k_1 ; whence $r(v_1) < r(v)$. Therefore, we cannot apply upliftings of rivers indefinitely to u. We must stop only when we get a landscape with no river, that is, a landscape from $L_2^+ \cup L_1 \cup G'$. Note that these landscapes give us *j*-tuples with all entries equal to 0. Thus R(u) must always contain the *j*-tuple with all entries equal to 0, and this *j*-tuple is obviously the smallest element of R(u).

It is also easy to see that the uplifting of rivers is a commutative operation. In other words, if g_i and g_j are two rivers of $u \in L$, then the uplifting of g_i followed by the uplifting of g_j gives the same landscape as the uplifting of g_j followed by the uplifting of g_i . Hence, the uplifting of rivers is a system of rules commonly known as a noetherian locally confluent system of rules. An important property of this kind of systems is that, independently of the order of the rules that we choose to apply to an element, we must always stop after a finite number of steps with the same 'reduced' element (see [5]). In the case considered here, this means that, independently of the order of upliftings of rivers that we choose to apply to $u \in L$, we will always end up with the same landscape from U(u) with no rivers. In particular, U(u) has a unique landscape with no rivers. We designate it by $\beta_2(u)$. Note further that $\beta_2(u) = \beta_2(v)$ if $u \in L$ and $u \xrightarrow{*} v$.

Corollary 3.4. $\beta_2(u) \approx u$ for all $u \in L$.

Proof. This corollary is an obvious consequence of Lemma 3.3.

Note that the set of all mountain ranges is closed for $\stackrel{*}{\rightarrow}$, that is, $v \in MR$ if $u \in MR$ and $u \stackrel{*}{\rightarrow} v$. Hence, $U(u) \subseteq MR$ if $u \in MR$. Since the only mountain ranges with no rivers are the mountains, the landscape $\beta_2(u)$ must be a mountain if u is a mountain range. Thus, if we set $\beta(v) = \beta_2(\beta_1(v))$ for any word $v \in G^+$, then $\beta(v) \in M^1$. The previous corollary and Lemma 3.2 also allow us to conclude that $\beta(v) \approx v$. We leave this observation registered in the next corollary for future reference.

Corollary 3.5. $\beta(v) \approx v$ for all $v \in G^+$.

Let us take a pause and look more carefully to the structure of the mountains. First, they are words $u = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n}$ of length 4n + 1 $(n \in \mathbb{N}_0)$, $g_0 = g_{2n} = 1$, and peak g_n . If $n \geq 1$, then also $g_1 = g_{2n-1} = g_{xx'} \in \mathcal{G}_e$. In fact, if g_i is the highest letter of $\lambda_l(u)$ belonging to \mathcal{G}_e , then the subsequence $g_0 g_1 \cdots g_i$ is completely determined by $g_i: g_k = g_i^{l^{i-k}} = g_i^{r^{i-k}} \in \mathcal{G}_e$; whence $g_0 g_1 \cdots g_i \in \mathcal{G}_e^+$. In contrast, the subsequence $a_1 \cdots a_i$ may vary: a_k can be any anchor from $\{1, x\}$ if k even, or from $\{1, x'\}$ if k odd.

If we look now to the upper part of the uphill $\lambda_l(u)$ (above g_i), quite the opposite occurs. Each g_{k-1} is either the left or the right entry of g_k , for $i < k \leq n$. However, the side of g_{k-1} inside g_k completely determines a_k : a_k is the anchor of g_k corresponding to the side of g_{k-1} . Therefore, while the (possible empty) subsequence $g_i \cdots g_n$ may vary (although the peak g_n

is fixed), the subsequence $a_{i+1} \cdots a_n$ is completely determined by $g_i \cdots g_n$. Note also that $g_{i+1} \cdots g_n \in \mathbf{G}_d^+$ if it is not empty.

As we saw, the uphill $\lambda_l(u)$ is divided into two sections with very distinct properties. The same conclusions can be drawn for the downhill $\lambda_r(u)$. If g_j is the first letter from G_e in $\lambda_r(u)$, then $g_k = g_j^{l^{k-j}} = g_k^{r^{k-j}} \in G_e$ and g_j fixes the subsequence $g_j \cdots g_{2n} \in G_e^+$. In contrast, for each k > j, $a_k \in \{1, x\}$ if k even, or $a_k \in \{1, x'\}$ if k odd; whence the subsequence $a_{j+1} \cdots a_{2n}$ does not depend of g_j . In the upper part of $\lambda_r(u)$, each g_k can be either the left or the right entry of g_{k-1} , for $n < k \leq j$. However, for $n < k \leq j$, a_k is the inverse anchor of the anchor of g_k determined by the side of g_{k-1} .

From the conclusions described above, it is now obvious that the number of uphills and downhills between 1 and some $g \in G_n$ is 2^n in both cases. We leave this conclusion registered in the following corollary for future reference.

Corollary 3.6. If $g \in G_n$, then there are 2^n uphills from 1 to g and 2^n downhills from g to 1.

3.6. The word problem for $\langle G, R \rangle$. In the next sequence of results we show that $\beta(u)$ completely characterizes the ρ -class of $u \in G^+$. We will prove that [u] = [v] if and only if $\beta(u) = \beta(v)$, for all $u, v \in G^+$. This result solves the word problem for $\langle G, R \rangle$: we just need to compute both $\beta(u)$ and $\beta(v)$ and check if we get the same mountain to know if [u] = [v]. So, let

$$\rho_1 = \{(u, v) \in \mathbf{G}^+ \times \mathbf{G}^+ \mid \beta(u) = \beta(v)\}.$$

We want to show that $\rho = \rho_1$, but we already know that $\rho_1 \subseteq \rho$ by Corollary 3.5. The next couple of results show us that $\rho_e \subseteq \rho_1$.

Lemma 3.7. Let $g \in G'$. Then:

(i)
$$\beta_1(xx') \stackrel{*}{\to} \beta_1(g_{xx'}), \ \beta_1(xx'x) \stackrel{*}{\to} \beta_1(x) \text{ and } \beta_1(x'xx') \stackrel{*}{\to} \beta_1(x').$$

(ii) $\lambda_r(g) * \lambda_l(g) \stackrel{*}{\to} g \text{ and } \beta_1(g^2) \stackrel{*}{\to} \beta_1(g).$

Proof. (i). Clearly

$$\beta_1(xx') = \beta_1(x) * \beta_1(x') = 11g_{xx'}x1x'g_{xx'}11 \xrightarrow{1} 11g_{xx'}11 = \beta_1(g_{xx'}).$$

Now,

$$\beta_1(xx'x) \to \beta_1(g_{xx'}) * \beta_1(x) = 11g_{xx'}111g_{xx'}x1 \xrightarrow{1} 11g_{xx'}x1 = \beta_1(x) \,.$$

We show $\beta_1(x'xx') \xrightarrow{*} \beta_1(x')$ similarly.

(*ii*). The first part of (*ii*) is proved by induction on v(g). Clearly $\lambda_r(1) * \lambda_l(1) = 1$ and $\lambda_r(g_{xx'}) * \lambda_l(g_{xx'}) = g_{xx'} \cdot 111g_{xx'} \xrightarrow{1} g_{xx'}$. Let $g \in G_i$ for $i \ge 2$ and assume that $\lambda_r(h) * \lambda_l(h) \xrightarrow{*} h$ for all $h \in G_j$ with j < i. Since

$$\lambda_r(g) * \lambda_l(g) = g(g^{ra})' g^r (g^{raa})' \big(\lambda_r(g^c) * \lambda_l(g^c)\big) g^{laa} g^l g^{la} g,$$

we conclude that

$$\lambda_r(g) * \lambda_l(g) \xrightarrow{*} g(g^{ra})'g^r(g^{raa})'g^c g^{laa}g^l g^{la}g$$

by the induction hypothesis. But

 $g(g^{ra})'g^r(g^{r_aa})'g^cg^{l_aa}g^lg^{la}g \xrightarrow{g^c} g(g^{ra})'g^rg^{ra}g(g^{la})'g^lg^{la}g \xrightarrow{g^r,g^l} g.$

We have shown that $\lambda_r(g) * \lambda_l(g) \xrightarrow{*} g$ and proved the first part of (*ii*). Now,

$$\beta_1(g^2) = \lambda_l(g) * \lambda_r(g) * \lambda_l(g) * \lambda_r(g) \xrightarrow{*} \lambda_l(g) * g * \lambda_r(g) = \beta_1(g),$$

and we have proved the second part of (ii) also.

Corollary 3.8. $\rho_e \subseteq \rho_1$.

Proof. The pairs (xx'x, x), (x'xx', x'), $(xx', g_{xx'})$ and (g^2, g) , for $g \in G'$, belong to ρ_1 by Lemma 3.7. Then $\rho_e \subseteq \rho_1$ since it is also obvious that $\beta(g) = \beta(g1) = \beta(1g)$.

The next result contains technical details needed to prove that $\rho_s \subseteq \rho_1$.

Lemma 3.9. Let
$$g \in G'$$
 and let $u = g_0 a_1 g_1 \cdots a_n g_n \in L$. Then:

(i) If $v(g) \geq 2$ and $s \in \{l, r\}$, then $\lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) \xrightarrow{*} g^s g^{sa}g$ and $\lambda_r(g) * \beta_1((g^{sa})') * \lambda_l(g^s) \xrightarrow{*} g(g^{sa})'g^s$. (ii) $\beta_1(u) \xrightarrow{*} \lambda_l(g_0) * u * \lambda_r(g_n)$. (iii) $\beta(u) = \beta_2(u)$ if $u \in MR$. (iv) $\beta(u) = u$ if $u \in M^1$.

Proof. (i). We use induction again on v(g) to prove (i). We need, in fact, to prove both statements of (i) simultaneously by induction on v(g). For v(g) = 2, recall first the two elements of $G_2 = G_{2,e}$ described on page 7. Then $g^s = g_{xx'}$, $\lambda_r(g^s) = g_{xx'}$ 11 and

$$\lambda_l(g) = \begin{cases} 11g_{xx'}1g_{2,e,1} & \text{if } g = g_{2,e,1}, \\ 1x'g_{xx'}xg_{2,e,2} & \text{if } g = g_{2,e,2}. \end{cases}$$

Consider first the case s = l. If $g = g_{2,e,1}$, then $\beta_1(g^{sa}) = \beta_1(1) = 1$ and

$$\lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) = g_{xx'} 111g_{xx'} 1g \xrightarrow{1} g_{xx'} 1g = g^s g^{sa} g$$

If $g = g_{2,e,2}$, then $\beta_1(g^{sa}) = \beta_1(x) = 11g_{xx'} x1$ and

 $\lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) = g_{xx'} 111 g_{xx'} x 1 x' g_{xx'} xg \xrightarrow{1,1} g_{xx'} xg = g^s g^{sa} g.$

Now, consider the case s = r. If $g = g_{2,e,1}$, then $\beta_1(g^{sa}) = \beta_1(x) = 11g_{xx'}x_1^{a}$ and

$$\begin{array}{rcl} \lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) &=& g_{xx'} 111 g_{xx'} x 11 g_{xx'} 1g \xrightarrow{1} g_{xx'} x 11 g_{xx'} 1g \xrightarrow{1} \\ &\xrightarrow{1} g_{xx'} xg 1 g_{xx'} 1g \xrightarrow{g_{xx'}} g_{xx'} xg &=& g^s g^{sa} g \,. \end{array}$$

If $g = g_{2,e,2}$, then $\beta_1(g^{sa}) = \beta_1(1) = 1$ and

$$\begin{array}{rcl} \lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) &=& g_{xx'} 11x' g_{xx'} xg \xrightarrow{1} \\ & \stackrel{1}{\to} & g_{xx'} 1gx' g_{xx'} xg \xrightarrow{g_{xx'}} & g_{xx'} 1g &=& g^s g^{sa} g xg^{sa} g g_{xx'} gg_{xx'} gg_$$

We have proved the statement $\lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) \xrightarrow{*} g^s g^{sa}g$ for $\upsilon(g) = 2$. Similarly, we prove the statement $\lambda_r(g) * \beta_1((g^{sa})') * \lambda_l(g^s) \xrightarrow{*} g(g^{sa})'g^s$ for $\upsilon(g) = 2$.

Let $g \in G_i$ with i > 2, and assume that $\lambda_r(h^t) * \beta_1(h^{ta}) * \lambda_l(h) \xrightarrow{*} h^t h^{ta} h$ and $\lambda_r(h) * \beta_1((h^{ta})') * \lambda_l(h^t) \xrightarrow{*} h(h^{ta})'h^t$ for all $h \in G_j$ with $2 \le j < i$ and $t \in \{l, r\}$. Let $s \in \{l, r\}$ and recall that $g^{sa} = (g^{saa})'$ and $g^{sa} = g^c$. Then

$$\begin{split} \lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) &= \lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g^c) g^{l_a a} g^l g^{l a} g \\ &= \lambda_r(g^s) * \beta_1((g^{s_a a})') * \lambda_l(g^c) g^{l_a a} g^l g^{l a} g \\ &\stackrel{*}{\to} g^s(g^{s_a a})' g^c g^{l_a a} g^l g^{l a} g \,, \end{split}$$

where $\stackrel{*}{\rightarrow}$ follows by the induction hypothesis. Note now that if s = l, then

$$g^{s}(g^{s_{a}a})'g^{c}g^{l_{a}a}g^{l}g^{la}g = g^{l}(g^{l_{a}a})'g^{c}g^{l_{a}a}g^{l}g^{la}g \xrightarrow{g^{c}} g^{l}g^{la}g = g^{s}g^{sa}g;$$

and if
$$s = r$$
, then

$$g^{s}(g^{s_{a}a})'g^{c}g^{l_{a}a}g^{l}g^{la}g = g^{r}(g^{r_{a}a})'g^{c}g^{l_{a}a}g^{l}g^{la}g$$
$$\xrightarrow{g^{c}} g^{r}(g^{r_{a}a})'gg^{l_{a}a}g^{l}g^{la}g$$
$$= g^{r}g^{ra}g(g^{la})'g^{l}g^{la}g$$
$$\xrightarrow{g^{l}} g^{r}g^{ra}g = g^{s}g^{sa}g.$$

We have proved that $\lambda_r(g^s) * \beta_1(g^{sa}) * \lambda_l(g) \xrightarrow{*} g^s g^{sa}g$. The proof of $\lambda_r(g) * \beta_1((g^{sa})') * \lambda_l(g^s) \xrightarrow{*} g(g^{sa})'g^s$ is similar, and so (i) is proved by mathematical induction.

(*ii*). For $i \in \{0, \dots, n\}$, let $u_i = g_0 a_1 g_1 \cdots a_i g_i$. Thus all u_i are prefixes of u and $u = u_n$. Consequently, (*ii*) becomes proved once we show that $\beta_1(u_i) \xrightarrow{*} \lambda_l(g_0) * u_i * \lambda_r(g_i)$ by induction on i. Clearly

$$\beta_1(u_0) = \lambda_l(g_0) * u_0 * \lambda_r(g_0)$$

since $u_0 = g_0$. Assume that $\beta_1(u_{i-1}) \xrightarrow{*} \lambda_l(g_0) * u_{i-1} * \lambda_r(g_{i-1})$. By definition of landscape and (i), we have also

$$\lambda_r(g_{i-1}) * \beta_1(a_i) * \lambda_l(g_i) \xrightarrow{*} g_{i-1}a_ig_i.$$

Now, using the induction hypothesis, we obtain

$$\begin{split} \beta_1(u_i) &= \beta_1(u_{i-1}) * \beta_1(a_i) * \beta_1(g_i) \\ &\stackrel{*}{\to} \lambda_l(g_0) * u_{i-1} * \lambda_r(g_{i-1}) * \beta_1(a_i) * \lambda_l(g_i) * \lambda_r(g_i) \\ &\stackrel{*}{\to} \lambda_l(g_0) * u_{i-1} * (g_{i-1}a_ig_i) * \lambda_r(g_i) = \lambda_l(g_0) * u_i * \lambda_r(g_i) \end{split}$$

We have finished the proof of (ii).

(*iii*) follows easily from (*ii*) since $g_0 = 1 = g_n$ if $u \in MR$:

$$\beta(u) = \beta_2(\beta_1(u)) = \beta_2(\lambda_l(g_0) * u * \lambda_r(g_n)) = \beta_2(u);$$

and (iv) is obvious from (iii).

Proposition 3.10. Let $g \in G^5$ with $v(g) \ge 2$ and $u, v \in G^+$.

(i) $\beta(g) = \beta(g^c g^L g) = \beta(gg^R g^c)$ and $\beta(g^r \cdot g^c \cdot g^l) = \beta(g^r \cdot g \cdot g^l)$. (ii) $\beta(u) = \beta(v)$ if and only if [u] = [v].

Proof. (i). By Lemma 3.9.(ii),

 $\beta_1(g^cg^Lg) \xrightarrow{*} \lambda_l(g^c) * (g^cg^Lg) * \lambda_r(g) = \lambda_l(g) * \lambda_r(g) = \beta_1(g) \,.$

Thus $\beta(g^c g^L g) = \beta(g)$. We have also $\beta(gg^R g^c) = \beta(g)$ by duality. Again by Lemma 3.9.(*ii*), we have $\beta_1(g^r \cdot g \cdot g^l) \xrightarrow{*} \lambda_l(g^r) * (g^r \cdot g \cdot g^l) * \lambda_r(g^l)$ and

$$\beta_1(g^r \cdot g^c \cdot g^l) \xrightarrow{*} \lambda_l(g^r) * (g^r(g^{r_aa})'g^c g^{l_aa}g^l) * \lambda_r(g^l)$$
$$\xrightarrow{g^c} \lambda_l(g^r) * (g^r(g^{r_aa})'gg^{l_aa}g^l) * \lambda_r(g^l)$$
$$= \lambda_l(g^r) * (g^r g^{ra}g(g^{la})'g^l) * \lambda_r(g^l)$$
$$= \lambda_l(g^r) * (g^r \cdot g \cdot g^l) * \lambda_r(g^l).$$

Thereby $\beta(g^r \cdot g^c \cdot g^l) = \beta(g^r \cdot g \cdot g^l).$

(*ii*). We already know that [u] = [v] if $\beta(u) = \beta(v)$ by Corollary 3.5. From Corollary 3.8, we have also $\rho_e \subseteq \rho_1$. Now, we proved in (*i*) that $\rho_s \subseteq \rho_1$; whence $R \subseteq \rho_1$. To finish the proof of (*ii*), we just need to conclude that ρ_1 is a congruence on G⁺. But this is obvious since $\beta(u_1u_2) = \beta(\beta(u_1)u_2) = \beta(u_1\beta(u_2))$.

Although G and $\rho_e \cup \rho_s$ are infinite sets, Proposition 3.10.(*ii*) gives us a solution for the word problem for F_1^1 as explained before: to see if $u \approx v$, we just need to compute both $\beta(u)$ and $\beta(v)$, and check if we get the same mountain. Thus:

Corollary 3.11. The word problem for F_1^1 is decidable.

Proposition 3.10.(*ii*) tells us also that each ρ -class [u] has a unique mountain, namely $\beta(u)$. We will call $\beta(u)$ the *canonical form* of $u \in G^+$. We introduce the operation \odot on both M^1 and M as follows:

$$u_1 \odot u_2 = \beta(u_1 * u_2) = \beta_2(u_1 * u_2),$$

for any $u_1, u_2 \in \mathbf{M}^1$.

Proposition 3.12. (M^1, \odot) and (M, \odot) are models for F_1^1 and F_1 , respectively. Moreover, if $u, v, w \in M^1$ are such that $w = u \odot v$, then

- (i) $\lambda_l(u)$ is a prefix of $\lambda_l(w)$, while $\lambda_r(v)$ is a suffix of $\lambda_r(w)$.
- (ii) $v(w) \ge \max\{v(u), v(v)\}$, and v(w) = v(u) [v(w) = v(v)] if and only if $\kappa(w) = \kappa(u)$ $[\kappa(w) = \kappa(v)]$.

Proof. The first part is just a consequence of Proposition 3.10.(*ii*), while (*ii*) follows obviously from (*i*). Hence, we only need to prove (*i*). But notice that, by definition of uplifting of rivers, if $u_1 \rightarrow u_2$ for a mountain range u_1 , then $\lambda_l(u_1)$ is a prefix of $\lambda_l(u_2)$ and $\lambda_r(u_1)$ is a suffix of $\lambda_r(u_2)$. Applying several times the previous observation, we conclude that $\lambda_l(u) = \lambda_l(u * v)$

is a prefix of $\lambda_l(\beta(u * v)) = \lambda_l(u \odot v)$, while $\lambda_r(v) = \lambda_r(u * v)$ is a suffix of $\lambda_r(\beta(u * v)) = \lambda_r(u \odot v)$.

4. The semigroup F_1

In the previous section we constructed F_1 . In this section we prove that F_1 is a regular semigroup weakly generated by [x]. We begin by showing that F_1 is regular. To prove that F_1 is weakly generated by [x], we need first to describe the Green's relations on F_1 and then prove that [g] is the only element in $S([g^Rg^c], [g^cg^L])$. This latter fact will imply that any regular subsemigroup of F_1 containing [x], will contain also G^5 . Using this information, we will conclude that F_1 is weakly generated by [x].

Let $u = g_0 a_1 g_1 \cdots g_{n-1} a_n g_n$ be a landscape. Note that $g_n a_n g_{n-1} \cdots g_1 a_1 g_0$ is not a landscape in general but $g_n a'_n g_{n-1} \cdots g_1 a'_1 g_0$ is. We define the *reverse* of u as the landscape $\bar{u} = g_n a'_n g_{n-1} \cdots g_1 a'_1 g_0$. Clearly the reverse of \bar{u} is uagain.

Lemma 4.1. If $u = g_0 a_1 g_1 \cdots a_n g_n \in L_1^-$, then $u \overline{u} \approx u * \overline{u} \approx g_0$.

Proof. Note that $u * \overline{u}$ is a valley with river g_n and clearly $u\overline{u} \approx u * \overline{u}$ since $[g_n]$ is an idempotent of F_1^1 . Let $i \in \{1, \dots, n\}$. Then $g_i = g_{i-1}^{s_i}$ for some $s_i \in \{l, r\}$ since $u \in L_1^-$. In fact, we know also that $a_i = (g_{i-1}^{s_i a})'$. Hence

$$g_{i-1}a_ig_ia'_ig_{i-1} = g_{i-1} \cdot g_{i-1}^{s_i} \cdot g_{i-1} \approx g_{i-1}$$

by Lemma 3.1.(i). Now, applying several times this observation, we obtain

$$\begin{split} u\bar{u} &\approx u * \bar{u} = g_0 a_1 \cdots g_{n-2} a_{n-1} g_{n-1} a_n g_n a'_n g_{n-1} a'_{n-1} g_{n-2} \cdots a'_1 g_0 \\ &\approx g_0 a_1 \cdots g_{n-2} a_{n-1} g_{n-1} a'_{n-1} g_{n-2} \cdots a'_1 g_0 \\ &\vdots \\ &\approx g_0 a_1 g_1 a'_1 g_0 \approx g_0 \,. \end{split}$$

We have proved that $u\bar{u} \approx u * \bar{u} \approx g_0$ as wanted.

Next, we prove that F_1^1 is a regular monoid.

Proposition 4.2. The monoid F_1^1 is regular and $[\overline{u}]$ is an inverse of [u] for all $u \in L_2^+$.

Proof. By Corollary 3.5, $v \approx \beta(v)$ for all $v \in G^+$. Since $\beta(v) \in M^1 \subseteq L_2^+ \cup \{1\}$, it is enough to prove the second part of this proposition to conclude also the first. Let $u \in L_2^+$ with peak $g \in G^5$. Then $u = u_1 * u_2$ for some $u_1 \in L_1^+$ and $u_2 \in L_1^-$ such that $\tau(u_1) = g = \sigma(u_2)$. By Lemma 4.1, $u_2 \bar{u}_2 \approx g \approx \bar{u}_1 u_1$ (note that $\bar{u}_1 \in L_1^-$ and its reverse is u_1). Hence

$$uuu = u_1 * u_2 u_2 * u_1 u_1 * u_2 \approx u_1 * g * g * u_2 = u_1 * u_2 = u_1$$

We show similarly that $\bar{u}u\bar{u} \approx \bar{u}$. Therefore, $[\bar{u}]$ is an inverse of [u].

The ground $\epsilon(g)$ of a letter $g \in G'$ is defined recursively as follows:

$$\epsilon(1) = \{1\}$$
 and $\epsilon(g) = \epsilon(g^l) \cup \{g\} \cup \epsilon(g^r)$

if $v(g) \geq 1$. Thus $v(g_1) < v(g)$ for any $g_1 \in \epsilon(g) \setminus \{g\}$. We define the relation \preceq on G' by setting $h \preceq g$ if $h \in \epsilon(g)$. Due to statement (i) of the following result, \preceq is a partial order on G'. We extend the notion of ground to any landscape by setting, for $u = g_0 a_1 g_1 \cdots a_n g_n \in L$, $\epsilon(u) = \bigcup_{i=0}^n \epsilon(g_i)$. Again by statement (i) of the following result, $\epsilon(u)$ is the union of the grounds of its ridges, and the highest letters of $\epsilon(u)$ are the peaks of u. In particular, $\epsilon(u) = \epsilon(\kappa(u))$ if u is a mountain.

Lemma 4.3. Let $g, h \in G'$ and $u, v, w \in M^1$ such that $w = u \odot v$. Then:

- (i) $h \in \epsilon(g)$ if and only if $\epsilon(h) \subseteq \epsilon(g)$.
- (ii) If $h \in \epsilon(g) \setminus \{g\}$, then there exists $u = g_0 a_1 g_1 \cdots a_n g_n \in L_1^+$ such that n = v(g) v(h), $g_i \in \epsilon(g)$ for all $0 \le i \le n$, $g_0 = h$ and $g_n = g$. (iii) $\epsilon(u) \cup \epsilon(v) \subseteq \epsilon(w)$.

Proof. (i). Of course, we only need to prove that $\epsilon(h) \subseteq \epsilon(g)$ if $h \in \epsilon(g)$. This statement is proved by induction on v(g), and it is trivially true for v(g) = 0, that is, for g = 1. Assume that $\epsilon(h_1) \subseteq \epsilon(g_1)$ for all $g_1 \in G'$ such that $v(g_1) < v(g)$ and all $h_1 \in \epsilon(g_1)$. If $h \in \epsilon(g)$, then either h = g, or $h \in \epsilon(g^l)$, or $h \in \epsilon(g^r)$. In the latter two cases, we have either $\epsilon(h) \subseteq \epsilon(g^l)$ or $\epsilon(h) \subseteq \epsilon(g^r)$, respectively, by the induction hypothesis. Hence,

$$\epsilon(h) \subseteq \epsilon(g^r) \cup \{g\} \cup \epsilon(g^l) = \epsilon(g)$$

as wanted.

(*ii*). Let $h \in \epsilon(g) \setminus \{g\}$. We prove (*ii*) by induction on n = v(g) - v(h). If n = 1, then $h = g^s$ for some $s \in \{l, r\}$. Hence $u = hg^{sa}g$ is an uphill satisfying the conditions of (*ii*). Assume now that $n \ge 2$. Then $h \in \epsilon(g^s)$ for some $s \in \{l, r\}$. Without loss of generality, we assume that $h \in \epsilon(g^l)$. Since $v(g^l) - v(h) = n - 1$, there exists $g_0 a_1 g_1 \cdots a_{n-1} g_{n-1} \in L_1^+$ such that all $g_i \in \epsilon(g^l)$ for $0 \le i \le n - 1$, $g_0 = h$ and $g_{n-1} = g^l$, by the induction hypothesis. Clearly $u = g_0 a_1 g_1 \cdots a_{n-1} g_{n-1} g^{la}g$ is now an uphill satisfying the conditions stated in (*ii*).

(*iii*). Note that $\kappa(u), \kappa(v) \in \epsilon(w)$ since $\lambda_l(u)$ is a prefix of $\lambda_l(w)$ and $\lambda_r(v)$ is a suffix of $\lambda_r(w)$ (Proposition 3.12.(*i*)). (*iii*) follows now from (*i*).

We describe next the partial orders $\leq_{\mathscr{R}}, \leq_{\mathscr{L}}$ and $\leq_{\mathscr{I}}$ on (M^1, \odot) .

Proposition 4.4. Let $u, v \in M^1$. Then:

- (i) $u \leq_{\mathscr{R}} v$ if and only if $\lambda_l(v)$ is a prefix of $\lambda_l(u)$.
- (ii) $u \leq_{\mathscr{L}} v$ if and only if $\lambda_r(v)$ is a suffix of $\lambda_r(u)$.
- (iii) $u \leq \mathcal{J} v$ if and only if $\kappa(v) \preceq \kappa(u)$.

Proof. We prove only (i) and (iii) since (ii) is the dual of (i).

(i). Assume that $u \leq_{\mathscr{R}} v$. Thus $u = v \odot w$ for some $w \in M^1$. By Proposition 3.12.(i), $\lambda_l(v)$ is a prefix of $\lambda_l(u)$. Assume now that $\lambda_l(v)$ is a

prefix $\lambda_l(u)$. Then $u = \lambda_l(v) * u_1$ for some landscape u_1 with $\tau(\lambda_l(v)) = \kappa(v) = \sigma(u_1)$. Let $v_1 = \lambda_r(v)$ and consider $w = \overleftarrow{v_1} * u_1$. Clearly $w \in M^1$. Further, $v_1 * \overleftarrow{v_1} \xrightarrow{*} \kappa(v)$ by Lemma 4.1, and so

$$v * w = \lambda_l(v) * v_1 * \overleftarrow{v_1} * u_1 \xrightarrow{*} \lambda_l(v) * \kappa(v) * u_1 = \lambda_l(v) * u_1 = u,$$

that is, $u = v \odot w$ and $u \leq_{\mathscr{R}} v$.

(*iii*). If $u \leq \mathscr{J} v$, then $u = w_1 \odot v \odot w_2$ for two mountains w_1 and w_2 . By Lemma 4.3.(*iii*), $\epsilon(\kappa(v)) = \epsilon(v) \subseteq \epsilon(u) = \epsilon(\kappa(u))$ and $\kappa(v) \preceq \kappa(u)$. Conversely, if $\kappa(v) \preceq \kappa(u)$, then let u_1 be an uphill such that $\sigma(u_1) = \kappa(v)$ and $\tau(u_1) = \kappa(u)$, whose existence is guaranteed by Lemma 4.3.(*ii*). Observe that $w_1 = \lambda_l(u) * \dot{u_1} * \dot{\lambda_l(v)}$ and $w_2 = \lambda_r(v) * u_1 * \lambda_r(u)$ are well defined mountains. Further,

$$w_1 * v * w_2 = \lambda_l(u) * \overleftarrow{u_1} * (\overleftarrow{\lambda_l(v)} * \lambda_l(v)) * (\lambda_r(v) * \overleftarrow{\lambda_r(v)}) * u_1 * \lambda_r(u)$$

$$\xrightarrow{*} \lambda_l(u) * \overleftarrow{u_1} * \kappa(v) * \kappa(v) * u_1 * \lambda_r(u)$$

$$\xrightarrow{*} \lambda_l(u) * \kappa(u) * \lambda_r(u) = u,$$

and $w_1 \odot v \odot w_2 = u$. Thus $u \leq_{\mathscr{J}} v$.

The next two results are corollaries of the previous proposition. The first one is an obvious consequence. For the second one, we need to say something more about the \mathscr{H} and the \mathscr{D} relations.

Corollary 4.5. Let $u, v \in M^1$. Then:

- (i) v covers u for $\leq_{\mathscr{R}}$ if and only if $\lambda_l(u) = \lambda_l(v)a\kappa(u)$ for some $a \in A$.
- (ii) v covers u for $\leq_{\mathscr{L}}$ if and only if $\lambda_r(u) = \kappa(u) a \lambda_r(v)$ for some $a \in A$.
- (iii) v covers u for $\leq_{\mathscr{I}}$ if and only if $\kappa(v) \in \{(\kappa(u))^l, (\kappa(u))^r\}$.

Corollary 4.6. Let $u, v \in M^1$. Then:

- (i) $u \mathscr{R} v$ if and only if $\lambda_l(u) = \lambda_l(v)$.
- (ii) $u \mathscr{L} v$ if and only if $\lambda_r(u) = \lambda_r(v)$.
- (iii) $u \not j v$ if and only if $\kappa(u) = \kappa(v)$.
- (iv) $u \mathscr{H} v$ if and only if u = v.
- $(v) \mathcal{D} = \mathcal{J}.$

Proof. The first three statements follow from the corresponding statements of Proposition 4.4. Then (iv) is a consequence of (i) and (ii). So, we only need to prove that $\mathscr{J} \subseteq \mathscr{D}$. Assume that $u \not \mathscr{J} v$. Then $w = \lambda_l(u) * \lambda_r(v) \in M^1$ since $\kappa(u) = \kappa(v)$ by (iii). Now, by (i) and (ii), we conclude that $u \mathscr{R} w \mathscr{L} v$, whence $u \mathscr{D} v$.

The statements (iii) and (v) of the previous corollary tell us that the $\mathscr{D} = \mathscr{J}$ -classes of F_1^1 are in one-to-one correspondence with the elements of G', and that the set $\{[g] \mid g \in G'\}$ is a transversal (or cross-section) for the set of \mathscr{D} -classes of F_1^1 . The next result gives us the size of each \mathscr{R} , \mathscr{L} and \mathscr{D} -class of F_1^1 .

Corollary 4.7. If $g \in G_n$ for $n \ge 1$, then $|R_{[g]}| = 2^n = |L_{[g]}|$ and $|D_{[g]}| = 2^{2n}$. Further, $D_{[g]}$ has $2^n \mathscr{R}$ -classes and $2^n \mathscr{L}$ -classes.

Proof. We only need to prove that $|\mathbf{R}_{[g]}| = 2^n$ since $|\mathbf{L}_{[g]}| = 2^n$ follows by duality and the statements about \mathscr{D} follow from the statements about \mathscr{R} and \mathscr{L} and from Corollary 4.6.(*iv*). But Corollary 4.6.(*i*) tells us that the size of $\mathbf{R}_{[g]}$ is equal to the number of downhills from g to 1, and this number is 2^n due to Corollary 3.6. Hence, $|\mathbf{R}_{[g]}| = 2^n$.

We know that $[g] \in S([g^R g^c], [g^c g^L])$ by Lemma 3.1.(vi). In the next result we prove that [g] is the only element of $S([g^R g^c], [g^c g^L])$ in F_1^1 .

Lemma 4.8. For each $g \in G^5$ with $v(g) \ge 2$, $S([g^R g^c], [g^c g^L]) = \{[g]\}$ in F_1^1 .

Proof. Let $u \in M^1$ be such that $[u] \in S([g^R g^c], [g^c g^L])$. Then $\kappa(u) = g$ by Corollary 4.6 since $[u] \mathscr{D}[g]$. Let v be the mountain $\lambda_l(g^c)g^{l_a a}\lambda_r(g^l)$. By Lemma 3.9.(*ii*), $\beta_1(g^c g^{l_a a}g^l) \xrightarrow{*} v$, and so $g^c g^{l_a a}g^l \approx v$. Hence

$$u \approx g^c g^L u \approx v g^{la} u \approx v * \beta_1(g^{la}) * u \approx v \odot \beta_1(g^{la}) \odot u \,,$$

and so $u = v \odot \beta_1(g^{la}) \odot u$. By Proposition 3.12.(*i*), $\lambda_l(v) = \lambda_l(g^c)g^{l_aa}g^l$ is a prefix of $\lambda_l(u)$. In a similar way, we conclude also that $g^r(g^{r_aa})'\lambda_r(g^c)$ is a suffix of $\lambda_r(u)$.

If $g \in G_d$, then $g^l \neq g^r$. Hence, for this case, $\lambda_l(g)$ is the only uphill from 1 to g with prefix $\lambda_l(g^c)g^{l_a a}g^l$ and $\lambda_r(g)$ is the only downhill from gto 1 with suffix $g^r(g^{r_a a})'\lambda_r(g^c)$. We have concluded that $\lambda_l(g) = \lambda_l(u)$ and $\lambda_r(g) = \lambda_r(u)$ since $\kappa(u) = g$. Thus, $u = \beta_1(g)$ and [u] = [g] if $g \in G_d$.

The case $g \in G_e$ is more complex since $g^l = g^r$, and thus there are two distinct uphills from 1 to g, namely $\lambda_l(g)$ and $\lambda_l(g^c)g^{l_aa}g^lg^{ra}g$. There are also two distinct downhills form g to 1, namely $\lambda_r(g)$ and $g(g^{l_a})'g^r(g^{r_aa})'\lambda_r(g^c)$. We have concluded that

$$u = \lambda_l(g^c)g^{l_a a}g^l a_1 g a'_2 g^r(g^{r_a a})' \lambda_r(g^c),$$

for $a_1, a_2 \in \{g^{ra}, g^{la}\}$. However, [u] is also an idempotent of F_1^1 . Hence,

$$\begin{split} u &\approx u^2 \;\;=\;\; \lambda_l(u) * \lambda_r(u) * \lambda_l(u) * \lambda_r(u) \\ &\stackrel{*}{\to}\;\; \lambda_l(u) * ga'_2 g^r(g^{r_a a})' g^c g^{l_a a} g^l a_1 g * \lambda_r(u) \\ &\stackrel{g^c}{\longrightarrow}\;\; \lambda_l(u) * ga'_2 g^r g^{r a} g(g^{l a})' g^l a_1 g * \lambda_r(u) \,. \end{split}$$

If $a_2 = g^{la}$, then $g_1 = (g, (g^{ra})', g^r, (g^{la})', g) \in \mathbf{G}_e$ and

$$ga_2'g^rg^{ra}g \xrightarrow{g^r} ga_2'g_1g^{ra}g$$

So, $g_1 \in \epsilon(\beta(u^2)) = \epsilon(u) = \epsilon(g)$ and $v(g_1) = v(g) + 1$, which is a contradiction. Therefore a_2 must be equal to g^{ra} . Similarly, we conclude also that $a_1 = g^{la}$. Consequently $\lambda_l(u) = \lambda_l(g)$ and $\lambda_r(u) = \lambda_r(g)$, that is, $u = \beta_1(g)$ and [u] = [g] if $g \in G_e$ also. We have proved the statement of this proposition.

We have now all the ingredients necessary to show that F_1 is weakly generated by [x].

Proposition 4.9. The regular semigroup F_1 is weakly generated by [x].

Proof. Let S be a regular subsemigroup of F_1 containing [x]. By Corollary 4.6,

$$D_{[x]} = \{ [x], [x'], [g_{xx'}], [x'x] \},\$$

and [x'] is the only inverse of [x]. Hence, S contains $D_{[x]}$. If $g \in G_2$, then $g^c = 1$ and [g] is the only element of $S([g^Rg^c], [g^cg^L]) = S([g^R], [g^L])$. Note that both $[g^R]$ and $[g^L]$ belong to S since $g^r = g^l = g_{xx'}$. Since S is regular, it must contain also an element from $S([g^R], [g^L])$, that is, it must contain [g]. Next, we show that S contains the set $\{[g] \mid g \in G^5\}$ by induction on v(g).

Let $g \in G_i$ for $i \geq 3$, and assume that [h] belongs to S for all $h \in G_j$ with j < i. Then $[g^l]$, $[g^r]$ and $[g^c]$ belong to S by the induction hypothesis (note that $g^c \neq 1$ because $v(g) \geq 3$). Since [x] and [x'] also belong to S, we conclude that $[g^R g^c] \in S$ and $[g^c g^L] \in S$. Again, we know that

$$S([g^Rg^c], [g^cg^L]) \cap S \neq \emptyset$$

because S is regular; whence $[g] \in S$ due to Lemma 4.8. We have shown that S contains

$$\{[x], [x']\} \cup \{[g] \mid g \in \mathbf{G}^5\}.$$

Since this latter set generates F_1 , we must have $S = F_1$. Therefore, F_1 is weakly generated by [x].

In this section we have already described the Green's relations on F_1 . We end it describing the idempotents, the inverses and the natural partial order on F_1 . A canyon is a valley w with $\sigma(w) = \tau(w)$ and a gorge is a canyon such that $w \xrightarrow{*} \sigma(w) = \tau(w)$. Thus $\beta_2(w) = \sigma(w)$ for any gorge w, and $w_1 \xrightarrow{*} \sigma(w)$ for any $w_1 \in G^+$ such that $w \xrightarrow{*} w_1$. Note however that w_1 may not be a gorge since we cannot guarantee that w_1 is even a valley. We will use gorges to characterize the idempotents, the inverses and the natural partial order, but this characterization will be just theoretical. For practical purposes, it is hard to identify the gorges, and consequently also the idempotents, the inverses and the natural partial order.

Proposition 4.10. Let $u, v \in M^1$.

- (i) u is an idempotent if and only if the canyon $\lambda_r(u) * \lambda_l(u)$ is a gorge.
- (ii) v is an inverse of u if and only if $\lambda_r(u) * \lambda_l(v)$ and $\lambda_r(v) * \lambda_l(u)$ are both gorges.
- (iii) v < u if and only if $\lambda_l(v) = \lambda_l(u) a_1 u_1$ and $\lambda_r(v) = u_2 a_2 \lambda_r(u)$ for some $u_1 \in L_1^+ \cup G^5$, $u_2 \in L_1^- \cup G^5$ and $a_1, a_2 \in A$ such that $u_2 a_2 \kappa(u) a_1 u_1$ is a gorge.

Proof. If $\lambda_r(u) * \lambda_l(u) \xrightarrow{*} \kappa(u)$, then

 $u^2 \approx \lambda_l(u) * \lambda_r(u) * \lambda_l(u) * \lambda_r(u) \xrightarrow{*} \lambda_l(u) * \kappa(u) * \lambda_r(u) = u$

and u is an idempotent. Conversely, if u is an idempotent, then $u^2 \xrightarrow{*} u$. By definition of uplifting of rivers, it is obvious that we must have $\lambda_r(u) * \lambda_l(u) \xrightarrow{*} \kappa(u)$. We have proved (*i*). The proof of (*ii*) is similar and as obvious as the proof of (*i*). Let us prove (*iii*) now.

Assume first that v < u. In particular $v \leq_{\mathscr{R}} u$ and $v \leq_{\mathscr{L}} u$. Since $u \neq v$,

$$\lambda_l(v) = \lambda_l(u) a_1 u_1$$
 and $\lambda_r(v) = u_2 a_2 \lambda_r(u)$,

for some $u_1 \in L_1^+ \cup G^5$, $u_2 \in L_1^- \cup G^5$ and $a_1, a_2 \in A$, by Proposition 4.4. But, there exists also another mountain $w \in M^1$ such that

$$w \in E(\mathbf{M}^1), \quad w \mathscr{L} v \quad \text{and} \quad v = u \odot w$$

as v < u. Hence, $\lambda_r(w) * \lambda_l(w)$ is a gorge by (i), $\lambda_r(w) = \lambda_r(v)$ by Corollary 4.6.(ii), and $\lambda_r(u) * \lambda_l(w) \xrightarrow{*} \kappa(u) a_1 u_1$. Then

$$\lambda_r(w) * \lambda_l(w) = \lambda_r(v) * \lambda_l(w) = u_2 a_2 \lambda_r(u) * \lambda_l(w) \xrightarrow{*} u_2 a_2 \kappa(u) a_1 u_1.$$

We can now conclude that $u_2 a_2 \kappa(u) a_1 u_1$ is a gorge because $\lambda_r(w) * \lambda_l(w)$ is a gorge too (note that $u_2 a_2 \kappa(u) a_1 u_1$ is a canyon).

Assume now that $\lambda_l(v) = \lambda_l(u) a_1 u_1$ and $\lambda_r(v) = u_2 a_2 \lambda_r(u)$ for some $u_1 \in L_1^+ \cup G^5$, $u_2 \in L_1^- \cup G^5$ and $a_1, a_2 \in A$ such that $u_2 a_2 \kappa(u) a_1 u_1$ is a gorge. In particular $u \neq v$. Note also that

$$w_r = \lambda_r(u) a_1 u_1 * \lambda_r(v)$$
 and $w_l = \lambda_l(v) * u_2 a_2 \lambda_l(u)$

are well defined mountains. In fact, they are both idempotents of M^1 . For example,

$$\lambda_r(v) * \lambda_r(u) a_1 u_1 = u_2 a_2 \lambda_r(u) * \lambda_r(u) a_1 u_1 \xrightarrow{*} u_2 a_2 \kappa(u) a_1 u_1;$$

thus $\lambda_r(v) * \lambda_r(u) a_1 u_1$ is a gorge and so w_r is an idempotent. Finally, observe that $u \odot w_r = v$ and $w_l \odot u = v$, whence v < u.

5. The universal property of F_1

In this section we prove that all regular semigroups weakly generated by x are homomorphic images of F_1 under a homomorphism that sends [x] into x. A crucial concept is the notion of skeleton. A *skeleton mapping* for a regular semigroup S is a mapping $\phi : G \to S^1$ satisfying

(i)
$$x\phi \in S, x'\phi \in V(x\phi), g_{xx'}\phi = (x\phi)(x'\phi) \text{ and } 1\phi \in E(S^1);$$

(ii) $(1\phi)(x\phi) = x\phi = (x\phi)(1\phi) \text{ and } (1\phi)(x'\phi) = x'\phi = (x'\phi)(1\phi);$
(iii) $g\phi \in S(g^{\phi,r}, g^{\phi,l})$ for all $g \in G_i$ with $i \ge 2$, where

$$g^{\phi,l} = (g^c \phi)((g^{la})'\phi)(g^l \phi)(g^{la}\phi)$$

and

$$g^{\phi,r} = ((g^{ra})'\phi)(g^r\phi)(g^{ra}\phi)(g^c\phi).$$

In this definition we are using the non-idempotent version of the definition of sandwich set since we cannot guarantee at this point that $g^{\phi,l}$ and $g^{\phi,r}$ are idempotents of S. However, after the following observations, we will prove that these elements are indeed idempotents of S. From that point on, we will consider here also the usual definition of sandwich set for idempotents.

It is easy to see that $(1\phi)(g\phi) = g\phi = (g\phi)(1\phi)$ for all $g \in G$ by induction on v(g). In fact, by (*iii*), for all $g \in G_i$ with $i \ge 2$, we just need to observe that

$$g\phi = g^{\phi,l}y(g\phi) = (g\phi)zg^{\phi,r}$$

for any $y \in V(g^{\phi,l})$ and $z \in V(g^{\phi,r})$, and then use the induction hypothesis. This same observation gives us that $(g^c\phi)(g\phi) = (g\phi) = (g\phi)(g^c\phi)$ since $g^c\phi$ is an idempotent of S. Further, (i) guarantees that $(G \setminus \{1\})\phi \subseteq S$. In this paper, a *skeleton* of S is the image of $G \setminus \{1\}$ under some skeleton mapping. Thus, except for $x\phi$ and $x'\phi$, which are mutually inverse elements with $g_{xx'}\phi = (x\phi)(x'\phi)$, all other elements of a skeleton are idempotents of S.

Lemma 5.1. Let S be a regular semigroup and $\phi : \mathbf{G} \to S^1$ be a skeleton mapping. Then $g^{\phi,l}$ and $g^{\phi,r}$ are idempotents of S for all $g \in \mathbf{G}_i$ with $i \geq 2$.

Proof. We need to prove that $g^{\phi,l}$ and $g^{\phi,r}$ are idempotents of S simultaneously by induction on i. For i = 2, note that $g^l = g_{xx'} = g^r$ and $g^c = 1$. Thus, we can easily check that $g^{\phi,l}$ and $g^{\phi,r}$ belong to the set $\{aa', a'a\}$ for $a = x\phi$ and $a' = x'\phi$; whence they are idempotents.

Assume now that i > 2. By definition of skeleton mapping,

$$g^l \phi \in S((g^l)^{\phi,r}, (g^l)^{\phi,l})$$

and we may assume that $(g^l)^{\phi,l}$ and $(g^l)^{\phi,r}$ are idempotents by the induction hypothesis. Thus $(g^l)^{\phi,l}(g^l\phi) = g^l\phi$ and $(g^l\phi)(g^l)^{\phi,r} = g^l\phi$. Note also that $g^c = g^{l^2}$ and $g^{la} = (g^{l^2a})'$, or $g^c = g^{lr}$ and $g^{la} = (g^{lra})'$. If $g^c = g^{l^2}$ and $g^{la} = (g^{l^2a})'$, then

$$\begin{aligned} (g^{l}\phi)(g^{la}\phi)(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) &= (g^{l}\phi)(g^{lc}\phi)(g^{la}\phi)(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) \\ &= (g^{l}\phi)(g^{lc}\phi)((g^{l^{2}a})'\phi)(g^{l^{2}}\phi)(g^{l^{2}a}\phi)(g^{l}\phi) \\ &= (g^{l}\phi)((g^{l})^{\phi,l})(g^{l}\phi) = g^{l}\phi. \end{aligned}$$

If $g^c = g^{lr}$ and $g^{la} = (g^{lra})'$, then

$$\begin{aligned} (g^{l}\phi)(g^{la}\phi)(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) &= (g^{l}\phi)(g^{la}\phi)(g^{c}\phi)((g^{la})'\phi)(g^{lc}\phi)(g^{l}\phi) \\ &= (g^{l}\phi)((g^{lra})'\phi)(g^{lr}\phi)(g^{lra}\phi)(g^{lc}\phi)(g^{l}\phi) \\ &= (g^{l}\phi)((g^{l})^{\phi,r})(g^{l}\phi) = g^{l}\phi \,. \end{aligned}$$

It is easy to see now that $g^{\phi,l} = (g^c \phi)((g^{la})'\phi)(g^l \phi)(g^{la}\phi)$ is an idempotent of S. The proof that $g^{\phi,r}$ is also an idempotent of S is similarly. \Box

If a is an element of a regular semigroup S, we can easily construct recursively a skeleton mapping such that $x\phi = a$. We can choose an inverse a'of a in S and define $x'\phi = a'$, $g_{xx'}\phi = aa'$ and $1\phi = 1 \in S^1$. Then, assuming that $g \in G_i$ with $i \ge 2$ and ϕ is defined for all $h \in G_j$ with j < i, we choose

$$y \in S(g^{\phi,r}, g^{\phi,l}),$$

and set $g\phi = y$. Note that $S(g^{\phi,r}, g^{\phi,l}) \neq \emptyset$ since S is regular. We cannot however guarantee that the skeleton mapping ϕ is unique. In terms of skeletons, each element $a \in S$ belongs to at least one skeleton of S, but it may belong to many distinct skeletons.

In the next result we show that every skeleton mapping for a regular semigroup S can be uniquely extended into a homomorphism from F_1^1 to S^1 .

Proposition 5.2. If $\phi : \mathbf{G} \to S^1$ is a skeleton mapping, then there is a unique (semigroup) homomorphism $\varphi : \mathbf{F}_1^1 \to S^1$ extending ϕ (that is, such that $g\phi = [g]\varphi$ for all $g \in \mathbf{G}$). Furthermore, $\langle \mathbf{G}\phi \rangle = \mathbf{F}_1^1\varphi$ is a regular subsemigroup of S^1 and a monoid with identity element 1ϕ .

Proof. Since G^+ is freely generated by G, let $\phi_1 : G^+ \to S^1$ be the only homomorphism that extends ϕ . By definition of skeleton mapping,

$$\rho_e \cup \left\{ (g^c g^L g, g), (g g^R g^c, g) | g \in \mathcal{G} \text{ with } v(g) \ge 2 \right\}$$

is contained in the kernel of ϕ_1 . To conclude that ρ is contained in the kernel of ϕ_1 , it is now enough to show that $(g^r \cdot g \cdot g^l, g^r \cdot g^c \cdot g^l)$ belongs to that kernel for all $g \in G_i$ with $i \ge 2$.

Due to space constrains, we denote by w and z the expressions

$$(g^l\phi)(g^{la}\phi)(g^c\phi)((g^{la})'\phi)(g^l\phi)$$
 and $(g^r\phi)(g^{ra}\phi)(g^c\phi)((g^{ra})'\phi)(g^r\phi)$,

respectively. In the proof of Proposition 5.1 we showed that $g^l \phi = w$. Similar arguments allow us to conclude that $g^r \phi = z$. We need also the equality $g\phi = (g^c \phi)(g\phi)(g^c \phi)$ that follows from an observation made prior to Lemma 5.1. So,

$$\begin{split} (g^{r} \cdot g \cdot g^{l})\phi_{1} &= (g^{r}\phi)(g^{ra}\phi)(g\phi)((g^{la})'\phi)(g^{l}\phi) \\ &= z \, (g^{ra}\phi)(g^{c}\phi)(g\phi)(g^{c}\phi)((g^{la})'\phi) \, w \\ &= (g^{r}\phi)(g^{ra}\phi)(g^{c}\phi)g^{\phi,r}(g\phi)g^{\phi,l}(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) \\ &= (g^{r}\phi)(g^{ra}\phi)(g^{c}\phi)g^{\phi,r}g^{\phi,l}(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) \\ &= z \, (g^{ra}\phi)(g^{c}\phi)((g^{la})'\phi) \, w \\ &= (g^{r}\phi)(g^{ra}\phi)(g^{c}\phi)((g^{la})'\phi)(g^{l}\phi) = (g^{r} \cdot g^{c} \cdot g^{l})\phi_{1} \,, \end{split}$$

where the fourth "=" sign is due to $g\phi \in S(g^{\phi,r}, g^{\phi,l})$.

So, we have shown that ρ is contained in the kernel of ϕ_1 . Thus, if $\varphi_1 : \mathbf{G}^+ \to \mathbf{F}_1^1$ denotes the natural quotient homomorphism, then there exists a homomorphism $\varphi : \mathbf{F}_1^1 \to S^1$ such that $\phi_1 = \varphi_1 \varphi$. Hence φ extends ϕ . If

 $\varphi': F_1^1 \to S^1$ is another homomorphism extending ϕ , then $\varphi_1 \varphi' = \phi_1 = \varphi_1 \varphi$. Consequently $\varphi' = \varphi$ since φ_1 is surjective. We proved there exists a unique homomorphism $\varphi: F_1^1 \to S^1$ extending ϕ .

Finally, the second part of this result follows immediately from the fact that F_1^1 is a regular monoid with identity element [1] (this property is inherited by homomorphic images) and it is generated by $\{[g] \mid g \in G\}$.

Note that, in the previous proposition, $F_1^1 \varphi$ may not be a submonoid of S^1 since $[1]\varphi$ may not be the identity of element of S^1 . Given a skeleton mapping ϕ , we can only guarantee that 1ϕ is an identity element for the subsemigroup $F_1^1 \varphi$. Also by the previous result, a skeleton of S generates always a regular subsemigroup of S. We leave this conclusion registered in the following corollary for future reference.

Corollary 5.3. If A is a skeleton of S, then $\langle A \rangle$ is a regular subsemigroup of S.

Let T be a regular subsemigroup of a (not necessarily regular) semigroup S. Next, we prove that if T is weakly generated by an element a, then there exists a homomorphism $\varphi : F_1 \to S$ such that $[x]\varphi = a$ and $F_1 \varphi = T$.

Corollary 5.4. Let T be a regular subsemigroup of a semigroup S. If T is weakly generated by a, then T is the image of F_1 under some homomorphism $\varphi: F_1 \to S$ such that $[x]\varphi = a$.

Proof. Let $\phi : \mathbf{G} \to T^1$ be a skeleton mapping such that $x\phi = a$. We have already observed that ϕ exists. Let $\varphi : \mathbf{F}_1^1 \to T^1$ be the unique homomorphism extending ϕ given by Proposition 5.2. Note that $\varphi_{|\mathbf{F}_1|}$ is a homomorphism from \mathbf{F}_1 to T. Thus $\mathbf{F}_1 \varphi$ is a regular subsemigroup of T containing a. Since T is weakly generated by a, we conclude that $\mathbf{F}_1 \varphi = T$. Finally, note that we can consider φ as a homomorphism from \mathbf{F}_1 to S, and $[x]\varphi = a$ as wanted.

The main result of this paper is now an obvious consequence of the previous corollary.

Theorem 5.5. All regular semigroups weakly generated by an element a are homomorphic images of F_1 under a homomorphism that sends [x] into a.

The reverse of this theorem is not true however, that is, there are homomorphic images of F_1 that are not weakly generated by the image of [x]. Next, we present one such example. This example is adapted from the one used in [6] to show that not all homomorphic images of FI_2 are weakly generated by two idempotents.

Example 1: We begin by defining a presentation $\langle X, R_1 \rangle$ for a semigroup S_1 . We set $X = \{x, x', g_1, g_2, h, 0\}$ and include in R_1 the relations corresponding to the information $e^0 = 0 = 0e$ for all $e \in X$, $x' \in V(x)$,

 $g_1 \in S(x'x, xx'), g_2 \in S(xx', x'x)$ and $h \in S(xg_2x', xg_1x')$. So 0 correspond to a zero element of S_1 as expected. Now, add to R_1 also the pairs (u, 0) for

 $u \in \{g_i g_j, g_i a g_j \mid \{i, j\} = \{1, 2\} \text{ and } a \in \{x, x'\}\} \cup \{x^2, (x')^2\}.$

The semigroup S_1 is the one obtained from F_1 by collapsing all \mathscr{D} -classes into the 0 element except for $D_{[x]}$, $D_{[g_{2,e,1}]}$, $D_{[g_{2,e,2}]}$ and $D_{[g_{3,d,2}]}$. In fact, the first set of relations identifies g_1 , g_2 and h with the elements $g_{2,e,1}$, $g_{2,e,2}$ and $g_{3,d,2}$, respectively, while the second set of relations collapses all other \mathscr{D} -classes into 0.

But the semigroup we are interested in is the semigroup S given by the presentation $\langle X, R \rangle$ where $R = R_1 \cup \{(g_1, g_1x'hxg_1), (g_2, g_2x'hxg_2)\}$. Since R contains R_1 , the semigroup S is also a homomorphic image of F_1 . The semigroup S is, in fact, obtained from S_1 by merging the \mathcal{D} -classes of g_1 and g_2 into distinct 'blocks' of the \mathcal{D} -class of h. We illustrate the "Egg-box" diagram of the \mathcal{D} -classes of S in Figure 5 (note that each congruence class is represented by one of its elements; we write also e = xx', f = x'x and $h_1 = x'hx$ due to space constrains).

x	e *
f *	x'

$g_1e *$	$g_1 x$	$g_1 x'$	$g_1 *$	$g_1 x' h *$	g_1h_1	$g_1 h_1 g_2$	$g_1h_1g_2x$
$x'g_1e$	$x'g_1x *$	$x'g_1x'$	$x'g_1$	$x'g_1x'h$	$x'g_1h_1 *$	$x'g_1h_1g_2$	$x'g_1h_1g_2x$
x^2x'	x^2	$xg_1x' *$	xg_1	h *	hx	hxg_2	hxg_2x
fe	fx	fg_1x'	$fg_1 *$	x'h	$h_1 *$	$h_1g_2 *$	h_1g_2x
xg_2fe	xg_2fx	xg_2fg_1x'	xg_2fg_1	$xg_2x' *$	$g_2 f$	xg_2	xg_2x
$g_2 f e$	$g_2 f x$	$g_2 f g_1 x'$	$g_2 f g_1$	$g_2 x'$	$g_2f *$	$g_2 *$	$g_2 x$
efe	efx	$efxg_1x'$	efg_1	ex'	ef	$eg_2 *$	eg_2x
x'fe	x'fx	$x'fg_1x'$	$x'fg_1$	$(x')^2$	x'f	$x'g_2$	$x'g_2x *$

0 *

FIGURE 5. "Egg-box" diagram of S (* - idempotent).

The Semigroup S is a homomorphic image of F_1 but it is not weakly generated by x. The subsemigroup T generated by the set $\{x, x', g_1 x' h x g_2\}$ is a proper regular subsemigroup of S. It is obtained from S by deleting the fifth and sixth rows and the third and fourth columns of D_h .

The fact that we were able to adapt the example for FI_2 to an example for F_1 shouldn't be so surprising. In the next section we show that FI_2 can be embedded into F_1 in a natural way. But before, let us end this section by proving that F_1 is unique up to isomorphism.

Proposition 5.6. Let T be a regular semigroup weakly generated by x such that all other regular semigroups weakly generated by x are homomorphic

images of T (under a homomorphism that fixes x). Then T and F_1 are isomorphic.

Proof. By the universal property of both T and F_1 , there are surjective homomorphisms $\varphi : F_1 \to T$ and $\psi : T \to F_1$ such that $([x])\varphi = x$ and $x\psi = [x]$. Thus $\varphi \circ \psi$ is an endomorphism of F_1 such that $([x])\varphi \circ \psi = [x]$. By Lemma 4.8, the restriction of $\varphi \circ \psi$ to the set $\{[g] \mid g \in G \setminus \{1\}\}$ must be the identity mapping. Therefore, $\varphi \circ \psi$ is the identity automorphism of F_1 since this semigroup is generated by the previous set. We have shown that φ and ψ are mutually inverse isomorphisms, whence T and F_1 are isomorphic.

6. Embedding FI_2 into F_1

In this section we prove that F_1 has a regular subsemigroup F_1° weakly generated by $\{[xx'], [x'x]\}$ isomorphic to FI_2 . The first step is to construct F_1° . After, we prove that F_1° is isomorphic to FI_2 . It is also important that the reader recalls the notation used in Section 3 for the elements of $G_{2,e}$ (page 7) and for some of the elements of $G_{3,d}$ (page 8).

6.1. The construction of F_1° . We begin by defining a subset $G^{\circ} = \bigcup_{i \in \mathbb{N}} G_i^{\circ}$ of G^5 . We set

$$G_1^{\circ} = G_1 = \{g_{xx'}\}$$
 and $G_2^{\circ} = G_2 = G_{2,e} = \{g_{2,e,1}, g_{2,e,2}\}.$

For i > 2, G_i° is always a proper subset of $G_{i,d}$:

$$\mathbf{G}_{3}^{\circ} = \{g_{3,d,1}, g_{3,d,2}, g_{3,d,3}, g_{3,d,4}\} \subset \mathbf{G}_{3,d}$$

and

$$\mathbf{G}_{i+1}^{\circ} = \{g \in \mathbf{G}_{i+1,d} \mid g^l, g^r \in \mathbf{G}_i^{\circ}\}.$$

Hence $G^{\circ} \subset G_1 \cup G_2 \cup G_d$. The next result is obvious by induction on the indexes *i*. We omit its proof.

Lemma 6.1. If $g \in G^{\circ}$, then $\epsilon(g) \subseteq G^{\circ}$.

Let MR[°] be the set of all nontrivial mountain ranges

$$u = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n}$$

such that:

(a)
$$g_0 g_1 \cdots g_{2n} \in (\mathbf{G}^{\circ} \cup \{1\})^+$$
; and
(b) if $v(g_i) = 1$, then $a_{i+1} = (a_i)'$.

The next result shows us that MR° is closed for uplifting of rivers.

Lemma 6.2. Let $u = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n} \in MR^\circ$ with a river g_i . If v is such that $u \xrightarrow{g_i} v$, then $v \in MR^\circ$.

Proof. Assume first that $g_{i-1} = g_{i+1}$ and $a_i = a'_{i+1}$. Then

$$v = g_0 \cdots g_{i-1} a_{i+2} g_{i+2} \cdots g_{2n}$$

Clearly (a) is satisfied, and if $v(g_{i-1}) \neq 1$, then also (b) is satisfied. If $v(g_{i-1}) = 1$, then $v(g_{i+1}) = 1$, and by condition (b) (applied to u) we know that $a_{i-1} = a'_i$ and $a_{i+1} = a'_{i+2}$. Thus $a_{i-1} = a'_{i+2}$ since $a_i = a'_{i+1}$, and v continues to satisfy (b) even in this case. In conclusion, $v \in MR^\circ$ if $g_{i-1} = g_{i+1}$ and $a_i = a'_{i+1}$.

Assume now that $g_{i-1} \neq g_{i+1}$ or $a_i \neq a'_{i+1}$. Then

$$h_i = (g_{i+1}, a'_{i+1}, g_i, a_i, g_{i-1}) \in \mathbf{G}$$

and

$$v = g_0 \cdots g_{i-1} a_i h_i a_{i+1} g_{i+1} \cdots g_{2n}.$$

In this case, v clearly satisfies (b) since the anchors subsequence does not change and $v(h_i) \neq 1$. If $g_i = 1$, then $h_i \in G_2 = G_2^{\circ}$ and v satisfies (a) also. If $g_i = g_{xx'}$, then $a_i = a'_{i+1}$ by condition (b) (applied to u) and so $g_{i-1} \neq g_{i+1}$. Consequently, $h_i \in G_3^{\circ}$ and v satisfies (a). Finally, if $v(g_i) \geq 2$, then $g_{i-1}, g_{i+1} \in G_d$. If $g_{i-1} = g_{i+1}$, then also $a_i = a'_{i+1}$ since $g_{i-1}^l \neq g_{i-1}^r$, which is a contradiction. Therefore, $g_{i-1} \neq g_{i+1}$ and, once more, v satisfies (a) since $h_i \in G^{\circ}$. Note that we have shown that $v \in MR^{\circ}$ if $g_{i-1} \neq g_{i+1}$ or $a_i \neq a'_{i+1}$. Thus, we have ended the proof of this result.

We denote by M° the set of all mountains from MR°. If $u \in M^{\circ}$, then (a) is equivalent to $\kappa(u) = g_n \in G^{\circ}$ by Lemma 6.1; while (b) is equivalent to the anchors subsequence $a_1a_2\cdots a_{2n-1}a_{2n}$ having a prefix and a suffix from the set $\{11, x'x\}$. In particular, the only two mountains of M° of height 1 are $\beta_1(xx') = 11g_{xx'}$ 11 and $\beta_1(x'x) = 1x'g_{xx'}x$ 1. In the next result we prove that M° is a regular subsemigroup of M.

Proposition 6.3. M° is a regular subsemigroup of M.

Proof. Let $u, v \in M^{\circ}$. By Lemma 6.2, $u \odot v$ is a mountain from MR[°] and M[°] is a subsemigroup of M. Now, M[°] is regular because $\tilde{u} \in M^{\circ}$ for every $u \in M^{\circ}$, and \tilde{u} is an inverse of u.

We denote by F_1° the regular subsemigroup of F_1 corresponding to M° . In the next subsection we prove that F_1° is isomorphic to FI₂.

6.2. An isomorphism from FI_2 to F_1° . It is important for this subsection that the reader recalls the construction of FI_2 described in Section 2. It should be clear at this point the connection between the theory developed here and the one developed in [6]. Roughly speaking, we can look to the theory developed in [6] as the simplified version for triples of the theory developed here, obtained by 'dropping' the information given by the anchors. So, the meaning of notions like *i-landscape* and *i-mountain range* (called landscape and mountain range in [6]) and of operations like uplifting of irivers should now be clear to the reader. Otherwise, we advice the reader to look for more information in [6] before she/he proceeds since a clear understanding of these notions is important for what follows.

35

We begin by defining recursively a mapping

$$\varphi: \mathbf{G}^{\circ} \setminus \{g_{xx'}\} \to \mathbf{H} \setminus \{1, e, f\}.$$

We start by setting

$$(g_{2,e,i})\varphi = h_{2,i}$$
 and $(g_{3,d,j})\varphi = h_{3,j}$,

for $i \in \{1,2\}$ and $j \in \{1,2,3,4\}$. Note that if $g_{3,d,j} = (g, a, g_{xx'}, a, g_1)$, for $G_2^{\circ} = \{g, g_1\}$ and $a \in \{1, x'\}$, then

$$(g_{3,d,i})\varphi = (g\varphi, e_1, g_1\varphi),$$

where $e_1 = e$ if a = 1, or $e_1 = f$ if a = x'. Extend φ to all $g \in G_i^{\circ}$ with i > 3 by setting

$$g\varphi = (g^l \varphi, g^c \varphi, g^r \varphi).$$

By the recursive definition, if $g^{ls} = g^c = g^{rt}$ for some $s, t \in \{l, r\}$, then $(g^l \varphi)^s = g^c \varphi = (g^r \varphi)^t$; whence $g\varphi \in H$ and φ is well defined. We will conclude next that φ is a bijection by defining its inverse mapping.

Consider now the mapping $\psi : H \setminus \{1, e, f\} \to G^{\circ} \setminus \{g_{xx'}\}$ obtained recursively as follows. Set

$$(h_{2,i})\psi = g_{2,e,i}$$
 and $(h_{3,j})\psi = g_{3,d,j}$

for $i \in \{1, 2\}$ and $j \in \{1, 2, 3, 4\}$. Observe that if $h_{3,j} = (h, e_1, h_1)$, for $H_2 = \{h, h_1\}$ and $e_1 \in H_1 = \{e, f\}$, then

$$(h_{3,j})\psi = (h\psi, a, g_{xx'}, a, h_1\psi),$$

where a = 1 if $e_1 = e$, or a = x' if $e_1 = f$. Let now $h \in H_i$ with i > 3 and define recursively

$$h\psi = (h^l\psi, a, h^c\psi, b, h^r\psi),$$

where $a' = (h^l \psi)^{sa}$ for $h^{ls} = h^c$ and $b' = (h^r \psi)^{ta}$ for $h^{rt} = h^c$. By the recursive definition and the choice of a and b, the 5-tuple $h\psi$ belongs to G° and ψ is well defined.

Clearly $\varphi_{|G_k^\circ}$ and $\psi_{|H_k}$ are mutually inverse bijections between G_k° and H_k for $k \in \{2,3\}$. If we now assume that $\varphi_{|G_j^\circ}$ and $\psi_{|H_j}$ are mutually inverse bijections between G_j° and H_j for all j < i, we can immediately conclude that $\varphi_{|G_i^\circ}$ and $\psi_{|H_i}$ are mutually inverse bijections between G_i° and H_i too. We just need to observe that the entries g^l , g^c and g^r of g completely determine the other entries g^{la} and g^{ra} for all $g \in G_i^\circ$. Therefore, we can conclude by induction that $\varphi_{|G_i^\circ}$ and $\psi_{|H_i}$ are mutually inverse bijections between G_i° and H_i for all $i \geq 2$. Thus φ and ψ are also mutually inverse bijections between $G^\circ \setminus \{g_{xx'}\}$ and $H \setminus \{1, e, f\}$. Note further that φ and ψ have the following property:

$$(g^s)\varphi = (g\varphi)^s$$
 and $(h^t)\psi = (h\psi)^t$

for any $s, t \in \{l, c, r\}$.

Denote by i-MR and i-M the sets of all nontrivial i-mountain ranges and i-mountains, respectively, from H^+ . Next, we define two mutually inverse

bijections $\overline{\varphi}$ and $\overline{\psi}$ between MR[°] and i-MR by applying naturally the mappings φ and ψ to each letter. There is however a problem with this approach: both φ and ψ are not yet defined for letters of height less than 2, and their definition for letters of height 1 is not so obvious since G_1° has only one letter, namely $g_{xx'}$, while H_1° has two letters, namely e and f.

Let $G^{\circ'} = (G^{\circ} \setminus \{g_{xx'}\}) \cup \{1, 1g_{xx'}1, x'g_{xx'}x\}$ and extend φ to a bijection from $G^{\circ'}$ onto H by setting

$$1\varphi = 1$$
, $(1g_{xx'}1)\varphi = e$, $(x'g_{xx'}x)\varphi = f$.

If $u = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n} \in MR^\circ$, define

$$u\overline{\varphi} = h_0 h_1 \cdots h_{2n-1} h_{2n} \,,$$

where $h_i = (a_i g_{xx'} a_{i+1}) \varphi$ for $g_i = g_{xx'}$, and $h_i = g_i \varphi$ for the other cases. The mapping $\overline{\varphi}$ is well defined since $a_i a_{i+1} \in \{11, x'x\}$ if $g_i = g_{xx'}$ by condition (b) of the definition of MR°. Also by definition of φ , the word $u\overline{\varphi}$ is clearly a nontrivial i-landscape; whence $\overline{\varphi}$ is a mapping from MR° to i-MR.

Extend now ψ to the inverse mapping of $\varphi : \mathbf{G}^{\circ'} \to \mathbf{H}$ and let $v = h_0 h_1 \cdots h_{2n-1} h_{2n}$ be a nontrivial i-mountain range. Set $g_i = g_{xx'}$ if $h_i \in \{e, f\}$, or $g_i = h_i \psi$ otherwise. If $h_i = e$ $[h_i = f]$, set also $a_i = 1 = a_{i+1}$ $[a_i = x' \text{ and } a_{i+1} = x]$; whence $h_i \psi = a_i g_i a_{i+1}$ if $h_i \in \{e, f\}$. If $v(h_{i-1}) \geq 2$ and $v(h_i) \geq 2$, then only one of the letters g_{i-1} or g_i is the left or the right entry of the other (and not both entries). Hence, there is only one anchor a_i that turns the triplet $g_{i-1}a_ig_i$ anchored. Define now

$$v\overline{\psi} = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n}$$

By the previous observations and the definition of ψ , $v\overline{\psi}$ is a mountain range from MR°. A close inspection allows us to conclude also that $\overline{\varphi}$ and $\overline{\psi}$ are inverse mappings since φ and ψ are inverse bijections too.

Note that both $u \in MR^{\circ}$ and $u\varphi$ have the same line graph structure: they differ only on the labels. In particular, $\overline{\varphi}_{|M^{\circ}}$ and $\overline{\psi}_{|i-M}$ are mutually inverse bijections between the sets M° and i-M. We leave these observations registered in the following lemma for future reference.

Lemma 6.4. The mappings $\overline{\varphi}$: MR[°] \rightarrow i-MR and $\overline{\psi}$: i-MR \rightarrow MR[°] are mutually inverse. Further, their restrictions $\overline{\varphi}_{|M^{°}}$ and $\overline{\psi}_{|i-M}$ to the sets of (nontrivial) mountains from MR[°] and (nontrivial) *i*-mountains, respectively, give us mutually inverse bijections between these sets.

It is not so hard to show now that $\overline{\varphi}_{|M^{\circ}}$ is an isomorphism. In fact, if we look carefully to the proof of Lemma 6.2, we see that if

$$u = g_0 a_1 g_1 \cdots g_{2n-1} a_{2n} g_{2n} \in MR^\circ, \quad u\overline{\varphi} = h_0 h_1 \cdots h_{2n-1} h_{2n},$$

and $u \xrightarrow{g_i} v$ for some river g_i , then $u\overline{\varphi} \xrightarrow{h_i} v\overline{\varphi}$. Thus

$$(u_1 \odot u_2)\overline{\varphi}_{|\mathsf{M}^\circ} = (u_1\overline{\varphi}_{|\mathsf{M}^\circ}) \odot (u_2\overline{\varphi}_{|\mathsf{M}^\circ})$$

by recursion. We have proved the following result:

Proposition 6.5. $\overline{\varphi}_{|M^{\circ}}: M^{\circ} \to i$ -M is an isomorphism with inverse isomorphism $\overline{\psi}_{|i-M}$.

Corollary 6.6. F_1° is a regular subsemigroup of F_1 weakly generated by the set of idempotents $\{[xx'], [x'x]\}$ and isomorphic to FI₂.

Proof. Since F_1° is isomorphic to M° and FI_2 is isomorphic to i-M, then F_1° is isomorphic to FI_2 by the previous proposition. In fact, the isomorphism $\overline{\psi}_{|i-M}$ induces an isomorphism $\phi: \operatorname{FI}_2 \to \operatorname{F}_1^\circ$ such that $(e\varrho)\phi = [xx']$ and $(f\varrho)\phi = [x'x]$. Since FI₂ is weakly generated by the set $\{e\varrho, f\varrho\}$, then F^o₁ is weakly generated by $\{[xx'], [x'x]\}$.

Since ϕ is an isomorphism, we know from [6] that F_1° is generated by the set $((H \setminus \{1\})\rho)\phi$. At first sight one may be lead to believe that $((H \setminus \{1\})\rho)\phi =$ $\{[g] \mid g \in G^{\circ}\} \cup \{[x'x]\}$ since $H\psi = (G^{\circ} \setminus \{g_{xx'}\}) \cup \{1, 1g_{xx'}1, x'g_{xx'}x\}$; and therefore one may be lead to think that F_1° is generated by the set $\{[g] \mid g \in$ $G^{\circ} \} \cup \{ [x'x] \}$. However, this is far from being true. In fact, we cannot even guarantee that all [g] belong to F_1° for $g \in G^{\circ}$. For example, if we take

$$g_{3,d,2} = (g_{2,e,1}, x', g_{xx'}, x, g_{2,e,2}) \in \mathbf{G}_3^\circ,$$

then $\beta(g_{3,d,2}) = \beta_1(g_{3,d,2})$ is the mountain

$$11g_{xx'}xg_{2,e,1}x'g_{3,d,2}xg_{2,e,2}x'g_{xx'}11,$$

which does not belong to M° since it fails to satisfy the condition (b). Hence $[g_{3,d,2}] \notin F_1^{\circ}$. Note that, in this case, $[x'g_{3,d,2}x] \in F_1^{\circ}$ as

$$eta(x'g_{3,d,2}x) = 1x'g_{xx'}xg_{2,e,1}x'g_{3,d,2}xg_{2,e,2}x'g_{xx'}x1 \in \mathrm{M}^\circ$$
 .

The example of the previous paragraph somehow illustrates what happens in general. A deeper analyzes into which [g] belong to F_1° , for $g \in G^{\circ}$ with $v(g) \geq 2$, leads us to three distinct cases. In two of them we conclude that [g]belongs to F_1° , but in the third is the ρ congruence class [x'gx] that belongs to F_1° (and not [g]). The three distinct cases are the following ones:

- (i) If v(g) is even, then $\beta_1(g) \in \mathcal{M}^\circ$ and $[g] \in \mathcal{F}_1^\circ$. (ii) If v(g) = 2n + 1, then $g^{c^{n-1}} \in \mathcal{G}_3^\circ$. If $g^{c^{n-1}} = g_{3,d,i}$ for i odd, then $11g_{xx'}1$ is a prefix of $\beta_1(g)$ and $1g_{xx'}11$ is a suffix of $\beta_1(g)$. We can now see that $\beta_1(g)$ also belongs to \mathcal{M}° , and so $[g] \in \mathcal{F}_1^\circ$.
- (*iii*) If v(g) = 2n+1 with $g^{c^{n-1}} = g_{3,d,i}$ for *i* even, then $11g_{xx'}x$ is a prefix of $\beta_1(g)$ and $x'g_{xx'}$ 11 is a suffix of $\beta_1(g)$. Hence $\beta_1(g) \notin M^\circ$ because it fails to satisfy the condition (b); whence $[g] \notin F_1^{\circ}$. However, $\beta(x'gx)$ is equal to $\beta(g)$ except for the first anchor, which becomes x', and the last anchor, which becomes x. So $\beta(x'gx) \in M^{\circ}$ and $[x'gx] \in F_1^{\circ}$.

Let $\overline{g} = g$ for the two first cases and $\overline{g} = x'gx$ for the last one. A careful analyzes to the definition of ϕ allows us to conclude that

$$((\mathrm{H} \setminus \{1\})\varrho)\phi = \{[xx'], [x'x]\} \cup \{[\overline{g}] \mid g \in \mathrm{G}^{\circ} \text{ with } \upsilon(g) \ge 2\}.$$

Consequently, F_1° is generated by the previous set.

In [6, Corollary 6.10] we proved that each regular semigroup weakly generated by a finite set of idempotents strongly divides FI_2 , that is, it is a homomorphic image of a regular subsemigroup of FI_2 . Using the Corollary 6.6 we now get the following result:

Corollary 6.7. All regular semigroups weakly generated by a finite set of idempotents strongly divide F_1 .

Note that the previous corollary applies in particular to all regular semigroup generated by a finite set of idempotents and to all finite idempotent generated semigroups. In [6, Corollary 6.11] we concluded that all finite semigroups divide FI_2 , that is, are homomorphic images of (not necessarily regular) subsemigroup of FI_2 . We now have also that:

Corollary 6.8. All finite semigroups divide F_1 .

Acknowledgments: This work was partially supported by CMUP, member of LASI, which is financed by (Portuguese) national funds through FCT - Fundação para a Ciência e a Tecnologia, I.P., under the project with reference UIDB/00144/2020. The author also would like to acknowledge the importance of the GAP software [1], and its Semigroup package [4], in the research presented in this paper: the several simulations computed in GAP allowed the emergence of the pattern used to construct the set G.

References

- The GAP Group, GAP Groups, Algorithms, and Programming, Version 4.10.0, 2018, http://www.gap-system.org.
- [2] T. Hall, Identities for existence varieties of regular semigroups, Bull. Austral. Math. Soc. 40 (1989), 59–77.
- [3] J. Kadourek and M. B. Szendrei, A new approach in the theory of orthodox semigroups, Semigroup Forum 40 (1990), 257–296.
- [4] J. Mitchell, Semigroups GAP package, Version 3.0.20, 2018, http://www.gap-system.org.
- [5] M. H. A. Newman, On theories with a combinatorial definition of equivalence, Ann. of Math. (2) 43 (1942), 223–243.
- [6] L. Oliveira, Regular semigroups weakly generated by idempotents, preprint.
- [7] Y. T. Yeh, The existence of e-free objects in e-varieties of regular semigroups, J. Algebra 164 (1992), 471–484.

CMUP, DEPARTAMENTO DE MATEMÁTICA, FACULDADE DE CIÊNCIAS, UNIVERSIDADE DO PORTO, RUA DO CAMPO ALEGRE S/N, 4169-007 PORTO, PORTUGAL

Email address: loliveir@fc.up.pt