

A possible chemical clock in high-mass star-forming regions: $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$?

Y. X. WANG(汪友鑫) ¹, J. S. ZHANG(张江水) ¹, H. Z. YU(余鸿智) ^{2,1}, Y. WANG(王尧) ³, Y. T. YAN(闫耀庭) ⁴,
J. L. CHEN(陈家梁) ¹, J. Y. ZHAO(赵洁瑜),¹ AND Y. P. ZOU(邹益鹏) ¹

¹Center for Astrophysics, Guangzhou University, Guangzhou, 510006, PR China

²Ural Federal University, 19 Mira Street, 620002 Ekaterinburg, Russia

³Purple Mountain Observatory and Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, 10 Yuanhua Road, Nanjing 210023, PR China

⁴Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

(Received **; Revised **; Accepted **)

Submitted to ApJ

ABSTRACT

We conducted observations of multiple HC_3N ($J = 10-9$, $12-11$ and $16-15$) lines and N_2H^+ ($J = 1-0$) line toward a large sample of 61 ultracompact (UC) HII regions, through the Institut de Radioastronomie Millimétrique 30 m and the Arizona Radio Observatory 12 m telescopes. The N_2H^+ $J = 1-0$ line is detected in 60 sources and HC_3N is detected in 59 sources, including 40 sources with three lines, 9 sources with two lines, and 10 sources with one line. Using the rotational diagram, the rotational temperature and column density of HC_3N were estimated toward sources with at least two HC_3N lines. For 10 sources with only one HC_3N line, their parameters were estimated, taking one average value of T_{rot} . For N_2H^+ , we estimated the optical depth of the N_2H^+ $J = 1-0$ line, based on the line intensity ratio of its hyperfine structure lines. Then the excitation temperature and column density were calculated. When combining our results in UC HII regions and previous observation results on high-mass starless cores (HMSCs) and high-mass protostellar cores (HMPOs), the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio clearly increases with the evolution of high mass star-forming regions (HMSFRs). Moreover, positive correlations between the ratio and other evolutionary indicators (dust temperature, bolometric luminosity, and luminosity-to-mass ratio) are found. Thus we propose the ratio of $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ as a reliable chemical clock of HMSFRs.

Keywords: astrochemistry — star, formation — ISM, molecules — ISM, column density

1. INTRODUCTION

High-mass stars ($>8 M_\odot$ and $>10^3 L_\odot$) play an important role in the dynamics and evolution of the interstellar medium. They play a major role in the energy budget of any galaxy, driving their evolution (Martins 1996; Kennicutt & Evans 2012). They also produce all heavy elements in the universe, which are fed back into the interstellar medium via feedback processes of high-mass stars, such as shocks, ultraviolet (UV) radiation, and stellar winds. The heavy elements will be used as the raw material to form new generations of stars and planets (see reviews by Zinnecker & Yorke 2007; Krumholz et al. 2014, and references therein). Therefore, it is extremely important to understand the high-mass star-formation process. However, high-mass stars are rare in galaxies, accounting for about 3% of the total number of stars (Kroupa 2001; Chabrier 2003). The majority of high-mass stars are at large distances away (> 2 kpc; e.g. Urquhart et al. 2018). They also evolve very quickly, reaching the main sequence while still deeply embedded (e.g. Motte et al. 2018; Padoan et al. 2020). All of these complications hinder the understanding of high-mass star formation processes (Urquhart et al. 2022).

So far, it is generally accepted that high-mass star-forming regions (HMSFRs) evolve from high-mass starless cores (HMSCs) to high-mass protostellar cores (HMPOs), then to ultracompact (UC) HII regions (e.g. Beuther et al. 2007; Gerner et al. 2014, 2015; Jin et al. 2015; Han et al. 2015). HMSCs, as the first stage of HMSFR, are formed by cold, dense gas and dust with a temperature of ~ 10 K. They primarily emit radiation at submillimeter wavelengths, while emitting almost nothing at mid-infrared wavelengths (e.g. Sridharan et al. 2005; Rathborne et al. 2006; Beuther et al. 2010; Yuan et al. 2017; Zhang et al. 2020). HMPOs are the second phase stage of HMSFRs, when an active protostar(s) is formed. Emission at mid-infrared wavelengths has been demonstrated, but no radio continuum emission. At this stage, there is a formed disk that transfers the infalling material from the envelope to the HMPO (e.g. Sridharan et al. 2002; Beuther et al. 2002; Ellingsen 2006; Cesaroni et al. 2007; Grave & Kumar 2009; Hsieh et al. 2021). Finally, an UC HII region (size ≤ 0.1 pc, electron density $>10^4$ cm $^{-3}$ and emission measure $\geq 10^7$ pc cm $^{-6}$) is formed when a central high-mass star begins to ionize the surrounding gas (e.g. Wood & Churchwell 1989; Kurtz & Hofner 2005; Hoare et al. 2007; Stéphan et al. 2018). A Lyman continuum flux would be derived from the free-free emission (bremsstrahlung emission) emitted by the hydrogen, which makes UC HII regions directly observable in radio continuum bands (e.g. Urquhart et al. 2013; Hu et al. 2016; Kalcheva et al. 2018; Djordjevic et al. 2019). However, this classification method, according to the physical conditions mentioned above, is extremely crude and may lead to potential overlap among these phases.

For detailed understanding of HMSFRs, it is necessary to determine the chemical composition of high-mass stars at different stages of evolution, owing to the fact that the chemical composition carries a large amount of information about the star-forming region (e.g. Mumma & Charnley 2011; Caselli & Ceccarelli 2012; Jørgensen et al. 2020; Öberg & Bergin 2021). Combining chemical features with physical conditions should be more helpful in distinguishing the different phases of HMSFRs.

To date, many attempts have been made to study the chemical evolution of HMSFRs, known as the chemical clock. The chemical clock is a parameter expressing the chemical evolution by ratios of the column density for two molecules (e.g. Wakelam et al. 2004; Tatematsu et al. 2017; Taniguchi et al. 2019b; Xie et al. 2021). The Millimeter Astronomy Legacy Team 90 GHz Survey (MALT90)¹ provides the largest sample of molecular lines in HMSFRs (Foster et al. 2011; Jackson et al. 2013). Based on the MALT90 survey data, a number of studies were conducted to investigate the chemical clock of HMSFRs. For example, Hoq et al. (2013) calculated the column density ratio between N₂H⁺ and HCO⁺, showing a slight increase from HMSCs to HMPOs, then to UC HII regions. Yu & Wang (2015) also carried out an investigation of the $N(\text{N}_2\text{H}^+)/N(\text{H}^{13}\text{CO}^+)$ ratio and the $N(\text{CCH})/N(\text{H}^{13}\text{CO}^+)$ ratio, which showed a marginal decrease from HMPOs to UC HII regions. Based on H¹³CN and HN¹³C line observations, Jin et al. (2015) found that a slight statistically increasing tendency in the HCN/HNC abundance ratio with the evolution of HMSFRs and suggested that the HCN/HNC abundance ratio can be used as a chemical clock. Moreover, Taniguchi et al. (2019b) carried out a survey observations of HC₃N ($J = 9-8$ and $10-9$), N₂H⁺ ($J = 1-0$) and CCS ($J_N = 6_7-5_6$) toward 17 HMSCs and 28 HMPOs using the Nobeyama 45 m telescope. They obtained that the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio clearly increases from HMSCs to HMPOs and therefore proposed that the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio may be used as a reliable chemical clock in HMSFRs. However, this survey was not sufficient to consider the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio as a good chemical clock due to the lack of observational evidence for the UC HII regions. Therefore, we have conducted survey observations of HC₃N and N₂H⁺ molecular line transitions toward UC HII regions.

In this paper, we presented results of the survey of HC₃N ($J = 10-9$, $12-11$, $16-15$) and N₂H⁺ ($J = 1-0$) toward 61 UC HII regions using the Institut de Radioastronomie Millimétrique (IRAM) 30 m telescope² and the Arizona Radio Observatory (ARO) 12 m telescope³. Our sample and observation information are summarized in Section 2. In Section 3, we present the fitting results of detected spectral lines and made the analyze on the spectral features (line wing, line width). The physical parameters (column density, temperature) are also estimated for our detections. In Section 4, we investigate the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio at different stages of HMSFRs and make a comparative analysis on the ratio and other evolutionary indicators, in order to check the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio as a reliable chemical clock of HMSFRs. A short summary is listed in Section 5.

2. SOURCE SELECTION AND OBSERVATIONS

2.1. Source selection

¹ <http://malt90.bu.edu/>

² <https://www.iram-institute.org/EN/content-page-55-7-55-0-0-0.html>

³ <https://aro.as.arizona.edu/?q=facilities/uarizona-aro-12-meter-telescope>

UC HII regions are the most luminous objects in the Milky Way at the far-infrared wavelengths, and have radio continuum emission. One large UC II region sample of about 600 sources has been established (Giveon et al. 2005a,b; Urquhart et al. 2007, 2009, 2013; Cesaroni et al. 2015; Hu et al. 2016; Kalcheva et al. 2018; Djordjevic et al. 2019), based on infrared survey (Price et al. 2001), radio surveys (Hoare et al. 2012; Purcell et al. 2013), and submillimeter surveys (Thompson et al. 2006; Schuller et al. 2009; Csengeri et al. 2014).

Toward a large sample of HMSFRs with accurate distances, thanks to the results of the Bar and Spiral Structure Legacy (BeSSeL⁴) project (Reid et al. 2014, 2019), we have performed one systematic survey on CS and its isotopic molecules, to make direct measurements of Galactic carbon and sulfur isotope ratios (Yan et al. 2022). Crossmatching this sample with the catalog of UC HII regions, we obtain 61 UC HII regions with accurate distances. These 61 UC HII regions serve as our targets to test the chemical clock $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ in HMSFRs. The detailed information of these UC HII regions are summarized in Table 1.

2.2. Observations

We carried out observations of multiple lines of HC_3N ($J = 10-9$, $12-11$, $16-15$) and N_2H^+ ($J = 1-0$) lines with the IRAM 30 m telescope and the ARO 12 m telescope. Table 2 lists the parameters of targeted lines, including the rest frequency ν , the upper energy level E_u , the product of the line strength, and the square of the electric dipole moment $S\mu^2$, which are taken from the Cologne Database for Molecular Spectroscopy (CDMS)⁵ catalog, and the half-power beamwidth (HPBW) for each transition.

2.2.1. IRAM 30 m observations

The observations of the HC_3N ($J = 10-9$ and $16-15$) and N_2H^+ ($J = 1-0$) lines were carried out toward our sample with IRAM 30 m telescope during August 18–24, 2020 within project 022-20 (PI Hongzhi Yu). The Eight Mixer Receiver (EMIR) heterodyne receivers (E090 and E150) were employed covering the frequency ranges of 90.5–98.2 GHz and 138.3–146 GHz. We applied a fast Fourier transform spectrometer (FTS) backend in the wideband mode (16 GHz, containing four IF subbands with 4 GHz each in dual polarization), with a velocity resolution of ~ 0.64 km s^{-1} at 90 GHz, ~ 0.62 km s^{-1} at 93 GHz, and ~ 0.40 km s^{-1} at 145 GHz. The standard position-switching mode and the chopper-wheel calibration method were adopted in our observations, with the off-source position set at $30'$ in azimuth. The typical system temperature was 92–150 K for HC_3N ($J = 10-9$) and 120–300 K for HC_3N ($J = 16-15$). The main-beam brightness temperature T_{mb} can be obtained from the antenna temperature T_{A}^* divided by the ratio of the main-beam efficiency B_{eff} and the forward efficiency F_{eff} . The ratio of $B_{\text{eff}}/F_{\text{eff}}$ is about 85% at 90 and 93 GHz, and 78% at 145 GHz⁶.

2.2.2. ARO 12 m observations

Toward those 49 sources with detections of both HC_3N lines ($J = 10-9$ and $16-15$) lines by the IRAM 30 m telescope, we performed complementary observations on their HC_3N $J = 12-11$ line in 2021 May, using the ARO 12 m telescope (Yu_20A.1, PI, Hongzhi Yu). The HC_3N $J = 10-9$ line of 33 sources among them was also observed, to check how much different beam sizes affect our results. A 3 mm sideband separating (SBS) and the dual polarization receiver and the two backends of the ARO Wideband Spectrometer (AROWS) mode were adopted, with a velocity resolution of ~ 0.52 km s^{-1} at 90 GHz and ~ 0.43 km s^{-1} at 109 GHz. The center frequencies were set at 97.75 GHz and 108.80 GHz, respectively. A standard position-switching mode was adopted in our observations by using $30'$ offset in azimuth. The typical system temperature was 85–175 K for HC_3N ($J = 10-9$) and 116–243 K for HC_3N ($J = 12-11$). Calibrated by the equation $T_{\text{mb}} = T_{\text{A}}^*/\eta_b$, T_{A}^* can be converted to the T_{mb} . Determined from continuum observations of the Jupiter, the T_{mb} is $\sim 94\%$ at 90 GHz and 92% at 109 GHz.

3. RESULTS AND ANALYSES

3.1. Spectra fitting results

The CLASS/GILDAS⁷ software was used to process the data. We first checked all spectra and then averaged the spectra for two polarizations of each transition for each source. A first-order linear fit was used to subtract the spectral baseline.

⁴ <http://bessel.vlbi-astrometry.org/>

⁵ <https://cdms.astro.uni-koeln.de/cdms/portal/>

⁶ <https://publicwiki.iram.es/Iram30mEfficiencies>

⁷ <https://www.iram.fr/IRAMFR/GILDAS/>

We set the criteria for line detection as a signal-to-noise ratio (S/N) above 5 ($T_{\text{mb}}/\text{rms} > 5$). Out of 61 UC HII regions observed by the IRAM 30 m telescope, N_2H^+ ($J = 1-0$) is detected in all sources, except G060.57+00.18. Forty-nine sources among them were detected in both HC_3N $J = 10-9$ and $J = 16-15$ lines and 10 sources among them were detected in HC_3N $J = 10-9$. Our complementary observations by the ARO 12 m telescope detected the HC_3N ($J = 12-11$) line in 40 sources out of those 49 sources.

For our sources, we fitted the spectra of HC_3N with a single Gaussian profile to obtain their spectral parameters, which are summarized in Table 3. The HC_3N $J = 16-15$ line in G033.39+00.00 and the HC_3N $J = 10-9$ line in G097.53+03.18 show a large velocity distribution that deviates from the Gaussian profile, so we use the "Print area" command in CLASS to obtain the total integrated intensity. The spectra of HC_3N lines of all 61 sources are shown in Figure 1. Our ARO 12 m observations also detected HC_3N ($J = 10-9$) in 33 sources. For comparison, their HC_3N $J = 10-9$ spectra from the IRAM 30 m telescope and the ARO 12 m telescope are displayed in Figure 2.

Although N_2H^+ ($J = 1-0$) theoretically has seven hyperfine transitions at different frequencies, the relatively broad line width ($\geq 2 \text{ km s}^{-1}$) observed toward HMSFRs results in the seven components blending into three groups with roughly Gaussian shapes (e.g. Purcell et al. 2009). Thus we tried to fit the spectra of N_2H^+ ($J = 1-0$) with three Gaussian profiles. For seven sources (G001.00+00.23, G028.39+00.08, G030.70+00.06, G030.81+00.05, G032.79+00.19, G045.49+00.12, and G097.53+03.18) with blending velocity components, the total integrated intensity of them was obtained by the "Print area" method. The N_2H^+ $J = 1-0$ spectra of our sample are presented in Figure 3 and the fitting results are summarized in Table 4.

3.2. Features of spectral lines

3.2.1. Line wing

Line wing features can be found clearly in the HC_3N spectra of 23 sources and most of them show both blue and red wings (for more details see Table 5). We crossmatched them with the outflow catalogs identified by the CO molecule (Wu et al. 2004; Cyganowski et al. 2008; Maud et al. 2015; Li et al. 2016; Yang et al. 2018; Li et al. 2018; Yang et al. 2022) and found that 20 out of 23 sources show CO molecular outflows. The detailed outflow information for these sources is also shown in Table 5.

Molecular outflows are one type of feedback of high-mass star formation to the surrounding molecular cloud, and often lead to the production of shock activities (e.g. Arce et al. 2007). To verify the presence of shock activities in these sources, we investigated observations of other shock tracers toward these sources. The H_2O maser, produced in the circumstellar disk, is caused by shocks from molecular outflows and stellar winds from young stars in HMSFRs (Torrelles et al. 2005). The Class I CH_3OH maser, associated with shocks, is known to be collision pumped (e.g. Leurini et al. 2016). We find that these sources mostly (19 of 23) show Class I CH_3OH (methanol) maser emission (e.g. Yang et al. 2017; Kim et al. 2019) and all of them are associated with H_2O maser emission except G034.79+01.38 (e.g. Breen et al. 2010; Titmarsh et al. 2014; Ladeyschikov et al. 2022; Breen & Ellingsen 2011; Xi et al. 2015; Szymczak et al. 2005; Kim et al. 2019; Cyganowski et al. 2013). Therefore, we proposed the existence of shock activities in these sources (Table 5), which needs further mapping observations with high resolution.

3.2.2. Line width

The line width from the Gaussian fit of observed lines gives the information about the state of turbulence and the region where the gas is mainly being emitting from (Beuther et al. 2002), and it consists of thermal and nonthermal components. The velocity dispersion of molecules attributed by the thermal motion can be estimated by the following formula:

$$\Delta V_{\text{thermal}} = \left(\frac{8kT \ln 2}{m} \right)^{1/2}, \quad (1)$$

where m is the molecular mass of the gas with a value of 51 amu and 29 amu for HC_3N and N_2H^+ , respectively, k is the boltzmann constant, and T is the gas temperature. Then the nonthermal line width can be obtained by

$$\Delta V_{\text{non-thermal}} = (\Delta V_{\text{FWHM}}^2 - \Delta V_{\text{thermal}}^2)^{1/2}, \quad (2)$$

where ΔV_{FWHM} is the observed line width.

Our calculated results show that the thermal line width can be negligible, with a maximum value of 0.17 km s^{-1} for HC_3N and 0.19 km s^{-1} for N_2H^+ , which is less than 10% of the total line width of our sources. Figure 4 shows plots of the comparison of the line width between HC_3N lines ($J = 10-9$, $12-11$ and $16-15$) and N_2H^+ ($J = 1-0$)

line. We used the FWHM of N_2H^+ $J = 1-0$, $F_1 = 0-1$ as the intrinsic FWHM of N_2H^+ , since this group spectra consists of only one component, without blending. It shows clearly that the line width of HC_3N tend to be larger than that of N_2H^+ toward our UC HII region sample. Feng et al. (2021) argued that the inner dense warm regions have more turbulence than the outer regions, by comparing the linewidths of the different transitions of HC_3N . Therefore, our results reflect that HC_3N is more likely to exist in inner and more active star-forming regions compared to N_2H^+ , considering the line width mainly from the nonthermal component caused by turbulence.

3.3. Physical parameters of HC_3N and N_2H^+

3.3.1. Rotational temperature and column density of HC_3N

Most of the lines of HC_3N are usually optically thin (Morris et al. 1976). Therefore, the optical depth effect is insignificant and is not used for further analysis. Under local thermodynamic equilibrium (LTE) assumption, the rotational temperature and the column density can be usually determined by the rotational diagram method with the following formula (Goldsmith & Langer 1999):

$$\ln \frac{3kW}{8\pi^3\nu S\mu^2} = \ln \frac{N}{Q(T_{rot})} - \frac{E_u}{kT_{rot}}, \quad (3)$$

where k and W are the Boltzmann constant and the integrated line intensity, respectively, $Q(T_{rot}) = 4.58 T_{rot} + 0.28$ is the partition function for HC_3N (Wang et al. 2022), T_{rot} and N are the rotational temperature and the column density, respectively, and the $S\mu^2$, ν and E_u values are taken from the CDMS catalog (see Section 2.2).

Since our data come from two different telescopes, the IRAM 30 m and ARO 12 m, with beam sizes of $\sim 27''$ and $\sim 69''$, the beam dilution effect should be investigated. On the IRAM 30 m and ARO 12 m HC_3N ($J = 10-9$) spectra of 33 sources is shown in Figure 2. It shows a stronger signal from the IRAM 30 m than that from the ARO 12 m, indicating that the source size should not be larger than the HPBW of ARO 12 m and there is nonnegligible beam dilution effect. Thus the brightness temperature (T_B) can be derived from the main-beam brightness temperature (T_{mb}) dilution:

$$T_B = T_{mb}/\eta_{BD} = T_{mb} \frac{(\theta_s^2 + \theta_{beam}^2)}{\theta_s^2}, \quad (4)$$

where η_{BD} is the beam-filling factor and θ_s and θ_{beam} are the source size and the HPBW, respectively (Zhang et al. 2017; Yu et al. 2020). According to T_{mb} values measured by the IRAM 30 m and ARO 12 m, the source size (HC_3N $J = 10-9$) is estimated for those 33 sources except G010.32-00.15 and G109.87+02.11 (unreasonable results of size, larger than the HPBW of ARO 12 m, may be nonuniform with a clumpy structure). The results show that the sizes of the sources are larger than the HPBW of the IRAM 30 m telescope and smaller than that of the ARO 12 m telescope. Thus our ARO 12 m HC_3N ($J = 12-11$) data needs to be corrected for the beam dilution effect. According to Equation (4), T_B for our sample is derived from T_{mb} divided by the beam-filling factor, which can be determined from the source size.

For 49 sources that have at least two HC_3N line detections, the rotational temperature and column density of HC_3N of them are derived by Equation (3), using the Levenberg-Marquardt algorithm provided in the python package *lmfit* (Newville et al. 2016). The rotational diagrams are shown in Figure 5 and the derived rotational temperature and column density of HC_3N are summarized in Table 6. For 10 sources with only HC_3N ($J = 10-9$) detection, the average T_{rot} value is used as their T_{rot} values. Then the column density of HC_3N is estimated by Equation (3) (see Table 6).

3.3.2. Excitation temperature and column density of N_2H^+

We used the line intensity ratio method to estimate the optical depth of N_2H^+ , following the procedure described by Purcell et al. (2009). The theoretical line-integrated intensity of Group 1/Group 2 (see details in Table 2) should be 1:5 under an optically thin condition, assuming equal line width for all individual hyperfine components. Assuming the beam-filling factor is 1, the optical depth of N_2H^+ can be determined with the following formula:

$$\frac{\int T_{B,group1} d\nu}{\int T_{B,group2} d\nu} = \frac{T_{mb,group1}}{T_{mb,group2}} = \frac{1 - e^{-0.2\tau_2}}{1 - e^{-\tau_2}}, \quad (5)$$

where τ_2 is the optical depth of N_2H^+ group 2. We then calculated the excitation temperature of N_2H^+ with the following equation:

$$T_{ex} = 4.47/\ln(1 + [\frac{T_{mb,group2}}{4.47(1 - e^{-\tau_2})} + 0.236]^{-1}). \quad (6)$$

Finally, the column density of N_2H^+ was derived using the following formula (Cummins et al. 1986):

$$N = \frac{3kW}{8\pi^3\nu S\mu^2} \left(\frac{T_{ex}}{T_{ex} - T_{bg}} \right) \left(\frac{\tau_2}{1 - \exp(-\tau_2)} \right) Q(T_{ex}) \exp(E_u/kT_{ex}), \quad (7)$$

where T_{bg} ($= 2.73$ K) and W are the background brightness temperature and the integrated intensity of N_2H^+ group 2, respectively. The partition function $Q(T_{ex})$ for N_2H^+ used in our calculations is $Q(T_{ex}) = 4.198T_{ex}$ (Purcell et al. 2009). The derived parameters of N_2H^+ , including the optical depth, the excitation temperature, and the column density, are also summarized in Table 6. We find that the excitation temperature of N_2H^+ is much lower than the rotational temperature of HC_3N . This indicates that N_2H^+ exists in outer and colder regions, while HC_3N traces inner and hotter ones, which is consistent with results in Section 3.2. However, N_2H^+ emission with a clumpy structure ($B_{\text{eff}} < 1$) cannot be ruled out. In this case, the derived T_{ex} is just a lower limit to the real one. Higher angular resolution or mapping observations would be needed to check it.

4. DISCUSSIONS

4.1. $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$, a possible chemical clock?

N_2H^+ is generally considered to be formed in cold and dense environments by the gas-phase reaction $\text{H}_3^+ + \text{N}_2 \rightarrow \text{N}_2\text{H}^+ + \text{H}_2$. When the dust temperature exceeds ~ 20 K, CO molecules will evaporate from the dust grains and react with N_2H^+ , resulting in the destruction of N_2H^+ in the gas phase through the reaction $\text{N}_2\text{H}^+ + \text{CO} \rightarrow \text{HCO}^+ + \text{N}_2$ (e.g. Tobin et al. 2013). The central protostar will release UV photons in the UC HII regions, which also contribute to the destruction of N_2H^+ by the dissociative recombination with electrons $\text{N}_2\text{H}^+ + e^- \rightarrow \text{N}_2 + \text{H}$ or $\text{NH} + \text{N}$ (e.g. Yu et al. 2015). The column density of N_2H^+ should decrease through the reaction with CO molecules or electrons induced by UV photons in the UC HII region phase. Using the N_2H^+ ($J = 1-0$) data of the Nobeyama 45 m telescope with a beam size of $17''$ in HMSCs and HMPOs from Taniguchi et al. (2019b), and following the procedure and equations in Section 3.3.2, we recalculated $N(\text{N}_2\text{H}^+)$ of HMSCs and HMPOs (see details in Table 7). Then we compared the N_2H^+ column density of our UC HII region sources with that of sources in HMSCs and HMPOs. The cumulative distribution function of $N(\text{N}_2\text{H}^+)$ for three samples was presented in Figure 6 (top-left panel). Significant differences on the distribution of $N(\text{N}_2\text{H}^+)$ between UC HII regions and other evolutionary stages (HMSCs and HMPOs) can be found, which is supported by Kolmogorov-Smirnov (K-S) test statistical results (see details in Table 8). This is also supported by the values of the mean $N(\text{N}_2\text{H}^+)$ within uncertainty ranges and corresponding t-test statistical results (Table 8). It indicates that the N_2H^+ column density remains basically stable in HMSCs and HMPOs but begins to increase in UC HII regions, inconsistent with previous theoretical expectations. More theoretical and observational works are needed to further investigate the chemical properties and evolution of N_2H^+ .

HC_3N is believed to be a typical hot-core tracer (e.g. Urquhart et al. 2019). Its precursor molecule, CH_4 , sublimates from dust grains and reacts with C^+ to produce C_2H_2 when the kinetic temperature reaches 25 K. It then reacts with CN in the gas phase through the following neutral-neutral reaction $\text{C}_2\text{H}_2 + \text{CN} \rightarrow \text{HC}_3\text{N} + \text{H}$, resulting in the formation of HC_3N (e.g. Hassel et al. 2008). As the temperature reaches $T \sim 55$ K, C_2H_2 in the dust-grain surface can sublime and further react with CN, leading to an increase in the abundance of HC_3N in the gas phase. When the temperature is greater than 90 K (above the sublimation temperature of HC_3N), HC_3N evaporates directly from the dust grains and its abundance reaches a peak (Taniguchi et al. 2019a). Thus the HC_3N abundance theoretically increases with the evolution of HMSFRs. Taniguchi et al. (2019b) derived $N(\text{HC}_3\text{N})$ in HMSCs and HMPOs with the rotational diagram method, using HC_3N ($J = 5-4, 9-8$ and $10-9$) data from the Nobeyama 45 m telescope with beam sizes of $37''$, $20''$, and $18''$ (Table 7). Comparisons on observed HC_3N abundance of those three samples (HMSCs and HMPOs from Taniguchi et al. (2019b) and UC HII regions from ours) show significant differences on their cumulative distributions (Figure 6, top-right panel), which is supported by K-S test statistical results (see details in Table 8). The difference in the mean value of HC_3N between three samples in different evolutionary stages is also significant, with a chance probability of less than 0.05 from the t-tests (Table 8). These results reflect that the column density of HC_3N gradually increases with evolution, i.e., from HMSCs to HMPOs to UC HII regions.

Urquhart et al. (2019) found that the integrated line intensity ratio of HC_3N and N_2H^+ increases with evolution based on a 3-mm molecular line survey toward 570 HMSFRs using the Mopra telescope, and thus suggested this ratio as a chemical clock. However, the line intensity ratio may be influenced by some factors, such as the optical depth. Compared with the line intensity ratio of HC_3N and N_2H^+ , the ratio of their column densities should be more appropriate for expressing their relative abundance. In Figure 6 (bottom-left panel), we plotted the $N(\text{HC}_3\text{N})$ against

the ratio. It clearly shows that the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio increases from HMSCs to HMPOs and then to UC HII regions, with an increasing value of the $N(\text{HC}_3\text{N})$. The difference in the ratio of the three samples can also be supported by the cumulative distribution results (Figure 6, bottom-right panel), and different mean ratio values (Table 8). All chance probabilities of tests (K-S test for distributions and t-test for the mean values) are less than 0.05 (Table 8), supporting a statistically significant difference in the $\text{HC}_3\text{N}/\text{N}_2\text{H}^+$ ratio among these three samples. It indicates that the ratio of $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ increases in those three evolutionary stages and thus it could be adopted as a chemical clock in HMSFRs.

The linear beam size is different for sources at different distances. A larger beam size of sources at larger distances may include more relatively diffused low-density gas, which could affect the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio results. To check this potential bias in our analysis, we plotted the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio against the heliocentric distance in Figure 7, where the heliocentric distance of UC HII regions is determined from trigonometric parallax measurements (Reid et al. 2014, 2019). HMSCs as well as HMPOs are determined from the latest parallax-based distance calculator V2⁸. It shows those three samples have a similar distance distribution (mainly from 2-11 kpc) and there is no systematic dependence between the ratio and the distance, which means that any observational bias related to the beam dilution is not significant for our ratio results.

4.2. The correlations between $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ and other evolutionary indicators

In the latest version of APEX Telescope Large Area Survey of the Galaxy (ATLASGAL) compact source catalog, Urquhart et al. (2022) classified 5007 HMSFRs into different evolutionary stages, based on multiwavelength (8–870 μm) images. And they found clear systematic increases in the dust temperature (T_{dust}), bolometric luminosity (L_{bol}), and luminosity-to-mass ($L_{\text{bol}}/M_{\text{clump}}$) ratio, which are consistent with the evolutionary stages identified. As discussed in Section 4.1, we find that the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio also increases with the evolution of HMSFRs. It is therefore reasonable to believe that positive correlations exist between the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio and other evolutionary indicators.

Toward those three evolutionary samples, we collected their physical parameters from Urquhart et al. (2018), including the dust temperature, bolometric luminosity, and luminosity-to-mass ratio. Using these data, we plotted the $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio against the dust temperature, bolometric luminosity, and luminosity-to-mass ratio for those samples (Figure 8). Positive correlations could be found in all plots, which supports our previous proposition (section 4.1), i.e., that $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ can be used as a reliable chemical clock in HMSFRs.

5. CONCLUSIONS

Using the IRAM 30 m and ARO 12 m telescopes, we have conducted systematic measurements on N_2H^+ ($J = 1-0$) line and multiple HC_3N ($J = 10-9$, $12-11$ and $16-15$) lines toward 61 UC HII regions. The main results in this work can be summarized as follows:

1. Out of 61 UC HII regions observed by the IRAM 30 m telescope, N_2H^+ ($J = 1-0$) is detected in all sources, except G060.57–00.18. Forty-nine sources among them were detected in both HC_3N $J = 10-9$ and $J = 16-15$ lines and 10 sources among them were detected in HC_3N $J = 10-9$. Our complementary observations by the ARO 12 m telescope detected the HC_3N ($J = 12-11$) line in 40 sources among those 49 sources.
2. The HC_3N spectra of 23 sources among those 61 sources show clear line wings, indicating the presence of molecular outflows. And the majority of these 23 sources show maser emission associated with shock, which reflects the existence of shock activities in these sources. Our calculations on the line width show that the thermal line width can be negligible in both HC_3N and N_2H^+ lines. Comparisons show that the line width of HC_3N is larger than that of N_2H^+ , suggesting that HC_3N is more likely from inner and more active star-forming regions compared to N_2H^+ .
3. Toward 49 sources with at least two HC_3N line detections, we determined the column density and the rotational temperature of HC_3N , using the rotational diagram method. For 10 sources with only one line detection, their $N(\text{HC}_3\text{N})$ was derived with the average T_{rot} . And we estimated the optical depth of N_2H^+ ($J = 1-0$) of this sample using the line intensity ratio method and then obtained their excitation temperatures and column densities of N_2H^+ .

⁸ <http://bessel.vlbi-astrometry.org/node/378>

4. Through comparative analysis on our data in UC HII regions and those in HMSCs and HMPOs, we found that the column density of HC_3N increases from HMSCs to HMPOs, and then to UC HII regions, while that of N_2H^+ stays basically stable. And the column density ratio of HC_3N and N_2H^+ was confirmed to increase with HMSFR evolution. Moreover, positive correlations were found between the ratio and other evolutionary indicators (the dust temperature, bolometric luminosity, and luminosity-to-mass ratio). This supports the proposal that the ratio of $N(\text{HC}_3\text{N})$ and $N(\text{N}_2\text{H}^+)$ can be a reliable chemical clock for HMSFRs.

ACKNOWLEDGMENTS

This work is supported by the Natural Science Foundation of China (Nos. 12041302, 11590782). We thank the operators and staff at the IRAM 30 m and ARO 12 m telescopes for their assistance during our observations. We also thank Dr. J.Z. Wang, and Dr. X. Chen for their nice comments and suggestions. Y.T.Y. is a member of the International Max Planck Research School (IMPRS) for Astronomy and Astrophysics at the Universities of Bonn and Cologne. Y.T.Y. would like to thank the China Scholarship Council (CSC) for support.

REFERENCES

- Arce, H. G., Shepherd, D., Gueth, F., et al. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 245. <https://arxiv.org/abs/astro-ph/0603071>
- Beuther, H., Churchwell, E. B., McKee, C. F., & Tan, J. C. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 165. <https://arxiv.org/abs/astro-ph/0602012>
- Beuther, H., Henning, T., Linz, H., et al. 2010, *A&A*, 518, L78, doi: [10.1051/0004-6361/201014532](https://doi.org/10.1051/0004-6361/201014532)
- Beuther, H., Schilke, P., Menten, K. M., et al. 2002, *ApJ*, 566, 945, doi: [10.1086/338334](https://doi.org/10.1086/338334)
- Breen, S. L., Caswell, J. L., Ellingsen, S. P., & Phillips, C. J. 2010, *MNRAS*, 406, 1487, doi: [10.1111/j.1365-2966.2010.16791.x](https://doi.org/10.1111/j.1365-2966.2010.16791.x)
- Breen, S. L., & Ellingsen, S. P. 2011, *MNRAS*, 416, 178, doi: [10.1111/j.1365-2966.2011.19020.x](https://doi.org/10.1111/j.1365-2966.2011.19020.x)
- Caselli, P., & Ceccarelli, C. 2012, *A&A Rv*, 20, 56, doi: [10.1007/s00159-012-0056-x](https://doi.org/10.1007/s00159-012-0056-x)
- Cesaroni, R., Galli, D., Lodato, G., Walmsley, C. M., & Zhang, Q. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 197. <https://arxiv.org/abs/astro-ph/0603093>
- Cesaroni, R., Pestalozzi, M., Beltrán, M. T., et al. 2015, *A&A*, 579, A71, doi: [10.1051/0004-6361/201525953](https://doi.org/10.1051/0004-6361/201525953)
- Chabrier, G. 2003, *ApJL*, 586, L133, doi: [10.1086/374879](https://doi.org/10.1086/374879)
- Chen, X., Gan, C.-G., Ellingsen, S. P., et al. 2013, *ApJS*, 206, 22, doi: [10.1088/0067-0049/206/2/22](https://doi.org/10.1088/0067-0049/206/2/22)
- Csengeri, T., Urquhart, J. S., Schuller, F., et al. 2014, *A&A*, 565, A75, doi: [10.1051/0004-6361/201322434](https://doi.org/10.1051/0004-6361/201322434)
- Cummins, S. E., Linke, R. A., & Thaddeus, P. 1986, *ApJS*, 60, 819, doi: [10.1086/191102](https://doi.org/10.1086/191102)
- Cyganowski, C. J., Koda, J., Rosolowsky, E., et al. 2013, *ApJ*, 764, 61, doi: [10.1088/0004-637X/764/1/61](https://doi.org/10.1088/0004-637X/764/1/61)
- Cyganowski, C. J., Whitney, B. A., Holden, E., et al. 2008, *AJ*, 136, 2391, doi: [10.1088/0004-6256/136/6/2391](https://doi.org/10.1088/0004-6256/136/6/2391)
- Djordjevic, J. O., Thompson, M. A., Urquhart, J. S., & Forbrich, J. 2019, *MNRAS*, 487, 1057, doi: [10.1093/mnras/stz1262](https://doi.org/10.1093/mnras/stz1262)
- Ellingsen, S. P. 2006, *ApJ*, 638, 241, doi: [10.1086/498673](https://doi.org/10.1086/498673)
- Feng, H., Wang, J., Li, S., et al. 2021, *PASJ*, 73, 467, doi: [10.1093/pasj/psab012](https://doi.org/10.1093/pasj/psab012)
- Foster, J. B., Jackson, J. M., Barnes, P. J., et al. 2011, *ApJS*, 197, 25, doi: [10.1088/0067-0049/197/2/25](https://doi.org/10.1088/0067-0049/197/2/25)
- Gerner, T., Beuther, H., Semenov, D., et al. 2014, *A&A*, 563, A97, doi: [10.1051/0004-6361/201322541](https://doi.org/10.1051/0004-6361/201322541)
- Gerner, T., Shirley, Y. L., Beuther, H., et al. 2015, *A&A*, 579, A80, doi: [10.1051/0004-6361/201423989](https://doi.org/10.1051/0004-6361/201423989)
- Giveon, U., Becker, R. H., Helfand, D. J., & White, R. L. 2005a, *AJ*, 129, 348, doi: [10.1086/426360](https://doi.org/10.1086/426360)
- . 2005b, *AJ*, 130, 156, doi: [10.1086/430459](https://doi.org/10.1086/430459)
- Goldsmith, P. F., & Langer, W. D. 1999, *ApJ*, 517, 209, doi: [10.1086/307195](https://doi.org/10.1086/307195)
- Grave, J. M. C., & Kumar, M. S. N. 2009, *A&A*, 498, 147, doi: [10.1051/0004-6361/200810902](https://doi.org/10.1051/0004-6361/200810902)
- Han, X. H., Zhou, J. J., Wang, J. Z., et al. 2015, *A&A*, 576, A131, doi: [10.1051/0004-6361/201423791](https://doi.org/10.1051/0004-6361/201423791)
- Hassel, G. E., Herbst, E., & Garrod, R. T. 2008, *ApJ*, 681, 1385, doi: [10.1086/588185](https://doi.org/10.1086/588185)
- Hoare, M. G., Kurtz, S. E., Lizano, S., Keto, E., & Hofner, P. 2007, in *Protostars and Planets V*, ed. B. Reipurth, D. Jewitt, & K. Keil, 181. <https://arxiv.org/abs/astro-ph/0603560>

- Hoare, M. G., Purcell, C. R., Churchwell, E. B., et al. 2012, *PASP*, 124, 939, doi: [10.1086/668058](https://doi.org/10.1086/668058)
- Hoq, S., Jackson, J. M., Foster, J. B., et al. 2013, *ApJ*, 777, 157, doi: [10.1088/0004-637X/777/2/157](https://doi.org/10.1088/0004-637X/777/2/157)
- Hsieh, T.-H., Takami, M., Connelley, M. S., et al. 2021, *ApJ*, 912, 108, doi: [10.3847/1538-4357/abee88](https://doi.org/10.3847/1538-4357/abee88)
- Hu, B., Menten, K. M., Wu, Y., et al. 2016, *ApJ*, 833, 18, doi: [10.3847/0004-637X/833/1/18](https://doi.org/10.3847/0004-637X/833/1/18)
- Jackson, J. M., Rathborne, J. M., Foster, J. B., et al. 2013, *PASA*, 30, e057, doi: [10.1017/pasa.2013.37](https://doi.org/10.1017/pasa.2013.37)
- Jin, M., Lee, J.-E., & Kim, K.-T. 2015, *ApJS*, 219, 2, doi: [10.1088/0067-0049/219/1/2](https://doi.org/10.1088/0067-0049/219/1/2)
- Jørgensen, J. K., Belloche, A., & Garrod, R. T. 2020, *ARA&A*, 58, 727, doi: [10.1146/annurev-astro-032620-021927](https://doi.org/10.1146/annurev-astro-032620-021927)
- Kalcheva, I. E., Hoare, M. G., Urquhart, J. S., et al. 2018, *A&A*, 615, A103, doi: [10.1051/0004-6361/201832734](https://doi.org/10.1051/0004-6361/201832734)
- Kennicutt, R. C., & Evans, N. J. 2012, *ARA&A*, 50, 531, doi: [10.1146/annurev-astro-081811-125610](https://doi.org/10.1146/annurev-astro-081811-125610)
- Kim, W.-J., Kim, K.-T., & Kim, K.-T. 2019, *ApJS*, 244, 2, doi: [10.3847/1538-4365/ab2fc9](https://doi.org/10.3847/1538-4365/ab2fc9)
- Kroupa, P. 2001, *MNRAS*, 322, 231, doi: [10.1046/j.1365-8711.2001.04022.x](https://doi.org/10.1046/j.1365-8711.2001.04022.x)
- Krumholz, M. R., Bate, M. R., Arce, H. G., et al. 2014, in *Protostars and Planets VI*, ed. H. Beuther, R. S. Klessen, C. P. Dullemond, & T. Henning, 243, doi: [10.2458/azu_uapress.9780816531240-ch011](https://doi.org/10.2458/azu_uapress.9780816531240-ch011)
- Kurtz, S., & Hofner, P. 2005, *AJ*, 130, 711, doi: [10.1086/431546](https://doi.org/10.1086/431546)
- Ladeyschikov, D. A., Sobolev, A. M., Bayandina, O. S., & Shakhvorostova, N. N. 2022, *AJ*, 163, 124, doi: [10.3847/1538-3881/ac480f](https://doi.org/10.3847/1538-3881/ac480f)
- Leurini, S., Menten, K. M., & Walmsley, C. M. 2016, *A&A*, 592, A31, doi: [10.1051/0004-6361/201527974](https://doi.org/10.1051/0004-6361/201527974)
- Li, F. C., Xu, Y., Wu, Y. W., et al. 2016, *AJ*, 152, 92, doi: [10.3847/0004-6256/152/4/92](https://doi.org/10.3847/0004-6256/152/4/92)
- Li, Q., Zhou, J., Esimbek, J., et al. 2018, *ApJ*, 867, 167, doi: [10.3847/1538-4357/aac2b8](https://doi.org/10.3847/1538-4357/aac2b8)
- Martins, M. A. P. 1996, *General Relativity and Gravitation*, 28, 1309, doi: [10.1007/BF02109523](https://doi.org/10.1007/BF02109523)
- Maud, L. T., Moore, T. J. T., Lumsden, S. L., et al. 2015, *MNRAS*, 453, 645, doi: [10.1093/mnras/stv1635](https://doi.org/10.1093/mnras/stv1635)
- Morris, M., Turner, B. E., Palmer, P., & Zuckerman, B. 1976, *ApJ*, 205, 82, doi: [10.1086/154252](https://doi.org/10.1086/154252)
- Motte, F., Bontemps, S., & Louvet, F. 2018, *ARA&A*, 56, 41, doi: [10.1146/annurev-astro-091916-055235](https://doi.org/10.1146/annurev-astro-091916-055235)
- Mumma, M. J., & Charnley, S. B. 2011, *ARA&A*, 49, 471, doi: [10.1146/annurev-astro-081309-130811](https://doi.org/10.1146/annurev-astro-081309-130811)
- Newville, M., Stensitzki, T., Allen, D. B., et al. 2016, *Lmfit: Non-Linear Least-Square Minimization and Curve-Fitting for Python*, Astrophysics Source Code Library, record ascl:1606.014. <http://ascl.net/1606.014>
- Öberg, K. I., & Bergin, E. A. 2021, *PhR*, 893, 1, doi: [10.1016/j.physrep.2020.09.004](https://doi.org/10.1016/j.physrep.2020.09.004)
- Padoan, P., Pan, L., Juvela, M., Haugbølle, T., & Nordlund, Å. 2020, *ApJ*, 900, 82, doi: [10.3847/1538-4357/abaa47](https://doi.org/10.3847/1538-4357/abaa47)
- Price, S. D., Egan, M. P., Carey, S. J., Mizuno, D. R., & Kuchar, T. A. 2001, *AJ*, 121, 2819, doi: [10.1086/320404](https://doi.org/10.1086/320404)
- Purcell, C. R., Longmore, S. N., Burton, M. G., et al. 2009, *MNRAS*, 394, 323, doi: [10.1111/j.1365-2966.2008.14283.x](https://doi.org/10.1111/j.1365-2966.2008.14283.x)
- Purcell, C. R., Hoare, M. G., Cotton, W. D., et al. 2013, *ApJS*, 205, 1, doi: [10.1088/0067-0049/205/1/1](https://doi.org/10.1088/0067-0049/205/1/1)
- Rathborne, J. M., Jackson, J. M., & Simon, R. 2006, *ApJ*, 641, 389, doi: [10.1086/500423](https://doi.org/10.1086/500423)
- Reid, M. J., Menten, K. M., Brunthaler, A., et al. 2014, *ApJ*, 783, 130, doi: [10.1088/0004-637X/783/2/130](https://doi.org/10.1088/0004-637X/783/2/130)
- . 2019, *ApJ*, 885, 131, doi: [10.3847/1538-4357/ab4a11](https://doi.org/10.3847/1538-4357/ab4a11)
- Schuller, F., Menten, K. M., Contreras, Y., et al. 2009, *A&A*, 504, 415, doi: [10.1051/0004-6361/200811568](https://doi.org/10.1051/0004-6361/200811568)
- Sridharan, T. K., Beuther, H., Saito, M., Wyrowski, F., & Schilke, P. 2005, *ApJL*, 634, L57, doi: [10.1086/498644](https://doi.org/10.1086/498644)
- Sridharan, T. K., Beuther, H., Schilke, P., Menten, K. M., & Wyrowski, F. 2002, *ApJ*, 566, 931, doi: [10.1086/338332](https://doi.org/10.1086/338332)
- Stéphan, G., Schilke, P., Le Bourlot, J., et al. 2018, *A&A*, 617, A60, doi: [10.1051/0004-6361/201730639](https://doi.org/10.1051/0004-6361/201730639)
- Szymczak, M., Pillai, T., & Menten, K. M. 2005, *A&A*, 434, 613, doi: [10.1051/0004-6361:20042437](https://doi.org/10.1051/0004-6361:20042437)
- Taniguchi, K., Herbst, E., Caselli, P., et al. 2019a, *ApJ*, 881, 57, doi: [10.3847/1538-4357/ab2d9e](https://doi.org/10.3847/1538-4357/ab2d9e)
- Taniguchi, K., Saito, M., Sridharan, T. K., & Minamidani, T. 2019b, *ApJ*, 872, 154, doi: [10.3847/1538-4357/ab001e](https://doi.org/10.3847/1538-4357/ab001e)
- Tatematsu, K., Liu, T., Ohashi, S., et al. 2017, *ApJS*, 228, 12, doi: [10.3847/1538-4365/228/2/12](https://doi.org/10.3847/1538-4365/228/2/12)
- Thompson, M. A., Hatchell, J., Walsh, A. J., MacDonald, G. H., & Millar, T. J. 2006, *A&A*, 453, 1003, doi: [10.1051/0004-6361:20054383](https://doi.org/10.1051/0004-6361:20054383)
- Titmarsh, A. M., Ellingsen, S. P., Breen, S. L., Caswell, J. L., & Voronkov, M. A. 2014, *MNRAS*, 443, 2923, doi: [10.1093/mnras/stu1346](https://doi.org/10.1093/mnras/stu1346)
- Tobin, J. J., Bergin, E. A., Hartmann, L., et al. 2013, *ApJ*, 765, 18, doi: [10.1088/0004-637X/765/1/18](https://doi.org/10.1088/0004-637X/765/1/18)
- Torrelles, J. M., Patel, N., Gómez, J. F., Anglada, G., & Uscanga, L. 2005, *Ap&SS*, 295, 53, doi: [10.1007/s10509-005-3659-0](https://doi.org/10.1007/s10509-005-3659-0)
- Urquhart, J. S., Busfield, A. L., Hoare, M. G., et al. 2007, *A&A*, 461, 11, doi: [10.1051/0004-6361:20065837](https://doi.org/10.1051/0004-6361:20065837)
- Urquhart, J. S., Hoare, M. G., Purcell, C. R., et al. 2009, *A&A*, 501, 539, doi: [10.1051/0004-6361/200912108](https://doi.org/10.1051/0004-6361/200912108)

- Urquhart, J. S., Thompson, M. A., Moore, T. J. T., et al. 2013, MNRAS, 435, 400, doi: [10.1093/mnras/stt1310](https://doi.org/10.1093/mnras/stt1310)
- Urquhart, J. S., König, C., Giannetti, A., et al. 2018, MNRAS, 473, 1059, doi: [10.1093/mnras/stx2258](https://doi.org/10.1093/mnras/stx2258)
- Urquhart, J. S., Figura, C., Wyrowski, F., et al. 2019, MNRAS, 484, 4444, doi: [10.1093/mnras/stz154](https://doi.org/10.1093/mnras/stz154)
- Urquhart, J. S., Wells, M. R. A., Pillai, T., et al. 2022, MNRAS, 510, 3389, doi: [10.1093/mnras/stab3511](https://doi.org/10.1093/mnras/stab3511)
- Wakelam, V., Caselli, P., Ceccarelli, C., Herbst, E., & Castets, A. 2004, A&A, 422, 159, doi: [10.1051/0004-6361:20047186](https://doi.org/10.1051/0004-6361:20047186)
- Wang, Y. X., Zhang, J. S., Yan, Y. T., et al. 2022, A&A, 663, A177, doi: [10.1051/0004-6361/202142450](https://doi.org/10.1051/0004-6361/202142450)
- Wood, D. O. S., & Churchwell, E. 1989, ApJ, 340, 265, doi: [10.1086/167390](https://doi.org/10.1086/167390)
- Wu, Y., Wei, Y., Zhao, M., et al. 2004, A&A, 426, 503, doi: [10.1051/0004-6361:20035767](https://doi.org/10.1051/0004-6361:20035767)
- Xi, H., Zhou, J., Esimbek, J., et al. 2015, MNRAS, 453, 4203, doi: [10.1093/mnras/stv1935](https://doi.org/10.1093/mnras/stv1935)
- Xie, J., Fuller, G. A., Li, D., et al. 2021, Science China Physics, Mechanics, and Astronomy, 64, 279511, doi: [10.1007/s11433-021-1695-0](https://doi.org/10.1007/s11433-021-1695-0)
- Yan, Y. T., Henkel, C., Kobayashi, C., et al. 2022, A&A, submitted
- Yang, A. Y., Thompson, M. A., Urquhart, J. S., & Tian, W. W. 2018, ApJS, 235, 3, doi: [10.3847/1538-4365/aaa297](https://doi.org/10.3847/1538-4365/aaa297)
- Yang, A. Y., Urquhart, J. S., Wyrowski, F., et al. 2022, A&A, 658, A160, doi: [10.1051/0004-6361/202142039](https://doi.org/10.1051/0004-6361/202142039)
- Yang, W., Xu, Y., Chen, X., et al. 2017, ApJS, 231, 20, doi: [10.3847/1538-4365/aa6ff3](https://doi.org/10.3847/1538-4365/aa6ff3)
- Yu, H. Z., Zhang, J. S., Henkel, C., et al. 2020, ApJ, 899, 145, doi: [10.3847/1538-4357/aba8f1](https://doi.org/10.3847/1538-4357/aba8f1)
- Yu, N., & Wang, J.-J. 2015, MNRAS, 451, 2507, doi: [10.1093/mnras/stv1058](https://doi.org/10.1093/mnras/stv1058)
- Yu, N., Wang, J.-J., & Li, N. 2015, MNRAS, 446, 2566, doi: [10.1093/mnras/stu2240](https://doi.org/10.1093/mnras/stu2240)
- Yuan, J., Wu, Y., Ellingsen, S. P., et al. 2017, ApJS, 231, 11, doi: [10.3847/1538-4365/aa7204](https://doi.org/10.3847/1538-4365/aa7204)
- Zhang, S., Zavagno, A., Yuan, J., et al. 2020, A&A, 637, A40, doi: [10.1051/0004-6361/201936792](https://doi.org/10.1051/0004-6361/201936792)
- Zhang, X.-Y., Zhu, Q.-F., Li, J., et al. 2017, A&A, 606, A74, doi: [10.1051/0004-6361/201730791](https://doi.org/10.1051/0004-6361/201730791)
- Zinnecker, H., & Yorke, H. W. 2007, ARA&A, 45, 481, doi: [10.1146/annurev.astro.44.051905.092549](https://doi.org/10.1146/annurev.astro.44.051905.092549)

Table 1. Detailed information of our UC HII region sample.

Source	R.A.	DEC.	D	CH ₃ OH	CH ₃ OH	H ₂ O	OH	Evolutionary stage	Ref.
	J2000	J2000	(kpc)	Class II	Class I				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
G359.13+00.03	17:43:25.61	−29:39:17.55	2.92	Y	N	Y	Y	UC HII	[1]
G000.31−00.20	17:47:09.11	−28:46:16.28	11.11	Y	N	Y	Y	UC HII	[1]
G001.00−00.23	17:48:55.28	−28:11:48.24	3.47	Y	...	Y	...	UC HII	[1]
G006.79−00.25	18:01:57.75	−23:12:34.25	20.41	Y	Y	Y	Y	UC HII	[1]
G007.47+00.05	18:02:13.18	−22:27:58.98	3.30	N	N	Y	...	UC HII	[2]
G009.21−00.20	18:06:52.84	−21:04:27.88	2.92	Y	N	Y	...	UC HII	[1]
G010.32−00.15	18:09:01.45	−20:05:07.85	2.76	Y	...	Y	...	UC HII	[1]
G010.62−00.33	18:10:17.98	−19:54:04.65	4.07	Y	N	Y	...	UC HII	[1]
G011.10−00.11	18:10:28.25	−19:22:30.22	3.79	Y	N	N	...	UC HII	[1]
G013.71−00.08	18:15:36.98	−17:04:32.11	4.55	Y	N	N	...	UC HII	[1]
G015.66−00.49	18:20:59.75	−15:33:09.80	2.35	Y	Y	Y	Y	UC HII	[1]
G016.86−02.15	18:29:24.41	−15:16:04.14	1.88	Y	Y	Y	...	UC HII	[1]
G017.02−02.40	18:30:36.29	−15:14:28.38	1.49	Y	Y	Y	...	UC HII	[1]
G017.63+00.15	18:22:26.38	−13:30:11.95	2.84	Y	N	Y	Y	UC HII	[1]
G019.36−00.03	18:26:25.78	−12:03:53.27	4.33	Y	Y	Y	Y	UC HII	[1]
G022.35+00.06	18:31:44.12	−09:22:12.34	5.92	Y	Y	Y	Y	UC HII	[1]
G023.25−00.24	18:34:31.24	−08:42:47.31	4.81	Y	...	Y	Y	UC HII	[1]
G023.38+00.18	18:33:14.32	−08:23:57.50	5.88	Y	Y	Y	Y	UC HII	[1]
G023.43−00.18	18:34:39.19	−08:31:25.41	6.67	Y	Y	Y	Y	UC HII	[3]
G024.78+00.08	18:36:12.56	−07:12:10.84	5.68	Y	Y	Y	...	UC HII	[1]
G024.85+00.08	18:36:18.39	−07:08:50.83	4.52	Y	N	Y	...	UC HII	[1]
G028.30−00.38	18:44:21.97	−04:17:39.90	4.33	Y	...	Y	Y	UC HII	[1]
G028.39+00.08	18:42:51.98	−03:59:54.49	5.00	Y	Y	Y	Y	UC HII	[1]
G028.83−00.25	18:44:51.09	−03:45:48.38	3.52	Y	Y	Y	Y	UC HII	[1]
G030.22−00.18	18:47:08.30	−02:29:29.33	3.95	...	...	...	...	UC HII	[1]
G030.41−00.23	18:47:40.76	−02:20:30.91	6.54	Y	Y	Y	...	UC HII	[1]
G030.70−00.06	18:47:36.80	−02:00:54.34	3.07	Y	Y	Y	Y	UC HII	[1]
G030.74−00.04	18:47:39.72	−01:57:24.97	7.14	Y	...	Y	Y	UC HII	[1]
G030.78+00.20	18:46:48.09	−01:48:53.95	3.12	Y	Y	Y	Y	UC HII	[3]
G030.81−00.05	18:47:46.98	−01:54:26.42	13.16	Y	Y	Y	Y	UC HII	[1]
G031.24−00.11	18:48:45.08	−01:33:13.20	7.94	N	N	Y	Y	UC HII	[4]
G032.74−00.07	18:51:21.86	−00:12:06.22	9.71	Y	Y	Y	Y	UC HII	[1]
G032.79+00.19	18:50:30.73	−00:01:59.28	8.85	Y	N	Y	...	UC HII	[4]
G033.39+00.00	18:52:14.64	00:24:54.37	2.94	Y	Y	Y	Y	UC HII	[3]
G034.41+00.23	18:53:18.03	01:25:25.50	2.62	Y	Y	Y	Y	UC HII	[1]
G034.79−01.38	18:59:45.98	01:01:18.95	2.62	Y	...	N	...	UC HII	[1]
G035.79−00.17	18:57:16.89	02:27:58.01	4.07	Y	Y	Y	Y	UC HII	[1]

Table 1 *continued on next page*

Table 1 (*continued*)

Source	R.A.	DEC.	D	CH ₃ OH	CH ₃ OH	H ₂ O	OH	Evolutionary stage	Ref.
	J2000	J2000	(kpc)	Class II	Class I				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
G036.11+00.55	18:55:16.79	03:05:05.39	1.88	Y	Y	Y	Y	UC HII	[1]
G037.42+01.51	18:54:14.35	04:41:39.65	11.36	Y	Y	Y	...	UC HII	[1]
G037.47−00.10	19:00:07.14	03:59:52.98	4.13	Y	Y	Y	...	UC HII	[1]
G038.11−00.22	19:01:44.15	04:30:37.40	3.37	Y	N	N	...	UC HII	[1]
G040.28−00.21	19:05:41.21	06:26:12.70	12.82	Y	Y	Y	Y	UC HII	[1]
G040.42+00.70	19:02:39.62	06:59:09.05	12.50	Y	...	Y	Y	UC HII	[1]
G040.62−00.13	19:06:01.63	06:46:36.14	8.85	Y	N	Y	Y	UC HII	[1]
G041.22−00.19	19:07:21.38	07:17:08.12	7.69	Y	N	...	...	UC HII	[1]
G043.03−00.45	19:11:38.98	08:46:30.67	8.40	Y	Y	Y	Y	UC HII	[1]
G045.45+00.06	19:14:21.27	11:09:15.87	6.94	Y	...	Y	Y	UC HII	[4]
G045.49+00.12	19:14:11.35	11:13:06.37	5.62	Y	Y	Y	Y	UC HII	[2]
G048.99−00.29	19:22:26.13	14:06:39.13	6.10	Y	Y	Y	Y	UC HII	[4]
G049.04−01.07	19:25:22.25	13:47:19.53	8.85	Y	Y	Y	...	UC HII	[1]
G049.26+00.31	19:20:44.86	14:38:26.86	7.58	Y	Y	N	...	UC HII	[1]
G049.41+00.32	19:20:59.21	14:46:49.61	4.59	Y	N	Y	...	UC HII	[1]
G049.59−00.24	19:23:26.61	14:40:16.96	3.34	Y	N	Y	...	UC HII	[1]
G058.77+00.64	19:38:49.13	23:08:40.21	4.13	Y	...	Y	...	UC HII	[1]
G059.83+00.67	19:40:59.29	24:04:44.18	8.26	Y	Y	Y	Y	UC HII	[1]
G060.57−00.18	19:45:52.49	24:17:43.24	3.61	Y	N	Y	Y	UC HII	[1]
G071.52−00.38	20:12:57.89	33:30:27.08	5.85	Y	Y	Y	Y	UC HII	[1]
G090.92+01.48	21:09:12.97	50:01:03.66	7.52	Y	...	...	...	UC HII	[1]
G097.53+03.18	21:32:12.43	55:53:49.69	4.41	Y	Y	Y	Y	UC HII	[4]
G108.20+00.58	22:49:31.48	59:55:42.01	0.81	N	...	Y	...	UC HII	[4]
G109.87+02.11	22:56:18.06	62:01:49.56	6.06	Y	Y	Y	Y	UC HII	[1]

NOTE—(1): Source name; (2–3): Right ascension and Declination; (4): The heliocentric distance, which is taken from [Reid et al. \(2014, 2019\)](#); (5–8): Class II CH₃OH maser, Class I CH₃OH maser, H₂O maser and OH maser; (8): Evolutionary stage; (6): References. [1] [Hu et al. \(2016\)](#), [2] [Giveon et al. \(2005a\)](#), [3] [Kalcheva et al. \(2018\)](#), [4] [Urquhart et al. \(2009\)](#).

Table 2. Information of our targeted lines.

Telescope	Transition	Rest frequency (GHz)	E_u/k (K)	$S\mu^2$ (D)	HPBW ($''$)
(1)	(2)	(3)	(4)	(5)	(6)
IRAM 30 m	HC ₃ N (J = 10-9)	90.97902	24.01	139.25	27.6
	HC ₃ N (J = 16-15)	145.56096	59.38	222.79	17.3
	N ₂ H ⁺ (1 ₀ -0 ₁ , group 1) ^a	93.17188	4.47	37.25	27.0
	N ₂ H ⁺ (1 ₂ -0 ₁ , group 2) ^a	93.17370	4.47	62.09	27.0
	N ₂ H ⁺ (1 ₁ -0 ₁ , group 3) ^a	93.17613	4.47	12.42	27.0
ARO 12 m	HC ₃ N (J = 10-9)	90.97902	24.01	139.25	69.1
	HC ₃ N (J = 12-11)	109.17363	34.06	167.09	57.6

NOTE—(1): Telescope; (2): Transition; (3): Rest frequency; (4–5): The upper level energy and the product of the line strength and the square of the electric dipole moment, which are taken from CDMS; (6): Half power beam width.

^aThe seven hyperfine lines of the N₂H⁺ (J = 1–0) transition are blended into three groups: group 1 includes only the F₁F = 01–12 transition; group 2 includes the F₁F = 21–11, 23–12 and 22–11 transitions; and group 3 includes the F₁F = 11–10, 12–12 and 10–11 transitions (Chen et al. 2013).

Table 3. Spectral line parameters of HC₃N.

Source	Telescope	Transition HC ₃ N	Integrated time (min)	rms (K)	V_{LSR} (km s ⁻¹)	$\Delta\nu$ (km s ⁻¹)	$\int T_{\text{mb}} d\nu$ (K km s ⁻¹)	T_{mb} (K)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G000.31–00.20	IRAM 30 m	J = 10–9	3.9	0.06	18.43±0.02	1.91±0.05	4.99±0.11	2.46
	ARO 12 m	J = 10–9	10.0	0.02	18.43±0.02	2.18±0.06	1.58±0.04	0.68
	ARO 12 m	J = 12–11	6.0	0.05	18.34±0.04	1.85±0.09	1.58±0.06	0.80
	IRAM 30 m	J = 16–15	3.9	0.11	18.20±0.04	2.17±0.11	3.77±0.15	1.63
G006.79–00.25	IRAM 30 m	J = 10–9	61.8	0.01	21.10±0.01	2.62±0.02	7.43±0.04	2.66
	ARO 12 m	J = 10–9	10.0	0.03	21.11±0.03	2.81±0.08	2.07±0.05	0.69
	ARO 12 m	J = 12–11	6.0	0.06	20.95±0.07	2.57±0.17	1.97±0.11	0.72
	IRAM 30 m	J = 16–15	61.8	0.02	20.86±0.01	2.62±0.03	4.05±0.04	1.45
G010.32–00.15	IRAM 30 m	J = 10–9	61.8	0.02	12.02±0.01	2.72±0.04	3.96±0.04	1.37
	ARO 12 m	J = 10–9	10.0	0.02	12.22±0.02	2.40±0.06	1.87±0.04	0.73
	ARO 12 m	J = 12–11	6.0	0.06	12.10±0.09	2.51±0.21	1.39±0.10	0.52
	IRAM 30 m	J = 16–15	61.8	0.02	11.55±0.02	3.10±0.04	3.70±0.04	1.12
G015.66–00.49	IRAM 30 m	J = 10–9	3.9	0.04	−5.69±0.06	3.90±0.18	3.82±0.13	0.92
	ARO 12 m	J = 12–11	12.0	0.03	−5.85±0.14	4.32±0.50	0.97±0.08	0.21
	IRAM 30 m	J = 16–15	3.9	0.07	−6.00±0.11	3.61±0.31	2.11±0.14	0.55
G016.86–02.15	IRAM 30 m	J = 10–9	61.8	0.02	17.71±0.01	3.28±0.04	9.39±0.09	2.69

Table 3 continued on next page

Table 3 (*continued*)

Source	Telescope	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
		HC ₃ N	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G017.02−02.40	ARO 12 m	J = 10−9	210.0	0.01	18.56±0.02	4.24±0.05	4.46±0.05	0.99
	ARO 12 m	J = 12−11	6.0	0.04	18.18±0.06	4.28±0.14	4.05±0.11	0.89
	IRAM 30 m	J = 16−15	61.8	0.02	17.45±0.01	2.81±0.03	6.79±0.06	2.27
	IRAM 30 m	J = 10−9	120.0	0.01	20.22±0.02	2.89±0.05	3.70±0.05	1.21
	ARO 12 m	J = 10−9	30.0	0.01	19.88±0.04	3.04±0.11	1.13±0.03	0.35
	ARO 12 m	J = 12−11	6.0	0.06	19.93±0.13	2.66±0.33	1.01±0.10	0.36
G017.63+00.15	IRAM 30 m	J = 16−15	120.0	0.01	20.17±0.02	2.98±0.04	2.03±0.02	0.64
	IRAM 30 m	J = 10−9	3.9	0.05	22.52±0.02	2.10±0.05	4.21±0.08	1.89
	ARO 12 m	J = 10−9	10.0	0.02	22.23±0.02	2.22±0.05	1.67±0.03	0.71
	ARO 12 m	J = 12−11	6.0	0.06	22.09±0.05	1.75±0.14	1.35±0.09	0.72
	IRAM 30 m	J = 16−15	3.9	0.06	22.59±0.03	2.30±0.08	3.69±0.10	1.51
	IRAM 30 m	J = 10−9	61.8	0.01	26.65±0.02	3.27±0.05	5.37±0.07	1.54
G019.36−00.03	ARO 12 m	J = 10−9	30.0	0.01	26.65±0.04	3.26±0.09	1.29±0.03	0.37
	ARO 12 m	J = 12−11	6.0	0.07	26.51±0.11	3.18±0.38	1.71±0.14	0.50
	IRAM 30 m	J = 16−15	61.8	0.02	26.73±0.02	3.72±0.06	6.27±0.08	1.58
	IRAM 30 m	J = 10−9	3.9	0.04	84.32±0.05	2.03±0.11	1.58±0.08	0.73
	ARO 12 m	J = 12−11	12.0	0.03	84.21±0.12	1.99±0.26	0.40±0.05	0.19
	IRAM 30 m	J = 16−15	3.9	0.07	84.24±0.11	2.06±0.32	0.83±0.10	0.38
G023.38+00.18	IRAM 30 m	J = 10−9	3.9	0.04	75.69±0.03	2.43±0.09	2.63±0.08	1.02
	ARO 12 m	J = 12−11	12.0	0.03	75.63±0.13	3.24±0.43	0.64±0.06	0.19
	IRAM 30 m	J = 16−15	3.9	0.07	75.59±0.11	2.50±0.28	1.12±0.10	0.42
	IRAM 30 m	J = 10−9	32.8	0.03	101.79±0.03	4.82±0.08	10.95±0.14	2.13
	ARO 12 m	J = 10−9	10.0	0.02	101.28±0.05	5.66±0.13	3.55±0.06	0.59
	ARO 12 m	J = 12−11	6.0	0.04	101.37±0.08	6.43±0.23	3.99±0.11	0.58
G023.43−00.18	IRAM 30 m	J = 16−15	32.8	0.02	101.93±0.02	3.87±0.07	5.12±0.07	1.24
	IRAM 30 m	J = 10−9	61.8	0.03	110.41±0.03	4.87±0.07	17.29±0.20	3.34
	ARO 12 m	J = 10−9	80.0	0.02	110.16±0.03	4.45±0.09	4.54±0.07	0.96
	ARO 12 m	J = 12−11	6.0	0.04	110.18±0.04	4.44±0.12	4.75±0.10	1.01
	IRAM 30 m	J = 16−15	61.8	0.03	110.66±0.02	5.77±0.05	19.06±0.12	3.10
	IRAM 30 m	J = 10−9	3.9	0.04	109.24±0.07	2.42±0.18	1.54±0.09	0.60
G024.85+00.08	ARO 12 m	J = 10−9	20.0	0.01	109.18±0.05	2.18±0.11	0.39±0.02	0.17
	ARO 12 m	J = 12−11	12.0	0.02	109.05±0.11	2.18±0.29	0.37±0.04	0.16
	IRAM 30 m	J = 16−15	3.9	0.06	108.93±0.08	2.42±0.23	1.63±0.12	0.63
	IRAM 30 m	J = 10−9	3.9	0.04	85.46±0.09	3.05±0.22	1.42±0.09	0.44
	ARO 12 m	J = 12−11	6.0	0.03	84.92±0.15	2.73±0.40	0.53±0.06	0.18
	IRAM 30 m	J = 16−15	3.9	0.06	85.60±0.13	1.86±0.30	0.63±0.08	0.32
G028.30−00.38	IRAM 30 m	J = 10−9	61.8	0.01	78.32±0.02	3.83±0.05	7.23±0.07	1.77
	ARO 12 m	J = 10−9	10.0	0.02	78.29±0.04	3.96±0.12	2.24±0.05	0.53
	ARO 12 m	J = 12−11	6.0	0.04	78.00±0.08	4.26±0.24	1.97±0.08	0.43

Table 3 *continued on next page*

Table 3 (*continued*)

Source	Telescope	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
		HC ₃ N	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G028.83−00.25	IRAM 30 m	J = 16−15	61.8	0.01	78.25±0.02	4.32±0.05	5.50±0.05	1.20
	IRAM 30 m	J = 10−9	120.0	0.01	87.16±0.01	3.20±0.02	6.69±0.03	1.97
	ARO 12 m	J = 10−9	10.0	0.02	87.00±0.03	3.38±0.09	1.68±0.04	0.47
	ARO 12 m	J = 12−11	6.0	0.03	86.91±0.06	3.09±0.15	1.57±0.06	0.48
G030.22−00.18	IRAM 30 m	J = 16−15	120.0	0.01	87.26±0.01	3.45±0.02	3.70±0.02	1.01
	IRAM 30 m	J = 10−9	3.9	0.05	103.89±0.04	1.74±0.10	1.52±0.08	0.82
	ARO 12 m	J = 10−9	10.0	0.02	103.93±0.09	2.45±0.24	0.49±0.04	0.19
	ARO 12 m	J = 12−11	6.0	0.03	103.88±0.15	2.30±0.34	0.36±0.04	0.15
G030.41−00.23	IRAM 30 m	J = 16−15	3.9	0.06	103.53±0.08	1.94±0.21	1.09±0.09	0.53
	IRAM 30 m	J = 10−9	3.9	0.04	105.18±0.05	3.27±0.16	4.17±0.15	1.20
	ARO 12 m	J = 10−9	10.0	0.02	105.02±0.09	3.81±0.26	1.03±0.05	0.25
	ARO 12 m	J = 12−11	6.0	0.03	104.82±0.10	3.46±0.28	1.05±0.07	0.28
G030.70−00.06	IRAM 30 m	J = 16−15	3.9	0.06	104.87±0.09	4.76±0.33	3.84±0.18	0.76
	IRAM 30 m	J = 10−9	81.1	0.01	90.73±0.01	5.07±0.04	16.27±0.09	3.02
	ARO 12 m	J = 10−9	90.0	0.02	91.02±0.02	4.93±0.06	5.98±0.06	1.14
	ARO 12 m	J = 12−11	6.0	0.03	90.80±0.03	4.67±0.09	5.42±0.08	1.09
G030.74−00.04	IRAM 30 m	J = 16−15	81.1	0.02	90.21±0.01	6.08±0.03	15.73±0.06	2.43
	IRAM 30 m	J = 10−9	61.8	0.01	91.67±0.03	3.80±0.07	2.55±0.04	0.63
	ARO 12 m	J = 10−9	20.0	0.01	91.61±0.06	4.19±0.18	1.29±0.04	0.29
	ARO 12 m	J = 12−11	6.0	0.03	91.48±0.11	4.69±0.29	1.49±0.07	0.30
G030.78+00.20	IRAM 30 m	J = 16−15	61.8	0.01	91.46±0.02	4.07±0.05	2.89±0.03	0.67
	IRAM 30 m	J = 10−9	3.9	0.04	81.81±0.03	2.90±0.09	4.77±0.11	1.55
	ARO 12 m	J = 10−9	10.0	0.02	81.75±0.06	3.10±0.16	1.16±0.05	0.35
	ARO 12 m	J = 12−11	6.0	0.03	81.56±0.07	3.16±0.19	1.32±0.06	0.39
G030.81−00.05	IRAM 30 m	J = 16−15	3.9	0.06	81.83±0.05	3.16±0.12	3.95±0.12	1.18
	IRAM 30 m	J = 10−9	61.8	0.02	98.57±0.02	5.81±0.04	22.10±0.12	3.57
	ARO 12 m	J = 10−9	102.0	0.02	97.85±0.03	6.09±0.08	6.47±0.07	1.00
	ARO 12 m	J = 12−11	6.0	0.03	97.74±0.04	5.96±0.09	7.03±0.09	1.11
G032.74−00.07	IRAM 30 m	J = 16−15	61.8	0.03	99.13±0.01	5.96±0.03	23.51±0.11	3.70
	IRAM 30 m	J = 10−9	61.8	0.01	37.11±0.02	6.02±0.05	5.75±0.04	0.90
	ARO 12 m	J = 10−9	10.0	0.02	37.20±0.15	6.04±0.41	1.32±0.07	0.21
	ARO 12 m	J = 12−11	6.0	0.03	37.05±0.15	6.18±0.42	1.55±0.08	0.24
G032.79+00.19	IRAM 30 m	J = 16−15	61.8	0.01	36.60±0.02	7.25±0.05	5.88±0.04	0.76
	IRAM 30 m	J = 10−9	32.8	0.02	14.74±0.01	7.16±0.09	4.64±0.05	0.61
	ARO 12 m	J = 10−9	10.0	0.02	15.44±0.14	7.55±0.32	1.44±0.05	0.18
	ARO 12 m	J = 12−11	6.0	0.03	15.27±0.22	7.60±0.52	1.60±0.09	0.20
G033.39+00.00	IRAM 30 m	J = 16−15	32.8	0.03	14.24±0.05	7.76±0.11	6.56±0.08	0.79
	IRAM 30 m	J = 10−9	3.9	0.04	103.65±0.10	3.52±0.33	2.31±0.15	0.62
	ARO 12 m	J = 12−11	12.0	0.02	103.83±0.15	3.79±0.42	0.60±0.05	0.15

Table 3 *continued on next page*

Table 3 (*continued*)

Source	Telescope	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
		HC ₃ N	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G034.41+00.23	IRAM 30 m	J = 16–15	3.9	0.05	103.20	...	1.87	0.53
	IRAM 30 m	J = 10–9	61.8	0.02	57.81±0.02	3.53±0.07	11.11±0.16	2.95
	ARO 12 m	J = 10–9	120.0	0.02	57.82±0.03	3.67±0.08	3.94±0.07	1.01
	ARO 12 m	J = 12–11	6.0	0.04	57.70±0.04	3.57±0.10	3.98±0.09	1.05
G034.79–01.38	IRAM 30 m	J = 16–15	61.8	0.03	57.68±0.02	4.34±0.05	10.81±0.09	2.34
	IRAM 30 m	J = 10–9	32.8	0.01	45.95±0.01	2.28±0.03	6.21±0.06	2.56
	ARO 12 m	J = 10–9	10.0	0.02	45.88±0.02	2.38±0.06	1.83±0.04	0.72
	ARO 12 m	J = 12–11	6.0	0.04	45.83±0.04	2.29±0.11	1.82±0.07	0.75
G035.79–00.17	IRAM 30 m	J = 16–15	32.8	0.02	45.81±0.01	2.27±0.03	5.22±0.05	2.16
	IRAM 30 m	J = 10–9	3.9	0.04	61.66±0.07	3.78±0.23	2.92±0.13	0.73
	ARO 12 m	J = 10–9	20.0	0.01	61.74±0.07	3.50±0.20	0.67±0.03	0.18
	ARO 12 m	J = 12–11	18.0	0.02	61.38±0.15	4.70±0.44	0.66±0.05	0.13
G037.42+01.51	IRAM 30 m	J = 16–15	3.9	0.06	61.88±0.11	5.49±0.35	3.03±0.14	0.52
	IRAM 30 m	J = 10–9	61.8	0.01	43.95±0.01	2.83±0.02	5.40±0.03	1.79
	ARO 12 m	J = 10–9	10.0	0.02	43.79±0.03	2.75±0.07	1.36±0.03	0.46
	ARO 12 m	J = 12–11	6.0	0.04	43.54±0.06	2.90±0.14	1.60±0.07	0.52
G038.11–00.22	IRAM 30 m	J = 16–15	61.8	0.01	43.82±0.01	2.89±0.02	4.47±0.02	1.45
	IRAM 30 m	J = 10–9	3.9	0.04	83.20±0.04	1.97±0.10	1.70±0.07	0.81
	ARO 12 m	J = 10–9	10.0	0.02	83.02±0.09	2.00±0.23	0.32±0.03	0.15
	ARO 12 m	J = 12–11	6.0	0.03	82.78±0.10	1.61±0.24	0.03±0.05	0.22
G040.28–00.21	IRAM 30 m	J = 16–15	3.9	0.06	83.32±0.08	2.17±0.21	1.09±0.08	0.47
	IRAM 30 m	J = 10–9	3.9	0.04	73.91±0.06	4.60±0.17	6.10±0.17	1.25
	ARO 12 m	J = 10–9	10.0	0.02	73.73±0.06	4.38±0.17	1.46±0.04	0.31
	ARO 12 m	J = 12–11	6.0	0.04	73.60±0.15	5.23±0.47	1.54±0.10	0.28
G040.42+00.70	IRAM 30 m	J = 16–15	3.9	0.06	73.89±0.06	4.10±0.16	4.62±0.14	1.06
	IRAM 30 m	J = 10–9	3.9	0.03	12.99±0.05	1.97±0.13	1.26±0.07	0.60
	ARO 12 m	J = 12–11	24.0	0.02	12.59±0.11	2.85±0.27	0.37±0.03	0.12
	IRAM 30 m	J = 16–15	3.9	0.06	13.05±0.12	2.47±0.29	0.95±0.09	0.36
G043.03–00.45	IRAM 30 m	J = 10–9	4.0	0.04	57.80±0.04	3.55±0.10	4.08±0.10	1.08
	ARO 12 m	J = 10–9	20.0	0.02	57.65±0.07	3.40±0.17	0.81±0.03	0.22
	ARO 12 m	J = 12–11	6.0	0.04	57.51±0.13	3.72±0.28	0.98±0.07	0.25
	IRAM 30 m	J = 16–15	4.0	0.06	57.51±0.04	3.52±0.11	4.18±0.11	1.11
G045.45+00.06	IRAM 30 m	J = 10–9	3.9	0.04	58.56±0.14	4.02±0.37	1.53±0.11	0.36
	ARO 12 m	J = 10–9	10.0	0.02	59.07±0.11	4.39±0.29	0.81±0.04	0.17
	ARO 12 m	J = 12–11	6.0	0.04	58.53±0.18	3.51±0.51	0.68±0.07	0.18
	IRAM 30 m	J = 16–15	3.9	0.06	57.95±0.12	4.03±0.30	1.74±0.11	0.41
G048.99–00.29	IRAM 30 m	J = 10–9	3.9	0.04	67.90±0.02	3.25±0.04	11.83±0.12	3.42
	ARO 12 m	J = 10–9	80.0	0.01	67.90±0.02	3.45±0.04	3.24±0.03	0.88
	ARO 12 m	J = 12–11	12.0	0.02	67.81±0.02	3.27±0.04	3.89±0.04	1.12

Table 3 *continued on next page*

Table 3 (*continued*)

Source	Telescope	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
		HC ₃ N	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G049.26+00.31	IRAM 30 m	J = 16–15	3.9	0.05	67.79±0.02	3.38±0.04	13.34±0.14	3.70
	IRAM 30 m	J = 10–9	3.9	0.04	3.32±0.11	3.99±0.25	1.59±0.09	0.37
	ARO 12 m	J = 10–9	10.0	0.02	3.29±0.16	2.97±0.37	0.29±0.03	0.09
	ARO 12 m	J = 12–11	24.0	0.01	3.18±0.19	4.18±0.42	0.34±0.03	0.08
G058.77+00.64	IRAM 30 m	J = 16–15	3.9	0.06	3.18±0.16	4.17±0.38	1.43±0.11	0.32
	IRAM 30 m	J = 10–9	6.0	0.04	32.47±0.05	2.95±0.13	2.45±0.09	0.78
	ARO 12 m	J = 12–11	3.9	0.03	31.84±0.13	3.01±0.33	0.63±0.06	0.20
	IRAM 30 m	J = 16–15	3.9	0.05	32.29±0.05	2.58±0.14	2.12±0.09	0.77
G059.83+00.67	IRAM 30 m	J = 10–9	3.8	0.04	34.77±0.09	2.92±0.23	1.35±0.09	0.43
	ARO 12 m	J = 10–9	40.0	0.01	34.00±0.16	3.26±0.38	0.24±0.02	0.07
	ARO 12 m	J = 12–11	24.0	0.02	34.23±0.11	2.94±0.26	0.38±0.03	0.12
	IRAM 30 m	J = 16–15	3.8	0.05	34.62±0.06	2.97±0.17	2.05±0.09	0.65
G071.52–00.38	IRAM 30 m	J = 10–9	3.9	0.04	10.49±0.06	2.24±0.14	1.44±0.08	0.60
	ARO 12 m	J = 10–9	40.0	0.01	10.53±0.14	3.09±0.46	0.30±0.03	0.09
	ARO 12 m	J = 12–11	12.0	0.02	10.39±0.15	2.63±0.32	0.32±0.04	0.12
	IRAM 30 m	J = 16–15	3.9	0.06	10.55±0.06	2.10±0.15	1.39±0.08	0.62
G109.87+02.11	IRAM 30 m	J = 10–9	57.9	0.01	–10.77±0.01	3.61±0.01	8.13±0.03	2.12
	ARO 12 m	J = 10–9	162.0	0.01	–10.53±0.01	4.04±0.02	7.27±0.03	1.69
	ARO 12 m	J = 12–11	12.0	0.02	–10.46±0.02	3.46±0.04	4.40±0.04	1.20
	IRAM 30 m	J = 16–15	57.9	0.03	–10.80±0.04	3.58±0.09	2.89±0.06	0.76

NOTE—(1): Source name; (2): Telescope; (3): HC₃N transitions; (4–5): Integrated time and the corresponding root mean square (rms) value; (6): Local standard of rest (LSR) velocity; (7): Full width at half maximum (FWHM); (8): Integrated line intensity; (9): Peak T_{mb} value.

Table 4. Spectral line parameters of N_2H^+ .

Source	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
(1)	(2)	(min)	(K)	(km s $^{-1}$)	(km s $^{-1}$)	(K km s $^{-1}$)	(K)
G000.31–00.20	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.07	9.46 $\pm$ 0.07	2.16 $\pm$ 0.17	3.89 $\pm$ 0.25	1.70	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		17.66 $\pm$ 0.02	2.82 $\pm$ 0.05	17.36 $\pm$ 0.28	5.78	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		23.38 $\pm$ 0.04	2.48 $\pm$ 0.08	8.82 $\pm$ 0.25	3.34	
G006.79–00.25	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.01	12.20 $\pm$ 0.01	2.80 $\pm$ 0.02	5.33 $\pm$ 0.03	1.79	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		20.22 $\pm$ 0.01	3.38 $\pm$ 0.01	21.28 $\pm$ 0.03	5.91	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		26.01 $\pm$ 0.01	3.29 $\pm$ 0.01	12.12 $\pm$ 0.03	3.46	
G010.32–00.15	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.01	3.11 $\pm$ 0.01	2.17 $\pm$ 0.02	2.64 $\pm$ 0.02	1.15	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		11.40 $\pm$ 0.01	2.86 $\pm$ 0.01	12.03 $\pm$ 0.03	3.95	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		17.10 $\pm$ 0.01	2.68 $\pm$ 0.01	7.23 $\pm$ 0.03	2.53	
G015.66–00.49	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.05	–14.04 $\pm$ 0.04	2.81 $\pm$ 0.11	3.51 $\pm$ 0.10	1.17	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		–5.82 $\pm$ 0.01	3.75 $\pm$ 0.03	17.29 $\pm$ 0.12	4.33	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		–0.07 $\pm$ 0.02	3.48 $\pm$ 0.04	11.69 $\pm$ 0.12	3.16	
G016.86–02.15	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.01	8.48 $\pm$ 0.63	2.77 $\pm$ 0.63	6.39 $\pm$ 0.96	2.17	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		16.32 $\pm$ 0.63	2.98 $\pm$ 0.63	19.43 $\pm$ 0.96	6.12	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		22.11 $\pm$ 0.63	4.08 $\pm$ 0.63	16.45 $\pm$ 0.96	3.79	
G017.02–02.40	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.02	11.14 $\pm$ 0.63	2.08 $\pm$ 0.63	4.39 $\pm$ 0.75	1.99	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		19.19 $\pm$ 0.63	2.93 $\pm$ 0.63	16.46 $\pm$ 0.75	5.27	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		24.92 $\pm$ 0.63	2.57 $\pm$ 0.63	9.50 $\pm$ 0.75	3.48	
G017.63+00.15	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.04	13.17 $\pm$ 0.04	2.00 $\pm$ 0.10	1.79 $\pm$ 0.07	0.84	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		21.36 $\pm$ 0.01	2.69 $\pm$ 0.03	8.19 $\pm$ 0.08	2.86	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		27.08 $\pm$ 0.02	2.53 $\pm$ 0.05	4.72 $\pm$ 0.08	1.75	
G019.36–00.03	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.02	17.33 $\pm$ 0.01	2.10 $\pm$ 0.02	3.45 $\pm$ 0.02	1.54	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		25.54 $\pm$ 0.01	2.91 $\pm$ 0.01	14.61 $\pm$ 0.02	4.72	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		31.30 $\pm$ 0.01	2.69 $\pm$ 0.01	8.45 $\pm$ 0.02	2.96	
G022.35+00.06	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.04	75.20 $\pm$ 0.03	1.90 $\pm$ 0.07	1.81 $\pm$ 0.06	0.90	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		83.38 $\pm$ 0.01	2.72 $\pm$ 0.02	8.04 $\pm$ 0.06	2.78	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		89.14 $\pm$ 0.02	2.38 $\pm$ 0.04	4.29 $\pm$ 0.06	1.69	
G023.38+00.18	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.05	66.70 $\pm$ 0.04	1.90 $\pm$ 0.10	1.81 $\pm$ 0.08	0.89	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		75.02 $\pm$ 0.02	3.00 $\pm$ 0.05	8.50 $\pm$ 0.11	2.66	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		80.72 $\pm$ 0.03	2.91 $\pm$ 0.06	5.69 $\pm$ 0.10	1.84	
G023.43–00.18	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.01	92.53 $\pm$ 0.63	4.16 $\pm$ 0.63	5.51 $\pm$ 0.67	1.24	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		101.12 $\pm$ 0.63	5.51 $\pm$ 0.63	19.46 $\pm$ 0.67	3.32	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		107.15 $\pm$ 0.63	3.73 $\pm$ 0.63	9.32 $\pm$ 0.67	2.35	
G024.78+00.08	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.01	101.26 $\pm$ 0.63	3.93 $\pm$ 0.63	9.47 $\pm$ 1.25	2.26	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 2-1$		109.62 $\pm$ 0.63	5.09 $\pm$ 0.63	35.73 $\pm$ 1.25	6.60	
	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 1-1$		115.63 $\pm$ 0.63	4.01 $\pm$ 0.63	16.75 $\pm$ 1.25	3.93	
G024.85+00.08	$\text{N}_2\text{H}^+ \text{ J} = 1-0 \text{ F1} = 0-1$	0.04	100.06 $\pm$ 0.06	1.76 $\pm$ 0.14	0.93 $\pm$ 0.07	0.50	

Table 4 continued on next page

Table 4 (*continued*)

Source	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
(1)	(2)	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
		(3)	(4)	(5)	(6)	(7)	(8)
	N ₂ H ⁺ J = 1–0 F1 = 2–1		108.40±0.03	2.64±0.06	4.20±0.08	1.50	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		114.10±0.04	2.33±0.08	2.43±0.08	0.98	
G028.30–00.38	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	76.46±0.21	2.91±0.43	0.58±0.08	0.19	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		84.44±0.04	3.18±0.10	3.12±0.09	0.92	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		90.25±0.08	3.04±0.16	1.69±0.08	0.52	
G028.39+00.08	N ₂ H ⁺ J = 1–0	0.02	77.33	...	20.80	2.60	
G028.83–00.25	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	78.08±0.01	2.81±0.01	5.05±0.01	1.69	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		86.37±0.01	4.07±0.01	23.95±0.01	5.53	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		92.15±0.01	3.29±0.01	13.58±0.01	3.88	
G030.22–00.18	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	94.86±0.04	1.88±0.10	1.51±0.07	0.75	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		103.11±0.01	2.55±0.03	6.43±0.07	2.37	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		108.80±0.02	2.48±0.06	3.65±0.07	1.38	
G030.41–00.23	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	96.18±0.02	2.52±0.03	5.38±0.01	2.00	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		104.36±0.01	3.43±0.02	21.03±0.08	5.76	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		110.13±0.01	2.99±0.02	11.46±0.07	3.60	
G030.70–00.06	N ₂ H ⁺ J = 1–0	0.02	90.06	...	65.20	6.20	
G030.74–00.04	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	82.51±0.63	3.32±0.63	0.50±0.12	0.14	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		90.43±0.63	3.33±0.63	2.68±0.12	0.76	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		96.10±0.63	4.14±0.63	2.13±0.12	0.48	
G030.78+00.20	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.05	72.72±0.63	2.55±0.63	4.02±0.66	1.48	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		80.98±0.63	3.67±0.63	15.07±0.66	3.86	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		86.74±0.63	3.40±0.63	8.99±0.66	2.49	
G030.81–00.05	N ₂ H ⁺ J = 1–0	0.02	97.11	...	55.70	4.58	
G032.74–00.07	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	28.17±0.63	4.75±0.63	3.57±0.48	0.71	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		36.17±0.63	4.96±0.63	12.57±0.48	2.38	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		42.04±0.63	5.14±0.63	8.16±0.48	1.49	
G032.79+00.19	N ₂ H ⁺ J = 1–0	0.01	15.48	...	9.78	0.89	
G033.39+00.00	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.07	94.74±0.05	1.82±0.14	1.69±0.11	0.87	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		103.03±0.02	2.91±0.05	9.00±0.13	2.91	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		108.69±0.03	2.54±0.08	4.89±0.12	1.81	
G034.41+00.23	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	48.86±0.63	2.90±0.63	8.91±1.57	2.88	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		56.96±0.63	3.52±0.63	38.42±1.57	10.25	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		62.76±0.63	3.09±0.63	21.23±1.57	6.46	
G034.79–01.38	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.01	36.92±0.63	1.99±0.63	3.71±0.75	1.75	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		45.16±0.63	2.79±0.63	14.55±0.75	4.90	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		50.87±0.63	2.60±0.63	9.16±0.75	3.31	
G035.79–00.17	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.03	52.44±0.05	2.65±0.11	1.71±0.06	0.60	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		60.81±0.02	4.18±0.05	8.01±0.09	1.80	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		66.60±0.03	3.27±0.07	4.37±0.08	1.26	

Table 4 *continued on next page*

Table 4 (*continued*)

Source	Transition	Integrated time	rms	V_{LSR}	$\Delta\nu$	$\int T_{\text{mb}} d\nu$	T_{mb}
(1)	(2)	(min)	(K)	(km s ⁻¹)	(km s ⁻¹)	(K km s ⁻¹)	(K)
G037.42+01.51	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	34.81±0.63	2.43±0.63	2.93±0.64	1.13	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		43.11±0.63	3.08±0.63	14.67±0.64	4.47	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		48.81±0.63	3.19±0.63	8.99±0.64	2.65	
G038.11–00.22	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	74.15±0.04	1.87±0.10	1.29±0.06	0.65	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		82.40±0.02	2.79±0.04	6.54±0.07	2.20	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		88.08±0.03	2.69±0.06	3.78±0.07	1.32	
G040.28–00.21	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	65.14±0.63	4.63±0.63	4.27±0.67	0.87	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		72.81±0.63	3.79±0.63	14.46±0.67	3.58	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		78.56±0.63	4.87±0.63	12.91±0.67	2.49	
G040.42+00.70	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	3.76±0.10	2.00±0.22	0.68±0.07	0.32	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		12.09±0.04	2.95±0.08	3.52±0.08	1.12	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		17.82±0.06	2.59±0.13	1.80±0.08	0.65	
G043.03–00.45	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	49.06±0.07	4.07±0.18	2.70±0.09	0.62	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		57.07±0.03	4.31±0.06	10.10±0.12	2.20	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		62.85±0.04	3.75±0.09	5.60±0.11	1.40	
G045.45+00.06	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	50.18±0.14	2.95±0.35	1.10±0.10	0.35	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		58.34±0.04	3.62±0.11	4.27±0.10	1.11	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		64.15±0.06	3.08±0.16	2.29±0.10	0.70	
G048.99–00.29	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	58.81±0.63	3.06±0.63	7.05±1.26	2.17	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		67.02±0.63	4.19±0.63	31.15±1.26	6.99	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		72.84±0.63	3.54±0.63	18.53±1.26	4.92	
G049.26+00.31	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	–5.44±0.09	2.70±0.25	1.08±0.08	0.38	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		2.53±0.03	3.24±0.07	5.15±0.09	1.49	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		8.35±0.05	3.11±0.11	2.90±0.09	0.87	
G058.77+00.64	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	23.51±0.05	2.35±0.12	1.66±0.08	0.67	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		31.63±0.02	2.94±0.04	8.33±0.08	2.66	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		37.39±0.02	2.80±0.06	4.72±0.08	1.59	
G059.83+00.67	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.03	25.76±0.07	1.98±0.19	0.68±0.05	0.32	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		33.85±0.03	2.89±0.07	3.13±0.06	1.02	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		39.52±0.04	2.72±0.10	1.99±0.06	0.69	
G071.52–00.38	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.04	1.18±0.10	2.02±0.23	0.65±0.06	0.30	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		9.58±0.03	2.74±0.07	3.37±0.07	1.16	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		15.37±0.05	2.43±0.11	1.84±0.07	0.71	
G109.87+02.11	N ₂ H ⁺ J = 1–0 F1 = 0–1	0.02	–19.50±0.63	3.03±0.63	2.17±0.44	0.67	
	N ₂ H ⁺ J = 1–0 F1 = 2–1		–11.39±0.63	3.69±0.63	10.46±0.44	2.66	
	N ₂ H ⁺ J = 1–0 F1 = 1–1		–5.59±0.63	3.25±0.63	5.99±0.44	1.73	

NOTE—(1): Source name; (2): N₂H⁺ (J = 1–0) transitions; (3–4): Integrated time and the corresponding rms value; (5): The LSR velocity; (6): FWHM; (7): Integrated line intensity; (8): Peak T_{mb} value.

Table 5. Outflow and maser information of sources with line wings.

Source	Blue wing	Red wing	Outflow	Ref.	H ₂ O maser	Ref.	Class I CH ₃ OH maser	Ref.
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
G006.79−00.25	Y	Y	Y	[1]	Y	[a]	Y	[α]
G010.32−00.15	Y	Y	Y	[1]	Y	[a]	...	...
G015.66−00.49	Y	Y	...	...	Y	[b]	Y	[α]
G016.86−02.15	Y	Y	Y	[2]	Y	[b]	Y	[α]
G017.02−02.40	Y	Y	Y	[3]	Y	[b]	Y	[α]
G019.36−00.03	Y	Y	Y	[2]	Y	[c]	Y	[α]
G023.43−00.18	Y	Y	Y	[2]	Y	[d]	Y	[α]
G024.78+00.08	Y	Y	Y	[2]	Y	[d]	Y	[α]
G028.39+00.08	Y	Y	Y	[4]	Y	[e]	Y	[α]
G028.83−00.25	Y	Y	Y	[2]	Y	[c]	Y	[α]
G030.41−00.23	Y	Y	Y	[4]	Y	[e]	Y	[α]
G030.70−00.06	Y	Y	Y	[4]	Y	[c]	Y	[α]
G030.74−00.04	Y	Y	Y	[4]	Y	[e]	...	...
G030.78+00.20	Y	Y	Y	[4]	Y	[f]	Y	[α]
G030.81−00.05	Y	Y	Y	[4]	Y	[e]	Y	[α]
G033.39+00.00	Y	Y	Y	[5]	Y	[g]	Y	[α]
G034.41+00.23	Y	Y	Y	[4]	Y	[h]	Y	[α]
G034.79−01.38	Y	Y	...	...	...	...	...	...
G035.79−00.17	Y	Y	Y	[6]	Y	[d]	Y	[α]
G040.28−00.21	Y	...	Y	[6]	Y	[c]	Y	[α]
G048.99−00.29	Y	Y	Y	[7]	Y	[g]	Y	[β]
G097.53+03.18	N	Y	...	...	Y	[g]	Y	[β]
G109.87+02.11	Y	Y	Y	[2]	Y	[g]	Y	[β]

NOTE—(1): Source name; (2–3): Blue and red wings; (4–5): Dipolar CO outflows, with corresponding references. [1] Yang et al. (2022), [2] Li et al. (2016), [3] Wu et al. (2004), [4] Yang et al. (2018), [5] Li et al. (2018), [6] Cyganowski et al. (2008) and [7] Maud et al. (2015); (6–9): Water maser and Class I CH₃OH maser information, with corresponding references. [a] Breen et al. (2010), [b] Titmarsh et al. (2014), [c] Ladeyschikov et al. (2022), [d] Breen & Ellingsen (2011), [e] Xi et al. (2015), [f] Szymczak et al. (2005), [g] Kim et al. (2019) and [h] Cyganowski et al. (2013); [α] Yang et al. (2017) and [β] Kim et al. (2019).

Table 6. Derived parameters of HC₃N and N₂H⁺ in UC HII regions.

Source	HC ₃ N			N ₂ H ⁺		
	source size	T_{rot}	N	τ	T_{ex}	N
	(")	(K)	($\times 10^{13} \text{ cm}^{-2}$)		(K)	($\times 10^{13} \text{ cm}^{-2}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)
G000.31−00.20	42.71	28.27±2.78	2.10±0.14	1.08±0.14	11.90±0.02	7.73±0.13
G001.00−00.23	...	33.47**	1.03±0.05**	...	...	...
G006.79−00.25	40.85	22.49±1.47	3.07±0.20	1.17±0.02	11.72±0.01	9.78±0.01
G007.47+00.05	...	27.28*	0.20*	0.89±1.05	3.14±0.68	1.31±0.32
G009.21−00.20	...	33.47**	0.78±0.04**	2.99±0.17	6.83±0.01	12.35±0.10
G010.32−00.15	...	35.09*	1.67*	1.05±0.03	9.19±0.01	5.00±0.01
G010.62−00.33	...	33.47**	1.15±0.03**	2.05±0.09	8.73±0.01	10.34±0.06
G011.10−00.11	...	24.70*	0.89*	3.05±0.21	5.98±0.01	9.07±0.10
G013.71−00.08	...	33.47**	0.28±0.04**	4.05±0.50	4.51±0.03	6.55±0.15
G015.66−00.49	...	23.06*	1.51*	0.82±0.13	10.88±0.04	6.72±0.05
G016.86−02.15	52.59	27.08±3.48	4.00±0.37	1.72±0.02	10.59±0.01	10.73±0.53
G017.02−02.40	44.04	22.90±0.23	1.47±0.01	1.96±0.04	9.24±0.01	9.68±0.44
G017.63+00.15	53.34	34.91±8.12	1.60±0.17	1.08±0.16	7.40±0.04	3.48±0.04
G019.36−00.03	38.91	43.05±7.68	2.59±0.17	1.42±0.05	9.34±0.01	7.06±0.01
G022.35+00.06	...	22.33*	0.63*	1.40±0.16	6.73±0.03	3.98±0.03
G023.25−00.24	...	33.47**	0.40±0.03**	1.10±0.15	7.66±0.04	4.12±0.04
G023.38+00.18	...	19.72*	1.06*	1.51±0.21	6.45±0.04	4.46±0.06
G023.43−00.18	42.66	19.57±4.64	5.29±1.58	1.92±0.03	6.94±0.01	11.55±0.40
G024.78+00.08	43.81	44.42±10.37	7.39±0.61	1.59±0.02	11.44±0.01	19.20±0.67
G024.85+00.08	42.85	43.53±15.16	0.63±0.08	1.50±0.30	4.88±0.06	2.72±0.06
G028.30−00.38	...	20.18*	0.57*	0.08±0.55	15.15±1.78	0.98±0.12
G028.39+00.08	45.15	29.93±3.23	2.77±0.18	...	...	...
G028.83−00.25	38.42	22.83±0.99	2.72±0.11	1.20±0.04	11.05±0.01	10.95±0.01
G030.22−00.18	37.45	27.62±0.68	0.61±0.01	1.32±0.19	6.26±0.04	3.18±0.04
G030.41−00.23	35.67	33.71±3.46	1.81±0.09	1.64±0.08	10.27±0.01	11.20±0.04
G030.70−00.06	53.74	38.61±9.76	6.43±0.63	...	...	...
G030.74−00.04	63.27	41.80±6.20	1.21±0.06	...	...	...
G030.78+00.20	37.39	30.24±4.25	2.07±0.17	2.02±0.15	7.52±0.01	9.09±0.40
G030.81−00.05	42.92	40.08±0.77	9.82±0.07	...	...	...
G031.24−00.11	...	56.16*	0.46*	...	...	...
G032.74−00.07	37.68	37.93±2.42	2.55±0.06	1.12±0.09	6.57±0.03	5.62±0.22
G032.79+00.19	44.55	60.65±4.38	2.48±0.08	...	...	...
G033.39+00.00	...	21.71*	0.92*	1.13±0.27	7.36±0.07	3.91±0.07
G034.41+00.23	44.55	36.99±1.75	4.67±0.09	0.94±0.02	20.03±0.01	20.81±0.85
G034.79−01.38	43.35	31.88±0.47	2.53±0.02	1.75±0.02	9.04±0.01	7.93±0.41

Table 6 *continued on next page*

Table 6 (*continued*)

Source	HC ₃ N			N ₂ H ⁺		
	source size	T_{rot}	N	τ	T_{ex}	N
	(")	(K)	($\times 10^{13} \text{ cm}^{-2}$)		(K)	($\times 10^{13} \text{ cm}^{-2}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)
G035.79−00.17	39.53	41.64±10.54	1.20±0.11	1.50±0.19	5.30±0.04	4.76±0.06
G036.11+00.55	...	27.91*	0.34*	1.98±0.32	4.57±0.04	2.94±0.07
G037.42+01.51	40.75	30.66±2.63	2.28±0.11	0.62±0.05	12.83±0.02	5.50±0.24
G037.47−00.10	...	33.47**	0.27±0.04**	0.91±0.32	5.37±0.14	1.92±0.06
G038.11−00.22	32.99	24.60±3.65	0.73±0.09	1.09±0.20	6.34±0.06	2.93±0.04
G040.28−00.21	39.89	29.10±0.23	2.44±0.01	0.51±0.13	12.11±0.07	5.06±0.24
G040.42+00.70	...	28.93*	0.51*	0.99±0.40	4.72±0.17	1.94±0.09
G040.62−00.13	...	30.23*	0.73*	0.55±0.18	7.68±0.12	2.81±0.06
G041.22−00.19	...	...	...	1.24±0.29	4.54±0.09	1.90±0.06
G043.03−00.45	35.20	38.04±2.30	1.81±0.04	0.95±0.21	6.63±0.07	4.19±0.07
G045.45+00.06	67.08	47.23±15.78	0.66±0.08	1.31±0.40	4.44±0.11	2.93±0.10
G045.49+00.12	...	33.47**	1.05±0.05**	...	...	...
G048.99−00.29	40.67	41.61±5.94	5.57±0.29	1.25±0.06	12.96±0.01	3.47±0.62
G049.04−01.07	...	41.39*	0.45*	0.78±0.35	5.37±0.20	0.49±0.08
G049.26+00.31	39.35	35.68±7.90	0.61±0.06	0.65±0.31	6.14±0.20	0.41±0.07
G049.41+00.32	...	...	...	1.27±1.15	3.34±0.41	0.80±0.33
G049.59−00.24	...	33.47**	0.49±0.03**	0.93±0.14	8.34±0.04	0.64±0.03
G058.77+00.64	...	21.61*	1.01*	0.61±0.18	8.93±0.09	0.57±0.04
G059.83+00.67	30.04	57.29±35.39	0.83±0.21	1.29±0.33	4.32±0.10	0.48±0.07
G060.57−00.18	...	33.47**	0.14±0.02**	...	...	...
G071.52−00.38	29.18	33.40±10.58	0.69±0.10	0.69±0.40	5.31±0.26	1.43±0.08
G090.92+01.48	...	41.27*	0.36*	0.61±0.27	3.82±0.26	0.95±0.07
G097.53+03.18	...	35.19*	0.47*	...	...	...
G108.20+00.58	...	24.55*	0.26*	0.91±0.94	4.33±0.48	1.59±0.18
G109.87+02.11	...	33.63*	3.38*	0.61±0.09	8.93±0.05	3.59±0.15
G359.13+00.03	...	33.47**	0.90±0.04**	0.72±0.51	4.70±0.35	1.71±0.15

NOTE—(1): Source name; (2): Source size (HC₃N J = 10–9). (3–4): The rotational temperature and the column density of HC₃N. T_{rot} and N are obtained by the rotational diagram method with three HC₃N lines data. T_{rot} and N with * are calculated by the rotational diagram with two HC₃N (10–9 & 16–15) lines data. T_{rot} and N with ** represent the average value of T_{rot} of HC₃N and the corresponding column density of HC₃N; (5–7): The optical depth, the excitation temperature and the column density of N₂H⁺.

Table 7. Physical parameters of HC₃N and N₂H⁺ in HMSCs and HMPOs.

Evolutionary stage	Source	D	HC ₃ N		N ₂ H ⁺	
			N	τ	T_{ex}	N
		(kpc)	($\times 10^{13} \text{ cm}^{-2}$)		(K)	($\times 10^{13} \text{ cm}^{-2}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)
HMSC	18385-0512-3	10.96	0.49±0.15	2.20±0.16	3.66±0.02	3.57±0.11
	18445-0222-4	4.81	0.05±0.01	2.59±0.42	3.07±0.05	3.07±0.26
	18447-0229-4	6.88	0.19±0.06	1.45±0.33	3.23±0.10	1.92±0.12
	18447-0229-5	6.68	0.09±0.03	2.14±0.34	3.16±0.05	2.32±0.10
	18454-0158-1	7.27	0.18±0.06	2.78±0.14	3.77±0.01	5.65±0.16
	18454-0158-5	7.25	0.90±0.20	1.23±0.07	5.16±0.02	4.08±0.07
	18454-0158-8	7.26	0.27±0.08	2.76±0.11	4.07±0.01	7.37±0.16
	18454-0158-10	7.27	...	1.03±0.30	3.39±0.15	1.74±0.10
	19175+1357-3	10.21	0.11±0.03	1.98±0.32	3.19±0.05	2.52±0.11
	19175+1357-4	10.21	0.14±0.04	0.71±0.43	3.33±0.37	1.04±0.12
	19410+2336-2	2.16	0.80±0.20	0.97±0.05	6.71±0.02	4.93±0.05
	20081+2720-1	6.40	...	3.19±0.66	2.94±0.06	2.79±0.16
	22570+5912-3	2.84	0.09±0.03	0.45±0.34	3.84±0.49	0.91±0.10
	05358+3543	1.69	0.90±0.30	1.60±0.06	5.43±0.01	5.00±0.07
HMPO	05553+1631	1.55	1.36	1.36±0.71	2.96±0.24	1.42±0.18
	18447-0229	6.96	0.10±0.03	1.65±0.15	3.76±0.03	2.52±0.07
	19035+0641	1.92	0.44±0.14	0.72±0.20	3.98±0.16	1.74±0.07
	19074+0752	7.88	0.24±0.07	0.79±0.30	3.53±0.22	1.45±0.09
	19175+1357	10.06	0.44	3.90±0.47	3.06±0.03	5.97±0.25
	19217+1651	2.80	1.10±0.30	0.44±0.11	6.16±0.11	3.19±0.05
	19266+1745	4.60	0.41±0.13	0.62±0.33	3.61±0.33	1.18±0.09
	19282+1814	8.19	...	2.22±1.06	2.85±0.16	1.74±0.36
	19410+2336	2.16	2.30±0.70	1.88±0.03	8.33±0.00	13.44±0.15
	19411+2306	3.32	0.25±0.07	1.61±0.17	3.66±0.04	2.15±0.07
	19413+2332	2.16	0.15±0.05	2.85±0.59	2.97±0.06	2.71±0.20
	20051+3435	3.98	0.19±0.06	2.18±0.50	3.01±0.07	2.58±0.15
	20126+4104	1.12	...	2.86±0.05	5.64±0.00	14.74±0.10
	20216+4107	2.11	0.80±0.20	2.27±0.08	4.43±0.01	4.03±0.07
	20293+3952	6.00	0.64±0.19	0.66±0.11	5.24±0.08	2.22±0.06
	20319+3958	2.40	0.10±0.03	1.44±0.63	2.98±0.19	1.41±0.15
	20332+4124	2.67	0.32±0.09	1.64±0.22	3.43±0.05	2.51±0.08
	20343+4129	2.17	0.50±0.15	1.97±0.09	4.35±0.01	3.71±0.06
	22134+5834	2.61	0.80±0.20	0.80±0.28	3.59±0.19	1.01±0.06
	22570+5912	2.84	0.15±0.05	2.13±1.20	2.83±0.19	1.82±0.44
	23033+5951	2.73	1.50±0.40	0.78±0.07	6.25±0.03	3.99±0.06

Table 7 continued on next page

Table 7 (*continued*)

Evolutionary stage	Source	D (kpc)	HC ₃ N		N ₂ H ⁺	
			N	τ	T_{ex}	N
			($\times 10^{13} \text{ cm}^{-2}$)		(K)	($\times 10^{13} \text{ cm}^{-2}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)
	23139+5939	2.95	0.70 $\pm$ 0.20	1.32 $\pm$ 0.14	3.96 $\pm$ 0.04	2.68 $\pm$ 0.07

NOTE—(1): Evolutionary stage; (2): Source name; (3): The heliocentric distance, which is determined from the latest parallax-based distance calculator V2; (4): The column density of HC₃N, taken from Taniguchi et al. (2019b). (5–7): The optical depth, the excitation temperature and the column density of N₂H⁺, which are derived with the N₂H⁺ (J = 1–0) data of Taniguchi et al. (2019b) and the method of Purcell et al. (2009).

Table 8. Average $N(\text{N}_2\text{H}^+)$, $N(\text{HC}_3\text{N})$ and $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ values of HMSC, HMPO and UC HII region samples.

Evolutionary stage	$N(\text{N}_2\text{H}^+)$	$N(\text{HC}_3\text{N})$	$N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$
	($\times 10^{13} \text{ cm}^{-2}$)	($\times 10^{13} \text{ cm}^{-2}$)	
(5)	(6)	(7)	(8)
HMSC	3.39 $\pm$ 1.90	0.30 $\pm$ 0.28	0.09 $\pm$ 0.06
HMPO	3.18 $\pm$ 2.61	0.64 $\pm$ 0.54	0.24 $\pm$ 0.23
UC HII region	5.51 $\pm$ 4.55	1.65 $\pm$ 1.54	0.42 $\pm$ 0.41
K-S test p-value ^a	0.77	0.10	0.05
K-S test p-value ^b	0.01	0.02	0.04
K-S test p-value ^c	0.38	<0.01	<0.01
t Test p-value ^a	0.79	0.03	0.01
t Test p-value ^b	0.01	<0.01	0.02
t Test p-value ^c	0.02	<0.01	<0.01

NOTE—K-S test and t-test chance probabilities for comparison between: ^a HMSC and HMPO; ^b HMPO and UC HII region; ^c HMSC and UC HII region.




Figure 1. HC_3N spectra with fitting lines of those 40 sources with detections of all targeted lines, $J=10-9$ and $J=16-15$ IRAM 30 m spectra (top and bottom panels) and $J=12-11$ ARO 12 m spectra (middle panels).




Figure 1. Continued.




Figure 1. Continued.




Figure 1. Continued.




Figure 1. Continued.




Figure 2. $\text{HC}_3\text{N } J = 10-9$ spectra (upper panel: IRAM 30 m; lower panel: ARO 12 m) of 33 sources.




Figure 2. Continued.




Figure 3. N_2H^+ ($J=1-0$) spectra detected by the IRAM 30 m telescope.




Figure 3. Continued.




Figure 4. These panels show the comparison of line widths between HC_3N ($J = 10-9$, $12-11$ and $16-15$) and N_2H^+ ($J = 1-0$). The black dashed line means that both lines have the same line width.




Figure 5. Rotational diagrams of HC₃N for 40 sources with detections of all three HC₃N lines, where HC₃N ($J = 12-11$) was corrected by the beam dilution.




Figure 5. Continued.




Figure 5. Continued.




Figure 6. The cumulative distributions for $N(\text{N}_2\text{H}^+)$ (top left panel), $N(\text{HC}_3\text{N})$ (top right) and their ratio values (bottom right) for HMSC, HMPO and UC HII region samples. The $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ ratio is plotted against the $N(\text{HC}_3\text{N})$ (bottom left) and one clear evolution trend of the ratio can be found, i.e., from HMSC, HMPO to UC HII region stages (our data, red circles, half filling and empty ones for those sources where $N(\text{HC}_3\text{N})$ was derived using two HC_3N lines data and sources with only HC_3N $J = 10-9$ line, respectively). $N(\text{HC}_3\text{N})$ in HMSC and HMPO are taken from (Taniguchi et al. 2019b), $N(\text{N}_2\text{H}^+)$ in HMSC and HMPO are derived using the N_2H^+ ($J = 1-0$) data from Taniguchi et al. (2019b) and following the procedure of Purcell et al. (2009).




Figure 7. $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ against the heliocentric distance and no significant variation can be found between them.




Figure 8. The correlations between $N(\text{HC}_3\text{N})/N(\text{N}_2\text{H}^+)$ and other evolutionary indicators, including dust temperature, bolometric luminosity, and luminosity-to-mass ratio.