

The pride of lions around Messier 105

Igor D. Karachentsev,¹★ Elena I. Kaisina¹ and Valentina E. Karachentseva²

¹*Special Astrophysical Observatory of the Russian Academy of Sciences, Nizhnij Arkhyz, Karachai-Cherkessian Republic, 369167, Russia*

²*Main Astronomical Observatory, National Academy of Sciences of Ukraine, Kiev, 03143, Ukraine*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We undertook a search for new dwarf galaxies in the Leo-I group using the data from the DECaLS digital sky survey. Five new presumed members of this group have been found in a wide vicinity of M 105 (NGC 3379). Currently, the group has a population of 83 galaxies, 33 of which have measured radial velocities. More than half of the group members belong to early types with no signs of ongoing star formation. About a quarter of the galaxies are outside the group’s virial radius, $R_v = 385$ kpc. The presence of multiple systems with a size of about 15 kpc is evident in the group, but there are no noticeable global flat or filamentary substructures. The luminosity function of the group looks to be deficient in galaxies with absolute magnitudes in the interval $M_B = [-18, -15]$ mag. The M 105 group is characterized by a radial velocity dispersion of 136 km s^{-1} , orbital mass estimate $(5.76 \pm 1.32) \times 10^{12} M_\odot$, and the total mass-to-K-band-luminosity ratio $(17.8 \pm 4.1) M_\odot/L_\odot$. The neighboring group of galaxies around M 66 (NGC 3627) has a similar virial radius, 390 kpc, velocity dispersion, 135 km s^{-1} , and total mass-to-luminosity ratio, $(15.6 \pm 3.9) M_\odot/L_\odot$. Both groups in the Leo constellation are approaching the Local Group with a velocity of about 100 km s^{-1} . In the background of the M 105 group, we noted a group of 6 galaxies with an unusually low virial mass-to-luminosity ratio, $M_T/L_K = (4.1 \pm 2.2) M_\odot/L_\odot$.

Key words: galaxies: dwarf – galaxies: groups: individual (M 105, M 66) – surveys – catalogs

1 INTRODUCTION

The Local Volume (LV), limited by a distance of $D \lesssim 11$ Mpc, hosts two dozen galaxies with the luminosity comparable to those of the Milky Way (MW) and the Andromeda galaxy (M 31). Being the dominating dynamic centers of low-population galaxy groups, most of them are surrounded by retinues of dwarf satellites. Measuring the distances and radial velocities of the satellites allows one to determine the total mass of the dark halo of the central galaxy. The median value of the total mass-to-K-band luminosity ratio for these groups is $M_T/L_K \approx 20 M_\odot/L_\odot$ (Karachentsev & Kashibadze 2021). In addition to the poor groups, a rich group around the M 105 (NGC 3379) galaxy in Leo constellation is located at the far boundary of the LV. It contains seven bright galaxies, the luminosity of which are comparable with each other. Early morphological type objects (E, S0), which have depleted their resources for further star formation, prevail among them. In terms of population and morphology, the group of galaxies around M 105 appears to be an intermediate link between the poor groups and the closest galaxy clusters Fornax and Virgo.

Searches for new, fainter galaxies in the M 105 group (it is also called the Leo-I group) were undertaken repeatedly (Ferguson & Sandage 1990; Schombert et al. 1997; Flint et al. 2001; Karachentsev & Karachentseva 2004; Müller et al. 2018). The fortunate position of the M 105 group in the Arecibo radio telescope survey zone made it possible to measure the radial velocities of dwarf irregular galaxies in the vicinity of M 105

(Stierwalt et al. 2009; Haynes et al. 2011). The LV galaxy database (Kaisina et al. 2012) and the Updated Neighboring Galaxy Catalog, UNGC (Karachentsev et al. 2013) include more than 60 M 105 group candidate members, 20 of which have measured radial velocities. These data were used to estimate the virial mass of the M 105 group, which turned out to be in the $(7\text{--}17) \times 10^{12} M_\odot$ range (Karachentsev & Karachentseva 2004; Makarov & Karachentsev 2011; Karachentsev & Kudrya 2014; Karachentsev et al. 2015).

In recent years, the region of the M 105 group has been covered by the DECaLS digital sky survey (Dey et al. 2019), which is approximately 1.5 mag deeper than the Sloan Digital Sky Survey (Abazajian et al. 2009). This made it possible to check the membership in the group of already known galaxies and to discover new probable members of the group (Carlsten et al. 2022). In combination with new measurements of the distances and velocities of galaxies in the group region, this gives reason to return to the estimate of the virial mass of the M 105 group again.

In Section 2, we report the discovery of five new peripheral candidate members of the M 105 group and provide an updated total list of 83 presumable members of the group. In Section 3, we discuss the structural properties of M 105 group. In Section 4, we re-estimate the virial mass of the M 105 group. In Section 5 we discuss structural properties and virial mass estimate for the M 66 group of galaxies neighboring to the M 105 group. Finally, we provide a brief summary of the main results in Section 6.

★ E-mail: idkarach@gmail.com

2 SEARCH FOR SATELLITES AROUND M105

The paper of [Carlsten et al. \(2022\)](#) is an important step in studying the Leo-I galaxy group. Based on the data of the new digital sky survey DECaLS ([Dey et al. 2019](#)) and deep images obtained on large telescopes (CFHT, Subaru, Gemini, Magellan), the authors carried out surface “*g*”- and “*r*”-band photometry for 60 presumable group members, estimated the distances to 32 galaxies using surface brightness fluctuations (SBF), and found 12 new candidate dwarf members of the M 105 group. Searches for new low-surface brightness objects were carried out in the virial zone of the group with a radius of about 350 kpc.

Evidently, some group members may be located outside the boundaries of its virial radius. We therefore repeated the search for dwarf galaxies in an area approximately four times larger. The search for dwarf members of the M 105 group was carried out visually with an emphasis on objects of low surface brightness. The search took into account the shape, structure, and color of galaxy images, which are characteristic of spheroidal and irregular dwarfs. In the virial zone of the group, we independently detected all 12 new dwarf galaxies classified by [Carlsten et al. \(2022\)](#) as “confirmed members” based on their distance measurements by the SBF- method. Among the 22 objects classified as “unconfirmed/possible candidate satellites” by [Carlsten et al. \(2022\)](#), we missed 12 cases. All of them turned out to be faint objects of small angular diameters with apparent magnitudes $g \approx 20 - 21$ mag. We assumed that these 12 objects are rather background galaxies.

Outside the virial radius of the group, we discovered five new dwarf galaxies: Dw 1033 + 1213, Dw 1038 + 1158, Dw 1053 + 1518, Dw 1100 + 1125, and Dw 1106 + 1250. Their DECaLS images are presented in Fig. 1. We measured the fluxes of these galaxies in *g* and *r* bands using the standard methods of processing in the ESO-MIDAS software package¹ for galaxy images from the DESI Legacy Imaging Surveys and obtained the apparent magnitudes according to their description $m = 22.5 - 2.5 \log_{10}(flux)$ ². The typical measurement error was about 0.15 mag.

A summary list of 83 presumable M 105 group members is presented in Table 1. The columns contain the following: name of the galaxy (the new dwarfs detected in this work are highlighted with bold); J2000 equatorial coordinates in degrees; *B*-band apparent magnitude of the galaxy from the LV galaxies database³ (LVGDB); the relation $B = g + 0.313(g - r) + 0.227$, proposed by Lupton⁴, was used for galaxies with measured “*g*” and “*r*” magnitudes; morphological type: Irr — irregular dwarf galaxy, Sph — spheroidal dwarf galaxy, Tr — intermediate between Irr and Sph, BCD — blue compact dwarf, dE — dwarf elliptical; radial velocity relative to the Local Group centroid; distance to the galaxy in Mpc; method used to estimate the distance: TRGB — by the tip of the red giant branch, Cep — by the luminosity of Cepheids, SBF — by the surface brightness fluctuations, TF — by the Tully-Fisher relation between the HI-line width and the galaxy luminosity, mem — by the assumed group membership of the galaxy; projected distance of the galaxy from the group center identified with M 105, in the assumption that all galaxies have an average distance of 10.8 Mpc; source of data on the distances of galaxies, reference to which are given at the note of Table 1.

Data on apparent magnitudes, morphological types, and radial velocities of galaxies are taken from the LVGDB, version of September 1, 2022.

It needs to be noted that apparent magnitudes of some faint galaxies in the LVGDB have been estimated roughly by eyeball. We made a comparison of visual *B*-magnitudes of galaxies from Table 1 with their apparent magnitudes, *g*, measured by [Carlsten et al. \(2020\)](#), based on archival CFHT/MegaCam imaging data. For 30 dwarf galaxies, the average difference in apparent magnitudes is $\langle B - g \rangle = (+0.38 \pm 0.09)$ mag, which is typical for a mixture of dIrr and dSph galaxies. The root-mean-square magnitude difference, to which the error of our visual estimates makes the main contribution, is $\sigma(B - g) = 0.46$ mag.

When compiling the list, we excluded several galaxies from M 105 group members: dw 1044+11 (104432.7+111610) as a tidal structure belonging to a distant elliptical galaxy, dw 1045 + 14a (104500.9 + 140619) as a background galaxy, LeG 23 (105009.1 + 132901) as a distant spiral galaxy and AGC 205268 (104252.4 + 134428) whose radial velocity, measured by the HI line, is a result of a mix-up with the HI flux from the neighboring spiral galaxy NGC 3338.

In the region of our survey, limited by the coordinates $RA = [10^h 30^m - 11^h 10^m]$, $DEC = [08^\circ 0' - 16^\circ 0']$, there are 37 other galaxies with radial velocities $V_{LG} < 2000$ km s⁻¹. Their list is given in Table 2. Some of these galaxies were previously considered to be members of the Leo-I group. Basic parameters of the galaxies are presented in Table 2 columns, their designations are the same as in Table 1. The data on apparent magnitudes and radial velocities are taken from NASA/IPAC Extragalactic Database⁵ (NED). The sources of data on galaxy distances are shown in Table 2 last column, reference to which are given at the note of the Table. The penultimate column of Table 2 gives the 21-cm line widths, W_{50} , at half-maximum intensity level ([Haynes et al. 2011](#)). As seen, the individual distances to most of the galaxies were measured via W_{50} using the Tully-Fisher relation in its usual (TF) or baryonic (bTF) version. The accuracy of the TF method is low for low-luminosity galaxies and gives a distance error of about 30%.

The Hubble “velocity–distance” diagram for galaxies in the area under consideration is shown in Fig. 2. It has a rather specific form, exhibiting a strong nonlinearity. The suggested group members around M 105 are characterized by an average velocity of $\langle V_{LG} \rangle = 654 \pm 24$ km s⁻¹ and average distance 10.8 Mpc. For the Hubble parameter $H_0 = 72$ km s⁻¹Mpc⁻¹ ([Freedman et al. 2001](#)), the M 105 group has a negative peculiar velocity of -124 ± 24 km s⁻¹ relative to the Local Group centroid. Galaxies with radial velocities $V_{LG} > 950$ km s⁻¹ from Table 2 are scattered across a wide range of distances from 16 Mpc to 40 Mpc with a median value of about 25 Mpc. Almost all of them have high negative peculiar velocities. A significant part of this effect is probably due to the TF-distance measurement errors, which scatter the galaxies horizontally by 5–8 Mpc. At the same time, the galaxy NGC 3389 with an accurate distance estimate of 25.2 ± 1.6 Mpc (marked by a star), measured by supernova luminosity ([Pejcha & Prieto 2015](#)), is present among them. It also has a high peculiar velocity, -652 ± 115 km s⁻¹. However, we should note that in an absolute frame of reference associated with the cosmic microwave background (CMB), the radial velocity of NGC 3389 amounts to 1656 km s⁻¹, i.e. the galaxy practically rests relative to the CMB reference frame. The kinematic situation

¹ <https://www.eso.org/sci/software/esomidas/>

² <https://www.legacysurvey.org/dr10/description/>

³ <http://www.sao.ru/lv/lvgdb>

⁴ <https://www.sdss3.org/dr10/algorithms/sdssUBVRITransform.php#Lupton2000>, <http://www.ned.ipac.caltech.edu>




Figure 1. Images of five new presumed M 105 satellites from the DECaLS sky survey. The size of the images is $2' \times 2'$. North is at the top, east is on the left.




Figure 2. Hubble diagram for galaxies in the vicinity of M 105. Galaxies with accurate distance estimates (TRGB and SBF) are shown by solid circles. Galaxies with less reliable distance estimates (TF and mem) are depicted by open circles. The distant background galaxy NGC 3389 with SN measured distance is marked by a star. The straight line corresponds to the unperturbed Hubble flow with the parameter $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

in this area of the sky can be presented in such a way so that the complex of distant galaxies with $D \gtrsim 20 \text{ Mpc}$ rests as a whole with respect to the CMB, while the LV, including the Local Group and

the Leo-I group moves in the direction of these distant galaxies with a velocity of about 600 km s^{-1} . This velocity, apparently, is due to the motion of the Local Volume directed away from the center of the expanding Local Void, as well as to the collective falling of galaxies towards the Great Attractor (Tully et al. 2008).

Although the group members around M 105 and the background galaxies do not significantly differ in their radial velocities, their separation by radial distance appears quite obvious.

The distribution of the presumed group members around M 105 in equatorial coordinates is presented in Fig. 3. Dwarf galaxies are shown by circles. Seven of the brighter galaxies with assigned NGC numbers are shown by squares. The circle centered on M 105 with a radius of $2^\circ 04'$ (or 385 kpc for an average 10.8 Mpc distance to the group) shows the group's virial region. The crosses show the positions of 37 background galaxies from Table. 2.

Among the distant background galaxies, a compact group of six galaxies can be isolated, located to the north-west of M 105. These galaxies are highlighted in bold in Table 2 and bolder crosses in Fig.3. The Sc galaxy NGC 3338 dominates in the group. The group is characterized by an average radial velocity $\langle V_{LG} \rangle = 1126 \text{ km s}^{-1}$, average distance 19.8 Mpc, average projected radius 150 kpc and an anomalously low radial velocity dispersion $\sigma_V = 23 \text{ km s}^{-1}$. The virial mass of the group is $(2.3 \pm 1.2) \times 10^{10} M_\odot$, and the K -band virial mass-to-luminosity ratio is equal to only $(4.1 \pm 2.2) M_\odot / L_\odot$, which indicates a deficit (or total absence) of dark matter in this group.

Note that groups of galaxies with a low M_T / L_K ratio are quite rare. Thus, in the list by Makarov & Karachentsev (2011) there are 215

groups of galaxies over the whole sky with an average radial velocity $V_{LG} < 3500 \text{ km s}^{-1}$ and the number of measured radial velocities $n_V > 5$. Among them, only 15 groups (i.e. 7%) have estimates $M_T/L_K < 10M_\odot/L_\odot$ and 5 groups have $M_T/L_K < 8M_\odot/L_\odot$. Such "skinny" groups deserve a more detailed study in order to understand the reason for the deficit of dark matter in them.

3 STRUCTURAL PROPERTIES OF THE GROUP OF GALAXIES AROUND M105

The presented data allow us to note the following features of the group of galaxies around M 105.

About a quarter of the galaxies among the presumable group members (21 out of 83) are located outside the virial radius. Their radial velocities and distances, where available, agree with the membership of these galaxies in the group. Therefore, the search area for new group members should not be limited to the region inside the virial radius. The radial velocities and distances of the periphery group members are especially important for estimating the total group mass.

More than half of the galaxies considered (47 out of 83) belong to early morphological types: Sph, E, S0, which have no signs of ongoing star formation. The group exhibits segregation by galaxy type: early-type objects have an average projected distance from the center $\langle R_p \rangle = 265 \pm 35 \text{ kpc}$, whereas in late-type galaxies (Irr, Tr, S) this quantity amounts to $400 \pm 45 \text{ kpc}$. Nonetheless, some early-type galaxies (NGC 3489, Dw 1033 + 1233) meet at distances $R_p > 600 \text{ kpc}$.

It is of interest to note that the M 105 group does not have any blue compact dwarfs (BCD), even though such objects, rich in gas, could be easily detected in the ALFALFA survey⁶. In some groups, for example, around M 81, BCD galaxies form a noticeable part of the population.

The distribution of the group galaxies by apparent and absolute B -magnitude is presented in the upper panel of Fig. 4. The solid line shows the luminosity function of the LV galaxies according to Driver & De Propriis (2003), which is well described by the standard Schechter (1976) function

$$F(M) = c[10^{0.4(M^*-M)}]^{1+\alpha} \times \exp[-10^{0.4(M^*-M)}] \quad (1)$$

with parameters $\alpha = -1.2$ and $M^* = -21.5 \text{ mag}$. In the range of $M_B < -11.0 \text{ mag}$, where the incompleteness of the sample is not yet affected, the probability of agreement between the observed and expected distributions is $p = 0.08$ according to the χ^2 test. The observed distribution shows a marked deficit of intermediate luminosity galaxies (similar to Magellanic Clouds) with absolute magnitudes $[-18^m, -15^m]$. Such a gap may be related to the peculiarities of the formation of this group or the galaxy merging process in it.

Flat or filamentary substructures were detected in the nearby groups around the Milky Way, M 31, Centaurus-A galaxies, consisting of dwarf satellites (Pawlowski & Kroupa 2013; Ibata et al. 2014; Libeskind et al. 2015). The panorama of the Leo-I group (Fig. 3) exhibits a chain of eight galaxies, located across the diagonal in the NW direction from the NGC 3351 ($161^\circ 0 + 11^\circ 7$) spiral. However, the radial velocities known for five of them do not show any specific alignment that could indicate a physical nature of this filament. The group exhibits also several dwarf pairs and triplets with mutual projected separations of the components $3'-5'$, i.e. $10-15 \text{ kpc}$. Some of them have close radial velocities (KK 94 and LeG 21) or a mutual

morphological similarity (dw 1052+1352 and dw 1050+1353). Such substructures deserve a more detailed investigation.

Schneider (1989) studied the central region of this group in the HI line and discovered nine intergalactic hydrogen clouds, which form a giant ring-like structure with a diameter of about 1° or approximately 200 kpc . The radial velocities of the HI clouds are in the $V_{LG} = 600-900 \text{ km s}^{-1}$ range, which implies their Leo-I group membership. Mihos et al. (2018) obtained deep frames of one HI cloud near a Sab galaxy NGC 3368 and detected a diffuse blue object of very low surface brightness (about $29^m/\text{sq.arcsec}$). This object, BST 1047 + 1156, is also noticeable in the ultraviolet images, taken with GALEX space telescope. The unusual dwarf system with a hydrogen mass of about $4.5 \times 10^7 M_\odot$ has a low star formation rate of about $5 \times 10^{-5} M_\odot/\text{year}$. About 99% of its mass is in gaseous state.

Sameer et al. (2022) studied spectra of 11 quasars located behind the M 105 group in a 7×6 square degree area. From the absorption lines corresponding to the radial velocity range of the Leo-I group, they determined that the intergalactic medium of the group has a high metallicity of the order of or higher than the solar metallicity. Based on this, the authors conclude that the huge ring-like structure made of HI clouds does not have the relic origin but was formed as a result of tidal interaction of the brightest group members. This result, as well as the presence of morphological segregation of galaxies, indicates that the Leo-I group is at a rather advanced stage of its dynamic evolution.

4 VIRIAL MASS OF THE M105 GROUP

Several methods of estimating the total mass of a galaxy group can be encountered in the literature. They are all based on the virial balance of kinetic and gravitational energy of the system. The differences between them are associated with different assumptions about the character of galaxy motion and different considerations of the projection factors.

Limber & Mathews (1960) proposed estimating the virial mass of a group consisting of n galaxies as

$$M_V = (3\pi/2) \times n \times (n-1)^{-1} \times G^{-1} \times \sigma_V^2 \times R_h, \quad (2)$$

where σ_V is the galaxy radial velocity dispersion relative to the center, R_h is the average harmonic distance between the galaxies in sky projection, and G is the gravitational constant. For the group around M 105 we obtained $\sigma_V = 136 \text{ km s}^{-1}$ and $R_h = 191 \text{ kpc}$ from 33 galaxies with measured radial velocities, which gives a group virial mass estimate of $M_V = 3.88 \times 10^{12} M_\odot$. Heisler et al. (1985) performed numerical n -body simulations of groups of galaxies with equal masses and found that for groups with $n \sim 10$, the virial mass estimator gives a typical error of mass about $10^{0.15}$.

According to Tully (2015), the total virial mass of a rich group may be determined as

$$M_T = (\beta \times \pi/2) \times G^{-1} \times \sigma_V^2 \times \langle R_p \rangle, \quad (3)$$

where $\langle R_p \rangle$ is the average projected group radius, and $\beta \approx 2.5 \pm 0.1$ addresses the transition from radial velocity dispersion to volumetric dispersion in the assumption of a weak anisotropy. For $\sigma_V = 136 \text{ km s}^{-1}$ and $\langle R_p \rangle = 318 \pm 35 \text{ kpc}$ we derive a mass estimate of $M_T = (5.35 \pm 1.25) \times 10^{12} M_\odot$.

Another "orbital" mass estimation method (Bahcall & Tremaine 1981), (Karachentsev & Kudrya 2014) assumes that small group members move around the massive center by arbitrarily oriented elliptical orbits with an average eccentricity of $\langle e^2 \rangle = 1/2$

⁶ <http://egg.astro.cornell.edu/>




Figure 3. Distribution of galaxies in the vicinity of M 105 in equatorial coordinates. The suggested M 105 group members are shown by circles. Seven of the brightest group members with NGC numbers are marked by squares. The crosses show background galaxies with radial velocities less than 2000 km s^{-1} . The 385 kpc ($2^{\circ}04$) circle corresponds to the group virial radius.

(Barber et al. 2014). In this case,

$$M_{\text{orb}} = (16/\pi) \times G^{-1} \times \langle \Delta V^2 \times R_p \rangle. \quad (4)$$

Based on 33 galaxies with known radial velocities we get $M_{\text{orb}} = (5.76 \pm 1.32) \times 10^{12} M_{\odot}$.

Note that in the case of a random Kepler orbit orientation with an eccentricity of $e^2 = 1/2$, the relative M_{orb} estimate variation due to the projection factors amounts to

$$\sigma(M_{\text{orb}})/\langle M_{\text{orb}} \rangle = [(256 - 10\pi^2)/10\pi^2]^{1/2} = 1.2625. \quad (5)$$

This value is in wonderful agreement with the relative variation of M_{orb} obtained by the scatter of individual M_{orb} estimates made for different group members: $\sigma(M_{\text{orb}}) = 7.48 \times 10^{12} M_{\odot}$ and $\sigma(M_{\text{orb}})/M_{\text{orb}} = 1.2986$.

According to the LV galaxies database, the total K_s -band luminosity of the M 105 group galaxies is $L_K = 3.23 \times 10^{11} L_{\odot}$. Adopting the obtained M_{orb} values as an optimal virial mass estimate of the group, we have the total mass-to-luminosity ratio $M_{\text{orb}}/L_K = (17.8 \pm 4.1) M_{\odot}/L_{\odot}$.

Tully (2015) found that the group virial mass and virial radius are connected through an empirical relation

$$(R_V/215) = (M_V/10^{12})^{1/3}, \quad (6)$$

where the radius R_V is expressed in kpc, and the mass is in solar units. For the derived orbital mass estimate, the virial radius is $R_V = 385 \text{ kpc}$ or $2^{\circ}04$ (shown by the circle in Fig. 3).

5 NEIGHBORING GROUP OF GALAXIES AROUND M66

Considering the distribution of galaxies in the local universe, Tully (1988) noted that the galaxy group associated with M 105 is part of the diffuse extended structure named Leo Spur. This cloud also hosts close groups around galaxies NGC 3521 ($D = 10.7 \text{ Mpc}$) and NGC 3115 ($D = 10.2 \text{ Mpc}$), the structures and populations of which are described in Karachentsev et al. (2022). These groups are separated from M 105 by angular distances 13° and 23° , correspondingly. The closest neighbor of M 105 is the group of galaxies around M 66 (NGC 3627), the distance to which is $11.12 \pm 0.56 \text{ Mpc}$ (Hoyt et al. 2019). The angular distance between the centers of these groups is equal to $7^{\circ}9$.

The group M 66, as is the case for group M 105, is located in the ALFALFA HI survey zone (Haynes et al. 2011), which contributed to measuring the radial velocities for many irregular dwarf members of the group. Searches for dwarf galaxies around M 66 were undertaken by various authors (Müller et al. 2018; Carlsten et al. 2022; Karachentsev & Kaisina 2022). To date, 35 M 66 candidate members, 22 of which have measured radial velocities, were detected in the RA = $167^{\circ}5$ – $174^{\circ}0$, DEC = $10^{\circ}0$ – $16^{\circ}0$ region.

Data on the presumable M 66 group members are given in Table 3. Its structure and designations of the presented parameters are the same as in Table 1. Data on apparent magnitudes, morphological types, and radial velocities of galaxies are taken from the latest version of the LV database. The sources of data on galaxy distances are shown in the Table last column, reference to which are given at the note of the Table.

The distribution of galaxies in the area recently searched by us




Figure 4. Luminosity function for the suggested members of M105 (upper panel) and M66 (lower panel). The solid line on the upper panel indicates the luminosity function for the Local volume population.

(Karachentsev & Kaisina 2022) for new group members is shown in Fig. 5. Four bright galaxies having NGC numbers are marked by squares, the rest of the dwarf galaxies are shown by circles. The fact that this group has been studied in much less detail than M 105 comes to attention. Only one galaxy, M 66 itself, has an accurate distance estimate obtained by the TRGB method (there are 15 such galaxies in the M 105 group), although both groups are located practically at the same distance from the observer. A deficit of intermediate luminosity galaxies like the Magellanic Clouds, with the absolute magnitude interval $M_B = [-16^m, -18^m]$, is evident in the luminosity function for group M 66 (see lower panel in Fig. 4). Early type galaxies (Sph, dE, S0) make up 40% of the total population. Unlike M 105, the M 66 group has four blue compact dwarfs (BCDs).

In the considered region, 18 background galaxies are known, with radial velocities $V_{LG} < 2000 \text{ km s}^{-1}$. These galaxies are marked by crosses in Fig. 5. Distance estimates were obtained for 13 of them based on the Tully-Fisher relation (Karachentsev et al. 2015). Given the large scatter of distance estimates for the background galaxies, their median value is equal to 22 Mpc. Comparing it to the median radial velocities $\langle V_{LG} \rangle = 1150 \text{ km s}^{-1}$ and $\langle V_{CMD} \rangle = 1590 \text{ km s}^{-1}$, we conclude that the conglomerate of galaxies behind the M 66 group approaches the Local Volume with a typical velocity of about 500 km s^{-1} , but rests relative to the cosmic microwave background. A similar situation also takes place for the galaxies behind the M 105 group, as was noted above.

According to the data in Table 3, the M 66 group is characterized by the following parameters: average radial velocity $\langle V_{LG} \rangle = 646 \text{ km s}^{-1}$, radial velocity dispersion $\sigma_V = 135 \text{ km s}^{-1}$, average projected separation of the members $\langle R_p \rangle = 270 \text{ kpc}$, and the total orbital mass estimate $M_{orb} = (5.94 \pm 1.50) \times 10^{12} M_\odot$. For the total group luminosity $L_K = 3.83 \times 10^{11} L_\odot$, the total mass-to-

luminosity ratio is $M_T/L_K = (15.6 \pm 3.9) M_\odot/L_\odot$, which coincides within the margin of error with the same ratio for group M 105. The virial radius of the M 66 group determined from expression (6) is $R_V = 390 \text{ kpc}$ or $2^\circ.01$. It is shown in Fig. 5 by a circle centered on M 66.

Besides the virial radius, the zero velocity sphere radius R_0 , where the dynamic influence of the group dominates over general cosmological expansion, is an important group parameter. For a typical relation $R_0 \approx 3.5 \times R_V$, the radius R_0 for both neighboring groups is $7'.1$. A fragment of an arc of radius R_0 around the M 105 group center is shown in Fig. 5 in dashes. The mutual distance between the centers of these groups, $7'.9$, turns out to be somewhat larger than their R_0 radii. Therefore, future merging of both groups into a single system appears unlikely.

We should note that the M 66 group, as is the case with M 105, shows signs of active galaxy interaction. In the M 66 group, the second brightest spiral galaxy NGC 3628 exhibits strong distortions at the periphery. A tidal tail of very low surface brightness stretches to the east of NGC 3628 (Jennings et al. 2015), the length of which reaches $42'$ or 135 kpc . This structure is well seen also in the HI line (Stierwalt et al. 2009). The dwarf galaxy dw 1120 + 1337 = NGC 3628, UCD 1 is the brightest part of this tail and was probably destroyed by the tidal influence of NGC 3628. The far end of the eastern tail hosts an irregular dwarf galaxy with low surface brightness AGC 215414, the hydrogen mass of which is 25 times higher than its stellar mass (Nikiel-Wroczyński et al. 2014).

The main parameters of the M 66 group in comparison with those for the M 105 group are given in Table 4.

6 CONCLUDING REMARKS

The group of galaxies around M 105 is one of the richest groups in the Local Volume. Unlike most other LV groups, not one but several galaxies dominate in it with a luminosity comparable to that of the Milky Way. The group is conveniently located in the “blind” HI strip survey carried out with the Arecibo radio telescope, which allowed radial velocities to be measured for many dwarf members of this system. In recent years the population of the group has significantly increased due to searches for new dwarf galaxies in the images of the DECaLS multi color digital sky survey. In addition to the searches for new objects in the group virial zone (Carlsten et al. 2022), we extended the search area to distant outskirts of the group and discovered another five candidate group members.

Currently the Leo-I group includes 83 presumable members, 33 of which have measured radial velocities. Using the projected separations and radial velocities of these galaxies, we determined their total group mass using four different methods. The estimates turned out to be in the $(3.9\text{--}5.8) \times 10^{12} M_\odot$ range. Early type galaxies without signs of star formation exhibit a known effect of stronger concentration towards the group center. The presence of small multiple systems is noticeable in Leo-I, but no large co-planar structure has been detected.

Another group of galaxies around M 66 neighbors the M 105 group. Both the groups are separated from the observer by similar distances, $10.8 \pm 0.5 \text{ Mpc}$ and $11.1 \pm 0.5 \text{ Mpc}$, correspondingly. The similarity of groups M 105 and M 66 also manifests itself in other parameters: the average radial velocity, the radial velocity dispersion, the average projected radius, the total luminosity, and the orbital mass estimate. Both the groups have a luminosity function gap in the absolute magnitude interval $[-15^m, -18^m]$, corresponding to the luminosity of the Magellanic Clouds. At the same time, the




Figure 5. Distribution of galaxies in the vicinity of M 66. The symbols for the suggested group members and background galaxies are the same as in Fig. 3. The circle corresponds to the virial radius 390 kpc ($2^{\circ}.01$). The dashed arc on the right shows the zero velocity sphere radius for the neighboring M 105 group.

number population of the M 66 group turned out to be twice smaller than that of M 105, which is in no way due to observational selection.

In the background of the M 105 group, we noted a group of 6 galaxies with an unusually low virial mass-to-luminosity ratio, $M_T/L_K = (4.1 \pm 2.2)M_{\odot}/L_{\odot}$, which indicates a deficit or total absence of dark matter in this group.

The conglomerate of distant background galaxies with $V_{LG} = 1000\text{--}2000 \text{ km s}^{-1}$ radial velocities behind both groups has a median distance of $D \sim 25 \text{ Mpc}$ and 22 Mpc, correspondingly. The collection of distant galaxies behind the group is in a state of rest relative to the cosmic microwave background, and both the M 105 and M 66 groups move towards the distant background with a characteristic peculiar velocity of about 600 km s^{-1} .

ACKNOWLEDGEMENTS

We are grateful to the anonymous referee for his/her constructive comments that helped to improve the paper. This work has made use of the DECaLS sky survey, The Arecibo Legacy Fast ALFA (ALFALFA), the NASA/IPAC Extragalactic Database, and the revised version of the Local Volume galaxy database. The Local Volume galaxies database has been updated within the framework of grant 075 – 15 – 2022 – 262 (13.MNPMU.21.0003) of the Ministry of Science and Higher Education of the Russian Federation.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY

The data underlying this article are available in the Local Volume galaxy database <https://www.sao.ru/lv/lvgdb/>. The additional data will be shared on reasonable request to the corresponding author.

REFERENCES

- Abazajian, K.N., Adelman-McCarthy, J.K., Agüeros, M.A., Allam, S.S., et al. 2009, *ApJS*, 182, 543
- Bahcall, J. N. & Tremaine, S., 1981, *ApJ*, 244, 805
- Barber, C., Starkenburg, E., Navarro, J.F., et al. 2014, *MNRAS*, 437, 959, <https://doi.org/10.1093/mnras/stt1959>
- Bottinelli, L., Gouguenheim, L., Paturel, G., & de Vaucouleurs, G., 1984, *A&A*, 56, 381
- Carlsten, S.G., Greene, J.E., & Beaton, R.L., 2022, *ApJ*, 933, id.47
- Carlsten, S.G., Greco, J.P., Beaton, R.L., & Greene, J.E., 2020, *ApJ*, 891, 144
- Cohen, Y., van Dokkum, P., Danieli, S., Romanowsky A.J., et al. 2018, *ApJ*, 868, 96
- Dey, A., Schlegel, D.J., Lang, D., et al., 2019, *AJ*, 157, 168
- Driver, S. & De Propris, R., 2003, *ASpSci*, 285, 175
- Ferrarese, L., Mould, J.R., Kennicutt, R.C., Jr., Huchra, J., et al., 2000, *ApJ*, 529, 745
- Ferguson, H.C. & Sandage, A., 1990, *AJ*, 100, 1
- Flint, K., Metevier, A.J., & Bolte, M., 2001, *ApJS*, 134, 53
- Freedman W.L., Madore, B.F., Gibson B.K., Ferrarese L., et al. 2001, *ApJ*, 553, 47
- Haynes, M.P., Giovanelli, R., Martin, A.M., et al., 2011, *AJ*, 142, 170
- Heisler, J., Tremaine, S., & Bahcall, J.N., 1985, *ApJ*, 298, 8
- Hoyt, T.J., Freedman, W.L., Madore, B.F., et al. 2019, *ApJ*, 882, 150
- Ibata, R.A., Ibata, N.G., Lewis, G.F., et al. 2014, *ApJ*, 784L, 6
- Jennings, Z.G., Romanowsky, A.J., Brodie, J.P., et al. 2015, *ApJ*, 812L, 10
- Kaisina, E.I., Makarov, D.I., Karachentsev, I.D., & Kaisin, S.S., 2012, *AstBu*, 67, 115
- Karachentsev, I.D., & Kaisina, E.I., 2022, *AstBu*, 77, 372

- Karachentsev, I.D., Makarova, L.N., Anand, G.S., & Tully, R.B., 2022, *AJ*, 163, 234
- Karachentsev, I., & Kashibadze, O., 2021, *Astr. Nachr.*, 342, 999
- Karachentsev, I.D., Nasonova, O.G., & Karachentseva, V.E., 2015, *AstBu*, 70, 1
- Karachentsev, I.D., & Kudrya, Y.N., 2014, *AJ*, 148, 50
- Karachentsev, I.D., Makarov, D.I., & Kaisina, E.I., 2013, *AJ*, 145, 101
- Karachentsev, I.D., & Karachentseva, V.E., 2004, *Astronomy Reports*, 48, 267
- Lee, M.G., & Jang I.S., 2016, *ApJ*, 822, 70
- Libeskind, N.I., Hoffman, Y., Tully, R.B., et al. 2015, *MNRAS*, 452, 1052
- Limber, D.N., & Mathews, W.G., 1960, *ApJ*, 132, 286
- Makarov, D.I., & Karachentsev, I.D., 2011, *MNRAS*, 412, 2498
- Mihos, J.C., Carr, C.T., Watkins, A.E., et al. 2018, *ApJ*, 863L, 7
- Müller, O., Jerjen, H., & Binggeli, B., 2018, *A& A*, 615A, 105
- Nikiel-Wroczyński, B., Soida, M., Bomans, D.J., & Urbanik, M., 2014, *ApJ*, 786, 144
- Pawlowski, M.S., & Kroupa, P., 2013, *MNRAS*, 435, 2116
- Pejcha, O., & Prieto, J.L., 2015, *ApJ*, 799, 215
- Sameer, Charlton, J.C., Kacprzak, G.G., et al. 2022, *MNRAS*, 510, 5796
- Schechter, P., 1976, *ApJ*, 203, 297
- Schneider, S.E., 1989, *ApJ*, 343, 94
- Schombert, J.M., Pildis, R.A., & Eder, J.A., 1997, *ApJS*, 111, 233
- Shaya, E.J., Tully, R.B., Hoffman Y., & Pomarède D., 2017, *ApJ*, 850, 207
- Sorce, J. G., Tully, R. B., Courtois, H. M., Jarrett, T. H., et al. 2014, *MNRAS*, 444, 527
- Springob, C.M., Masters, K.L., Haynes, M.P., Giovanelli, R., et al. 2009, *ApJS*, 182, 474
- Stierwalt, S., Haynes, M.P., Giovanelli, R., et al. 2009, *AJ*, 138, 338
- Tonry, J.L., Dressler, A., Blakeslee, J.P., Ajhar, E.A., et al. 2001, *ApJ*, 546, 681
- Tully, R.B., 2015, *AJ*, 149, 54
- Tully, R.B., Courtois, H.M., Dolphin, A.E., Fisher, J.R., et al. 2013, *AJ*, 146, 86
- Tully, R.B., Shaya, E.J., Karachentsev, I.D., et al. 2008, *ApJ*, 676, 184
- Tully, R.B., 1988, *Nearby Galaxies Catalog*, Cambridge University Press, Cambridge

Table 1: Suggested members of the M 105 group.

Name	RA J2000.0 Dec. deg	B_I mag	Type	V_{LG} km s ⁻¹	D Mpc	Method	R_p kpc	Refs
Dw1033+1213	158.372 +12.229	22.1	Sph	-	10.8	mem	662	
NGC3299	159.099 +12.707	13.3	Sdm	453	10.8	mem	525	
AGC205165	159.270 +15.337	16.4	Irr	586	10.8	mem	716	
dw1037+09	159.420 +09.105	17.9	Irr	-	10.8	mem	804	
Dw1038+1158	159.520 +11.973	20.5	Sph	-	10.8	mem	462	
LeG04	159.917 +12.734	18.7	Tr	-	10.8	mem	376	
LeG05	159.929 +12.635	16.8	Tr	629	10.8	mem	370	
LeG06	159.982 +13.908	18.3	Irr	863	10.8	mem	440	
UGC05812	160.236 +12.473	15.5	Im	857	10.8	mem	314	
FS04	160.502 +12.335	15.7	Im	621	10.8	mem	269	
LeG09	160.643 +12.151	17.4	Sph	-	10.19	TRGB	253	(1)
dw1042+1359	160.689 +13.990	21.2	Sph	-	10.51	SBF	350	(2)
dw1043+1410	160.795 +14.180	20.1	Sph	-	11.92	SBF	366	(2)
dw1043+1415	160.957 +14.255	20.2	Sph	-	12.24	SBF	362	(2)
LeG10	160.979 +12.133	19.2	Irr	-	10.8	mem	197	
NGC3351 (M95)	160.990 +11.704	10.6	Sb	623	9.95	TRGB	241	(3)
LeG11	161.008 +15.589	18.8	Tr	-	10.8	mem	593	
LeG12	161.031 +11.534	18.5	Irr	-	10.8	mem	259	
dw1044+1359	161.108 +13.999	21.0	Sph	-	10.60	SBF	306	(2)
AGC205445	161.147 +13.940	16.4	Tr	490	11.13	SBF	293	(2)
dw1044+1351	161.159 +13.859	19.9	Sph	-	11.49	SBF	279	(2)
LeG13	161.239 +11.916	17.7	Irr	718	9.99	SBF	180	(2)
dw1045+14b	161.484 +14.227	20.0	Sph	-	10.8	mem	319	
dw1045+13	161.492 +13.548	19.8	Sph	-	10.8	mem	199	
LeG14	161.559 +12.961	18.7	Sph	740	10.00	TRGB	101	(1)
KK93	161.603 +14.024	18.0	Sph	-	9.51	TRGB	277	(1)
LeG16	161.626 +11.756	18.4	Sph	-	11.02	SBF	166	(2)
LeG17	161.672 +12.327	17.0	Sph	880	10.99	SBF	70	(2)
NGC3368 (M96)	161.691 +11.820	10.1	Sab	740	10.42	Cep	150	(4)
LeG18	161.720 +12.742	18.2	Irr	488	10.19	TRGB	52	(1)
LeG19	161.728 +12.789	18.8	Sph	-	11.68	SBF	57	(2)
KK94	161.739 +12.998	17.5	Tr	684	10.19	TRGB	87	(1)
dw1047+16	161.754 +16.146	18.3	Sph	-	10.8	mem	673	
LeG21	161.752 +12.960	19.2	Irr	696	11.56	SBF	80	(2)
dw1047+1153	161.775 +11.878	21.4	Sph	-	9.35	SBF	136	(2)
N3368dwTBG	161.784 +11.285	19.6	Sph	-	10.8	mem	244	
M96-DF7	161.805 +12.802	18.9	Sph	-	10.19	TRGB	50	(1)
DDO088	161.843 +14.069	14.4	Sm	431	10.00	TRGB	281	(5)
M96-DF2	161.919 +12.049	20.0	Sph	-	10.62	TRGB	100	(1)
dw1047+1258	161.926 +12.975	20.0	Sph	-	11.00	SBF	74	(2)
NGC3377	161.926 +13.986	11.2	E	536	10.42	TRGB	262	(3)
BST1047+1156	161.932 +11.934	20.0	Irr	818	10.8	mem	122	
NGC3379 (M105)	161.957 +12.582	10.2	E	774	10.23	TRGB	0	(6)
M96-DF1	162.055 +11.968	19.8	Sph	-	10.42	TRGB	116	(1)
NGC3384	162.070 +12.629	10.9	S0	556	9.42	TRGB	23	(3)
dw1048+1121	162.096 +11.352	20.5	Sph	-	10.8	mem	231	
dw1048+1154	162.104 +11.912	19.7	Sph	-	10.75	SBF	128	(2)
dw1048+13	162.149 +13.060	20.1	Sph	-	10.62	TRGB	96	(1)
dw1048+1259	162.217 +12.995	20.3	Sph	-	11.81	SBF	91	(2)
AGC200596	162.224 +14.124	15.7	dE	400	12.67	SBF	292	(2)
dw1049+12a	162.298 +12.792	19.8	Tr	-	11.91	SBF	74	(2)
dw1049+15	162.310 +15.973	18.8	Tr	-	10.8	mem	642	
dw1049+12b	162.358 +12.552	19.3	Sph	-	10.81	TRGB	73	(1)
LeG22	162.404 +11.351	18.0	Tr	-	10.8	mem	244	
dw1050+1352	162.515 +13.889	19.6	Sph	-	10.8	mem	264	
UGC05944	162.579 +13.272	15.6	dE	928	11.07	SBF	172	(2)
dw1050+1353	162.585 +13.896	19.2	Sph	-	10.8	mem	271	

Table 1 – continued

Name	RA J2000.0 Dec. deg	B_I mag	Type	V_{LG} km s ⁻¹	D Mpc	Method	R_p kpc	Refs
dw1050+1213	162.600 +12.218	21.3	Sph	-	11.19	SBF	136	(2)
KK96	162.613 +12.360	17.3	Sph	-	10.00	TRGB	127	(1)
dw1050+1236	162.649 +12.609	20.6	Tr	-	11.25	SBF	126	(2)
NGC3412	162.722 +13.412	11.4	S0	702	11.32	SBF	208	(7)
LSBC D640-11	162.756 +13.334	16.7	Sph	557	10.89	SBF	202	(2)
dw1051+11	162.766 +11.020	18.1	Sph	-	10.8	mem	327	
LeG26	162.838 +12.849	16.6	Sph	483	10.34	SBF	168	(2)
AGC205540	162.881 +14.115	18.0	Tr	691	10.8	mem	332	
AGC205544	163.020 +15.030	17.1	dE	692	10.8	mem	501	
AGC202456	163.081 +11.043	16.2	dE	669	10.8	mem	353	
LeG27	163.082 +14.708	18.6	Tr	-	10.8	mem	451	
LeG28	163.252 +10.378	18.3	Sph	-	10.8	mem	479	
Dw1053+1518	163.443 +15.311	21.3	Tr	-	10.8	mem	582	
dw1055+11	163.930 +11.969	17.9	Sph	-	10.8	mem	380	
LSBC D640-12	163.981 +12.338	18.4	Tr	699	10.8	mem	374	
LSBC D640-13	164.059 +12.009	16.9	Irr	840	10.8	mem	401	
LSBC D640-14	164.543 +11.998	18.5	Sph	-	10.8	mem	487	
AGC205278	164.717 +14.130	17.3	Irr	548	11.80	TF	585	(8)
dw1059+11	164.962 +11.427	19.2	Tr	-	10.8	mem	593	
NGC3489	165.077 +13.901	11.1	S0	556	12.08	SBF	624	(7)
Dw1100+1125	165.131 +11.431	21.5	Tr	-	10.8	mem	622	
LeG33	165.188 +14.172	18.6	Tr	-	10.8	mem	596	
LSBC D640-08	165.216 +13.881	17.0	Sph	-	10.8	mem	646	
dw1101+11	165.344 +11.753	19.7	Irr	-	10.8	mem	641	
CGCG 066-109	166.109 +11.756	16.2	Irr	629	10.30	TF	778	(8)
Dw1106+1250	166.541 +12.845	18.8	Tr	-	10.8	mem	843	

Note. (1) Cohen et al. (2018); (2) Carlsten et al. (2022); (3) Shaya et al. (2017); (4) Ferrarese et al. (2000); (5) LVGDB; (6) Lee & Jang (2016); (7) Tonry et al. (2001); (8) Karachentsev et al. (2015)

Table 2. Background galaxies with $V_{LG} < 2000 \text{ km s}^{-1}$ in the same area outlined in Fig. 3

Name	RA (J2000.0 DEC)	B, mag	V_{LG} , km s^{-1}	D , Mpc	Method	W_{50} km s^{-1}	Refs
AGC202244	157.920 +13.835	16.5	1141	34.3	TF	102	(1)
AGC202016	158.330 +10.190	19.1	1270	-		32	
AGC205161	158.524 +15.780	17.9	1081	-		114	
NGC 3279	158.678 +11.197	13.9	1236	32.2	TF	347	(2)
AGC202248	158.734 +11.492	17.5	1020	28.2	TF	62	(3)
LeG 03	158.953 +08.481	17.3	987	21.9	TF	70	(3)
FGC125a	159.370 +12.396	17.4	1178	25.0	TF	59	(1)
CGCG065-074	159.533 +10.381	14.4	1013	25.1	TF	178	(1)
AGC203080	160.421 +13.825	17.5	1127	23.1	mem	-	
NGC 3338	160.531 +13.747	11.4	1155	23.1	TF	339	(2)
AGC203082	160.610 +13.957	17.8	1137	17.5	TF	41	(1)
UGC 5832	160.702 +13.460	14.3	1081	18.6	TF	102	(1)
AGC200543	160.773 +13.510	16.2	1117	18.1	TF	70	(1)
NGC 3346	160.912 +14.872	12.6	1135	18.5	TF	162	(4)
AGC205270	161.291 +15.450	17.1	1083	-		51	
NGC 3389	162.116 +12.533	12.5	1159	25.2	SN	266	(5)
Mrk 1263	162.237 +12.195	15.7	1171	34.2	TF	125	(1)
AGC200600	162.249 +10.835	16.2	1782	38.4	TF	120	(2)
AGC200603	162.322 +12.422	15.7	1234	15.6	TF	68	(3)
PGC032376	162.327 +12.378	18.0	1200	-		25	
AGC202253	162.361 +12.258	17.5	1188	-		-	
AGC205197	162.428 +13.828	19.0	1190	22.7	bTF	42	(3)
AGC205198	162.507 +13.785	17.2	1168	19.1	TF	53	(1)
UGC 5948	162.658 +15.763	16.6	987	26.3	bTF	106	(1)
UGC 6014	163.428 +09.728	15.0	965	17.0	TF	94	(1)
SDSSJ10540	163.500 +09.831	17.5	1023	17.0	mem	-	
AGC202033	163.765 +14.093	18.9	1968	-		40	
AGC202260	164.409 +13.979	17.5	1078	28.8	bTF	92	(1)
NGC 3485	165.010 +14.842	12.7	1301	-		135	
AGC202040	165.757 +08.048	18.1	1194	37.6	bTF	96	(1)
AGC215256	165.860 +16.016	16.9	1102	37.5	TF	105	(1)
AGC219117	165.945 +08.572	18.7	1575	-		68	
NGC 3524	166.634 +11.385	13.4	1216	-		415	
AGC215262	166.648 +12.231	18.2	1461	36.0	bTF	63	(3)
UGC 6169	166.764 +12.060	14.6	1405	34.6	TF	241	(1)
CGCG 067-014	167.347 +10.834	15.3	1404	21.1	mem	66	
NGC 3547	167.483 +10.721	13.2	1438	21.1	TF	204	(6)

Note. (1) Karachentsev et al. (2015); (2) Springob et al. (2009); (3) this work;
 (4) Bottinelli et al. (1984); (5) Pejcha & Prieto (2015); (6) Tully et al. (2013)

Table 3. Suggested members of the M 66 group.

Name	RA(J2000.0)DEC deg	B mag	Type	V_{LG} km s ⁻¹	D Mpc	Method	R_p kpc	Refs
Dw1111+1338	167.896 +13.644	20.5	Sph		11.12	mem	430	
AGC215282	168.605 +15.534	16.5	Im	731	11.12	mem	565	
NGC3593	168.654 +12.818	11.9	S0a	492	10.80	TF	269	(1)
AGC202256	168.688 +12.648	17.5	Irr	490	11.00	TF	269	(1)
dw1114+1307	168.713 +13.132	20.1	Sph		11.12	mem	256	
IC2684	169.254 +13.099	16.2	Tr	451	11.12	mem	154	
dw1117+1250	169.435 +12.837	20.6	Sph		11.12	mem	123	
dw1118+1330	169.566 +13.514	20.6	Sph		11.12	mem	137	
dw1118+1233	169.676 +12.561	17.5	Sph		9.67	SBF	111	(2)
dw1118+1348	169.723 +13.805	19.2	Sph		9.86	SBF	170	(2)
NGC3623 (M65)	169.733 +13.092	10.3	Sa	666	12.19	TF	66	(3)
AGC215286	169.803 +14.327	16.8	Tr	867	11.70	TF	263	(4)
AGC202257	169.810 +11.952	16.5	Irr	719	11.70	bTF	206	(4)
AGC215354	169.816 +14.290	17.4	BCD	659	11.12	mem	256	
dw1119+1404	169.840 +14.076	16.9	Sph	354	9.48	SBF	214	(2)
Dw1119+1011	169.878 +10.199	18.1	BCD		11.12	mem	544	
NGC3627 (M66)	170.062 +12.992	9.7	Sb	579	11.12	TRGB	0	(5)
dw1120+1332	170.067 +13.549	18.3	Im		11.43	SBF	108	(2)
NGC3628	170.071 +13.590	10.3	Sb	710	10.52	TF	116	(6)
dw1120+1337	170.255 +13.607	14.9	Tr	681	11.12	mem	125	
dw1121+1411	170.372 +14.197	20.1	Sph		11.12	mem	240	
dw1121+1326	170.404 +13.447	19.9	Irr		8.69	SBF	109	(2)
AGC213436	170.600 +12.980	16.9	dE	491	11.66	SBF	101	(2)
IC2782	170.731 +13.441	15.0	dE	719	10.46	SBF	153	(2)
AGC215414	170.806 +13.714	17.8	Irr	746	11.12	mem	198	
IC2787	170.830 +13.630	15.3	BCD	576	11.06	SBF	190	(2)
IC2791	170.907 +12.896	17.1	Im	530	9.95	SBF	161	(2)
dw1123+1346	170.985 +13.778	20.1	Sph		11.12	mem	232	
AGC215415	171.144 +12.674	18.6	Tr	868	11.12	mem	213	
dw1125+1316	171.479 +13.269	17.3	dE		11.12	mem	273	
Dw1127+1346	171.804 +13.781	20.1	Sph		11.12	mem	357	
AGC213091	172.394 +10.810	18.5	Sph	600	11.12	mem	612	
KKH68	172.722 +14.145	16.6	Irr	753	8.50	TF	552	(1)
AGC215248	173.462 +14.054	17.9	BCD	808	9.90	mem	676	
KKH69	173.721 +11.021	16.6	Irr	741	7.40	TF	791	(1)

Note. (1) [Karachentsev et al. \(2013\)](#); (2) [Carlsten et al. \(2022\)](#); (3) [Tully et al. \(2013\)](#); (4) LVGDB; (5) [Hoyt et al. \(2019\)](#); (6) [Sorce et al. \(2014\)](#)

Table 4. Basic parameters of M 105 and M 66 group of galaxies.

Parameter	M 105 group	M 66 group
Number of known members	83	35
Average radial velocity, km s ⁻¹	654±24	646±9
Radial velocity dispersion, km s ⁻¹	136	135
Distance, Mpc	10.8±0.5	11.1±0.5
Average projected radius, $\langle R_p \rangle$, kpc	318	270
Total K -band luminosity, $10^{11} L_\odot$	3.2	3.8
M_{orb} , $10^{12} M_\odot$	5.76±1.32	5.94±1.50
R_{vir} , kpc	385	390
Total mass-to-luminosity ratio, $M_\odot/L_\odot$	17.8±4.1	15.6±3.9