

A search for transients in the Reionization Lensing Cluster Survey (RELICS): Three new supernovae

Miriam Golubchik^{1*}, Adi Zitrin¹, Justin Pierel², Lukas J. Furtak¹, Ashish K. Meena¹, Or Graur^{3,4}, Patrick L. Kelly⁵, Dan Coe^{2,6,7}, Felipe Andrade-Santos^{8,9}, Maor Asif¹, Larry D. Bradley², Wenlei Chen⁵, Brenda L. Frye¹⁰, Sebastian Gomez², Saurabh Jha¹¹, Guillaume Mahler^{12,13}, Mario Nonino¹⁴, Louis-Gregory Strolger², Yuanyuan Su¹⁵

¹Physics Department, Ben-Gurion University of the Negev, P.O. Box 653, Be'er-Sheva 84105, Israel

²Space Telescope Science Institute (STScI), 3700 San Martin Drive, Baltimore, MD 21218, USA

³Institute of Cosmology & Gravitation, University of Portsmouth, Dennis Sciama Building, Portsmouth, PO1 3FX, UK

⁴Department of Astrophysics, American Museum of Natural History, Central Park West and 79th Street, New York, NY 10024, USA

⁵School of Physics and Astronomy, University of Minnesota, 116 Church Street SE, Minneapolis, MN 55455, USA

⁶Association of Universities for Research in Astronomy (AURA) for the European Space Agency (ESA), STScI, Baltimore, MD, USA

⁷Center for Astrophysical Sciences, Department of Physics and Astronomy, The Johns Hopkins University, 3400 N Charles St. Baltimore, MD 21218, USA

⁸Department of Liberal Arts and Sciences, Berklee College of Music, 7 Haviland Street, Boston, MA 02215, USA

⁹Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

¹⁰Department of Astronomy/Steward Observatory, University of Arizona, 933 N. Cherry Avenue, Tucson, AZ 85721, USA

¹¹Department of Physics and Astronomy, Rutgers, the State University of New Jersey, 136 Frelinghuysen Road, Piscataway, NJ 08854-8019, USA

¹²Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE, UK

¹³Centre for Extragalactic Astronomy, Durham University, South Road, Durham DH1 3LE, UK

¹⁴INAF-Trieste Astronomical Observatory, Via Bassoni 2, 34124, Trieste, Italy

¹⁵University of Kentucky, 505 Rose Street, Lexington, KY 40506, USA

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

The Reionization Cluster Survey (RELICS) imaged 41 galaxy clusters with the *Hubble Space Telescope* (HST), in order to detect lensed and high-redshift galaxies. Each cluster was imaged to about 26.5 AB mag in three optical and four near-infrared bands, taken in two distinct visits separated by varying time intervals. We make use of the multiple near-infrared epochs to search for transient sources in the cluster fields, with the primary motivation of building statistics for bright *caustic crossing events* in gravitational arcs. Over the whole sample, we do not find any significant ($\geq 5\sigma$) caustic crossing events, in line with expectations from semi-analytic calculations but in contrast to what may be *naïvely* expected from previous detections of some bright events, or from deeper transient surveys that do find high rates of such events. Nevertheless, we find six prominent *supernova* (SN) candidates over the 41 fields: three of them were previously reported and three are new ones reported here for the first time. Out of the six candidates, four are likely core-collapse (CC) SNe – three in cluster galaxies, and among which only one was known before, and one slightly behind the cluster at $z \sim 0.6 - 0.7$. The other two are likely Ia – both of them previously known, one probably in a cluster galaxy, and one behind it at $z \approx 2$. Our study supplies empirical bounds for the rate of caustic crossing events in galaxy cluster fields to typical HST magnitudes, and lays the groundwork for a future SN rate study.

Key words: supernovae: general – galaxies: clusters: general – gravitational lensing: strong – stars: massive

1 INTRODUCTION

Lensing of transient sources, whether theoretically or observationally, has risen in interest in recent years due to the ability to teach us about the lensed sources, but also constrain the dark matter (DM) composition of the lens (see Oguri 2019 for review). Among these transient sources, for example, are Fast Radio Bursts (FRBs; e.g. Muñoz et al. 2016), Gamma Ray Bursts (e.g. Paynter et al. 2021), Supernovae (SNe; e.g. Kelly et al. 2015; Goobar et al. 2017; Rod-

ney et al. 2021), Gravitational Waves (e.g. Dai et al. 2020; Broadhurst et al. 2022), and Caustic Crossing Events of lensed stars (e.g. Miralda-Escude 1991; Venumadhav et al. 2017; Kelly et al. 2018; Diego et al. 2018; Meena et al. 2022b). Among the several transients described above, the most natural transients to look for in optical and near-infrared imaging of galaxy clusters are lensed supernovae and caustic crossing events.

Lensed SNe have been attracting much interest, in part since they are expected to be found to larger redshifts than non magnified SNe, thus potentially improving the constraints on the cosmological parameters through their contribution to the Hubble diagram. Several

* E-mail: golubmir@post.bgu.ac.il

SNe lensed by galaxy clusters have been found to date. The first few examples were found sufficiently far from the center of the lens or at a low enough redshift behind it so they were magnified but not multiply imaged. For example, [Patel et al. \(2014\)](#); [Nordin et al. \(2014\)](#) found three lensed SNe in the 25 cluster fields of the *Cluster Lensing And Supernova survey with Hubble* (CLASH; [Postman et al. 2012](#), see also [Graur et al. 2014](#)). In another example, [Rodney et al. \(2015\)](#) found a Type Ia SN behind the *Hubble Frontier Fields* (HFF; [Lotz et al. 2017](#)) cluster Abell 2744, which was then confronted with magnification estimates from various lens models (see also [Mahler et al. 2018](#)). The past few years have seen also the first examples of several multiply imaged SNe, long anticipated ([Refsdal 1964](#)). *Refsdal*, the first multiply imaged SN, was detected by [Kelly et al. \(2015\)](#) as an Einstein cross around a cluster galaxy in the CLASH/HFF cluster MACS J1149.5+2223 ([Ebeling et al. 2007](#)). Another image of the SN appeared a year later in a counter-image of the SN's host galaxy ([Kelly et al. 2016](#)), enabling a measurement of the expansion rate of the universe from the time delays ([Vega-Ferrero et al. 2018](#); [Grillo et al. 2018](#)). About a year later, another lensed SN was found, multiply imaged by a field galaxy ([Goobar et al. 2017](#)). [Rodney et al. \(2021\)](#) have found another SN multiply imaged by a foreground massive cluster lens, MACS J0138.0-2155 ([Ebeling et al. 2010](#)), with the next image predicted to appear in about two decades. More recently, [Chen et al. \(2022a\)](#) found a multiply imaged SN around a galaxy in the HFF cluster Abell 370. At a redshift of $z \sim 3$, this SN is the farthest one known ($z \approx 3$), allowing also an early view of the SN's explosion thanks to the time delays. In addition to the physics they enable us to study, such multiply imaged transients are also important as they allow us to probe and re-calibrate our lens modeling techniques (e.g. [Treu et al. 2016](#); [Zitrin 2021](#)).

The rate of SNe in galaxy clusters has also spurred much interest (e.g. [Gal-Yam et al. 2003](#); [Graham et al. 2008](#); [Mannucci et al. 2008](#); [Dilday et al. 2010](#); [Sharon et al. 2010](#); [Sand et al. 2011](#); [Maoz & Mannucci 2012](#); [Maoz & Graur 2017](#); [Freundlich & Maoz 2021](#)). The process of chemical or metal enrichment by SNe is not very well constrained observationally ([Sarkar et al. 2022](#)). One of the crucial ingredients to characterise it is the distribution of delay time between a burst of star formation and the explosions as SNe, especially for Type Ia – for which the delay-time distribution spans Myr to Gyr time scales ([Maoz & Graur 2017](#)), unlike core-collapse (CC) SNe, for example, which result from massive, short lived stars. Because most galaxies in clusters are red elliptical galaxies, most SNe we expect to see in clusters are thus Type Ia, with a higher CC to Ia fractions towards higher redshifts around when the clusters formed ($z \sim 3 - 4$). That said, [Graham et al. \(2012\)](#), for example, found several CC SNe even in low-redshift cluster galaxies showing that at least some low level star-formation is taking place also in these so called red-and-dead galaxies. In addition, the progenitor of Type Ia SNe is not unambiguously known. In particular, it is not clear if most Type Ia result from single or double degenerate CO white dwarfs (e.g. [Hillebrandt & Niemeyer 2000](#)). Since different progenitor scenarios involve different timescales that control the production rate of SN Ia events, the rate, and its dependence on the host stellar-population age, can help discriminate between these models ([Maoz & Mannucci 2012](#)). The rate of different types of SNe in cluster fields, as a function of redshift, is thus of high importance.

The detection of the first multiply imaged SN *Refsdal* ([Kelly et al. 2015](#)) led also to the serendipitous detection of another type of transient, namely, a *caustic crossing event*. In follow-up observations of the SN, a new transient was seen atop the expected position of the lensing critical curve, where the magnification gets extremely high ([Kelly et al. 2018](#)). The Spectral Energy Distribution (SED) and

properties of the transient matched well that of a star crossing the caustic, getting temporarily, extremely magnified ([Miralda-Escude 1991](#); [Diego et al. 2018](#)). This event then opened the door to observing cosmological stars throughout the universe. Indeed, a growing number of lensed stars have been observed since (e.g. [Chen et al. 2019](#); [Kurov et al. 2019](#); [Meena et al. 2022a](#); [Kelly et al. 2022](#); [Diego et al. 2022b](#)), with the highest redshift example being Earendel at $z \approx 6.2$ ([Welch et al. 2022a](#)). JWST now offers a deeper window to observing and studying such stars through cosmic time, and has already revealed various types of stars in its first few months of operation (e.g. [Chen et al. 2022b](#); [Diego et al. 2022a](#); [Meena et al. 2022b](#); [Pascale et al. 2022](#); [Welch et al. 2022b](#)).

In this work we report the results from a search for transients in *Hubble Space Telescope* (HST) images taken for the RELICS program. In RELICS, 41 clusters were imaged in seven optical and near-infrared bands, to about 26.5 AB magnitudes per band. The four-band near-infrared (NIR) imaging was repeated in two different epochs (see section 2), separated by different periods of time (usually a few weeks or months), allowing us to search for transient sources. Our main motivation for the work is to estimate the rate of caustic crossing events in the survey. Although dedicated surveys with HST (e.g. the Flashlights program; [Kelly et al. 2022](#)) and the first clusters imaged with JWST (see works mentioned above) imply relatively high rates to $\sim 29 - 30$ AB, the first lensed stars were discovered at a level of 26-27 AB magnitudes ([Kelly et al. 2018](#); [Chen et al. 2019](#); [Kurov et al. 2019](#)), suggesting that RELICS observations should be sufficient for detecting at least *some* bright caustic crossing events. The observed rate depends, however, on various observational and physical properties such as filter choice, the background stellar mass density, the source radius-, luminosity-, and mass-functions, as well as properties of the lenses; which is why it is important to constrain.

The paper is organised as follows. We describe the data and observations in §2, and the methods in §3. The results are presented and discussed in §4, and the work is concluded in §5. Throughout we use a standard flat Λ CDM cosmology with $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_\Lambda = 0.7$, and $\Omega_m = 0.3$. All magnitudes quoted are in the AB system ([Oke & Gunn 1983](#)) and all quoted uncertainties represent 1σ ranges unless stated otherwise.

2 OBSERVATIONS AND DATA

Throughout this work we use data obtained in the framework of the RELICS program. In this program 41 massive galaxy clusters were observed with the HST (PI: D. Coe), and the *Spitzer Space Telescope* (PI: M. Bradac), with the goal of detecting gravitationally lensed arcs, bright high-redshift galaxies ([Salmon et al. 2020](#); [Strait et al. 2021](#)), as well as various transients (e.g. see the SNe listed in [Coe et al. 2019](#) and the first spectacular detection of a lensed star at $z \approx 6.2$ in [Welch et al. 2022a](#)). All targets were observed (from two separate epochs combined) to about 26.5 magnitudes in 7 HST bands: F435W, F606W, F814W with the *Advanced Camera for Surveys* (ACS), and F105W, F125W, F140W, and F160W, with the *Wide Field Camera Three* (WFC3). For some clusters, previous HST observations were used as well, as detailed in [Coe et al. \(2019\)](#).

The RELICS data products include reduced and colour images, photometric catalogs generated with SExtractor ([Bertin & Arnouts 1996](#)) and photometric redshifts computed with the Bayesian Photometric Redshifts tool (BPZ; [Benítez et al. 2004](#); [Coe et al.](#)




Figure 1. Examples of caustic crossing arcs out of 47 arcs inspected in RELICS in which one may expect to observe lensed stars. For each example arc we present, from left to right, a colour image, a combined image of all WFC3 bands for the first epoch, a similar image for the second epoch, and their difference image. The white bar in each image represents a scale of 2 arcsec. We find no clear detection of caustic crossing events in the RELICS survey. Left (from top to bottom): ACT-CL J0102–49151, A2813, ACT-CL J0102–49151, MACS J0417.5–1154, PLCK G004.5–19.5. Right (from top to bottom): AS295, A2813, MACS J0035.4–2015, CL J0152.7–1357, MACS J0553.4–3342.

2006). These are publicly available on the RELICS website¹. We refer the reader to Coe et al. (2019) for additional details on the HST data reduction and catalog construction.

3 METHODS

3.1 Image Subtraction

Due to the fact that the final drizzled images of each epoch in each band are not part of the RELICS data products, we assemble an image for each epoch in each of the bands using the raw ‘.flt’ files. For each cluster we use observations made with the WFC3 camera in four different bands (F105W, F125W, F140W, F160W). For each epoch, all exposures of the same band are aligned using TweakReg and combined into a final image using AstroDrizzle (Koekemoer et al. 2011). Both functions are part of the DrizzlePac software package available online. An automatic procedure is used to determine optimal values for the ‘conv_width’ and ‘threshold’ parameters of the TweakReg function, minimizing the *rms* of the offsets in *x* and *y*. In several cases the initial offsets of the data were large and hence we used the ‘search_radius’ parameter for optimal alignment. We

refer the reader to the DrizzlePac documentation² for further details about these functions. The two final images are subtracted, resulting in a difference image in which transients are searched. We use once more the AstroDrizzle function to calculate the total errors for each epoch, and finally a signal-to-noise map is calculated.

Note that for each cluster and for each epoch we also create a deeper (~ 27 AB) image by a weighted sum of the four WFC3/IR filters, and generate difference images from these “IR-combined” images as well.

3.2 Transients Detection

We search for transients in the difference images by eye. The systematic search is initially done on the F140W difference image for all RELICS clusters. Using a simple script, each difference image (typically $2' \times 2'$) is split into squares with an overlap of a few arcseconds between sub-frames. For each initial candidate identified, zoomed-in stamps in all available bands, including the IR-combined images, from both epochs and their difference images, are then generated for further inspection. Note that the search is done with a pixel scale of $\sim 0.12''$. This pixel scale is comparable to the WFC3

¹ <https://relics.stsci.edu/>

² <https://drizzlepac.readthedocs.io/en/latest/tweakreg.html>

Table 1. The six supernova candidates found in this work. *Column 1:* Candidate ID, indicating also the abbreviated name of the cluster (see §4 for full cluster names); *Columns 2 & 3:* RA and Dec of the supernova, in J2000.0 *Column 4:* Cluster redshift as listed in Coe et al. (2019); *Column 5:* Photo- z from the RELICS catalog for the most-probable host galaxy; *Columns 6 & 7:* Dates of first and second epoch; *Column 8:* Name of supernova from Coe et al. (2019) and references therein. The three supernovae for which no names are designated are the three newly found supernovae.

Candidate ID	R.A. (J2000)	Dec. (J2000)	Cluster Redshift	Host z_{phot}^a	First Epoch	Second Epoch	Name
RXCJ0142-SN1	25.740828	44.641610	0.341	3.53 [3.43–3.55]	2015-12-04	2016-01-14	—
AS295-SN1	41.3927500	−53.0308231	0.300	0.67 [0.53–0.72]	2016-08-30	2016-10-09	—
PLCKG171-SN1	48.2464374	8.3786862	0.270	2.71 [2.59–2.76]	2016-09-10	2016-10-21	Kukulkan
RXCJ0600-SN1	90.0510314	−20.1233171	0.460	0.38 [0.30–0.41]	2017-01-11	2017-02-15	William
A1763-SN1	203.8130885	41.0043610	0.228	1.75 [1.65–1.79]	2016-05-08	2016-06-17	Nebra
PLCKG287-SN1	177.7162202	−28.0932911	0.390	0.35 [0.30 – −0.37]	2017-02-21	2017-03-18	—

^a - Note this is the host photo- z as output by the RELICS pipeline. The final deduced type and redshift for each supernova, as well as an indication whether it is in or behind the cluster, is given in Table 2 and §4.

NIR point-spread-function (PSF) and is nominally double both the RELICS pixel scale and the pixel scale we use here for photometry and measurement (see §3.3 below). This means that the search was made on, effectively, somewhat smoothed images, which can help in detecting some events.

We also perform another manual search on the deeper, IR-combined images using SAOImageDS9, especially around gravitationally lensed arcs, in a more focused attempt to detect caustic crossing events. In practice we go over 47 arcs over all clusters (see Fig. 1 for examples). The mean photometric redshift of the arcs is $z \sim 1.8$, with a standard deviation of ~ 1.2 .

The above searches are made on both the difference image and on its negative, to minimise biases related to white-versus-black detections. For each cluster we then perform a more detailed inspection of all candidates that survived the previous steps. We accept a candidate as a reliable transient if it appears in the difference images of all four NIR bands and in the relevant epoch in all available bands, and corresponds in practice to a signal-to-noise ratio higher than about ~ 5 . This level was estimated by planting some point sources with different signal-to-noise ratios in some images, convolved with the PSF, and repeating the detection process. We also check whether the transient appears in RELICS colour images. In the scrutinizing process, the shape of the transient is also considered, as well as its location in the field, e.g., some events in the very edges were discarded as likely artefacts.

3.3 Photometry measurements

We register our WFC3/IR images for the two epochs, and the RELICS optical images to the *Gaia* DR3 (Gaia Collaboration et al. 2022) astrometry using Scamp (Bertin 2006) and re-sample the images with Swarp (Bertin 2010) onto the same $0.06''/\text{pix}$ grid as is used in RELICS. The photometry is measured on these images with the photutils package (v1.5.0; Bradley et al. 2022) in a circular aperture, with a radius of ~ 3 pixels of $0.06''$, and corrected for local background flux measured in a circular annulus around the source. Exact apertures and background radii were slightly refined manually for each candidate, to minimise contamination by the host galaxy. We also run the same photometry on the difference images themselves. This constitutes an important consistency check as the difference in flux between the photometry of the first and second epochs should match the fluxes measured in the difference image. All photometry is summarised in Table 3. The images, including the difference images,

as well as the photometry with relevant Julian date for each band, are made available online³.

3.4 Supernovae light curve fitting

We attempt to classify each of the SN candidates using the STARDUST2 Bayesian light curve classification tool (Rodney et al. 2014), which is built on the underlying SNCosmo framework and designed for classifying SNe using *HST*. STARDUST2 uses the SALT3-NIR model to represent Type Ia SNe (Pierel et al. 2022) and a collection of 42 spectrophotometric time series templates to represent CC SNe (27 Type II and 15 Type Ib/c). These CCSN templates comprise all of the templates developed for the SN analysis software SNANA (Kessler et al. 2009), derived from the SN samples of the Sloan Digital Sky Survey (Frieman et al. 2008; Sako et al. 2008; D’Andrea et al. 2010), *Supernova Legacy Survey* (Astier et al. 2006), and *Carnegie Supernova Project* (Hamuy et al. 2006; Stritzinger et al. 2009; Morrell 2012), and extended to the NIR by Pierel et al. (2018). With STARDUST2 we use a nested sampling algorithm to measure likelihoods over the SN simulation parameter space, including priors on dust parameters described in Rodney et al. (2014). Nested sampling is a Monte Carlo method that traverses the likelihood space in a manner that samples the Bayesian likelihood (Skilling 2004). The results of the fitting and classification procedure are summarised by Table 2 and Fig. 2. We find two likely SNe Ia, one in a cluster member and one the previously discovered SN *Nebra* (see Table 1), two CC SNe with 75% and 84% probabilities of being SNe Ib/c, and two likely SNe II. Of the CC SNe, all but one appear to be cluster members. These are further detailed in the results section §4.

4 RESULTS AND DISCUSSION

In our transient search we find no strong caustic crossing event candidates. This may be surprising at first sight, given that the first couple of lensed stars were detected to roughly similar magnitudes as those reached by RELICS. However, as we show below in section 4.1, this is in broad agreement with a rate expectation based on an order-of-magnitude calculation using some simple assumptions. We do find, however, six other prominent transients, likely SNe, detailed in section 4.2.

³ <https://www.dropbox.com/s/cl/fo/5yfm78d1kwizylh01gone/h?dl=0&r1key=k00fqpuj4yynamatfq182117a>

Table 2. Summary of the SN light-curve fits. *Column 1:* ID; *Column 2:* Redshift of the cluster; *Column 3:* Redshift range allowed in the light-curve fitting. In case the fit was performed using the cluster’s redshift a single value is shown. *Column 4:* Best-fit redshift from the light-curve fit, in case a range was allowed; *Column 5:* SN classification; *Column 6:* Probability of each SN type; *Column 7:* Time of peak (MJD); *Column 8:* Reduced chi-square; *Column 9:* Absolute B-band magnitude. For further details on the light-curve fitting procedure see section 3.4.

Candidate ID	Cluster z	z bounds	Fitted z	Classification	Probability	Peak	Reduced chi-square	Absolute mag (B-band)
					Ia–II–Ib/c			
RXCJ0142-SN1	0.341	0.341	–	II	0.0–1.0–0.0	57305	11.2	–17.2
AS295-SN1	0.300	[0.53–0.72]	0.64 ± 0.02	Ib/c	0.25–0.0–0.75	57542	1.2	–18.4
PLCKG171-SN1	0.270	0.270	–	II	0.0–1.0–0.0	57671	13.2	–16.0
RXCJ0600-SN1	0.460	0.460	–	Ia	1.0–0.0–0.0	57727	9.6	–18.9
A1763-SN1	0.228	[1.65–2.0]	1.96 ± 0.04	Ia	0.79–0.21–0.0	57553	1.0	–20.1
PLCKG287-SN1	0.390	0.390	–	Ib/c	0.0–0.23–0.77	57728	0.6	–18.3




Figure 2. The best fit model for each SN light curve. Each colour corresponds to a different *HST* filter, with measurements shown as points with errors (or upper-limit arrows) and maximum likelihood model realizations shown as solid lines. Shaded regions of the same colour correspond to 1σ uncertainties from the fitting procedure. The SN type shown is the same as in Table 2, and corresponds to each plotted model.

4.1 Caustic Crossing Events

We now briefly estimate the expected rate of events one should detect to the depth reached by RELICS, over 41 cluster fields.

To do this, we start by noting that caustic crossing events are biased towards bright and luminous stars. For an O/B-type star at redshift $z = 2$, with effective temperatures of $T_{\text{eff}} = 12,000 - 45,000$ K, a magnification in the range $\sim 20,000 - 50,000$ is needed for it to be visible at an apparent magnitude of $26.5 - 27.0$ AB in the *HST* filters. In the corrugated network forming around the macro-critical curve, thanks to point masses in the lens such as stars etc., the typical peak magnification for a stellar source of radius R is expected to be (e.g., Venumadhav et al. 2017; Oguri et al.

2018) $[10^4, 10^5] \times (R/10R_{\odot})^{-1/2}$, with the exact value depending on the macro-convergence, micro-lens density, micro-lens mass function etc., with an average frequency of peaks of about ~ 1 per year, per source star. For the above we assume a typical stellar mass density of $10 M_{\odot}/\text{pc}^2$. For a typical lensed arc we adopt a typical apparent brightness of ~ 26 AB/arcsec 2 . This is equivalent to $\sim 10^9 L_{\odot}/\text{arcsec}^2$ for a source at $z \sim 2$, which roughly translates into $\sim 1,000$ massive stars per arcsec 2 . For this order-of-magnitude calculation we assume a Salpeter IMF ranging between 0.1 and a $100 M_{\odot}$. Given most lensed arcs are typically blue star-forming galaxies, we neglect here the time evolution of the stellar population. It should however be acknowledged that the age, as well as other factors such as metallicity, may be important for assessing the true number of

expected stars, and our calculation remains crude in that sense. In particular, a more evolved stellar population will include less massive stars so our estimation is in that aspect, an upper limit. The typical relevant area of the corrugated network in the image plane is 0.1 arcsec^2 , equivalent to 0.0005 arcsec^2 in the source plane, assuming a magnification of 200 within the corrugated network. Assuming a velocity of 500 km s^{-1} , this implies that we expect roughly one crossing every two years in an arc.

Assuming that these events follow a Poissonian distribution, with an event lasting for ~ 3 days, we can estimate the probability of detecting at least one event in a given arc, in one visit, to be $\sim 0.4\%$. The probability of detecting at least one event over all the RELICS clusters, assuming one caustic-crossing arc per cluster, is $\sim 16\%$. Note that this is in principle, an upper limit: stars in the corrugated network that are sufficiently close to the main caustic, such that they are constantly sufficiently highly magnified, will only show minor fluctuations and will not be detected as transient sources in our difference images.

Outside the corrugated network, the magnification needed for O/B-type stars is typically too high to be reached in an individual micro-caustic crossing. Still, such a magnification can be reached if several microlenses sit near each other so that we have overlapping micro-caustics. Microlensing simulations (Meena et al. 2022c), show that the frequency of bright enough events in this regime is 1-2 orders of magnitude lower than on the corrugated network, hence contributing only little to the chances of seeing an event in RELICS.

Note however that some stars, such as various super- and hypergiants, may be intrinsically brighter than what we assume above, so they could be observed even with an almost order-of-magnitude lower magnification than considered here. One such example is Icarus, for which a peak magnification of ~ 3000 was sufficient for its detection. This means that in practice, there may be somewhat more events expected than estimated above, although due to their low numbers (or the small area in which such magnifications can be reached), we do not expect a significant contribution from such stars.

4.2 SN candidates

Our study yields 6 strong SN candidates summarised in Table 1. Three of the candidates were previously reported in Coe et al. (2019). We present the SNe candidates in Fig. 3. Four of the transients are detected to be fading and two are getting brighter between the first and second epoch. The SN light-curve fits are summarised in Table 2 and shown in Fig. 2. While in principle SNe in galaxy clusters can appear in the diffuse intracluster light, all six SNe we find here seem to lie in potential host galaxies. We in addition use the Bayesian Analysis of GaLaxy sEds tool (BEAGLE; Chevallard & Charlot 2016) to infer the properties of the host galaxies based on their broad-band photometry presented in Table 3. Note that we assume the same redshifts for both the light-curve fitting of the SNe and the SED-fitting of their host galaxies.

4.2.1 RXCJ0142.9+4438

We detect a transient in the cluster RXC J0142.9+4438 ($z = 0.341$; Fig. 3). The transient is detected to be fading between the first and the second epoch. The apparent host has a photometric redshift of $z_{\text{phot}} \approx 3.53$ in the RELICS catalog. If this redshift is correct, this would make the SN the farthest one detected by *HST*. Indeed, the host galaxy does seem to be bluish-greenish in the RELICS colour image (Fig. 3) suggesting it is not a typical cluster member, but

possibly behind the cluster. Our lens model implies that the galaxy would not be multiply imaged at that redshift, and indeed we do not identify any possible multiple images. The host galaxy does not seem to be stretched tangentially, as would be expected by the lensing shear in this case, thus disfavoring the high-redshift solution. For second redshift estimate, we also run an EAZY (Brammer et al. 2008) photometric-redshift fit, using the RELICS photometry for the host. We obtain a lower redshift solution, similar to the cluster redshift, namely a best-fit redshift and 68% confidence interval of $0.34 [0.26-0.43]$, which would suggest the host is in the cluster. The SN light curve fit is thus run with the cluster's redshift as input (see also Table 2). The photometry fits well a CC SN of Type II, with more than 90% probability. While future spectroscopy of the host would be useful for securing the host's redshift, we conclude that it is likely a CC SN of Type II at the redshift of the cluster. The SN was not known before and reported here for the first time, to the best of our knowledge. As for the host galaxy, we find it is relatively massive with $\log(M_{\star}/M_{\odot}) = 9.50 \pm 0.02$, relatively young with $t_{\text{age}} \approx 270 \text{ Myr}$, and has a moderate star-formation rate (SFR) of $\log(\psi/M_{\odot} \text{ yr}^{-1}) = 0.3 \pm 0.2$.

4.2.2 Abell S295

In the cluster Abell S295 ($z = 0.300$) we detect a transient fading between the first and the second epoch. The transient appears to be inside a host galaxy (see Fig. 3) with a photometric redshift of $z_{\text{phot}} \approx 0.67$. A photometric-redshift EAZY fit yields a similar redshift of 0.67, with a narrow 68% confidence interval of $[0.61-0.71]$. From the light curve fit, in which the SN's redshift is free to vary between ~ 0.5 and ~ 0.7 , we obtain it is likely (with 75% probability) a CC of Type Ib/c, at an approximated redshift of 0.64 ± 0.02 . The SN was not known before and reported here for the first time, to our knowledge. Our SED fit to the host galaxy suggests a massive ($\log(M_{\star}/M_{\odot}) = 10.4 \pm 0.1$), relatively young ($t_{\text{age}} \approx 160 \text{ Myr}$), dusty ($A_V = 3.3^{+0.4}_{-0.3}$) and heavily star-forming spiral with an SFR of $\log(\psi/M_{\odot} \text{ yr}^{-1}) = 1.5 \pm 0.3$.

4.2.3 PLCK G171.9-40.7

The transient detected in PLCK G171.9-40.7 ($z = 0.270$) gets brighter between the first and the second epoch and appears to lie inside a very faint host galaxy, with $z_{\text{phot}} \approx 2.71$ (see Fig. 3). Based on other multiple images known in this cluster (Cerny et al. 2018), we would expect the host to be multiply imaged, were it at $z \sim 2$ or above. However, we do not detect any counter image in the expected position based on the lensing symmetry, which suggests the galaxy is probably at a lower redshift. Preliminary investigation of this candidate in (Coe et al. 2019) classified the host as a cluster member galaxy. We run an EAZY photometric-redshift fit to the host photometry in the first epoch, where the SN's contribution is negligible, and obtain a relatively wide range of possible redshifts, with a single-template best-fit and 68% confidence interval of $0.737 [0.05-2.11]$, leaving the redshift ambiguous. For this redshift, our lens model suggests a magnification of $\mu \sim 2.3$. Given the wide photometric redshift range, for simplicity we run the light-curve fit assuming the cluster's redshift. The fit suggests that it is a CC SN of Type II, with over 90% probability. For the host galaxy, we find a low stellar mass of $\log(M_{\star}/M_{\odot}) = 7.58^{+0.08}_{-0.07}$, a very low SFR of $\log(\psi/M_{\odot} \text{ yr}^{-1}) = -1.61 \pm 0.3$ and a very young stellar age of $t_{\text{age}} \approx 13 \text{ Myr}$. Note however that given the faintness of the object and the relatively large uncertainties on the photometry, BEAGLE does

not find a very good fit at the cluster redshift assumed for the light-curve fit of the SN. In addition, since it is fit at a very low redshift, this galaxy does not have any rest-frame UV photometry to properly constrain its current population of massive stars which produce CC SNe.

4.2.4 RXC J0600.1-2007

The transient detected in the cluster RXC J0600.1-2007 ($z = 0.460$) seems to be fading between the first and the second epoch and appears to have exploded in the outskirts of a host galaxy (see Fig. 3) with a BPZ photometric redshift and 95% confidence interval of 0.383 [0.30–0.41], not too far from the cluster’s redshift. This candidate has been previously reported in Coe et al. (2019), where it was mentioned that the host was probably a cluster member. We perform the light-curve fit using the cluster redshift as input and obtain that this is likely a Type Ia SN, with more than 90% probability. The host galaxy is a massive ($\log(M_{\star}/M_{\odot}) = 10.96 \pm 0.02$), quiescent ($\log(\psi/M_{\odot} \text{ yr}^{-1}) = 0.1_{-1.3}^{+0.5}$) cluster galaxy of $t_{\text{age}} \approx 960$ Myr.

4.2.5 Abell 1763

We detect a transient in the cluster Abell 1763 ($z = 0.228$), which seems to get brighter between the first and the second epoch, and appears to lie inside a very faint host galaxy ($z_{\text{phot}} \approx 1.75$) see Fig. 3. This SN had been followed up with further observations, and was reported in some more detail in Rodney et al. (2016); Coe et al. (2019) where it was classified as a lensed Type Ia SN at $z \sim 2$. Initial magnification estimate yielded $\mu \sim 2$ (private communication). In this work, similar to all other cluster fields, we only use the first two epochs for our detection and analysis. Nevertheless, the light-curve fit, allowing a redshift range of [1.65–2] and based only on those two epochs, suggests a Type Ia SN at a redshift of $z \approx 2$, with 80% probability and in agreement with the previous analysis. We find the host galaxy has a relatively high stellar mass of $\log(M_{\star}/M_{\odot}) = 9.0 \pm 0.2$ and a low SFR $\log(\psi/M_{\odot} \text{ yr}^{-1}) = -0.7_{-0.9}^{+1.0}$ and stellar age of $t_{\text{age}} \approx 120$ Myr.

4.2.6 PLCK G287.0+32.9

We detect a transient in the cluster PLCK G287.0+32.9 ($z = 0.390$). The transient appears to be fading between the first and the second epoch and apparently lies in the halo of a cluster member host galaxy with $z_{\text{phot}} = 0.35$ [0.30 – 0.37] (see Fig. 3). This candidate has not been previously reported, to our knowledge. We adopt the redshift of the cluster for the light-curve fit and obtain that this is most likely a Type Ib/c CC SN (with over 75% probability). In this case, the host galaxy is a very massive ($\log(M_{\star}/M_{\odot}) = 11.76 \pm 0.01$), quiescent ($\log(\psi/M_{\odot} \text{ yr}^{-1}) = -1.1_{-0.6}^{+0.7}$) and old ($t_{\text{age}} \approx 10$ Gyr) cluster member galaxy.

5 CONCLUSION

In this work we present results from a search for transients in the 41 RELICS cluster fields. Motivated by the discovery of various caustic crossing events of lensed stars in imaging depths broadly similar to those obtained in RELICS (e.g., Kelly et al. 2018; Chen et al. 2019), and the rapidly increasing numbers of events detected in deeper observations (e.g., Kelly et al. 2022; Meena et al. 2022b; Pascale et al. 2022), our main goal was to characterise the caustic-crossing event rate in lensed arcs in the RELICS survey.

We utilise the fact that the NIR imaging took place in two distinct epochs to create difference images for each cluster and search for transients. We do not detect any prominent caustic-crossing event. We calculate the expected rate of events given the observational depth and find that the probability to detect at least one caustic crossing event in a cluster is $\sim 0.04\%$ per visit. Assuming one lensed arc per cluster we finally conclude a probability of $\sim 16\%$ to detect at least one caustic crossing event in RELICS. Our study thus supplies an empirical limit on the rate of bright caustic crossing events, to typical HST magnitudes, suggesting that indeed, a depth of ~ 26.5 AB is insufficient for detecting substantial numbers of lensed stars and that deeper observations are needed (as indeed successfully demonstrated in the references above).

Note that some lensed stars may also appear as quasi-persistent sources, in case they are sufficiently close to the caustic, in the so-called corrugated caustic network. One such famous example is Earendel, the farthest known lensed star detected at redshift $z \approx 6.2$ whose observed brightness has remained approximately constant over several years (Welch et al. 2022a), with only mild fluctuations. As another example, some types of stars such as Luminous Blue Variables can be bright enough for long periods of time at redshifts of $z \gtrsim 1$, even if farther away from the caustic. One such prominent example is *Godzilla* in the Sunrise Arc (Diego et al. 2022b). Since these are not expected to appear as transient sources we also search for possible point-sources near to where the critical curves pass in lensed arcs. We find no additional, prominent point sources in caustic-crossing arcs in RELICS, although we note also that this perhaps merits a further, designated examination.

While we do not find any lensed stars, our search yields six SN candidates. Three were previously known and reported in Coe et al. (2019), and three are, to the best of our knowledge, new ones presented here for the first time. Note also that Coe et al. (2019) found a few other candidates that are however mostly smaller or fainter-looking than our candidates, and not retained in our search (such detection differences can be attributed to different image-subtraction and source-identification procedures). The SN candidates we find here are classified using the Bayesian light-curve fitting code STARDUST2, and we obtain that two of the six SNe are Type Ia candidates, and four are core collapse: two Type II candidates, and two candidates that are most probably Type Ib/c. Four of the SNe go off in the cluster, likely in cluster galaxies, where two seem to be lensed and lie behind the cluster, the farthest of them at $z \approx 2$. In terms of type, three of the four SNe found inside the clusters are CC, and one is Ia. This ratio is perhaps a bit surprising given that the delay times from stellar formation to explosion is much longer for Ia than for CC, and that cluster galaxies are relatively early type, so-called red-and-dead galaxies with very little new star-formation. This may be partially explained by the fact that some of these SN host galaxies do not seem to be typical cluster members (i.e., red passive ellipticals). Additionally, the SZ-selected RELICS sample may be biased towards relatively massive, younger clusters in which galaxies possibly have some more star-formation compared to the average cluster galaxy at similar redshifts. We run BEAGLE to extract the physical parameters of each host, and obtain that for three of the CC SN hosts the SFR is between a few and a few dozen M_{\odot}/yr , although for the fourth it is very low ($\sim 0.03 M_{\odot}/\text{yr}$). For the two Ia SN hosts, the SFR is between ~ 0.1 and $\sim 3 M_{\odot}/\text{yr}$, i.e., on average lower than for the CC hosts, as may be expected. Given the low number of SNe in our sample, we however do not attempt to draw general conclusions regarding this ratio and the link to the host’s properties.

The SN rate in galaxy clusters, particularly as a function of redshift and type, is of high importance for a variety of studies from

characterising the metal enrichment history in the cluster, through estimating the quenching and ICL distribution timescales (Maoz & Graur 2017), to shedding light on the SN progenitor problem (Maoz & Mannucci 2012). The detection of several SNe in this work thus calls for a rate calculation. In a follow-up work we aim to estimate the completeness of SNe detection in our method, extend the search to the parallel fields accompanying each RELICS cluster field, and calculate the resulting SN rate.

ACKNOWLEDGEMENTS

We wish to thank the RELICS collaboration for data products that enabled this work. The BGU group acknowledges support by Grant No. 2020750 from the United States-Israel Binational Science Foundation (BSF) and Grant No. 2109066 from the United States National Science Foundation (NSF), and by the Ministry of Science & Technology, Israel. P.L.K. acknowledges support through NSF grant AST-1908823.

This work is based on observations made with the NASA/ESA Hubble Space Telescope obtained from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555. These observations are associated with program ID 15959. Support for program 15959 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. This work is also based on observations made with ESO Telescopes at the La Silla Paranal Observatory obtained from the ESO Science Archive Facility.

This research made use of *Astropy*,⁴ a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013; Price-Whelan et al. 2018) as well as the packages NumPy (van der Walt et al. 2011), SciPy (Virtanen et al. 2020), matplotlib (Hunter 2007), specutils (Earl et al. 2021), spectral-cube (Ginsburg et al. 2014) and some of the astronomy MATLAB packages (Ofek 2014).

⁴ <http://www.astropy.org>

Table 3. Aperture photometry for the six SN candidates found in this work and their host galaxies. For each SN we list the photometry in both the first epoch and the second epoch, as well as the photometry on the difference images in the near-IR for consistency. For the host galaxy we use the default (isophotal) photometry from the RELICS catalogs. In cases where the transient is not detected we note a lower bound corresponding to 1σ point-source AB magnitude limit.

AB magnitude in:	F435W	F606W	F814W	F105W	F125W	F140W	F160W
RXCJ0142-SN1							
Difference Image	–	–	–	24.42 ± 0.02	24.20 ± 0.03	24.29 ± 0.03	24.91 ± 0.03
Epoch 1	26.19 ± 0.07	–	23.42 ± 0.01	23.74 ± 0.02	23.46 ± 0.02	23.61 ± 0.02	23.90 ± 0.03
Epoch 2	–	24.76 ± 0.02	–	24.39 ± 0.04	24.25 ± 0.04	24.48 ± 0.06	24.40 ± 0.04
RXCJ0142-SN1-Host							
RELICS Data	23.34 ± 0.03	22.23 ± 0.01	21.63 ± 0.01	21.63 ± 0.01	21.63 ± 0.01	21.56 ± 0.02	21.50 ± 0.01
AS295-SN1							
2015 Jan. 23	> 28.95	> 29.35	> 28.17	–	–	–	–
Difference Image	–	–	–	25.72 ± 0.06	25.63 ± 0.09	25.74 ± 0.08	25.30 ± 0.04
Epoch 1	–	–	–	24.94 ± 0.04	25.12 ± 0.06	25.20 ± 0.06	24.97 ± 0.04
Epoch 2	–	–	–	25.69 ± 0.07	26.38 ± 0.19	26.23 ± 0.14	26.00 ± 0.09
AS295-SN1-Host							
RELICS Data	22.33 ± 0.02	21.64 ± 0.01	20.85 ± 0.01	20.59 ± 0.01	20.46 ± 0.01	20.41 ± 0.01	20.36 ± 0.01
PLCKG171-SN1							
Difference Image	–	–	–	24.94 ± 0.04	24.81 ± 0.06	24.84 ± 0.05	25.23 ± 0.06
Epoch 1	28.78 ± 0.94	–	26.81 ± 0.15	26.13 ± 0.10	26.49 ± 0.21	25.66 ± 0.09	25.96 ± 0.09
Epoch 2	–	24.77 ± 0.02	–	24.63 ± 0.03	24.69 ± 0.05	24.42 ± 0.04	24.81 ± 0.04
PLCKG171-SN1-Host							
RELICS Data	23.40 ± 0.13	22.65 ± 0.03	23.05 ± 0.01	22.94 ± 0.04	22.74 ± 0.05	22.70 ± 0.04	22.74 ± 0.03
RXCJ0600-SN1							
Difference Image	–	–	–	25.49 ± 0.05	25.67 ± 0.10	25.26 ± 0.06	25.67 ± 0.06
Epoch 1	27.76 ± 0.26	–	26.93 ± 0.12	24.89 ± 0.03	25.08 ± 0.06	24.95 ± 0.04	25.11 ± 0.04
Epoch 2	–	26.09 ± 0.04	–	25.80 ± 0.06	25.94 ± 0.11	26.42 ± 0.14	26.10 ± 0.09
RXCJ0600-SN1-Host							
RELICS Data	22.34 ± 0.03	20.514 ± 0.004	19.602 ± 0.002	19.204 ± 0.003	18.990 ± 0.004	18.850 ± 0.003	18.725 ± 0.002
A1763-SN1							
Difference Image	–	–	–	25.26 ± 0.04	24.92 ± 0.06	25.13 ± 0.06	25.33 ± 0.04
Epoch 1	26.78 ± 0.15	–	27.38 ± 0.19	26.35 ± 0.09	26.70 ± 0.21	26.25 ± 0.12	26.40 ± 0.09
Epoch 2	–	26.97 ± 0.09	–	24.94 ± 0.03	24.69 ± 0.04	24.77 ± 0.03	24.97 ± 0.03
A1763-SN1-Host							
RELICS Data	24.72 ± 0.06	24.75 ± 0.04	24.75 ± 0.06	24.22 ± 0.05	23.55 ± 0.02	23.76 ± 0.06	23.70 ± 0.02
PLCKG287-SN1							
2016 Aug. 3	–	> 29.35	> 27.99	–	–	–	–
Difference Image	–	–	–	25.96 ± 0.11	25.22 ± 0.07	25.98 ± 0.12	26.50 ± 0.13
Epoch 1	28.62 ± 0.66	–	–	25.36 ± 0.22	24.93 ± 0.18	25.60 ± 0.34	26.10 ± 0.61
Epoch 2	–	–	–	26.33 ± 0.45	26.63 ± 0.84	26.82 ± 1.07	27.64 ± 2.56
PLCKG287-SN1-Host							
RELICS Data	21.06 ± 0.02	20.67 ± 0.01	19.435 ± 0.003	18.575 ± 0.002	18.194 ± 0.002	18.045 ± 0.001	17.931 ± 0.003




Figure 3. The six SN detections. For each one, at the top row we present three colour images, one for each epoch from the WFC3 filters, and one including all ACS+WFC3 filters; Second row displays the difference image for each filter; Next two rows present the images from the two epochs. A scale of $1''$ is marked upon the image. The red arrow in each stamp points to the exact candidate location. In parentheses we designate for each SN candidate the best fitted type. The orientation of the figures is arbitrary.

DATA AVAILABILITY

The data used in this work are publicly available on the MAST archive and the RELICS website.

REFERENCES

- Astier P., et al., 2006, *A&A*, **447**, 31
- Astropy Collaboration et al., 2013, *A&A*, **558**, A33
- Benítez N., et al., 2004, *ApJS*, **150**, 1
- Bertin E., 2006, Automatic Astrometric and Photometric Calibration with SCAMP, p. 112
- Bertin E., 2010, SWarp: Resampling and Co-adding FITS Images Together, Astrophysics Source Code Library, record ascl:1010.068 (ascl:1010.068)
- Bertin E., Arnouts S., 1996, *A&AS*, **117**, 393
- Bradley L., et al., 2022, *astropy/photutils*: 1.6.0, doi:10.5281/zenodo.7419741, <https://doi.org/10.5281/zenodo.7419741>
- Brammer G. B., van Dokkum P. G., Coppi P., 2008, *ApJ*, **686**, 1503
- Broadhurst T., Diego J. M., Smoot G. F., 2022, arXiv e-prints, p. arXiv:2202.05861
- Cerny C., et al., 2018, *ApJ*, **859**, 159
- Chen W., et al., 2019, *ApJ*, **881**, 8
- Chen W., et al., 2022a, *Nature*, **611**, 256
- Chen W., et al., 2022b, *ApJ*, **940**, L54
- Chevallard J., Charlot S., 2016, *MNRAS*, **462**, 1415
- Coe D., Benítez N., Sánchez S. F., Jee M., Bouwens R., Ford H., 2006, *AJ*, **132**, 926
- Coe D., et al., 2019, *ApJ*, **884**, 85
- D'Andrea C. B., et al., 2010, *ApJ*, **708**, 661
- Dai L., Zackay B., Venumadhav T., Roulet J., Zaldarriaga M., 2020, arXiv e-prints, p. arXiv:2007.12709
- Diego J. M., et al., 2018, *ApJ*, **857**, 25
- Diego J. M., et al., 2022a, arXiv e-prints, p. arXiv:2210.06514
- Diego J. M., Pascale M., Kavanagh B. J., Kelly P., Dai L., Frye B., Broadhurst T., 2022b, *A&A*, **665**, A134
- Dilday B., et al., 2010, *ApJ*, **715**, 1021
- Earl N., et al., 2021, *astropy/specutils*: V1.5.0, doi:10.5281/zenodo.5721652, <https://doi.org/10.5281/zenodo.5721652>
- Ebeling H., Barrett E., Donovan D., Ma C., Edge A. C., van Speybroeck L., 2007, *ApJ*, **661**, L33
- Ebeling H., Edge A. C., Mantz A., Barrett E., Henry J. P., Ma C. J., van Speybroeck L., 2010, *MNRAS*, **407**, 83
- Freundlich J., Maoz D., 2021, *MNRAS*, **502**, 5882
- Frieman J. A., et al., 2008, *AJ*, **135**, 338
- Gaia Collaboration et al., 2022, arXiv e-prints, p. arXiv:2208.00211
- Gal-Yam A., Maoz D., Guhathakurta P., Filippenko A. V., 2003, *AJ*, **125**, 1087
- Ginsburg A., Robitaille T., Beaumont C., ZuHone J., 2014, Release Candidate 2 - includes yt interop, doi:10.5281/zenodo.11485, <https://doi.org/10.5281/zenodo.11485>
- Goobar A., et al., 2017, *Science*, **356**, 291
- Graham M. L., et al., 2008, *AJ*, **135**, 1343
- Graham M. L., et al., 2012, *ApJ*, **753**, 68
- Graur O., et al., 2014, *ApJ*, **783**, 28
- Grillo C., et al., 2018, *ApJ*, **860**, 94
- Hamuy M., et al., 2006, *PASP*, **118**, 2
- Hillebrandt W., Niemeyer J. C., 2000, *ARA&A*, **38**, 191
- Hunter J. D., 2007, *Computing in Science & Engineering*, **9**, 90
- Kaurov A. A., Dai L., Venumadhav T., Miralda-Escudé J., Frye B., 2019, *ApJ*, **880**, 58
- Kelly P. L., et al., 2015, *Science*, **347**, 1123
- Kelly P. L., et al., 2016, *ApJ*, **819**, L8
- Kelly P. L., et al., 2018, *Nature Astronomy*, **2**, 334
- Kelly P. L., et al., 2022, arXiv e-prints, p. arXiv:2211.02670
- Kessler R., et al., 2009, *PASP*, **121**, 1028
- Koekemoer A. M., et al., 2011, *ApJS*, **197**, 36
- Lotz J. M., et al., 2017, *ApJ*, **837**, 97
- Mahler G., et al., 2018, *MNRAS*, **473**, 663
- Mannucci F., Maoz D., Sharon K., Botticella M. T., Della Valle M., Gal-Yam A., Panagia N., 2008, *MNRAS*, **383**, 1121
- Maoz D., Graur O., 2017, *ApJ*, **848**, 25
- Maoz D., Mannucci F., 2012, *Publ. Astron. Soc. Australia*, **29**, 447
- Meena A. K., et al., 2022a, arXiv e-prints, p. arXiv:2211.01402
- Meena A. K., et al., 2022b, arXiv e-prints, p. arXiv:2211.13334
- Meena A. K., Arad O., Zitrin A., 2022c, *MNRAS*, **514**, 2545
- Miralda-Escudé J., 1991, *ApJ*, **379**, 94
- Morrell N. I., 2012, in Roming P., Kawai N., Pian E., eds, Vol. 279, Death of Massive Stars: Supernovae and Gamma-Ray Bursts. pp 361–362, doi:10.1017/S174392131201335X
- Muñoz J. B., Kovetz E. D., Dai L., Kamionkowski M., 2016, *Phys. Rev. Lett.*, **117**, 091301
- Nordin J., et al., 2014, *MNRAS*, **440**, 2742
- Ofek E. O., 2014, MAAT: MATLAB Astronomy and Astrophysics Toolbox, Astrophysics Source Code Library, record ascl:1407.005 (ascl:1407.005)
- Oguri M., 2019, *Reports on Progress in Physics*, **82**, 126901
- Oguri M., Diego J. M., Kaiser N., Kelly P. L., Broadhurst T., 2018, *Phys. Rev. D*, **97**, 023518
- Oke J. B., Gunn J. E., 1983, *ApJ*, **266**, 713
- Pascale M., et al., 2022, *ApJ*, **938**, L6
- Patel B., et al., 2014, *ApJ*, **786**, 9
- Paynter J., Webster R., Thrane E., 2021, *Nature Astronomy*, **5**, 560
- Pierel J. D. R., et al., 2018, *PASP*, **130**, 114504
- Pierel J. D. R., et al., 2022, *ApJ*, **939**, 11
- Postman M., et al., 2012, *ApJS*, **199**, 25
- Price-Whelan A. M., et al., 2018, *AJ*, **156**, 123
- Refsdal S., 1964, *MNRAS*, **128**, 307
- Rodney S. A., et al., 2014, *AJ*, **148**, 13
- Rodney S. A., et al., 2015, *ApJ*, **811**, 70
- Rodney S., et al., 2016, The Astronomer's Telegram, **9224**, 1
- Rodney S. A., Brammer G. B., Pierel J. D. R., Richard J., Toft S., O'Connor K. F., Akhshik M., Whitaker K. E., 2021, *Nature Astronomy*, **5**, 1118
- Sako M., et al., 2008, *AJ*, **135**, 348
- Salmon B., et al., 2020, *ApJ*, **889**, 189
- Sand D. J., et al., 2011, *ApJ*, **729**, 142
- Sarkar A., Su Y., Truong N., Randall S., Mernier F., Gastaldello F., Biffi V., Kraft R., 2022, *MNRAS*, **516**, 3068
- Sharon K., et al., 2010, *ApJ*, **718**, 876
- Skilling J., 2004, in Fischer R., Preuss R., Toussaint U. V., eds, American Institute of Physics Conference Series Vol. 735, Bayesian Inference and Maximum Entropy Methods in Science and Engineering: 24th International Workshop on Bayesian Inference and Maximum Entropy Methods in Science and Engineering. pp 395–405, doi:10.1063/1.1835238
- Strait V., et al., 2021, *ApJ*, **910**, 135
- Stritzinger M., et al., 2009, *ApJ*, **696**, 713
- Treu T., et al., 2016, *ApJ*, **817**, 60
- Vega-Ferrero J., Diego J. M., Miranda V., Bernstein G. M., 2018, *ApJ*, **853**, L31
- Venumadhav T., Dai L., Miralda-Escudé J., 2017, *ApJ*, **850**, 49
- Virtanen P., et al., 2020, *Nature Methods*, **17**, 261
- Welch B., et al., 2022a, *Nature*, **603**, 815
- Welch B., et al., 2022b, *ApJ*, **940**, L1
- Zitrin A., 2021, *ApJ*, **919**, 54
- van der Walt S., Colbert S. C., Varoquaux G., 2011, *Computing in Science Engineering*, **13**, 22

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.