

A note on real similarity to a diagonal dominant matrix

Zhiyong Sun, Brian D. O. Anderson, Wei Chen

^aDepartment of Electrical Engineering, Technische Universiteit Eindhoven, the Netherlands

^bSchool of Engineering, The Australian National University, Canberra ACT 2601, Australia

^cDepartment of Mechanics and Engineering Science, Peking University, Beijing 100871, China

Abstract

This note presents several conditions to characterize **real matrix similarity** between a Hurwitz matrix (and then more generally, a real square matrix) and a **diagonal dominant matrix**.

Keywords:

Hurwitz matrix, diagonal dominant matrix, real similarity.

1. Definitions and notations

- A real square matrix $A = \{a_{ij}\} \in \mathbb{R}^{n \times n}$ is **Hurwitz** if all its eigenvalues have negative real parts.
- *Row-diagonal dominant matrix:*
 - A real matrix $A = \{a_{ij}\} \in \mathbb{R}^{n \times n}$ is (non-strict) **row-diagonal dominant**, if the absolute value of each diagonal entry is greater than or equal to the sum of the absolute values of off-diagonal entries in that row; i.e., $|a_{ii}| \geq \sum_{j=1, j \neq i}^n |a_{ij}|, \forall i = 1, 2, \dots, n$.
 - If the absolute value of each diagonal entry is **strictly** greater than the sum of the absolute values of off-diagonal entries in that row, i.e., $|a_{ii}| > \sum_{j=1, j \neq i}^n |a_{ij}|, \forall i = 1, 2, \dots, n$, then we call the matrix A **strictly row-diagonal dominant**.
- Similarly, we can define a **column-diagonal dominant** matrix by following the above definition but using the column sums of off-diagonal entries. In this note, we call a matrix A **diagonal dominant** if it is either row-diagonal dominant or column-diagonal dominant.¹
- A real square matrix is a **Z-matrix** if it has nonpositive off-diagonal elements.

¹For the case of a complex matrix, the definition of a diagonal dominant complex matrix follows similarly but replacing ‘absolute value’ by ‘magnitude’ in the definition.

- A **Metzler** matrix is a real matrix in which all the off-diagonal components are nonnegative (equal to or greater than zero).
- A matrix $A \in \mathbb{R}^{n \times n}$ is called an **M-matrix**, if it is a Z-matrix and its eigenvalues have positive real parts.²
- For a real matrix $A = \{a_{ij}\} \in \mathbb{R}^{n \times n}$, we associate it with a **comparison matrix** $M_A = \{m_{ij}\} \in \mathbb{R}^{n \times n}$, defined by

$$m_{ij} = \begin{cases} |a_{ij}|, & \text{if } j = i; \\ -|a_{ij}|, & \text{if } j \neq i. \end{cases}$$

A given matrix A is called an **H-matrix** if its comparison matrix M_A is an M-matrix.

2. Problem of interest

In this note, we are interested in the following problem:

- When is a real Hurwitz matrix A real similar to a real diagonal dominant matrix B ?

and a more general problem on real similarity:

- When is a real square matrix A real similar to a diagonal dominant matrix B ?

By “a real matrix A is real similar to a real matrix B ” we mean there exists a **real** non-singular matrix P such that $B = PAP^{-1}$.

3. Real similarity with 2×2 real matrix

3.1. Case I: real matrix with real eigenvalues

We first give the following conditions to characterize real similarity for 2×2 real Hurwitz matrix with real eigenvalues.

Lemma 1. *Consider a 2×2 real Hurwitz matrix A and suppose its two eigenvalues are real, denoted by $\lambda_1 < 0$ and $\lambda_2 < 0$. Then there exists a real non-singular matrix P_1 such that A is real similar to a real matrix $B_1 = PAP^{-1}$:*

$$B_1 = \begin{bmatrix} \lambda_1 & \star \\ 0 & \lambda_2 \end{bmatrix} \quad (1)$$

²In this note by convention an M-matrix is meant a *non-singular* M-matrix. We will make it clear to distinguish non-singular M-matrix and *singular* M-matrix in the context (A typical example of a singular M-matrix is graph Laplacian matrix).

where $\star$ denotes a ‘don’t-care’ term. Furthermore, there exists a real non-singular matrix P_2 such that A is real similar to a strictly diagonal dominant matrix $B_2 = P_2 A P_2^{-1}$:

$$B_2 = \begin{bmatrix} \lambda_1 & \beta \\ 0 & \lambda_2 \end{bmatrix} \quad (2)$$

where $|\lambda_1| > |\beta|$.

Proof. The proof of the first statement follows similarly the standard procedure of real Jordan decomposition (e.g., (Horn and Johnson, 2012)) and is omitted here. We proceed with the proof of the second statement, which is equivalent to proving that ‘ B_1 is real similar to B_2 ’. Consider a real diagonal matrix $\bar{P} = \text{diag}\{k_1, k_2\}$ with $k_1 > 0, k_2 > 0$. Then

$$\bar{P} B_1 \bar{P}^{-1} = \begin{bmatrix} \lambda_1 & (k_1 \star)/k_2 \\ 0 & \lambda_2 \end{bmatrix} \quad (3)$$

Since both λ and the ‘don’t-care’ term $\star$ are bounded, there exist k_1 and k_2 such that $|\lambda_1| > |(k_1 \star)/k_2| := \beta$. By letting $P_2 := \bar{P} P$, we complete the proof. $\square$

Note the proof of the above lemma does not require the condition that A is Hurwitz. The statements still hold with the condition that the matrix A has two real and **non-zero** eigenvalues. In fact, we conclude:

- Every real 2×2 matrix with non-zero real eigenvalues is real similar to a strictly diagonal dominant real matrix.

3.2. Case II: real matrix with complex eigenvalues

We then consider conditions for real matrix with complex eigenvalues. To begin, the Hurwitz constraint will not be imposed.

Lemma 2. Consider the following 2×2 real matrices

$$J = \begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix} \quad \beta \neq 0 \quad (4)$$

and

$$K = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \quad (5)$$

Then J and K are similar if and only if there exist real x and $y \neq 0$ such that

$$\begin{aligned} a &= \alpha - x \\ d &= \alpha + x \\ b &= y^{-1} \sqrt{\beta^2 + x^2} \\ c &= -y \sqrt{\beta^2 + x^2} \end{aligned} \quad (6)$$

Proof. Suppose first that J and $x, y \neq 0$ are given and the quantities a, b, c, d are then defined as above. It is easily seen that $a + d = 2\alpha$ and $ad - bc = \alpha^2 + \beta^2$, ensuring that J and K have the same trace and determinant. Hence they are similar.

Conversely, suppose J and K are given and known to be similar. Since the traces must be equal, there holds $a + d = 2\alpha$, and then there must be an x satisfying the first two equations of (6); we can define it as $\alpha - a$ and there follows $x = d - \alpha$.

We next argue that b and c must be nonzero. For a contradiction, we suppose to the contrary. Then the determinant of K would be $\alpha^2 - x^2$ while the determinant of J is $\alpha^2 + \beta^2$. Equality of the two determinants would imply that $x = 0, \beta = 0$, which contradicts the theorem hypothesis that $\beta \neq 0$.

Next, define y by $y = \sqrt{\beta^2 + x^2}/b$, and observe that $y \neq 0$. Using again the determinantal equality $\alpha^2 + \beta^2 = ad - bc$, and the expressions for a, d and b in (6), the equality for c in (6) is immediate. $\square$

Given a matrix J as in (4) with $|\alpha| < |\beta|$, we now show that there is no choice of x, y for which we can even get non-strict diagonal dominance of K .

Lemma 3. *Consider the matrix J of (4) with $|\alpha| < |\beta|$. Then there is no choice of $x, y \neq 0$ such that the matrix $L := K$ as below*

$$L = \begin{bmatrix} \alpha - x & y^{-1}\sqrt{\beta^2 + x^2} \\ -y\sqrt{\beta^2 + x^2} & \alpha + x \end{bmatrix} \quad (7)$$

is non-strictly diagonal dominant, and therefore no non-singular similarity transformation of J for which the transformed matrix is non-strictly diagonal dominant.

Proof. Suppose, to obtain a contradiction, that $|\alpha| < |\beta|$ and there is a choice of $x, y \neq 0$ ensuring that L is non-strictly diagonal dominant. Then there holds

$$\begin{aligned} |\alpha - x| &\geq |y^{-1}|\sqrt{\beta^2 + x^2} \\ |\alpha + x| &\geq |y|\sqrt{\beta^2 + x^2} \end{aligned} \quad (8)$$

Multiplying the two inequalities together gives

$$|\alpha^2 - x^2| = |\alpha - x||\alpha + x| \geq \beta^2 + x^2 \quad (9)$$

Since $\beta^2 > \alpha^2$, there exists a nonzero γ^2 such that $\beta^2 = \alpha^2 + \gamma^2$, and then the above inequality becomes

$$|\alpha^2 - x^2| \geq \alpha^2 + \gamma^2 + x^2 \quad (10)$$

However, this inequality can never be satisfied: the inequality (10) implies $\alpha^2 + x^2 \geq |\alpha^2 - x^2| \geq \alpha^2 + \gamma^2 + x^2$ which is impossible with $\gamma^2 > 0$. Therefore, there does not exist a non-singular real transformation of J for which the transformed matrix is non-strictly diagonal dominant. This completes the proof. $\square$

We now summarize the above results and present the following main conditions.

Theorem 1. Consider a 2×2 real Hurwitz matrix A and suppose its two eigenvalues are complex, denoted by $\lambda_{1,2} = \alpha \pm \beta j$ with $\alpha < 0, \beta \neq 0$. Then there exists a real non-singular matrix P_1 such that A is real similar to a real matrix B_1

$$B_1 = \begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix} \quad (11)$$

Furthermore,

- If $|\alpha| > |\beta|$, then there exists a real non-singular matrix P such that A is real similar to a **strictly** diagonal dominant real matrix B .
- If $|\alpha| = |\beta|$, then there exists a real non-singular matrix P such that A is real similar to a **non-strict** diagonal dominant real matrix B .
- If $|\alpha| < |\beta|$, then there **does not** exist a real non-singular matrix P such that A is real similar to a diagonal dominant real matrix B .

Proof. The proof of the first statement follows similarly the standard procedure of real Jordan decomposition (e.g., (Horn and Johnson, 2012)) and is omitted here. We proceed with the proof of the second set of three statements. If $|\alpha| > |\beta|$, then the matrix B_1 in (11) is a **strictly** diagonal dominant real matrix. If $|\alpha| = |\beta|$, then the matrix B_1 in (11) is a **non-strict** diagonal dominant real matrix. By letting $B_1 := B$ we conclude the statements of these two cases. For the third case that $|\alpha| < |\beta|$, Lemma 3 states that there is no non-singular real transformation matrix P such that A is real similar to a diagonal dominant real matrix B . $\square$

Note the above lemmas and Theorem 1 do not require the condition that A is Hurwitz. The statements on real similarity in Theorem 1 still hold under the condition that the complex eigenvalues of the matrix A satisfy $|\alpha| > |\beta|$. In particular, we conclude:

- Every real 2×2 matrix with complex eigenvalues $\lambda_{1,2} = \alpha \pm \beta j$ and with $|\alpha| > |\beta|$ is real similar to a strictly diagonal dominant real matrix.

As a side note, we also consider the problem of complex similarity of a 2×2 real matrix to a complex diagonal dominant matrix.

Lemma 4. Every 2×2 real Hurwitz matrix is complex similar to a strictly diagonal dominant complex matrix; i.e., for any 2×2 real Hurwitz matrix A , there exists a **complex** non-singular matrix, such that the real matrix A is similar to a (possibly complex) strictly diagonal dominant matrix.

Proof. For the case that the Hurwitz matrix A has two real eigenvalues, the proof follows similarly to that of Lemma 1 while we can take the real matrix P as the transformation matrix. For the case that the Hurwitz matrix A has complex eigenvalues, it is real similar to the real matrix B_1 in (11). Taking a non-singular complex matrix

$$P_3 = \begin{bmatrix} -j & -j \\ 1 & -1 \end{bmatrix} \quad (12)$$

we can obtain

$$P_3^{-1}B_1P_3 = \begin{bmatrix} \alpha + bj & 0 \\ 0 & \alpha - bj \end{bmatrix} \quad (13)$$

which is complex and diagonal. This completes the proof. $\square$

Note: the statement in this lemma (for the complex eigenvalue case) is a special case of the general result: a matrix that has distinct eigenvalues is diagonalizable.

Remark 1. *The above results treat similarity to a row-diagonal dominant matrix; while evidently, all main statements can be extended to a column-diagonal dominant matrix. This is due to the following well-known result: a real square matrix A is real similar to A^T .*

4. Intuitive interpretation by Gershgorin's circles

The Gershgorin circle theorem (Varga, 2010) gives a convenient method for characterizing locations of eigenvalues for a (real or complex) matrix: all eigenvalues of a matrix are located in the union of all Gershgorin circles, while each circle is centered at each diagonal entry with radius defined by the sum of the magnitude of all off-diagonal entries (in the corresponding row or column). We now provide a Gershgorin circle interpretation to the main results of Lemmas 1-4.

For Lemma 1 when the Hurwitz matrix A has real non-zero eigenvalues: the real matrix P serves as a real scaling matrix such that the transformed real matrix $B_2 = P_2^{-1}AP_2$ has two scaled Gershgorin circles; each circle is centered at one real eigenvalue and does not touch upon the origin point. In fact, the radius of each circle can be made arbitrarily small by a carefully chosen transformation matrix P , and the diagonal dominant property of B_2 is ensured.

For Theorem 1 when the matrix A has complex eigenvalues and is real similar to the matrix the matrix B_1 in (11): the real similarity transformation via P results in another real matrix (in particular, the diagonal entries are real), and the centers of all Gershgorin circles are restricted to lying on the real line. For the matrix B_1 in (11) and its real similarity classes, the **smallest** Gershgorin circle with a center on the real line that includes both complex eigenvalues is the circle centered at α with a radius $|\beta|$. For $|\beta| > |\alpha|$, such a Gershgorin circle encircles the origin and always touches upon the complex line. There is no hope of finding a real transformation matrix P_2 to further scale the circles so that they do not encircle the origin, which implies that A cannot be real similar to a diagonal dominant matrix.

For Lemma 4 with complex matrix similarity: when the transformation matrix P is allowed to be a complex matrix, the transformed matrix can also be complex (in particular, the diagonal entries can be complex). The Gershgorin circles are not necessarily centered on the real line but can be relocated in the complex plane by the complex matrix P . In this case, when the eigenvalues have a non-zero real part, each Gershgorin circle can be shifted and centered at an eigenvalue, and its size can be rescaled with an arbitrarily small radius so that it does not encircle the origin. Thus, in this case, any real matrix with each eigenvalue of non-zero real part is complex similar to a (possibly complex) diagonal dominant matrix.

5. Real similarity with general matrix

We are now ready to show the following general results. We note the following theorem does not require the Hurwitz condition.

Theorem 2. *Consider a real square non-singular matrix $A \in \mathbb{R}^{n \times n}$. Then the following statements hold.*

- *Suppose that all eigenvalues of A are real. Then A is real similar to a strictly diagonal dominant real matrix.*
- *Suppose that some eigenvalues of A are complex and for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ there holds $|\alpha_i| > |\beta_i|, \forall i$. Then A is real similar to a strictly diagonal dominant real matrix.*
- *Suppose that some eigenvalues of A are complex and for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ there holds $|\alpha_i| \geq |\beta_i|, \forall i$. Furthermore, for any complex eigenvalue pair $\lambda_i = \alpha_i \pm \beta_i j$ with $|\alpha_i| = |\beta_i|$, suppose its geometric multiplicity equals its algebraic multiplicity. Then A is real similar to a non-strict diagonal dominant real matrix.*
- *For all other cases not covered by the above conditions, there **does not** exist a real non-singular matrix P such that A is real similar to a diagonal dominant real matrix.*

Proof. First, we recall the fact that any real square matrix $A \in \mathbb{R}^{n \times n}$ is real similar to the following real Jordan normal form (cf. (Horn and Johnson, 2012, Chapter 3))

$$J = PAP^{-1} = \text{blk-diag}\{J_1, J_2, \dots, J_i, \dots, J_k\} \quad (14)$$

where P is a real non-singular matrix, J_i is the i -th **real** Jordan block associated with the eigenvalues of A .

- If all eigenvalues of A are real, then the diagonal entries of J are the set of eigenvalues of A (counting multiplicities). Following a similar proof to that of Lemma 1, one can find a real non-singular diagonal matrix $\bar{P}$ such that $\bar{P}J\bar{P}^{-1}$ is strictly diagonal dominant.
- If some eigenvalues of A are complex, denoted by $\lambda_i = \alpha_i \pm \beta_i j$, then the 2×2 block matrix

$$B_i = \begin{bmatrix} \alpha_i & \beta_i \\ -\beta_i & \alpha_i \end{bmatrix} \quad (15)$$

appears in the i -th real Jordan block that takes the following form

$$J_i = \begin{bmatrix} B_i & I_2 & & \\ & B_i & \ddots & \\ & & \ddots & I_2 \\ & & & B_i \end{bmatrix} \quad (16)$$

We now consider the following cases that for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ there hold $|\alpha_i| \geq |\beta_i|, \forall i$.

- Case I: for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ there hold $|\alpha_i| > |\beta_i|, \forall i$. Then following a similar proof to that of Lemma 1, one can find a real non-singular matrix $\bar{P}_i$ such that $\bar{P}_i J_i \bar{P}_i^{-1}$ is strictly diagonal dominant, which in turn ensures that A is real similar to a strictly diagonal dominant real matrix.
- Case II: for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ with $|\alpha_i| = |\beta_i|$, suppose the geometric multiplicity equals the algebraic multiplicity. Then the real Jordan block J_i for such an eigenvalue pair λ_i with $|\alpha_i| = |\beta_i|$ degenerates to $J_i = B_i$. According to Theorem 1, this ensures that A is real similar to a non-strict diagonal dominant real matrix.
- Case III: for all complex eigenvalue pairs $\lambda_i = \alpha_i \pm \beta_i j$ with $|\alpha_i| = |\beta_i|$, suppose for at least one such pair, the algebraic multiplicity is greater than the geometric multiplicity. Then the real Jordan block J_i in (16) for such an eigenvalue pair λ_i has at least one off-diagonal identity block I_2 . As a consequence, there **does not** exist a real non-singular matrix $\bar{P}_i$ such that $\bar{P}_i J_i \bar{P}_i^{-1}$ is diagonal dominant, and A is not real similar to a diagonal dominant matrix.
- For all other cases with complex eigenvalue pairs (i.e., there exist at least one complex eigenvalue with $|\alpha_i| < |\beta_i|$), according to Theorem 1, the associated real Jordan block J_i cannot be made to be real similar to a diagonal dominant matrix. As a consequence, there **does not** exist a real non-singular matrix P such that A is real similar to a diagonal dominant real matrix.

□

To complete the story, we further discuss complex similarity of a general real square matrix to a diagonal dominant matrix.

Theorem 3. *Every real Hurwitz matrix is complex similar to a (possibly complex) strictly diagonal dominant matrix.*

A more general result on complex similarity is shown below.

Theorem 4. *Every real non-singular matrix is complex similar to a (possibly complex) strictly diagonal dominant matrix.*

The proofs follow similarly the proof of the 2×2 matrix case and the real similarity in Theorem 2, which are omitted here.

6. Special cases

Some special cases are recalled here for further references (Horn and Johnson, 2012; Berman and Plemmons, 1994).

Lemma 5. *Consider a real Hurwitz matrix A and suppose its off-diagonal entries are non-negative (i.e., $-A$ is an M -matrix). Then there exists a real **diagonal** matrix K such that KAK^{-1} is strictly diagonal dominant with negative diagonal entries.*

Lemma 6. *Consider a real Hurwitz matrix and suppose it is also an H-matrix. Then there exists a real **diagonal** matrix K such that $KA K^{-1}$ is strictly diagonal dominant with negative diagonal entries.*

References

- Berman, A., Plemmons, R.J., 1994. Nonnegative matrices in the mathematical sciences. SIAM.
- Horn, R.A., Johnson, C.R., 2012. Matrix analysis. Cambridge University Press.
- Varga, R.S., 2010. Geršgorin and his circles. volume 36. Springer Science & Business Media.