ON DETECTING THE TRIVIAL RATIONAL 3-TANGLE

BO-HYUN KWON

ABSTRACT. An important issue in classifying the rational 3-tangle is how to know whether or not the given tangle is the trivial rational 3-tangle called ∞ -tangle. The author[2] provided a certain algorithm to detect the ∞ -tangle. In this paper, we give a much simpler method to detect the ∞ -tangle by using the *bridge arc replacement*. We hope that this method can help prove many application problems such as a classification of 3-bridge knots.

1. INTRODUCTION

A classification of rational 2-tangle is provided by Conway [1] in 1970. It is a perfect classification of them since he had shown that each rational 2-tangle is presented by the corresponding rational number. The author [2] gave an algorithm to classify the rational 3-tangles. Unfortunately, the algorithm is not enough to classify the rational 3-tangle completely since it is only used to distinguish given two rational 3-tangles not assigning them to a certain natural group. The main part of the algorithm is to show whether or not the modified rational 3-tangle from the given two rational 3-tangles is the trivial rational 3-tangle called the ∞ -tangle. There is a basic way to detect ∞ -tangle by the fundamental group argument, but the algorithm is much better than the traditional way. In this paper, we provide a new method to detect ∞ -tangle.

2. The trivial rational 3-tangle, ϵ



FIGURE 1. Three compressing disks in ∞ -tangle, (B, ϵ)

Consider an ordered pair $T = (B, \epsilon = \epsilon_1 \cup \epsilon_2 \cup \epsilon_3)$ as in Figure 1, where B is the unit 3-ball. This trivial rational 3-tangle in B is called an ∞ tangle or ϵ . Let E_1, E_2 and E_3 be

Date: Monday, March 13, 2023.

BO-HYUN KWON

the three pairwise disjoint, non-parallel compressing disks in $B - \epsilon$ as in Figure 1. Then E_1, E_2 and E_3 separate B into four components. Let H_i be the component which contains ϵ_i and $H = cl(B - (H_1 \cup H_2 \cup H_3))$, where cl(*) means that the closure of *. Let E'_i be the 2-punctured disk in $\Sigma_{0,6}$ so that $\partial E'_i = \partial E_i$ and $E_i \cup E'_i$ bounds the ball H_i in B. Let $E = E_1 \cup E_2 \cup E_3$, $E' = E'_1 \cup E'_2 \cup E'_3$ and $\partial E = \partial E' = \partial E_1 \cup \partial E_2 \cup \partial E_3$. Let $\Sigma_{0,6} = \partial B - \epsilon$ and $P = cl(\Sigma_{0,6} - (E'))$ which is a pair of pants.

3. BRIDGE ARC REPLACEMENT

Let $\tau = \tau_1 \cup \tau_2 \cup \tau_3$ be a rational 3-tangle in B, where τ_i be a properly embedded arc in B. A disk D in B is called a *bridge disk* if $D^\circ \cap \tau = \emptyset$ and $\partial D = \tau_i \cup \beta$ and $\tau_i \cap \beta = \partial \tau_i$ for some $i \in \{1, 2, 3\}$, where β is a simple arc in S^2 so that $\beta^\circ \cap \tau = \emptyset$. The simple arc β in $\Sigma_{0,6}$ is called a *bridge arc* if it cobounds a bridge disk with a component of τ . We note that there is a collection of disjoint three bridge arcs for a rational 3-tangle. However, we also note that the collection is not unique (up to isotopy) for the same rational 3-tangle. Actually, there are infinitely many different bridge arcs for the same string of the rational 3-tangle. We say that the simple closed curve γ is *obtained from* the bridge arc β if the boundary of a small regular neighborhood of β in $\Sigma_{0,6}$ is isotopic to γ .

Lemma 1. Let β_1 and β_2 be two disjoint bridge arcs on $\Sigma_{0,6}$ of (B, τ) , where τ is a rational 3-tangle in B. Let β_3 be an arc connecting the two remaining punctures on $\Sigma_{0,6}$ which is disjoint with $\beta_1 \cup \beta_2$. Then γ_3 is also a bridge arc on $\Sigma_{0,6}$ of (B, τ) .

Proof. We first claim that there are two disjoint bridge disks so that each of them contains the bridge arcs β_1 and β_2 respectively. We take a bridge disk bounded by the union of β_1 and a string of τ . Then cut *B* along the bridge disk and open it along the two cut disks to have a rational 2-tangle. Since β_2 is disjoint with β_1 , we can take a bridge disk bounded by β_2 and a string of the obtained rational 2-tangle. We glue the two cut disks back to have the original rational 3-tangle. This implies the existence of two disjoint bridge disks satisfying the condition. Moreover, from the rational 2-tangle, if we cut the tangle along the bridge disk bounded by β_2 and a string then we have a rational 1-tangle. It is clear that any arc connecting the two endpoint of the 1-tangle on $\Sigma_{0,2}$ without intersecting the opened two disks along the bridge disks cobounds a bridge disk in the 3-ball containing the rational 1-tangle. By gluing the two pairs of bridge disks back, we can guarantee the lemma above.

Let $\{\beta_1, \beta_2, \beta_3\}$ be a collection of disjoint three bridge arcs on $\Sigma_{0,6}$ for $T = (B, \tau)$. Now, we consider the subarcs of $\beta := \beta_1 \cup \beta_2 \cup \beta_3$ in P. We assume that there is no bigon so that one of the sides is a subarc of β in P. If there is a such bigon then we can isotope the subarc to vanish the bigon. We say that β is *dense* in P if none of two adjacent endpoints of the component of β in ∂P belongs to the same β_i for i = 1, 2, 3. We assume that there are at least two successive adjacent endpoints of the components of β in ∂P contained in the same β_i to not satisfy the definition of dense as in Figure 2. We note that two of the successive endpoints connect the puncture directly without passing through the other successive endpoints. Let c_1 and c_2 be the two subarc of β_i explained above. We define the two directions (+) and (-) as in the first diagram of Figure 2. If the two punctures appear when the two subarcs go to the opposite direction we take the new arc instead of β_i as in the second diagram of Figure 2. By Lemma 1, the new arc is also a bridge arc. Moreover, it also clear that the newly selected bridge arc is disjoint with $\beta - \beta_i$. Similarly, if the two punctures appear when they go to the same direction as in the third diagram, we take the new bridge arc which is disjoint with the remaining two bridge arcs as in the fourth diagram of Figure 2. This procedure is called the *bridge arc replacement*, briefly **BR**.

Lemma 2. Let $\{\beta_1, \beta_2, \beta_3\}$ be a collection of disjoint three bridge arcs on $\Sigma_{0,6}$ of $T = (B, \tau)$. Then there exist a collection of disjoint three bridge arcs $\{\beta'_1, \beta'_2, \beta'_3\}$ on $\Sigma_{0,6}$ of T which is dense in P.



FIGURE 2. Bridge arc replacement, **BR**

Proof. By repeating the procedure **BR**, we would get the collection which is dense in P since **BR** reduces the number of endpoints if the collection is not dense and the number of endpoints are finite.

We claim that the procedure **BR** can detect the ∞ -tangle. The following theorem would cover two of the three possible cases.



FIGURE 3. A collection of three disjoint simple arcs which is dense in P.

Theorem 3. Let $\mathcal{B} = \{\beta_1, \beta_2, \beta_3\}$ be a collection of disjoint three bridge arcs on $\Sigma_{0,6}$ of $T = (B, \epsilon)$ which is dense in P. Then \mathcal{B} is unique up to isotopy. Especially, the three simple closed curves obtained from $\{\beta_1, \beta_2, \beta_3\}$ respectively are $\partial E_1, \partial E_2, \partial E_3$ up to isotopy.

Proof of the first two cases. We note that if a simple closed curve obtained from β_i is isotopic to ∂E_j then β_i is the straight arc connecting the two punctures in E'_j up to isotopy. We first assume that two of the simple closed curves obtained from $\{\beta_1, \beta_2, \beta_3\}$ are isotopic to ∂E_i and ∂E_j for $i \neq j$. Then we have the first diagram in Figure 3. It is clear that the remaining bridge arc is also isotopic to the straight arc connecting the two punctures in E'_k if the collection is dense in P. Secondly, we assume that one of them is isotopic to ∂E_i for some *i*. Then we have the second diagram of Figure 3. We note that it is impossible to have

BO-HYUN KWON

a wave in terms of ∂E if the collection is dense in P in this case. This implies that there is no wave. So, every simple closed curves obtained from $\{\beta_1, \beta_2, \beta_3\}$ are isotopic to one of ∂E_j for j = 1, 2, 3.

For the last case, we assume that none of them is isotopic to ∂E_i for some *i*. In order to prove the last case, we use the arguments when we made the algorithm to classify rational 3-tangles.

3.1. Dehn's parametrization of a simple closed curve in $\Sigma_{0.6}$.



FIGURE 4. Standard arc in P

Let γ be a simple closed curve in $\Sigma_{0,6}$. Assume that γ meets ∂E_i . Then we can isotope γ so that it meets with ∂E_i in a special subarc ω_i of ∂E_i only. We call ω_i the window in ∂E_i . Then γ has parallel arcs which are the same type in P. Let l_{ij} be the arc type of them as in Figure 4 which is called the *standard* arc type. (Refer to [3].) Let x_{ij} be the number of parallel arcs of the type l_{ij} which is called the *weight* of l_{ij} for γ . There is a relation between x_{ij} and the geometric intersection number $|\gamma \cap \omega_s|$ as follows.

Lemma 4 ([2]). Let $I_s = |\gamma \cap \omega_s|$, where $s \in \{1, 2, 3\}$. Then I_s determine x_{ij} . Especially, if $x_{ii} > 0$ for some i, then $x_{ij} = I_j, x_{ik} = I_k$ and $x_{ii} = \frac{I_i - I_j - I_k}{2}$. Also, if $x_{ii} = 0$ for all i, then $x_{ij} = \frac{I_i + I_j - I_k}{2}$ for distinct i, j, k.

We note that γ can be determined by x_{ij} and the patterns in E'_i . So, we can parametrize γ in $\Sigma_{0,6}$ as follows.



FIGURE 5. parameterization in two punctured disk

Theorem 5 (Special case of Dehn's Theorem [3]). There is a parametrization of closed essential 1-manifolds in $\Sigma_{0,6}$ by ordered sequences of nine parameters $(p_1, q_1, t_1, p_2, q_2, t_2, p_3, q_3, t_3)$. (where $p_i, q_i \in \mathbb{Z}^+ \cup \{0\}$ and $t_i \in \mathbb{Z}$ for i = 1, 2, 3.), where $2p_i = I_i$ and $2p_i = J_i$.

In the diagrams in Figure 5, E''_i is the inner two punctured disk in E'_i to have only parallel arcs. Then q_i and t_i determine all the arc types and weights of them in E'_i by determining the connecting pattern between b_i and s_j .

We are dealing with the simple arc instead of the simple closed curve for proving the main theorem. We note that the argument in this section is in terms of simple closed curves not a simple arc. The simple closed curves obtained from the simple arcs can be parameterized by the argument in this section. Moreover, a simple arc β_i is a bridge arc if and only if γ_i obtained from β_i bounds a compressing disk in $B - \epsilon$. So, we can discuss the all argument in terms of simple arcs instead of simple closed curves if we consider the exceptional arc types having a puncture as one of the endpoints of them.

4. Standard diagram and the proof of the last case

Recall that $\{\beta_1, \beta_2, \beta_3\}$ is a collection of disjoint bridge arcs on $\Sigma_{0,6}$ for $T = (B, \epsilon)$. In order to analyse β_i , we set $\Sigma_{0,6}$ with the three disjoint non-parallel simple closed curves $\partial E_1, \partial E_2$ and ∂E_3 . Let $P' = \partial B - (\bigcup_{i=1}^2 E''_i)$. Now, take equators e_i for each E''_i as in Figure 6. Then e_i divides E''_i into E''_i and E''_i as in Figure 6. We note that ∂E_i is isotopic to $\partial E''_i$ for i = 1, 2, 3. Now, we introduce a lemma which helps us to realize β in a certain formed diagram called *standard diagram*.

Lemma 6 (Lemma 6.1 in [2]). Let N be a compressing disk in $B - \epsilon$ and h be the homeomorphism obtained from the clockwise half Dehn twist supported on N' which is the 2-punctured disk in $\Sigma_{0,6}$ so that $\partial N = \partial N'$. Then a simple closed curve δ bounds a compressing disk in $B - \epsilon$ if and only if $h(\delta)$ bounds a compressing disk in $B^3 - \epsilon$.

We note that Lemma 6 implies that a simple arc β_i is a bridge arc in $B - \epsilon$ if and only if $h(\beta_i)$ is a bridge arc in $B - \epsilon$.



FIGURE 6. Standard diagram

Let γ_i be the simple closed curve obtained from β_i which can be parametrized by an ordered sequence of integers $(p_1, q_1, t_1, p_2, q_2, t_2, p_3, q_3, t_3)$. We now investigate the simple closed curve γ'_i (it may be in a different isotopy class in $\Sigma_{0,6}$) so that every component of $\gamma'_i \cap P'$ is isotopic to one of the given real arc types in P' of one of the two diagrams as in Figure 6 by taking a properly chosen t'_i instead of t_i by the following lemma.

Lemma 7 (Lemma 8.2 in [2]). Suppose that γ is parametrized by $(p_1, q_1, t_1, p_2, q_2, t_2, p_3, q_3, t_3)$ and it has l_{11} arc type in P. Then there exist simple closed curves γ' which is parameterized by $(p_1, q_1, t'_1, p_2, q_2, t'_2, p_3, q_3, t'_3)$ for some $t'_i \in \mathbb{Z}$ so that all components of $\gamma' \cap P'$ are isotopic to one of the arcs in the diagram.

We say that a simple closed curve γ is in standard position if all componet of $\gamma \cap P'$ are isotopic to one of the arcs in the diagram of Figure 6. The diagram in Figure 6 is called standard diagram. Let β'_i be the simple arc so that γ'_i is the simple closed curve obtained from β'_i . We note that the components of $\beta'_i \cap P'$ are also carried by the arc types in the standard diagram. In order to see this, it is enough to investigate the component in P' which connects the puncture in E''_i directly as in Figure 7. We note that γ has the two components (a) and (b) and β has the component (b') which is parallel to the component (b). The original lemma is for a simple closed curve not a multiple of simple closed curves. However, there is no difference between a simple closed curve and multiple simple closed curves when we apply the above lemma. It works for $\beta' = \beta'_1 \cup \beta'_2 \cup \beta'_3$ as well. We note that the changes of the twisting number t_i would preserve the density since the half Dehn twist along E''_i just rotate the position of the all endpoints on $\partial E''_i$ by 180°. We also note that the trivial rational 3-tangle ϵ in B is preserved when we apply the half Dehn twists supported on N' defined in Lemma 6. For additional modifications of β , we consider the following two lemmas and the theorem.

Lemma 8 (Lemma 9.1 in [2]). If A is a compressing disk in $B^3 - \epsilon$, then $(\delta_1 \delta_2^{-1})^{\pm 1}(\partial A)$ also bounds a compressing disk in $B^3 - \epsilon$. (where δ_1 and δ_2 are the clockwise half Dehn twists supported on two punctured disks C_1 and C_2 respectively as in Figure 8.)







FIGURE 8. Half Dehn twists to preserve the boundness of a compressing disk in $B^3 - \epsilon$

Lemma 9 (Lemma 9.2 in [2]). Suppose that A is a compressing disk in $B^3 - \epsilon$. Let δ_3 be the counter clockwise half Dehn twist supported on the 2-punctured disk E'_4 as in Figure 9. Then $\delta_3(\partial A)$ also bounds a compressing disk $\delta_3(A)$ in $B^3 - \epsilon$.

Theorem 10 (Theorem 9.3 in [2]). Suppose that A is a compressing disk in $B^3 - \epsilon$ and ∂A is in standard position in P'. Then one of the half Dehn twists $(\delta_1 \delta_2^{-1})^{\pm 1}$ and $\delta_3^{\pm 1}$ reduces the minimal intersection number of ∂A with ∂E .

The following lemma is the main idea to prove the last case of the main theorem.



FIGURE 9. A half Dehn twists to preserve the boundedness of a compressing disk in $B^3-\epsilon$

Lemma 11. Let $\{\beta_1, \beta_2, \beta_3\}$ be the collection of disjoint simple arcs which are bridge arcs in $B - \epsilon$. We assume that $\beta = \beta_1 \cup \beta_2 \cup \beta_3$ is in standard position and all of them have a wave in terms of $\partial E''$. Then, there exist two adjacent endpoints of $\beta \cap P'$ in $\partial E''_i$ for some *i* which belong to the same β_j for j = 1, 2, 3.



FIGURE 10.



FIGURE 11.

Proof. By the assumption that all of β_i have a wave, without loss of generality, we assume that β has subarcs of type 1 or type 3 as in the first diagram of Figure 10 and Figure 12. By applying the homeomorphism δ_3 in Lemma 9 to β , we can remove the subarcs of type 2 as in the second diagram of Figure 10 and Figure 12 respectively. Let m_k be the weights of each type k. In the second diagram of Figure 10, we note that $\mathbf{A} = m_2, \mathbf{B} = m_3 - m_2$ and $\mathbf{C} = m_{10} + m_{11}$. In the right diagram of Figure 12, we have $\mathbf{A} = m_2$ and $\mathbf{B} = m_3 - m_2$. Also, in the bottom diagram of Figure 12, we have $\mathbf{A} = m_2 - m_3, \mathbf{B} = m_3$ and $\mathbf{C} = m_2, \mathbf{D} = m_1 - m_2, \mathbf{E} = m_{72} + m_3 - m_2$, where 7₂ is lower track of the type 7's in the standard diagram. For the detail of this weight changes in Figure 10 and Figure 12, refer to Theorem 9.3 in [2]. We note that the homeomorphism in each step reduces the total weight of the subarcs in P' by Theorem 10.

We claim that the collection of the new bridge arcs is dense in P'. In order to show the claim for the case described in Figure 10, consider the diagrams in Figure 11. It is enough to investigate the pattern of endpoints in $\partial E''_1$ since the endpoints in $\partial E''_2$ and $\partial E''_3$ are fixed. Since β is dense in P', the depicted two adjacent arcs in the three positions (1), (2) and (3) should be a part of distinct bridge arcs. This implies that the right diagram does not violate the rule to satisfy the density as well. It is not difficult to see that the other adjacent endpoints in $\partial E''_1$ except the indicated adjacent endpoints by the parallel arcs also keep the rule for the density since the patterns are the same with the previous corresponding patterns.



FIGURE 12.

Similarly, we note that the collections of the modified bridge arcs described in Figure 12 are dense by Figure 13. Finally, we would get the four diagrams as in Figure 14 which do not have the type 2. First, we assume that both of m_1 and m_3 are positive as in the diagram





(A). Then, there is two adjacent endpoints in $\partial E_1''$ since the two punctures in E_1'' should be the two endpoints of the same β_i for some i = 1, 2, 3.



FIGURE 14.

For the rest of the cases, we apply a multiple of the homeomorphism $(\delta_1^{-1}\delta_2)^{\pm 1}$ to have a simpler diagram as follows. For the diagram (B), we apply a multiple of the homeomorphism $(\delta_1^{-1}\delta_2)$ to have the rightmost diagram in Figure 11. We note that the homeomorphism $(\delta_1^{-1}\delta_2)$ preserves the density.



FIGURE 15.





Now, consider the cases (C) and (D). Since the cases are symmetric, it is enough to figure out the case (D). We apply the homeomorphism $(\delta_1^{-1}\delta_2)$ to the diagram (D) until we get the rightmost diagrams in Figure 17. We note that the homeomorphism $(\delta_1^{-1}\delta_2)$ preserves



FIGURE 17.





the density as in Figure 18. If we have the diagram having the rightmost two diagrams, the diagram has the type 2. We repeat the procedure to remove the type 2. We note that the procedure preserves the density. Since the total weights of the subarcs in P' is finite and the procedures mentioned in this proof reduce the total weights, we eventually get a collection of bridge arcs so that at least one of them is isotopic to the straight arc connecting the two punctures in E''_i for some $i \in \{1, 2, 3\}$. So, it is enough to consider the diagrams right before the final diagram. Then we have the two diagrams (a) and (b) as in Figure 19 if we ignore symmetric cases. We note that if we apply $(\delta_1^{-1}\delta_2)$ to the diagram (a) then we have the bridge arc which connects the two punctures in E''_3 . Also, we apply δ_3^{-1} to the diagram (b)

then we have the bridge arc which connects the two punctures in E''_1 . In (a), assume that β_1 is isotopic to the red arc connecting the two punctures in E''_1 . Since β_2 and β_3 also have a wave, it is clear that there are two adjacent endpoints in $\partial E''_2$ which belong to the same β_j for some j = 2, 3. Similarly, in (b), there are two adjacent endpoints in $\partial E''_2$ which belong to the same β_j for some j = 2, 3. This implies that either $\delta_3^{\pm 1}$ or $(\delta_1^{-1}\delta_2)^{\pm 1}$ cannot preserve the density. This violates the argument above.



FIGURE 19.

Now, we can complete the proof of Theorem 3.

Proof of the last case. By Lemma 11, it is not possible to have a collection of disjoint three bridge arcs so that at least one of them has a wave. Therefore, if β is dense in P' then none of β_i has a wave. This implies that the collection of the bridge arcs should be isotopic to the collection of ∂E_1 , ∂E_2 and ∂E_3 .

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the the Ministry of Education (NRF-2020R1I1A1A01052279 and NRF-2020R1A2C1A01009184)

References

- [1] J. Conway, An enumeration of knots and links, and some of their algebraic properties, in Computational Problems in Abstract Algebra (Pergamon, Oxford, Press, 1970), pp. 329–358.
- [2] B. Kwon, An algorithm to classify rational 3-tangles, J. Knot Theory Ramifications 24 (2015), no. 1, 1550004 (62 pages).
- [3] R. C. Penner; J. L. Harer Combinatorics of Train Tracks, Annals of Mathematics Studies No. 125, Princeton University Press, Princeton, NJ, 1992. xii+216 pp.