

A stable supermassive charged gravitino?

Krzysztof A. Meissner¹ and Hermann Nicolai²

¹*Faculty of Physics, University of Warsaw
Pasteura 5, 02-093 Warsaw, Poland*

²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut)
Mühlenberg 1, D-14476 Potsdam, Germany*

Some time ago it was suggested that dark matter may consist in part of an extremely dilute gas of supermassive gravitinos with fractional charge $2e/3$ [1]. This scheme makes the definite (and falsifiable) prediction that massive gravitinos are the *only* new fermionic degrees of freedom beyond the known three generations of quarks and leptons of the Standard Model of Particle Physics. In this note we re-examine one special outlier event reported and subsequently discarded by the MACRO collaboration [2] in the light of this proposal and point out the possibility of an alternative interpretation of this event supporting the above hypothesis, whose confirmation (or refutation) would, however, require an independent dedicated experimental effort.

I. INTRODUCTION.

The nature of dark matter (DM) continues to be one of the most vexing questions of modern physics. While current DM scenarios are usually based on the assumption of ultralight constituents (such as axions) or TeV scale WIMPs, the possibility has been raised in recent work [1] that DM could consist at least in part of an extremely dilute gas of supermassive stable gravitinos with charge $q = \pm \frac{2}{3}$ in units of the elementary charge e . This proposal has its roots in Gell-Mann's old observation that the fermion content of the Standard Model of Particle Physics (SM) with three, and only three, generations of quarks and leptons (including right-chiral neutrinos) can be matched with the spin- $\frac{1}{2}$ content of the maximal $N=8$ supermultiplet after removal of eight Goldstinos [3, 4]. The only extra fermions beyond the known three generations of SM fermions would thus be eight massive gravitinos, but nothing else. Crucially, the matching of $U(1)_{em}$ charges requires a 'spurion shift' of $\delta q = \pm \frac{1}{6}$ [3] that is not part of $N=8$ supergravity and that, in terms of the original spin- $\frac{1}{2}$ fermions of $N=8$ supergravity, takes the very special form given in [5, 6]. The main new step taken in [1, 6] consisted in extending these considerations to the eight massive gravitinos which split as

$$\left(\mathbf{3}, \frac{1}{3}\right) \oplus \left(\bar{\mathbf{3}}, -\frac{1}{3}\right) \oplus \left(\mathbf{1}, \frac{2}{3}\right) \oplus \left(\mathbf{1}, -\frac{2}{3}\right) \quad (1)$$

under $SU(3) \times U(1)_{em}$. All gravitinos would thus carry *fractional electric charges*. If one identifies the $SU(3)$ in (1) with $SU(3)_c$ as in [1], a complex triplet of gravitinos would be subject to strong interactions. Importantly, the $U(1)_{em}$ charge assignments for the gravitinos include the spurion shift needed for matching the spin- $\frac{1}{2}$ sectors [1]. The explanation of how to extend the spurion shift to the gravitinos requires a detour via $D=11$ supergravity and the enlargement of the $SU(8)$ R-symmetry of $N=8$ supergravity to $K(E_{10})$ [7]. Although the combined spin- $\frac{1}{2}$ and spin- $\frac{3}{2}$ content would thus coincide with the fermionic part of the $N=8$ supergravity multiplet, the underlying theory would need to be a very specific, but as yet unknown, extension of (gauged) $N=8$ supergravity, which in its original form [8, 9] cannot be correct for reasons that have been known for more than 40 years. Such a theory would almost certainly require a framework beyond space-time based quantum field theory, but most likely different from conventional string theory.

In this note we would like to follow up on this line of thought and further explore options towards linking $N=8$ supergravity to real physics, beyond matching its spin- $\frac{1}{2}$ fermion content with that of the SM. More specifically, we would like to search for possible manifestations and experimental signatures in the spin- $\frac{3}{2}$ sector which could support the above scenario. In comparison with currently popular proposals for extending the SM, which usually come with a lot of extra new particles (especially in the context of low energy supersymmetry), the present scheme is thus very minimalistic. Nevertheless we believe that such a novel approach, also to the DM problem, is amply justified in view of strong indications that the SM could simply survive more or less *as is* all the way up to the Planck scale, and in view of decades of failed attempts to discover any signs of new physics beyond the SM with more conventional approaches. As we will argue here, the MACRO experiment [10, 11], which was originally set up for a very different purpose, may offer some interesting new perspectives in this direction that merit further investigation.

II. SUPERHEAVY GRAVITINOS AS DARK MATTER CANDIDATES

An important consequence of (1) is that, due to their fractional charges the gravitinos cannot decay into SM fermions, and are therefore stable independently of their mass. Their stability against decays makes them natural candidates for DM [1] (let us note that exotic fractionally charged stable states can also appear in certain supersymmetric GUT-type string compactifications [12]). While a very large mass is strongly suggested by the absence of low energy supersymmetry at LHC, an equally compelling argument for large mass are the bounds on the charge q of any putative DM particle of mass m_X derived in [13–15]. These bounds were derived with a perspective different from the present one, namely bounding the charge of ‘low mass’ (in comparison with the Planck scale) DM candidates, by considering various astrophysical constraints (structure formation and CMB, relic density, DM halo constraints, *etc.*); for instance, [13] considers only DM candidates of mass $m_X \lesssim 10^4$ GeV, for which $q < 10^{-4}$. Taking the opposite view of this analysis as providing a *lower* bound on the mass of supermassive particles with given $\mathcal{O}(1)$ charges therefore constitutes an extrapolation well beyond the mass range considered in these references, and therefore the inequalities derived there must be taken with a grain of salt. Nevertheless, extrapolating the plots in figure 1 of [13] to larger mass values one sees that the bounds are actually much less constraining for masses close to the Planck scale. It then follows that gravitinos with charges (1) can be comfortably within the allowed range to be viable DM candidates if their mass is close to, but still below, the Planck scale (if the Schwarzschild radius of the particle were to exceed its Compton wave length, we would be dealing not with a stable particle, but with a mini-black hole which would immediately decay by Hawking radiation). As we argued in [1] the strongly interacting gravitinos would have mostly disappeared during the cosmic evolution (but could play a role in explaining ultra-high energy cosmic rays and the predominance of heavy ions in such events [16]). By contrast, the abundance of the color singlet gravitinos cannot be estimated since they were never in thermal equilibrium, but one can plausibly assume their abundance in first approximation to be given by the average DM density inside galaxies [17], *viz.*

$$\rho_{DM} \lesssim 0.3 \cdot 10^6 \text{ GeV} \cdot \text{m}^{-3} \quad (2)$$

(the average in the Universe is a million times smaller). If DM were entirely made out of nearly Planck mass particles with an assumed mass of 10^{18} GeV, this would amount to $\sim 3 \cdot 10^{-13}$ particles per cubic meter within galaxies. Furthermore assuming an average velocity of $30 \text{ km} \cdot \text{s}^{-1}$ this yields the flux estimate

$$\Phi \sim 0.03 \text{ m}^{-2} \text{ yr}^{-1} \text{ sr}^{-1} \quad (3)$$

We emphasize that, in addition to the unknown mass of the particle, there remain important uncertainties which could shift this estimate in either direction. One concerns possible inhomogeneities in the DM distribution within galaxies or stellar systems, which could lead to either a local depletion or to a local enhancement of the DM density (2) in the vicinity of the earth (see *e.g.* [18] for an early discussion of this point). The other is, of course, the average velocity of superheavy DM particles w.r.t. the earth (which is $\sim 30 \text{ km} \cdot \text{s}^{-1}$ if the DM particles are bound to the Solar System, and $\sim 300 \text{ km} \cdot \text{s}^{-1}$ if they are bound to our Galaxy). In the end this will be a matter for observation to decide.

A distinctive feature of the present proposal is that the DM gravitinos *do* participate in SM interactions with couplings of order $\mathcal{O}(1)$. In this sense, our DM candidates are not dark at all, but simply too faint to be seen directly because of their extremely low abundance! This is in contrast to other scenarios involving supermassive DM particles which are assumed to have only (super-)weak and gravitational interactions with SM matter [19–29], and which are mainly motivated by inflationary cosmology, whence the mass of those DM constituents would still be well below the Planck scale, on the order of the scale of inflation $\lesssim 10^{16}$ GeV. By contrast the gravitinos in (1) could in principle be detected if a way could be found to overcome their low abundance. A superheavy electrically charged particle could easily pass through the earth without deflection, leaving a very straight but tiny ionized track in the earth’s crust.

This leaves us basically with two options for discovery. Either one searches for traces of such tracks in old and very stable rock with a paleodetector, or otherwise one sets up an underground detector with sufficiently large fiducial area/volume and waits for the candidate particle to come by. The paleodetector option has been tried in the past with MICA samples [30, 31], again to search for magnetic monopoles; a general difficulty here is that the tracks would have to remain unaffected by geological processes over very long times, and the detection technique must be such as not to destroy the tracks (this favors MICA which comes with a naturally layered structure). The other and perhaps more promising option is to look for ionized tracks with suitable underground detectors and time of flight measurements, focusing on *slow* ionizing particles. The main background would come from cosmic ray muons, but a possible way to rule those out would be to look for slow particles *moving bottom up* which must have traversed a substantial part of the earth before being registered by the detector. In this context, the possible relevance of the MACRO experiment [10, 11] with its very large exposure time and surface area was already pointed out in [1], where it was suggested to have a second look at the data collected over many years. This is what we will now do, focusing on one special event.




FIG. 1: Energy loss [...] for the LIP events that passed the track quality and geometry cuts and satisfied the requirement of a maximum energy loss rate. [...] The signal region is in the $[0, 1.35]$ MeV/cm interval. Figure copied from [2].

III. THE MACRO EXPERIMENT AND A SPECIAL EVENT

The MACRO experiment [10, 11] was originally designed to search for magnetic monopoles, finishing with a null result after several years of taking data. We refer to the summary paper [11] for a detailed description of the experiment and of what the detector was capable of doing, as well as a summary of the collected results. The search covered a large part of parameter space, including the full range of velocities from relativistic particles down to ‘slow’ particles with $\beta \sim 4 \times 10^{-5}$, coming in from all directions. In this way the detector was able to search not only for magnetic monopoles, but also for other, and unknown kinds of ionizing particles, including fractionally charged particles. Results of the latter search which concentrated specifically on *lightly ionizing particles* (LIPs) appeared in a separate publication [2].

While [11] mentions 40 events (out of a total of about 35 000) that were subsequently discarded as spurious and not further discussed, Ref. [2] reports one special event of a type different from an expected monopole signal. The relevant information about this event is contained in figure 3 of [2] which we here reproduce for the reader’s convenience, together with its figure caption. Let us also quote excerpts from the accompanying part of the text in [2] which says: “As one can see [...] there is one event (run 15871, event 5649) that appears in the signal region. It corresponds to a maximum energy loss of 0.66 MeV/cm, i.e., about 20% lower than expected for a particle of charge $2e/3$ and about a factor of 3 higher than what expected for a particle of charge $e/3$. Three scintillator counters were involved in this trigger; the first in one of the upper vertical layers, the second in the central horizontal layer and the third in the lower horizontal layer. [...] The position along the counter for this particular box measured by the PHRASE and by the streamer tube track geometry were in agreement (within 15 cm). We have examined this event by hand relying primarily on the wave forms as recorded for all the counters involved in the trigger. The apparent amplitude of the recorded wave forms was consistent with the energy thresholds [...]. Having three scintillator counters involved in the trigger we have checked for a consistency in the relative timing of them with the crossing of a single particle of constant velocity. The relative timing between the counter in the upper part of the detector and that in the central part was consistent with the passage of a relativistic particle coming from above while the relative timing between the

box in the lower part of the detector and any of the other two hits was consistent with a slowly moving upward-going particle. We thus discarded this event from the signal region.”

The main reason for discarding this event was therefore not some obvious instrumental malfunction, but rather the fact that there appears to be no way to reconcile all three scintillator signals (as well as the fact that the signal does not conform with expectations for a magnetic monopole [10, 11]), whence one concludes that one of the three signals must be ascribed to a different origin. A crucial additional fact is that there is unmistakable and independent evidence from the streamer tubes for a single particle track running through the whole detector. The ‘obvious’ interpretation of this event would seem to be in terms of a relativistic particle corresponding to a cosmic ray muon coming from above, and this remains perhaps the more plausible explanation. However, this interpretation would not only require disregarding the earlier scintillator signal in the lower part of the detector, but also explaining why a third signal in the same scintillator, coincident with the central and upper scintillators, is missing. It would furthermore require an explanation why the energy loss rate is well below the allowed minimum in Fig. 1 corresponding to the minimum ionization point of the Bethe-Bloch curve [32] (which is reached for $\beta\gamma \sim 5$). A charge $|q| = \frac{2}{3}$ (or less) is simply not compatible with a muon.

By contrast, the alternative second interpretation with a slow particle moving upward would require only the signal in the upper scintillator to be due to some other cause. While it appears that the question cannot be finally resolved on the basis of the existing MACRO data, we here wish to raise attention to the possibility that the signal in the upper layer could have masked the presumed later arrival of the slow particle in the upper detector. As we argue below, a slow particle would not have been registered at all without such an accompanying signal, on account of the trigger vetoes imposed for the LIPs search! The event could then correspond to the passage of a ‘slow’ charged particle through the MACRO detector. To see why such a particle would be only lightly ionizing we recall that the ionization rate increases with decreasing β like $\propto \beta^{-2}$, but then drops again rapidly for $\beta < 0.01$ (Lindhard-Scharff regime) [32]. Even without a full fledged quantum mechanical calculation, and perhaps oversimplifying things a bit, this can be understood by means of a very simple classical argument. For a head-on collision of a supermassive particle of speed β with an electron, the maximum velocity change imparted to the electron would be 2β , whence the maximum energy carried away by the electron would be $2m_e\beta^2 \sim 1 \text{ MeV} \times \beta^2$. For $\beta \lesssim 10^{-3}$ this is much less than the minimum ionization energy required to free the electron from the atom (a proper quantum mechanical treatment, which is currently not available, would need to take into account the interaction of the supermassive particle with the whole cloud of electrons). Besides, for this range of β the loss of energy by elastic collisions gets more and more important with decreasing β . However, for a slow positively charged supermassive particle another process could become relevant: due to its large mass its motion is not affected at all by the atom, while thanks to its slow motion it can spend enough time in the electron cloud to ‘drag along’ an electron, forming a lightly bound fractionally charged state that moves along with unchanged speed. Having removed the electron from the atom, it thus leaves behind an ion that can be detected by a streamer tube or a drift chamber, but not a scintillator. This might explain why all streamer tubes registered the track, but not all scintillators produced accompanying signals for the event in question. Otherwise, in this regime, the energy losses would be dominated by nuclear recoils (which is not what MACRO was looking for), which incidentally offers another possible avenue for testing our hypothesis. Furthermore, in the absence of more detailed calculations, the charge of the particle can no longer be straightforwardly deduced.

If we therefore proceed with this assumption we have two additional indications for the correctness of our hypothesis, as emphasized in the above quote from [2], namely

- the track as seen by the streamer tubes was consistent with the tracks in the lower and central scintillators; and
- the time of flight was consistent with a slow particle moving bottom up in the lower and central scintillators

With these additional consistency checks let us re-iterate that a ‘slow’ particle moving bottom up would be difficult to explain in terms of known physics. First of all, going up, it obviously cannot be a muon. Second, it cannot be a magnetic monopole, nor a dyon, because monopoles generally are not expected to be lightly ionizing because of their large magnetic charge, although they may in principle be able to traverse the earth [33]. This is also the energy range where the relation between the energy loss and the light yield (which is what is measured by the photomultipliers) is no longer linear [34]. The same comment applies to dyons for which the energy loss would be even bigger (in principle an electrically neutral monopole can acquire a very small electric charge proportional to the CP violating θ -angle by means of the Witten effect [35]; however, given the known upper limits on the value of θ the ionization would be dominated by magnetic interactions). Third, the full track as reconstructed from the streamer tubes cannot be the result of a radioactive decay in the surrounding rock, since such products have energies of at most several MeV, so they could not penetrate the scintillator more deeply than a few centimeters. Therefore a superheavy fractionally charged particle seems to be the most plausible explanation if one adopts our hypothesis.

While [2] does not explicitly quantify what ‘slow’ means a more precise knowledge of the velocity would be useful as it would enable us to make a first estimate of the expected gravitino flux. If the MACRO detector had been set up to

search for slow particles such as a supermassive gravitino, the expected event rate according to the estimate (3) would amount to several events per year. However, the trigger used in [2] was sensitive only to *fast* lightly ionizing particles. As stated in [2]: “Particles which have a velocity lower than $0.25c$ are not guaranteed to pass through the detector quickly enough to insure that the LIP trigger will detect a coincidence in the faces of the scintillator system”. Hence without an accompanying coincident triggering signal any slower particle would have escaped detection, and thus possible events involving *only* a slow particle could have been missed. Therefore the rate of one event for the five-year cycle covered by [2] could be a significant underestimate of the actual abundance and flux rate for supermassive gravitinos. Conversely, the trigger settings used for the original monopole search [10, 11] which did include velocity ranges down to $\beta \sim 4 \cdot 10^{-5}$, did not allow for lightly ionizing particles, thus explaining why no further events of this type were observed, even for a flux close to the estimated value (3). For this reason we cannot at this point reliably deduce the actual gravitino flux in the vicinity of the Earth from the given data. This, as well as the confirmation (or refutation) of our hypothesis, would require a dedicated new experiment, concentrating on slow and lightly ionizing particles.

IV. CONCLUSIONS

Re-inspection of the special MACRO event reported in [2] has revealed the possibility of an explanation different from the ‘obvious’ one in terms of a cosmic ray muon, namely

- a slow particle *moving bottom up*, which thus must have traversed a substantial part of the earth; and
- partial evidence that this particle carries fractional charge.

Nevertheless, in order to find out whether this outlier event is real physics or just a fluke, and to proceed from circumstantial evidence to an actual discovery, further and independent confirmation is obviously needed, and we hope that the present paper can motivate a dedicated effort that will decide the issue in the not-so-distant future.

On the other hand, corroboration of our new interpretation of this event would have dramatic implications. In particular, it would bring $N = 8$ supergravity back into focus for unification, although in an unexpected way. We emphasize that the considerations leading to (1) are so far purely kinematical, and that the dynamics underlying the present scheme remains unknown, possibly requiring a framework beyond space-time based quantum field theory. Nevertheless our findings may indicate that $N = 8$ supergravity could be closer to the truth than is widely thought (as is also suggested by the finiteness properties of the theory [36] and various anomaly cancellations [37–40]). We also note that the spurion shift required to match the spin- $\frac{1}{2}$ sectors of the theory with three generations of quarks and leptons [3], and here extended to the gravitinos [6], appears to be incompatible with space-time supersymmetry. This could mean that, contrary to many expectations, (maximal) space-time supersymmetry might not be a relevant concept for unification after all, but, through its fermionic (spin- $\frac{1}{2}$ and spin- $\frac{3}{2}$) content, merely a theoretical crutch to guide us to the right answer.

Acknowledgments: We would like to thank Barry Barish for helpful correspondence. Furthermore, we are greatly indebted to Erik Katsavounidis for alerting us to [2] and the strange outlier event, and for many explanations and clarifications concerning the MACRO experiment without which this article would not have seen the light of the day. K.A. Meissner was partially supported by the Polish National Science Center grant UMO-2020/39/B/ST2/01279. The work of H. Nicolai has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 740209).

-
- [1] K.A. Meissner and H. Nicolai, Phys. Rev. **D100** (2019) 035001
 - [2] M. Ambrosio *et al*, *Final search for lightly ionizing particles with the MACRO detector*, [arXiv:hep-ex/0402006](https://arxiv.org/abs/hep-ex/0402006)
 - [3] M. Gell-Mann, in Proceedings of the 1983 Shelter Island Conference on Quantum Field Theory and the Fundamental Problems of Physics, eds. R. Jackiw, N.N. Khuri, S. Weinberg and E. Witten, Dover Publications, Mineola, New York (1985)
 - [4] H. Nicolai and N.P. Warner, Nucl. Phys. **B259** (1985) 412
 - [5] K.A. Meissner and H. Nicolai, Phys. Rev. **D91** (2015) 065029
 - [6] K.A. Meissner and H. Nicolai, Phys. Rev. Lett. **121** (2018) 091601
 - [7] A. Kleinschmidt and H. Nicolai, Phys. Lett. **B747** (2015) 251

- [8] E. Cremmer and B. Julia, Nucl. Phys. **B159** (1979) 141
- [9] B. de Wit and H. Nicolai, Nucl. Phys. **B208** (1982) 323
- [10] M. Ambrosio et al. (MACRO coll.), Phys. Rev **D62** (2000) 052003
- [11] M. Ambrosio et al. (MACRO coll.), Eur.Phys.J. **C25** (2002) 511
- [12] S. Chang, C. Coriano and A.E. Faraggi, Phys. Lett. **B397** (1997), 76;
L. Delle Rose, A.E. Faraggi, C. Marzo, J. Rizos, Phys.Rev. **D96** (2017) 055025
- [13] S. D. McDermott, H. B. Yu and K. M. Zurek, Phys. Rev. **D83**(2011) 063509
- [14] A. D. Dolgov, S. L. Dubovsky, G. I. Rubtsov and I. I. Tkachev, Phys. Rev. **D88**(2013)117701
- [15] E. Del Nobile, M. Nardecchia and P. Panci, JCAP **1604** (2016) no.04, 048
- [16] K.A. Meissner and H. Nicolai, JCAP **09** (2019) 041
- [17] M. Weber and W. de Boer, Astron.Astrophys. **509** (2010) A25
- [18] T. Damour and L.M. Krauss, Phys. Rev. **D59** (1999) 063509
- [19] A. De Rújula, S.L. Glashow and U. Sarid, Nucl.Phys. **B333** (1990) 173
- [20] C. Burrage, J. Jaeckel, J. Redondo and A. Ringwald, JCAP **0911** (2009) 002
- [21] L. Chuzhoy and E.W. Kolb, JCAP **0907** (2009) 014
- [22] M.A. Markov, JETP **24** (1967) 584
- [23] Y. Aharonov, A. Casher and S. Nussinov, Phys.Lett **B191** (1987) 51
- [24] A.M. Srivastava, Phys.Rev. **D36** (1987) 2368
- [25] D.J.H. Chung, E.W. Kolb and A. Riotto, Phys. Rev. **D59** (1998) 023501; D.J.H. Chung, P. Crotty, E.W. Kolb and A. Riotto, Phys. Rev. **D64** (2001) 043503
- [26] E.W. Kolb, A.A. Starobinsky and I.I. Tkachev, JCAP **0707** (2007) 005
- [27] M. Garny, McCullen Sandora and M.S. Sloth, Phys. Rev. Lett. **116** (2016) 101302
- [28] M. Garny, McCullen Sandora and M.S. Sloth, [arXiv:1709.09688](#)
- [29] K. Kannike, A. Racioppi and M. Raidal, [arXiv:1605.09378 \[hep-ph\]](#)
- [30] P.B. Price and M.H. Salamon, Phys. Rev. Lett. **56** (1986) 1226
- [31] J.F. Acevedo, J. Bramante and A. Goodman, [arXiv:2205.06473 \[hep-ph\]](#)
- [32] Particle Data Group, Nucl. and Part. Phys. **33** (July 2006) pp. 258-270
- [33] J. Derkaoui et al., Astroparticle Physics **9** (1998) 173
- [34] J. Derkaoui et al., Astroparticle Physics **10** (1999) 339
- [35] E. Witten, Phys. Lett. **B86** (1979) 283
- [36] Z. Bern, J.J. Carrasco, W.M. Chen, A. Edison, H. Johansson, J. Parra-Martinez, R. Roiban and M. Zeng, Phys. Rev. **D98** (2018) 086021
- [37] N. Marcus, Phys.Lett. **B157** (1985) 383
- [38] J.J.M. Carrasco, R. Kallosh, R. Roiban and A.A. Tseytlin, JHEP **1307** (2013) 029
- [39] Z. Bern, S. Davies and T. Dennen, Phys.Rev. **D90** (2014) 105011
- [40] K.A. Meissner, and H. Nicolai, Phys. Lett. **B772** (2017) 169